

ENTANGLED QUANTUM STATES OF CAUSAL FERMION SYSTEMS AND UNITARY GROUP INTEGRALS

FELIX FINSTER, NIKY KAMRAN, AND MORITZ REINTJES

JULY 2023

ABSTRACT. This paper is dedicated to a detailed analysis and computation of quantum states of causal fermion systems. The mathematical core is to analyze integrals over the unitary group asymptotically for a large dimension of the group, for various integrands with a specific scaling behavior in this dimension. It is shown that, in a well-defined limiting case, the localized refined pre-state is positive and allows for the description of general entangled states.

CONTENTS

1. Introduction and Overview of Main Results	2
2. Preliminaries	5
2.1. Basics on Causal Fermion Systems	5
2.2. Notions of Quantum States of Causal Fermion Systems	17
2.3. Basics on Fock Spaces and Entanglement	22
3. Quantum States of Causal Fermion Systems and Entanglement	23
3.1. General Considerations on the Quantum State	23
3.2. Why Entanglement Requires the Refined Pre-State	25
3.3. Why it is Preferable to Work with Localized States	28
3.4. The Precise Form of the Localized Refined Pre-State	28
4. Computation of Integrals over the Unitary Group	30
4.1. Preliminary Considerations	30
4.2. Integrals over Functions on a Subspace	33
4.3. A Few Simple Examples	37
4.4. The Gaussian Asymptotics	40
4.5. Treating Integrands which Depend on the Dimension	41
4.6. Asymptotic Decoupling on Orthogonal Subspaces	44
5. Model Examples	48
5.1. An Exponential of a Linear Functional	49
5.2. An Exponential of a Quartic Functional	54
6. Computation of the Localized Refined Pre-State	62
6.1. The Saddle Point Asymptotics for the High-Energy Wave Functions	62
6.2. Introducing a Weight for the Size of the Subsystems	71
6.3. The Phase Freedom of the Saddle Points	73
6.4. Saddle Points for the Low-Energy Wave Functions	74
6.5. The Structure of the Saddle Points	77
6.6. Construction of the Insertions	78
6.7. Positivity of the Localized Refined Pre-State	80

7. Description of Entanglement	82
Appendix A. Diagrammatic Derivation of the Gaussian Asymptotics	83
References	87

1. INTRODUCTION AND OVERVIEW OF MAIN RESULTS

The theory of *causal fermion systems* is a recent approach to fundamental physics (see the basics in Section 2, the reviews [24, 14, 17], the textbooks [13, 23] or the website [1]). In this approach, spacetime and all objects therein are described by a measure ρ on a set \mathcal{F} of linear operators on a Hilbert space $(\mathcal{H}, \langle \cdot | \cdot \rangle_{\mathcal{H}})$. The physical equations are formulated by means of the so-called *causal action principle*, a nonlinear variational principle where an action \mathcal{S} is minimized under variations of the measure ρ . The present paper is a major step in an ongoing research program aimed at giving a conceptually clear and mathematically sound *derivation of quantum field theory* within the theory of causal fermion systems. In simple terms, the goal of this program is to show that, in a well-defined limiting case, the dynamics of a causal fermion system can be described as in quantum field theory by a unitary time evolution on bosonic and fermionic Fock spaces. This program was initiated in [12], where quantum electrodynamics was obtained from the causal action principle under certain simplifying assumptions and approximations, which were stated but not yet justified in a fully satisfying way. The first step in the detailed treatment based on the mathematical structures of causal fermion systems is [19], where a connection between causal variational principles and a dynamics on bosonic Fock spaces was established. More recently, in [20] it was shown that a causal fermion system in a Minkowski-type spacetime gives rise to a distinguished quantum state. Moreover, various modifications and refinements of this construction were given. The aim of the present paper is to analyze the different definitions in detail and to identify the construction which is most suitable for the description of entanglement. To this end, we need to delve deeper into the question of how to compute these quantum states.

Our main objective is to show that general entangled states can indeed be described with the help of the *localized refined partition function* Z_V^t , which can be written as

$$Z_V^t(\alpha, \beta, \tilde{\rho}) := \int_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_{<}) \int_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_{>}) e^{\alpha N \mathcal{T}_V^t(\tilde{\rho}, T\mathcal{U}_{<}, \mathcal{U}_{>} \rho)}. \quad (1.1)$$

Before giving an overview of our results, we make a few comments on the structure of Z_V^t . In the above formula, $\tilde{\rho}$ is the measure describing the interacting spacetime, whereas ρ describes the Minkowski vacuum. The partition function is a double integral over a compact group $\mathcal{G} \simeq \mathrm{U}(N)$ of unitary operators on the Hilbert space \mathcal{H} (here $\mu_{\mathcal{G}}$ is the Haar measure on this group, and by \int we always denote a normalized integral). The integrand involves the exponential of the functional \mathcal{T}_V^t which relates the interacting and vacuum measures at time t within a bounded spatial region described by a subregion V of the vacuum spacetime (see Figure 1, where the support of the measure $M := \mathrm{supp} \rho$ is the vacuum spacetime, and Ω^t is the region in the past of time t ; for details see Section 2 below). The functional \mathcal{T}_V^t has the mathematical structure of a *nonlinear surface layer integral* (see (2.42) and the preliminaries in Section 2.1.12).




FIGURE 1. Decomposition of the vacuum spacetime in the localized state.

The *localized refined pre-state* ω_V^t can be written symbolically as

$$\omega_V^t(\dots) := \frac{1}{Z_V^t(\alpha, \beta, \tilde{\rho})} \oint_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_{<}) \oint_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_{>}) e^{\alpha N \mathcal{T}_V^t(\tilde{\rho}, T_{\mathcal{U}_{<}, \mathcal{U}_{>}} \rho)}(\dots), \quad (1.2)$$

where the dots on the left stand for an operator in the observable algebra \mathcal{A} formed of the linearized fields in the vacuum spacetime. The dots on the right, on the other hand, stand for suitable surface layer integrals which again involve the linearized fields in the vacuum. The structure of (1.2) has some resemblance with the path integral formulation of quantum theory, where the n -point functions are obtained by integrating an exponential of the classical action over field configurations, taking the fields as insertions. In this formalism, even the density operator has been constructed in [45]. However, the similarity to the path integral formalism does not seem to extend beyond a formal analogy, because in (1.2) we integrate over unitary operators, not over field configurations. Moreover, the surface layer integral \mathcal{T}_V^t in the exponent is not the classical action (or the causal action), but instead a surface layer integral which can be understood as a device for “comparing” the measures $\tilde{\rho}$ and ρ . In this “comparison” it is important that the unitary operators $\mathcal{U}_{<}$ and $\mathcal{U}_{>}$ come up in a nonlinear way.

In general terms, the localized refined pre-state describes the interacting measure $\tilde{\rho}$ using the familiar objects of the vacuum spacetime (linear bosonic and fermionic fields). The measure $\tilde{\rho}$ should be thought of as having a very complicated structure, both on microscopic and macroscopic scales. More specifically, all the objects of the interacting physical system are encoded in the *physical wave functions*, being a family of spinorial wave functions in spacetime. The collective behavior of all these wave functions gives rise to the usual spacetime structures (causality, metric, particles, fields, etc.). Due to the mutual interaction of all the wave functions, this collective behavior can be intricate, including *long-range correlations* and *dephasing effects* between sub-families of waves propagating in different spatial directions. The role of the integration over the unitary group is to detect all these phenomena as encoded in the interacting measure $\tilde{\rho}$ and to quantify them in the familiar language of quantum field theory. The interplay of these effects gives rise to entanglement, as will be worked out and made precise in this paper (see the discussion at the end of Section 7).

In order to get into a well-defined limiting case in which the above integrals over the unitary group can be computed, we consider the following asymptotics:

- (i) The causal fermion systems involve an ultraviolet cutoff on a length scale ε (which can be thought of as the Planck length). We consider the asymptotics $\varepsilon \searrow 0$, while keeping the length scales of macroscopic physics fixed.
- (ii) For any given $\varepsilon > 0$, we also consider the asymptotics $N \rightarrow \infty$ where the dimension of the Hilbert space \mathcal{H} tends to infinity. Since the parameter N also appears in the

exponent of Z_V^t and ω_V^t , this also amounts to a suitable rescaling of the nonlinear surface layer integral. Since \mathcal{H} describes the system also outside the spacetime region V , the limit $N \rightarrow \infty$ can be thought of as the infinite volume limit when the size of the whole system tends to infinity, while keeping the bounded spacetime region V fixed.

In physical terms, the quantum state tells us about the outcome of measurements performed at time t in the spacetime region V . With this in mind, the above asymptotics reflect the physical facts that the Planck length is much smaller than all other relevant length scales (point (i)) and that a physical measurement takes place in a spatial region which is typically much smaller than the size of the universe (point (ii)). This asymptotics will be worked out in the main sections of our paper using Gaussian integrals and saddle points techniques (Sections 4–6).

We now outline our main results. We first prove that, in the above limiting case, the localized refined pre-state is positive, i.e.

$$\omega_V^t : \mathcal{A} \rightarrow \mathbb{C} \quad \text{with} \quad \omega_V^t(A^*A) \geq 0 \quad \text{for all } A \in \mathcal{A} \quad (1.3)$$

(see Theorem 6.8 in Section 6.7). Thus it is a *quantum state* as used in the algebraic formulation of quantum field theory. Next, we show that our quantum state allows for the general description of *entanglement* (Section 7). We even get a concrete prescription for how to encode a given entangled Fock state in the measure $\tilde{\rho}$ (see Section 7). We remark that the usual notion of a quantum state described by a density operator acting on a Hilbert space is obtained from the algebraic formulation (1.3) by constructing representations of the field algebra (for details see [20, Section 4.5]). Then the quantum state ω^t is obtained from the density operator by taking the expectation value, i.e.

$$\omega^t(A) = \text{tr}_{\mathcal{F}}(\sigma^t A) \quad \text{for all } A \in \mathcal{A}.$$

This density operator gives all the familiar structures of quantum field theory. In particular, the density operator gives rise to notions of *entropy* like the von Neumann entropy $S = -\text{tr}_{\mathcal{F}}(\sigma^t \log \sigma^t)$, the relative entropy and the entanglement entropy of a spatial subregion. We remark that the only but very important point which at this stage is still missing compared to standard quantum field theory is the *dynamics* of the quantum state. This will be worked out in detail in the upcoming paper [9].

The mathematical core of our analysis is the computation of specific integrals over the unitary group, asymptotically when the dimension of the group gets large. Similar integrals have been studied in the context of *random matrix theory* and *lattice gauge theories* (see for example the standard textbooks [41, 46]). Despite similarities to problems studied in this context (like the Harish-Chandra integral [41, 40], the Itzykson-Zuber model [39], the Gaussian asymptotics of group integrals in [49, 6, 38] and diagrammatic approaches [7]), these results do not immediately apply to our problem. The main difference is that, in our case, the integrand has an explicit dependence on the dimension of the group, changing the asymptotics as this dimension tends to infinity. We take advantage of the fact that our integrand only depends on the matrix entries on a fixed subspace. This makes it possible to carry out the integral over all other matrix entries, giving the reduction formula in Theorem 4.2 (for related results see [42, 32, 33, 3]). From this formula one can deduce the Gaussian asymptotics (see Proposition 4.10) as well as the leading asymptotics if the integrand is a product depending on the matrix entries of two orthogonal subspaces (see Propositions 4.11

and 4.12). We then apply these results to specific integrands involving exponentials of the matrix elements (see Proposition 5.1 and Theorem 5.6).

We finally comment on the physical picture behind our constructions. In the usual description of quantum theory, the expectation value of a measurement is given by the quantum state applied to the corresponding observables. With this in mind, the construction of a quantum state of a causal fermion system can be understood as the preparation of a measurement device which can be used for performing certain measurements, where the notion of a “measurement device” is meant more generally not for an experimental apparatus, but rather for a mathematical procedure for extracting information from the causal fermion system. In this analogy, the freedom in the construction of the state is not problematic or surprising; it can be understood similar to the fact that different experimental setups can be used to measure the same physical quantity. From this perspective, we need to address the question which construction captures the physical essence of quantum fields, including entanglement. Answering this question also shows that the theory of causal fermion systems is indeed capable of describing the effects of quantum field theory.

The paper is organized as follows. After giving the necessary preliminaries on causal fermion systems, quantum states and Fock spaces (Section 2), in Section 3 we begin with qualitative considerations which explain why it is preferable to consider the localized refined pre-state (Sections 3.1–3.3). We also specify the precise mathematical setup (Section 3.4). In Section 4 we review the general methods which will be used later in this paper for the computation of the group integrals. In Section 5 we consider group integrals of exponentials with a specific scaling in the dimension N . These integrals have a similar structure as the refined partition function (1.1). They can be used for the detailed analysis of the localized refined pre-state (1.2), as is worked out in Section 6. In Section 7 it is shown how entangled states can be described. In Appendix A an alternative method is given for computing group integrals. This method is superseded by the stronger and more suitable methods in Section 4.4. We explain it nevertheless, because it gives an alternative way of understanding why group integrals simplify when evaluating them asymptotically for large N .

2. PRELIMINARIES

2.1. Basics on Causal Fermion Systems. This section provides the necessary background on causal fermion systems. Here we introduce all the basic structures needed later on. We keep the presentation brief and refer for details to corresponding articles and textbooks.

2.1.1. Causal Fermion Systems and the Causal Action Principle. We begin with the abstract definitions.

Definition 2.1. (*causal fermion systems*) Given a separable complex Hilbert space \mathcal{H} with scalar product $\langle \cdot | \cdot \rangle_{\mathcal{H}}$ and a parameter $n \in \mathbb{N}$ (the “*spin dimension*”), we let $\mathcal{F} \subset L(\mathcal{H})$ be the set of all symmetric¹ operators on \mathcal{H} of finite rank, which (counting multiplicities) have at most n positive and at most n negative eigenvalues. On \mathcal{F} we are given a positive measure ρ (defined on a σ -algebra of subsets of \mathcal{F}). We refer to $(\mathcal{H}, \mathcal{F}, \rho)$ as a *causal fermion system*.

¹Here by a symmetric operator A we mean that $\langle Au | v \rangle_{\mathcal{H}} = \langle u | Av \rangle_{\mathcal{H}}$ for all $u, v \in \mathcal{H}$. For bounded operators as considered here, the notions “symmetric” and “selfadjoint” coincide.

A causal fermion system describes a spacetime together with all structures and objects therein. In order to single out the physically admissible causal fermion systems, one must formulate physical equations. To this end, we impose that the measure ρ should be a minimizer of the causal action principle, which we now introduce. For any $x, y \in \mathcal{F}$, the product xy is an operator of rank at most $2n$. However, in general it is no longer a symmetric operator because $(xy)^* = yx$, and this is different from xy unless x and y commute. As a consequence, the eigenvalues of the operator xy are in general complex. We denote these eigenvalues counting algebraic multiplicities by $\lambda_1^{xy}, \dots, \lambda_{2n}^{xy} \in \mathbb{C}$ (more specifically, denoting the rank of xy by $k \leq 2n$, we choose $\lambda_1^{xy}, \dots, \lambda_k^{xy}$ as all the non-zero eigenvalues and set $\lambda_{k+1}^{xy}, \dots, \lambda_{2n}^{xy} = 0$). We introduce the Lagrangian and the causal action by

$$\text{Lagrangian:} \quad \mathcal{L}(x, y) = \frac{1}{4n} \sum_{i,j=1}^{2n} \left(|\lambda_i^{xy}| - |\lambda_j^{xy}| \right)^2 \quad (2.1)$$

$$\text{causal action:} \quad \mathcal{S}(\rho) = \iint_{\mathcal{F} \times \mathcal{F}} \mathcal{L}(x, y) d\rho(x) d\rho(y). \quad (2.2)$$

The *causal action principle* is to minimize \mathcal{S} by varying the measure ρ under the following constraints,

$$\text{volume constraint:} \quad \rho(\mathcal{F}) = \text{const} \quad (2.3)$$

$$\text{trace constraint:} \quad \int_{\mathcal{F}} \text{tr}(x) d\rho(x) = \text{const} \quad (2.4)$$

$$\text{boundedness constraint:} \quad \iint_{\mathcal{F} \times \mathcal{F}} |xy|^2 d\rho(x) d\rho(y) \leq C, \quad (2.5)$$

where C is a given parameter, tr denotes the trace of a linear operator on \mathcal{H} , and the absolute value of xy is the so-called spectral weight,

$$|xy| := \sum_{j=1}^{2n} |\lambda_j^{xy}|. \quad (2.6)$$

This variational principle is mathematically well-posed if \mathcal{H} is finite-dimensional. For the existence theory and the analysis of general properties of minimizing measures we refer to [11, 4] and [23, Chapter 12]. In the existence theory one varies in the class of regular Borel measures (with respect to the topology on $L(\mathcal{H})$ induced by the operator norm), and the minimizing measure is again in this class. With this in mind, here we always assume that ρ is a *regular Borel measure*.

2.1.2. Spacetime and Physical Wave Functions. Let ρ be a *minimizing measure*. *Spacetime* is defined as the support of this measure,

$$M := \text{supp } \rho.$$

Thus the spacetime points are symmetric linear operators on \mathcal{H} . On M we consider the topology induced by \mathcal{F} (generated by the operator norm on $L(\mathcal{H})$). Moreover, the measure $\rho|_M$ restricted to M gives a volume measure on spacetime. This makes spacetime into a *topological measure space*.

The operators in M contain a lot of information which, if interpreted correctly, gives rise to spacetime structures like causal and metric structures, spinors and interacting

fields (for details see [13, Chapter 1]). Here we restrict attention to those structures needed in this paper. We begin with a basic notion of causality:

Definition 2.2. (*causal structure*) For any $x, y \in \mathcal{F}$, the product xy is an operator of rank at most $2n$. We denote its non-trivial eigenvalues (counting algebraic multiplicities) by $\lambda_1^{xy}, \dots, \lambda_{2n}^{xy}$. The points x and y are called *spacelike* separated if all the λ_j^{xy} have the same absolute value. They are said to be *timelike* separated if the λ_j^{xy} are all real and do not all have the same absolute value. In all other cases (i.e. if the λ_j^{xy} are not all real and do not all have the same absolute value), the points x and y are said to be *lightlike* separated.

Restricting the causal structure of \mathcal{F} to M , we get causal relations in spacetime.

Next, for every $x \in \mathcal{F}$ we define the *spin space* S_x by $S_x = x(\mathcal{H})$; it is a subspace of \mathcal{H} of dimension at most $2n$. It is endowed with the *spin inner product* $\prec \cdot | \succ_x$ defined by

$$\prec u | v \succ_x = -\langle u | xv \rangle_{\mathcal{H}} \quad (\text{for all } u, v \in S_x).$$

A *wave function* ψ is defined as a function which to every $x \in M$ associates a vector of the corresponding spin space,

$$\psi : M \rightarrow \mathcal{H} \quad \text{with} \quad \psi(x) \in S_x M \quad \text{for all } x \in M.$$

A wave function ψ is said to be *continuous* at x if for every $\varepsilon > 0$ there is $\delta > 0$ such that

$$\|\sqrt{|y|}\psi(y) - \sqrt{|x|}\psi(x)\|_{\mathcal{H}} < \varepsilon \quad \text{for all } y \in M \text{ with } \|y - x\| \leq \delta \quad (2.7)$$

(where $|x|$ is the absolute value of the symmetric operator x on \mathcal{H} , and $\sqrt{|x|}$ is the square root thereof). Likewise, ψ is said to be continuous on M if it is continuous at every $x \in M$. We denote the set of continuous wave functions by $C^0(M, SM)$ (where $SM := \cup_{x \in M} S_x M$ generalizes the spinor bundle; for details see [18, Section 3]).

It is an important observation that every vector $u \in \mathcal{H}$ of the Hilbert space gives rise to a unique wave function. To obtain this wave function, denoted by ψ^u , we simply project the vector u to the corresponding spin spaces,

$$\psi^u : M \rightarrow \mathcal{H}, \quad \psi^u(x) = \pi_x u \in S_x M.$$

We refer to ψ^u as the *physical wave function* of $u \in \mathcal{H}$. A direct computation shows that the physical wave functions are continuous (in the sense (2.7)). Associating to every vector $u \in \mathcal{H}$ the corresponding physical wave function gives rise to the *wave evaluation operator*

$$\Psi : \mathcal{H} \rightarrow C^0(M, SM), \quad u \mapsto \psi^u. \quad (2.8)$$

Evaluating at a fixed spacetime point $x \in M$, we obtain a corresponding mapping

$$\Psi(x) : \mathcal{H} \rightarrow S_x M, \quad u \mapsto \psi^u(x).$$

Every $x \in M$ can be written as (for the derivation see [13, Lemma 1.1.3])

$$x = -\Psi(x)^* \Psi(x). \quad (2.9)$$

In words, every spacetime point operator is the *local correlation operator* of the wave evaluation operator at this point (for details see [13, §1.1.4 and Section 1.2]).

2.1.3. *The Kernel of the Fermionic Projector.* For computations, it is most convenient to work with the *kernel of the fermionic projector* $P(x, y)$ which can be defined in terms of the wave evaluation operator (2.8) by

$$P(x, y) := -\Psi(x) \Psi(y)^* : S_y \rightarrow S_x . \quad (2.10)$$

It can be regarded as the basic object of the theory because the Lagrangian as well as the constraints can be computed from it. This is based on the observation that the non-trivial eigenvalues of the operator product xy coincide with the eigenvalues of the *closed chain* A_{xy} defined by

$$A_{xy} := P(x, y) P(y, x) : S_x \rightarrow S_x$$

(for details see for example [13, §1.1.3]). First variations of the Lagrangian can be written as (for details see [13, §1.4.1])

$$\delta \mathcal{L}(x, y) = 2 \operatorname{Re} \operatorname{Tr}_{S_y} (Q(y, x) \delta P(x, y)) , \quad (2.11)$$

where $Q(x, y)$ is a symmetric kernel, i.e.

$$Q(x, y) : S_y \rightarrow S_x \quad \text{and} \quad Q(x, y)^* = Q(y, x) . \quad (2.12)$$

We will use this variational formula in Section 6.4.

2.1.4. *Causal Fermion Systems in Minkowski Space.* The prime examples of causal fermion systems are constructed from Dirac wave functions in Minkowski space. For these systems, the causal action principle has been studied in detail in [13]. In particular, it is known that in a specific limiting case, referred to as the *continuum limit*, the measure describing the Minkowski vacuum is indeed a minimizer of the causal action principle. Moreover, the continuum limit analysis developed in [13] associates a specific class of critical measures with systems of Dirac particles in Minkowski space which interact via classical bosonic fields which satisfy classical field equations.

We now recall a few basics on causal fermion systems in Minkowski space, which will be needed in the later computations in Section 6. More detailed introductions can be found in [13, Section 1.2] or [23, Chapter 5]. In order to describe the Minkowski vacuum, one considers the Hilbert space $(\mathcal{H}, \langle \cdot | \cdot \rangle_m)$ of solutions of the Dirac equation $(i\partial\!\!\!/ - m)\psi = 0$ of mass m with the scalar product

$$(\phi | \psi)_m := \int_{\mathbb{R}^3} \prec \phi | \gamma^0 \psi \succ(t, \vec{x}) d^3x$$

(where $\prec \cdot | \cdot \succ$ is the spin inner product of signature $(2, 2)$; this scalar product does not depend on time due to current conservation). Next, we choose $(\mathcal{H}, \langle \cdot | \cdot \rangle_{\mathcal{H}})$ as the subspace of \mathcal{H} of all solutions of negative energy (i.e. negative frequency) with the induced scalar product. For the construction of a corresponding causal fermion system, one needs to introduce an *ultraviolet regularization* on a microscopic length scale ε , which can be thought of as the Planck length. To this end, one introduces a so-called *regularization operator* \mathfrak{R}_ε as an operator which maps \mathcal{H} to the continuous wave functions,

$$\mathfrak{R}_\varepsilon : \mathcal{H} \rightarrow C^0(\mathcal{M}, S\mathcal{M}) .$$

Next, for any $x \in \mathcal{M}$ one introduces the *local correlation operator* F^ε by

$$F^\varepsilon(x) := -\mathfrak{R}_\varepsilon(x)^* \mathfrak{R}_\varepsilon(x) : \mathcal{H} \rightarrow \mathcal{H} .$$

Taking into account that the inner product on the Dirac spinors at x has signature $(2, 2)$, the local correlation operator $F^\varepsilon(x)$ is a symmetric operator on \mathcal{H} of rank

at most four, which has at most two positive and at most two negative eigenvalues. We thus obtain a mapping

$$F^\varepsilon : \mathcal{M} \rightarrow \mathcal{F},$$

where $\mathcal{F} \subset \mathcal{L}(\mathcal{H})$ is the set of all symmetric operators on \mathcal{H} of finite rank, which (counting multiplicities) have at most two positive and at most two negative eigenvalues. Finally, we introduce the measure ρ^ε on \mathcal{F} as the push-forward of the volume measure on \mathcal{M}

$$\rho^\varepsilon := F_*^\varepsilon \mu.$$

We thus obtain a causal fermion system of spin dimension two.

For the purposes of this paper, it suffices to consider the simplest regularization, the so-called $i\varepsilon$ -regularization (for details see for example [13, §2.4.1]). Its effect is seen most easily in the kernel of the fermionic projector (2.10), which can be computed to be the integral over the lower mass shell with a convergence-generating factor $e^{\varepsilon k^0}$, i.e.

$$P^\varepsilon(x, y) = \int \frac{d^4 k}{(2\pi)^4} (\not{k} + m) \delta(k^2 - m^2) \Theta(-k^0) e^{\varepsilon k^0} e^{-ik(x-y)}.$$

For any $\varepsilon > 0$, the kernel $P^\varepsilon(x, y)$ is a smooth function. However, in the limit $\varepsilon \rightarrow 0$, this kernel converges to a distribution which has singularities on the light cone (i.e. if x and y have lightlike separation). In a naive computation, these singularities give rise to divergent contributions to the causal action. The continuum limit analysis gives a systematic method for studying the causal action principle asymptotically for small ε . In this paper, we shall not need the details. Instead, we can make do with the scaling behavior in ε as captured by the following simple formalism, which was introduced in [15]. The singularity of a composite expression in the kernel of the fermionic projector (like the Lagrangian or variational derivatives thereof, the spectral weight $|xy|$, etc.) is described by the *degree on the light cone*. The scaling behavior of a term of degree L on the light cone is given by

$$\sim \frac{1}{(\varepsilon |\vec{\xi}|)^L} \quad \text{if } |\vec{\xi}| \gg \varepsilon \text{ and } ||t| - |\vec{\xi}|| \lesssim \varepsilon, \quad (2.13)$$

where $\xi := y - x$ with components $\xi = (t, \vec{\xi})$ (thus t is the time distance between x and y). When integrating over spacetime, the t -integration can be carried out across the light cone, compensating one factor of ε . Thus the term (2.13) can be written as a distribution supported on the light cone of the form

$$\simeq \frac{1}{\varepsilon^{L-1} |t|^L} \delta(|t| - |\vec{\xi}|) \simeq \frac{1}{(\varepsilon |t|)^{L-1}} \delta(\xi^2), \quad (2.14)$$

where \simeq means “up to a smooth prefactor” (here ξ^2 is again the Minkowski inner product). The general *formalism of the continuum limit* (see [10, Chapter 4] or [13, Section 2.4]) gives a systematic method of computing the smooth prefactors in (2.14) including precise error terms.

2.1.5. Connection to the Setting of Causal Variational Principles. For the analysis of the causal action principle it is most convenient to get into the simpler setting of *causal variational principles*. In this setting, \mathcal{F} is a (possibly non-compact) smooth manifold of dimension $m \geq 1$ and ρ a positive Borel measure on \mathcal{F} . Moreover, we are given a non-negative function $\mathcal{L} : \mathcal{F} \times \mathcal{F} \rightarrow \mathbb{R}_0^+$ (the *Lagrangian*) with the following properties:

- (i) \mathcal{L} is symmetric: $\mathcal{L}(x, y) = \mathcal{L}(y, x)$ for all $x, y \in \mathcal{F}$.

- (ii) \mathcal{L} is lower semi-continuous, i.e. for all sequences $x_n \rightarrow x$ and $y_{n'} \rightarrow y$,

$$\mathcal{L}(x, y) \leq \liminf_{n, n' \rightarrow \infty} \mathcal{L}(x_n, y_{n'}) .$$

The *causal variational principle* is to minimize the action

$$\mathcal{S}(\rho) = \int_{\mathcal{F}} d\rho(x) \int_{\mathcal{F}} d\rho(y) \mathcal{L}(x, y) \quad (2.15)$$

under variations of the measure ρ , keeping the total volume $\rho(\mathcal{F})$ fixed (*volume constraint*). If the total volume $\rho(\mathcal{F})$ is finite, one minimizes (2.15) over all regular Borel measures with the same total volume. If the total volume $\rho(\mathcal{F})$ is infinite, however, it is not obvious how to implement the volume constraint, making it necessary to proceed as follows. We need the following additional assumptions:

- (iii) The measure ρ is *locally finite* (meaning that any $x \in \mathcal{F}$ has an open neighborhood U with $\rho(U) < \infty$).
- (iv) The function $\mathcal{L}(x, \cdot)$ is ρ -integrable for all $x \in \mathcal{F}$, giving a lower semi-continuous and bounded function on \mathcal{F} .

Given a regular Borel measure ρ on \mathcal{F} , we then vary over all regular Borel measures $\tilde{\rho}$ with

$$|\tilde{\rho} - \rho|(\mathcal{F}) < \infty \quad \text{and} \quad (\tilde{\rho} - \rho)(\mathcal{F}) = 0$$

(where $|\cdot|$ denotes the total variation of a measure). These variations of the causal action are well-defined. The existence theory for minimizers is developed in [29].

There are several ways to get from the causal action principle to causal variational principles, as we now recall. If the Hilbert space \mathcal{H} is *finite-dimensional* and the total volume $\rho(\mathcal{F})$ is finite, one can proceed as follows: As a consequence of the trace constraint (2.4), for any minimizing measure ρ the local trace is constant in spacetime, i.e. there is a real constant $c \neq 0$ such that (see [4, Theorem 1.3] or [13, Proposition 1.4.1])

$$\text{tr } x = c \quad \text{for all } x \in M .$$

Restricting attention to operators with fixed trace, the trace constraint (2.4) is equivalent to the volume constraint (2.3) and may be disregarded. The boundedness constraint, on the other hand, can be treated with a Lagrange multiplier. More precisely, in [4, Theorem 1.3] it is shown that for every minimizing measure ρ , there is a Lagrange multiplier $\kappa > 0$ such that ρ is a critical point of the causal action with the Lagrangian replaced by

$$\mathcal{L}_\kappa(x, y) := \mathcal{L}(x, y) + \kappa |xy|^2 ,$$

leaving out the boundedness constraint. Having treated the constraints, the difference to causal variational principles is that in the setting of causal fermion systems, the set of operators $\mathcal{F} \subset \text{L}(\mathcal{H})$ does not necessarily have the structure of a manifold. In order to give this set a manifold structure, we need to assume that a given minimizing measure ρ (for the Lagrangian \mathcal{L}_κ) is *regular* in the sense that all operators in its support have exactly n positive and exactly n negative eigenvalues. This leads us to introduce the set \mathcal{F}^{reg} as the set of all operators F on \mathcal{H} with the following properties:

- (i) F is symmetric, has finite rank and (counting multiplicities) has exactly n positive and n negative eigenvalues.
- (ii) The trace is constant, i.e.

$$\text{tr}(F) = c > 0 .$$

The set \mathcal{F}^{reg} has a smooth manifold structure (see the concept of a flag manifold in [35] or the detailed construction in [22, Section 3]). In this way, the causal action principle becomes an example of a causal variational principle.

This finite-dimensional setting has the drawback that the total volume $\rho(\mathcal{F})$ of spacetime is finite, which is not suitable for describing asymptotically flat spacetimes or spacetimes of infinite lifetime like Minkowski space. Therefore, it is important to also consider the *infinite-dimensional setting* where $\dim \mathcal{H} = \infty$ and consequently also $\rho(\mathcal{F}) = \infty$ (see [13, Exercise 1.3]). In this case, the set \mathcal{F}^{reg} has the structure of an infinite-dimensional Banach manifold (for details see [30]). Here we shall not enter the subtleties of this infinite-dimensional analysis. Instead, we get by with the following simple method: Given a minimizing measure ρ , we choose \mathcal{F}^{reg} as a finite-dimensional manifold which contains $M := \text{supp } \rho$. We then restrict attention to variations of ρ in the class of regular Borel measures on \mathcal{F}^{reg} . In this way, we again get into the setting of causal variational principles. We refer to this method by saying that we *restrict attention to locally compact variations*. Keeping in mind that the dimension of \mathcal{F}^{reg} can be chosen arbitrarily large, this method seems like a sensible technical simplification.

For ease in notation, in what follows we will omit the superscript “reg.” Thus \mathcal{F} stands for a smooth (in general non-compact) manifold which contains the support M of the minimizing measure ρ .

2.1.6. The Euler-Lagrange Equations and Jet Spaces. A minimizer of a causal variational principle satisfies the following *Euler-Lagrange (EL) equations*. For a suitable value of the parameter $\mathfrak{s} > 0$, the lower semi-continuous function $\ell : \mathcal{F} \rightarrow \mathbb{R}_0^+$ defined by

$$\ell(x) := \int_M \mathcal{L}(x, y) d\rho(y) - \mathfrak{s}$$

is minimal and vanishes on spacetime $M := \text{supp } \rho$,

$$\ell|_M \equiv \inf_{\mathcal{F}} \ell = 0. \quad (2.16)$$

The parameter \mathfrak{s} can be understood as the Lagrange parameter corresponding to the volume constraint. For the derivation and further details we refer to [26, Section 2].

The EL equations (2.16) are nonlocal in the sense that they make a statement on the function ℓ even for points $x \in \mathcal{F}$ which are far away from the spacetime M . It turns out that for the applications we have in mind, it is preferable to evaluate the EL equations only locally in a neighborhood of M . This leads to the *restricted EL equations* introduced in [26, Section 4]. Here we give a slightly less general version of these equations which is sufficient for our purposes. In order to explain how the restricted EL equations come about, we begin with the simplified situation in which the function ℓ is smooth. In this case, the minimality of ℓ implies that the derivative of ℓ vanishes on M , i.e.

$$\ell|_M \equiv 0 \quad \text{and} \quad D\ell|_M \equiv 0 \quad (2.17)$$

(where $D\ell(p) : T_p \mathcal{F} \rightarrow \mathbb{R}$ is the derivative). In order to combine these two equations in a compact form, it is convenient to consider a pair $\mathbf{u} := (a, \mathbf{u})$ consisting of a real-valued function a on M and a vector field \mathbf{u} on $T\mathcal{F}$ along M , and to denote the combination of multiplication and directional derivative by

$$\nabla_{\mathbf{u}} \ell(x) := a(x) \ell(x) + (D_{\mathbf{u}} \ell)(x). \quad (2.18)$$

Then the equations (2.17) imply that $\nabla_{\mathbf{u}}\ell(x)$ vanishes for all $x \in M$. The pair $\mathbf{u} = (a, \mathbf{u})$ is referred to as a *jet*.

In the general lower-continuous setting, one must be careful because the directional derivative $D_{\mathbf{u}}\ell$ in (2.18) need not exist. Our method for dealing with this problem is to restrict attention to vector fields for which the directional derivative is well-defined. Moreover, we must specify the regularity assumptions on a and u . To begin with, we always assume that a and \mathbf{u} are *smooth* in the sense that they have a smooth extension to the manifold \mathcal{F} (for more details see [19, Section 2.2]). Thus the jet \mathbf{u} should be an element of the jet space

$$\mathfrak{J}_\rho := \{ \mathbf{u} = (a, \mathbf{u}) \text{ with } a \in C^\infty(M, \mathbb{R}) \text{ and } \mathbf{u} \in \Gamma(M, T\mathcal{F}) \} ,$$

where $C^\infty(M, \mathbb{R})$ and $\Gamma(M, T\mathcal{F})$ denote the space of real-valued functions and vector fields on M , respectively, which admit a smooth extension to \mathcal{F} .

Clearly, the fact that a jet \mathbf{u} is smooth does not imply that the functions ℓ or \mathcal{L} are differentiable in the direction of \mathbf{u} . This must be ensured by additional conditions which are satisfied by suitable subspaces of \mathfrak{J}_ρ which we now introduce. First, we let $\Gamma_\rho^{\text{diff}}$ be those vector fields for which the directional derivative of the function ℓ exists,

$$\Gamma_\rho^{\text{diff}} = \{ \mathbf{u} \in C^\infty(M, T\mathcal{F}) \mid D_{\mathbf{u}}\ell(x) \text{ exists for all } x \in M \} .$$

This gives rise to the jet space

$$\mathfrak{J}_\rho^{\text{diff}} := C^\infty(M, \mathbb{R}) \oplus \Gamma_\rho^{\text{diff}} \subset \mathfrak{J}_\rho .$$

For the jets in $\mathfrak{J}_\rho^{\text{diff}}$, the combination of multiplication and directional derivative in (2.18) is well-defined. We choose a linear subspace $\mathfrak{J}_\rho^{\text{test}} \subset \mathfrak{J}_\rho^{\text{diff}}$ with the property that its scalar and vector components are both vector spaces,

$$\mathfrak{J}_\rho^{\text{test}} = C^{\text{test}}(M, \mathbb{R}) \oplus \Gamma_\rho^{\text{test}} \subseteq \mathfrak{J}_\rho^{\text{diff}} ,$$

and the scalar component is nowhere trivial in the sense that

$$\text{for all } x \in M \text{ there is } a \in C^{\text{test}}(M, \mathbb{R}) \text{ with } a(x) \neq 0 . \quad (2.19)$$

Moreover, compactly supported jets are always denoted by a subscript zero, like for example

$$\mathfrak{J}_{\rho,0}^{\text{test}} := \{ \mathbf{u} \in \mathfrak{J}_\rho^{\text{test}} \mid \mathbf{u} \text{ has compact support} \} .$$

Then the *restricted EL equations* read (for details cf. [26, (eq. (4.10))])

$$\nabla_{\mathbf{u}}\ell|_M = 0 \quad \text{for all } \mathbf{u} \in \mathfrak{J}_\rho^{\text{test}} . \quad (2.20)$$

Before going on, we point out that the restricted EL equations (2.20) do not hold only for minimizers, but also for critical points of the causal action. With this in mind, all the methods and results of this paper do not apply only to minimizers, but more generally to critical points of the causal variational principle. For brevity, we also refer to a measure which satisfies the restricted EL equations (2.20) as a *critical measure*.

2.1.7. Minkowski-Type Spacetimes. For technical simplicity, in this paper we restrict attention to spacetimes which are smooth and have the same topology as Minkowski space. This will make it possible to work with the usual coordinates (t, \vec{x}) and corresponding foliations by hypersurfaces $t = \text{const}$. Let $(\mathcal{H}, \mathcal{F}, \rho)$ be a causal fermion system, which may be thought of as describing either the vacuum or the interacting physical system. We assume that ρ is a critical point of the causal action principle.

Moreover, we assume that the corresponding spacetime $M := \text{supp } \rho$ is diffeomorphic to a four-dimensional spacetime with trivial topology, i.e.

$$M \simeq \mathcal{M} := \mathbb{R}^4.$$

Next, we assume that, using this identification, the measure ρ is absolutely continuous with respect to the Lebesgue measure with a smooth weight function, i.e.

$$d\rho = h(x) d^4x \quad \text{with} \quad h \in C^\infty(\mathcal{M}, \mathbb{R}^+). \quad (2.21)$$

We also assume that h is bounded from above and below, i.e. there should be a constant $C > 1$ with

$$\frac{1}{C} \leq h(x) \leq C \quad \text{for all } x \in \mathcal{M}.$$

We also denote the coordinate x^0 as time function T ,

$$T : M \rightarrow \mathbb{R}, \quad (t, \mathbf{x}) \mapsto t. \quad (2.22)$$

For any $t \in \mathbb{R}$, we let Ω^t be the past of t ,

$$\Omega^t := \{x \in \mathcal{M} \mid T(x) \leq t\}. \quad (2.23)$$

2.1.8. Surface Layer Integrals for Jets. Surface layer integrals were first introduced in [25] as double integrals of the general form

$$\int_{\Omega} \left(\int_{M \setminus \Omega} (\cdots) \mathcal{L}_{\kappa}(x, y) d\rho(y) \right) d\rho(x), \quad (2.24)$$

where (\cdots) stands for a suitable differential operator formed of jets. A surface layer integral generalizes the concept of a surface integral over $\partial\Omega$ to the setting of causal fermion systems. The connection can be understood most easily in the case when $\mathcal{L}_{\kappa}(x, y)$ vanishes unless x and y are close together. In this case, we only get a contribution to (2.24) if both x and y are close to the boundary of Ω . A more detailed explanation of the idea of a surface layer integral is given in [25, Section 2.3].

In the present paper, we always choose the set Ω according to (2.23) as the past of a time t . We now recall those surface layer integrals for jets which will be of relevance in this paper.

Definition 2.3. *We define the following surface layer integrals,*

$$\begin{aligned} \gamma_{\rho}^t : \mathfrak{J}_{\rho, \text{sc}} &\rightarrow \mathbb{R} && (\text{conserved one-form}) \\ \gamma_{\rho}^t(\mathbf{v}) &= \int_{\Omega} d\rho(x) \int_{M \setminus \Omega} d\rho(y) (\nabla_{1, \mathbf{v}} - \nabla_{2, \mathbf{v}}) \mathcal{L}(x, y) \end{aligned} \quad (2.25)$$

$$\begin{aligned} \sigma_{\rho}^t : \mathfrak{J}_{\rho, \text{sc}} \times \mathfrak{J}_{\rho, \text{sc}} &\rightarrow \mathbb{R} && (\text{symplectic form}) \\ \sigma_{\rho}^t(\mathbf{u}, \mathbf{v}) &= \int_{\Omega} d\rho(x) \int_{M \setminus \Omega} d\rho(y) (\nabla_{1, \mathbf{u}} \nabla_{2, \mathbf{v}} - \nabla_{2, \mathbf{u}} \nabla_{1, \mathbf{v}}) \mathcal{L}(x, y) \end{aligned} \quad (2.26)$$

$$\begin{aligned} (\cdot, \cdot)_{\rho}^t : \mathfrak{J}_{\rho, \text{sc}} \times \mathfrak{J}_{\rho, \text{sc}} &\rightarrow \mathbb{R} && (\text{surface layer inner product}) \\ (\mathbf{u}, \mathbf{v})_{\rho}^t &= \int_{\Omega} d\rho(x) \int_{M \setminus \Omega} d\rho(y) (\nabla_{1, \mathbf{u}} \nabla_{1, \mathbf{v}} - \nabla_{2, \mathbf{u}} \nabla_{2, \mathbf{v}}) \mathcal{L}(x, y). \end{aligned} \quad (2.27)$$

Here $\mathfrak{J}_{\rho, \text{sc}}$ denotes the jets in $\mathfrak{J}_{\rho}^{\text{vary}}$ with spatially compact support (for details see [8, Section 5.3]), where $\mathfrak{J}_{\rho}^{\text{vary}}$ is a suitably chosen subspace of $\mathfrak{J}_{\rho}^{\text{test}}$ (for details see [8, Section 3.2]).

2.1.9. The Linearized Field Equations. In simple terms, the linearized field equations describe variations of the measure ρ which preserve the EL equations. More precisely, we consider variations where we multiply ρ by a non-negative function and take the push-forward with respect to a mapping from M to \mathcal{F} . Thus we consider families of measures $(\tilde{\rho}_\tau)_{\tau \in (-\delta, \delta)}$ of the form

$$\tilde{\rho}_\tau = (F_\tau)_*(f_\tau \rho),$$

where the f_τ and F_τ are smooth,

$$f_\tau \in C^\infty(M, \mathbb{R}^+) \quad \text{and} \quad F_\tau \in C^\infty(M, \mathcal{F}),$$

depend smoothly on the parameter τ and have the properties $f_0(x) = 1$ and $F_0(x) = x$ for all $x \in M$ (moreover, the star denotes the push-forward measure, which is defined for a subset $\Omega \subset \mathcal{F}$ by $((F_\tau)_*\mu)(\Omega) = \mu(F_\tau^{-1}(\Omega))$; see for example [5, Section 3.6]). We assume that the measures $(\tilde{\rho}_\tau)_{\tau \in (-\delta, \delta)}$ satisfy the EL equations (2.16) for all τ . Then the infinitesimal generator of the variation denoted by

$$\mathfrak{v}(x) := \left. \frac{d}{d\tau} (f_\tau(x), F_\tau(x)) \right|_{\tau=0},$$

satisfies the *linearized field equations*

$$0 = \langle \mathfrak{u}, \Delta \mathfrak{v} \rangle(x) := \nabla_{\mathfrak{u}} \left(\int_M (\nabla_{1, \mathfrak{v}} + \nabla_{2, \mathfrak{v}}) \mathcal{L}_\kappa(x, y) d\rho(y) - \nabla_{\mathfrak{v}} \mathfrak{s} \right),$$

which hold for all $\mathfrak{u} \in \mathfrak{J}_\rho^{\text{test}}$ and all $x \in M$ (for details see [16, Section 3.3]). We denote the vector space of all solutions of the linearized field equations by $\mathfrak{J}_\rho^{\text{lin}}$.

The linearized field equations harmonize with the structure of surface layer integrals in the sense that linearized solutions give rise to the following *conservation laws* or almost conserved surface layer integrals (for details see [25, 26, 27] or [23, Chapter 9]):

- (a) The conserved one-form is time independent if \mathfrak{v} is a linearized solution with vanishing scalar component.
- (b) The symplectic form is independent of t for all linearized solutions \mathfrak{u} and \mathfrak{v} .
- (c) The surface layer inner product is conserved up to quadratic corrections to the linearized field equations.

We remark that the quadratic corrections will not be of relevance in this paper because we shall consider the linearized fields only in the non-interacting vacuum spacetime.

2.1.10. Inner Solutions, Arranging Jets without Scalar Components. We now briefly recall the definition of inner solutions as introduced in [19, Section 3].

Definition 2.4. An inner solution is a jet \mathfrak{v} of the form

$$\mathfrak{v} = (\text{div } \mathbf{v}, \mathbf{v}) \quad \text{with} \quad \mathbf{v} \in \Gamma(M, TM),$$

where the divergence is taken with respect to the measure in (2.21),

$$\text{div } \mathbf{v} := \frac{1}{h} \partial_j (h \mathbf{v}^j).$$

Under suitable regularity and decay assumptions, an inner solution solves the linearized field equations (for details see [19, Section 3.1]). We denote these inner solutions by $\mathfrak{J}_\rho^{\text{in}}$. In [19, Proposition 3.6] it is shown that inner solutions can be used for testing. With this in mind, we always assume that

$$\mathfrak{J}_\rho^{\text{in}} \subset \mathfrak{J}_\rho^{\text{test}}.$$

Inner solutions can be regarded as infinitesimal generators of transformations of M which leave the measure ρ unchanged. Therefore, inner solutions do not change the causal fermion system, but merely describe symmetry transformations of the measure. With this in mind, we can modify solutions of the linearized field equations by adding inner solutions. For conceptual clarity, it is preferable that the diffeomorphism generated by an inner solution does not change the global time function T in (2.22). Therefore, we consider vector fields which are tangential to the hypersurfaces $N_t := T^{-1}(t)$. As shown in [31, Lemma 2.7], the divergence of such a vector field can be arranged to be any given function $a \in C^\infty(M, \mathbb{R})$. Thus, by adding the corresponding inner solutions we can achieve that all linearized solutions have no scalar components. Therefore, in what follows we may restrict attention to linearized solutions $\mathbf{v} \in \mathfrak{J}^{\text{lin}}$ with vanishing scalar component. We also write these jets as

$$\mathbf{v} = (0, \mathbf{v}) \quad \text{with} \quad \mathbf{v} \in \Gamma_\rho^{\text{lin}}.$$

2.1.11. The Dynamical Wave Equation and the Extended Hilbert Space. The restricted EL equations (2.20) can be expressed in terms of the physical wave equations. This gives rise to the *dynamical wave equation* as introduced in [21]. The solutions of this wave equation form a Hilbert space, the *extended Hilbert space*. We now recall a few concepts and results from [21]. Our starting point is the formula (2.9) which expresses the spacetime point operator as a local correlation operator. Varying the wave evaluation operator gives a vector field \mathbf{u} on \mathcal{F} along M ,

$$\mathbf{u}(x) = -\delta\Psi(x)^* \Psi(x) - \Psi(x)^* \delta\Psi(x). \quad (2.28)$$

In order to make mathematical sense of this formula in agreement with the concept of restricting attention to locally compact variations, we choose a finite-dimensional subspace $\mathcal{H}^f \subset \mathcal{H}$, i.e.

$$f^f := \dim \mathcal{H}^f < \infty \quad (2.29)$$

and impose the following assumptions on $\delta\Psi$ (similar variations were first considered in [16, Section 7]):

- (a) The variation is trivial on the orthogonal complement of \mathcal{H}^f ,

$$\delta\Psi|_{(\mathcal{H}^f)^\perp} = 0.$$

- (b) The variations of all physical wave functions are continuous and compactly supported, i.e.

$$\delta\Psi : \mathcal{H} \rightarrow C_0^0(M, SM).$$

Before going on, we point out that the choice of \mathcal{H}^f is *not canonical*. Ultimately, one would like to exhaust \mathcal{H} by a sequence of finite-dimensional subspaces $\mathcal{H}_1^f \subset \mathcal{H}_2^f \subset \dots$ and take the limit. Here we shall not enter this analysis, but for technical simplicity we rather choose \mathcal{H}^f as a finite-dimensional subspace of sufficiently large dimension.

We choose $\Gamma_{\rho,0}^f$ as a space of vector fields of the form (2.28). For convenience, we identify the vector field with the first variation $\delta\Psi$ and write $\delta\Psi \in \Gamma_{\rho,0}^f$ (this representation of \mathbf{u} in terms of $\delta\Psi$ may not be unique, but this is of no relevance for what follows). Choosing trivial scalar components, we obtain a corresponding space of jets $\mathfrak{J}_{\rho,0}^f$, referred to as the *fermionic jets*. We always assume that the fermionic jets are admissible for testing, i.e.

$$\mathfrak{J}_{\rho,0}^f := \{0\} \oplus \Gamma_{\rho,0}^f \subset \mathfrak{J}_{\rho,0}^{\text{test}}.$$

Moreover, in analogy to the condition (2.19) for the scalar components of the test jets, we assume that the variation can have arbitrary values at any spacetime point, i.e.

for all $x \in M$, $\chi \in S_x$ and $\phi \in \mathcal{H}^f$ there is $\delta\Psi \in \Gamma_{\rho,0}^f$ with $\delta\Psi(x)\phi = \chi$.

Evaluating first variations of the causal action in direction of fermionic jets, one obtains the *EL equation for the physical wave functions*

$$\int_M Q(x, y) \Psi^f(y) d\rho(y) = \mathfrak{r} \Psi^f(x) \quad \text{for all } x \in M, \quad (2.30)$$

where $\mathfrak{r} \in \mathbb{R}$ is the Lagrange parameter of the trace constraint, $Q(x, y)$ is the kernel (2.12) (with \mathcal{L} replaced by the κ -Lagrangian), and $\Psi^f := \Psi|_{\mathcal{H}^f}$ denotes the restriction of the wave evaluation operator to the finite-dimensional subspace \mathcal{H}^f . More details on this formulation of the EL equations and many computations can be found in [13, Sections 1.4 and 2.6 as well as Chapters 3-5]. One way of looking at these equations is to regard (2.30) for given $Q(x, y)$ as a linear equation describing the dynamics of the physical wave functions. In [21], this wave equation was extended to more general solutions which form the so-called *extended Hilbert space* $(\mathcal{H}_\rho^{f,t}, \langle \cdot | \cdot \rangle_\rho^t)$. The dynamics in $\mathcal{H}_\rho^{f,t}$ is described by the so-called *dynamical wave equation*

$$Q^{\text{dyn}}\psi = 0,$$

where Q^{dyn} is an integral operator with a symmetric integral kernel

$$Q^{\text{dyn}}(x, y) : S_y \rightarrow S_x.$$

The scalar product $\langle \cdot | \cdot \rangle_\rho^t$ at time t has the form

$$\begin{aligned} \langle \psi | \phi \rangle_\rho^t &= -2i \left(\int_{\Omega^t} d\rho(x) \int_{M \setminus \Omega^t} d\rho(y) - \int_{M \setminus \Omega^t} d\rho(x) \int_{\Omega^t} d\rho(y) \right) \\ &\quad \times \prec \psi(x) | Q^{\text{dyn}}(x, y) \phi(y) \succ_x. \end{aligned}$$

This scalar product is conserved, i.e. time independent. With this in mind, we can drop to upper index t and denote the extended Hilbert space simply by $(\mathcal{H}_\rho^f, \langle \cdot | \cdot \rangle_\rho)$.

2.1.12. A Conserved Nonlinear Surface Layer Integral. In [19, Section 4] a nonlinear surface layer integral $\gamma^t(\rho, \tilde{\rho})$ was introduced which can be used for comparing the measure $\tilde{\rho}$ describing the interacting system with the vacuum measure ρ . As the starting point, we let ρ and $\tilde{\rho}$ be two critical measures on \mathcal{F} , which describe the vacuum and the interacting system, respectively. In order to relate the interacting spacetime $\tilde{M} := \text{supp } \tilde{\rho}$ with the vacuum spacetime, we choose a mapping

$$\Phi : \tilde{M} \rightarrow M,$$

which we assume to be measurable (in the sense that $\Phi^{-1}(U)$ is $\tilde{\rho}$ -measurable for every ρ -measurable set $U \subset M$). Choosing a foliation $(N_t)_{t \in \mathbb{R}}$ of M , the past sets $\Omega^t \subset M$ give rise to corresponding past sets $\tilde{\Omega}^t := \Phi^{-1}(\Omega^t) \subset \tilde{M}$. Then the nonlinear surface layer integral at time t is defined by (see [19, Definition 4.1] and Figure 2)

$$\gamma^t(\tilde{\rho}, \rho) = \int_{\tilde{\Omega}^t} d\tilde{\rho}(x) \int_{M \setminus \Omega^t} d\rho(y) \mathcal{L}(x, y) - \int_{\tilde{M} \setminus \tilde{\Omega}^t} d\tilde{\rho}(x) \int_{\Omega^t} d\rho(y) \mathcal{L}(x, y). \quad (2.31)$$

In order to ensure that the nonlinear surface layer integral is well-defined and finite, we need to make the following assumption.




FIGURE 2. The nonlinear surface layer integral.

Definition 2.5. *The measures ρ and $\tilde{\rho}$ are **surface layer admissible** if for all $t \in \mathbb{R}$, both integrals in (2.31) are finite,*

$$\int_{\tilde{\Omega}^t} d\tilde{\rho}(x) \int_{M \setminus \Omega^t} d\rho(y) \mathcal{L}(x, y) < \infty, \quad \int_{\tilde{M} \setminus \tilde{\Omega}^t} d\tilde{\rho}(x) \int_{\Omega^t} d\rho(y) \mathcal{L}(x, y) < \infty.$$

By a suitable choice of the mapping $\Phi : \tilde{M} \rightarrow M$ one can arrange that the nonlinear surface layer integral is conserved. In order to construct Φ , we introduce a measure ν on M and a measure $\tilde{\nu}$ on $\tilde{M} := F(M)$ (the so-called *correlation measures*) by

$$d\nu(x) := \left(\int_{\tilde{M}} \mathcal{L}(x, y) d\tilde{\rho}(y) \right) d\rho(x) \quad \text{and} \quad d\tilde{\nu}(x) := \left(\int_M \mathcal{L}(x, y) d\rho(y) \right) d\tilde{\rho}(x).$$

As shown in [19, Appendix A], the conservation law is related to properties of the correlation measures:

Proposition 2.6. *The surface layer integral (2.31) vanishes for every compact $\Omega \subset M$ if and only if $\nu = \Phi_* \tilde{\nu}$.*

As worked out in detail again in [19, Appendix A], the existence of such a diffeomorphism Φ follows from a general result in [34].

2.2. Notions of Quantum States of Causal Fermion Systems. We now recall the definition of the quantum state and various refinements of the construction as given in [20]. Our presentation is very brief and is limited to listing all the objects and definitions. We will come back to these notions in more detail in Section 3, where we will discuss them with regard to entanglement and explain how the quantum states can be computed.

2.2.1. The Field Algebra of the Vacuum Spacetime. We let $(\mathcal{H}, \mathcal{F}, \rho)$ be the causal fermion system describing the vacuum. We now introduce the algebra of observables, denoted by \mathcal{A} . It is the $*$ -algebra generated by the bosonic and fermionic field operators defined as follows. The *fermionic creation and annihilation operators* are denoted by

$$\Psi^\dagger(\phi) \quad \text{and} \quad \Psi(\bar{\phi}) \quad \text{for} \quad \phi \in \mathcal{H}_{\rho, \text{sc}}^{\text{f}}.$$

They satisfy the canonical anti-commutation relations

$$\{\Psi(\bar{\phi}), \Psi^\dagger(\phi')\} = \langle \phi | \phi' \rangle_\rho^t,$$

and all other operators anti-commute,

$$\{\Psi(\bar{\phi}), \Psi(\bar{\phi}')\} = 0 = \{\Psi^\dagger(\phi), \Psi^\dagger(\phi')\}.$$

The definition of the *bosonic field operators* involves the choice of a *complex structure*. We always work with the canonical complex structure J is induced on $\Gamma_{\rho, \text{sc}}^{\text{lin}}$ by the

surface layer integrals $(\cdot, \cdot)_\rho^t$ and σ_ρ^t (see (2.27) and (2.26); in the considered vacuum spacetime these surface layer integrals are both time independent). The construction, which is carried out in detail in [19, Section 6.3], is summarized as follows. We assume that the surface layer integral $(\cdot, \cdot)_\rho^t$ restricted to $\Gamma_{\rho, \text{sc}}^{\text{lin}} \times \Gamma_{\rho, \text{sc}}^{\text{lin}}$ is positive semi-definite. Dividing out the null space and forming the completion, we obtain a real Hilbert space denoted by $(\mathfrak{h}^{\mathbb{R}}, (\cdot, \cdot)_\rho^t)$. For the construction of J , one assumes that σ_ρ^t is bounded relative to the scalar product $(\cdot, \cdot)_\rho^t$. Then we can represent σ_ρ^t as

$$\sigma_\rho^t(u, v) = (u, \mathcal{T}v)_\rho^t,$$

where \mathcal{T} is a uniquely determined bounded operator on the Hilbert space $\mathfrak{h}^{\mathbb{R}}$. Assuming that \mathcal{T} is invertible, we set

$$J := -(-\mathcal{T}^2)^{-\frac{1}{2}} \mathcal{T}.$$

Next, we complexify the Hilbert space $\mathfrak{h}^{\mathbb{R}}$ and denote its complexification by $\mathfrak{h}^{\mathbb{C}}$. On this complexification, the operator J has eigenvalues i and $-i$. The corresponding eigenspaces are referred to as the *holomorphic* and *anti-holomorphic subspaces*, respectively. We write the decomposition into holomorphic and anti-holomorphic components as

$$\mathbf{v} = \mathbf{v}^{\text{hol}} + \mathbf{v}^{\text{ah}}.$$

We also complexify the symplectic form to a *sesquilinear* form on $\mathfrak{h}^{\mathbb{C}}$ (i.e. anti-linear in its first and linear in its second argument). On the holomorphic jets we introduce a scalar product $(\cdot, \cdot)_\rho^t$ by

$$(\cdot, \cdot)_\rho^t := \sigma_\rho^t(\cdot, J \cdot) : \Gamma_\rho^{\text{hol}} \times \Gamma_\rho^{\text{hol}} \rightarrow \mathbb{C}.$$

Taking the completion gives a Hilbert space, which we denote by $(\mathfrak{h}, (\cdot, \cdot)_\rho^t)$. This scalar product has the useful property that

$$\text{Im}(u|v)_\rho^t = \text{Im} \sigma_\rho^t(u, Jv) = \text{Re} \sigma_\rho^t(u, v).$$

The *bosonic creation and annihilation operators* are denoted by

$$a^\dagger(z) \quad \text{and} \quad a(\overline{z}) \quad \text{with} \quad z \in \mathfrak{h}$$

(the overline in $a(\overline{z})$ serves as a reminder that this operator is anti-linear). They satisfy the canonical commutation relations

$$[a(\overline{z}), a^\dagger(z')] = (z|z')_\rho^t, \tag{2.32}$$

and all other operators commute,

$$[a(\overline{z}), a(\overline{z'})] = 0 = [a^\dagger(z), a^\dagger(z')].$$

We define \mathcal{A} as the unital $*$ -algebra generated by the above field operators.

2.2.2. The Partition Function. Our starting point consists of two causal fermion systems $(\mathcal{H}, \mathcal{F}, \rho)$ (describing the vacuum) and $(\tilde{\mathcal{H}}, \tilde{\mathcal{F}}, \tilde{\rho})$ (describing the interacting system). We assume that both measures ρ and $\tilde{\rho}$ are critical points of the causal action. Moreover, we assume that both spacetimes M and \tilde{M} are of Minkowski type (see Section 2.1.7) with given time functions T and \tilde{T} .

The idea for constructing a quantum state is “compare” these two systems at a given fixed time and to try to describe the interacting system in terms of linearized fields and wave functions in the vacuum spacetime. This “comparison” can also be understood as a “measurement” performed in the interacting spacetime using objects from the vacuum spacetime as “measurement devices.” In more technical terms, we want to work

with the nonlinear surface layer integral (see Section 2.1.12) and variations thereof. When working with causal fermion systems, however, there is the complication that the two causal fermion systems are defined on two different Hilbert spaces \mathcal{H} and $\tilde{\mathcal{H}}$. Therefore, in order to make sense of the nonlinear surface layer integral, we need to identify the Hilbert spaces \mathcal{H} and $\tilde{\mathcal{H}}$ by a unitary transformation denoted by V ,

$$V : \mathcal{H} \rightarrow \tilde{\mathcal{H}} \quad \text{unitary} .$$

Then operators in $\tilde{\mathcal{F}}$ can be identified with operators in \mathcal{F} by the unitary transformation,

$$\mathcal{F} = V^{-1} \tilde{\mathcal{F}} V .$$

An important point to keep in mind is that this identification is not canonical, but that it leaves the freedom to transform the operator V according to

$$V \rightarrow V\mathcal{U} \quad \text{with} \quad \mathcal{U} \in \text{L}(\mathcal{H}) \text{ unitary} .$$

For ease of notation, in what follows we always identify \mathcal{H} and $\tilde{\mathcal{H}}$ via V , making it possible to always work in the Hilbert space \mathcal{H} . Then the non-uniqueness of the identification still shows up in the unitary transformation of the vacuum measure

$$\rho \mapsto \mathcal{U}\rho , \quad (2.33)$$

where $\mathcal{U}\rho$ is defined by

$$(\mathcal{U}\rho)(\Omega) := \rho(\mathcal{U}^{-1} \Omega \mathcal{U}) \quad \text{for} \quad \Omega \subset \mathcal{F} . \quad (2.34)$$

The method for treating the unitary freedom (2.33) is to integrate over the transformation \mathcal{U} . This leads to the definition of the partition function, obtained by integrating the exponential of the nonlinear surface layer integral.

Definition 2.7. *The partition function is defined by*

$$Z^t(\beta, \tilde{\rho}) = \int_{\mathcal{G}} \exp \left(\beta \gamma^t(\tilde{\rho}, \mathcal{U}\rho) \right) d\mu_{\mathcal{G}}(\mathcal{U}) , \quad (2.35)$$

where \mathcal{G} is chosen as a compact Lie subgroup of the unitary group,

$$\mathcal{G} \subset \text{U}(\mathcal{H}) \quad \text{compact} .$$

Here $\gamma^t(\tilde{\rho}, \mathcal{U}\rho)$ is the nonlinear surface layer integral (2.31) for the unitarily transformed measure (2.34), i.e.

$$\begin{aligned} \gamma^t(\tilde{\rho}, \mathcal{U}\rho) &= \int_{\tilde{\Omega}^t} d\tilde{\rho}(x) \int_{M \setminus \Omega^t} d\rho(y) \mathcal{L}(x, \mathcal{U}y\mathcal{U}^{-1}) \\ &\quad - \int_{\tilde{M} \setminus \tilde{\Omega}^t} d\tilde{\rho}(x) \int_{\Omega^t} d\rho(x) \mathcal{L}(x, \mathcal{U}y\mathcal{U}^{-1}) . \end{aligned} \quad (2.36)$$

It is one of the main goals of the present paper to compute integrals over the unitary group as in (2.35).

We finally comment on the choice of the group \mathcal{G} . The most natural choice is the whole unitary group $\mathcal{G} = \text{U}(\mathcal{H})$. But clearly, this choice is possible only if the Hilbert space \mathcal{H} is finite-dimensional. For this reason, in [20] we chose $\mathcal{G} = \text{U}(\mathcal{H}^f)$, where \mathcal{H}^f is the finite-dimensional subspace of \mathcal{H} already introduced in the context of the dynamical wave equation (2.29). In this paper, it will be preferable to choose \mathcal{H} as a finite-dimensional Hilbert space and $\mathcal{G} = \text{U}(\mathcal{H})$ as the whole unitary group; this will be discussed and explained in detail in Section 3.4.

2.2.3. The Quantum State of a Causal Fermion System. The general idea behind the construction of the quantum state is to probe the interacting spacetime with the objects of the non-interacting spacetime. More specifically, it is a linear mapping from the algebra \mathcal{A} (the field algebra of the vacuum spacetime; see Section 2.2.1) to the complex numbers which is positive, i.e.

$$\omega : \mathcal{A} \rightarrow \mathbb{C} \quad \text{with} \quad \omega(A^*A) \geq 0 \quad \text{for all } A \in \mathcal{A}.$$

It has the general structure

$$\omega^t(A) = \frac{1}{Z^t(\beta, \tilde{\rho})} \int_{\mathcal{G}} (\dots) e^{\beta \gamma^t(\tilde{\rho}, \mathcal{U}\rho)} d\mu_{\mathcal{G}}(\mathcal{U}), \quad (2.37)$$

where (\dots) stands for insertions formed of surface layer integrals which depend on A . In more detail,

$$\begin{aligned} & \omega^t \left(a^\dagger(z'_1) \cdots a^\dagger(z'_p) \Psi^\dagger(\phi'_1) \cdots \Psi^\dagger(\phi'_{r'}) a(\bar{z}_1) \cdots a(\bar{z}_q) \Psi(\bar{\phi}_1) \cdots \Psi(\bar{\phi}_r) \right) \\ &:= \frac{1}{Z^t(\beta, \tilde{\rho})} \delta_{r'r} \frac{1}{r!} \sum_{\sigma, \sigma' \in S_r} (-1)^{\text{sign}(\sigma) + \text{sign}(\sigma')} \\ & \quad \times \int_{\mathcal{G}} e^{\beta \gamma^t(\tilde{\rho}, \mathcal{U}\rho)} \langle \phi_{\sigma(1)} | \pi^{f,t} \phi'_{\sigma'(1)} \rangle_{\rho}^t \cdots \langle \phi_{\sigma(r)} | \pi^{f,t} \phi'_{\sigma'(r)} \rangle_{\rho}^t \\ & \quad \times D_{z'_1} \gamma^t(\tilde{\rho}, \mathcal{U}\rho) \cdots D_{z'_p} \gamma^t(\tilde{\rho}, \mathcal{U}\rho) D_{\bar{z}_1} \gamma^t(\tilde{\rho}, \mathcal{U}\rho) \cdots D_{\bar{z}_q} \gamma^t(\tilde{\rho}, \mathcal{U}\rho) d\mu_{\mathcal{G}}(\mathcal{U}), \end{aligned} \quad (2.38)$$

where the bosonic insertions are derivatives of the nonlinear surface layer integral, and $\pi^{f,t}$ is a projection operator which, in non-technical terms, tells us how the physical wave functions of the interacting spacetime look like in the non-interacting spacetime (for details see [20, Section 4.3] and the discussion in Remark 6.6).

2.2.4. The Refined Pre-State. The *refined pre-state* was introduced in [20, Section 5.3] with the intention of getting more detailed phase information on the interacting measure. We now briefly recall the definition. In Section 3.2, we will come back to the refined pre-state and explain why it is needed in order to account for the physical effect of entanglement. The basic idea is to work, instead of \mathcal{U} , with two unitary transformations $\mathcal{U}_<$ and $\mathcal{U}_>$, and to replace the unitarily transformed measure $\mathcal{U}\rho$ in (2.34) by the measure $T_{\mathcal{U}_<, \mathcal{U}_>} \rho$ defined by

$$(T_{\mathcal{U}_<, \mathcal{U}_>} \rho)(\Omega) := \rho(\mathcal{U}_>^{-1} \Omega \mathcal{U}_<)$$

(note that this measure may be supported on non-symmetric operators). For the refined partition function Z_{ref}^t and the refined pre-state ω_{ref}^t one modifies (2.35) and (2.37) by working with a double integral over the unitary group \mathcal{G} , i.e. symbolically

$$Z_{\text{ref}}^t(\beta, \tilde{\rho}) := \int_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_<) \int_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_>) e^{\beta \gamma^t(\tilde{\rho}, T_{\mathcal{U}_<, \mathcal{U}_>} \rho)} \quad (2.39)$$

$$\omega_{\text{ref}}^t(\dots) := \frac{1}{Z^t(\beta, \tilde{\rho})} \int_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_<) \int_{\mathcal{G}} d\mu_{\mathcal{G}}(\dots) e^{\beta \gamma^t(\tilde{\rho}, T_{\mathcal{U}_<, \mathcal{U}_>} \rho)}. \quad (2.40)$$

Similar to (2.38), the insertions are formed of suitable surface layer integral. We postpone the details to the thorough construction and discussion in Section 6.6.

We finally remark that by a *pre-state* we mean a linear functional from \mathcal{A} to \mathbb{C} , which does not necessarily need to be positive. Therefore, before applying the refined pre-state in physical applications, we need to show positivity (see Section 6.7).

2.2.5. The Localized Refined Pre-State. The quantum state defined in Section 2.2.3 was defined globally in space. For most applications, however, it seems more suitable to define a state describing only a *bounded spatial region*, which can be thought of as our laboratory or the subsystem of our present universe accessible to measurements. This picture will also be very helpful for the constructions in the present paper, because it will make it possible to take the limit when the dimension of the group \mathcal{G} tends to infinity, while leaving our laboratory unchanged (for more details see Section 3.3 below). In order to implement this picture, we choose a subset $V \subset M$ of the vacuum spacetime such that its intersection with the surface layer contains the spatial region of interest (see Figure 1). Correspondingly, the set $\tilde{V} := \Phi^{-1}(V) \subset \tilde{M}$ is the region of the interacting spacetime region which is probed by the laboratory. We define the *localized nonlinear surface layer integral* γ_V^t by restricting all the integrals in the nonlinear surface layer integral (2.36) to V respectively \tilde{V} ,

$$\gamma_V^t(\tilde{\rho}, \rho) = \int_{\tilde{\Omega}^t \cap \tilde{V}} d\tilde{\rho}(x) \int_{V \setminus \Omega^t} d\rho(y) \mathcal{L}(x, y) - \int_{\tilde{V} \setminus \tilde{\Omega}^t} d\tilde{\rho}(x) \int_{\Omega^t \cap V} d\rho(y) \mathcal{L}(x, y), \quad (2.41)$$

and similarly for the unitarily transformed measure as used in the refined state,

$$\begin{aligned} \gamma_V^t(\tilde{\rho}, T_{\mathcal{U}_<, \mathcal{U}_>} \rho) &= \int_{\tilde{\Omega}^t \cap \tilde{V}} d\tilde{\rho}(x) \int_{V \setminus \Omega^t} d\rho(y) \mathcal{L}(x, \mathcal{U}_> y \mathcal{U}_<^{-1}) \\ &\quad - \int_{\tilde{V} \setminus \tilde{\Omega}^t} d\tilde{\rho}(y) \int_{\Omega^t \cap V} d\rho(x) \mathcal{L}(x, \mathcal{U}_> y \mathcal{U}_<^{-1}). \end{aligned}$$

Next, we replace all surface layer integrals in (2.39) and (2.40) by the corresponding localized surface layer integrals. After doing so, however, one must take into account that the unitary group \mathcal{G} typically includes the unitary transformations describing the symmetries of Minkowski space like time translations and spatial translations. In order to remove the freedom of performing translations in Minkowski space, it is convenient to insert the factor $e^{\alpha \mathcal{T}_V^t}(\tilde{\rho}, \mathcal{U}\rho)$ into the integrand of the partition function and the state, where $\mathcal{T}_V^t(\tilde{\rho}, \mathcal{U}\rho)$ involves two spacetime integrals,

$$\begin{aligned} \mathcal{T}_V^t(\tilde{\rho}, T_{\mathcal{U}_<, \mathcal{U}_>} \rho) \\ := \left(\int_{\tilde{\Omega}^t \cap \tilde{V}} d\tilde{\rho}(x) \int_{\Omega^t \cap V} d\rho(y) + \int_{(\tilde{M} \setminus \tilde{\Omega}^t) \cap \tilde{V}} d\tilde{\rho}(x) \int_{(M \setminus \Omega^t) \cap V} d\rho(y) \right) |x \mathcal{U}_> y \mathcal{U}_<^{-1}|^2. \end{aligned} \quad (2.42)$$

(here the absolute value is again the spectral weight (2.6)). Thus the *refined localized partition function* Z_V and the *refined localized state* ω_V^t are defined similar to (2.39) and (2.40) by

$$Z_V^t(\alpha, \beta, \tilde{\rho}) := \int_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_<) \int_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_>) e^{\alpha \mathcal{T}_V^t(\tilde{\rho}, T_{\mathcal{U}_<, \mathcal{U}_>} \rho) + \beta \gamma_V^t(\tilde{\rho}, T_{\mathcal{U}_<, \mathcal{U}_>} \rho)} \quad (2.43)$$

$$\begin{aligned} \omega_V^t(\cdots) &:= \frac{1}{Z_V^t(\alpha, \beta, \tilde{\rho})} \int_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_<) \int_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_>) \\ &\quad \times e^{\alpha \mathcal{T}_V^t(\tilde{\rho}, T_{\mathcal{U}_<, \mathcal{U}_>} \rho) + \beta \gamma_V^t(\tilde{\rho}, T_{\mathcal{U}_<, \mathcal{U}_>} \rho)} (\cdots), \end{aligned} \quad (2.44)$$

where we again abbreviated the insertions by (\cdots) . The point is that, in contrast to γ_V^t , the functional \mathcal{T}_V^t is *not* a surface layer integral (due to the plus sign in (2.42)). The general picture is that, choosing α sufficiently large (and V as a set of finite measure), the main contribution to the group integral is obtained when the functional \mathcal{T}_V^t is maximal. This maximum should be attained when the sets $\Omega^t \cap V$ and $(M \setminus \Omega^t) \cap V$

coincide corresponding sets of the interacting spacetime. In this way, the freedom in performing translations in Minkowski is fixed. This general picture will be confirmed and worked out in detail in the present paper.

2.3. Basics on Fock Spaces and Entanglement. In this section, we recall the basics on Fock spaces and explain the phenomenon of entanglement.

For the construction of the *bosonic Fock space*, we take *symmetrized* products of the Hilbert space $(\mathfrak{h}, \langle \cdot | \cdot \rangle_\rho^t)$ of complexified linearized solutions as introduced in Section 2.2.1. We let $\mathfrak{h}^n = \mathfrak{h} \otimes \cdots \otimes \mathfrak{h}$ be the k -fold tensor product, endowed with the natural scalar product

$$(\psi_1 \otimes \cdots \otimes \psi_k | \phi_1 \otimes \cdots \otimes \phi_k)_\rho^t := (\psi_1 | \phi_1)_\rho^t \cdots (\psi_k | \phi_k)_\rho^t.$$

We denote total symmetrization by an index s , i.e.

$$(\psi_1 \otimes \cdots \otimes \psi_k)_s := \frac{1}{k!} \sum_{\sigma \in S_k} \psi_{\sigma(1)} \otimes \cdots \otimes \psi_{\sigma(k)},$$

where S_k denotes the group of all permutations. These totally symmetrized products, referred to as *product states* or *factorizable states*, do in general not form a vector space. But they generate a linear subspace denoted by

$$\mathcal{F}_{\rho,k}^b := \overline{(\mathfrak{h}^k)_s} \subset \mathfrak{h}^k.$$

The *bosonic Fock space* $(\mathcal{F}_\rho^b, \langle \cdot | \cdot \rangle_{\mathcal{F}_\rho^b})$ is the direct sum of the n -particle spaces,

$$\mathcal{F}_\rho^b = \bigoplus_{k=0}^{\infty} \mathcal{F}_{\rho,k}^b.$$

Likewise, the *fermionic Fock space* is constructed by taking totally *anti-symmetric* products of the one-particle Hilbert space $(\mathcal{H}_\rho^f, \langle \cdot | \cdot \rangle_\rho)$. We let $(\mathcal{H}_\rho^f)^k = \mathcal{H}_\rho^f \otimes \cdots \otimes \mathcal{H}_\rho^f$ be the k -fold tensor product, endowed with the natural scalar product

$$\langle \psi_1 \otimes \cdots \otimes \psi_k | \phi_1 \otimes \cdots \otimes \phi_k \rangle_{\mathcal{H}_\rho^f} := \langle \psi_1 | \phi_1 \rangle_\rho \cdots \langle \psi_k | \phi_k \rangle_\rho.$$

Totally anti-symmetrizing the tensor product gives the wedge product

$$\psi_1 \wedge \cdots \wedge \psi_k := \frac{1}{k!} \sum_{\sigma \in S_k} (-1)^{\text{sign}(\sigma)} \psi_{\sigma(1)} \otimes \cdots \otimes \psi_{\sigma(k)}$$

(here S_k denotes the set of all permutations and $\text{sign}(\sigma)$ is the sign of the permutation σ). The wedge product gives rise to a mapping

$$\Lambda_k : \underbrace{\mathcal{H}_\rho^f \times \cdots \times \mathcal{H}_\rho^f}_{k \text{ factors}} \rightarrow (\mathcal{H}_\rho^f)^k : (\psi_1, \dots, \psi_k) \mapsto \psi_1 \wedge \cdots \wedge \psi_k.$$

The vectors in the image of the mapping Λ_k are the k -particle *Hartree-Fock states* or *factorizable states*. Again, these states do in general not form a vector space. We denote the vector space generated by the k -particle Hartree-Fock states by

$$\mathcal{F}_{\rho,k}^f := \overline{\langle \Lambda_k((\mathcal{H}_\rho^f)^k) \rangle} \subset (\mathcal{H}_\rho^f)^k.$$

Taking the direct sum of these spaces gives the *fermionic Fock space* denoted by

$$\mathcal{F}_\rho^f := \bigoplus_{k=0}^{\infty} \mathcal{F}_{\rho,k}^f.$$

Finally, the Fock space \mathcal{F}_ρ is defined as the tensor product of the bosonic and fermionic Fock spaces,

$$\mathcal{F}_\rho := \mathcal{F}_\rho^{\text{b}} \otimes \mathcal{F}_\rho^{\text{f}}.$$

It is an important observation that the factorizable states do *not* form a vector space. A Fock vector which is not factorizable is said to be *entangled*. The simplest and most prominent entangled state is the spin singlet state of two observers, usually referred to as ALICE and BOB,

$$\Phi := \frac{1}{\sqrt{2}} (\phi_A^\uparrow \otimes \phi_B^\downarrow - \phi_A^\downarrow \otimes \phi_B^\uparrow)_s, \quad (2.45)$$

where \uparrow and \downarrow denote the polarizations (for ease of presentation of bosonic particles), and A and B refer to ALICE and BOB, respectively. Entanglement is a basic phenomenon of quantum physics, with important applications to quantum information theory and quantum computing. We refer the reader interested in the physical background to standard textbooks like [48, Section 20.4], [43, Section 2.2.8], [47, Section 3.10] or [36, 2].

3. QUANTUM STATES OF CAUSAL FERMION SYSTEMS AND ENTANGLEMENT

In [20], the interacting spacetime at time t was described by a *quantum state*, being a positive linear functional on the field algebra \mathcal{A} of the non-interacting spacetime (see the preliminaries in Sections 2.2.1–2.2.3). Moreover, various refinements of the constructions were given (in particular the *refined pre-state* and the *localized refined pre-state*; see the preliminaries in Sections 2.2.4 and 2.2.5). In this section, we reconsider these notions with the aim of determining whether they are capable of describing entanglement. Moreover, we shall set up the problem such as to obtain a clear mathematical setting which can be analyzed explicitly in a concise limiting case. In preparation, we begin with a few general considerations on the group integrals which appear in the definition of the partition function and the state (see for example (2.35) and (2.39)). These considerations partly anticipate results of the later analysis. They have the purpose of conveying the correct physical picture in nontechnical terms.

3.1. General Considerations on the Quantum State. In this section we give a few qualitative considerations on the structure of our quantum state and its various refinements. We begin with the partition function Z^t defined in (2.35). The main task of the present paper is to unravel how the integrand behaves as a function of \mathcal{U} . In particular, we need to quantify the main contributions to the integral. Let us now explain the qualitative picture for specific choices of $\tilde{\rho}$. In the case $\tilde{\rho} = \rho$ of the Minkowski vacuum, we expect that the integrand is largest if we choose $\mathcal{U} = \mathbb{1}$. In this case, the integrand can be computed rather explicitly in the formalism of the continuum limit (as developed in [10, 13]; see also Section 6.1 below). In particular, a direct computation shows that the surface layer integral γ^t in (2.35) has the scaling $\gamma^t \sim \varepsilon^{-4}$ and thus diverges in the limit $\varepsilon \searrow 0$ when the ultraviolet regularization is removed. This divergence can be understood from the fact that in the limit $\varepsilon \searrow 0$ the number of physical wave functions tends to infinity, and that the contributions by all these wave functions add up. More specifically, the contributions by all the wave functions to the kernel of the fermionic projector $P(x, y)$ are “in phase” on the light cone (i.e. if x and y are lightlike separated), giving rise to singularities as $\varepsilon \searrow 0$ (here by “in phase” we mean that all the summands have the same sign, and no relative phases appear). This

explains why the integrand in (2.35) is very large if $\mathcal{U} = \mathbb{1}$. For generic \mathcal{U} , however, the matrix elements of \mathcal{U} involve many phases. As a consequence, the contributions by the physical wave functions no longer add up, but instead we get sums of terms involving oscillatory phase factors. Similar to the effect of destructive interference, many summands will cancel each other. As a result, the integrand in (2.35) will be very small. Here the notions “very large” and “very small” will be made more precise in terms of a scaling behavior in ε (this is worked out in Section 6.1).

We next consider the situation when $\tilde{\rho}$ describes Minkowski space involving a *classical bosonic field* (for example a plane electromagnetic wave). In this case, the unitary operator \mathcal{U} can compensate for gauge phases, which means that the maximum of the integrand in (2.35) will be attained for a certain unitary operator \mathcal{U} for which the gauge phases are compensated in the optimal way. Then the surface layer integral can be computed again in the formalism of the continuum limit, giving a large contribution to the partition function. Moreover, the insertions in the quantum state (2.37) (as shown in more detail in (2.38)) become surface layer integrals which involve the electromagnetic field. In this way, the state ω^t makes it possible to probe the interacting spacetime using the free fields of the vacuum spacetime.

We now move on to the situation of a general minimizing measure $\tilde{\rho}$. This measure can have a very complicated structure. In particular, we cannot expect that the integrand in (2.35) will have a single maximum. Instead, the integrand will have many maxima, and the integral will give rise to a sum over contributions near the maxima (this picture will be made precise in Section 6). In simple terms, this means that $\tilde{\rho}$ can no longer be described by a single spacetime involving classical fields. Instead, it could be a “mixture” of different spacetimes, possibly involving many different classical fields. Even more, the measure ρ could have a structure which can no longer be described by classical fields or a classical spacetime. The main point is that, even in such very non-classical situations, the quantum state (2.37) makes it possible to probe the measure $\tilde{\rho}$ in terms of familiar objects in Minkowski space. It turns out that this probing gives rise to a quantum state in the sense familiar from the algebraic formulation of *quantum field theory*. In this way, a general measure $\tilde{\rho}$ can be described in the language of quantum field theory. This description does not only allow for the description of quantum fields, but it even gives words like “quantum spacetime” and “quantum geometry” a well-defined meaning in the realm of quantum field theory.

Exactly as explained for the vacuum state, for a generic unitary operator \mathcal{U} the integrand in (2.35) will typically be very small due to destructive interference. In other words, the integrand should be close to zero, except on a set of very small measure, where the integrand should be very large. A general conclusion from these qualitative consideration is that we must face the difficulty that the integrand in (2.43) (and similar in the various states and pre-states) will have a highly nonlinear dependence on \mathcal{U} . In particular, we cannot expect that a power expansion in \mathcal{U} will be sufficient for our purposes. Instead, we need to develop methods for evaluating group integrals in the non-perturbative regime.

Before entering a more detailed discussion of the state and its various refinements, we comment on a question which the critical reader may ask: *Why is there a freedom in introducing the quantum state? Should the quantum state as a physical object not be given canonically in a unique way?* In order to answer this question, we return to our motivation and our general procedure for introducing the quantum state: We wanted to probe the interacting measure $\tilde{\rho}$ using the objects of the vacuum spacetime. The

fact that the measure $\tilde{\rho}$ can have a very complicated structure can also be expressed by saying that this measure contains a lot of information. Only a small portion of this information should be relevant for describing macroscopic phenomena (for example, the measure $\tilde{\rho}$ also encodes the detailed structure of spacetime on the Planck scale, which cannot be tested in current experiments). Therefore, our task is to extract the portion of the information on $\tilde{\rho}$ which is relevant for describing quantum physics. Thinking of the problem in this way, it is no longer surprising that there is not a canonical way to retrieve this information. Similar to the usual situation in physics that a physical quantity can be determined alternatively in different types of experiments, there may be many different ways to retrieve the relevant information from $\tilde{\rho}$. We can make use of this freedom to our advantage. Similar to setting up an optimal experiment, we can try to come up with a definition of a quantum state which has all desirable properties and is easiest to analyze in detail. The following considerations can be regarded as consecutive steps towards this goal.

3.2. Why Entanglement Requires the Refined Pre-State. Clearly, entanglement is a fundamental feature of quantum physics. Therefore, any physically sensible notion of quantum state should be capable of describing entanglement. We now explain why this requirement makes it necessary to consider the *refined* pre-state.

Our arguments are very general and do not make use of the specific form of the interacting measure. Let us assume that we want to realize an entangled Fock state, which can be written as a linear combination of product states. For simplicity, we consider only a bosonic Fock vector,

$$\Phi = \sum_{\alpha=1}^K \Phi_{\alpha} \quad \text{with} \quad \Phi_{\alpha} = (\phi_{1,\alpha} \otimes \cdots \otimes \phi_{p_{\alpha},\alpha})_s. \quad (3.1)$$

Then the expectation value of an observable \mathcal{O} can be written as a double sum,

$$\langle \Phi | \mathcal{O} | \Phi \rangle_{\mathcal{F}_{\rho}^b} = \sum_{\alpha, \beta=1}^K \langle \Phi_{\alpha} | \mathcal{O} | \Phi_{\beta} \rangle_{\mathcal{F}_{\rho}^b}.$$

As a typical example of a p -particle measurement we choose \mathcal{O} as a product of bosonic field operators,

$$\mathcal{O} = a^{\dagger}(z_1^t) \cdots a^{\dagger}(z_p^t) a(\bar{z}_1^t) \cdots a(\bar{z}_q^t).$$

In this case, the expectation value can be written as

$$\langle \Phi | \mathcal{O} | \Phi \rangle_{\mathcal{F}_{\rho}^b} = \sum_{\alpha, \beta=1}^K \langle a(\bar{z}_1^t) \cdots a(\bar{z}_p^t) \Phi_{\alpha} | a(\bar{z}_1^t) \cdots a(\bar{z}_q^t) \Phi_{\beta} \rangle_{\mathcal{F}_{\rho}^b}.$$

Using that the Φ_{α} are product states (3.1), on the left one gets products of scalar products $(z_{\ell}^t | \phi_{\ell',\alpha})_{\rho}^t$ with $\ell, \ell' \in \{1, \dots, p\}$. Similarly, on the right one gets the scalar products $(z_{\ell}^t | \phi_{\ell',\beta})_{\rho}^t$. The point is that all the factors on the left involve the same index α (describing the *bra*), whereas all the factors on the right involve the same index β (describing the *ket*).

Our goal is to realize this expectation value by associating to the observable \mathcal{O} suitable insertions in (1.2). These insertions are complex-valued functions which depend on the corresponding linearized solutions z_{ℓ}^t or \bar{z}_{ℓ}^t . Moreover, the insertions clearly

depend on the unitary operator \mathcal{U} . Denoting the insertions by $b^t(z, \mathcal{U})$, the state takes the form

$$\begin{aligned} & \omega^t \left(a^\dagger(z_1^t) \cdots a^\dagger(z_p^t) a(\bar{z}_1^t) \cdots a(\bar{z}_q^t) \right) \\ &:= \frac{1}{Z^t(\beta, \tilde{\rho})} \int_{\mathcal{G}} b^t(z_1^t, \mathcal{U}) \cdots b^t(z_p^t, \mathcal{U}) \overline{b^t(z_1^t, \mathcal{U}) \cdots b^t(z_q^t, \mathcal{U})} e^{\beta \gamma^t(\tilde{\rho}, \mathcal{U}\rho)} d\mu_{\mathcal{G}}(\mathcal{U}). \end{aligned} \quad (3.2)$$

In order to encode the Fock components, we write every insertion as a sum of terms indexed by α . This leads us to the general ansatz

$$b^t(z, \mathcal{U}) = \sum_{\alpha=1}^K b_{\alpha}^t(z, \mathcal{U}). \quad (3.3)$$

Ultimately, the insertions with index α should encode the form of the α -component of the Fock vector (3.1). At this stage, we do not need to specify what the α -dependence of the insertions really means (this will be made precise in Section 6.5 and explained in Section 7). Instead, for explaining the basic mechanism it is preferable to work with (3.3) symbolically. Using this ansatz in (3.2) and multiplying out, we obtain

$$\begin{aligned} & \omega^t \left(a^\dagger(z_1^t) \cdots a^\dagger(z_p^t) a(\bar{z}_1^t) \cdots a(\bar{z}_q^t) \right) = \frac{1}{Z^t(\beta, \tilde{\rho})} \sum_{\alpha_1, \dots, \alpha_p=1}^K \sum_{\beta_1, \dots, \beta_q=1}^K \\ & \times \int_{\mathcal{G}} b_{\alpha_1}^t(z_1^t, \mathcal{U}) \cdots b_{\alpha_p}^t(z_p^t, \mathcal{U}) \overline{b_{\beta_1}^t(z_1^t, \mathcal{U}) \cdots b_{\beta_q}^t(z_q^t, \mathcal{U})} e^{\beta \gamma^t(\tilde{\rho}, \mathcal{U}\rho)} d\mu_{\mathcal{G}}(\mathcal{U}). \end{aligned}$$

Here the insertions and the exponential factor depend nonlinearly on the unitary operator \mathcal{U} . As a consequence, the integral in general does not split into a product of sums. Instead, integrating over \mathcal{U} will give rise to correlations between the different insertions. These correlations are crucial for getting a connection to the interacting many-particle picture of quantum field theory. More specifically, in order to describe entangled states, we would like that all the holomorphic insertions should be in the same component, and similarly for the anti-holomorphic insertions. Therefore, the state should be approximately of the form

$$\begin{aligned} & \omega^t \left(a^\dagger(z_1^t) \cdots a^\dagger(z_p^t) a(\bar{z}_1^t) \cdots a(\bar{z}_q^t) \right) \approx \frac{1}{Z^t(\beta, \tilde{\rho})} \sum_{\alpha, \beta=1}^K \\ & \times \int_{\mathcal{G}} b_{\alpha}^t(z_1^t, \mathcal{U}) \cdots b_{\alpha}^t(z_p^t, \mathcal{U}) \overline{b_{\beta}^t(z_1^t, \mathcal{U}) \cdots b_{\beta}^t(z_q^t, \mathcal{U})} e^{\beta \gamma^t(\tilde{\rho}, \mathcal{U}\rho)} d\mu_{\mathcal{G}}(\mathcal{U}). \end{aligned} \quad (3.4)$$

Following a notion introduced in [12], we also say that *bra* and *ket* should be *synchronized*.

In order to clarify this construction, we work it out in more detail for the spin singlet state

Example 3.1. For the spin singlet state (2.45), we have two components $\alpha = 1, 2$ and

$$\Phi_1 = \frac{1}{\sqrt{2}} (\phi_A^\uparrow \otimes \phi_B^\downarrow)_s \quad \text{and} \quad \Phi_2 = -\frac{1}{\sqrt{2}} (\phi_A^\downarrow \otimes \phi_B^\uparrow)_s.$$

ALICE and BOB choose one-particle measurement operators, which we write in *bra/ket*-notation as

$$\mathcal{O}_A := |z'_A\rangle\langle z_A| \quad \text{and} \quad \mathcal{O}_B := |z'_B\rangle\langle z_B|.$$

Then the two-particle measurement is described by the observable $\mathcal{O} = \mathcal{O}_A \otimes \mathcal{O}_B$. We thus obtain

$$\begin{aligned} \langle \Phi | \mathcal{O} | \Phi \rangle_{\mathcal{F}_\rho^b} &= \sum_{\alpha, \beta=1}^K \langle \Phi_\alpha | z'_A \otimes z'_B \rangle \langle z_A \otimes z_B | \Phi_\alpha \rangle \\ &= \frac{1}{8} \left((\phi_A^\uparrow | z'_A)_\rho^t (\phi_B^\downarrow | z'_B)_\rho^t (z_A | \phi_A^\uparrow)_\rho^t (z_B | \phi_B^\downarrow)_\rho^t - (\phi_A^\downarrow | z'_A)_\rho^t (\phi_B^\uparrow | z'_B)_\rho^t (z_A | \phi_A^\uparrow)_\rho^t (z_B | \phi_B^\downarrow)_\rho^t \right. \\ &\quad \left. - (\phi_A^\uparrow | z'_A)_\rho^t (\phi_B^\downarrow | z'_B)_\rho^t (z_A | \phi_A^\downarrow)_\rho^t (z_B | \phi_B^\uparrow)_\rho^t + (\phi_A^\downarrow | z'_A)_\rho^t (\phi_B^\uparrow | z'_B)_\rho^t (z_A | \phi_A^\downarrow)_\rho^t (z_B | \phi_B^\uparrow)_\rho^t \right). \end{aligned}$$

If a synchronization is present, these expectation values can be realized by the quantum state (3.4) if we choose $p = q = 2$ and

$$z'_1 = z'_A, \quad z'_2 = z'_B \quad \text{and} \quad z_1 = z_A, \quad z_2 = z_B.$$

Moreover, we need to arrange that the insertions are the corresponding one-particle expectation values, i.e.

$$\begin{aligned} b_1^t(z_A^t) &= c(\phi_A^\uparrow | z'_A)_\rho^t, & b_1^t(z_B^t) &= c(\phi_B^\downarrow | z'_B)_\rho^t \\ b_2^t(z_A^t) &= -c(\phi_A^\downarrow | z'_A)_\rho^t, & b_2^t(z_B^t) &= c(\phi_B^\uparrow | z'_B)_\rho^t \end{aligned}$$

with $c = 8^{-\frac{1}{4}}$, and similarly without the primes.

Without synchronization, however, the many-particle measurement reduces to a product of one-particle measurements. A direct computation (similar to the original Einstein-Podolsky-Rosen argument) shows that it is indeed impossible to realize the expectation values of the spin-singlet state. \diamond

In view of this example, the basic question is whether (3.4) can hold, i.e. whether by a suitable choice of the insertions we can arrange a synchronization of *bra* and *ket*. The answer is no, as the next lemma shows.

Lemma 3.2. *In the case $p = q = 2$ and choosing $z_\ell^t = z_\ell^t = z^t$, the bosonic state (3.2) satisfies the inequality*

$$\begin{aligned} &\sum_{\alpha, \beta=1}^K \left| \int_{\mathcal{G}} b_\alpha^t(z^t) b_\alpha^t(z^t) \overline{b_\beta^t(z^t) b_\beta^t(z^t)} e^{\beta \gamma^t(\bar{\rho}, \mathcal{U}\rho)} d\mu_{\mathcal{G}}(\mathcal{U}) \right| \\ &\leq \sum_{\alpha, \beta=1}^K \int_{\mathcal{G}} b_\alpha^t(z^t) b_\beta^t(z^t) \overline{b_\alpha^t(z^t) b_\beta^t(z^t)} e^{\beta \gamma^t(\bar{\rho}, \mathcal{U}\rho)} d\mu_{\mathcal{G}}(\mathcal{U}). \end{aligned}$$

Proof. Using that the exponential factor is positive, we obtain with the help of the Schwarz inequality

$$\begin{aligned} &\left| \int_{\mathcal{G}} b_\alpha^t(z^t) b_\alpha^t(z^t) \overline{b_\beta^t(z^t) b_\beta^t(z^t)} e^{\beta \gamma^t(\bar{\rho}, \mathcal{U}\rho)} d\mu_{\mathcal{G}}(\mathcal{U}) \right| \\ &\leq \int_{\mathcal{G}} |b_\alpha^t(z^t)|^2 |b_\beta^t(z^t)|^2 e^{\beta \gamma^t(\bar{\rho}, \mathcal{U}\rho)} d\mu_{\mathcal{G}}(\mathcal{U}). \end{aligned}$$

Using that the anti-holomorphic insertion is the complex conjugate of the holomorphic insertion, the result follows. \square

This lemma shows that, no matter how the insertions are chosen, the main contribution to the state is *not* of the form (3.4), because the contributions with synchronizations between *bra* and *ket* dominate. We conclude that the state ω^t is not capable

of capturing essential features of the quantum dynamics of the causal fermion system. The *refined* pre-state (2.40), however, does *not* suffer from this shortcoming. This will be worked out in detail in Section 7 later in this paper. In simple terms, the double integral in (2.44) gives rise to separate phases for the holomorphic and anti-holomorphic insertions, so that the counter argument of Lemma 3.2 no longer applies.

3.3. Why it is Preferable to Work with Localized States. As explained qualitatively in Section 3.1, we need to compute integrals over the unitary group for integrands with a highly nonlinear dependence on the unitary operator \mathcal{U} . In order to get into the position where such integrals can be analyzed explicitly, it is very helpful to work with the *localized* states, as we now explain. Since we already know from the previous section that we must consider refined states, we consider the localized refined pre-state (see (2.43) and (2.44) in Section 2.2.5). In this setting, the pre-state is computed in a subset V of our spacetime. This subset can be chosen arbitrarily large with the only constraint that it should be small compared to the size of the whole spacetime. Clearly, if the whole system is Minkowski space, this constraint is unproblematic because as the whole spacetime we can consider an arbitrarily large box in Minkowski space (more details and the relevant scalings will be given in Section 6 below).

Associating to V a subspace $\mathcal{H}^{\text{lab}} \subset \mathcal{H}$ of the whole Hilbert space, the integrand of the group integrals depends only on the restriction of the unitary operator to this subspace, i.e. on the operator

$$\pi_{\mathcal{H}^{\text{lab}}} \mathcal{U} \pi_{\mathcal{H}^{\text{lab}}}$$

(where $\pi_{\mathcal{H}^{\text{lab}}} : \mathcal{H} \rightarrow \mathcal{H}^{\text{lab}}$ denotes the orthogonal projection). Therefore, we can carry out the integrals over all the other matrix elements of \mathcal{U} . Doing so simplifies the structure of the remaining integrals considerably (for details see Theorem 4.2 in Section 4.2). Moreover, we can consider the limiting case when the dimension of \mathcal{H} tends to infinity, whereas the dimension of \mathcal{H}^{lab} remains fixed. This simplifies the formulas further, making it possible to analyze the group integrals in detail.

We point out that working with localized states is not preferable for principal or conceptual reasons, but it only has the computational benefit that it becomes possible to evaluate the group integrals in a mathematically precise limiting case.

3.4. The Precise Form of the Localized Refined Pre-State. We now make the setting of the localized refined pre-state more concrete. In order to describe the vacuum, we let $(\mathcal{H}, \mathcal{F}, \rho)$ be a causal fermion system constructed from a vacuum Dirac system in a three-dimensional box of length $2L$, i.e. in the spacetime cylinder

$$\mathcal{M} = \mathbb{R} \times [-L, L]^3 \subset \mathbb{R}^{1,3} \quad (3.5)$$

with periodic boundary conditions (similar to the standard construction in Minkowski space as given in [13, Section 1.2] or in more detail in [44]). Fixing the scale $\varepsilon > 0$ of the ultraviolet regularization, the Dirac sea is built up of a finite number f of wave functions with the scaling

$$\dim \mathcal{H} = f \sim \left(\frac{L}{\varepsilon} \right)^3. \quad (3.6)$$

The set V is chosen as

$$V = [-T, T] \times [-\ell, \ell]^3 \quad \text{with} \quad \ell \ll L. \quad (3.7)$$

Thus ℓ can be thought of as the size of the physical system of interest. Moreover, the parameter T is the time scale on which the measurement takes place. The three-dimensional box $[-\ell, \ell]^3$ can be associated to a subspace of \mathcal{H} consisting of all wave functions which at time t are localized inside the box (except for exponentially decaying tails which arise due to the incompatibility of spatial localization and frequency splitting; this effect will not be of relevance for our computations). We denote this subspace by $\mathcal{H}^{\text{lab}} \subset \mathcal{H}$. Its dimension scales like

$$\dim \mathcal{H}^{\text{lab}} \sim \left(\frac{\ell}{\varepsilon}\right)^3. \quad (3.8)$$

We are interested in the asymptotics $L \rightarrow \infty$ and $\varepsilon \searrow 0$ for fixed ℓ and T .

We choose the group \mathcal{G} as the unitary group of the whole Hilbert space,

$$\mathcal{G} = \text{U}(\mathcal{H}). \quad (3.9)$$

This is different from [20], where we wanted to allow for the possibility that the Hilbert space \mathcal{H} is infinite dimensional, making it necessary to choose \mathcal{G} as the unitary group of a finite-dimensional subspace \mathcal{H}^f . In order to have a simple and clean setting, we here decided to choose \mathcal{H} to be finite-dimensional (3.6). In this setting, the choice (3.9) is most natural. But clearly, the dimension of \mathcal{H} tends to infinity if we let $L \rightarrow \infty$.

For the later computations, we set $N = f = \dim \mathcal{H}$ and often identify \mathcal{H} with \mathbb{C}^N . Apart from the subspace \mathcal{H}^{lab} , we also need to consider the subspace $\mathcal{H}^f \subset \mathcal{H}$ of *low-energy wave functions*. Here by “low energy” we mean that their energy is much smaller than the Planck energy. In technical terms, the wave functions in \mathcal{H}^f must satisfy the conditions needed for the construction of the extended Hilbert space \mathcal{H}_ρ^f (see the preliminaries in Section 2.1.11). For the constructions of the present paper, we do not need to specify \mathcal{H}^f . We recall from (2.29) that its dimension is denoted by f^f . Finally, we need to decompose the Hilbert space of the laboratory into low- and high-energy wave functions. To this end, we set

$$\mathcal{H}^{\text{le}} := \mathcal{H}^f \cap \mathcal{H}^{\text{lab}} \quad \text{and} \quad \mathcal{H}^{\text{he}} := (\mathcal{H}^f)^\perp \cap \mathcal{H}^{\text{lab}},$$

so that

$$\mathcal{H}^{\text{lab}} = \mathcal{H}^{\text{le}} \oplus \mathcal{H}^{\text{he}}.$$

Before developing the methods and entering the detailed analysis, we make two remarks. First, in order to get into the regime where the integrand depends highly nonlinearly on \mathcal{U} (as described qualitatively in Section 3.1), we need to insert a factor N into the exponents in (2.43) and (2.44). Thus, when letting the dimension of \mathcal{H} tend to infinity, we also increase the nonlinearity of the exponential. The reason for the linear scaling in N will be explained in Example 4.9; it will also become clear in a more general setting in the model examples in Section 5. The second remark is that we set the parameter β equal to zero. This leads us to the definition of the refined localized state $\omega_V^t(\cdots)$ as stated in the introduction (see (1.2) and (1.1)). The precise definition of the insertions requires more preparations and is therefore postponed to Section 6.6.

We conclude this section by explaining how the simplification of setting β to zero is to be understood. In preparation, we recall how the exponential factors $\exp(\beta\gamma_V^t)$ and $\exp(\alpha\mathcal{T}_V^t)$ came about. The factor $\exp(\beta\gamma^t)$ was introduced in [19] motivated by the fact that it coincides with the norm of the corresponding bosonic Fock vector (see [19, Sections 7 and 8]). Moreover, in [20] this factor has the benefit that the insertions can be understood as variational derivatives of the partition function (see [20,

Section 4.6]). The factor $\exp(\alpha\mathcal{T}_V^t)$, on the other hand, was introduced in the context of localized states in order to fix the freedom to perform translations in M (see [20, Section 5.3] or the preliminaries in Section 2.2.5). The reason why this freedom cannot be fixed with the factor $\exp(\beta\gamma_V^t)$ is that the surface layer integral γ_V^t is not necessarily positive. But \mathcal{T}_V^t is not the only positive functional. In particular, one could replace the spectral weight $|x\mathcal{U}_{>}y\mathcal{U}_{<}^{-1}|$ in (2.42) by the Lagrangian,

$$\left(\int_{\tilde{\Omega}^t \cap \tilde{V}} d\tilde{\rho}(x) \int_{\Omega^t \cap V} d\rho(y) + \int_{(\tilde{M} \setminus \tilde{\Omega}^t) \cap \tilde{V}} d\tilde{\rho}(x) \int_{(M \setminus \Omega^t) \cap V} d\rho(y) \right) \mathcal{L}(x, \mathcal{U}_{>} y \mathcal{U}_{<}^{-1}). \quad (3.10)$$

Working with the Lagrangian seems more natural because of the closer connection to the causal action principle. However, the Lagrangian is more difficult to compute, because the summands of different signs in (2.1) may and typically will cancel each other, making it rather subtle to determine the leading contributions. In the spectral weight (2.6), on the other hand, all summands are positive, so that it suffices to compute the leading contributions to the eigenvalues. This is the reason why we here prefer to work with the functional \mathcal{T}_V^t . The last consideration also shows that \mathcal{T}_V^t will typically be much larger than the surface layer integral γ_V^t constructed out of the Lagrangian. This justifies why we can leave γ_V^t out of the exponent by setting $\beta = 0$.

To summarize, at present the detailed form of the functional in the exponent of the localized refined quantum state is not completely fixed. We here choose the functional which is most suitable for the computations. The differences to other possible choices are mainly technical and should not leave the basic mechanisms and results of this paper unchanged.

4. COMPUTATION OF INTEGRALS OVER THE UNITARY GROUP

4.1. Preliminary Considerations. In general terms, our task is to analyze integrals over the unitary group $U(N)$ asymptotically as $N \rightarrow \infty$. In order to motivate our methods and to explain how they come about, we now begin with a few elementary considerations. We always denote the unitary group by $\mathcal{G} = U(N)$ and let $\mu_{\mathcal{G}}$ be the normalized Haar measure on \mathcal{G} . For notational clarity, integrals with respect to such a normalized measure will always be denoted by $\int_{\mathcal{G}}$. The simplest integrals are obtained by choosing the integrand as a monomial in the matrix entries and their complex conjugates, i.e.

$$\int_{\mathcal{G}} u_{j_1}^{i_1} \cdots u_{j_p}^{i_p} \overline{u_{l_1}^{k_1} \cdots u_{l_q}^{k_q}} d\mu_{\mathcal{G}}(u) \quad (4.1)$$

with $p, q \in \mathbb{N}_0$ and arbitrary indices in $\{1, \dots, N\}$. The easiest method is to make use of the invariance of the Haar measure under transformations

$$u \rightarrow V u W \quad \text{with} \quad V, W \in \mathcal{G}. \quad (4.2)$$

In particular, choosing $V = e^{i\varphi}$ as a phase factor and $W = \mathbf{1}$, one sees that the group integral (4.1) vanishes unless in the case $p = q$ where the number of conjugated and unconjugated factors coincides. Therefore, the simplest non-trivial integral is obtained in the case of two factors by

$$\int_{\mathcal{G}} |u_j^i|^2 d\mu_{\mathcal{G}}(u) \quad (4.3)$$

for given indices i, j . Applying the invariance (4.2) again for $W = 1$ and a unitary operator V which flips the components i and k , i.e.

$$V_b^a = \delta_b^a - \delta^{ai} \delta_{bi} - \delta^{ak} \delta_{bk} + \delta^{ai} \delta_{bk} + \delta^{ak} \delta_{bi} ,$$

one sees that the integral (4.3) is the same for every choice of i . In particular,

$$\oint_{\mathcal{G}} |\mathcal{U}_j^i|^2 d\mu_{\mathcal{G}}(\mathcal{U}) = \frac{1}{N} \sum_{k=1}^N \oint_{\mathcal{G}} |\mathcal{U}_j^k|^2 d\mu_{\mathcal{G}}(\mathcal{U}) .$$

The components $\mathcal{U}_j^1, \dots, \mathcal{U}_j^N$ form the j^{th} row of the matrix \mathcal{U} . For any unitary matrix, this row vector is a unit vector, i.e.

$$\sum_{k=1}^N |\mathcal{U}_j^k|^2 = 1 .$$

We thus obtain

$$\oint_{\mathcal{G}} |\mathcal{U}_j^i|^2 d\mu_{\mathcal{G}}(\mathcal{U}) = \frac{1}{N} \oint_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}) = \frac{1}{N} .$$

More generally, the above method yields

$$\oint_{\mathcal{G}} \mathcal{U}_j^i (\mathcal{U}^{-1})_k^l d\mu_{\mathcal{G}}(\mathcal{U}) = \oint_{\mathcal{G}} \mathcal{U}_j^i \overline{\mathcal{U}_k^l} d\mu_{\mathcal{G}}(\mathcal{U}) = \frac{1}{N} \delta_k^i \delta_j^l .$$

We now move on to the integral (4.1) with four unitary factors. Using the unitary invariance (4.2), we get zero unless $p = q = 2$ and each row and column index appears twice, once with and once without conjugation. Thus, after commuting the factors, it suffices to consider the integral

$$\oint_{\mathcal{G}} \mathcal{U}_{j_1}^{i_1} \mathcal{U}_{j_2}^{i_2} \overline{\mathcal{U}_{j_1}^{k_1}} \overline{\mathcal{U}_{j_2}^{k_2}} d\mu_{\mathcal{G}}(\mathcal{U})$$

with either $i_1 = k_1, i_2 = k_2$ or $i_1 = k_2, i_2 = k_1$. The corresponding row vectors

$$u^i := \mathcal{U}_{j_1}^i \quad \text{and} \quad v^i := \mathcal{U}_{j_2}^i$$

are again unit vectors. Moreover, in the case $j_1 \neq j_2$ they are orthogonal. This orthogonality condition makes the computation more difficult. But the following qualitative argument suggests that the situation should simplify asymptotically for large N : The scalar product

$$\langle u, v \rangle_{\mathbb{C}^N} = \sum_{i=1}^n \overline{u^i} v^i \tag{4.4}$$

involves a sum over the components. Each summand can be positive or negative, so that in the sum there may be cancellations of terms with different signs. These cancellations will be more efficient if N is large. In other words, taking randomly chosen unit vectors u and v , their scalar product (4.4) typically becomes very small if N gets large. Asymptotically, the integral (4.4) should factor into two integrals over u and v , both over the unit sphere in \mathbb{C}^N .

This qualitative argument can be made precise, giving rise to the following result.

Lemma 4.1. *Given $p, N \in \mathbb{N}$, we let $\mathcal{G} = \mathrm{U}(N)$ be the unitary group with normalized Haar measure $d\mu_{\mathcal{G}}$. Then, asymptotically for large N ,*

$$\begin{aligned} & \int_{\mathcal{G}} \mathcal{U}_{j_1}^{i_1} \cdots \mathcal{U}_{j_p}^{i_p} (\mathcal{U}^{-1})_{l_1}^{k_1} \cdots (\mathcal{U}^{-1})_{l_p}^{k_p} d\mu_{\mathcal{G}}(\mathcal{U}) \\ &= \frac{1}{N^p} \sum_{\sigma \in S(p)} \delta_{l_{\sigma(1)}}^{i_1} \delta_{j_1}^{k_{\sigma(1)}} \cdots \delta_{l_{\sigma(p)}}^{i_p} \delta_{j_p}^{k_{\sigma(p)}} + \mathcal{O}\left(\frac{1}{N^{p+1}}\right), \end{aligned} \quad (4.5)$$

where $S(p)$ denotes the symmetric group.

This result can be understood in analogy to Gaussian integrals or, similarly, to Wick's rule for free bosonic quantum fields: Integrating over the unitary group gives rise to pairings of the unitary matrices, and we need to sum over all possible pairings. Each pair involves a factor \mathcal{U} and a factor \mathcal{U}^{-1} . A pairing consists in replacing the two matrices by two Kronecker deltas which connect the indices of \mathcal{U} and \mathcal{U}^{-1} crosswise. We refer to asymptotic formulas of type (4.5) as the *Gaussian asymptotics*. There are various ways to prove the above lemma. One method is to analyze properties of tensor representations of the unitary group, as inspired by Michael Creutz [7]. In order not to distract from the main ideas, the resulting proof of Lemma 4.1 is given in Appendix A. Another method for proving formulas similar to (4.5) is to get a direct connection to Gaussian integrals; this will be explained in Section 4.4 below. Other methods can be found in [49, 6, 38].

It is an important observation that, for our purposes, the Gaussian asymptotics is *not* sufficient. Namely, as already described qualitatively in Section 3.1, we have the situation in mind where the integrand is very small, except on a set of very small measure, where the integrand is very large. Such a “spiky landscape” is in conflict with the Gaussian asymptotics, where we are only allowed to evaluate polynomials, and where the factorization via Wick rules means that the components of the unitary matrices are correlated only weakly in the sense that every pairing gives a small factor $1/N$.

In order to evaluate group integrals in the strongly correlated regime, we make essential use of the fact that the *localized* quantum state only depends on the physical wave functions inside the laboratory as described by the subsets $V \subset M$ and $\tilde{V} \subset \tilde{M}$ (see (2.41)). More technically, we may choose a subspace $I \subset \mathbb{C}^N$ of dimension q and write the group integral as

$$\int_{\mathcal{G}} f(N, A) d\mu_{\mathcal{G}}(\mathcal{U}) \quad \text{with} \quad A := \pi_I \mathcal{U} \pi_I \quad (4.6)$$

(and $\pi_I : \mathbb{C}^N \rightarrow I$ is the orthogonal projection). Now we can take the limit

$$N \rightarrow \infty \quad \text{for fixed } q = \dim I.$$

In other words, we let the whole system get larger and larger while keeping the size of our laboratory unchanged. In view of the size of the whole universe compared to the size of a typical measurement apparatus, this limiting case seems physically sensible for most applications in mind. It turns out that, in this limiting case, the group integral becomes tractable, even in the case that the function f in (4.6) is highly nonlinear and cannot be expanded in a power series or has an explicit N -dependence. This technical simplification was even one of our motivations for introducing the localized state. In the case that the function f is a monomial, we shall recover the Gaussian asymptotics similar to that in (4.5).

After these introductory words, we now enter the analysis of group integrals of the form (4.6) and work out the different limiting cases needed for our purposes.

4.2. Integrals over Functions on a Subspace. The following formula is the core of our method for computing the quantum states.

Theorem 4.2. *Let $I \subset \mathcal{H}$ with $\dim I = q$, $\dim \mathcal{H} = N$ and $N \geq 2q$. Let $f \in C^0(\mathcal{A}_q, \mathbb{C})$ be a function of $A := \pi_I \mathcal{U}|_I$ only (where $\pi_I : \mathcal{H} \rightarrow I$ is the orthogonal projection). Then its group integral over $\mathcal{G} = \mathrm{U}(N)$ with respect to the normalized Haar measure $\mu_{\mathcal{G}}$ can be written as*

$$\begin{aligned} \int_{\mathcal{G}} f(A) d\mu_{\mathcal{G}}(\mathcal{U}) &= \frac{1}{\pi^{(q^2)}} \frac{(N-1)!(N-2)! \cdots (N-q)!}{(N-q-1)!(N-q-2)! \cdots (N-2q)!} \\ &\quad \times \int_{\mathcal{A}_q} f(A) (\det(\mathbf{1} - A^* A))^{N-2q} d\mu_{\mathbb{C}^{q \times q}}(A). \end{aligned} \quad (4.7)$$

Here \mathcal{A}_q is the set of matrices

$$\mathcal{A}_q := \{A \in \mathbb{C}^{q \times q} \mid A^* A < \mathbf{1}_{\mathbb{C}^q}\} \subset \mathbb{C}^{q \times q}, \quad (4.8)$$

and $\mu_{\mathbb{C}^{q \times q}}$ is the Lebesgue measure on $\mathbb{C}^{q \times q} \simeq \mathbb{R}^{2q^2}$, i.e.

$$d\mu_{\mathbb{C}^{q \times q}}(A) = \prod_{j,k=1}^q dx_k^j dy_k^j \quad \text{with} \quad x_k^j := \operatorname{Re} A_k^j, \quad y_k^j := \operatorname{Im} A_k^j.$$

Similar results can be found in the literature (see [42, Theorem 1.3] or [32, 33, 3]). For completeness, and in order to get the prefactors right, we shall give a detailed proof of this theorem. Before beginning, we state two Corollaries. The first is the special case when f is chosen as the constant function one.

Corollary 4.3. *Under the assumptions of Theorem 4.2,*

$$\int_{\mathcal{A}_q} (\det(\mathbf{1} - A^* A))^{N-2q} d\mu_{\mathbb{C}^{q \times q}}(A) = \pi^{(q^2)} \frac{(N-q-1)!(N-q-2)! \cdots (N-2q)!}{(N-1)!(N-2)! \cdots (N-q)!}.$$

Next, expanding the factorials in (4.7), one immediately gets the following result.

Corollary 4.4. *The integral (4.7) has the leading asymptotics for large N*

$$\int_{\mathcal{G}} f(A) d\mu_{\mathcal{G}}(\mathcal{U}) = \left(\frac{N}{\pi}\right)^{(q^2)} \int_{\mathcal{A}_q} f(A) (\det(\mathbf{1} - A^* A))^{N-2q} d\mu_{\mathbb{C}^{q \times q}}(A) \left(1 + \mathcal{O}\left(\frac{1}{N}\right)\right).$$

The remainder of this section is devoted to the proof of Theorem 4.2. We let e_1, \dots, e_q be an orthonormal basis of I and extend it to an orthonormal basis of \mathcal{H} . Then for the evaluation of the integral in (4.7) it suffices to restrict attention to the first q columns of the unitary matrix \mathcal{U} . The corresponding column vectors, denoted by

$$\mathcal{U}_i \in \mathbb{C}^N \quad \text{with } i \in \{1, \dots, q\},$$

are orthonormal. For computational purposes, it is most convenient to realize the orthonormality by integrating over all matrix entries and inserting δ -distributions. This leads us to introduce on $\mathbb{C}^{N \times q}$ the measure

$$d\mu_{\mathcal{G}|_I} := \prod_{j=1}^q \delta(\|\mathcal{U}_j\|^2 - 1) \prod_{i < j} \delta(\operatorname{Re} \langle \mathcal{U}_i, \mathcal{U}_j \rangle) \delta(\operatorname{Im} \langle \mathcal{U}_i, \mathcal{U}_j \rangle) d\mathcal{U}_i, \quad (4.9)$$

where $d\mathcal{U}_i \in \mathbb{C}^N \simeq \mathbb{R}^{2N}$ is the Lebesgue measure on the i^{th} column,

$$\prod_{k=1}^N d\operatorname{Re} \mathcal{U}_i^k d\operatorname{Im} \mathcal{U}_i^k.$$

Since the measure (4.9) is invariant under unitary transformations, and the Haar measure is uniquely determined by unitary invariance and the normalization, these two measures coincide up to a constant, i.e.

$$\int_{\mathcal{G}} f(A) d\mu_{\mathcal{G}}(\mathcal{U}) = \frac{1}{c} \int_{\mathbb{C}^{N \times q}} f(A) d\mu_{\mathcal{G}|_I}. \quad (4.10)$$

We first compute the normalization constant.

Lemma 4.5. *The constant c in (4.10) is given by*

$$c = \int_{\mathbb{C}^{N \times q}} d\mu_{\mathcal{G}|_I} = \frac{\pi^{\frac{q(2N+1-q)}{2}}}{(N-1)! \cdots (N-q)!}.$$

Proof. We carry out the integrals iteratively choosing polar coordinates. The integral over the first column is computed by

$$\begin{aligned} \int_{\mathbb{C}^N} \delta(\|\mathcal{U}_1\|^2 - 1) d\mathcal{U}_1 &= \int_0^\infty R^{2N-1} dR \int_{S^{2N-1}} d\omega \delta(R^2 - 1) \\ &= \frac{\mu(S^{2N-1})}{2} = \frac{\pi^N}{(N-1)!}. \end{aligned} \quad (4.11)$$

Here we carried out the R -integral using the well-known formula

$$\int f(x) \delta(g(x)) dx = \sum_x \frac{f(x)}{|g'(x)|}, \quad (4.12)$$

where the sum goes over all the zeros of the function g . This identity will also be used frequently in the following computations. Moreover, in the last step in (4.11) we used that the volume of an odd-dimensional unit sphere is given by (see for example [37])

$$\mu(S^{2n-1}) = \frac{2\pi^n}{(n-1)!}$$

for $n \in \mathbb{N}$.

Having carried out the integral over \mathcal{U}_1 , in the next step we can assume by spherical symmetry of the measure that $\mathcal{U}_1 = (1, 0, \dots, 0)$. Hence

$$\begin{aligned} &\int_{\mathbb{C}^N} \delta(\|\mathcal{U}_2\|^2 - 1) \delta(\operatorname{Re}\langle \mathcal{U}_1, \mathcal{U}_2 \rangle) \delta(\operatorname{Im}\langle \mathcal{U}_1, \mathcal{U}_2 \rangle) d\mathcal{U}_2 \\ &= \int_{\mathbb{C}^N} \delta(\|\mathcal{U}_2\|^2 - 1) \delta(\operatorname{Re} \mathcal{U}_2^1) \delta(\operatorname{Im} \mathcal{U}_2^1) d\mathcal{U}_2 = \int_{\mathbb{C}^{N-1}} \delta(\|u\|^2 - 1) du \\ &= \int_0^\infty r^{2N-3} dr \int_{S^{2N-3}} d\omega \delta(r^2 - 1) = \frac{\mu(S^{2N-3})}{2} = \frac{\pi^{N-1}}{(N-2)!}. \end{aligned}$$

Proceeding inductively, we conclude that

$$\int_{\mathbb{C}^{N \times q}} d\mu_{\mathcal{U}|_I} = \frac{\pi^N \pi^{N-1} \cdots \pi^{N+1-q}}{(N-1)! (N-2)! \cdots (N-q)!} = \frac{\pi^{\frac{q(2N+1-q)}{2}}}{(N-1)! \cdots (N-q)!},$$

giving the result. \square

We next compute the integral on the right side of (4.10).

Lemma 4.6.

$$\int_{\mathbb{C}^{N \times q}} f(A) d\mu_{\mathcal{G}|_I} = \frac{\pi^{\frac{q(2N+1-3q)}{2}}}{(N-q-1)!(N-q-2)! \cdots (N-2q)!} \times \int_{\mathcal{A}_q} f(A) (\det(\mathbb{1}_{\mathbb{C}^q} - A^*A))^{N-2q} d\mu_{\mathbb{C}^q \times q}(A).$$

Proof. Given $\mathcal{U} \in \mathrm{U}(N)$ we choose an “adapted” orthonormal basis f_1, \dots, f_{N-q} of $J := I^\perp$ such that the lower left block matrix entry of \mathcal{U} is an upper triangular matrix, i.e.

$$B := \pi_J \mathcal{U}|_I = \begin{pmatrix} R_1 & m_2^1 & m_3^1 & \cdots & m_q^1 \\ 0 & R_2 & m_3^2 & \cdots & m_q^2 \\ 0 & 0 & R_3 & \cdots & m_q^3 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & R_q \\ 0 & 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ 0 & 0 & 0 & \cdots & 0 \end{pmatrix} \quad (4.13)$$

with $R_k \geq 0$ (in other words, B is the matrix R in the QR -decomposition of the lower left block matrix entry of \mathcal{U}). It suffices to consider the case that all the R_k are strictly positive, because the unitary matrices \mathcal{U} for which one of the R_k is zero form a set of measure zero. All the matrix entries of this matrix can be computed uniquely from the matrix A using the fact that the columns of a unitary matrix are orthonormal. For example,

$$R_1 = \sqrt{1 - \|Ae_1\|^2} \\ m_2^1 = -\frac{1}{R_1} \langle Ae_1, Ae_2 \rangle_{\mathbb{C}^q}, \quad R_2 = \sqrt{1 - \|Ae_2\|^2 - |m_2^1|^2},$$

and so on.

Let us verify that this procedure of constructing B from A works if and only if the matrix $\mathbb{1}_{\mathbb{C}^q} - A^*A$ is positive definite. Clearly, starting from a unitary matrix \mathcal{U} , the corresponding matrix $A = \pi_I \mathcal{U}|_I$ has the property

$$\mathbb{1}_{\mathbb{C}^q} = \pi_I \mathcal{U}^* \mathcal{U}|_I = A^*A + B^*B, \quad (4.14)$$

showing that $A^*A \leq \mathbb{1}_{\mathbb{C}^q}$. Moreover, the condition that all R_k are strictly positive is equivalent to B^*B being positive definite. Conversely, assume that $\mathbb{1}_{\mathbb{C}^q} - A^*A > 0$. Then its square root $\sqrt{\mathbb{1}_{\mathbb{C}^q} - A^*A}$ is again positive definite. We make the ansatz

$$B = V \sqrt{\mathbb{1}_{\mathbb{C}^q} - A^*A}$$

with a unitary matrix V which is chosen such that B is an upper triangular matrix. Choosing

$$\mathcal{U}|_I = \begin{pmatrix} A \\ B \end{pmatrix}, \quad (4.15)$$

the columns of this matrix are orthonormal. Extending this orthonormal system to an orthonormal basis of \mathcal{H} , one can extend (4.15) to a unitary operator \mathcal{U} simply by choosing the additional columns as the additional basis vectors.

It remains to compute the integral. We again work with the block representation (4.15) and integrate out the columns of B after each other. In the first column, we obtain

$$\begin{aligned} \int_{\mathbb{C}^{N-q}} \delta(\|\mathcal{U}_1\|^2 - 1) dB_1 &= \int_{\mathbb{C}^{N-q}} \delta(\|A_1\|^2 + \|B_1\|^2 - 1) dB_1 \\ &= \int_0^\infty r^{2N-2q-1} dr \int_{S^{2N-2q-1}} d\omega \delta(\|A_1\|^2 + r^2 - 1) \\ &= \frac{\mu(S^{2N-2q-1})}{2} R_1^{2N-2q-2} = \frac{\pi^{N-q}}{(N-q-1)!} R_1^{2N-2q-2}, \end{aligned}$$

where $A_1 := Ae_1$ and $B_1 := Be_1$ are the first column vectors of the corresponding matrices, and R_1 coincides with the corresponding matrix entry in (4.13). In the following computation we can assume by symmetry that $B_1 = (R_1, 0, \dots, 0)$ (in other words, we work in a basis where the first column of B has the same form as in (4.13)). We then obtain in the second column

$$\begin{aligned} \int_{\mathbb{C}^{N-q}} \delta(\|\mathcal{U}_2\|^2 - 1) \delta(\operatorname{Re}\langle \mathcal{U}_1, \mathcal{U}_2 \rangle) \delta(\operatorname{Im}\langle \mathcal{U}_1, \mathcal{U}_2 \rangle) dB_2 \\ &= \int_{\mathbb{C}^{N-q}} \delta(\|A_2\|^2 + \|B_2\|^2 - 1) \\ &\quad \times \delta(\operatorname{Re}\langle A_1, A_2 \rangle + R_1 \operatorname{Re} B_2^1) \delta(\operatorname{Im}\langle A_1, A_2 \rangle + R_1 \operatorname{Im} B_2^1) dB_2 \\ &= \frac{1}{R_1^2} \int_{\mathbb{C}^{N-q-1}} \delta(\|A_2\|^2 + |B_2^1|^2 + \|u\|^2 - 1) du \\ &= \frac{1}{R_1^2} \int_0^\infty r^{2N-2q-3} dr \int_{S^{2N-2q-3}} d\omega \delta(r^2 - R_2^2) \\ &= \frac{\mu(S^{2N-2q-3})}{2R_1^2} R_2^{2N-2q-4} = \frac{\pi^{N-q-1}}{(N-q-2)! R_1^2} R_2^{2N-2q-4}, \end{aligned}$$

where we decomposed B_2 as $B_2 = (B_2^1, u)$ and carried out the integrals over the real and imaginary parts of B_2^1 with the help of (4.12) (the parameter R_2 is again the corresponding matrix entry in (4.13)). Proceeding inductively, we conclude that

$$\begin{aligned} \int_{\mathcal{U}|_I} f(A) d\mu_{\mathcal{U}|_I} &= \int_{A_q} d\mu_{\mathbb{C}^q \times q}(A) f(A) \frac{1}{R_1^{2N-4q} \dots R_q^{2N-4q}} \\ &\quad \times \frac{\pi^{N-q} \pi^{N-q-1} \dots \pi^{N+1-2q}}{(N-q-1)! (N-q-2)! \dots (N-2q)!}. \end{aligned}$$

Now, we apply (4.14) to obtain

$$R_1^2 \dots R_q^2 = |\det B|^2 = \det(B^* B) = \det(\mathbb{1} - A^* A).$$

We finally use that

$$\pi^{N-q} \pi^{N-q-1} \dots \pi^{N+1-2q} = \pi^{\frac{1}{2} q(2N+1-3q)}.$$

This gives the result. \square

Combining the formulas of Lemmas 4.5 and 4.6, one immediately gets the statement of Theorem 4.2.

4.3. A Few Simple Examples. In this section we illustrate and verify the result of Corollary 4.3 in the cases $q = 1$ and $q = 2$.

Example 4.7. (q=1) In the case $q = 1$, Corollary 4.3 states that

$$\int_{\mathcal{A}_1} (\det(\mathbf{1} - A^* A))^{N-2} d\mu_{\mathbb{C}^{q \times q}}(A) = \frac{\pi}{N-1}.$$

On the other hand, the integral on the left can be computed directly,

$$\begin{aligned} & \int_{\mathcal{A}_1} (\det(\mathbf{1} - A^* A))^{N-2} d\mu_{\mathbb{C}}(A) \\ &= \int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} dy \Theta(1 - x^2 - y^2) (1 - x^2 - y^2)^{N-2} = 2\pi \int_0^1 r dr (1 - r^2)^{N-2} \\ &= \pi \int_0^1 d(r^2) (1 - r^2)^{N-2} = -\pi \frac{(1 - r^2)^{N-1}}{N-1} \Big|_{r^2=0}^{r^2=1} = \frac{\pi}{N-1}, \end{aligned}$$

as desired. ◇

Example 4.8. (q=2) In the case $q = 2$, we write A as

$$A = \alpha_0 \mathbf{1} + \vec{\alpha} \vec{\sigma} + i\beta_0 \mathbf{1} + i\vec{\beta} \vec{\sigma}$$

with $\alpha = (\alpha_0, \vec{\alpha}), \beta = (\beta_0, \vec{\beta}) \in \mathbb{R}^4$. Then

$$\begin{aligned} A^* &= \alpha_0 \mathbf{1} + \vec{\alpha} \vec{\sigma} - i\beta_0 \mathbf{1} - i\vec{\beta} \vec{\sigma} \\ A^* A &= (\alpha_0^2 + \beta_0^2 + |\vec{\alpha}|^2 + |\vec{\beta}|^2) \mathbf{1} + 2\alpha_0 \vec{\alpha} \vec{\sigma} + 2\beta_0 \vec{\beta} \vec{\sigma} - 2(\vec{\alpha} \wedge \vec{\beta}) \cdot \vec{\sigma}. \end{aligned}$$

We want to compute for $p = 0, 1, 2, \dots$ the integral

$$\int_{\mathcal{A}_2} (\det(\mathbf{1} - A^* A))^p d\mu_{\mathbb{C}^{2 \times 2}}(A).$$

To this end, we first transform to the integration variables $\alpha, \beta \in \mathbb{R}^4$,

$$\int_{\mathcal{A}_2} (\det(\mathbf{1} - A^* A))^p d\mu_{\mathbb{C}^{2 \times 2}}(A) = 16 \int_{\mathbb{R}^4} d^4 \alpha \int_{\mathbb{R}^4} d^4 \beta (\det(\mathbf{1} - A^* A))^p.$$

Introducing polar coordinates in the plane,

$$\alpha_0 = r \cos \phi, \quad \beta_0 = r \sin \phi,$$

we obtain

$$\int_{\mathcal{A}_2} (\det(\mathbf{1} - A^* A))^p d\mu_{\mathbb{C}^{2 \times 2}}(A) = 32\pi \int_0^\infty r dr \int_{\mathbb{R}^3} d^3 \alpha \int_{\mathbb{R}^3} d^3 \beta \chi_{\mathcal{A}_2}(A) (\det(\mathbf{1} - A^* A))^p.$$

In view of rotational symmetry in the angle ϕ , we may assume that $\alpha_0 = r$ and $\beta_0 = 0$. Moreover, we introduce spherical coordinates with

$$R := |\vec{\alpha}|, \quad \hat{R} := |\vec{\beta}|,$$

and denote the angle between $\vec{\alpha}$ and $\vec{\beta}$ by ϑ . We thus obtain

$$\begin{aligned} & \int_{\mathcal{A}_2} (\det(\mathbf{1} - A^* A))^p d\mu_{\mathbb{C}^{2 \times 2}}(A) \\ &= 32 \pi \int_0^\infty r dr \, 4\pi \int_0^\infty R^2 dR \, 2\pi \int_0^\infty \hat{R}^2 d\hat{R} \int_{-1}^1 d \cos \vartheta \, \chi_{\mathcal{A}_2}(A) (\det(\mathbf{1} - A^* A))^p \\ &= 256 \pi^3 \int_0^\infty dr \int_0^\infty dR \int_0^\infty d\hat{R} \int_{-1}^1 d \cos \vartheta \, \chi_{\mathcal{A}_2}(A) r R^2 \hat{R}^2 (\det(\mathbf{1} - A^* A))^p, \end{aligned}$$

where, due to the symmetries, the matrix A can be arranged to be of the form

$$A = \begin{pmatrix} r + R + ir \cos \vartheta & ir \sin \vartheta \\ ir \sin \vartheta & r + R - ir \cos \vartheta \end{pmatrix}.$$

A direct computation yields for the eigenvalues λ_\pm of the matrix $A^* A$

$$\lambda_\pm = R^2 + r^2 + \hat{R}^2 \pm 2R \sqrt{r^2 + \hat{R}^2 \sin^2 \vartheta}$$

and thus

$$\det(\mathbf{1} - A^* A) = (1 - \lambda_+)(1 - \lambda_-) = (1 - R^2 + r^2 + \hat{R}^2)^2 - 4R^2(r^2 + \hat{R}^2 \sin^2 \vartheta).$$

Next, we need to parametrize the set \mathcal{A}_2 . To this end, we need to ensure that the largest eigenvalue of $A^* A$ is at most one, i.e.

$$f(r, R, \hat{R}, \vartheta) := R^2 + r^2 + \hat{R}^2 + 2R \sqrt{r^2 + \hat{R}^2 \sin^2 \vartheta} \leq 1.$$

Obviously, the function f is monotone increasing in R . Its minimal value is attained at $R = 0$,

$$f(r, 0, \hat{R}, \vartheta) = r^2 + \hat{R}^2.$$

Therefore, we may parametrize the integration domain \mathcal{A}_2 as follows,

$$\begin{aligned} & \int_0^\infty dr \int_0^\infty dR \int_0^\infty d\hat{R} \int_{-1}^1 d \cos \vartheta \, \chi_{\mathcal{A}_2}(A) \cdots \\ &= \int_0^1 dr \int_0^{\sqrt{1-r^2}} d\hat{R} \int_{-1}^1 d \cos \vartheta \int_0^{R_{\max}} dR \cdots, \end{aligned}$$

where R_{\max} is determined by the equation $f(r, R_{\max}, \hat{R}, \vartheta) = 1$ to be

$$R_{\max} = \sqrt{1 - \hat{R}^2 \cos^2 \vartheta} - \sqrt{r^2 + \hat{R}^2 \sin^2 \vartheta}.$$

A straightforward computation (which we carried with the help of MAPLE) gives

$$\int_{\mathcal{A}_2} (\det(\mathbf{1} - A^* A))^p d\mu_{\mathbb{C}^{2 \times 2}}(A) = \pi^4 \frac{(p+1)! p!}{(p+3)! (p+2)!}$$

for $p = 0, \dots, 3$. This is in agreement with the general formula from Corollary 4.3 if we choose $N = 4, \dots, 7$. \diamond

We conclude this section with another example, where the function f in the integrand is chosen as an exponential. This example serves several purposes: First, it illustrates the Gaussian asymptotics of Lemma 4.1, which will be developed further in the following section (Section 4.4). Second, it shows that by adding a factor N into the exponent, one can get into a nonlinear regime where the Gaussian asymptotics no longer applies, but where the group integral can be computed instead using saddle

point methods. It is a simple example of N -dependent integrands as will be analyzed in Section 4.5. Moreover, it is a preparation for the saddle point methods to be used in greater generality in Section 5.

Example 4.9. (an exponential integrand) Let $I \subset \mathcal{H}$ be a one-dimensional subspace of \mathcal{H} . We denote the orthogonal projection to I by π_I . We choose $\mathcal{G} = \mathrm{U}(\mathcal{H}) \simeq \mathrm{U}(N)$ as the group of all unitary transformations on \mathcal{H} and consider the integral

$$\mathcal{J} := \int_{\mathcal{G}} \exp \left(\beta \operatorname{tr}_{\mathcal{H}} (\pi_I \mathcal{U} \pi_I \mathcal{U}^{-1}) \right) d\mu_{\mathcal{G}}(\mathcal{U}), \quad (4.16)$$

where β is a real parameter. This is indeed a special case of the Harish-Chandra integral [40]. In the present one-dimensional situation, the integral can be computed explicitly as follows. Applying Theorem 4.2, we obtain

$$\mathcal{J} = \frac{1}{\pi} \frac{(N-1)!}{(N-2)!} \int_{\mathcal{A}_1} e^{\beta |A|^2} (\det(\mathbf{1} - A^* A))^{N-2} d\mu_{\mathbb{C}}(A)$$

Using a polar decomposition $A = r e^{i\varphi}$ and integrating out the angular dependence, we obtain $d\mu_{\mathbb{C}}(A) = r d\varphi dr$ and thus

$$\mathcal{J} = 2(N-1) \int_0^1 e^{\beta r^2} (1-r^2)^{N-2} r dr. \quad (4.17)$$

Introducing the integration variable $u := r^2$, we obtain

$$\mathcal{J} = (N-1) \int_0^1 e^{\beta u} (1-u)^{N-2} du.$$

Asymptotically for large N , the weight function $(1-u)^{N-2}$ is very small except near $u = 0$. Therefore, we may expand the exponential in a power series and carry out the integral term by term,

$$\begin{aligned} \mathcal{J} &= (N-1) \sum_{p=0}^{\infty} \int_0^1 \frac{(\beta u)^p}{p!} (1-u)^{N-2} du = (N-1) \sum_{p=0}^{\infty} \beta^p \frac{(N-2)!}{(N+p-1)!} \\ &= \sum_{p=0}^{\infty} \beta^p \frac{(N-1)!}{(N+p-1)!} = \sum_{p=0}^{\infty} \frac{\beta^p}{N^p} (1 + \mathcal{O}(N^{-1})). \end{aligned}$$

This formula is obtained alternatively if we expand the exponential in (4.16) in a power series and compute term by term with the help of Lemma 4.1. In particular, we conclude that \mathcal{J} tends to one as $N \rightarrow \infty$.

In order to obtain a nontrivial limit as $N \rightarrow \infty$, one must increase β with N . The right scaling is to choose β as a linear function in N . In order to see how this comes about, we now insert a factor N into the exponent in (4.16),

$$\begin{aligned} \int_{\mathcal{G}} \exp \left(\beta N \operatorname{tr}_{\mathcal{H}} (\pi_I \mathcal{U} \pi_I \mathcal{U}^{-1}) \right) d\mu_{\mathcal{G}}(\mathcal{U}) &= (N-1) \int_0^1 e^{\beta N u} (1-u)^{N-2} du \\ &= (N-1) \int_0^1 e^{N(\beta u + \log(1-u))} (1-u)^{-2} du. \end{aligned} \quad (4.18)$$

We point out that, due to this additional factor N , it is not clear how the large- N -asymptotics can be obtained from the Harish-Chandra integral. But we can use that,

for large N , the integrand is peaked near the maximum of the function

$$\beta u + \log(1 - u) .$$

If $\beta \leq 1$, this function has its maximum at $u_0 = 0$, whereas in the case $\beta > 1$ it has a maximum at some $u_0 > 0$. In both cases, the function has a unique maximum, which means that the main contribution to the integral (4.18) comes from a neighborhood of one point u_0 .

We finally compute the integral (4.18) asymptotically as $N \rightarrow \infty$, for brevity only in the case $\beta < 1$. In this case, the main contribution to the integral comes from the region near $u = 0$. Expanding the logarithm as

$$\log(1 - u) = -u + \mathcal{O}(u^2) ,$$

we obtain

$$\begin{aligned} (N-1) \int_0^1 e^{-N(1-\beta)u} (1-u)^{-2} du \left(1 + \mathcal{O}\left(\frac{1}{N}\right)\right) \\ = (N-1) \int_0^\infty e^{-N(1-\beta)u} du \left(1 + \mathcal{O}\left(\frac{1}{N}\right)\right) \end{aligned} \quad (4.19)$$

$$= (N-1) \frac{1}{N(1-\beta)} \left(1 + \mathcal{O}\left(\frac{1}{N}\right)\right) = \frac{1}{1-\beta} + \mathcal{O}\left(\frac{1}{N}\right) , \quad (4.20)$$

valid if $\beta < 1$. More systematically, integrals of this type can be computed asymptotically for large N by applying the saddle point method at the maximum of the integrand (to this end, it is preferable to work as in (4.17) with the coordinate r , in which case the exponential in (4.19) becomes a Gaussian). The saddle point method will be developed systematically in Section 5. \diamond

4.4. The Gaussian Asymptotics. In the next proposition it is shown that for large N , the integral over the unitary group goes over to a Gaussian integral. Choosing orthonormal bases, we use the identification $I \simeq \mathbb{C}^q$.

Proposition 4.10. *Let $I \subset \mathcal{H}$ with $\dim I = q$ and $\dim \mathcal{H} = N$. Assume that $f \in C^\infty(\mathbb{C}^{q \times q}, \mathbb{C})$ is homogeneous of degree $d \in \mathbb{N}_0$. Then, asymptotically for large N ,*

$$\oint_{\mathcal{G}} f(A) d\mu_{\mathcal{G}}(\mathcal{U}) = \frac{1}{\pi^{(q^2)}} \frac{1}{N^{\frac{d}{2}}} \int_{\mathbb{C}^{q \times q}} f(\tilde{A}) e^{-\text{Tr}(\tilde{A}^* \tilde{A})} d\mu_{\mathbb{C}^{q \times q}}(\tilde{A}) + \mathcal{O}\left(\frac{1}{N^{\frac{d}{2}+1}}\right) . \quad (4.21)$$

Related results on the Gaussian asymptotics can be found in the literature; see for example [38, Section II]. We remark that, if f is a homogeneous polynomial, the Gaussian can be computed in a straightforward way. Since the covariance is the identity matrix, we get back the formula in Lemma 4.1 (the condition that f is a function of A only is not a restriction because in order to deduce the formula (4.5) from (4.21) one simply chooses I as the subspace spanned by the basis vectors corresponding to all the indices in (4.5)).

We also remark for clarity that the formula (4.21) applies more generally if f is real analytic or smooth. This can be proved by approximation. However, in these cases it is more subtle to specify the error term. Working with homogeneous functions will be sufficient for the purpose of this paper.

Proof of Proposition 4.10. According to Corollary 4.4,

$$\oint_{\mathcal{G}} f(A) d\mu_{\mathcal{G}}(\mathcal{U}) = \left(\frac{N}{\pi}\right)^{(q^2)} \int_{\mathcal{A}_q} f(A) (\det(\mathbf{1} - A^* A))^{N-2q} d\mu_{\mathbb{C}^{q \times q}}(A) \left(1 + \mathcal{O}(N^{-1})\right) .$$

We transform to the new integration variable $\tilde{A} = \sqrt{N}A$,

$$\begin{aligned} & \int_{\mathcal{G}} f(A) d\mu_{\mathcal{G}}(\mathcal{U}) \\ &= \frac{1}{\pi^{(q^2)}} \int_{\sqrt{N}\mathcal{A}_q} f\left(\frac{\tilde{A}}{\sqrt{N}}\right) \left(\det\left(\mathbf{1} - \frac{\tilde{A}^* \tilde{A}}{N}\right)\right)^{N-2q} d\mu_{\mathbb{C}^{q \times q}}(\tilde{A}) \left(1 + \mathcal{O}(N^{-1})\right). \end{aligned}$$

We now expand the determinant using the well-known formula

$$\frac{d}{d\tau} \det B(\tau) = \det B(\tau) \operatorname{tr} (B(\tau)^{-1} \dot{B}(\tau)) \quad (4.22)$$

(this formula can be derived for example by differentiating the Laplace expansion).

$$\det\left(\mathbf{1} - \frac{\tilde{A}^* \tilde{A}}{N}\right) = 1 - \frac{1}{N} \operatorname{Tr}(\tilde{A}^* \tilde{A}) + \mathcal{O}(N^{-2}),$$

we obtain

$$\begin{aligned} & \int_{\mathcal{G}} f(A) d\mu_{\mathcal{G}}(\mathcal{U}) \\ &= \frac{1}{\pi^{(q^2)}} \int_{\sqrt{N}\mathcal{A}_q} f\left(\frac{\tilde{A}}{\sqrt{N}}\right) \left(1 - \frac{1}{N} \operatorname{Tr}(\tilde{A}^* \tilde{A})\right)^{N-2q} d\mu_{\mathbb{C}^{q \times q}}(\tilde{A}) \left(1 + \mathcal{O}(N^{-1})\right), \end{aligned}$$

where, from now on, the error term denotes the leading order as specified in the statement of the proposition above. Applying Gauss' formula

$$\left(1 - \frac{x}{N}\right)^N = e^{-x} \left(1 + \mathcal{O}\left(\frac{1}{N}\right)\right) \quad (4.23)$$

gives

$$\begin{aligned} & \int_{\mathcal{G}} f(A) d\mu_{\mathcal{G}}(\mathcal{U}) \\ &= \frac{1}{\pi^{(q^2)}} \int_{\sqrt{N}\mathcal{A}_q} f\left(\frac{\tilde{A}}{\sqrt{N}}\right) e^{-\operatorname{Tr}(\tilde{A}^* \tilde{A})} d\mu_{\mathbb{C}^{q \times q}}(\tilde{A}) \left(1 + \mathcal{O}(N^{-1})\right). \end{aligned}$$

Using that f is homogeneous of degree d gives the result. \square

4.5. Treating Integrands which Depend on the Dimension. If the integrand f depends on N , the behavior of the integral for large N may change completely. In this case, the asymptotics will depend sensitively on the detailed form of the N -dependence. But we can treat the case when the integrand is a product of two functions, one of which is independent of N . More precisely, we choose two orthogonal subspaces I and J of \mathcal{H} ,

$$I \perp J \quad \text{with} \quad \dim I = q, \dim J = p \quad \text{and} \quad 2(p+q) \leq N. \quad (4.24)$$

Choosing orthonormal bases, we use the identifications

$$I \simeq \mathbb{C}^q \quad \text{and} \quad J \simeq \mathbb{C}^p.$$

Moreover, we use the notation

$$A := \pi_I \mathcal{U}|_I \quad \text{and} \quad D := \pi_J \mathcal{U}|_J. \quad (4.25)$$

Here and in what follows, we always compute Gaussian integrals with the well-known formula

$$\int_{-\infty}^{\infty} e^{-\alpha x^2} dx = \sqrt{\frac{\pi}{\alpha}} \quad (\text{for } \alpha > 0) \quad (4.26)$$

Proposition 4.11. *Given $p, q \in \mathbb{N}$, we assume that $g \in C^\infty(\mathbb{C}^{p \times p}, \mathbb{C})$ is smooth and homogeneous of degree $d \in \mathbb{N}_0$ and that $f \in C^\infty(\mathbb{R} \times A_q, \mathbb{C})$ is smooth. Then, asymptotically for large N ,*

$$\begin{aligned} \oint_{\mathcal{G}} f(N, A) g(D) d\mu_{\mathcal{G}}(\mathcal{U}) &= \left(\frac{N}{\pi}\right)^{(q^2)} \int_{A_q} f(N, A) (\det(\mathbf{1} - A^* A))^{N-2q} d\mu_{\mathbb{C}^{q \times q}}(A) \\ &\times \left\{ \frac{1}{\pi(p^2)} \frac{1}{N^{\frac{d}{2}}} \int_{\mathbb{C}^{p \times p}} g(\tilde{D}) e^{-\text{Tr}(\tilde{D}^* \tilde{D})} d\mu_{\mathbb{C}^{p \times p}}(\tilde{D}) + \mathcal{O}\left(\frac{1}{N^{\frac{d}{2}+1}}\right) \right\} \left(1 + \mathcal{O}\left(\frac{1}{N}\right)\right). \end{aligned}$$

In words, this formula shows that the integral factorizes into an integral over A (which is similar to the general formula of Corollary 4.4) and an integral over D (where the Gaussian asymptotics of Proposition 4.10 applies). In order to verify the prefactors, it is instructive to consider the case that g is the constant function one (being homogeneous of degree $d = 0$). Then, carrying out the $2q^2$ -dimensional Gaussian integral with the help of (4.26) gives us back the formula in Corollary 4.4.

Proof of Proposition 4.11. We choose the subspace $\hat{I} = I \oplus J$. For the matrix $\hat{A} := \pi_{\hat{I}} \mathcal{U}|_{\hat{I}}$ we use the block matrix notation

$$\hat{A} = \begin{pmatrix} A & B \\ C & D \end{pmatrix}. \quad (4.27)$$

We apply Corollary 4.4 for the subspace \hat{I} and the integrand $\hat{f}(\hat{A}) = f(N, A) g(D)$,

$$\begin{aligned} \oint_{\mathcal{G}} f(N, A) g(D) d\mu_{\mathcal{G}}(\mathcal{U}) &= \left(\frac{N}{\pi}\right)^{(p+q)^2} \\ &\times \int f(N, A) g(D) (\det(\mathbf{1} - \hat{A}^* \hat{A}))^{N-2p-2q} d\mu_{\mathbb{C}^{p+q \times p+q}}(\hat{A}) \left(1 + \mathcal{O}\left(\frac{1}{N}\right)\right). \end{aligned} \quad (4.28)$$

For ease in notation, we write the integration measure as

$$d\mu_{\mathbb{C}^{p+q \times p+q}}(\hat{A}) = dA dB dC dD.$$

Our next task is to compute the factor $\det(\mathbf{1} - \hat{A}^* \hat{A})$. Again in block matrix notation,

$$\begin{aligned} \hat{A}^* &= \begin{pmatrix} A^* & C^* \\ B^* & D^* \end{pmatrix} \\ \hat{A}^* \hat{A} &= \begin{pmatrix} A^* A + C^* C & A^* B + C^* D \\ B^* A + D^* C & B^* B + D^* D \end{pmatrix} \\ \mathbf{1} - \hat{A}^* \hat{A} &= \begin{pmatrix} \mathbf{1} - A^* A - C^* C & -A^* B - C^* D \\ -B^* A - D^* C & \mathbf{1} - B^* B - D^* D \end{pmatrix} \\ &= \begin{pmatrix} \mathbf{1} - A^* A & 0 \\ 0 & \mathbf{1} \end{pmatrix} - \begin{pmatrix} C^* C & A^* B + C^* D \\ B^* A + D^* C & B^* B + D^* D \end{pmatrix} \\ &= \begin{pmatrix} \mathbf{1} - A^* A & 0 \\ 0 & \mathbf{1} \end{pmatrix} (\mathbf{1} - X) \end{aligned}$$

with the matrix X given by

$$X := \begin{pmatrix} (\mathbb{1} - A^*A)^{-1} C^*C & (\mathbb{1} - A^*A)^{-1} (A^*B + C^*D) \\ B^*A + D^*C & B^*B + D^*D \end{pmatrix}.$$

Hence we can factorize the determinant as

$$\det(\mathbb{1} - \hat{A}^* \hat{A}) = \det(\mathbb{1} - A^*A) \det(\mathbb{1} - X). \quad (4.29)$$

Similar as in the proof of Proposition 4.10, the integrals over B , C and D can be computed as Gaussian integrals,

$$\begin{aligned} & \int g(D) \det(\mathbb{1} - X)^{N-2p-2q} dB dC dD = \left\{ B, C, D = \frac{1}{\sqrt{N}} (\tilde{B}, \tilde{C}, \tilde{D}) \right\} \\ &= \frac{1}{N^{2pq+p^2}} \int g\left(\frac{\tilde{D}}{\sqrt{N}}\right) \det \left\{ \mathbb{1} - \frac{1}{N} \begin{pmatrix} (\mathbb{1} - A^*A)^{-1} \tilde{C}^* \tilde{C} & (\mathbb{1} - A^*A)^{-1} \tilde{C}^* \tilde{D} \\ \tilde{D}^* \tilde{C} & \tilde{B}^* \tilde{B} + \tilde{D}^* \tilde{D} \end{pmatrix} \right. \\ & \quad \left. - \frac{1}{\sqrt{N}} \begin{pmatrix} 0 & (\mathbb{1} - A^*A)^{-1} A^* \tilde{B} \\ \tilde{B}^* A & 0 \end{pmatrix} \right\}^{N-2p-2q} d\tilde{B} d\tilde{C} d\tilde{D}. \end{aligned}$$

Again using (4.22), we now expand the determinant to second order,

$$\begin{aligned} & \frac{d}{d\tau} \det(\mathbb{1} - \tau Y) = -\det(\mathbb{1} - \tau Y) \operatorname{tr}((\mathbb{1} - \tau Y)^{-1} Y) \\ & \frac{d}{d\tau} \det(\mathbb{1} - \tau Y) \Big|_{\tau=0} = -\operatorname{tr}(Y) \\ & \frac{d^2}{d\tau^2} \det(\mathbb{1} - \tau Y) \Big|_{\tau=0} = \operatorname{tr}(Y)^2 - \operatorname{tr}(Y^2) \\ & \implies \det(1 - \tau Y) = 1 - \tau \operatorname{tr}(Y) + \frac{\tau^2}{2} \left(\operatorname{tr}(Y)^2 - \operatorname{tr}(Y^2) \right) + \mathcal{O}(\tau^3). \end{aligned}$$

We thus obtain

$$\begin{aligned} & \int g(D) \det(\mathbb{1} - X)^{N-2p-2q} dB dC dD \\ &= \frac{1}{N^{2pq+p^2}} \left(1 + \mathcal{O}\left(\frac{1}{N}\right) \right) \int g\left(\frac{\tilde{D}}{\sqrt{N}}\right) \left\{ 1 - \frac{1}{N} \operatorname{Tr} \begin{pmatrix} (\mathbb{1} - A^*A)^{-1} \tilde{C}^* \tilde{C} & 0 \\ 0 & \tilde{B}^* \tilde{B} + \tilde{D}^* \tilde{D} \end{pmatrix} \right. \\ & \quad \left. - \frac{1}{2N} \operatorname{Tr} \begin{pmatrix} (\mathbb{1} - A^*A)^{-1} A^* \tilde{B} \tilde{B}^* A & 0 \\ 0 & \tilde{B}^* A (\mathbb{1} - A^*A)^{-1} A^* \tilde{B} \end{pmatrix} \right\}^{N-2p-2q} d\tilde{B} d\tilde{C} d\tilde{D} \\ &= \frac{1}{N^{2pq+p^2}} \left(1 + \mathcal{O}\left(\frac{1}{N}\right) \right) \int d\tilde{B} d\tilde{C} d\tilde{D} g\left(\frac{\tilde{D}}{\sqrt{N}}\right) \\ & \quad \times \exp \left(-\operatorname{Tr} \left((\mathbb{1} - A^*A)^{-1} \tilde{C}^* \tilde{C} + \tilde{B}^* \tilde{B} + \tilde{D}^* \tilde{D} + (\mathbb{1} - A^*A)^{-1} A^* \tilde{B} \tilde{B}^* A \right) \right), \end{aligned}$$

where in the last step we again used (4.23). The trace of the last summand can be simplified as follows,

$$\begin{aligned} & \operatorname{Tr} \left((\mathbb{1} - A^*A)^{-1} A^* \tilde{B} \tilde{B}^* A \right) = \operatorname{Tr} \left(A (\mathbb{1} - A^*A)^{-1} A^* \tilde{B} \tilde{B}^* \right) \\ &= \operatorname{Tr} \left(A A^* (\mathbb{1} - A A^*)^{-1} \tilde{B} \tilde{B}^* \right) = -\operatorname{Tr}(\tilde{B} \tilde{B}^*) + \operatorname{Tr} \left((\mathbb{1} - A A^*)^{-1} \tilde{B} \tilde{B}^* \right). \end{aligned}$$

We conclude that

$$\begin{aligned} & \int g(D) \det(\mathbf{1} - X)^{N-2p-2q} dB dC dD \\ &= \frac{1}{N^{2pq+p^2}} \left(1 + \mathcal{O}\left(\frac{1}{N}\right)\right) \int d\tilde{B} d\tilde{C} d\tilde{D} g\left(\frac{\tilde{D}}{\sqrt{N}}\right) \\ & \quad \times \exp\left(-\operatorname{Tr}\left((\mathbf{1} - A^*A)^{-1} \tilde{C}^* \tilde{C} + (\mathbf{1} - AA^*)^{-1} \tilde{B} \tilde{B}^* + \tilde{D}^* \tilde{D}\right)\right). \end{aligned}$$

Now we can carry out the integrals over \tilde{B} and \tilde{C} , which are both Gaussian. We begin with the integral of \tilde{C} . Using (4.26) we obtain

$$\begin{aligned} & \int \exp\left(-\operatorname{Tr}\left((\mathbf{1} - A^*A)^{-1} \tilde{C}^* \tilde{C}\right)\right) d\tilde{C} \\ &= \frac{\pi^{pq}}{\det((\mathbf{1} - A^*A)^{-1})^p} = \pi^{pq} (\det(\mathbf{1} - A^*A))^p. \end{aligned}$$

The power $-p$ of the determinant can be derived as follows: The Gaussian integral is of dimension $2pq$. If we diagonalize the $q \times q$ -matrix $(\mathbf{1} - A^*A)^{-1}$, every eigenspace gives rise to a Gaussian integral of dimension $2pq/q = 2p$. Therefore, Gaussian integration yields the corresponding eigenvalue to the power $-p$. In total, we get the factor $\det(\mathbf{1} - A^*A)^{-p}$.

The integral over \tilde{B} is computed similarly. We conclude that

$$\begin{aligned} & \int g(D) \det(\mathbf{1} - X)^{N-2p-2q} dB dC dD \\ &= \frac{\pi^{2pq}}{N^{2pq+p^2}} (\det(\mathbf{1} - A^*A))^{2p} \int g\left(\frac{\tilde{D}}{\sqrt{N}}\right) \exp^{-\operatorname{Tr}(\tilde{D}^* \tilde{D})} d\tilde{D} \left(1 + \mathcal{O}\left(\frac{1}{N}\right)\right). \end{aligned}$$

After applying (4.29) in (4.28), we can use the last formula to obtain the result. \square

4.6. Asymptotic Decoupling on Orthogonal Subspaces. In this section we again consider the situation of two orthogonal subspaces (4.24) and two function $f(A)$ and $g(B)$ depending on the sub-matrices in (4.25). But now both functions may depend on N . Clearly, we no longer get the Gaussian asymptotics. But the resulting formula makes it possible to analyze whether and in which sense the integral factorizes into separate integrals over A and B . We first state our result and briefly discuss it afterward.

Proposition 4.12. *Given $p, q \in \mathbb{N}$, we assume that $g \in C^\infty(\mathbb{R} \times \mathcal{A}_p, \mathbb{C})$ and $f \in C^\infty(\mathbb{R} \times \mathcal{A}_q, \mathbb{C})$ are smooth. Then, asymptotically for large N ,*

$$\begin{aligned} & \int_{\mathcal{G}} f(N, A) g(N, D) d\mu_{\mathcal{G}}(\mathcal{U}) = \left(\frac{N}{\pi}\right)^{(p^2+q^2)} \\ & \quad \times \int_{\mathcal{A}_q} d\mu_{\mathbb{C}^{q \times q}}(A) \int_{\mathcal{A}_p} d\mu_{\mathbb{C}^{p \times p}}(D) f(N, A) g(N, D) \\ & \quad \times \frac{(\det(\mathbf{1} - A^*A))^{N-2q} (\det(\mathbf{1} - D^*D))^{N-2p}}{\det(\mathbf{1} - A^*A \otimes D^*D)} \left(1 + \mathcal{O}\left(\frac{1}{N}\right)\right). \end{aligned} \quad (4.30)$$

Here the tensor product is a $(pq \times pq)$ -matrix with entries

$$(A^*A \otimes D^*D)_{jl}^{ik} = (A^*A)_j^i (D^*D)_l^k.$$

These matrix entries are very small, provided that either A or D is small. In these cases, we know that

$$\det(\mathbb{1} - A^*A \otimes D^*D) \approx 1, \quad (4.31)$$

and the integral factorizes. In particular, this will be the case if g is homogeneous of degree d , and applying Proposition 4.10, we get back the formula of Proposition 4.11. The main improvement of Proposition 4.12 is that, by estimating the determinant in (4.31) from above, one can quantify the error of the factorization.

Proof of Proposition 4.12. We begin similar as in the proof of Proposition 4.11. Using again the notation (4.27), we obtain

$$\begin{aligned} \hat{A}^* \hat{A} &= \begin{pmatrix} A^*A + C^*C & A^*B + C^*D \\ B^*A + D^*C & B^*B + D^*D \end{pmatrix} \\ \mathbb{1} - \hat{A}^* \hat{A} &= \begin{pmatrix} \mathbb{1} - A^*A - C^*C & -A^*B - C^*D \\ -B^*A - D^*C & \mathbb{1} - B^*B - D^*D \end{pmatrix} \\ &= \begin{pmatrix} \mathbb{1} - A^*A & 0 \\ 0 & \mathbb{1} - D^*D \end{pmatrix} - \begin{pmatrix} C^*C & A^*B + C^*D \\ B^*A + D^*C & B^*B \end{pmatrix} \\ &= \begin{pmatrix} \mathbb{1} - A^*A & 0 \\ 0 & \mathbb{1} - D^*D \end{pmatrix} (\mathbb{1} - X) \end{aligned}$$

where now the matrix X is given by

$$X := \begin{pmatrix} (\mathbb{1} - A^*A)^{-1} C^*C & (\mathbb{1} - A^*A)^{-1} (A^*B + C^*D) \\ (\mathbb{1} - D^*D)^{-1} (B^*A + D^*C) & (\mathbb{1} - D^*D)^{-1} B^*B \end{pmatrix}.$$

Hence we can factorize the determinant as

$$\det(\mathbb{1} - \hat{A}^* \hat{A}) = \det(\mathbb{1} - A^*A) \det(\mathbb{1} - D^*D) \det(\mathbb{1} - X).$$

Our task is to carry out the integrals over B and C . Compared to the proof of Proposition 4.11, the formulas are more involved. For this reason, it is useful to simplify the formulas as follows. We take polar decompositions of A and D ,

$$A = U_L \mathcal{D}_A U_R \quad \text{and} \quad D = V_L \mathcal{D}_B V_R$$

with unitary matrices $U_L, U_R \in \mathbb{C}^{q \times q}$ and $V_L, V_R \in \mathbb{C}^{p \times p}$. Here \mathcal{D}_A and \mathcal{D}_B are diagonal matrices

$$\mathcal{D}_A = \text{diag}(a_1, \dots, a_q) \quad \text{and} \quad \mathcal{D}_B = \text{diag}(d_1, \dots, d_p),$$

whose entries take values in the interval $(0, 1)$. It follows that

$$\hat{A} = \begin{pmatrix} U_L & 0 \\ 0 & V_L \end{pmatrix} \begin{pmatrix} \mathcal{D}_A & U_L^{-1} B V_R^{-1} \\ V_L^{-1} C U_R^{-1} & \mathcal{D}_B \end{pmatrix} \begin{pmatrix} U_R & 0 \\ 0 & V_R \end{pmatrix}.$$

The unitary transformations at the very left and right drop out when forming $\det(\mathbb{1} - \hat{A}^* \hat{A})$. The unitary transformations in the off-diagonal entries

$$B \rightarrow U_L^{-1} B V_R^{-1} \quad \text{and} \quad C \rightarrow V_L^{-1} C U_R^{-1}, \quad (4.32)$$

on the other hand, do not change the integration measures dB and dC . Indeed, these integration measures transform with the Jacobian determinant of the transformation matrices, if we write out the real and imaginary parts separately. As is shown in

Lemma 4.13 below, the resulting Jacobian determinants are equal to one. Therefore, the unitary transformations in (4.32) indeed preserve the integration measures.

After these transformations, we can assume without loss of generality that A and D coincide with the diagonal matrices \mathcal{D}_A and \mathcal{D}_D , respectively. After this simplification, the integrals over B and C can be computed as follows. We first rewrite the integrals as

$$\begin{aligned} \int \det(\mathbf{1} - X)^{N-2p-2q} dB dC &= \left\{ B, C = \frac{1}{\sqrt{N}} (\tilde{B}, \tilde{C}) \right\} \\ &= \frac{1}{N^{2pq}} \int \det \left\{ \mathbf{1} - \frac{1}{N} \begin{pmatrix} (\mathbf{1} - A^2)^{-1} \tilde{C}^* \tilde{C} & 0 \\ 0 & (\mathbf{1} - D^2)^{-1} \tilde{B}^* \tilde{B} \end{pmatrix} \right. \\ &\quad \left. - \frac{1}{\sqrt{N}} \begin{pmatrix} 0 & (\mathbf{1} - A^2)^{-1} (A\tilde{B} + \tilde{C}^* D) \\ (\mathbf{1} - D^2)^{-1} (\tilde{B}^* A + D\tilde{C}) & 0 \end{pmatrix} \right\}^{N-2p-2q} d\tilde{B} d\tilde{C}. \end{aligned}$$

Now we can again expand the determinant with the help of (4.22) to second order to obtain

$$\begin{aligned} &\int \det(\mathbf{1} - X)^{N-2p-2q} dB dC \\ &= \frac{1}{N^{2pq}} \left(1 + \mathcal{O}\left(\frac{1}{N}\right) \right) \int \left\{ 1 - \frac{1}{N} \operatorname{Tr} \begin{pmatrix} (\mathbf{1} - A^2)^{-1} \tilde{C}^* \tilde{C} & 0 \\ 0 & (\mathbf{1} - D^2)^{-1} \tilde{B}^* \tilde{B} \end{pmatrix} \right. \\ &\quad \left. - \frac{1}{2N} \operatorname{Tr} \begin{pmatrix} 0 & (\mathbf{1} - A^2)^{-1} (A\tilde{B} + \tilde{C}^* D) \\ (\mathbf{1} - D^2)^{-1} (\tilde{B}^* A + D\tilde{C}) & 0 \end{pmatrix}^2 \right\}^{N-2p-2q} d\tilde{B} d\tilde{C} \\ &= \frac{1}{N^{2pq}} \left(1 + \mathcal{O}\left(\frac{1}{N}\right) \right) \int d\tilde{B} d\tilde{C} \\ &\quad \times \exp \left(- \operatorname{Tr} \left((\mathbf{1} - A^2)^{-1} \tilde{C}^* \tilde{C} + (\mathbf{1} - D^2)^{-1} \tilde{B}^* \tilde{B} \right. \right. \\ &\quad \left. \left. + (\mathbf{1} - A^2)^{-1} (A\tilde{B} + \tilde{C}^* D) (\mathbf{1} - D^2)^{-1} (\tilde{B}^* A + D\tilde{C}) \right) \right). \end{aligned}$$

The trace can be simplified using the formulas

$$\begin{aligned} &\operatorname{Tr} \left((\mathbf{1} - A^2)^{-1} (A\tilde{B}) (\mathbf{1} - D^2)^{-1} (\tilde{B}^* A) \right) \\ &= \operatorname{Tr} \left(A^2 (\mathbf{1} - A^2)^{-1} \tilde{B} (\mathbf{1} - D^2)^{-1} \tilde{B}^* \right) \\ &= - \operatorname{Tr} \left(\tilde{B} (\mathbf{1} - D^2)^{-1} \tilde{B}^* \right) + \operatorname{Tr} \left((\mathbf{1} - A^2)^{-1} \tilde{B} (\mathbf{1} - D^2)^{-1} \tilde{B}^* \right). \end{aligned}$$

and similarly

$$\begin{aligned} &\operatorname{Tr} \left((\mathbf{1} - A^2)^{-1} (\tilde{C}^* D) (\mathbf{1} - D^2)^{-1} (D\tilde{C}) \right) \\ &= - \operatorname{Tr} \left((\mathbf{1} - A^2)^{-1} \tilde{C}^* \tilde{C} \right) + \operatorname{Tr} \left((\mathbf{1} - A^2)^{-1} \tilde{C}^* (\mathbf{1} - D^2)^{-1} \tilde{C} \right). \end{aligned}$$

This gives

$$\begin{aligned}
& \int \det(\mathbb{1} - X)^{N-2p-2q} dB dC \\
&= \frac{1}{N^{2pq}} \left(1 + \mathcal{O}\left(\frac{1}{N}\right) \right) \int d\tilde{B} d\tilde{C} \\
& \quad \times \exp \left(-\operatorname{Tr} \left((\mathbb{1} - A^2)^{-1} \tilde{B} (\mathbb{1} - D^2)^{-1} \tilde{B}^* + (\mathbb{1} - A^2)^{-1} \tilde{C}^* (\mathbb{1} - D^2)^{-1} \tilde{C} \right. \right. \\
& \quad \left. \left. + (\mathbb{1} - A^2)^{-1} A \tilde{B} (\mathbb{1} - D^2)^{-1} D \tilde{C} \right. \right. \\
& \quad \left. \left. + (\mathbb{1} - A^2)^{-1} \tilde{C}^* D (\mathbb{1} - D^2)^{-1} \tilde{B}^* A \right) \right).
\end{aligned}$$

Using that A and D are diagonal, we obtain

$$\begin{aligned}
& \int \det(\mathbb{1} - X)^{N-2p-2q} dB dC \\
&= \frac{1}{N^{2pq}} \left(1 + \mathcal{O}\left(\frac{1}{N}\right) \right) \int d\tilde{B} d\tilde{C} \\
& \quad \times \exp \left(- \sum_{i=1}^q \sum_{j=1}^p \left(\frac{1}{1-a_i^2} \frac{1}{1-d_j^2} (|\tilde{B}_{ij}|^2 + |\tilde{C}_{ij}^*|^2) \right. \right. \\
& \quad \left. \left. + \frac{a_i}{1-a_i^2} \frac{d_j}{1-d_j^2} (\tilde{B}_{ij} \overline{\tilde{C}_{ij}^*} + \overline{\tilde{B}_{ij}} \tilde{C}_{ij}^*) \right) \right).
\end{aligned}$$

This integral can be computed separately for every i and j , giving

$$X := \int_{\mathbb{C}} db \int_{\mathbb{C}} dc \exp \left(- \left\langle \begin{pmatrix} b \\ c \end{pmatrix}, Y \begin{pmatrix} b \\ c \end{pmatrix} \right\rangle_{\mathbb{C}^2} \right)$$

with

$$Y := \frac{1}{1-a_i^2} \frac{1}{1-d_j^2} \begin{pmatrix} 1 & a_i d_j \\ a_i d_j & 1 \end{pmatrix}.$$

Next,

$$\det Y = \left(\frac{1}{1-a_i^2} \frac{1}{1-d_j^2} \right)^2 (1 - a_i^2 d_j^2) = \frac{1 - a_i^2 d_j^2}{(1-a_i^2)^2 (1-d_j^2)^2},$$

and carrying out the Gaussian integrals with the help of (4.26) gives

$$\begin{aligned}
X &= \frac{\pi^2}{\det Y} = \pi^2 \frac{(1-a_i^2)^2 (1-d_j^2)^2}{1-a_i^2 d_j^2} \\
& \int \det(\mathbb{1} - X)^{N-2p-2q} dB dC \\
&= \frac{1}{N^{2pq}} \left(1 + \mathcal{O}\left(\frac{1}{N}\right) \right) \prod_{i=1}^q \prod_{j=1}^p \pi^2 \frac{(1-a_i^2)^2 (1-d_j^2)^2}{1-a_i^2 d_j^2} \\
&= \frac{\pi^{2pq}}{N^{2pq}} \det(\mathbb{1} - A^2)^{2p} \det(\mathbb{1} - D^2)^{2q} \left(1 + \mathcal{O}\left(\frac{1}{N}\right) \right) \prod_{i=1}^q \prod_{j=1}^p \frac{1}{1-a_i^2 d_j^2}.
\end{aligned}$$

The last term tells us about the “coupling” of A and B . It can be written as

$$\prod_{i=1}^q \prod_{j=1}^p \frac{1}{1 - a_i^2 d_j^2} = \frac{1}{\det(\mathbb{1} - A^2 \otimes D^2)}.$$

This gives the result. \square

It remains to show that the unitary transformations in (4.32) are measure-preserving. This follows immediately from the following lemma.

Lemma 4.13. *Let X be an invertible complex quadratic matrix. Then*

$$\det \begin{pmatrix} \operatorname{Re} X & -\operatorname{Im} X \\ \operatorname{Im} X & \operatorname{Re} X \end{pmatrix} = |\det X|^2.$$

Proof. By direct computation, one finds that

$$\begin{pmatrix} \operatorname{Re} X & -\operatorname{Im} X \\ \operatorname{Im} X & \operatorname{Re} X \end{pmatrix} \begin{pmatrix} \operatorname{Re} Y & -\operatorname{Im} Y \\ \operatorname{Im} Y & \operatorname{Re} Y \end{pmatrix} = \begin{pmatrix} \operatorname{Re}(XY) & -\operatorname{Im}(XY) \\ \operatorname{Im}(XY) & \operatorname{Re}(XY) \end{pmatrix}.$$

Proceeding inductively, we deduce that for any polynomial g ,

$$g \left(\begin{pmatrix} \operatorname{Re} X & -\operatorname{Im} X \\ \operatorname{Im} X & \operatorname{Re} X \end{pmatrix} \right) = \begin{pmatrix} \operatorname{Re}(g(X)) & -\operatorname{Im}(g(X)) \\ \operatorname{Im}(g(X)) & \operatorname{Re}(g(X)) \end{pmatrix}.$$

The continuous functional calculus allows us to extend this formula to any continuous function g . In particular,

$$\log \begin{pmatrix} \operatorname{Re} X & -\operatorname{Im} X \\ \operatorname{Im} X & \operatorname{Re} X \end{pmatrix} = \begin{pmatrix} \operatorname{Re} \log X & -\operatorname{Im} \log X \\ \operatorname{Im} \log X & \operatorname{Re} \log X \end{pmatrix}. \quad (4.33)$$

We now take the trace,

$$\begin{aligned} \log \det \begin{pmatrix} \operatorname{Re} X & -\operatorname{Im} X \\ \operatorname{Im} X & \operatorname{Re} X \end{pmatrix} &= \operatorname{Tr} \left(\log \begin{pmatrix} \operatorname{Re} X & -\operatorname{Im} X \\ \operatorname{Im} X & \operatorname{Re} X \end{pmatrix} \right) \\ &\stackrel{(4.33)}{=} \operatorname{Tr} \begin{pmatrix} \operatorname{Re} \log X & -\operatorname{Im} \log X \\ \operatorname{Im} \log X & \operatorname{Re} \log X \end{pmatrix} = 2 \operatorname{Tr} \operatorname{Re} \log X. \end{aligned}$$

Taking the exponential gives

$$\det \begin{pmatrix} \operatorname{Re} X & -\operatorname{Im} X \\ \operatorname{Im} X & \operatorname{Re} X \end{pmatrix} = e^{2 \operatorname{Tr} \operatorname{Re} \log X} = e^{2 \operatorname{Re} \operatorname{Tr} \log X} = |e^{\operatorname{Tr} \log X}|^2 = |\det X|^2.$$

This concludes the proof. \square

5. MODEL EXAMPLES

In this section we compute group integrals for two model examples in which the integrand is the exponential of a specific potential. As we shall see in Section 6, the localized refined pre-state can be computed by suitably combining the results of these two model examples.

5.1. An Exponential of a Linear Functional. We let $\mathcal{H} = \mathbb{C}^N$ with the canonical scalar product and $I \subset \mathcal{H}$ a q -dimensional subspace. Choosing an orthonormal basis, we identify I with \mathbb{C}^q . As in (4.25) we use the notation

$$A := \pi_I \mathcal{U}|_I.$$

We let $Y \in L(I)$ be a Hermitian $q \times q$ -matrix. In this section, we shall compute the integral

$$\oint_{\mathcal{G}} e^{N \operatorname{Re} \operatorname{Tr}(AY)} d\mu_{\mathcal{G}}(\mathcal{U})$$

asymptotically for large N . We first state our main result and discuss and prove it afterward.

Proposition 5.1. *Given a symmetric operator $Y \in L(I)$, we define the operator A_0 by*

$$A_0 := \frac{Y}{\sqrt{Y^2 + \mathbf{1}} + \mathbf{1}}. \quad (5.1)$$

Then, asymptotically for large N ,

$$\oint_{\mathcal{G}} e^{N \operatorname{Re} \operatorname{Tr}(AY)} d\mu_{\mathcal{G}}(\mathcal{U}) = e^{N \operatorname{Tr}(A_0 Y)} \frac{(\det(\mathbf{1} - A_0^2))^N}{\sqrt{\det(\mathbf{1} - A_0^2 \otimes A_0^2)}} (1 + \mathcal{O}(N^{-1})). \quad (5.2)$$

If $Y \neq 0$, the inequality

$$\operatorname{Tr}(A_0 Y) + \log \det(\mathbf{1} - A_0^2) > 0 \quad (5.3)$$

holds, showing that, for large N , the right side of (5.2) grows exponentially in N .

Before coming to the proof, we illustrate the statement of this proposition by proving a simple special case.

Example 5.2. ($q=1$) In the case $q = 1$, by applying Corollary (4.4), we obtain the leading contribution

$$\begin{aligned} & \oint_{\mathcal{G}} e^{N \operatorname{Re} \operatorname{Tr}(AY)} d\mu_{\mathcal{G}}(\mathcal{U}) (1 + \mathcal{O}(N^{-1})) \\ &= \mathcal{J} := \frac{N}{\pi} \int_{B_1} e^{Ny \operatorname{Re} a} (1 - |a|^2)^{N-2} d\mu_{\mathbb{C}}(a), \end{aligned}$$

where we write $Y = y\mathbf{1}$ (with $y \in \mathbb{R}$), and $B_1 \subset \mathbb{C}$ denotes the unit disc. Choosing polar coordinates, we obtain

$$\mathcal{J} = \frac{N}{\pi} \int_0^1 r dr \int_0^{2\pi} d\varphi e^{Ny r \cos \varphi} (1 - r^2)^{N-2} = \frac{N}{\pi} \int_0^1 dr \int_0^{2\pi} d\varphi e^{N g(r, \varphi)} \frac{r}{(1 - r^2)^2}$$

with the function g defined by

$$g(r, \varphi) := y r \cos \varphi + \log(1 - r^2).$$

Our strategy for evaluating this integral asymptotically for large N is to compute the maxima of the function g and to use the saddle point approximation. Without loss of generality, it suffices to consider the case $y > 0$. Then the maximum of g is attained at $\varphi = 0$ and the radius r with

$$0 = \frac{\partial}{\partial r} g(r, 0) = y - \frac{2r}{1 - r^2}.$$

This equation has the two solutions

$$r = \lambda_{\pm} := -\frac{1}{y} \pm \sqrt{1 + \frac{1}{y^2}}.$$

Clearly, only the solution λ_+ lies in the interval $[0, 1]$. We can simplify it to

$$\lambda_+ = \frac{\sqrt{y^2 + 1} - 1}{y} = \frac{\sqrt{y^2 + 1} - 1}{y} \frac{\sqrt{y^2 + 1} + 1}{\sqrt{y^2 + 1} + 1} = \frac{y}{\sqrt{y^2 + 1} + 1},$$

giving agreement with (5.1). Since $g(0, 0) = 0$ and $\lim_{r \nearrow 1} g(r, 0) = -\infty$, the point $(\lambda_+, \varphi = 0)$ is the unique maximum.

Having a unique maximum, we can compute the integral asymptotically for large N with the saddle point approximation: The Hessian of g at the maximum is the diagonal matrix

$$D^2g(\lambda_+, 0) = -2 \operatorname{diag} \left(\frac{1 + \lambda_+^2}{(1 - \lambda_+^2)^2}, \frac{\lambda_+^2}{1 - \lambda_+^2} \right).$$

Its determinant is computed by

$$\det D^2g(\lambda_+, 0) = 4 \frac{\lambda_+^2 (1 + \lambda_+^2)}{(1 - \lambda_+^2)^3} = 4 \frac{\lambda_+^2 (1 - \lambda_+^4)}{(1 - \lambda_+^2)^4}.$$

Hence, using (4.26), the saddle point approximation gives

$$\begin{aligned} \mathcal{J} &= \frac{e^{g(\lambda_+, 0)}}{\sqrt{\det D^2g(\lambda_+, 0)/2}} \frac{\lambda_+}{(1 - \lambda_+^2)^2} \left(1 + \mathcal{O}(N^{-1}) \right) \\ &= e^{Ny\lambda_+} (1 - \lambda_+^2)^N \sqrt{\frac{(1 - \lambda_+^2)^4}{\lambda_+^2 (1 - \lambda_+^4)}} \frac{\lambda_+}{(1 - \lambda_+^2)^2} \left(1 + \mathcal{O}(N^{-1}) \right) \\ &= e^{Ny\lambda_+} \frac{(1 - \lambda_+^2)^N}{\sqrt{1 - \lambda_+^4}} \left(1 + \mathcal{O}(N^{-1}) \right). \end{aligned}$$

This agrees with (5.2) in the special case $q = 1$. ◇

We now enter the proof of Proposition 5.1, which will be completed at the end of this section. Applying Corollary (4.4), we obtain the leading contribution

$$\begin{aligned} &\oint_{\mathcal{G}} e^{N \operatorname{Re} \operatorname{Tr}(AY)} d\mu_{\mathcal{G}}(\mathcal{U}) \left(1 + \mathcal{O}(N^{-1}) \right) \\ &= \left(\frac{N}{\pi} \right)^{(q^2)} \int_{\mathcal{A}_q} e^{N \operatorname{Re} \operatorname{Tr}(AY)} (\det(\mathbf{1} - A^* A))^{N-2q} d\mu_{\mathbb{C}^{q \times q}}(A). \end{aligned}$$

We rewrite the integrand as

$$e^{N \operatorname{Re} \operatorname{Tr}(AY)} (\det(\mathbf{1} - A^* A))^{N-2q} = e^{Ng(A)} (\det(\mathbf{1} - A^* A))^{-2q}$$

with

$$g : \mathcal{A}_q \rightarrow \mathbb{R}, \quad g(A) = \operatorname{Re} \operatorname{Tr}(AY) + \log \det(\mathbf{1} - A^* A). \quad (5.4)$$

Lemma 5.3. *The function $g(A)$ has a unique maximum at $A = A_0$ with A_0 according to (5.1).*

Proof. At the boundary of \mathcal{A}_q , the operator A^*A has eigenvalues which are equal to one, implying that the factor $\log \det(\mathbb{1} - A^*A)$ tends to minus infinity on this boundary. Therefore, it suffices to consider the interior of \mathcal{A}_q .

The determinant can be expanded linearly again using the formula (4.22). We thus obtain for first variations of g

$$\begin{aligned} \delta g(A) &= \operatorname{Re} \operatorname{Tr} ((\delta A) Y) - \operatorname{Tr} \left((\mathbb{1} - A^*A)^{-1} ((\delta A^*) A + A^*(\delta A)) \right) \\ &= \operatorname{Tr} \left(\delta A^* \left(\frac{Y}{2} - A (\mathbb{1} - A^*A)^{-1} \right) \right) + \operatorname{Tr} \left(\delta A \left(\frac{Y}{2} - (\mathbb{1} - A^*A)^{-1} A^* \right) \right). \end{aligned}$$

At the maximum, this variation vanishes for any choice of δA . The freedom in inserting a phase factor $\delta A \rightarrow e^{i\varphi} \delta A$ in the above equation shows that the summands involving δA and $(\delta A)^*$ must vanish separately. Thus

$$\operatorname{Tr} \left(\delta A^* \left\{ \frac{Y}{2} - A (\mathbb{1} - A^*A)^{-1} \right\} \right) = 0.$$

Choosing δA equal to the matrix in the curly brackets, we conclude that

$$A (\mathbb{1} - A^*A)^{-1} = \frac{Y}{2} = (\mathbb{1} - A^*A)^{-1} A^*. \quad (5.5)$$

We now form a polar decomposition of A , i.e.

$$A = US$$

with $U \in \operatorname{U}(\mathbb{C}^q)$ unitary and $S \in \operatorname{L}(\mathbb{C}^q)$ positive semi-definite with spectrum in the half-open interval $[0, 1)$. Then (5.5) can be written as

$$U T = Y = T U^{-1}, \quad (5.6)$$

where the operator T is defined via the spectral calculus as the positive semi-definite operator

$$T := 2S (\mathbb{1} - S^2)^{-1} = \frac{2S}{\mathbb{1} - S^2} \geq 0. \quad (5.7)$$

Applying (5.6) iteratively, it follows that $U^p T = T U^{-p}$. The continuous functional calculus yields

$$f(U) T = T f(U^{-1}) \quad (5.8)$$

for any continuous function f . Forming a spectral decomposition of U ,

$$U = \sum_k \lambda_k E_k \quad \text{with} \quad \lambda_k \in \mathbb{C}, |\lambda_k| = 1,$$

and evaluating (5.8) for functions f which vanish on the spectrum except at single eigenvalues, we conclude that for every eigenvalue $\lambda \in \sigma(U)$, also its complex conjugate $\bar{\lambda}$ is an eigenvalue of U and

$$E_\lambda T = T E_{\bar{\lambda}}.$$

Let λ with $\lambda \neq \bar{\lambda}$ be a non-real eigenvalue of U . Then

$$\operatorname{Tr} (E_\lambda T) = \operatorname{Tr} (E_\lambda (E_\lambda T)) = \operatorname{Tr} (E_\lambda (T E_{\bar{\lambda}})) = \operatorname{Tr} (E_{\bar{\lambda}} E_\lambda T) = 0.$$

Since T is positive semi-definite, it follows that

$$E_\lambda T = 0 = T E_\lambda.$$

In other words, the operator T vanishes on all the eigenspaces of U corresponding to non-real eigenvalues. Since the spectral calculus (5.7) maps the kernel of T to the

kernel of S , the same is true for the operator S . Therefore, by changing U to be the identity on the kernel of T , we can arrange the polar decomposition (5.6) with a unitary operator U with purely real eigenvalues.

After this construction, the operator U in (5.6) has the properties

$$U = U^* \quad \text{and} \quad [U, T] = 0 = [U, S].$$

Moreover, the operator A is symmetric. Therefore, the equations (5.5) simplify to

$$\frac{2A}{\mathbf{1} - A^2} = Y.$$

The corresponding scalar quadratic equation (which can be viewed as an equation for the eigenvalues of the corresponding matrices) takes the form

$$a^2 + \frac{2a}{y} - 1 = 0;$$

it has a unique root in the interval $(0, 1)$ provided that $y \neq 0$ given by

$$a = -\frac{1}{y} + \epsilon(y) \sqrt{1 + \frac{1}{y^2}} = \frac{y}{\sqrt{y^2 + 1} + 1},$$

where ϵ is the sign function $\epsilon(x) = 1$ for $x \geq 0$ and $\epsilon(x) = -1$ otherwise. The last formula also applies if $y = 0$. Therefore, the spectral calculus determines A uniquely by (5.1). \square

We next compute the Hessian at the maximum. It is most convenient to work in an eigenvector basis of the matrix A_0 .

Lemma 5.4. *Assume that the matrix A_0 in (5.1) is diagonal,*

$$A_0 = \text{diag}(\lambda_1, \dots, \lambda_q).$$

Then the function $g(A)$ defined in (5.4) has the quadratic expansion

$$\begin{aligned} g(A) = g(A_0) - \frac{1}{2} \sum_{i,j=1}^q \frac{1}{1 - \lambda_i^2} \frac{1}{1 - \lambda_j^2} \left\langle \begin{pmatrix} \delta A_j^i \\ \delta A_i^j \end{pmatrix}, \begin{pmatrix} 1 & \lambda_i \lambda_j \\ \lambda_i \lambda_j & 1 \end{pmatrix} \begin{pmatrix} \delta A_j^i \\ \delta A_i^j \end{pmatrix} \right\rangle_{\mathbb{C}^2} \\ + \mathcal{O}((\Delta A)^3), \end{aligned}$$

where δA denotes a linear variation, whereas $\Delta A := A - A_0$ is a small but finite perturbation.

Proof. The determinant can be expanded iteratively using the formula (4.22) together with

$$\frac{d}{d\tau} A(\tau)^{-1} = -A(\tau)^{-1} \dot{A}(\tau) A(\tau)^{-1}.$$

We thus obtain

$$\begin{aligned} A^* A &= A_0^2 + (\Delta A^*) A_0 + A_0 (\Delta A) + (\Delta A^*)(\Delta A) \\ \log \det(\mathbf{1}_{\mathbb{C}^q} - A^* A) &= \log \det(\mathbf{1} - A_0^2) \\ &\quad - \text{tr} \left((\mathbf{1} - A_0^2)^{-1} ((\Delta A^*) A_0 + A_0 (\Delta A)) \right) - \text{tr} \left((\mathbf{1} - A_0^2)^{-1} (\Delta A^*)(\Delta A) \right) \\ &\quad - \frac{1}{2} \text{tr} \left((\mathbf{1} - A_0^2)^{-1} ((\Delta A^*) A_0 + A_0 (\Delta A)) (\mathbf{1} - A_0^2)^{-1} ((\Delta A^*) A_0 + A_0 (\Delta A)) \right) \\ &\quad + \mathcal{O}((\Delta A)^3). \end{aligned}$$

Using (5.1), the term linear in ΔA cancels with the variation $\text{Re Tr}(Y \Delta A)$ of the first summand in (5.4). Using that A_0 is diagonal, we obtain

$$\begin{aligned} & g(A) - g(A_0) + \mathcal{O}((\delta A)^3) \\ &= - \sum_{i,j=1}^q \frac{1}{1-\lambda_j^2} |\delta A_j^i|^2 - \frac{1}{2} \sum_{i,j=1}^q \frac{1}{1-\lambda_i^2} \frac{1}{1-\lambda_j^2} (\overline{\delta A_j^i} \lambda_j + \lambda_i \delta A_j^i) (\overline{\delta A_j^i} \lambda_i + \lambda_j \delta A_j^i) \\ &= - \frac{1}{2} \sum_{i,j=1}^q \frac{1}{1-\lambda_i^2} \frac{1}{1-\lambda_j^2} \left(2 |\delta A_j^i|^2 + \lambda_i \lambda_j \left((\delta A_j^i)(\delta A_i^j) + \overline{(\delta A_j^i)(\delta A_i^j)} \right) \right). \end{aligned}$$

Writing the bilinear terms as a matrix expectation value gives the result. \square

Proof of Proposition 5.1. We now compute the Gaussian integrals. In the case $i = j$, setting $z = \delta A_i^i$, the matrix expectation value becomes

$$\begin{aligned} & \left\langle \begin{pmatrix} z \\ \bar{z} \end{pmatrix}, \begin{pmatrix} 1 & \lambda_i^2 \\ \lambda_i^2 & 1 \end{pmatrix} \begin{pmatrix} z \\ \bar{z} \end{pmatrix} \right\rangle_{\mathbb{C}^2} = 2|z|^2 + \lambda_i^2 (z^2 + \bar{z}^2) \\ &= 2(\text{Re}^2 z + \text{Im}^2 z) + \lambda_i^2 (2\text{Re}^2 z - 2\text{Im}^2 z) = 2(1 + \lambda_i^2) \text{Re}^2 z + 2(1 - \lambda_i^2) \text{Im}^2 z. \end{aligned}$$

Thus

$$\begin{aligned} & \int_{\mathbb{C}} \exp \left(- \frac{N}{2} \frac{1}{(1-\lambda_i^2)^2} \left\langle \begin{pmatrix} z \\ \bar{z} \end{pmatrix}, \begin{pmatrix} 1 & \lambda_i^2 \\ \lambda_i^2 & 1 \end{pmatrix} \begin{pmatrix} z \\ \bar{z} \end{pmatrix} \right\rangle_{\mathbb{C}^2} \right) dz \\ &= \int_{-\infty}^{\infty} d\text{Re } z \int_{-\infty}^{\infty} d\text{Im } z \exp \left(- \frac{N}{(1-\lambda_i^2)^2} \left((1 + \lambda_i^2) \text{Re}^2 z - (1 - \lambda_i^2) \text{Im}^2 z \right) \right) \\ &= \frac{\pi}{N} \frac{(1 - \lambda_i^2)^2}{\sqrt{(1 + \lambda_i^2)(1 - \lambda_i^2)}} = \frac{\pi}{N} \frac{(1 - \lambda_i^2)^2}{\sqrt{1 - \lambda_i^4}}, \end{aligned}$$

where we again carried out the Gaussian integrals with the help of (4.26).

Given $i < j$, we get a four-dimensional Gaussian integral over $z := \delta A_j^i$ and $\hat{z} := \overline{\delta A_i^j}$,

$$\begin{aligned} & \int_{\mathbb{C}} dz \int_{\mathbb{C}} d\hat{z} \exp \left(- N \frac{1}{1-\lambda_i^2} \frac{1}{1-\lambda_j^2} \left\langle \begin{pmatrix} \delta A_j^i \\ \delta A_i^j \end{pmatrix}, \begin{pmatrix} 1 & \lambda_i \lambda_j \\ \lambda_i \lambda_j & 1 \end{pmatrix} \begin{pmatrix} \delta A_j^i \\ \delta A_i^j \end{pmatrix} \right\rangle_{\mathbb{C}^2} \right) \\ &= \frac{\pi^2}{N^2} \frac{(1 - \lambda_i^2)^2 (1 - \lambda_j^2)^2}{1 - \lambda_i^2 \lambda_j^2}. \end{aligned}$$

Putting these results together, we obtain

$$\begin{aligned} & \oint_{\mathcal{G}} e^{N \text{Re Tr}(AY)} d\mu_{\mathcal{G}}(\mathcal{U}) \\ &= e^{N \text{Tr}(A_0 Y)} (\det(\mathbf{1} - A_0^2))^{N-2q} \prod_{i,j=1}^q \frac{(1 - \lambda_i^2)(1 - \lambda_j^2)}{\sqrt{1 - \lambda_i^2 \lambda_j^2}} (1 + \mathcal{O}(N^{-1})) \end{aligned}$$

(note that $\text{Tr}(A_0 Y)$ is real, making it possible to leave out the real part in the exponent). Writing the products with determinants gives (5.2).

In order to prove (5.3), we recall that in Lemma 5.3 we showed that the function g has a unique maximum at A_0 . Since $g(0) = 0$, this uniqueness statement gives rise to the implication

$$A_0 \neq 0 \implies g(A_0) > 0.$$

We finally note that, in view of (5.1), A_0 vanishes only if $Y = 0$. This concludes the proof of (5.3). \square

5.2. An Exponential of a Quartic Functional. We let $\mathcal{H} = \mathbb{C}^N$ with the canonical scalar product. We consider the finite one-dimensional set X formed of q points

$$X := \{1, \dots, q\}. \quad (5.9)$$

We let $I \simeq \mathbb{C}^q$ be the space of complex-valued functions on X . We choose the canonical orthonormal basis $(e_y)_{y \in X}$ by

$$e_y : X \rightarrow \mathbb{C}, \quad (e_y)(x) = \delta_{x,y}.$$

We regard I as a subspace of the Hilbert space \mathcal{H} . The wave evaluation operator is introduced for any $x \in X$ by

$$\Psi(x) : I \rightarrow \mathbb{C}, \quad e_y \mapsto e_y(x). \quad (5.10)$$

The computation

$$\langle \Psi(x)^* z | e_y \rangle = \bar{z} (\Psi(x) e_y) = \bar{z} e_y(x) = \left\langle \sum_{a=1}^q \overline{e_a(x)} e_a z \middle| e_y \right\rangle$$

(with $z \in \mathbb{C}$) shows that the adjoint of the wave evaluation operator is given by

$$\Psi(x)^* = \sum_{a=1}^q \overline{e_a(x)} e_a. \quad (5.11)$$

We want to model the surface layer integral (2.42), keeping $\xi = y - x$ fixed, in the simplified situation that all the wave functions are “in phase” along the light cone, meaning that $\Psi(x)$ and $\Psi(x + \xi)$ coincide. This leads us to considering the expression

$$\mathcal{T}(A_<, A_>) := \sum_{x \in X} \left| \text{tr}_I (\mathcal{U}_> \Psi(x)^* \Psi(x) \mathcal{U}_<^{-1} \Psi(x)^* \Psi(x)) \right|^2. \quad (5.12)$$

Substituting the form of the wave evaluation operator (5.10) and its adjoint (5.11), we obtain

$$\mathcal{T}(A_<, A_>) = \sum_{x \in X} \left| \text{tr}_I (\mathcal{U}_> \Psi(x)^* \Psi(x) \mathcal{U}_<^* \Psi(x)^* \Psi(x)) \right|^2 \quad (5.13)$$

$$= \sum_{x \in X} (A_>)_x^x (A_<^*)_x^x (A_>^*)_x^x (A_<)_x^x. \quad (5.14)$$

We want to compute the integral

$$\oint_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_>) \oint_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_<) e^{\beta N \mathcal{T}(A_<, A_>)}. \quad (5.15)$$

Before entering the detailed computation, we illustrate the structure of this expression by considering the case of one point.

Example 5.5. (q=1) In the case $q = 1$, the functional \mathcal{T} in (5.14) simplifies to

$$\mathcal{T} = |a_{<}|^2 |a_{>}|^2.$$

Hence, using Theorem 4.2 in the case $q = 1$, we obtain

$$\begin{aligned} & \oint_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_{>}) \oint_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_{<}) e^{\beta N \mathcal{T}(A_{<}, A_{>})} \\ &= \frac{(N-1)^2}{\pi^2} \int_{B_1} da_{<} \int_{B_1} da_{>} e^{\beta N |a_{<}|^2 |a_{>}|^2} (1 - |a_{<}|^2)^{N-2} (1 - |a_{>}|^2)^{N-2}, \end{aligned}$$

where $B_1 \subset \mathbb{C}$ denotes the unit disc. Choosing polar coordinates, we obtain

$$\begin{aligned} & \oint_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_{>}) \oint_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_{<}) e^{\beta N \mathcal{T}(A_{<}, A_{>})} \\ &= 4(N-1)^2 \int_0^1 r_{<} dr_{<} \int_0^1 r_{>} dr_{>} e^{\beta N r_{<}^2 r_{>}^2} (1 - r_{<}^2)^{N-2} (1 - r_{>}^2)^{N-2} \\ &= 4(N-1)^2 \int_0^1 r_{<} dr_{<} \int_0^1 r_{>} dr_{>} e^{N g(r_{<}, r_{>})} \frac{1}{(1 - r_{<}^2)^2} \frac{1}{(1 - r_{>}^2)^2}, \end{aligned}$$

where

$$g(r_{<}, r_{>}) := \beta r_{<}^2 r_{>}^2 + \log(1 - r_{<}^2) + \log(1 - r_{>}^2).$$

Our strategy for evaluating this integral asymptotically for large N is to compute the maxima of the function g and to use the saddle point approximation. In order to compute the maxima, we set the gradient of g to zero. This gives the equations

$$\beta r_{<}^2 = \frac{1}{1 - r_{>}^2} \quad \text{and} \quad \beta r_{>}^2 = \frac{1}{1 - r_{<}^2}.$$

Solving the first equation for $r_{>}^2$ and substituting it into the second equation, we obtain the quadratic equation

$$(r_{>}^2)^2 - r_{>}^2 + \frac{1}{\beta} = 0.$$

If $\beta < 4$, this quadratic polynomial has no roots. This means that the function g has no interior maxima. Its global maximum is at the origin,

$$g(0, 0) = 0.$$

In the case $\beta \geq 4$, however, the quadratic polynomial has two roots. We thus obtain the local extrema at

$$r_{>}^2 = r_{<}^2 = \lambda_{-} := \frac{1}{2} - \frac{1}{2} \sqrt{1 - \frac{4}{\beta}} \quad \text{and} \quad r_{>}^2 = r_{<}^2 = \lambda_{+} := \frac{1}{2} + \frac{1}{2} \sqrt{1 - \frac{4}{\beta}}.$$

The first extremum is a minimum, whereas the second is a local maximum. The value of g at the local maximum is given by

$$g(c, c) = \frac{c^2}{1 - c^2} + 2 \log(1 - c^2) \quad \text{with} \quad c := \sqrt{\lambda_{+}}.$$

For sufficiently large β (more precisely, if $\beta > 4.911\dots$), the local maximum is strictly positive and thus the global maximum. This makes it possible to compute the integral with the saddle point approximation. To this end, one computes the Hessian of g to

$$D^2 g(c, c) = -\frac{4}{(1 - c^2)^2} \begin{pmatrix} c^2 & -(1 - c^2) \\ -(1 - c^2) & c^2 \end{pmatrix}.$$

Hence the saddle point approximation gives

$$\begin{aligned}
& \oint_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_{>}) \oint_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_{<}) e^{\beta N \mathcal{T}(A_{<}, A_{>})} \left(1 + \mathcal{O}(N^{-1})\right) \\
&= 4(N-1)^2 \frac{c^2}{(1-c^2)^4} \int_0^1 dr_{<} \int_0^1 dr_{>} e^{Ng(c,c)} \frac{\pi}{N} \frac{1}{\sqrt{\det(-D^2 g(c,c)/2)}} \\
&= 4\pi N \frac{c^2}{(1-c^2)^4} e^{Ng(c,c)} \frac{(1-c^2)^2}{2} \frac{1}{\sqrt{2c^2-1}} \\
&= 2\pi N \frac{1}{(1-c^2)^2} \frac{c^2}{\sqrt{2c^2-1}} \exp \left\{ N \left(\frac{c^2}{1-c^2} + 2 \log(1-c^2) \right) \right\}.
\end{aligned}$$

In this way, we have computed the double integral asymptotically for large N . \diamond

We now state our main result for general q . Exactly as in the last example, we only get local maxima if $\beta > 4$. In this case, we can again use the saddle point approximation to obtain the following result.

Theorem 5.6. *Assume that $\beta > 4$. Then the integrand of the group integral (5.15) with \mathcal{T} according to (5.12) has saddle points at*

$$A_{<} = c \operatorname{diag} (e^{i\varphi_{<}^1}, \dots, e^{i\varphi_{<}^q}), \quad A_{>} = c \operatorname{diag} (e^{i\varphi_{>}^1}, \dots, e^{i\varphi_{>}^q}), \quad (5.16)$$

where $\varphi_{<}^1, \dots, \varphi_{<}^q$ and $\varphi_{>}^1, \dots, \varphi_{>}^q$ are arbitrary real phase angles, and c is given in terms of β by

$$c^2 = \frac{1}{2} + \frac{1}{2} \sqrt{1 - \frac{4}{\beta}}. \quad (5.17)$$

The corresponding contribution to the group integral (5.15) has the large- N -asymptotics

$$\begin{aligned}
& \oint_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_{>}) \oint_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_{<}) e^{\beta N \mathcal{T}(A_{<}, A_{>})} \\
& \asymp (2\pi N)^q (1-c^4)^{-q^2} \left(\frac{1+c^2}{1-c^2} \frac{c^2}{\sqrt{2c^2-1}} \right)^q \\
& \quad \times \exp \left\{ Nq \left(\frac{c^2}{1-c^2} + 2 \log(1-c^2) \right) \right\} \left(1 + \mathcal{O}\left(\frac{1}{N}\right) \right) \quad (5.18)
\end{aligned}$$

(here the symbol \asymp means that we restrict attention to the contribution near the saddle points).

The remainder of this section is devoted to the proof of this theorem. Applying Theorem 4.2, the double integral (5.15) can be rewritten as

$$\begin{aligned}
& \oint_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_{>}) \oint_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_{<}) e^{\beta N \mathcal{T}(A_{<}, A_{>})} \\
&= C \int_{\mathcal{A}_q} d\mu_{\mathbb{C}^{q \times q}}(A_{<}) \int_{\mathcal{A}_q} d\mu_{\mathbb{C}^{q \times q}}(A_{>}) \\
& \quad \times e^{\beta N \mathcal{T}(A_{<}, A_{>})} (\det(\mathbf{1} - A_{<}^* A_{<}))^{N-2q} (\det(\mathbf{1} - A_{>}^* A_{>}))^{N-2q} \\
&= C \int_{\mathcal{A}_q} d\mu_{\mathbb{C}^{q \times q}}(A_{<}) \int_{\mathcal{A}_q} d\mu_{\mathbb{C}^{q \times q}}(A_{>}) \\
& \quad \times e^{Ng(A_{<}, A_{>})} (\det(\mathbf{1} - A_{<}^* A_{<}))^{-2q} (\det(\mathbf{1} - A_{>}^* A_{>}))^{-2q} \quad (5.19)
\end{aligned}$$

with the function $g(A_<, A_>)$ and the normalization constant C given by

$$g(A_<, A_>) := \beta \mathcal{T}(A_<, A_>) + \log \det(\mathbf{1} - A_<^* A_<) + \log \det(\mathbf{1} - A_>^* A_>) \quad (5.20)$$

$$C := \left(\frac{1}{\pi^{(q^2)}} \frac{(N-1)! \cdots (N-q)!}{(N-q-1)! \cdots (N-2q)!} \right)^2 = \left(\frac{N}{\pi} \right)^{2q^2} \left(1 + \mathcal{O}(N^{-1}) \right). \quad (5.21)$$

The first step is to determine the maxima of the function g .

Lemma 5.7. *Assume that $\beta > 4$. Then the function g has a global maximum at $(A_<, A_>)$ if and only if both matrices $A_<$ and $A_>$ are, up to phase factors on the diagonal, the same multiple of the identity matrix, i.e.*

$$A_< = c \operatorname{diag} (e^{i\varphi_<^1}, \dots, e^{i\varphi_<^q}), \quad A_> = c \operatorname{diag} (e^{i\varphi_>^1}, \dots, e^{i\varphi_>^q}) \quad (5.22)$$

with phase angles $\varphi_{<,>}^x \in \mathbb{R}$. Moreover, the parameter $c > 0$ is related to β by

$$c^2 = \frac{1}{2} + \frac{1}{2} \sqrt{1 - \frac{4}{\beta}} \quad \text{or} \quad \beta = \frac{1}{c^2(1 - c^2)}. \quad (5.23)$$

The maximum of g is given by

$$g(c\mathbf{1}_{\mathbb{C}^q}, c\mathbf{1}_{\mathbb{C}^q}) = \frac{q c^2}{1 - c^2} + 2q \log(1 - c^2). \quad (5.24)$$

Proof. Obviously, the functional $\det(\mathbf{1} - A^* A)$ does not change if A is multiplied from the left or right by a unitary matrix, i.e. it is invariant under the transformation

$$A \rightarrow U A V \quad \text{with} \quad U, V \in \mathbf{U}(I). \quad (5.25)$$

With this in mind, we first compute the maximum of the function g under such transformations. In view of (5.20), we need to maximize the functional $\mathcal{T}(U A_< V, A_>)$. We begin by varying $A_<$, keeping $A_>$ fixed. We set

$$s_x := |(A_>)_x^x|^2 \geq 0.$$

By renaming the points in X we can arrange that

$$0 \leq s_1 \leq \cdots \leq s_q.$$

Using this notation in (5.14), we obtain

$$\mathcal{T}(U A_< V, A_>) = \sum_{x \in X} s_x \left| (U A_< V)_x^x \right|^2 \leq \sum_{x, y \in X} s_x \left| (U A_< V)_x^y \right|^2 \quad (5.26)$$

$$= \sum_{x \in X} s_x \left((U A_< V) (V^* A_<^* U^*) \right)_x^x = \sum_{x \in X} s_x (U A_< A_<^* U^*)_x^x =: G(U) \quad (5.27)$$

(in the inequality in (5.26) we increased the sum by taking more non-negative summands).

Let us prove the estimate

$$G(U) \leq \sum_{\ell \in X} s_\ell \lambda_{<}^\ell, \quad (5.28)$$

where $\lambda_{<}^1, \dots, \lambda_{<}^q$ are the eigenvalues of the symmetric matrix $A_< A_<^*$ in increasing order, i.e.

$$0 \leq \lambda_{<}^1 \leq \cdots \leq \lambda_{<}^q. \quad (5.29)$$

To this end, let U be a maximum of G (this maximum is attained due to continuity of G and compactness of the group $U(I)$). We consider a first order variation $\delta U = iBU$ with a symmetric operator B . Rewriting $G(U)$ as a trace according to

$$G(U) = \text{tr}_I (S U A_{<} A_{<}^* U^*)$$

with $S := \text{diag}(s_1, \dots, s_q)$, maximality implies that

$$0 = \delta G(U) = i \text{tr}_I (S B U A_{<} A_{<}^* U^* - S U A_{<} A_{<}^* U^* B^*) = \text{tr}_I (B i [U A_{<} A_{<}^* U^*, S]).$$

Since B is arbitrary, it follows that the commutator vanishes,

$$[U A_{<} A_{<}^* U^*, S] = 0.$$

This shows that the matrix $U A_{<} A_{<}^* U^*$ is invariant on the eigenspaces of S . This in turns implies that G has the form (5.28) with λ_ℓ the eigenvalues of $A_{<} A_{<}^*$, in an arbitrary order. For the maximum, we obviously need to reorder the eigenvalues according to (5.29). This completes the proof of (5.28) and (5.29).

We next study how to arrange equality in all these estimates. To this end, we form a polar decomposition of $A_{<}$ by choosing unitary operators U and V such that

$$U A_{<} V = \text{diag} \left(\sqrt{\lambda_{<}^1}, \dots, \sqrt{\lambda_{<}^q} \right).$$

In this way, the upper bound (5.28) is realized. This diagonal matrix also realizes equality in (5.26). Therefore, it is not only a maximizer of the functional G , but it is even a desired maximizer of the functional $\mathcal{T}(U A_{<} V, A_{>})$.

It remains to determine the general maximizer. To this end, we note that the inequality in (5.26) becomes an equality if and only if $(U A_{<} V)_x^y$ vanishes for all $x \neq y$. In other words, the matrix $U A_{<} V$ must be diagonal. By direct computation, one sees that each diagonal entry must be a phase factor times the square root of the $\lambda_{<}^\ell$. We conclude that the maximum of $\mathcal{T}(U A_{<} V, A_{>})$ under variations of U and V is attained if and only if

$$U A_{<} V = \text{diag} \left(e^{i\varphi_{<}^1} \sqrt{\lambda_{<}^1}, \dots, e^{i\varphi_{<}^q} \sqrt{\lambda_{<}^q} \right) \quad (5.30)$$

with arbitrary phase angles $\varphi_{<}^1, \dots, \varphi_{<}^q$.

We next interchange the roles of $A_{<}$ and $A_{>}$ and vary $A_{>}$ while keeping $A_{<}$ fixed. We thus obtain that the maximum of g is attained for matrices $U A_{>} V$ again of the form (5.30), but with $<$ replaced by $>$. We conclude that, using the unitary invariance of the functional $\det(\mathbb{1} - A^* A)$ under the transformation (5.25), the maxima of g are attained if the matrices $A_{>}$ and $A_{<}$ are both diagonal. Restricting attention to such diagonal matrices, the functional $g(A_{<}, A_{>})$ simplifies to

$$g(A_{<}, A_{>}) = \sum_{x=1}^q \left(\beta \lambda_{<}^x \lambda_{>}^x + \log(1 - \lambda_{<}^x) + \log(1 - \lambda_{>}^x) \right). \quad (5.31)$$

Now we can maximize for every x separately. Similar as in Example 5.5, the criticality conditions are

$$\beta \lambda_{>}^x = \frac{1}{1 - \lambda_{<}^x}, \quad \beta \lambda_{<}^x = \frac{1}{1 - \lambda_{>}^x}.$$

Solving the first equation for $\lambda_{>}^x$ and substituting it into the second equation, we obtain a quadratic equation for $\lambda_{<}^x$, having the two roots

$$\lambda_{\pm} = \frac{1}{2} \pm \frac{1}{2} \sqrt{1 - \frac{4}{\beta}}.$$

By direct computation one sees that the plus sign gives the maximum. Furthermore, one readily finds that $\lambda_{>}^x = \lambda_{<}^x = \lambda_+ = c^2$ with c^2 according to (5.23). Moreover, each summand in (5.31) takes the value

$$\begin{aligned} & \beta \lambda_{<}^x \lambda_{>}^x + \log(1 - \lambda_{<}^x) + \log(1 - \lambda_{>}^x) \\ &= \beta c^4 + 2 \log(1 - c^2) = \frac{c^2}{1 - c^2} + 2 \log(1 - c^2), \end{aligned}$$

where in the last step we used that β can be expressed in terms of c by the right equation in (5.23). Using this formula in (5.31) gives (5.24). This concludes the proof. \square

We next expand the function g near its maximum. In view of the gauge freedom in (5.22), it suffices to consider the maximum at

$$A_{<} = A_{>} = c \mathbf{1} \quad (5.32)$$

with c as given by (5.23). It suffices to expand the matrices $A_{<}$ and $A_{>}$ to first order, i.e.

$$A_{<,>} = c \mathbf{1} + \Delta A_{<,>} \quad \text{with} \quad \Delta A_{<,>} = \tau \delta A_{<,>} + \mathcal{O}(\tau^2),$$

where τ denotes the perturbation parameter.

Lemma 5.8. *Near the maximum (5.32), the function $g(A_{<}, A_{>})$ has the quadratic expansion*

$$g(A_{<}, A_{>}) = \frac{qc^2}{1 - c^2} + 2q \log(1 - c^2) \quad (5.33)$$

$$+ \frac{4\tau^2}{1 - c^2} \sum_{x \in X} \left(\operatorname{Re} \left((\delta A_{>})_x^x \right) \operatorname{Re} \left((\delta A_{<})_x^x \right) \right) \quad (5.34)$$

$$- \frac{2\tau^2 c^2}{(1 - c^2)^2} \sum_{x \in X} \left(\left| \operatorname{Re}(\delta A_{<})_x^x \right|^2 + \left| \operatorname{Re}(\delta A_{>})_x^x \right|^2 \right) \quad (5.35)$$

$$- \frac{\tau^2}{1 - c^2} \sum_{x \neq y} \left(\left| (\delta A_{<})_y^x \right|^2 + \left| (\delta A_{>})_y^x \right|^2 \right) \quad (5.36)$$

$$- \frac{\tau^2 c^2}{2(1 - c^2)^2} \sum_{x \neq y} \left(\left| (\delta A_{<})_y^x + \overline{(\delta A_{<})_x^y} \right|^2 + \left| (\delta A_{>})_y^x + \overline{(\delta A_{>})_x^y} \right|^2 \right) + \mathcal{O}(\tau^3). \quad (5.37)$$

Proof. We first expand the function $\mathcal{T}(A_{<}, A_{>})$ in (5.14),

$$\begin{aligned} \mathcal{T}(A_{<}, A_{>}) &= \sum_{x \in X} \left| (A_{<})_x^x \right|^2 \left| (A_{>}^*)_x^x \right|^2 \\ &= qc^4 + 2c^3 \sum_{x \in X} \operatorname{Re} \left((\Delta A_{<})_x^x + (\Delta A_{>})_x^x \right) \end{aligned} \quad (5.38)$$

$$+ \tau^2 c^2 \sum_{x \in X} \left(\left| (\delta A_{<})_x^x \right|^2 + \left| (\delta A_{>})_x^x \right|^2 \right) \quad (5.39)$$

$$+ 4\tau^2 c^2 \sum_{x \in X} \left(\operatorname{Re} \left((\delta A_{>})_x^x \right) \operatorname{Re} \left((\delta A_{<})_x^x \right) \right) + \mathcal{O}(\tau^3). \quad (5.40)$$

The last two summands in (5.20), on the other hand, can be expanded as in the proof of Lemma 5.4. We thus obtain

$$\begin{aligned}
A^*A &= c^2 \mathbf{1} + c(\Delta A^* + \Delta A) + (\Delta A^*)(\Delta A) \\
\log \det(\mathbf{1} - A^*A) &= \log \det((1 - c^2) \mathbf{1}_{\mathbb{C}^q}) \\
&\quad - \operatorname{tr} \left((\mathbf{1} - A^*A)^{-1} c(\Delta A^* + \Delta A) \right) - \operatorname{tr} \left((\mathbf{1} - A^*A)^{-1} (\Delta A^*)(\Delta A) \right) \\
&\quad - \frac{1}{2} \operatorname{tr} \left((\mathbf{1} - A^*A)^{-1} c(\Delta A^* + \Delta A)(\mathbf{1} - A^*A)^{-1} c(\Delta A^* + \Delta A) \right) + \mathcal{O}((\Delta A)^3) \\
&= q \log(1 - c^2) - \frac{c}{1 - c^2} \operatorname{tr}(\Delta A^* + \Delta A) - \frac{\tau^2}{1 - c^2} \operatorname{tr}((\delta A^*)(\delta A)) \\
&\quad - \frac{\tau^2 c^2}{2(1 - c^2)^2} \operatorname{tr}((\delta A^* + \delta A)(\delta A^* + \delta A)) + \mathcal{O}(\tau^3) \\
&= q \log(1 - c^2) - \frac{2c}{1 - c^2} \sum_{x \in X} \operatorname{Re}(\Delta A)_x^x - \frac{\tau^2}{1 - c^2} \sum_{x \in X} |(\delta A)_x^x|^2 - \frac{\tau^2}{1 - c^2} \sum_{x \neq y} |(\delta A)_y^x|^2 \\
&\quad - \frac{2\tau^2 c^2}{(1 - c^2)^2} \sum_{x \in X} |\operatorname{Re}(\delta A)_x^x|^2 - \frac{\tau^2 c^2}{2(1 - c^2)^2} \sum_{x \neq y} \left| (\delta A)_y^x + \overline{(\delta A)_x^y} \right|^2 + \mathcal{O}(\tau^3),
\end{aligned}$$

where in the last step we wrote the contributions by the diagonal and off-diagonal matrix entries separately.

Adding all the contributions and using the formula for β in (5.23) gives the result. \square

One complication in applying the saddle point approximation is that there is an underlying *local gauge symmetry*. This gauge symmetry is apparent in (5.13), showing that \mathcal{T} is invariant under local phase transformations of $\mathcal{U}_<$ and $\mathcal{U}_>$,

$$(A_<)_y^x \rightarrow e^{i\tau\varphi_<(x)} (A_<)_y^x \quad \text{and} \quad (A_>)_y^x \rightarrow e^{i\tau\varphi_>(x)} (A_>)_y^x,$$

where $\varphi_<$ and $\varphi_>$ are two arbitrary real-valued functions on X . Likewise, the global maximum constructed in Lemma 5.7 is unique only up to the gauge phases in (5.22). Moreover, the gauge invariance becomes apparent in the quadratic expansion in Lemma 5.8 in that the imaginary parts of the diagonal matrix entries

$$\operatorname{Im} \delta A_x^x \quad \text{with} \quad x \in X$$

do not enter in the expansion. The method to deal with the gauge invariance is to integrate over the resulting compact gauge group before applying the saddle-point approximation. Then we can work with arbitrary representatives of the gauge orbits. This justifies the following *gauge-fixing procedure*. Perturbing to first order around the maximum at $A_{<,>} = c\mathbf{1}_{\mathbb{C}^q}$, we obtain

$$A_{<,>} = c\mathbf{1}_{\mathbb{C}^q} + i\tau c \varphi_{<,>}(x) + \mathcal{O}(\tau^2).$$

Thus we can use the linear gauge freedom in every order of perturbation theory in order to arrange that the finite perturbations $\Delta A_{<,>}$ satisfy for all $k \in \{1, \dots, q\}$ the constraints

$$\operatorname{Im}(\Delta A_{<,>})_x^x = 0 \quad \text{for all } a \in \{1, \dots, q\}.$$

After this gauge fixing, we are ready to compute the Gaussian integral in the saddle point approximation.

Proof of Theorem 5.6. Our strategy is to carry out the Gaussian integral step by step using the relation (4.26). Before beginning, we integrate over the gauge phases of $A_{<}$. This amounts to q integrals over a circle of radius c , giving rise to the factor

$$(2\pi c)^q.$$

Clearly, integrating over the gauge phases of $A_{>}$ gives the same factor. The terms in (5.34) and (5.35) give, for each x , a two-dimensional Gaussian integral with the integrand

$$\exp \left(-N \left\langle \begin{pmatrix} \text{Re}(\delta A_{>}^x_x) \\ \text{Re}(\delta A_{<}^x_x) \end{pmatrix}, E \begin{pmatrix} \text{Re}(\delta A_{>}^x_x) \\ \text{Re}(\delta A_{<}^x_x) \end{pmatrix} \right\rangle_{\mathbb{C}^2} \right)$$

and with the covariance matrix

$$E := \frac{2}{(1-c^2)^2} \begin{pmatrix} c^2 & -(1-c^2) \\ -(1-c^2) & c^2 \end{pmatrix}.$$

Gaussian integration gives a factor

$$\frac{\pi}{\sqrt{\det(NE)}} = \frac{\pi}{N} \frac{(1-c^2)^2}{\sqrt{8c^2-4}}.$$

In this way, for the resulting $2q$ -dimensional Gaussian integrals over $\text{Re}(\delta A_{<}^x_x)$ and $\text{Re}(\delta A_{>}^x_x)$ we obtain

$$\left(\frac{\pi}{N} \frac{(1-c^2)^2}{\sqrt{8c^2-4}} \right)^q.$$

We next consider the off-diagonal matrix entries $(\delta A_{<}^x_y)$ for $x \neq y$. Multiplying out (5.37), one sees that for each $x \neq y$ one gets a Gaussian integral with the covariance matrix

$$\exp \left(-N \left\langle \begin{pmatrix} \text{Re}(\delta A_{>}^x_y) \\ \text{Re}(\delta A_{<}^y_x) \end{pmatrix}, F \begin{pmatrix} \text{Re}(\delta A_{>}^x_y) \\ \text{Re}(\delta A_{<}^y_x) \end{pmatrix} \right\rangle_{\mathbb{C}^2} \right)$$

with the covariance matrix

$$F := \frac{1}{(1-c^2)^2} \begin{pmatrix} 1 & c^2 \\ c^2 & 1 \end{pmatrix}.$$

Thus Gaussian integration gives a factor

$$\frac{\pi}{\sqrt{\det(NF)}} = \frac{\pi}{N} \frac{(1-c^2)^2}{\sqrt{1-c^4}}.$$

The integral over $\text{Im}(\delta A_{<}^y_x)$ and $\text{Im}(\delta A_{<}^y_x)$ gives the same factor. Moreover, the integrals over the off-diagonal terms of $A_{>}$ can be treated in the same way. Keeping in mind that we have $q(q-1)/2$ ways to choose indices x, y with $x < y$, we end up with the factor

$$\left(\frac{\pi}{N} \frac{(1-c^2)^2}{\sqrt{1-c^4}} \right)^{2q(q-1)}.$$

Finally, the additional factors in (5.19) need to be evaluated at the maximum. We thus obtain the factor

$$(\det(\mathbf{1} - A^*A))^{-2q}|_{A=c\mathbf{1}} = ((1-c^2)^q)^{-2q} = (1-c^2)^{-2q^2},$$

again for $A_{<}$ and $A_{>}$.

Combining all the terms gives

$$\begin{aligned}
& \int_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_{>}) \int_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_{<}) e^{\beta N \mathcal{T}(A_{<}, A_{>})} \\
&= C (2\pi c)^{2q} \left(\frac{\pi}{N} \frac{(1-c^2)^2}{\sqrt{8c^2-4}} \right)^q \left(\frac{\pi}{N} \frac{(1-c^2)^2}{\sqrt{1-c^4}} \right)^{2q(q-1)} (1-c^2)^{-4q^2} \\
&\quad \times \exp \left\{ Nq \left(\frac{c^2}{1-c^2} + 2 \log(1-c^2) \right) \right\} \left(1 + \mathcal{O}\left(\frac{1}{N}\right) \right) \\
&= C 2^q c^{2q} \frac{\pi^{2q^2+q}}{N^{2q^2-q}} \left(\frac{1}{\sqrt{2c^2-1}} \right)^q \left(\frac{1}{\sqrt{1-c^4}} \right)^{2q(q-1)} (1-c^2)^{-2q} \\
&\quad \times \exp \left\{ Nq \left(\frac{c^2}{1-c^2} + 2 \log(1-c^2) \right) \right\} \left(1 + \mathcal{O}\left(\frac{1}{N}\right) \right) \\
&= C 2^q c^{2q} \frac{\pi^{2q^2+q}}{N^{2q^2-q}} \left(\frac{1}{\sqrt{2c^2-1}} \right)^q (1-c^4)^{-q^2} \frac{(1-c^4)^q}{(1-c^2)^{2q}} \\
&\quad \times \exp \left\{ Nq \left(\frac{c^2}{1-c^2} + 2 \log(1-c^2) \right) \right\} \left(1 + \mathcal{O}\left(\frac{1}{N}\right) \right)
\end{aligned}$$

with C according to (5.21). We conclude that

$$\begin{aligned}
& \int_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_{>}) \int_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_{<}) e^{\beta N \mathcal{T}(A_{<}, A_{>})} \\
&= (2\pi N)^q c^{2q} \left(\frac{1}{\sqrt{2c^2-1}} \right)^q (1-c^4)^{-q^2} \frac{(1-c^4)^q}{(1-c^2)^{2q}} \\
&\quad \times \exp \left\{ Nq \left(\frac{c^2}{1-c^2} + 2 \log(1-c^2) \right) \right\} \left(1 + \mathcal{O}\left(\frac{1}{N}\right) \right).
\end{aligned}$$

This gives the result. \square

6. COMPUTATION OF THE LOCALIZED REFINED PRE-STATE

6.1. The Saddle Point Asymptotics for the High-Energy Wave Functions.

We now want to extend the construction of the previous section to the situation in four-dimensional Minkowski space. Our goal is to compute the localized refined pre-state as introduced in Section 3.4. We choose sets $\tilde{\Omega} \subset \tilde{M}$ and $\Omega \subset M$ as well as two sets $\tilde{V} \subset \tilde{M}$ and $V \subset M$ of the same finite volume,

$$\rho(V) = \tilde{\rho}(\tilde{V}) < \infty,$$

and consider the partition function (1.1), where the functional $\mathcal{T}_V^t(\tilde{\rho}, T_{\mathcal{U}_{<}, \mathcal{U}_{>}} \rho)$ is given by (2.42). This functional has the advantage that it is relatively easy to compute. Therefore, it is a good starting point for the detailed computation of quantum states.

In order to find the saddle point and to determine their scaling behavior, it suffices to consider the case that both ρ and $\tilde{\rho}$ describe the Minkowski vacuum with a regularization on the scale ε . We again work in finite spatial volume as introduced in Section 3.3. Thus, instead of Minkowski space, as in (3.5) we consider again the spacetime cylinder $\mathcal{M} = \mathbb{R} \times [-L, L]^3$. We consider solutions of the Dirac equation $(i\tilde{\not{D}} - m)\psi = 0$ in

this cylinder with periodic boundary conditions with the usual scalar product

$$(\psi|\phi)_m := \int_{[-L,L]^3} \prec \psi(t, \vec{x}) | \gamma^0 \phi(t, \vec{x}) \succ d^3x .$$

We denote the resulting Hilbert space of Dirac solutions by $(\mathcal{H}_m, (|\cdot\rangle)_m)$. Next we choose $\mathcal{H} = \tilde{\mathcal{H}}$ as the subspace of all negative energy solutions with a cutoff on the scale $1/\varepsilon$, i.e.

$$-\frac{1}{\varepsilon} < \omega < 0 .$$

Denoting the local correlation map by (see also (2.9))

$$F^\varepsilon : \mathcal{M} \rightarrow \mathcal{F} , \quad x \mapsto -\Psi^\varepsilon(x)^* \Psi^\varepsilon(x) ,$$

we choose

$$\tilde{\rho} = \rho = (F^\varepsilon)_* \mu ,$$

where $d\mu = d^4x$ is the volume measure on \mathcal{M} . The local correlation map makes it possible to identify Minkowski space \mathcal{M} with the spacetimes M and \tilde{M} . For convenience, in what follows we shall work in Minkowski space. Next, we choose the sets Ω and $\tilde{\Omega}$ as the past of the Cauchy surface of time zero, i.e.

$$\tilde{\Omega} = \Omega = \{(t, \vec{x}) \in \mathcal{M} \mid t < 0\} .$$

Next, we choose the sets $\tilde{V} = V = [-T, T] \times [-\ell, \ell]^3$ which localize the state again as in (3.7). The wave functions which are localized mainly inside the three-dimensional box $[-\ell, \ell]^3$ can be associated to a subspace $\mathcal{H}^{\text{lab}} \subset \mathcal{H}$ whose dimension has the scaling behavior (3.8). In this setting, the functional \mathcal{T}_V^t can be written as

$$\begin{aligned} \mathcal{T}_V^t(\tilde{\rho}, T_{\mathcal{U}_<, \mathcal{U}_>}) &= \left(\int_{\Omega \cap V} d^4x \int_{\Omega \cap V} d^4y + \int_{(M \setminus \Omega) \cap V} d^4x \int_{(M \setminus \Omega) \cap V} d^4y \right) \\ &\quad \times |F^\varepsilon(x) \mathcal{U}_> F^\varepsilon(y) \mathcal{U}_<^{-1}|^2 . \end{aligned} \quad (6.1)$$

In order to understand the structure of the functional \mathcal{T}_V^t in (6.1), we note that the unitary operators $\mathcal{U}_>$ and $\mathcal{U}_<$ involve many phases, typically leading to destructive interference. If $\mathcal{U}_<$ and $\mathcal{U}_>$ are chosen as a multiple of the identity, then no destructive interference occurs, making the functional \mathcal{T}_V^t large. On the other hand, the set of unitary operators which are close to a multiple of the identity has a very small measure, making the contribution to the partition function smaller. This suggests that, in order to identify the leading contributions to the partition function, one should consider configurations where the unitary operators $\mathcal{U}_<$ and $\mathcal{U}_>$ are close to multiples of the identity on a suitable *subspace* of \mathcal{H}^{lab} , which we denote by \mathcal{H}^{sp} . Using a notation similar to (4.15), we set

$$A_{<,>} := \pi_{\mathcal{H}^{\text{sp}}} \mathcal{U}_{<,>} \big|_{\mathcal{H}^{\text{sp}}} .$$

We expect a saddle point if these matrices are multiples of the identity matrix, e.g.

$$A_< = A_> = c \mathbf{1}_{\mathcal{H}^{\text{sp}}} \quad (6.2)$$

for suitable $c \in \mathbb{R}$. We make the simplifying assumption that, on the orthogonal complement of \mathcal{H}^{sp} , destructive interference does occur, making it possible to restrict attention in the computation to the subspace \mathcal{H}^{sp} . Proceeding in this way, we can compute the group integrals again with saddle point methods. Since there are many possible choices of \mathcal{H}^{sp} , there will also be many saddle points. Our task is to determine their scaling behavior and combinatorics.




FIGURE 3. The subsystems $\mathcal{H}^{\text{sp}} \subset \mathcal{H}^{\text{he}} \subset \mathcal{H}^{\text{lab}}$ in momentum space.

The main step in the quantitative analysis is to compute how the contribution of the saddle point to the partition function depends on the dimension of the subspace \mathcal{H}^{sp} . We begin this analysis by restricting attention to the wave functions with energy on the Planck scale,

$$|\omega| \simeq \frac{1}{\varepsilon}.$$

We also refer to these wave functions as the *high-energy wave functions*. We denote all the high-energy wave functions of our system by

$$\mathcal{H}^{\text{he}} \subset \mathcal{H}^{\text{lab}}.$$

This analysis will make it possible to determine the scaling behavior in ε (the low-energy wave functions will not change this scaling behavior, but they will modify the structure of the saddle points; this will be worked out in Section 6.4 below.)

Next, in order to maximize the contribution to \mathcal{T}_V^t , we choose \mathcal{H}^{sp} as the subset of wave functions whose spatial momenta lie in a cone with opening angle $\vartheta_0 \in [0, \pi)$, i.e.

$$p = (\omega, \vec{k}) \quad \text{with} \quad \vec{k} \in C_{\vartheta_0}(\vec{k}_0) := \{\vec{k} \in \mathbb{R}^3 \mid \angle(\vec{k}, \vec{k}_0) \leq \vartheta_0\}. \quad (6.3)$$

This subset is illustrated in Figure 3. The dimension of \mathcal{H}^{sp} scales like

$$\dim \mathcal{H}^{\text{sp}} \sim \sin^2 \vartheta_0 \dim \mathcal{H}^{\text{he}} \sim \vartheta_0^2 \left(\frac{\ell}{\varepsilon}\right)^3. \quad (6.4)$$

Having chosen \mathcal{H}^{sp} , we next compute the form of the resulting saddle point. To this end, we can use methods and results of the light-cone expansion and the continuum limit analysis (see the preliminaries in Section 2.1.4 and [13]) and combine them with methods introduced in [15] for the computation of surface layer integrals. We begin with the regularized kernel of the fermionic projector in the vacuum, for simplicity with $i\varepsilon$ -regularization,

$$P^\varepsilon(x, y) = \int_{\mathbb{R}^4} \frac{d^4 p}{(2\pi)^4} (\not{p} + m) \delta(p^2 - m^2) \Theta(-p^0) e^{\varepsilon p^0} e^{-ip(x-y)}. \quad (6.5)$$

In the limit $\varepsilon \searrow 0$, this distribution develops singularities on the light cone. Likewise, for small $\varepsilon > 0$, this distribution has large contributions near the light cone. These contributions can be analyzed in detail with the so-called regularized light-cone expansion and the formalism of the continuum limit (for details see the preliminaries in Section 2.1.4 and [13, Sections 2.2. and 2.4] or [28]). Here we are interested only in the scaling behavior in ε , making it possible to work with the following simple argument. Let $\xi := y - x$ be a vector on the light cone (i.e. $\xi \neq 0$ and $\xi^2 = 0$). The Fourier

integral (6.5) involves the oscillating phase factor $\exp(ip\xi)$. The oscillations give rise to destructive interference. This is why the leading contribution to the Fourier integral is obtained simply by restricting attention to those momenta for which $p\xi$ is small. For determining the scaling behavior in ε , it suffices to integrate over the set

$$\Phi(\xi) := \{p \in \mathbb{R}^4 \mid |p\xi| \leq 1\}$$

and leave out the phase factor. We thus obtain

$$P^\varepsilon(x, y) \simeq \int_{\Phi(\xi)} \frac{d^4 p}{(2\pi)^4} (\not{p} + m) \delta(p^2 - m^2) \Theta(-p^0) e^{\varepsilon p^0} (1 + \mathcal{O}(\varepsilon/t)). \quad (6.6)$$

We compute this integral in order to illustrate that this computation is indeed compatible with the scaling obtained in the formalism of the continuum limit. Without loss of generality, we only consider the case that ξ is on the upper light cone, i.e.

$$t := \xi^0 = |\vec{\xi}|.$$

Moreover, we set $p = (\omega, \vec{k})$ and introduce polar coordinates (k, ϑ, φ) with

$$k := |\vec{k}| \quad \text{and} \quad \vec{\xi} \vec{k} = -tk \cos \vartheta$$

(here the minus sign has the advantage that for small ϑ , the vector p is in $\Phi(\xi)$). Then the set $\Phi(\xi)$ is characterized by the inequality

$$-\omega t + k |\vec{\xi}| \cos \vartheta \leq 1.$$

Again restricting attention to large energies, we can set $\omega \approx -k$. Moreover, we can expand for small angles to obtain

$$kt - kt \left(1 - \frac{\vartheta^2}{2}\right) \leq 1 + kt \mathcal{O}(\vartheta^4)$$

and thus

$$\vartheta \leq \hat{\vartheta} (1 + \mathcal{O}(\hat{\vartheta}^2)) \quad \text{with} \quad \hat{\vartheta}^2 := \frac{1}{kt}. \quad (6.7)$$

Using this formula in (6.6) gives

$$\begin{aligned} P^\varepsilon(x, y) &\simeq \int_0^\infty k^2 dk \int_0^{\hat{\vartheta}} \sin \vartheta d\vartheta \frac{1}{|\omega|} (\not{p} + m) e^{\varepsilon \omega} \Big|_{\omega=-k} (1 + \mathcal{O}(\hat{\vartheta}^2)) \\ &\simeq \int_0^\infty k^2 dk \frac{1}{kt} \frac{1}{|\omega|} (\not{p} + m) e^{\varepsilon \omega} \Big|_{\omega=-k} (1 + \mathcal{O}(\hat{\vartheta}^2)) \\ &\simeq \frac{1}{t} \int_0^\infty k e^{-\varepsilon k} dk (1 + \mathcal{O}(\hat{\vartheta}^2)) \simeq \frac{1}{\varepsilon^2 t} (1 + \mathcal{O}(\hat{\vartheta}^2)). \end{aligned}$$

This is compatible with the formalism of the continuum limit, where

$$P^\varepsilon(x, y) \simeq \not{\xi} T^{(-1)} + (\deg < 2) \simeq \frac{\not{\xi}}{\varepsilon^2 t^2} + (\deg < 2) \simeq \frac{1}{\varepsilon^2 t} + (\deg < 2). \quad (6.8)$$

Before going on, we slightly simplify the setup. Using that the leading terms for small ε come from the wave functions of energy $k \sim \omega \sim \varepsilon^{-1}$, we may replace the factor k in the definition of $\hat{\vartheta}$ in (6.7) by $1/\varepsilon$,

$$\hat{\vartheta}^2 := \frac{\varepsilon}{t}. \quad (6.9)$$

This changes the computation of the Fourier integral to

$$\begin{aligned} P^\varepsilon(x, y) &\simeq \int_0^\infty k^2 dk \int_0^{\hat{\vartheta}} \sin \vartheta d\vartheta \frac{1}{|\omega|} (\not{p} + m) e^{\varepsilon\omega} \Big|_{\omega=-k} (1 + \mathcal{O}(\varepsilon/t)) \\ &\simeq \frac{\varepsilon}{t} \int_0^\infty k^2 e^{-\varepsilon k} dk (1 + \mathcal{O}(\varepsilon/t)) \simeq \frac{1}{\varepsilon^2 t} (1 + \mathcal{O}(\varepsilon/t)) , \end{aligned} \quad (6.10)$$

without any effect on the scalings. The choice (6.9) will simplify the following considerations.

The next step is to adapt the above method to the case where we only take into account the wave functions in the subspace $\mathcal{H}^{\text{sp}} \subset \mathcal{H}^{\text{he}}$. This simply corresponds to replacing the integration domain $\Phi(\xi)$ in (6.6) by

$$\Phi_{\text{sp}}(\xi) := \{p \in \mathbb{R}^4 \mid |p\xi| \leq 1 \text{ and } \vec{k} \in C_{\vartheta_0}(\vec{k}_0)\} ,$$

where $C_{\vartheta_0}(\vec{k}_0)$ is again the cone in (6.3). For the computation of the resulting Fourier integral, it is convenient to distinguish the cases where $\hat{\vartheta}$ is larger or smaller than ϑ_0 . This corresponds to the two cases

$$t < t_0 \quad \text{respectively} \quad t \geq t_0$$

with

$$t_0 := \frac{\varepsilon}{\vartheta_0^2} .$$

In the case $t \gg t_0$, the computation (6.10) as well as (6.8) remain valid, but only if $\vec{\xi}$ lies inside the cone around \vec{k}_0 with opening angle ϑ_0 , i.e.

$$P^\varepsilon(x, y) \simeq \frac{g}{\varepsilon^2 t^2} \chi_{C_{\vartheta_0}(\vec{k}_0)}(\vec{\xi}) \quad \text{if } t \gg t_0 .$$

In the opposite case $t \ll t_0$, we get a contribution whenever $\vec{\xi}$ lies inside the cone around \vec{k}_0 with opening angle $\hat{\vartheta}$. In this case, all the vectors in \mathcal{H}^{sp} are “in phase.” However, these are fewer wave functions than in (6.10). Since the number of wave functions scales quadratically in the opening angle of the cone, we obtain

$$P^\varepsilon(x, y) \simeq \frac{g}{\varepsilon^2 t^2} \frac{\vartheta_0^2}{\hat{\vartheta}^2} \chi_{C_{\hat{\vartheta}}(\vec{k}_0)}(\vec{\xi}) \simeq \frac{g}{\varepsilon^2 t t_0} \chi_{C_{\hat{\vartheta}}(\vec{k}_0)}(\vec{\xi}) \quad \text{if } t \ll t_0 . \quad (6.11)$$

We next compute the closed chain $A_{xy} := P^\varepsilon(x, y) P^\varepsilon(y, x)$. In the case $t \gg t_0$, we can use the formalism of the continuum limit to obtain (more precisely, we apply (2.13) with $L = 3$)


$$A_{xy} \simeq \frac{1}{\varepsilon^3 t^3} \chi_{C_{\vartheta_0}(\vec{k}_0)}(\vec{\xi}) \quad \text{if } t \gg t_0 .$$

In the case $t \ll t_0$, however, the kernel of the fermionic projector no longer depends on t (see (6.11); intuitively speaking, this is because all the wave functions of \mathcal{H}^{sp} are “in phase”). Therefore, the closed chain is obtained by taking the closed chain at $t = t_0$,

$$A_{xy} \simeq \frac{1}{\varepsilon^3 t_0^3} \chi_{C_{\vartheta_0}(\vec{k}_0)}(\vec{\xi}) \quad \text{if } t \ll t_0 .$$

In order to determine the scaling behavior, it suffices to interpolate between the asymptotics for large and small t . We thus obtain

$$A_{xy} \simeq \frac{1}{\varepsilon^3 t^3 t_0^3} (1 + \mathcal{O}(\varepsilon/t)) \times \begin{cases} t_0^3 \chi_{C_{\vartheta_0}(\vec{k}_0)}(\vec{\xi}) & \text{if } t \geq t_0 \\ t^3 \chi_{C_{\hat{\vartheta}}(\vec{k}_0)}(\vec{\xi}) & \text{if } t < t_0 . \end{cases}$$

FIGURE 4. The support of the closed chain as a function of ξ .

The support of this function is shown in Figure 4.

Now we can proceed in the formalism of the continuum limit. More precisely, using (2.14), we obtain

$$\begin{aligned} |\lambda_i^{xy}|^2 &\simeq \frac{1}{\varepsilon^6 t_0^6} (1 + \mathcal{O}(\varepsilon/t)) \times \begin{cases} t_0^6 \chi_{C_{\vartheta_0}(\vec{k}_0)}(\vec{\xi}) & \text{if } t \geq t_0 \\ t^6 \chi_{C_{\hat{\vartheta}}(\vec{k}_0)}(\vec{\xi}) & \text{if } t < t_0 \end{cases} \\ &\simeq \frac{1}{\varepsilon^5 t_0^6} \delta(t - |\vec{\xi}|) (1 + \mathcal{O}(\varepsilon/t)) \times \begin{cases} t_0^6 \chi_{C_{\vartheta_0}(\vec{k}_0)}(\vec{\xi}) & \text{if } t \geq t_0 \\ t^6 \chi_{C_{\hat{\vartheta}}(\vec{k}_0)}(\vec{\xi}) & \text{if } t < t_0. \end{cases} \end{aligned}$$

We thus obtain the following contribution to the functional \mathcal{T}_V^t in (6.1) (in the special case $\tilde{\rho} = \rho$ and $\mathcal{U}_< = \mathcal{U}_> = \mathbf{1}$ for the states in \mathcal{H}^{sp}),

$$\begin{aligned} \mathcal{T}_V^t(\rho, \rho) &\simeq T \ell^3 \int_{\mathbb{R}^4} |\lambda_i^{xy}|^2 d^4 y & (6.12) \\ &\simeq T \ell^3 \int_0^\infty r^2 dr \frac{1}{\varepsilon^5 r^6} \frac{1}{t_0^6} (1 + \mathcal{O}(\varepsilon/r)) \times \begin{cases} t_0^6 \vartheta_0^2 & \text{if } r \geq t_0 \\ r^6 \hat{\vartheta}^2 & \text{if } r < t_0 \end{cases} \\ &\simeq T \ell^3 \int_0^\infty r^2 dr \frac{1}{\varepsilon^4 r^6} \frac{1}{t_0^6} (1 + \mathcal{O}(\varepsilon/r)) \times \begin{cases} t_0^5 & \text{if } r \geq t_0 \\ r^5 & \text{if } r < t_0 \end{cases} \\ &\simeq T \ell^3 \frac{1}{\varepsilon^4 t_0^3} \frac{1}{t_0^6} t_0^5 (1 + \mathcal{O}(\vartheta_0^2)) = T \ell^3 \frac{1}{\varepsilon^4 t_0^4} (1 + \mathcal{O}(\vartheta_0^2)) = \frac{T \ell^3}{\varepsilon^8} \vartheta_0^8 (1 + \mathcal{O}(\vartheta_0^2)) \\ &= \frac{T \ell^3}{\varepsilon^8} \left(\frac{\dim \mathcal{H}^{\text{sp}}}{\dim \mathcal{H}^{\text{he}}} \right)^4 (1 + \mathcal{O}(\vartheta_0^2)) \stackrel{(6.4)}{=} \frac{T \ell^3}{\varepsilon^8} (\dim \mathcal{H}^{\text{sp}})^4 \left(\frac{\varepsilon}{\ell} \right)^{12} (1 + \mathcal{O}(\vartheta_0^2)) \\ &= T \frac{\varepsilon^4}{\ell^9} (\dim \mathcal{H}^{\text{sp}})^4 (1 + \mathcal{O}(\vartheta_0^2)). & (6.13) \end{aligned}$$

After these approximations and computations, we are in a setting which is very similar to the one-dimensional example (5.15). In order to see the similarity, we first make the simplifying assumption that there is a basis of \mathcal{H}^{sp} consisting of wave packets having mutually disjoint supports in the spacetime region V . Identifying each wave packet with a point $x \in X$ in (5.9), we are precisely in the setting of the example (5.15) with

$$q = \dim \mathcal{H}^{\text{sp}} \quad \text{and} \quad \beta = \alpha T \frac{\varepsilon^4}{\ell^9} q^3$$

(compare (5.15) and the linear scaling in q of (5.14) with (1.1) and the quartic scaling in q in (6.13)). Clearly, due to dispersion effects and the fact that we only have solutions

of negative energies, there is no basis consisting of wave packets with mutually disjoint supports. Nevertheless, we can apply the result of the model example studied in Section 5.2 in various situations, as we now explain. The simplest method is to choose the width T of the spacetime region V so small that wave packets do not get dispersed in time. Then the remaining error terms are of higher order in $\varepsilon\ell$, εm and $(T\vartheta_0/\varepsilon)^2$ (these error terms will be explained for specific wave packets in Section 6.3 below). In this setting, we can apply Theorem 5.6. Noting that saddle points exist only if β is larger than four, it is most convenient to write β in Theorem 5.6 as

$$\beta = 4 \frac{q^3}{q_{\min}^3},$$

which means that the prefactor α in (1.1) is chosen as

$$\alpha = \frac{\ell^9}{T \varepsilon^4} \frac{4}{q_{\min}^3}. \quad (6.14)$$

Note that α is a constant independent of q of length dimension four. We thus obtain the following result.

Proposition 6.1. *We consider the functional \mathcal{T}_V^t for V a four-dimensional cuboid (3.7) inside a spacetime cylinder (3.5) with*

$$T \lesssim \frac{\varepsilon}{\vartheta_0^{\frac{5}{3}}}. \quad (6.15)$$

Then the subsystem formed by the wave functions \mathcal{H}^{sp} inside a cone of opening angle ϑ_0 (see (6.3)) gives rise to the saddle point contribution

$$\begin{aligned} & \int_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_{>}) \int_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_{<}) e^{\alpha N \mathcal{T}_V^t(\rho, \rho)} \asymp (2\pi N)^q (1 - c^4)^{-q^2} \left(\frac{1 + c^2}{1 - c^2} \frac{c^2}{\sqrt{2c^2 - 1}} \right)^q \\ & \times \exp \left\{ N q \left(\frac{c^2}{1 - c^2} + 2 \log(1 - c^2) \right) \right\} \left(1 + \mathcal{O}\left(\frac{1}{N}\right) + \mathcal{O}\left(\frac{T^2}{\varepsilon^2} \vartheta_0^{\frac{10}{3}}\right) \right), \end{aligned} \quad (6.16)$$

provided that $q > q_{\min}$. Here α is chosen according to (6.14) and

$$N = \dim \mathcal{H} \sim \left(\frac{L}{\varepsilon} \right)^3 \quad (6.17)$$

$$q = \dim \mathcal{H}^{\text{sp}} \sim \vartheta_0^2 \left(\frac{\ell}{\varepsilon} \right)^3 \quad (6.18)$$

$$c(q)^2 = \frac{1}{2} + \frac{1}{2} \sqrt{1 - \frac{q_{\min}^3}{q^3}}. \quad (6.19)$$

Proof. The previous computation gives (6.16), except for the last error term. In order to determine the scaling of this error term, we form q wave packets localized evenly in the box of size ℓ . According to (6.18), the Euclidean distance d between the wave packets scales like $d \simeq \varepsilon \vartheta_0^{-2/3}$. The wave packets disperse with an opening angle ϑ_0 (see Figure 3). In order for this effect to be negligible, we need to assume that $d \gtrsim T \sin \vartheta_0 \simeq T \vartheta_0$, giving (6.15). The resulting overlap of the wave packets is integrated over a set of dimension at least two. Therefore, we may square the corresponding error term. This gives the result. \square

We point out that the last error term in (6.16) makes it necessary to choose T very small, as quantified by (6.15). This does not cause any problems, because at this stage we are free to choose T arbitrarily. Nevertheless, in order to model a realistic measurement device, it may be preferable to choose T on the Compton scale or even larger. Before explaining how to treat this situation, we briefly discuss what the result of Proposition 6.1 means for the refined localized state (1.1). In this proposition we computed the contribution of one saddle point corresponding to the subsystem \mathcal{H}^{sp} . Clearly, there are many possible choices for \mathcal{H}^{sp} , giving rise to many saddle points. We do not need to work out the combinatorics of these saddle points, because the relative size of the different contributions can be determined with the following scaling argument. We consider the asymptotics for small ε in the limit when N tends to infinity. The combinatorics of the saddle points depend on ℓ and ε , but it is independent of the size L of the spacetime cylinder. Therefore, the asymptotics for large N can be studied by taking the limit $L \rightarrow \infty$ (see (6.17)). Thus the leading contributions to the refined localized state are obtained by those subsystems for which the function in the exponent

$$h(q) := q \left(\frac{c(q)^2}{1 - c(q)^2} + 2 \log(1 - c(q)^2) \right) \quad (6.20)$$

with $c(q)$ according to (6.19) is largest.

Lemma 6.2. *The function $h(q)$ is strictly monotone increasing for $q \geq q_{\min}$.*

Proof. A direct computation yields

$$\begin{aligned} h'(q_{\min}) &= 4 - 2 \log 2 > 0 \\ h''(q) &= \frac{3(2x - 2\sqrt{x}\sqrt{1+x} + 3)}{q_{\min}\sqrt{x}(1+x)^{\frac{1}{3}}(\sqrt{1+x} - \sqrt{x})^3} \quad \text{for } q > q_{\min}, \end{aligned}$$

where we parametrized q in terms of $x > 0$ by

$$q = q_{\min}(1+x)^{\frac{1}{3}}.$$

Clearly, the denominator of our formula for $h''(q)$ is positive. The same is true for the numerator in view of the Cauchy-Schwarz inequality

$$2\sqrt{x}\sqrt{1+x} \leq (\sqrt{x})^2 + (\sqrt{1+x})^2 = 2x + 2 < 2x + 3.$$

Therefore, the function h is convex on (q_{\min}, ∞) . Using that $h'(q_{\min})$ is strictly positive, we conclude that h is strictly monotone on $[q_{\min}, \infty)$. \square

This lemma shows that in the considered limiting case $N \rightarrow \infty$, the main contribution to the dominant saddle points come from the large systems. The largest possible system is that formed of all wave functions in \mathcal{H}^{he} , i.e.

$$\mathcal{H}^{\text{sp}} = \mathcal{H}^{\text{he}} \quad \text{and} \quad \vartheta_0 = 2\pi.$$

We finally explain how one can relax the condition (6.15). If this condition is violated, we cannot choose a basis of \mathcal{H}^{sp} consisting of wave packets whose supports are mutually disjoint in V up to small error terms. Instead, these wave packets will have substantial overlaps due to *dispersion*. The main observation is that this has no effect on the exponential factor in (6.16), but it may change the prefactors.

Corollary 6.3. *Consider the setting of Proposition 6.1, but now without assuming (6.15). Then the subsystem formed by the wave functions \mathcal{H}^{sp} inside a cone of opening angle ϑ_0 (see (6.3)) gives rise to the saddle point contribution*

$$\int_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_{>}) \int_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_{<}) e^{\alpha N \mathcal{T}_V^t(\rho, \rho)} \asymp \exp \left\{ N h(q) + \mathcal{O}(\log N) \right\},$$

provided that $q > q_{\min}$, where α , q , c and $h(q)$ are again given by (6.14), (6.17)–(6.19) and (6.20).

Proof. We denote the localized wave packets by e_x with $x \in X := \{1, \dots, q\}$. Compared to the model example (5.12), we need to take into account that for every x , also other waves e_y with $y \neq x$ contribute. In order to describe this situation in the most general setting, we consider a functional \mathcal{T}^{dis} of the general form

$$\mathcal{T}^{\text{dis}}(A_{<}, A_{>}) = \sum_{x \in X} \sum_{\alpha=1}^{p_x} \left| \text{tr}_I (\mathcal{U}_{>} \Psi(x)^* \Psi(x) \mathcal{U}_{<}^{-1} P_{x,\alpha}) \right|^2 \quad (6.21)$$

with parameters $p_1, \dots, p_q \geq 0$ and positive operators $P_{x,1}, \dots, P_{x,p_x}$ on I . These operators can be thought of as being non-zero on all wave packets which have a substantial overlap with e_x . Since the wave function e_x clearly has an overlap, we may assume that the new functional bounds the original functional from below, i.e.

$$\mathcal{T}(A_{<}, A_{>}) \leq \mathcal{T}^{\text{dis}}(A_{<}, A_{>}) \quad \text{for all } A_{<}, A_{>} \in \mathcal{A}_q \quad (6.22)$$

(and \mathcal{A}_q as introduced in (4.8)).

Similar to (5.14), we write (6.21) as

$$\mathcal{T}^{\text{dis}}(A_{<}, A_{>}) = \sum_{x \in X} \sum_{\alpha=1}^{p_x} \left| \text{tr}_I (A_{>} \pi_x A_{<}^* P_{x,\alpha}) \right|^2$$

(where π_x is the orthogonal projection to the subspace spanned by e_x). Given $\mathcal{U}_{<}$ and $\mathcal{U}_{>}$, we now consider the integral over all matrices obtained by multiplying from the left by unitary operators acting on $U(I)$,

$$\int_{U(I)} d\mu_{U(I)}(V_{<}) \int_{U(I)} d\mu_{U(I)}(V_{>}) e^{\alpha N \mathcal{T}^{\text{dis}}(V_{<} A_{<}, V_{>} A_{>})} \quad (6.23)$$

(here $d\mu_{U(I)}$ denotes again the normalized Haar measure). Integrating subsequently over the whole group $\mathcal{G} \times \mathcal{G}$ gives the partition function.

In order to estimate (6.23), we first write the functional $\mathcal{T}^{\text{dis}}(V_{<} A_{<}, V_{>} A_{>})$ as

$$\begin{aligned} \mathcal{T}^{\text{dis}}(V_{<} A_{<}, V_{>} A_{>}) &= \sum_{x \in X} \sum_{\alpha=1}^{p_x} \left| \text{tr}_I (V_{>} A_{>} \pi_x A_{<}^* V_{<}^* P_{x,\alpha}) \right|^2 \\ &= \sum_{x \in X} \sum_{\alpha=1}^{p_x} \left| \langle e_x | A_{<}^* V_{<}^* P_{x,\alpha} V_{>} A_{>} e_x \rangle \right|^2. \end{aligned}$$

For any x and α , we let u be a normalized eigenvector of $P_{x,\alpha}$ corresponding to the largest eigenvalue. We choose $V_{>}$ such that it maps the vector $A_{>} e_x$ to a multiple of u . Likewise, we choose $V_{<}$ such that it maps the vector $A_{<} e_x$ to a multiple of u . Then

$$\left| \langle e_x | A_{<}^* V_{<}^* P_{x,\alpha} V_{>} A_{>} e_x \rangle \right| = \|A_{>} e_x\| \|A_{<}^* e_x\| \|P_{x,\alpha}\|.$$

Using continuity and the Schwarz inequality, we conclude that the double integral (6.23) can be estimated from above by the corresponding integral with \mathcal{T}^{dis} replaced by our original functional \mathcal{T} in (5.14). The point is that the constant involved in this estimate depends only on q , but not on N (this could be worked out in more detail with a covering argument, using that the set \mathcal{A}_q in (4.8) is relatively compact). Now we can compute the saddle point contribution for the upper bound. Moreover, (6.22) gives a lower bound for the saddle point contribution. Clearly, the above estimates change the q -dependent prefactors in (6.16). The exponential factor in (6.16), however, remains unchanged, because it is determined solely by the value of the integrand at the saddle point (see (6.13)) and the Haar measure on \mathcal{G} . \square

6.2. Introducing a Weight for the Size of the Subsystems. We now explain a method which allows us to introduce a weight for the size of the subsystems. In particular, this makes it possible to get finer information on the smaller subsystems formed by wave functions pointing in different spatial directions. To this end, we choose a bounded cutoff function η supported in a neighborhood of the origin, i.e.

$$\eta \in L_0^\infty([0, 1], \mathbb{R}^+) \quad \text{with} \quad \eta|_{[0, \delta]} \equiv 1$$

for some $\delta > 0$. We now modify the functional \mathcal{T}_V^t in (6.1) by inserting the cutoff function

$$\eta\left(\frac{\|xy\|}{\|x\|\|y\|}\right) \tag{6.24}$$

into the integrand (here $\|\cdot\|$ is the operator norm on $L(\mathcal{H})$),

$$\begin{aligned} \mathcal{T}_V^t(\tilde{\rho}, T_{\mathcal{U}_<, \mathcal{U}_> \rho}) = & \left(\int_{\Omega \cap V} d^4x \int_{\Omega \cap V} d^4y + \int_{(M \setminus \Omega) \cap V} d^4x \int_{(M \setminus \Omega) \cap V} d^4y \right) \\ & \times \eta\left(\frac{\|xy\|}{\|x\|\|y\|}\right) |F^\varepsilon(x) \mathcal{U}_> F^\varepsilon(y) \mathcal{U}_<^{-1}|^2. \end{aligned} \tag{6.25}$$

As a result, the contribution for small ξ in Figure 4 is suppressed. In order to clarify the effect of the cutoff function on the scaling behavior, we consider the particularly simple choice of a Heaviside function, i.e.


$$\eta(\tau) = \Theta(\tau_0 - \tau) \tag{6.26}$$

with a dimensionless constant $\tau_0 > 0$. As a consequence, in (6.12) we integrate only over the region

$$|\xi^0| > \underline{t}, \tag{6.27}$$

where \underline{t} is a function of τ_0 with the scaling

$$\underline{t} \simeq \frac{\varepsilon}{\tau_0}. \tag{6.28}$$

FIGURE 5. The function $h(q)$ for η a sharp cutoff function.

Therefore, we can choose \underline{t} arbitrary by a suitable choice of the cutoff function. This changes the computation of \mathcal{T}_V^t after (6.12) to

$$\begin{aligned}
 \mathcal{T}_V^t(\rho, \rho) &\simeq T\ell^3 \int_{\mathbb{R}^4} |\lambda_i^{xy}|^2 d^4y \\
 &\simeq T\ell^3 \int_{\underline{t}}^{\infty} r^2 dr \frac{1}{\varepsilon^4 r^6} \frac{1}{t_0^6} (1 + \mathcal{O}(\varepsilon/r)) \times \begin{cases} t_0^5 & \text{if } r \geq t_0 \\ r^5 & \text{if } r < t_0 \end{cases} \\
 &\simeq \frac{T\ell^3}{\varepsilon^4} \frac{1}{t_0 \max(t_0, \underline{t})^3} \left(1 + \mathcal{O}\left(\frac{\varepsilon}{\max(t_0, \underline{t})}\right)\right) \\
 &= \frac{T\ell^3}{\varepsilon^8} \vartheta_0^2 \min(\vartheta_0^2, \underline{\vartheta}^2)^3 \left(1 + \mathcal{O}\left(\min(\vartheta_0^2, \underline{\vartheta}^2)\right)\right) \\
 &= T \frac{\varepsilon^4}{\ell^9} q \min(q, \underline{q})^3 \left(1 + \mathcal{O}\left(\min(\vartheta_0^2, \underline{\vartheta}^2)\right)\right),
 \end{aligned}$$

where we used the notation

$$\underline{\vartheta}^2 := \frac{\varepsilon}{\underline{t}} \quad \text{and} \quad \underline{q} := \underline{\vartheta}^2 \left(\frac{\ell}{\varepsilon}\right)^3 = \frac{\varepsilon}{\underline{t}} \left(\frac{\ell}{\varepsilon}\right)^3. \quad (6.29)$$

Thus for $q > \underline{q}$, the functional \mathcal{T}_V^t scales only linearly in q . This can be taken into account simply by modifying the function $c(q)$ in (6.19) to

$$c(q)^2 = \frac{1}{2} + \frac{1}{2} \sqrt{1 - \frac{q_{\min}^3}{\min(q, \underline{q})^3}}. \quad (6.30)$$

On the interval $[q_{\min}, \underline{q}]$, the function $h(q)$ is unchanged. In particular, we know from (6.2) that this function is strictly monotone increasing. For $q > \underline{q}$, however, $h(q)$ simply is a linear function. Its monotonicity properties are determined by the slope

$$\frac{c(\underline{q})^2}{1 - c(\underline{q})^2} + 2 \log(1 - c(\underline{q})^2).$$

If this slope is arranged to be negative, then the function $h(q)$ has a maximum at \underline{q} (see Figure 5). If this is done, the systems with $q = \underline{q}$ give the main contribution to the saddle points.

More generally, with the cutoff function η in (6.24) one can introduce an arbitrary weight function which determines to what extent subsystems of a certain size contribute to the quantum state. We finally give a possible interpretation for the parameter \underline{t} .

Remark 6.4. (potential significance of \underline{t}) In the above construction, the parameter τ_0 in (6.26), and consequently also the parameter \underline{t} in (6.28), were introduced

ad-hoc as a free parameter which can be used in order to introduce a weight for the size of the subsystems forming the saddle points. Ultimately, it might be preferable conceptually to replace the ad-hoc functional (6.25) by another functional which arises more naturally in the theory of causal fermion systems. The most obvious candidate would be the functional (3.10) involving the Lagrangian mentioned in Section 3.4. Indeed, the Lagrangian involves a natural cutoff for small distances (6.27) which comes from the fact that the formalism of the continuum limit applies only away from the origin (i.e. if $|\xi^0|$ is much larger than the Planck scale; for details see [10, Chapter 4] or [13, Section 2.4]). It seems a good idea to identify the corresponding length scale with \underline{t} . Here we shall not enter the details because, as mentioned in Section 3.4, the computation of the Lagrangian is more subtle due to cancellations of summands in (2.1). \diamond

6.3. The Phase Freedom of the Saddle Points. For the model example in Section 5.2, the saddle point is unique only up to the phase freedom described by the functions $\phi^1_{<}, \dots, \phi^q_{<}$ and $\phi^1_{>}, \dots, \phi^q_{>}$ in (5.16). In Section 6.1 we saw that, in a suitable approximation where we took into account only the leading order in $\varepsilon/\ell_{\text{macro}}$, the high-energy wave functions in Minkowski space could be described by this model example. It remains to clarify how the phase freedom in (5.16) translates to the saddle point in Minkowski space. As we will now explain, this saddle point has indeed a phase freedom, but only if we restrict attention to the leading order in $\varepsilon/\ell_{\text{macro}}$. This phase freedom is in general broken if

$$\text{higher orders in } \frac{\varepsilon}{\ell} \quad \text{and} \quad m\varepsilon$$

are taken into account. This means more concretely that all the computations of Sections 6.1 and 6.2 do involve the phase freedom, but the error terms do not.

In order to derive this result, we decompose the system $[-\ell, \ell]^3 \subset \mathbb{R}^3$ describing our laboratory into strips in x -direction of width d with the scaling

$$\varepsilon \ll d \lesssim \ell.$$

We consider solutions of the scalar wave equation $\square\phi = 0$ which do not depend on the coordinates y and z . Then ϕ is a solution of the two-dimensional wave equation

$$(\partial_t^2 - \partial_x^2)\phi = 0.$$

These solutions can be decomposed into the left- and right-moving solutions. We consider for example the right-moving solutions $\phi(t-x)$. We multiply the initial data by phase factors $e^{i\varphi_k}$ and solve the Cauchy problem. In this way, the solution is multiplied in each right-moving strip by a corresponding phase factor; see Figure 6. Clearly, we are interested in solutions of the Dirac equation. They are obtained by acting with the operator $i\partial\!\!\!/$, without changing the picture of the right-moving waves propagating independently in the right-moving time strips. But we obtain error terms for the following reasons:

- (I) The waves must be formed of wave functions contained in \mathcal{H}^{sp} . To this end, we choose a smooth cutoff function $\eta(k)$ which vanishes outside \mathcal{H}^{sp} (i.e. the space of high-energy wave functions inside the cone with opening angle ϑ_0 ; see Figure 3). This cutoff function varies on the scale ε^{-1} . Therefore, multiplying by this cutoff function in momentum spaces gives rise to a convolution in position space with a kernel which decays on the regularization scale ε , as is indicated in Figure 6 for




FIGURE 6. Phase freedom of the saddle points.

the solutions in the time strip with phase $e^{i\varphi^3}$ by the dark shaded regions near the boundary of the strip.

- (II) The non-zero mass m gives rise to contributions which violate the strong Huygens principle and propagate into other time strips (see the light gray region in Figure 6).

In order to describe the waves in the different strips, we introduce symmetric linear operators E_k acting on \mathcal{H}^{lab} . Disregarding the convolution in position space (I), these operators could be chosen as orthogonal projection operators. Taking these convolutions into account, we can nevertheless choose the E_k as orthogonal projection operators outside the transition regions of size $\sim \epsilon$. Next, in analogy to (5.16) we introduce the unitary operators

$$A_{<} = c \exp \left(i \sum_k \varphi_{<}^k E_k \right) \quad \text{and} \quad A_{>} = c \exp \left(i \sum_k \varphi_{>}^k E_k \right)$$

(with c as in (6.19) or (6.30)).

Let us consider how the phases $\varphi_{<}^k$ and $\varphi_{>}^k$ enter the functional $\mathcal{T}_V^t(\tilde{\rho}, T_{\mathcal{U}_{<}, \mathcal{U}_{>}} \rho)$. Clearly, if both x and y lie inside the same time strip, then the phases drop out. Therefore, we only need to consider the case that x and y are in different time strips. The contribution of the error terms (I) from the convolution by the cutoff function η to the integrand in (6.1) are not small. But they vanish unless y lies in a strip of width $\sim \epsilon$. Therefore, after integrating over y , the resulting contribution to $\mathcal{T}_V^t(\tilde{\rho}, T_{\mathcal{U}_{<}, \mathcal{U}_{>}} \rho)$ is of higher order in ϵ/ℓ . The error terms (II) due to the rest mass, on the other hand, do not affect the leading degree of the singularity on the light cone (as defined before (2.13)). Therefore, their contribution to $\mathcal{T}_V^t(\tilde{\rho}, T_{\mathcal{U}_{<}, \mathcal{U}_{>}} \rho)$ involves a scaling factor $m\epsilon$. This proves the claim.

6.4. Saddle Points for the Low-Energy Wave Functions. In the previous sections (Sections 6.1 and (6.2)) we saw that the high-energy wave functions give rise to saddle points. If $\mathcal{U}_{<}$ and $\mathcal{U}_{>}$ are near these saddle points, the surface layer integrals can be computed using the formalism of the continuum limit; in particular, they have a well-defined scaling in negative powers of ϵ . By introducing a parameter \underline{t} with the dimension of length into the functional \mathcal{T}_V^t , we could prescribe the number of wave functions forming the saddle points. This procedure leads to many saddle points, which come with a certain combinatorics. Before entering this combinatorics

(see Section 6.5 below), we now consider a single saddle point and analyze how it is modified by the low-energy wave functions.

In the following consideration, we may disregard the space $(\mathcal{H}^{\text{sp}})^\perp \cap \mathcal{H}^{\text{he}}$ formed of high-energy wave functions which are not part of the saddle point (because if they had an effect, we would have included them in \mathcal{H}^{sp}). Therefore, it remains to consider the subspace

$$\mathcal{H}^{\text{sp}} \oplus \mathcal{H}^{\text{le}} \subset \mathcal{H}^{\text{lab}}. \quad (6.31)$$

We denote the restriction of a unitary operator \mathcal{U} to this subspace by \hat{A} . Using a block matrix notation in the decomposition (6.31), this matrix has the form

$$\hat{A} = \begin{pmatrix} A & * \\ * & * \end{pmatrix}, \quad (6.32)$$

where $A : \mathcal{H}^{\text{sp}} \rightarrow \mathcal{H}^{\text{sp}}$ is the matrix used in the computations of Sections 6.1 and 6.2, and the stars denote the other (still undetermined) matrix entries. The saddle point of the high-energy wave functions is again characterized by (6.2). Being close to this saddle point means that the functional \mathcal{T}_V^t can be computed using the formalism of the continuum limit, exactly as explained in Section 6.1. Therefore, we can expand this functional in powers of individual matrix entries in (6.32). Clearly, expanding in matrix elements of A gives us back the expansion of Lemma 5.8. But we can also expand in the matrix elements of the “starred” block matrices in (6.32). Similar to (2.11), the first variation of the integrand in (2.42) can be expressed with a trace involving the variation of the kernel of the fermionic projector, i.e.

$$\delta(|x \mathcal{U}_> y \mathcal{U}_<^{-1}|^2) = 2 \operatorname{Re} \operatorname{Tr}_{S_y} \left(R_<(y, x) \delta P_<(x, y) + R_>(y, x) \delta P_>(x, y) \right) \quad (6.33)$$

with integral kernels $R_<(y, x)$ and $R_>(y, x)$. Here the kernel of the fermionic projector with index $>$ and $<$ is defined in analogy of (2.10) by

$$P_>(x, y) = -\Psi(x) \mathcal{U}_> \Psi(y)^*, \quad P_<(x, y) = -\Psi(x) \mathcal{U}_< \Psi(y)^*.$$

The kernels $R_<(y, x)$ and $R_>(y, x)$ have singularities on the light cone which are generated by the wave functions in \mathcal{H}^{sp} and thus depend on A alone. The variations δP , on the other hand, are formed of individual wave functions which are bounded in space-time. Therefore, an expansion in powers of δP amounts to an expansion in orders on the light cone (see the preliminaries at the end of Section 2.1.4). After carrying out the integrals in (2.42), we thus obtain an expansion of the form

$$\Delta \mathcal{T}_V^t(\tilde{\rho}, T_{\mathcal{U}_<, \mathcal{U}_>} \rho) = \operatorname{Re} \Phi_<(\delta P_<) + \operatorname{Re} \Phi_>(\delta P_>) + (\text{higher orders in } \varepsilon/\ell_{\text{macro}}), \quad (6.34)$$

where $\Phi_<$ and $\Phi_>$ are complex-linear functionals, and ℓ_{macro} denotes the length scale characterizing the low-energy wave functions. Note that $\Phi_<$ and $\Phi_>$ are determined by the saddle point computation in Section 6.1.

Varying more specifically the unitary operators on $\mathcal{H}^{\text{sp}} \oplus \mathcal{H}^{\text{le}}$, we obtain the formula

$$\begin{aligned} \mathcal{T}_V^t(\tilde{\rho}, T_{\mathcal{U}_<, \mathcal{U}_>} \rho) &= \mathcal{T}_V^t(c\mathbf{1}) + \operatorname{Re} \operatorname{Tr}(\delta \hat{A}_< Y_<) + \operatorname{Re} \operatorname{Tr}(\delta \hat{A}_> Y_>) \\ &\quad + (\text{higher orders in } \varepsilon/\ell_{\text{macro}}), \end{aligned} \quad (6.35)$$

for suitable symmetric matrices $Y_<$ and $Y_>$, where $\mathcal{T}_V^t(c\mathbf{1})$ denotes the functional at the saddle point. Since first variations of $A_<$ and $A_>$ (where we again use the block

matrix form (6.32)) give us back the results of Lemma 5.8, we know that $Y_{<}$ and $Y_{>}$ have the block matrix form

$$Y_{<,>} = \begin{pmatrix} d_{<,>} \mathbf{1} & * \\ * & * \end{pmatrix},$$

where the parameters $d_{<,>}$ describe the first variation of \mathcal{T} as computed in (5.38).

Next, we diagonalize $Y_{<}$ and $Y_{>}$ by a unitary transformation. This unitary transformation can be regarded as a redefinition of the subspaces \mathcal{H}^{le} and \mathcal{H}^{sp} . With this in mind, for ease in notation we do not write out the unitary transformation, but simply replace $Y_{<}$ and $Y_{>}$ by the unitarily transformed matrices. Using that the dimension of \mathcal{H}^{le} is independent of ε , whereas the dimension of \mathcal{H}^{sp} is $\sim \varepsilon^{-3}$, we conclude that, to leading order in $\varepsilon/\ell_{\text{macro}}$, this diagonalization does not change the upper left block matrix entries, i.e.

$$Y_{<,>} = \begin{pmatrix} d_{<,>} \mathbf{1} & 0 \\ 0 & Z_{<,>} \end{pmatrix} + (\text{higher orders in } \varepsilon/\ell_{\text{macro}}) \quad (6.36)$$

with Z a symmetric operator on \mathcal{H}^{le} . Here the error term can be understood in more detail if one proceeds in the following two steps. In the first step, one chooses an orthonormal basis of \mathcal{H}^{sp} such that the matrices $Y_{<}$ and $Y_{>}$ take the form

$$Y_{<,>} = \begin{pmatrix} d_{<,>} \mathbf{1} & 0 & 0 \\ 0 & d_{<,>} \mathbf{1} & * \\ 0 & * & * \end{pmatrix},$$

where we decomposed the first block component into two blocks, where the second block refers to a subspace of \mathcal{H}^{sp} of dimension $d := \dim \mathcal{H}^{\text{le}}$. In the second step we diagonalize the lower $(2d \times 2d)$ -block. This diagonalization redefines \mathcal{H}^{sp} . Consequently, it also changes the computation of the saddle point of the high-energy wave functions. More precisely, we change at most $2d$ physical wave functions. Since each change of physical wave function can be described perturbatively as explained after (6.34) above, one sees that it only affects the higher orders in $\varepsilon/\ell_{\text{macro}}$. Using that the dimension d is independent of ε , we obtain the desired error term in (6.36).

After these preparations, the exponential in (1.1) can be written as a product of a function depending only on $A_{<}$ and $A_{>}$, a function of $\pi_{\mathcal{H}^{\text{le}}} \hat{A}_{<} \pi_{\mathcal{H}^{\text{le}}}$ and a function of $\pi_{\mathcal{H}^{\text{le}}} \hat{A}_{>} \pi_{\mathcal{H}^{\text{le}}}$. Consequently, the $\mathcal{U}_{<}$ -integral can be analyzed with the help of Proposition 4.12 if we set

$$A = A_{<} \quad \text{and} \quad D = \pi_{\mathcal{H}^{\text{le}}} \hat{A}_{<} \pi_{\mathcal{H}^{\text{le}}}.$$

Now we argue as follows. Using that

$$A^* A \otimes D^* D \leq A^* A \otimes \mathbf{1}_{\mathbb{C}^p},$$

the factor in the denominator in (4.30) can be estimated by

$$\det(\mathbf{1} - A^* A)^p \leq \det(\mathbf{1} - A^* A \otimes D^* D) \leq 1.$$

This shows that we can again apply the saddle point computation of Section 5.2. An additional factor $\det(\mathbf{1} - A^* A)^{-p}$ in the integrand changes the formula (5.18) in Theorem 5.6 only by a prefactor $(1 - c^2)^{-p}$, which is independent of N . Therefore, independent of the choice of D , we get a saddle point in the high-energy region, exactly as explained in Sections 6.1 and 6.2. At the saddle point, we know that

$$\det(\mathbf{1} - A^* A \otimes D^* D) = \det(\mathbf{1} - c^2 D^* D)^q$$

with the parameter c given by (5.17), being bounded below by $1/2$ and bounded from above away from one. Therefore, we can again apply Proposition 4.12 to conclude that the saddle point for the low-energy wave functions can be described by applying Proposition 5.1. The denominator in (4.30) simply gives rise to the prefactor

$$\frac{1}{\det(\mathbb{1} - c^2 A_0^2)^q}$$

with A_0 as in (5.1) and $Y = Z_{<,>}$.

This consideration shows that, taking the low-energy wave functions into account, the saddle points can be computed as follows: We first compute the saddle points of the high-energy wave functions as described in Sections 6.1 and (6.2). For each of the resulting high-energy saddle points, we obtain a unique corresponding saddle point for the low-energy wave functions, which can be computed with the help of Proposition 5.1. Due to the linear expansion in (6.33), the dependence on the low-energy wave functions decouples into a product, making it possible to integrate separately over $\pi_{\mathcal{H}^{\text{le}}} \mathcal{U}_{<} \pi_{\mathcal{H}^{\text{le}}}$ and $\pi_{\mathcal{H}^{\text{le}}} \mathcal{U}_{>} \pi_{\mathcal{H}^{\text{le}}}$.

6.5. The Structure of the Saddle Points. In the previous sections, we located and computed various saddle points. We now analyze the saddle points systematically and work out the combinatorics.

The saddle points for the high-energy wave functions were computed starting from a subspace $\mathcal{H}^{\text{sp}} \subset \mathcal{H}^{\text{lab}}$. Clearly, there are many possible choices for this subspace, which can be described by a combinatorics depending on the dimensions of \mathcal{H}^{lab} and \mathcal{H}^{sp} . We point out that this combinatorics is independent of N . This observation is very helpful because, as computed in Proposition 6.1, the contribution by each saddle point involves the exponential factor

$$\exp \{N h(q)\} \tag{6.37}$$

with $h(q)$ defined by (6.20). This shows that, asymptotically for large N , those saddle points will be dominant for which the function h is maximal. The combinatorics describing the possible choices of the subspace $\mathcal{H}^{\text{sp}} \subset \mathcal{H}^{\text{lab}}$ only gives prefactors which are independent of N and will therefore be dominated by the exponential (6.37). In the setting of Section 6.1, we saw in Lemma 6.2 that the function $h(q)$ is monotone increasing. Therefore, as already explained after the statement of Proposition 6.1, in this setting it suffices to take into account one saddle point corresponding to choosing $\mathcal{H}^{\text{sp}} = \mathcal{H}^{\text{lab}}$ formed by all high-energy wave functions. In the setting of Section 6.2, on the other hand, we arranged that the function $h(q)$ has one maximum at $q = \underline{q}$ (see Figure 5). Consequently, we need to take into account all the saddle points involving approximately \underline{q} wave functions. The combinatorics of these saddle points simply corresponds to the possible choices of cones of opening angle $\underline{\vartheta}$ defined by (6.29). For our purposes, it is unnecessary to work out the detailed combinatorics of these saddle points. Instead, we simply label them by an index $a \in \mathfrak{S}$, where \mathfrak{S} is an abstract index set. For notational simplicity, we treat \mathfrak{S} as a discrete set and sum over $a \in \mathfrak{S}$ with weights $c_a \geq 0$.

An important point for what follows is that the saddle point of the model example in Section 5.2 is not unique, but involves the freedom in choosing the phases $\phi_{<}^1, \dots, \phi_{<}^q$ and $\phi_{>}^1, \dots, \phi_{>}^q$ in (5.16). In this way, every a does not stand for a single saddle point, but instead for a whole family of saddle points. For a convenient notation, we denote

the phases by $\phi_\alpha^<$ (for *bra*) and $\phi_\beta^>$ (for *ket*), i.e.

$$e^{i\varphi_\alpha^<} = \phi_\alpha^< \quad \text{and} \quad e^{i\varphi_\beta^>} = \phi_\beta^> \quad \text{with} \quad \alpha, \beta \in \{1, \dots, q\}.$$

As explained in detail in Section 6.3, there is also a phase freedom for the saddle points formed of the high-energy wave functions in Minkowski space, which can be expressed as the freedom in performing transformations of the form

$$P_<(x, y) \rightarrow e^{i\varphi_\alpha^<(x, y)} P_<(x, y) \quad \text{and} \quad P_>(x, y) \rightarrow e^{i\varphi_\beta^>(x, y)} P_>(x, y) \quad (6.38)$$

for suitable functions $\varphi_\alpha^<$ and $\varphi_\beta^>$. As the details are not needed later on, we simply work with these phases abstractly and label them by indices $\alpha, \beta \in \mathfrak{T}_a$, where \mathfrak{T}_a is any index set which may depend on $a \in \mathfrak{S}$. For notational simplicity, we again treat \mathfrak{T} as a discrete set and sum over $\alpha, \beta \in \mathfrak{T}_a$. In cases when \mathfrak{T}_a is continuous, the sum must be replaced by corresponding integrals over phases.

We now come to the saddle points for the low-energy wave functions. We saw in Section 6.4 that these saddle points can be described with the help of Proposition 5.1, where A_0 is determined by the saddle point (a, α, β) (with $a \in \mathfrak{S}$ and $\alpha, \beta \in \mathfrak{T}_a$) of the high-energy wave functions. Proposition 5.1 gives a unique saddle point. Moreover, from (5.3) we know that the contribution by the saddle point in (5.2) grows exponentially in N . Therefore, for large N , it suffices to consider the contribution at this unique saddle point. The interesting issue is the dependence on α and β . Clearly, the phase factors in (6.38) drop out of the integrand in (2.42) when taking the spectral weight. However, being created by unitary transformations acting on the high-energy wave functions, these phases do not appear in the variations of the low-energy wave functions, i.e.

$$\delta P_<(x, y) \rightarrow \delta P_<(x, y) \quad \text{and} \quad \delta P_>(x, y) \rightarrow \delta P_>(x, y),$$

where we vary $A_>$ and $A_<$. As a consequence, the first variation of the integrand in (2.42) does involve the inverse phases. More precisely, (6.33) transforms to

$$2 \operatorname{Re} \operatorname{Tr}_{S_x} \left(R_<(y, x) e^{-i\varphi_\alpha^<(x, y)} \delta P_<(x, y) + R_>(y, x) e^{-i\varphi_\alpha^>(x, y)} \delta P_>(x, y) \right). \quad (6.39)$$

Integrating over x and y , in (6.34) we obtain a dependence on α and β , which can be described most conveniently by adding indices to the matrices $Y_<$ and $Y_>$ in (6.35),

$$\begin{aligned} \mathcal{T}_V^t(\tilde{\rho}, T_{\mathcal{U}_<, \mathcal{U}_>} \rho) &= \mathcal{T}_V^t(c\mathbf{1}) + \operatorname{Re} \operatorname{Tr}(\delta \hat{A}_< Y_<^{a, \alpha}) + \operatorname{Re} \operatorname{Tr}(\delta \hat{A}_> Y_>^{a, \beta}) \\ &\quad + (\text{higher orders in } \varepsilon/\ell_{\text{macro}}), \end{aligned}$$

We point out that $Y_<$ depends only on α , whereas $Y_>$ depends only on β . This is a consequence of how the phase factors show up in (6.39). The index a clarifies that all the terms also depend on the saddle point $a \in \mathfrak{S}$ of the high-energy wave functions under consideration.

6.6. Construction of the Insertions. Our next task is to construct the insertions to be placed into the integrand of the localized refined pre-state. Given $\mathcal{U}_<$ and $\mathcal{U}_>$, we form the *bosonic insertions* exactly as explained in [20, Section 5.3] as variational derivatives of the nonlinear surface layer integral. More precisely, given holomorphic linearized solution z'_1, \dots, z'_p and z_1, \dots, z_q , we work with the bosonic insertion

$$D_{z'_1}^< \gamma^t(\tilde{\rho}, T_{\mathcal{U}_<, \mathcal{U}_>} \rho) \cdots D_{z'_p}^< \gamma^t(\tilde{\rho}, T_{\mathcal{U}_<, \mathcal{U}_>} \rho) D_{z_1}^> \gamma^t(\tilde{\rho}, T_{\mathcal{U}_<, \mathcal{U}_>} \rho) \cdots D_{z_q}^> \gamma^t(\tilde{\rho}, T_{\mathcal{U}_<, \mathcal{U}_>} \rho).$$

For the *fermionic insertions*, we improve the construction given in [20]. We first give the improved construction and explain the differences to the earlier construction in [20] afterward (see Remark 6.6 below). Given unitary operators $\mathcal{U}_<, \mathcal{U}_>$ as well as a vector $e_\ell \in \mathcal{H}^f$, we consider the anti-linear mapping

$$b_\ell^> : \mathcal{H}_\rho^f \rightarrow \mathbb{C}, \quad b_\ell^>(\phi) = D_{|e_\ell\rangle\bar{\phi}}^> \gamma^t(\tilde{\rho}, T_{\mathcal{U}_<, \mathcal{U}_>} \rho).$$

We represent this anti-linear mapping by a vector $\psi_\ell^>(\mathcal{U}_<, \mathcal{U}_>)$ in the Hilbert space \mathcal{H}_ρ^f , i.e.

$$b_\ell^>(\phi) = \langle \phi | \psi_\ell^> \rangle_\rho \quad \text{for all } \phi \in \mathcal{H}_\rho^f.$$

Intuitively speaking, the wave function $\psi_\ell^>(\mathcal{U}_<, \mathcal{U}_>)$ tells us how the physical wave corresponding to e_ℓ of the *interacting* spacetime looks like for an observer in the vacuum. Next, we let $(e_\ell)_{\ell=1, \dots, f_f}$ be an orthonormal basis of \mathcal{H}_ρ^f and form the Hartree-Fock state

$$\Phi^>(\mathcal{U}_<, \mathcal{U}_>) := \psi_1^>(\mathcal{U}_<, \mathcal{U}_>) \wedge \dots \wedge \psi_{f_f}^>(\mathcal{U}_<, \mathcal{U}_>) \in \mathcal{F}_{\rho, f_f}^f.$$

Similarly, we define the Hartree-Fock state $\Phi^<$ by representing the linear mapping

$$b_\ell^< : \mathcal{H}_\rho^f \rightarrow \mathbb{C}, \quad b_\ell^<(\phi) = D_{\phi\langle e_\ell|}^< \gamma^t(\tilde{\rho}, T_{\mathcal{U}_<, \mathcal{U}_>} \rho)$$

with a vector $\psi_\ell^<(\mathcal{U}_<, \mathcal{U}_>)$,

$$b_\ell^<(\phi) = \langle \psi_\ell^< | \phi \rangle_\rho \quad \text{for all } \phi \in \mathcal{H}_\rho^f,$$

and by taking the resulting Hartree-Fock state,

$$\Phi^<(\mathcal{U}_<, \mathcal{U}_>) := \psi_1^<(\mathcal{U}_<, \mathcal{U}_>) \wedge \dots \wedge \psi_{f_f}^<(\mathcal{U}_<, \mathcal{U}_>) \in \mathcal{F}_{\rho, f_f}^f.$$

The fermionic insertion is introduced as the expectation value of these Hartree-Fock states,

$$\langle \Phi^<(\mathcal{U}_<, \mathcal{U}_>) | \Psi^\dagger(\phi'_1) \dots \Psi^\dagger(\phi'_{r'}) \Psi(\bar{\phi}_1) \dots \Psi(\bar{\phi}_r) | \Phi^>(\mathcal{U}_<, \mathcal{U}_>) \rangle_{\mathcal{F}_{\rho, f_f}^f}. \quad (6.40)$$

This leads us to the following definition.

Definition 6.5. *The localized refined pre-state ω_V^t at time t is defined by*

$$\begin{aligned} \omega_V^t & \left(a^\dagger(z'_1) \dots a^\dagger(z'_p) \Psi^\dagger(\phi'_1) \dots \Psi^\dagger(\phi'_{r'}) a(\bar{z}_1) \dots a(\bar{z}_q) \Psi(\bar{\phi}_1) \dots \Psi(\bar{\phi}_r) \right) \\ & := \frac{1}{Z_V^t(\beta, \tilde{\rho})} \int_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_<) \int_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_>) e^{\alpha N \mathcal{T}_V^t(\tilde{\rho}, T_{\mathcal{U}_<, \mathcal{U}_>} \rho)} \\ & \quad \times \langle \Phi^<(\mathcal{U}_<, \mathcal{U}_>) | \Psi^\dagger(\phi'_1) \dots \Psi^\dagger(\phi'_{r'}) \Psi(\bar{\phi}_1) \dots \Psi(\bar{\phi}_r) | \Phi^>(\mathcal{U}_<, \mathcal{U}_>) \rangle_{\mathcal{F}_{\rho, f_f}^f} \\ & \quad \times D_{z'_1}^< \gamma^t(\tilde{\rho}, T_{\mathcal{U}_<, \mathcal{U}_>} \rho) \dots D_{z'_p}^< \gamma^t(\tilde{\rho}, T_{\mathcal{U}_<, \mathcal{U}_>} \rho) D_{\bar{z}_1}^> \gamma^t(\tilde{\rho}, T_{\mathcal{U}_<, \mathcal{U}_>} \rho) \dots D_{\bar{z}_q}^> \gamma^t(\tilde{\rho}, \mathcal{U} \rho) \end{aligned}$$

with the normalization constant given by

$$Z_V^t := \int_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_<) \int_{\mathcal{G}} d\mu_{\mathcal{G}}(\mathcal{U}_>) e^{\alpha N \mathcal{T}_V^t(\tilde{\rho}, T_{\mathcal{U}_<, \mathcal{U}_>} \rho)} \langle \Phi^<(\mathcal{U}_<, \mathcal{U}_>) | \Phi^>(\mathcal{U}_<, \mathcal{U}_>) \rangle_{\mathcal{F}_{\rho, f_f}^f}.$$

Remark 6.6. (discussion of fermionic insertions) We now compare the insertion (6.40) with the construction used in [20]. For clarity, we begin with the case $\mathcal{U}_< = \mathcal{U}_> = \mathcal{U}$ without refinement. In this case, we can leave out all indices $<$ and $>$, so that (6.40) simplifies to

$$\langle \Phi(\mathcal{U}) | \Psi^\dagger(\phi'_1) \dots \Psi^\dagger(\phi'_{r'}) \Psi(\bar{\phi}_1) \dots \Psi(\bar{\phi}_r) | \Phi(\mathcal{U}) \rangle_{\mathcal{F}_{\rho, f_f}^f}.$$

This expectation value can be computed directly with the Wick rules. The result becomes particularly simple in the case that the effective one-particle wave functions ψ_ℓ are orthonormal

$$\langle \psi_\ell | \psi_{\ell'} \rangle_\rho = \delta_{\ell, \ell'}.$$

In this case, the Hartree-Fock wave function $\Phi(\mathcal{U})$ is normalized. Moreover, the projection operator $\pi^{f,t}$ to the image of the effective one-particle wave functions ψ_ℓ can be written as

$$\pi^{f,t} := \sum_{\ell=1}^{f_f} |\psi_\ell\rangle_\rho \langle \psi_\ell|.$$

A straightforward computation yields

$$\begin{aligned} & \langle \Phi(\mathcal{U}) | \Psi^\dagger(\phi'_1) \cdots \Psi^\dagger(\phi'_{r'}) \Psi(\overline{\phi_1}) \cdots \Psi(\overline{\phi_r}) | \Phi(\mathcal{U}) \rangle_{\mathcal{F}_{\rho, f_f}^f} \\ &= \delta_{r'r} \frac{1}{r!} \sum_{\sigma, \sigma' \in S_r} (-1)^{\text{sign}(\sigma) + \text{sign}(\sigma')} \langle \phi_{\sigma(1)} | \pi^{f,t} \phi'_{\sigma'(1)} \rangle_\rho^t \cdots \langle \phi_{\sigma(r)} | \pi^{f,t} \phi'_{\sigma'(r)} \rangle_\rho^t, \end{aligned} \quad (6.41)$$

giving agreement with the fermionic insertions in [20, Definition 4.1].

Clearly, in typical situations the effective one-particle wave functions will *not* be orthonormal. The construction in [20, Section 4.3] amounts to making these wave functions orthonormal with a Gram-Schmidt procedure. After this has been done, the relation (6.41) again holds. This consideration shows that the construction in [20] differs from Definition 6.5 precisely by the orthonormalization of the effective one-particle wave functions. This raises the question whether such an orthonormalization is desirable or physically preferable. The only advantage of the orthonormalization is that a p -particle measurement only involves p one-particle wave functions (and not all wave functions via the normalization factors coming up in the expectation value). On the other hand, this advantage disappears if one keeps in mind that through the Gram-Schmidt procedure, these p wave functions depend on all the other wave functions as well. More importantly, the orthonormalization seems unnatural, because if the one-particle wave functions are small, then they should also give a small contribution to the expectation value. Finally, Definition 6.5 has the advantage that the positivity properties are more apparent, as will be explained in Section 6.7.

These considerations apply analogously to the refined state. In particular, the suggestion for the fermionic insertions made for the refined state in [20, Section 5] seems superseded by Definition 6.5. \diamond

6.7. Positivity of the Localized Refined Pre-State. Having defined the localized refined pre-state (see Definition 6.5), the next step is to compute it in more detail and to prove positivity. As explained in Section 6.5, to leading order in $1/N$ it suffices to evaluate the integrand at the saddle points. Moreover, we may expand the integrand in powers of the operators $A_<$ and $A_>$. To leading order on the light cone, the insertions are linear in these operators and thus depend only on one of these operators. We take these dependencies into account in our notation by evaluating at the saddle points using the notation

$$\Phi(a, \alpha) := \Phi^<(\mathcal{U}_<, \mathcal{U}_>), \quad \Phi(a, \beta) := \Phi^>(\mathcal{U}_<, \mathcal{U}_>),$$

and similarly for the bosonic insertions. Evaluating the group integrals at the saddle points as explained in Section 6.5 gives the following result.

Theorem 6.7. *To leading order in $1/N$ and $\varepsilon/\ell_{\text{macro}}$, the localized refined pre-state ω_V^t takes the form*

$$\begin{aligned} & \omega_V^t \left(a^\dagger(z'_1) \cdots a^\dagger(z'_p) \Psi^\dagger(\phi'_1) \cdots \Psi^\dagger(\phi'_{r'}) a(\bar{z}_1) \cdots a(\bar{z}_q) \Psi(\bar{\phi}_1) \cdots \Psi(\bar{\phi}_r) \right) \\ &= \sum_{a \in \mathfrak{S}} c_a \sum_{\alpha, \beta \in \mathfrak{T}_a} \langle \Phi(a, \alpha) | \Psi^\dagger(\phi'_1) \cdots \Psi^\dagger(\phi'_{r'}) \Psi(\bar{\phi}_1) \cdots \Psi(\bar{\phi}_r) | \Phi(a, \beta) \rangle_{\mathcal{F}_{\rho, f_f}^f} \\ & \quad \times D_{z'_1}^< \gamma^t(a, \alpha) \cdots D_{z'_p}^< \gamma^t(a, \alpha) D_{\bar{z}_1}^> \gamma^t(a, \beta) \cdots D_{\bar{z}_q}^> \gamma^t(a, \beta) \\ & \quad + \text{higher orders in } 1/N \text{ and } \varepsilon/\ell_{\text{macro}} \end{aligned}$$

with non-negative weights c_a .

To avoid confusion, we point out that, for notational convenience, the factor $1/Z_V^t(\beta, \tilde{\rho})$ was absorbed into the weights c_a .

Theorem 6.8. *To leading order in $1/N$ and $\varepsilon/\ell_{\text{macro}}$, the localized refined pre-state ω_V^t is positive, thereby defining a quantum state (1.3).*

Proof. Before beginning, we note that, using linearity together with the canonical anti-commutation relations, one sees that the formula in Proposition 6.7 holds more generally for any combination B_{fermi} of fermionic field operators, i.e.

$$\omega_V^t \left(a^\dagger(z'_1) \cdots a^\dagger(z'_p) a(\bar{z}_1) \cdots a(\bar{z}_q) B_{\text{fermi}} \right) \quad (6.42)$$

$$\begin{aligned} &= \sum_{a \in \mathfrak{S}} c_a \sum_{\alpha, \beta \in \mathfrak{T}_a} \langle \Phi(a, \alpha) | B_{\text{fermi}} | \Phi(a, \beta) \rangle_{\mathcal{F}_{\rho, f_f}^f} \\ & \quad \times D_{z'_1}^< \gamma^t(a, \alpha) \cdots D_{z'_p}^< \gamma^t(a, \alpha) D_{\bar{z}_1}^> \gamma^t(a, \beta) \cdots D_{\bar{z}_q}^> \gamma^t(a, \beta) \\ & \quad + \text{higher orders in } 1/N \text{ and } \varepsilon/\ell_{\text{macro}} . \end{aligned} \quad (6.43)$$

Given $A \in \mathcal{A}$, our task is to show that $\omega_V^t(A^* A) \geq 0$. We first need to order the bosonic field operators in $A^* A$ as in (6.42). To this end, we use the canonical commutation relations (2.32) to bring all the creation operators to the left and all the annihilation operators to the right. This gives rise to pairings according to

$$a(\bar{z}) a^\dagger(z') \text{ gives } (z|z')_\rho^t ,$$

where always one annihilation operator in A^* is combined with one creation operator in A . After having performed all these commutations, we end up with a product of bosonic field operators of the form as in Definition 6.5, where each creation and annihilation operator gives rise to an insertion $D_z \gamma^t(a, \alpha)$ and $D_{\bar{z}} \gamma^t(a, \beta)$, respectively. After these transformations, the expectation value can be written as follows,

$$\begin{aligned} \omega_V^t(A^* A) &= \sum_{a \in \mathfrak{S}} c_a \sum_{r=0}^{\infty} \sum_{\alpha, \beta \in \mathfrak{T}_a} \sum_{k_1, \dots, k_r} \sum_{n_1, \dots, n_r} \overline{T_{k_1 \dots k_r}(a, \alpha)} T_{n_1 \dots n_r}(a, \beta) \\ & \quad \times (z_{k_1} | z_{n_1})_\rho^t \cdots (z_{k_r} | z_{n_r})_\rho^t \langle \Phi(a, \alpha) | A_{\text{fermi}}^*(r, \alpha) A_{\text{fermi}}(r, \beta) | \Phi(a, \beta) \rangle_{\mathcal{F}_{\rho, f_f}^f} . \end{aligned}$$

In order to clarify the structure of this formula, we introduce the bosonic Fock vector

$$Z_r(T) := \sum_{k_1, \dots, k_r} T_{k_1 \dots k_r} z_{k_1} \otimes \cdots \otimes z_{k_r} .$$

We thus obtain

$$\begin{aligned} \omega_V^t(A^*A) = \sum_{a \in \mathfrak{S}} c_a \sum_{r=0}^{\infty} \sum_{\alpha, \beta \in \mathfrak{I}_a} \langle Z_r(T(a, \alpha)) \otimes \Phi(a, \alpha) \mid A_{\text{fermi}}^*(r, \alpha) \\ \times A_{\text{fermi}}(r, \beta) \mid Z_r(T(a, \beta)) \otimes \Phi(a, \beta) \rangle_{\mathcal{F}_\rho} . \end{aligned}$$

Bringing the operator $A_{\text{fermi}}^*(r, \alpha)$ to the left and introducing the vector

$$\Psi(a, r) = \sum_{\alpha \in \mathfrak{I}_a} Z_r(T(a, \alpha)) \otimes A_{\text{fermi}}(r, \alpha) \Phi(a, \alpha) ,$$

we conclude that

$$\omega_V^t(A^*A) = \sum_{a \in \mathfrak{S}} c_a \sum_{r=0}^{\infty} \langle \Psi(a, r) \mid \Psi(a, r) \rangle_{\mathcal{F}_\rho} .$$

This is obviously non-negative, concluding the proof. \square

7. DESCRIPTION OF ENTANGLEMENT

The formula derived in Theorem 6.7 reveals that the refined state makes it possible to describe entanglement. Indeed, the state ω_V^t can be written as the expectation value of a density operator σ^t

$$\omega_V^t(A) = \text{tr}_{\mathcal{F}_\rho}(\sigma^t A) \quad \text{for all } A \in \mathcal{A} . \quad (7.1)$$

This density operator can be written in *bra/ket* notation as

$$\sigma^t = \sum_{a \in \mathfrak{S}} c_a \left| \sum_{\alpha \in \mathfrak{I}_a} \Psi_{a\alpha}^{\mathcal{F}} \right\rangle \left\langle \sum_{\beta \in \mathfrak{I}_a} \Psi_{a\beta}^{\mathcal{F}} \right| , \quad (7.2)$$

where the Fock vector $\Psi_{a\alpha}^{\mathcal{F}}$ is given as the tensor product of the fermionic Fock vector $\Phi(a, \beta)$ in the statement of Theorem 6.7 with a bosonic Fock vector which realizes the bosonic expectation values. From (7.2) one sees that, in the case of only one saddle point a , one gets a pure state, and the trace in (7.1) reduces to the usual expectation value,

$$\omega_V^t(A) = \left\langle \sum_{\beta \in \mathfrak{I}} \Psi_{\beta}^{\mathcal{F}} \mid A \mid \sum_{\alpha \in \mathfrak{I}} \Psi_{\alpha}^{\mathcal{F}} \right\rangle_{\mathcal{F}_\rho} .$$

If we have more than one saddle point, we can describe a mixed state. The sums over $\alpha, \beta \in \mathfrak{I}_a$ run over the phase freedom of the high-energy saddle point a . In this way, the state (7.2) allows for the description of general entangled states.

Based on these results, we can make the qualitative picture described in Section 3.2 precise. The index α in (3.1) and in the insertions (3.3) describes the phase freedom of our saddle points. This phase freedom appears separately for *bra* and *ket*, giving the synchronization aimed for in (3.4). In this way, we have bypassed the counter argument of Lemma 3.2, making it possible to describe entanglement.

These constructions and considerations even give some insight into the structure of the *long-range correlations* and *dephasing effects* mentioned in the paragraph after (1.2) in the introduction. Indeed, the saddle points are formed of wave functions in a cone in momentum space (see Figure 3). These wave functions are “in phase” even for large distances. Moreover, the phase freedom of each saddle point propagates in lightlike directions (see Figure 6). This phase freedom has the effect that every saddle point is formed of many components having different relative phases. Each component




FIGURE 7. Contractions among unitary factors.

encodes information on a specific Fock component, including both the fermionic and bosonic tensor factors. When integrating over the unitary group, dephasing effects (destructive interference) make it possible to detect and distinguish the different components. These wave components give rise to the summands α in the decomposition of a Fock vector into product states as in (7.2) and (3.1). In summary, we thus obtain nonlocal effects and long-range correlations, giving a natural explanation for Einstein's "spooky action at a distance."

APPENDIX A. DIAGRAMMATIC DERIVATION OF THE GAUSSIAN ASYMPTOTICS


In this appendix we give the proof of Lemma 4.1. Before beginning, we note that, since the group integral in (4.5) involves as many factors \mathcal{U} as \mathcal{U}^{-1} , phase factors in \mathcal{U} cancel out. Therefore, instead of considering the group $U(N)$, we can just as well work with the special unitary group $SU(N)$. We closely follow the procedure in [7], also using a similar graphical notation. We write each factor \mathcal{U} as an up arrow and each factor \mathcal{U}^{-1} as a down arrow (see the left of Figure 7). Next, we use Cramer's rule

$$(\mathcal{U}^{-1})^{ij} = \frac{1}{(N-1)!} \epsilon^{i i_1 \dots i_{N-1}} \epsilon^{j j_1 \dots j_{N-1}} \mathcal{U}_{i_1 j_1} \dots \mathcal{U}_{i_{N-1} j_{N-1}}$$

in order to rewrite the factors \mathcal{U}^{-1} in terms of \mathcal{U} (here ϵ is the totally anti-symmetric Levi-Civita symbol; moreover we use the Einstein summation convention). The resulting expression involves Np factors \mathcal{U} . Its group integral gives pairings of the indices with the Levi-Civita symbol (for details see [7, Figure 6]). Pairs of Levi-Civita symbols whose indices are contracted with each other can be rewritten as sums of products of Kronecker deltas (for details see [7, eq. (18)]). After these transformations, the up and down arrows on the left of Figure 7 are connected both at the top and the bottom. In order to simplify the graphical notation, we permute the down arrows in such a way that every up arrow is connected at the top to the down arrow at its right (these permutations must be taken into account in our end formula by a simultaneous symmetrization in the indices k_1, \dots, k_p and l_1, \dots, l_p). At the bottom, the arrows are connected as shown on the right of Figure 7, with a certain combinatorics which still needs to be specified.

Connecting the arrows in this way, we obtain closed lines connecting 4ℓ indices. We refer to such a closed line as an ℓ -chain (on the right of Figure 7 a three-chain is depicted). Using the freedom to permute the p pairs of arrows (these permutations must be taken into account in our end formula by a simultaneous symmetrization in the indices i_1, \dots, i_p and j_1, \dots, j_p), the resulting configuration of contractions is determined uniquely by chains of length $1 \leq \ell_1 \leq \ell_2 \leq \dots \leq \ell_K$, where K denotes the number of chains. Clearly,

$$\ell_1 + \dots + \ell_K = p.$$

FIGURE 8. The contractions in $[\ell_1, \dots, \ell_K]$.

For a convenient notation, we form the tuple

$$(\ell_k) = (\ell_1, \dots, \ell_K) \quad \text{and set} \quad |(\ell_k)| := \ell_1 + \dots + \ell_K.$$

Then the group integral (4.5) can be written as

$$\sum_{(\ell_k) \text{ with } |(\ell_k)| = p} c_{(\ell_k)}(N) [\ell_1, \dots, \ell_K], \quad (\text{A.1})$$

where $c_{(\ell_k)}(N)$ are combinatorial prefactors (which, due to the signs from the Levi-Civita symbol, could be positive or negative), and the tuple with square brackets stands for the contribution to the group integral

$$\begin{aligned} [\ell_1, \dots, \ell_K] = & \frac{1}{p!} \sum_{\sigma, \sigma' \in S(p)} \delta_{l_{\sigma'(1)}}^{i_{\sigma(1)}} \delta_{j_{\sigma(1)}}^{k_{\sigma'(1)}} \delta_{l_{\sigma'(2)}}^{i_{\sigma(2)}} \delta_{j_{\sigma(2)}}^{k_{\sigma'(2)}} \dots \delta_{l_{\sigma'(n_1)}}^{i_{\sigma(n_1)}} \delta_{j_{\sigma(n_1)}}^{k_{\sigma'(n_1)}} \\ & \times \delta_{l_{\sigma'(n_1+1)}}^{i_{\sigma(n_1+1)}} \delta_{j_{\sigma(n_1+1)}}^{k_{\sigma'(n_1+1)}} \delta_{l_{\sigma'(n_1+2)}}^{i_{\sigma(n_1+2)}} \delta_{j_{\sigma(n_1+2)}}^{k_{\sigma'(n_1+2)}} \\ & \times \dots \delta_{l_{\sigma'(n_1+2n_2+1)}}^{i_{\sigma(n_1+2n_2+1)}} \delta_{j_{\sigma(n_1+2n_2+1)}}^{k_{\sigma'(n_1+2n_2+1)}} \delta_{l_{\sigma'(n_1+2n_2+2)}}^{i_{\sigma(n_1+2n_2+2)}} \delta_{j_{\sigma(n_1+2n_2+2)}}^{k_{\sigma'(n_1+2n_2+2)}} \dots, \end{aligned}$$

where n_ℓ denotes the number chains of length ℓ . The contractions are shown graphically in Figure 8.

The remaining task is to estimate how the combinatorial factors $c_{(\ell_k)}$ scale in N . To this end, it is useful to denote the longest chain of a configuration (ℓ_1, \dots, ℓ_K) by

$$L(\ell_1, \dots, \ell_K) := \ell_K.$$

Next, we consider the combinatorial factors for all configurations for which this longest chain is longer than L and take their maximum,

$$C_L := \max \{ |c_{(\ell_1, \dots, \ell_K)}| \mid L(\ell_1, \dots, \ell_K) \geq L \}.$$


Clearly, these constants are decreasing in L ,

$$C_1 \geq C_2 \geq \dots \geq C_p \geq C_{p+1} = 0$$

(the last equality holds because all chains have length at most p). In the next lemma we estimate the constants C_ℓ . Since we are concerned only with the N -dependence, we want to disregard constants depending only on p , K and the chains (ℓ_1, \dots, ℓ_K) . For a short notation, we write \lesssim for smaller or equal up to a such constant, uniformly in N . Conversely, we write \gg if this uniform inequality does not hold, no matter how large the constant is chosen.

Lemma A.1. *For any $p \in \mathbb{N}$ and for all $L \in \{2, \dots, p\}$,*

$$C_L \lesssim \frac{C_1}{N}. \quad (\text{A.2})$$

FIGURE 9. Contracting the indices i_1 and k_1 .

Proof. We proceed by finite induction in L , beginning at $L = p + 1$ and decreasing L step by step. In the case $L = p + 1$, the inequality (A.2) holds trivially because $C_{p+1} = 0$. Assume that the inequality holds for $L + 1$, i.e.

$$C_{L+1}, \dots, C_{p+1} \lesssim \frac{C_1}{N}. \quad (\text{A.3})$$

Our task is to show that the inequality also holds for L , (A.2).

Following the procedure explained in the final example in [7], we contract the indices i_1 and l_1 . Using that the matrices are unitary, we get a factor $\delta_{j_1}^{k_1}$. The remainder is the group integral in the case $p - 1$. Denoting the contraction by an arrow, we write the resulting formula symbolically as

$$\sum_{(\ell_k) \text{ with } |(\ell_k)| = p} c_{(\ell_k)}(N) [\ell_1, \dots, \ell_K] \longrightarrow \delta_{j_1}^{k_1} \sum_{(\ell_k) \text{ with } |(\ell_k)| = p-1} c_{(\ell_k)}(N) [\ell_1, \dots, \ell_K]. \quad (\text{A.4})$$

On the other hand, the contraction of the indices i_1 and l_1 can be performed for each contribution $[\ell_1, \dots, \ell_K]$. This means symbolically that two arrows are connected at the top by an additional line. The lower points of these arrows become the “free” indices k_1 and j_1 . We denote the resulting configuration by $(q | \ell_1, \dots, \ell_k)$, where q is the length of the line connecting the two free indices, whereas ℓ_1, \dots, ℓ_k are the lengths of the closed chains. This construction is illustrated in Figure 9.

More concretely, we let (ℓ_1, \dots, ℓ_K) be a configuration with $L(\ell_1, \dots, \ell_K) = L \geq 2$. We assume that this configuration violates (A.2), i.e.

$$|c_{(\ell_1, \dots, \ell_K)}| \gg C_1/N. \quad (\text{A.5})$$

Our special attention are the factors N which arise when contracting i_1 and l_1 . Such a factor arises from the formula

$$\delta_j^i \delta_i^j = N$$

when a chain of length one is formed, as is depicted at the top of Figure 9 by a circle. In fact, a factor N arises only if such a circle is formed. Moreover, contracting i_1 and l_1 gives rise to at most one factor N . Next, a circle is formed only if two adjacent arrows are contracted (as shown in the example at the top of Figure 9). In this case, we get an open line whose length is the same as that of the corresponding chain. These findings are summarized by the formula

$$[\ell_1, \dots, \ell_K] \longrightarrow n_{\ell_K} \ell_K N [\ell_K | \ell_1, \dots, \ell_{K-1}] + \sum_{q < \ell_K} \alpha_{q, \tilde{\ell}_K} [q | \tilde{\ell}_1, \dots, \tilde{\ell}_{\tilde{K}}] \quad (\text{A.6})$$

with suitable combinatorial factors $\alpha_{q, \tilde{\ell}_K}$ (whose form will be irrelevant). For clarity, we note that the factor $n_{\ell_K} \ell_K$ arises when counting the number of possible contractions giving rise to a circle. Being independent of N , this combinatorial factor is irrelevant for our argument; we only need that this combinatorial factor is non-zero.

Comparing with (A.4), where the open line has length one, one concludes that the summand involving N in (A.6) must be compensated by other configurations. The configurations $(\tilde{\ell}_1, \dots, \tilde{\ell}_{\tilde{K}})$ with $L(\tilde{\ell}_1, \dots, \tilde{\ell}_{\tilde{K}}) > L$ are not relevant in this respect, because contracting them gives at most one factor N while by the induction hypothesis (A.3) all combinatorial factors involve a factor $1/N$. Therefore, all the contributions obtained by contracting configurations with $L(\tilde{\ell}_1, \dots, \tilde{\ell}_{\tilde{K}}) > L$ are of the order $C_1 \mathcal{O}(N^0)$ and thus cannot compensate the summand involving N in (A.6).

Next, configurations $(\tilde{\ell}_1, \dots, \tilde{\ell}_{\tilde{K}})$ with $L(\tilde{\ell}_1, \dots, \tilde{\ell}_{\tilde{K}}) = L$ cannot compensate the summand involving N in (A.6), simply because the condition $[\ell_K | \ell_1, \dots, \ell_{K-1}] \simeq [\tilde{\ell}_{\tilde{K}} | \tilde{\ell}_1, \dots, \tilde{\ell}_{\tilde{K}-1}]$ implies that the two configurations must be the same.

It remains to consider $(\tilde{\ell}_1, \dots, \tilde{\ell}_{\tilde{K}})$ with $L(\tilde{\ell}_1, \dots, \tilde{\ell}_{\tilde{K}}) < L$. In this case, the contractions do give rise to the configuration $(\ell_K | \ell_1, \dots, \ell_{K-1})$, but only if the indices i_1 and l_1 lie in different chains. This does not give rise to a factor N . We thus obtain the estimate

$$N |c_{(\ell_1, \dots, \ell_K)}| \lesssim C_1,$$

in contradiction to our assumption (A.5). This concludes the proof. \square

Using the estimate of this lemma in (A.1), the group integral can be expanded as

$$\begin{aligned} & \int_{\text{SU}(N)} u_{j_1}^{i_1} \dots u_{j_p}^{i_p} (u^{-1})_{l_1}^{k_1} \dots (u^{-1})_{l_p}^{k_p} d\mu_{\mathcal{G}}(u) = c_{(1, \dots, 1)} [1, \dots, 1] \left(1 + \mathcal{O}\left(\frac{1}{N}\right)\right) \\ &= c_{(1^p)} \frac{1}{p!} \sum_{\sigma, \sigma' \in S(p)} \delta_{l_{\sigma'(1)}}^{i_{\sigma(1)}} \delta_{j_{\sigma(1)}}^{k_{\sigma'(1)}} \dots \delta_{l_{\sigma'(p)}}^{i_{\sigma(p)}} \delta_{j_{\sigma(p)}}^{k_{\sigma'(p)}} \left(1 + \mathcal{O}\left(\frac{1}{N}\right)\right) \\ &= c_{(1^p)} \sum_{\sigma \in S(p)} \delta_{l_{\sigma(1)}}^{i_1} \delta_{j_1}^{k_{\sigma(1)}} \dots \delta_{l_{\sigma(p)}}^{i_p} \delta_{j_p}^{k_{\sigma(p)}} \left(1 + \mathcal{O}\left(\frac{1}{N}\right)\right), \end{aligned} \quad (\text{A.7})$$

where (1^p) is a short notation for the configuration with p chains of length one. Contracting the indices i_1 and l_1 gives

$$\begin{aligned} & \int_{\text{SU}(N)} u_{j_2}^{i_2} \dots u_{j_p}^{i_p} (u^{-1})_{l_2}^{k_2} \dots (u^{-1})_{l_p}^{k_p} d\mu_{\mathcal{G}}(u) \\ &= N c_{(1^p)} \delta_{j_1}^{k_1} \sum_{\sigma \in S(p-1)} \delta_{l_{\sigma(1)+1}}^{i_2} \delta_{j_1}^{k_{\sigma(1)+1}} \dots \delta_{l_{\sigma(p)+1}}^{i_p} \delta_{j_p}^{k_{\sigma(p)+1}} \left(1 + \mathcal{O}\left(\frac{1}{N}\right)\right). \end{aligned}$$

We thus obtain the inductive relation $c_{(1^p-1)} = N c_{(1^p)}$, which in view of $c_{()} = 1$ can be solved to obtain

$$c_{(1^p)} = \frac{1}{N^p}.$$

Using this formula in (A.7) gives (4.5). This concludes the proof of Lemma 4.1.

Acknowledgments: We would like to thank Gernot Akemann, Yan Fyodorov, Mario Kieburg and Tilo Wettig for helpful comments on the integration over the unitary group. We are grateful to the referees for valuable suggestions. We would like to thank the “Universitätsstiftung Hans Vielberth” for generous support. N.K.’s research was also supported by the NSERC grant RGPIN 105490-2018.

REFERENCES

- [1] *Link to web platform on causal fermion systems: www.causal-fermion-system.com.*
- [2] A. Afriat and F. Selleri, *The Einstein, Podolsky, and Rosen Paradox in Atomic, Nuclear, and Particle Physics*, Plenum Press, New York, 1999.
- [3] G. Akemann, Z. Burda, M. Kieburg, and T. Nagao, *Universal microscopic correlation functions for products of truncated unitary matrices*, arXiv:1310.6395 [math-ph], J. Phys. A **47** (2014), no. 25, 255202, 26.
- [4] Y. Bernard and F. Finster, *On the structure of minimizers of causal variational principles in the non-compact and equivariant settings*, arXiv:1205.0403 [math-ph], Adv. Calc. Var. **7** (2014), no. 1, 27–57.
- [5] V.I. Bogachev, *Measure Theory. Vol. I*, Springer-Verlag, Berlin, 2007.
- [6] B. Collins, *Moment methods on compact groups: Weingarten calculus and its applications*, arXiv:2207.08418 [math.OA] (2022).
- [7] M. Creutz, *On invariant integration over $SU(N)$* , J. Math. Phys. **19** (1978), no. 10, 2043–2046.
- [8] C. Dappiaggi and F. Finster, *Linearized fields for causal variational principles: Existence theory and causal structure*, arXiv:1811.10587 [math-ph], Methods Appl. Anal. **27** (2020), no. 1, 1–56.
- [9] C. Dappiaggi, F. Finster, N. Kamran, and M. Reintjes, *The Fock dynamics of causal fermion systems*, in preparation.
- [10] F. Finster, *The Principle of the Fermionic Projector*, hep-th/0001048, hep-th/0202059, hep-th/0210121, AMS/IP Studies in Advanced Mathematics, vol. 35, American Mathematical Society, Providence, RI, 2006.
- [11] ———, *Causal variational principles on measure spaces*, arXiv:0811.2666 [math-ph], J. Reine Angew. Math. **646** (2010), 141–194.
- [12] ———, *Perturbative quantum field theory in the framework of the fermionic projector*, arXiv:1310.4121 [math-ph], J. Math. Phys. **55** (2014), no. 4, 042301.
- [13] ———, *The Continuum Limit of Causal Fermion Systems*, arXiv:1605.04742 [math-ph], Fundamental Theories of Physics, vol. 186, Springer, 2016.
- [14] ———, *Causal fermion systems: A primer for Lorentzian geometers*, arXiv:1709.04781 [math-ph], J. Phys.: Conf. Ser. **968** (2018), 012004.
- [15] ———, *The causal action in Minkowski space and surface layer integrals*, arXiv:1711.07058 [math-ph], SIGMA Symmetry Integrability Geom. Methods Appl. **16** (2020), no. 091.
- [16] ———, *Perturbation theory for critical points of causal variational principles*, arXiv:1703.05059 [math-ph], Adv. Theor. Math. Phys. **24** (2020), no. 3, 563–619.
- [17] F. Finster and M. Jökel, *Causal fermion systems: An elementary introduction to physical ideas and mathematical concepts*, arXiv:1908.08451 [math-ph], Progress and Visions in Quantum Theory in View of Gravity (F. Finster, D. Giulini, J. Kleiner, and J. Tolksdorf, eds.), Birkhäuser Verlag, Basel, 2020, pp. 63–92.
- [18] F. Finster and N. Kamran, *Spinors on singular spaces and the topology of causal fermion systems*, arXiv:1403.7885 [math-ph], Mem. Amer. Math. Soc. **259** (2019), no. 1251, v+83 pp.
- [19] ———, *Complex structures on jet spaces and bosonic Fock space dynamics for causal variational principles*, arXiv:1808.03177 [math-ph], Pure Appl. Math. Q. **17** (2021), no. 1, 55–140.
- [20] ———, *Fermionic Fock spaces and quantum states for causal fermion systems*, arXiv:2101.10793 [math-ph], Ann. Henri Poincaré **23** (2022), no. 4, 1359–1398.

- [21] F. Finster, N. Kamran, and M. Oppio, *The linear dynamics of wave functions in causal fermion systems*, arXiv:2101.08673 [math-ph], J. Differential Equations **293** (2021), 115–187.
- [22] F. Finster and S. Kindermann, *A gauge fixing procedure for causal fermion systems*, arXiv:1908.08445 [math-ph], J. Math. Phys. **61** (2020), no. 8, 082301.
- [23] F. Finster, S. Kindermann, and J.-H. Treude, *An Introductory Course on Causal Fermion Systems*, in preparation, www.causal-fermion-system.com/intro-public.pdf (2023).
- [24] F. Finster and J. Kleiner, *Causal fermion systems as a candidate for a unified physical theory*, arXiv:1502.03587 [math-ph], J. Phys.: Conf. Ser. **626** (2015), 012020.
- [25] ———, *Noether-like theorems for causal variational principles*, arXiv:1506.09076 [math-ph], Calc. Var. Partial Differential Equations **55:35** (2016), no. 2, 41.
- [26] ———, *A Hamiltonian formulation of causal variational principles*, arXiv:1612.07192 [math-ph], Calc. Var. Partial Differential Equations **56:73** (2017), no. 3, 33.
- [27] ———, *A class of conserved surface layer integrals for causal variational principles*, arXiv:1801.08715 [math-ph], Calc. Var. Partial Differential Equations **58:38** (2019), no. 1, 34.
- [28] F. Finster and M. Kraus, *The regularized Hadamard expansion*, arXiv:1708.04447 [math-ph], J. Math. Anal. Appl. **491** (2020), no. 2, 124340.
- [29] F. Finster and C. Langer, *Causal variational principles in the σ -locally compact setting: Existence of minimizers*, arXiv:2002.04412 [math-ph], Adv. Calc. Var. **15** (2022), no. 3, 551–575.
- [30] F. Finster and M. Lottner, *Banach manifold structure and infinite-dimensional analysis for causal fermion systems*, arXiv:2101.11908 [math-ph], Ann. Global Anal. Geom. **60** (2021), no. 2, 313–354.
- [31] F. Finster and A. Platzer, *A positive mass theorem for static causal fermion systems*, arXiv:1912.12995 [math-ph], Adv. Theor. Math. Phys. **25** (2021), no. 7, 1735–1818.
- [32] W.A. Friedman and P.A. Mello, *Marginal distribution of an arbitrary square submatrix of the S -matrix for Dyson’s measure*, J. Phys. A **18** (1985), no. 3, 425–436.
- [33] Y.V. Fyodorov and H.-J. Sommers, *Random matrices close to Hermitian or unitary: Overview of methods and results*, arXiv:nlin/0207051 [nlin.CD], J. Phys. A **36** (2003), no. 12, 3303–3347.
- [34] R.E. Greene and K. Shiohama, *Diffeomorphisms and volume-preserving embeddings of noncompact manifolds*, Trans. Amer. Math. Soc. **255** (1979), 403–414.
- [35] S. Helgason, *Groups and Geometric Analysis*, Mathematical Surveys and Monographs, vol. 83, American Mathematical Society, Providence, RI, 2000, Integral geometry, invariant differential operators, and spherical functions, Corrected reprint of the 1984 original.
- [36] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, *Quantum entanglement*, arXiv:quant-ph/0702225, Rev. Modern Phys. **81** (2009), no. 2, 865–942.
- [37] G. Huber, *Notes: Gamma function derivation of n -sphere volumes*, Amer. Math. Monthly **89** (1982), no. 5, 301–302.
- [38] J.R. Ipsen and M. Kieburg, *Weak commutation relations and eigenvalue statistics for products of rectangular random matrices*, arXiv:1310.4154 [math-ph], Phys. Rev. E **89** (2014), 032106.
- [39] C. Itzykson and J.B. Zuber, *The planar approximation. II*, J. Math. Phys. **21** (1980), no. 3, 411–421.
- [40] C. McSwiggen, *The Harish-Chandra integral: An introduction with examples*, arXiv:1806.11155 [math-ph], Enseign. Math. **67** (2021), no. 3-4, 229–299.
- [41] M.L. Mehta, *Random Matrices*, third ed., Pure and Applied Mathematics (Amsterdam), vol. 142, Elsevier/Academic Press, Amsterdam, 2004.
- [42] Y.A. Neretin, *Hua-type integrals over unitary groups and over projective limits of unitary groups*, arXiv:math-ph/0010014, Duke Math. J. **114** (2002), no. 2, 239–266.
- [43] M.A. Nielsen and I.L. Chuang, *Quantum Computation and Quantum Information*, Cambridge University Press, Cambridge, 2000.
- [44] M. Oppio, *On the mathematical foundations of causal fermion systems in Minkowski space*, arXiv:1909.09229 [math-ph], Ann. Henri Poincaré **22** (2021), no. 3, 873–949.
- [45] D.N. Page, *Density matrix of the Universe*, Phys. Rev. D (3) **34** (1986), no. 8, 2267–2271.
- [46] H.J. Rothe, *Lattice Gauge Theories*, third ed., World Scientific Lecture Notes in Physics, vol. 74, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2005, An introduction.
- [47] J.J. Sakurai and J. Napolitano, *Advanced Quantum Mechanics*, second ed., Addison-Wesley Publishing Company, 1994.
- [48] F. Schwabl, *Quantum Mechanics*, third ed., Springer-Verlag, Berlin, 2002.

- [49] D. Weingarten, *Asymptotic behavior of group integrals in the limit of infinite rank*, J. Math. Phys. **19** (1978), no. 5, 999–1001.

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT REGENSBURG, D-93040 REGENSBURG, GERMANY
Email address: `finster@ur.de`

DEPARTMENT OF MATHEMATICS AND STATISTICS, MCGILL UNIVERSITY, MONTRÉAL, CANADA
Email address: `nkamran@math.mcgill.ca`

DEPARTMENT OF MATHEMATICS, CITY UNIVERSITY OF HONG KONG, SAR HONG KONG
Email address: `moritzreintjes@gmail.com`