Analyzing the Differentially Private Theil-Sen Estimator for Simple Linear Regression

Jayshree Sarathy Columbia University

Salil Vadhan Harvard University JS6514@COLUMBIA.EDU

SALIL_VADHAN@HARVARD.EDU

Abstract

In this paper, we study differentially private point and confidence interval estimators for simple linear regression. Motivated by recent work that highlights the strong empirical performance of an algorithm based on robust statistics, DPTheilSen, we provide a rigorous, finite-sample analysis of its privacy and accuracy properties, offer guidance on setting hyperparameters, and show how to produce differentially private confidence intervals to accompany its point estimates. **Keywords:** List of keywords

1. Introduction

Over the last several years, differential privacy (DP) (Dwork, 2006) has become a widely accepted standard for protecting the privacy of data subjects while releasing useful statistical information about datasets (Abowd, 2018; Altman et al., 2018; Biden, 2023). However, designing *usable* DP methods for common statistical inference tasks remains an ongoing challenge (Barrientos et al., 2021; Sarathy et al., 2023; Garrido et al., 2023). One example is simple (ie. one-dimensional) linear regression, which is one of the most fundamental tasks in data analysis. In 2018, the prominent economics research group, Opportunity Insights, found that there was a lack of guidance around choosing accurate DP algorithms for simple linear regression on regimes commonly used in practice (e.g. small-area analysis with 40 to 400 datapoints per regression) (Chetty and Friedman, 2019).

Motivated by this gap between theory and practice, Alabi et al. (2022) conducted an empirical evaluation of several DP algorithms for simple linear regression. They found that a suite of robust, median-based algorithms, DPTheilSen, based on the non-private Theil-Sen estimator developed by Theil (1950) and Sen (1968), performed better than standard OLS-based algorithms across a range of practical regimes. DPTheilSen operates in two steps: first, it computes the slope for some or all pairs of points and second, uses a DP median algorithm to output a single estimate of the slope.¹

While Alabi et al.'s empirical study was a valuable starting point, the authors stated that further theoretical understanding of the accuracy guarantees of DPTheilSen, as well as design of uncertainty estimates, would be needed to make this set of algorithms fully usable in practice. In this paper, we address these open questions by analyzing the accuracy guarantees of the DPTheilSen algorithms. Our contributions can be summarized as follows.

1. We provide a rigorous theoretical analysis of the DPTheilSen algorithms shown to perform strongly by Alabi et al. (2022). Our analyses offer finite-sample convergence bounds and shed light on *why* and *when* DPTheilSen outperforms other DP linear regression algorithms.

^{1.} The intercept can be computed either along with the slope or using the final estimate of the slope, as described in (Sen, 1968) and (Alabi et al., 2022).

2. We design and analyze DP confidence intervals for simple linear regression via DPTheilSen.

Although our work focuses on the one-dimensional linear regression setting, our approach is a crucial starting point to developing similar finite-sample analyses for higher-dimensional regimes.² Overall, we view our work as a contribution towards understanding private and robust regression more generally. Before providing an overview of our main results, we describe the problem of simple linear regression and the assumptions we make in this work, which are minimal.

1.1. Simple linear regression

We are given *n* values, x_1, \ldots, x_n , of the predictor variable *x*. For each x_i , we observe the corresponding value y_i of the response random variable *y*. The model is $y_i = \beta_0 + \beta_1 x_i + e_i$ for $i = 1, \ldots, n$, where β_0 and β_1 are unknown parameters. We make the following assumptions:

Assumption 1.1.1 x_1, \ldots, x_n are fixed and not all equal.

Assumption 1.1.2 Each e_i is sampled independently from the same continuous, symmetric, mean-0 distribution F_e .

A common formulation of linear regression is the Ordinary Least Squares (OLS) objective, which is characterized by the following optimization problem:

$$(\hat{\beta}_0, \hat{\beta}_1) = \arg\min_{\beta_0, \beta_1 \in \mathbb{R}} \|\mathbf{y} - \beta_1 \mathbf{x} - \beta_0 \mathbf{1}\|_2,$$

where $\mathbf{x} = (x_1, \dots, x_n)^T$, $\mathbf{y} = (y_1, \dots, y_n)^T$, and **1** is the all-ones vector. OLS has a simple closed form solution:

$$\hat{\beta_1} = rac{n \operatorname{cov}(\mathbf{x}, \mathbf{y})}{n \operatorname{var}(\mathbf{x})} \text{ and } \hat{\beta_0} = \bar{y} - \hat{\beta_1} \bar{x},$$

where $\bar{x} = \frac{1}{n} \sum_{i=1}^{n} x_i$, $\bar{y} = \frac{1}{n} \sum_{i=1}^{n} y_i$, $n \operatorname{cov}(\mathbf{x}, \mathbf{y}) = \langle \mathbf{x} - \bar{x}\mathbf{1}, \mathbf{y} - \bar{y}\mathbf{1} \rangle$, and $n \operatorname{var}(\mathbf{x}) = \langle \mathbf{x} - \bar{x}\mathbf{1}, \mathbf{x} - \bar{x}\mathbf{1} \rangle$. When **y** is generated according to the model $y_i = \beta_0 + \beta_1 \cdot x_i + e_i$, $\forall i \in [n]$ for i.i.d. Gaussian noise e_i , then the OLS solution is the maximum likelihood estimator. If we remove the assumption of Gaussian noise and add privacy constraints, however, robust estimators such as Theil-Sen have been shown to provide better accuracy (Dwork and Lei, 2009; Alabi et al., 2022). In this work, we analyze DP algorithms based on Theil-Sen, comparing them to the OLS-based approaches. Below, we provide brief descriptions of these two approaches:

• **OLS-based algorithms**: The DPSuffStats algorithm (Dwork et al., 2014; Wang, 2018; Alabi et al., 2022) follows the OLS approach closely, in that it involves perturbing the sufficient statistics $n \operatorname{cov}(\mathbf{x}, \mathbf{y})$ and $n \operatorname{var}(\mathbf{x})$. While this algorithm is computationally efficient and enables releasing the DP sufficient statistics at no extra privacy cost, it has been shown to not perform well in common regimes.³A second algorithm is DPOLSExp (Bassily et al., 2014; Alabi, 2022), which implements the exponential mechanism (McSherry and Talwar, 2007) with the OLS objective function.

^{2.} In particular, our analysis can be used as a black box to analyze Knop and Steinke (2022)'s approach of coordinate-wise medians for higher-dimensional linear regression. Some new ideas may be required to analyze approaches that use more sophisticated higher-dimensional aggregators based on Tukey depth (see Section 1.3), but again our analysis should provide guidance as it is the one-dimensional analogue of those methods.

^{3.} In particular, Alabi et al. (2022) showed that DPSuffStats performs poorly in the high privacy, small dataset, and/or clustered independent variable regime.

• Theil-Sen-based algorithms: Theil-Sen (Theil, 1950; Sen, 1968) is a family of robust linear regression estimators that proceed via two steps (as illustrated in Fig. 1) for estimating the slope β_1 : (1) compute the slope for some or all pairs of points, and (2) output the median of these estimates. To make this algorithm DP, we can simply replace the median with a DP median. Although there are many choices for the DP median, we consider a version of this algorithm, DPWideTS, which uses the *widened exponential mechanism* (Alabi et al., 2022; McSherry and Talwar, 2007) as the DP median sub-routine, as this version has been shown to exhibit strong empirical performance by Alabi et al. (2022).



Figure 1: Illustration of the standard non-private Theil-Sen algorithm (Theil, 1950; Sen, 1968), which (1) computes the slopes between all pairs of points, and (2) outputs the median slope in dark blue.

1.2. Overview of results

This work offers two main contributions: finite-sample guarantees and confidence intervals for DPTheilSen. First, we provide a finite-sample convergence bound for DPWideTS, the variant of DP Theil-Sen that was recommended by Alabi et al. (2022). We attain the convergence bound by developing a finite-sample analysis of non-i.i.d. *U-statistics* (Hoeffding, 1948; Chen et al., 2011), which may be of independent interest.

We state our convergence bound informally below. While our main analysis does not require any data assumptions beyond the ones stated above (Assumptions 1.1.1 and 1.1.2), the theorem is stated for a special case, where the independent variables x_1, \ldots, x_n are evenly split between the endpoints of an interval of length Δ_x (known as an *asymptotically optimal design* (Sen, 1968)), and the noise variables e_1, \ldots, e_n are drawn i.i.d. from $\mathcal{N}(0, \sigma_e^2)$.

Theorem 1.2.1 (Main result applied for case of asymptotically optimal design, informally stated) Let $\tilde{\beta}_1^{DPWideTS}$ be the DPWideTS estimator with privacy loss parameter ε , hyperparameter R for the range of the outputs, and hyperparameter θ for the granularity of the outputs. Assume that the true slope β_1 lies in the interval $[-R + \theta, R - \theta]$. Let τ be defined as follows.

$$\tau = \Phi^{-1} \left(1 - \frac{p}{8} \right) \cdot \sqrt{\frac{4}{3n}} + O\left(\frac{\ln(R/p\theta)}{\varepsilon n} \right)$$

where Φ^{-1} is the inverse standard normal distribution function. Then, for suff. large n, and suff. small τ and σ_e/Δ_x , we have that with probability at least 1 - p,

$$\widetilde{\beta_1}^{\text{DPWideTS}} \in [\beta_1 - z - \theta, \beta_1 + z + \theta]$$

for
$$z = \frac{\sqrt{\pi \cdot \sigma_e}}{\Delta_x} \cdot \left(\tau + O\left(\tau^{3/2}\right)\right)$$

In Table 1 below, we provide an informal comparison between the convergence bounds for DPWideTS (described above) and the OLS-based DPSuffStats algorithm for this setting (analyzed in Alabi (2022)). We also compare these DP estimators with their non-private counterparts. We can see that DPWideTS maintains the same leading constant of $2\sqrt{\pi/3}$ as that of non-private TS, while non-private OLS and DPSuffStats have a leading constant of 2, so the latter are slightly better in the asymptotic $n \to \infty$ regime. For DPWideTS, we see a logarithmic dependence on the output range R, as compared to the quadratic dependence shown by DPSuffStats on the input range r. Our comparison provides theoretical backing to the empirical finding of Alabi et al. (2022)—that the quantity $\varepsilon n \Delta_x^2$ is important in choosing between the standard DPSuffStats algorithm and the robust DPWideTS algorithm. In particular, when $\varepsilon n \Delta_x^2$ is small, which indicates a high privacy, small dataset size, and/or clustered independent variable regime, DPWideTS is the more accurate estimator. Finally, our convergence bound enables us to provide guidance on setting the θ hyperparameter (corresponding to the granularity of the widened exponential mechanism) for DPWideTS, which was a key open problem raised by Alabi et al. (2022) towards making this algorithm usable for practitioners.

Estimator	1-p Convergence Bound	
Non-priv OLS	$rac{2\sigma_e}{\Delta_x}\cdotrac{c_{p/2}}{\sqrt{n}}$	
Non-priv TS	$\sqrt{\frac{\pi}{3}} \cdot \frac{2\sigma_e}{\Delta_x} \cdot \frac{c_{p/4}}{\sqrt{n}}$	
DPSuffStats (Alabi, 2022)	$\frac{2\sigma_e}{\Delta_x} \cdot \frac{c_{p/6}}{\sqrt{n}} \cdot (1+\tau) + \tau \left(1+\tau + \beta_1 \right),$ $\tau \approx \frac{(1-1/n)r^2 \log(3/p)}{\varepsilon \cdot n \cdot \Delta_x^2}$	
DPWideTS (Thm. 1.2.1)	$ \sqrt{\frac{\pi}{3}} \cdot \frac{2\sigma_e}{\Delta_x} \cdot \left(\frac{c_{p/8}}{\sqrt{n}} + \gamma\right) (1 + o(1)) + \theta, \ \gamma \approx \frac{\ln(R/p \cdot \theta)}{\varepsilon n} $	

Table 1: 1 - p convergence bounds for point estimators for simple linear regression in special case of asymptotically optimal design. Note that r is range for both the input x_i, y_i datapoints, R is range for the output estimate of β_1 , and $c_{p/2} = \Phi^{-1}(1 - p/2) = \Theta(\log(1/p))$ for small p.

In addition to analyzing the DPWideTS point estimator, we design and analyze corresponding confidence interval estimators for simple linear regression. We describe two algorithms— DPWideTSCIUnion and DPWideTSCI—and analyze their privacy, coverage, and accuracy guarantees. We use the DP median estimators proposed by Drechsler et al. (2022) as sub-routines in our algorithms, making them additionally usable by analyzing the width of the confidence intervals they provide. We show that the confidence interval for DPWideTSCIUnion is approximately twice as large as the convergence bound for DPWideTS, and can be further improved using the algorithm DPWideTSCI.

1.3. Related work

This work draws on the rich connections between robust statistics and DP (Dwork and Lei, 2009; Asi et al., 2023; Hopkins et al., 2023). Dwork and Lei (2009) gave an asymptotic analysis of what they called the "Short-Cut Regression Method," which is similar to a simplified variant of DPTheilSen. However, they did not consider the more statistically efficient variants that we do in this work, provide finite sample guarantees, or offer measures of uncertainty for the estimates like the confidence intervals we provide. Other works (Couch et al., 2019; Avella-Medina, 2021) confirm the findings of Dwork and Lei in the context of hypothesis testing but similarly do not provide finite-sample analysis. A longer discussion of these works is provided in Appendix A.

In general, linear regression is one of the most fundamental tasks in statistics and yet has been a challenging problem in the DP literature (Zhang et al., 2012; Sheffet, 2017; Barrientos et al., 2017; Bernstein and Sheldon, 2018; Awan and Slavković, 2018; Wang, 2018; Cai et al., 2019; Evans and King, 2021; Ramsay and Chenouri, 2021; Knop and Steinke, 2022; Liu et al., 2022, 2023). Most prior works focus on non-robust methods, require additional assumptions on the data or model, or only provide asymptotic analysis. To address these gaps, Alabi et al. (2022) conducted an experimental evaluation of DP algorithms for simple linear regression in the *small dataset regime*, demonstrating that DP analogues of robust algorithms, such as DPTheilSen, perform better than non-robust methods in practical settings. However, Alabi et al. did not provide theoretical bounds or DP confidence intervals for this estimator, which are open questions we address in this work.

Our confidence interval algorithms build upon Drechsler et al. (2022)'s non-parametric DP confidence intervals for the median, but ours are more general in that they provide finite-sample validity for some forms of non-i.i.d. variables and characterizes the width of the confidence intervals. Other work produces DP confidence intervals using bootstrapping (D'Orazio et al., 2015; Barrientos et al., 2017; Brawner and Honaker, 2018; Ferrando et al., 2020), but these can be expensive to compute and rely on assumptions, such as normally distributed errors, that our work avoids.

Concurrent work. Since our results were first announced,⁴ there has been a flurry of work that uses median-based approaches for DP tasks (Brown et al., 2021; Liu et al., 2022; Ramsay and Chenouri, 2021; Cumings-Menon, 2022; Knop and Steinke, 2022) and clarifies the connections between privacy and robustness (Asi et al., 2023; Hopkins et al., 2023).⁵ None provides the analysis we offer in this paper; we touch on a few here but do not provide a comprehensive survey.

One relevant work by Amin et al. (2023) provides empirical evaluation of higher-dimensional variations of Theil-Sen, but unlike our work, does not offer theoretical bounds on accuracy or convergence. Knop and Steinke (2022), on the other hand, do provide theoretical privacy and utility analysis of some of the DP Theil-Sen algorithms in higher dimensions, as well as an experimental evaluation. They offer a finite-sample convergence bound that is consistent with our results, but their statement only applies for the highly simplified variant of DPTheilSen where each data point is used only once, so only n/2 slopes are computed and the slopes are independent of each other. Our results are more general and can handle cases that achieve provably stronger performance by reusing data points and introducing correlations. Examples include the asymptotically optimal design

^{4.} References for presentations of our results in 2021 have been removed for anonymity.

^{5.} See Appendix A for a longer discussion of related and concurrent work.

of Thm. 1.2.1 and the full Theil-Sen variant where all $\binom{n}{2}$ points are used.⁶ In addition, Knop and Steinke do not aim to match the leading constant of the non-private estimators, which is an important feature of our results.

2. Preliminaries

We consider datasets that are multisets. The space of datasets is denoted by Multisets(\mathcal{D}, n), where \mathcal{D} is the underlying set of elements and n is the cardinality of each multiset. Let $dist_{ms}(d, d')$ be the number of records that must be changed to transform d into d'.⁷

Since our algorithms include hyperparameters, we state a definition of DP for algorithms that take as input not only the dataset, but also the desired privacy parameters and any required hyperparameters. Two datasets $d, d' \in Multisets(\mathcal{D}, n)$ are *neighboring*, denoted $d \sim d'$, if dist(d, d') = 1. Let \mathcal{H} be a hyperparameter space and \mathcal{Y} be an output space.

Definition 2.0.1 (Differential Privacy (Dwork et al., 2006)) For $\varepsilon \in \mathbb{R}_{\geq 0}$, a randomized algorithm M: $Multisets(\mathcal{D}, n) \times \mathbb{R}_{\geq 0} \times \mathcal{H} \to \mathcal{Y}$ is ε -DP if and only if for all neighboring datasets $d \sim d' \in Multisets(\mathcal{D}, n)$ hyperparams $\in \mathcal{H}$, and sets $E \subseteq \mathcal{Y}$,

 $\begin{aligned} &\Pr[M(\boldsymbol{d},\varepsilon, hyperparams) \in E] \\ &\leq e^{\varepsilon} \cdot \Pr[M(\boldsymbol{d}',\varepsilon, hyperparams) \in E]. \end{aligned}$

where the probabilities are taken over the random coins of M.

Now, we will define the non-private Theil-Sen family of estimators.

Definition 2.0.2 (Theil-Sen Estimator (Theil, 1950; Sen, 1968)) Let $(x_1, y_1), \ldots, (x_n, y_n)$ be an arbitrary ordering of dataset $d \in Multisets(\mathbb{R} \times \mathbb{R}, n)$. Let S be a set of N unordered pairs of elements of $[n] = \{1, \ldots, n\}$ such that for each pair $\{i, j\} \in S$, $x_i \neq x_j$. Then, for each $\{i, j\} \in S$, compute the slope s_{ij} between the points (x_i, y_i) and (x_j, y_j) as follows: $s_{ij} = (y_j - y_i)/(x_j - x_i)$. Let $s = \{s_{ij}\}_{\{i,j\}\in S}$ denote the multiset of the slopes. The Theil-Sen estimator $\hat{\beta}_1^{\text{TS}}$ with corresponding set S is computed as follows: $\hat{\beta}_1^{\text{TS}} = median(s)$.

3. DPTheilSen Algorithm

In the differentially private version of Theil-Sen, which we call DPTheilSen (Algorithm 3.1), we similarly compute pairwise estimates of the slope. However, we replacing the computation of the median of the slopes with a differentially private median algorithm (denoted by DPmed, which can be one of several algorithms). DPmed takes as input the multiset of slopes, $\mathbf{s}_d \in$ Multisets ($\mathbb{R} \cup \{-\infty, \infty\}, N$), the scaled privacy parameter $\varepsilon/k \in \mathbb{R}_{\geq 0}$, where k is the max number of slopes computed using each datapoint, and the hyperparameters $\in \mathcal{H}$ for the given median algorithm.

Lemma 3.0.1 (Alabi et al. (2022)) Algorithm 3.1 (DPTheilSen) is ε -DP.

^{6.} Note that the simplified variant of Theil-Sen can be analyzed using a Chernoff-Hoeffding bound, while the versions of Theil-Sen with correlated slopes require more complex analytical tools such as U-statistics.

^{7.} I.e. "change-one distance" (Casacuberta et al., 2022)).

Algorithm 3.1 DPTheilSen: ε-DP Algorithm

Data: $d = (x_i, y_i)_{i=1}^n \in \text{Multisets}(\mathbb{R} \times \mathbb{R}, n),$ **Privacy params:** $\varepsilon \in \mathbb{R}_{\geq 0}$ **Hyperparams:** $S \in \binom{[n]}{2}$, DPmed, hyperparams $\in \mathcal{H}$ $\mathbf{s} = \{\}$ for each $\{i, j\} \in S$ (such that $x_i \neq x_j$) do $\begin{vmatrix} s_{ij} = (y_j - y_i)/(x_j - x_i) \\ \text{Add } s_{ij} \text{ to } \mathbf{s} \end{vmatrix}$ end Let $k = \max_{i \in [n]} \{ \#j \in [n] : \{i, j\} \in S \}$ $\widetilde{\beta_1}^{\text{TS}} = \text{DPmed}(\mathbf{s}, \varepsilon/k, \text{hyperparams})$ return $\widetilde{\beta_1}^{\text{TS}}$

In this work, we consider a version of DPTheilSen called DPWideTS, which uses for DPmed the widened exponential mechanism, DPWide (Alabi et al., 2022; McSherry and Talwar, 2007). This variant of DPTheilSen was found to have strong performance in the empirical work of Alabi et al. (2022). The DPWide algorithm (Alg. B.1) and utility theorem (Thm. B.0.3) are included in the Appendix.

4. Convergence Bound for DPWideTS

In this section, we provide a finite-sample analysis of DPWideTS. We look beyond asymptotics solely in *n*, as this will not explain the strong empirical performance of the DPWideTS algorithm compared to others such as DPSuffStats, as shown by (Alabi et al., 2022). In addition, finite-sample analysis will allow us to better understand the *conditions* under which DPWideTS outperforms non-robust algorithms.

The main challenge of analyzing DPWideTS is that the slopes computed from S are correlated.⁸ To deal with correlated slopes, Sen (1968) relies on the properties of *U*-statistics (Hoeffding, 1948) for an asymptotic analysis of the non-private Theil-Sen algorithm.

Definition 4.0.1 (U-statistic for simple linear regression (Sen, 1968)) Let x_1, \ldots, x_n satisfy Assumption 1.1.1, and let y_1, \ldots, y_n be the corresponding response variables under the model $y_i = \beta_0 + \beta_1 x_i + e_i$, where $\beta_0, \beta_1 \in \mathbb{R}$ and each e_i is sampled i.i.d from a continuous, symmetric, mean-0 distribution F_e . The U-statistic takes as input a "guess" $\hat{\beta}_1 \in \mathbb{R}$ for the true slope β_1 , as well as the datapoints $(x_i, y_i)_{i=1}^n$, indexed arbitrarily. Then, the U-statistic for simple linear regression is defined as follows.

$$U(\hat{\beta}_{1}, (x_{i}, y_{i})_{i=1}^{n}) = \frac{1}{N} \sum_{\{i, j\} \in S, i < j} sign\left(\frac{y_{j} - y_{i}}{x_{j} - x_{i}} - \hat{\beta}_{1}\right),$$
(1)

where S is the set of unordered pairs of datapoints used to compute the slopes (where each pair has distinct x values), N = |S| is the number of such pairs, and sign(q) = -1 if q < 0, 0 if q = 0, and 1

^{8.} We can avoid correlated slopes if we use an 'incomplete' version of the algorithm where each data point is only used in one slope, i.e. k = 1. Our general analysis applies to this simplified version, as well as to variants that compute a linear # of total slopes for computational efficiency.

if q > 0. For ease of notation, we will let $\hat{\beta}_1 = \beta_1 + z$, for some $z \in \mathbb{R}$, and use U_z as shorthand for $U\left(\hat{\beta}_1, (x_i, y_i)_{i=1}^n\right)$.

We build on this approach, adapting finite-sample 'Berry-Esseen-type' bounds for the convergence of the U-statistic in order to develop a finite-sample convergence bound for the DPWideTS estimate.

Theorem 4.0.2 (Convergence bound for $\beta_1^{\text{DPWideTS}}$) Let x_1, \ldots, x_n satisfy Assumption 1.1.1 and have empirical variance σ_x^2 . Let y_1, \ldots, y_n be the corresponding response variables under the model $y_i = \beta_0 + \beta_1 x_i + e_i$, where $\beta_0, \beta_1 \in \mathbb{R}$ and each e_i is sampled i.i.d from a continuous, symmetric, mean-0 distribution F_e . Let $\beta_1^{\text{DPWideTS}} = \text{DPTheilSen}(\{x_i, y_i\}_{i=1}^n, \varepsilon, (S, \text{DPWide}, \theta, -R, R))$, where $\varepsilon, R, \theta > 0, \beta_1 \in [-R + \theta, R - \theta]$, and $S \in {[n] \choose 2}$ is a set of unordered pairs with distinct x-values. Let U_z be the U-statistic defined with respect to S.

Then, there exists a constant c > 0 such that for sufficiently large n, with probability at least 1 - p, $\tilde{\beta_1}^{DPWideTS} \in [\beta_1 - z - \theta, \beta_1 + z + \theta]$ for every z that satisfies the following:

$$\frac{\mu(z) - \frac{4\ln(4R/p\theta)}{\varepsilon \cdot n}}{\sigma(z)} \ge \Phi^{-1} \left(1 - \frac{p}{2} + \frac{c}{n^2 \cdot \sigma^3(z)}\right)$$

where $\mu(z) = E[U_z], \sigma^2(z) = Var[U_z]$, and Φ is the standard normal cdf.

The quantities $\mu(z)$ and $\sigma^2(z)$ can be further described and evaluated, as shown in Sections C.1 and 5. In the next section, we provide an overview of the analysis and the use of the U-statistic. However, this result will be easier to interpret when applied to a special setting, as described in Section 5.

4.1. Overview of analysis

To analyze the performance of DPTheilSen, we further consider the U-statistic from Definition 4.0.1. In particular, we can rewrite statistic (1) in the form of a *Kendall's tau statistic* (Kendall, 1948) that measures the rank correlation between the x's and the residuals from a line of slope $\beta_1 + z$.

$$U_{z} = U(\beta_{1} + z, (x_{i}, y_{i})_{i=1}^{n}) = \frac{1}{N} \sum_{\substack{\{i, j\} \in S \\ i < j}} \operatorname{sign}(x_{j} - x_{i}) \cdot \operatorname{sign}(e_{j} - e_{i} - z \cdot (x_{j} - x_{i}))$$
(2)

Suppose z = 0. Then, we define the *null U-statistic* as

$$U_0 = U(\beta_1, (x_i, y_i)_{i=1}^n) = \frac{1}{N} \sum_{\{i, j\} \in S, i < j} \operatorname{sign}(x_j - x_i) \cdot \operatorname{sign}(e_j - e_i)$$
(3)

Furthermore, the distribution of U_0 matches the null distribution of the Kendall's τ -statistic, which is known to be asymptotically normal (Kendall, 1948). Observe that $E_{e_1,\ldots,e_n}[U_z] = 0$ since e_1,\ldots,e_n are i.i.d. In fact, we have:

Fact 4.1.1 (Sen (1968)) $E_{e_1,...,e_n}[U_z] = 0$ iff z = 0.

Using these properties of the U-statistic, we proceed with the algorithm analysis in two steps.

- By the utility theorem of the widened exponential mechanism (Theorem B.0.3), we show that with high probability, U (β₁^{DPWideTS}, (x_i, y_i)ⁿ_{i=1}) ≈ 0.
 Putting together U (β₁^{DPWideTS}, (x_i, y_i)ⁿ_{i=1}) ≈ 0, and the asymptotic normality of U_z for appropriate z (Theorem 4.2.1), we show that β₁^{DPWideTS} ≈ β₁ and characterize the finitesample convergence of the estimator.

Note that the second step requires showing that U_z , not just U_0 , is asymptotically normal, and we must characterize the convergence using the quantity z itself. We do so in the next section.

4.2. Finite-sample convergence of U-statistic

We develop a finite-sample bound for the absolute value difference between the distribution function of U_z and the standard normal distribution function Φ . To do so, we modify Berry-Esseen bounds for linear functions, such as U-statistics, of i.i.d mean-0 random variables (see, e.g., Chen et al. (2011)). In the case of the U-statistic U_z described in (2), the terms $sign(e_j - e_i - z \cdot (x_j - x_i))$ for $\{i, j\} \in S, i < j$ are non-identically distributed since $x_j - x_i$ may be different for different pairs (i, j). Therefore, we adapt the Berry-Esseen bounds to work for U-statistics of independent, non-identical random variables. Our finite-sample convergence bound is stated in Theorem 4.2.1 below.

Theorem 4.2.1 Let x_1, \ldots, x_n satisfy Assumption 1.1.1 (i.e. not all equal) and let y_1, \ldots, y_n be the corresponding response variables under the model $y_i = \beta_0 + \beta_1 x_i + e_i$, where $\beta_0, \beta_1 \in \mathbb{R}$ and each e_i is sampled i.i.d from a continuous, symmetric, mean-0 distribution. Let S be a set of unordered pairs of datapoints such that each pair has distinct x-values, and let U_z be defined with respect to S as in (2). Let $\mu(z) = \mathbb{E}_{e_1,\ldots,e_n}[U_z]$, and let $\sigma^2(z) = \operatorname{Var}_{e_1,\ldots,e_n}[U_z]$. Then, for sufficiently large n,

$$\sup_{t \in \mathbb{R}} \left| \Pr\left[\frac{U_z - \mu(z)}{\sigma(z)} \le t \right] - \Phi(t) \right| = O\left(\frac{1}{n^2 \cdot \sigma^3(z)} \right)$$

where Φ is the cdf of a standard normal distribution.

This result shows that the normalized U-statistic, U_z , converges to a standard normal distribution at a rate of $O(1/(n^2 \cdot \sigma^3(z)))$, where $\sigma^2(z) = \Theta(1/n)$ is the variance of the U-statistic.⁹

5. Evaluating Bound for Special Case

In this section, we will consider the special setting (called an asymptotically optimal design (Sen, (1968)) where the x-values are evenly split between the endpoints of an interval of length Δ_x . The DPWideTS algorithm computes $N = \lfloor n/2 \rfloor \cdot \lfloor n/2 \rfloor$ slopes¹⁰ from pairs of datapoints on opposite ends of the interval. In addition, we assume that the noise variables are sampled i.i.d from a normal distribution.

Assumption 5.0.1 $x_1, \ldots, x_{\lfloor n/2 \rfloor} = 0$ and $x_{\lfloor n/2 \rfloor + 1}, \ldots, x_n = \Delta_x$.

^{9.} For example, the variance of the U-statistic evaluated for the asymptotically optimal design is 4/3n + O(1/n), as shown in Lemma E.0.3. The general case is shown in Lemma C.1.2.

^{10.} Our analysis extends to efficient versions of this algorithm that compute a linear, rather than quadratic, number of slopes.

Assumption 5.0.2 $F_e = \mathcal{N}(0, \sigma_e^2)$.

Table 2 shows the DPWideTS convergence bound for this special setting, along with convergence bounds for the non-private algorithms OLS and Theil-Sen and the private algorithms DPSuffStats and DPOLSExp (Alabi, 2022) (these were described briefly in Section 1.1), all under the same assumptions.

Estimator	Size of $1 - p$ Convergence Bound	Constraints
OLS	$2 \cdot \frac{\sigma_e}{\Delta_x} \cdot \frac{c_{p/2}}{\sqrt{n}}$	
Theil-Sen (Sen, 1968)	$2 \cdot \sqrt{\frac{\pi}{3}} \cdot \frac{\sigma_e}{\Delta_x} \cdot \frac{c_{p/4}}{\sqrt{n}}$	
DPSuffStats (Alabi, 2022)	$2 \cdot \frac{\sigma_e}{\Delta_x} \cdot \frac{c_{p/6}}{\sqrt{n}} \cdot (1+\tau) + \tau \left(1+\tau + \beta_1 \right),$ $\tau \approx \frac{12(1-1/n)r^2 \log(3/p)}{\varepsilon \cdot n \cdot \Delta_x^2}$	$p \in (3\exp(-\varepsilon n\Delta_x^2/12), 1)$
DPOLSExp (Alabi, 2022)	$2 \cdot \frac{\sigma_e}{\Delta_x} \cdot \frac{c_{p/4}}{\sqrt{n}} + \sqrt{\frac{64R \cdot (3\log(3) + \log(2/p))}{\varepsilon n^2 \Delta_x^2}}$	
DPWideTS (Thm 1.2.1)	$2 \cdot \sqrt{\frac{\pi}{3}} \cdot \frac{\sigma_e}{\Delta_x} \cdot \left(\frac{c_{p/8}}{\sqrt{n}} + \frac{4\sqrt{3}\ln(4R/p\theta)}{\varepsilon \cdot n}\right) + \theta$	$p \in (0, 4R/\theta),$ $\frac{\sigma_e}{\Delta_x} \text{ suff. small, } n \text{ suff. large}$

Table 2: High-probability (1 - p) convergence bounds for estimators for special setting (Assumptions 5.0.1 and 5.0.2). Note that r is a hyperparameter range for both the input x_i, y_i datapoints, while R is a range for the output estimate of the slope.

The first two bounds in the table correspond to the non-private algorithms OLS and Theil-Sen. Both of these non-private algorithms have a convergence rate of $O(\sigma_e \cdot c_p/\Delta_x \cdot \sqrt{n})$. Looking at the leading constant, we see that Theil-Sen nearly recovers the accuracy of OLS, up to a factor of $\sqrt{\pi/3}$. The Theil-Sen bound has the term $c_{p/4}$ instead of $c_{p/2}$, which comes from the convergence of the U-statistic to a standard normal distribution at a rate of $O(1/\sqrt{n})$ (Theorem 4.2.1).

The bounds for DPSuffStats and DPTheilSen have the same constant factors for the highest order term as OLS and Theil-Sen, respectively, but they also contain two main differences from their non-private counterparts. First, the DP estimators have constant factor changes in p in the c_p terms compared to those in the non-private bounds; this comes from the DP estimators taking a union bound over both the sampling and privacy error, which splits up the failure probability p further. Note that $c_p = \Theta(\sqrt{\log(p)})$ for small p, so these constant factor changes in p are not impactful.

Second, the DP estimators include lower order terms corresponding to the noise due to privacy,¹¹ which provide insight into the relative performance of the DP algorithms in practical regimes. For

^{11.} For example, the $\ln(R/p\theta)/\varepsilon n$ term for DPWideTS corresponds to the noise due to privacy, which is $O(1/\varepsilon n)$ and thus overshadowed by the sampling error $O(1/\sqrt{n})$ for sufficiently large n.

example, we can see that DPSuffStats's lower order term τ is $O(r^2/\varepsilon \cdot n \cdot \Delta_x^2)$; the $\varepsilon n \Delta_x^2$ quantity in the denominator was highlighted in the empirical work of Alabi et al. (2022) as important for the performance of the algorithm but lacked theoretical backing prior to our work.¹² Meanwhile, the lower order term in DPOLSExp has the quantity $\sqrt{\varepsilon n \Delta_x}$ in the denominator, which indicates that it performs better than DPSuffStats for small n and for small ε . The lower order term in DPWideTS has the quantity $\varepsilon n \Delta_x$ in the denominator, which indicates that it performs better than DPSuffStats for small Δ_x , but not as well as DPOLSExp in the small ε regime.

Finally, we can compare the DPSuffStats, DPOLSExp, and DPWideTS bounds in terms of their dependence on hyperparameters. While DPSuffStats has a quadratic dependence on r, the range of the input datapoints, the other two algorithms avoid any dependence on the range of the inputs. Instead, DPOLSExp has a square root dependence on the range R of the output estimate, while DPWideTS has a milder logarithmic dependence on R. The bound for DPWideTS also includes the parameter θ (corresponding to the granularity of outputs in DPWide). In Section E.1, we discuss how to set this parameter, addressing an open question of (Alabi, 2022).

6. Confidence Intervals for DPWideTS

In addition to the DP point estimators, we also consider DP confidence intervals for the linear regression slope β_1 . We start with the non-private Theil-Sen estimator, described in Sen (1968). This estimator returns the (1/2 - b)th and (1/2 + b)th quantiles of the set of Theil-Sen slopes, s, where b is computed according to the distribution of the corresponding U-statistic.¹³

Now, we consider the differentially private confidence interval for β_1 , DPTSCI, which we describe in Algorithm F.1 (See Appendix). This algorithm is similar to the DPTheilSen point estimator (Alg. 3.1), except that we replace the call to DPMed with a call to an ε -DP confidence interval algorithm for the median, DPMedCI.

In Alg. 6.1, we describe one possible DPMedCI algorithm: DPWideCI (Union). This algorithm is based on a nonparametric confidence interval for the median designed by Drechsler et al. (2022). The idea is to run the DPWide point estimator twice such that with high probability, the two estimates capture the true slope β_1 . The DPWideCIUnion algorithm does so by outputting an interval that contains the non-private interval with high probability, while the DPWideCI algorithm outputs a tighter interval via more nuanced coverage analysis. While Drechsler et al. (2022) assume that the inputs to the confidence interval mechanism are sampled i.i.d. from a population distribution, here we only assume that we have a uniform bound on the convergence of the empirical distribution F_s to the standard normal distribution (which we develop in Thm. 4.2.1).

Let DPWideTSCIUnion = DPTSCI (d, ε , S, DPWideCIUnion, (α , r_{α} , θ , -R, R, Union=1)), where DPWideCIUnion is defined in Algorithm 6.1. Our main theorem for DPWideTSCIUnion, stated in the Appendix, describes the validity and width of this confidence interval. The proof of the validity relies on the validity of the non-private interval (Lemma F.1.2), along with the utility of DPWide (Thm. B.0.3). The width of the DPWideTSCIUnion confidence interval is approximately twice the size of the convergence bound of the point estimator (Thm. 4.0.2), which we expect even in the non-private setting. However, there is an additional factor of 4 in the term corresponding to privacy noise ($\ln(2R/\alpha\theta)$). This can be improved using a more nuanced coverage analysis.

^{12.} Alabi et al. specificially looked at the quantity $\varepsilon \cdot n \cdot \sigma_x^2$, where σ_x^2 is the empirical variance of the constants x_1, \ldots, x_n , but they did not provide theoretical basis for the importance of this quantity. In our special setting, $\Delta_x^2 = 4\sigma_x^2$.

^{13.} See the Appendix for more details on this interval estimator and a proof of its validity.

Algorithm 6.1 DPWideCI (Union): *ε*-DP Algorithm

Data: $\mathbf{s} = (s_1, \ldots, s_N) \in \mathbb{R}^N$ **Privacy params:** $\varepsilon \in \mathbb{R}_{>0}$ **Hyperparams:** $\alpha \in (0,1), r_{\alpha} \in (0,1), \theta \in \mathbb{R}_{>0}, [-R,R] \subset \mathbb{R}, \text{Union} \in \{0,1\}$ if Union then Let $\alpha_1 = r_{\alpha} \cdot \alpha$ and $\alpha_2 = (1 - r_{\alpha}) \cdot \alpha$ Let $b = \frac{1}{2} \cdot \Phi^{-1} (1 - \alpha_1/8) \cdot \sigma(0)$ // $\sigma^2(0)$ is the variance of the null U-statistic, U_0 , corresponding to the set of slopes **s**. As shown in Corollary C.1.3, it does not depend on the data. Let $c = 2 \ln(4R/\alpha_2\theta)/(\varepsilon N)$ Set $q_L = 1/2 - b - c$ and $q_U = 1/2 + b + c$ end else $q_L, q_U = \text{ComputeExpMechCITargets}(\mathbf{s}, \varepsilon, \alpha, -R, R, \theta)$ // Algorithm F.2 end $L_{\alpha}(\mathbf{s}) = \text{DPWide}(\mathbf{s}, \varepsilon/2, (q_L, R, \theta))$ // Algorithm B.1 $\widetilde{U_{\alpha}}(\mathbf{s}) = \text{DPWide}(\mathbf{s}, \varepsilon/2, (q_U, R, \theta))$ return $[L_{\alpha}(\mathbf{s}) - \theta, U_{\alpha}(\mathbf{s}) + \theta]$

To improve the width of the confidence interval, we apply the more sophisticated approach from Drechsler et al. (2022). In particular, we call DPWideCI with the Union flag set to 0, so that q_L, q_U are computed using ComputeExpMechCITargets (Alg. F.2). We offer a brief overview here: For a given $t_L, t_U \in [N], \varepsilon \in \mathbb{R}_{>0}$, let $\tilde{L}_{\alpha}(\mathbf{s}, t_L) = \text{DPWide}(\mathbf{s}, \varepsilon/2, (t_L/N, \theta, -R, R))$ and $\tilde{U}_{\alpha}(\mathbf{s}, t_U) = \text{DPWide}(\mathbf{s}, \varepsilon/2k, (t_U/N, \theta, -R, R))$. As before, the goal is to control the probability that the interval $[\tilde{L}_{\alpha}(\mathbf{s}, t_L) - \theta, \tilde{U}_{\alpha}(\mathbf{s}, t_U) + \theta]$ fails to contain the true slope β_1 . In particular, for $\alpha \in (0, 1)$, we would like to find the target ranks t_L and t_U closest to N/2 such that

$$\Pr_{\mathbf{s}, \text{DPWideCI}} \left[\tilde{L_{\alpha}}(\mathbf{s}, t_L) - \theta > \beta_1 \right] \le \alpha/2, \text{ and } \Pr_{\mathbf{s}, \text{DPWideCI}} \left[\tilde{U_{\alpha}}(\mathbf{s}, t_U) + \theta < \beta_1 \right] \le \alpha/2$$
(4)

In Algorithm F.2, we find these target ranks by first computing the probabilities above for all possible t_L and t_U 's, and then by numerically searching for the target ranks closest to N/2 such that the probabilities above are both within $\alpha/2$. This search can be implemented more efficiently by noting that t_L is greater than or equal to $N \cdot (1/2 - b - c)$ as defined in Algorithm 6.1, and similarly t_U is less than or equal to $N \cdot (1/2 + b + c)$. This also tell us that the width of the interval will always be the same or smaller than that of the DPWideTSCIUnion interval (Thm. F.3.1).

The validity analysis for Alg. F.2 (Thm. F.4.2) relies on characterizing the distribution of the rank of the population median in the dataset. In our setting, where the slopes **s** computed by DPTSCI are not necessarily i.i.d, we do so via the distribution of the corresponding U-statistic (Thm. 4.2.1).

7. Conclusion

In this work, we analyze the theoretical privacy and utility guarantees of DPTheilSen. We provide finite-sample convergence bounds, offer insight into hyperparameter selection, and show how to produce differentially private confidence intervals. Future work should analyze the optimality of these algorithms and extend the confidence intervals to multivariate settings.

Acknowledgments

We thank Daniel Alabi, Audra McMillan, and Adam Smith for discussions that motivated this work. We also thank Cynthia Dwork and James Honaker for helpful conversations at an early stage of this work, and Ryan Cumings-Menon and others at the Fields Institute Privacy Workshop in Summer 2022 for helpful comparisons with related work on robust regression.

SV was supported by a Simons Investigator Award and JS was supported by the Columbia Data Science Institute Fellowship. Both authors were supported by Cooperative Agreement CB20ADR0160001 with the Census Bureau. The views expressed in this paper are those of the authors and not those of the U.S. Census Bureau or any other sponsor.

References

- John M Abowd. The us census bureau adopts differential privacy. In *Proceedings of the 24th ACM SIGKDD international conference on knowledge discovery & data mining*, pages 2867–2867, 2018.
- Daniel Alabi, Audra McMillan, Jayshree Sarathy, Adam Smith, and Salil Vadhan. Differentially private simple linear regression. *Proceedings on Privacy Enhancing Technologies*, 2:184–204, 2022.
- Daniel Gbenga Alabi. *The Algorithmic Foundations of Private Computational Social Science*. PhD thesis, Harvard University, 2022.
- Micah Altman, Alexandra Wood, David R O'Brien, and Urs Gasser. Practical approaches to big data privacy over time. *International Data Privacy Law*, 8(1):29–51, 2018.
- Kareem Amin, Matthew Joseph, Mónica Ribero, and Sergei Vassilvitskii. Easy differentially private linear regression. In *The Eleventh International Conference on Learning Representations*, 2023.
- Hilal Asi, Jonathan Ullman, and Lydia Zakynthinou. From robustness to privacy and back. arXiv preprint arXiv:2302.01855, 2023.
- Marco Avella-Medina. Privacy-preserving parametric inference: a case for robust statistics. *Journal* of the American Statistical Association, 116(534):969–983, 2021.
- Jordan Awan and Aleksandra Slavković. Differentially private uniformly most powerful tests for binomial data. *Advances in Neural Information Processing Systems*, 31, 2018.
- Andr'es F. Barrientos, J. Reiter, Ashwin Machanavajjhala, and Yan Chen. Differentially private significance tests for regression coefficients. *Journal of Computational and Graphical Statistics*, 28:440 – 453, 2017.
- Andrés F Barrientos, Aaron R Williams, Joshua Snoke, and Claire McKay Bowen. A feasibility study of differentially private summary statistics and regression analyses for administrative tax data. *arXiv preprint arXiv:2110.12055*, 2021.
- Raef Bassily, Adam Smith, and Abhradeep Thakurta. Private empirical risk minimization: Efficient algorithms and tight error bounds. In 2014 IEEE 55th annual symposium on foundations of computer science, pages 464–473. IEEE, 2014.

- Garrett Bernstein and Daniel R Sheldon. Differentially private bayesian inference for exponential families. In S. Bengio, H. Wallach, H. Larochelle, K. Grauman, N. Cesa-Bianchi, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, volume 31. Curran Associates, Inc., 2018. URL https://proceedings.neurips.cc/paper/2018/file/ 08040837089cdf46631a10aca5258e16-Paper.pdf.
- Andrew C Berry. The accuracy of the gaussian approximation to the sum of independent variates. *Transactions of the american mathematical society*, 49(1):122–136, 1941.
- Joseph R Biden. Executive order on the safe, secure, and trustworthy development and use of artificial intelligence. 2023.
- Thomas W. Brawner and J. Honaker. Bootstrap inference and differential privacy: Standard errors for free. 2018.
- Gavin Brown, Marco Gaboardi, Adam Smith, Jonathan Ullman, and Lydia Zakynthinou. Covarianceaware private mean estimation without private covariance estimation. Advances in Neural Information Processing Systems, 34:7950–7964, 2021.
- T. Tony Cai, Yichen Wang, and Linjun Zhang. The cost of privacy: Optimal rates of convergence for parameter estimation with differential privacy. *CoRR*, abs/1902.04495, 2019. URL http://arxiv.org/abs/1902.04495.
- Sílvia Casacuberta, Michael Shoemate, Salil Vadhan, and Connor Wagaman. Widespread underestimation of sensitivity in differentially private libraries and how to fix it. In *Proceedings of the 2022 ACM SIGSAC Conference on Computer and Communications Security*, pages 471–484, 2022.
- Louis HY Chen, Larry Goldstein, and Qi-Man Shao. *Normal approximation by Stein's method*, volume 2. Springer, 2011.
- Raj Chetty and John N. Friedman. A practical method to reduce privacy loss when disclosing statistics based on small samples. *American Economic Review Papers and Proceedings*, 109: 414–420, 2019.
- Simon Couch, Zeki Kazan, Kaiyan Shi, Andrew Bray, and Adam Groce. Differentially private nonparametric hypothesis testing. *arXiv preprint arXiv:1903.09364*, 2019.
- Ryan Cumings-Menon. Differentially private estimation via statistical depth. *arXiv preprint arXiv:2207.12602*, 2022.
- Vito D'Orazio, J. Honaker, and G. King. Differential privacy for social science inference. *Alfred P. Sloan Foundation Economic Research Paper Series*, 2015.
- Joerg Drechsler, Ira Globus-Harris, Audra McMillan, Jayshree Sarathy, and Adam Smith. Nonparametric differentially private confidence intervals for the median. *Journal of Survey Statistics and Methodology*, 10:804–829, 2022.
- Cynthia Dwork. Differential privacy. In Automata, Languages and Programming, 33rd International Colloquium, ICALP 2006, Venice, Italy, July 10-14, 2006, Proceedings, Part II, pages 1–12, 2006.

- Cynthia Dwork and Jing Lei. Differential privacy and robust statistics. In *STOC*, volume 9, pages 371–380, 2009.
- Cynthia Dwork, Frank McSherry, Kobbi Nissim, and Adam D. Smith. Calibrating noise to sensitivity in private data analysis. In *Theory of Cryptography, Third Theory of Cryptography Conference, TCC 2006, New York, NY, USA, March 4-7, 2006, Proceedings*, pages 265–284, 2006.
- Cynthia Dwork, Kunal Talwar, Abhradeep Thakurta, and Li Zhang. Analyze gauss: optimal bounds for privacy-preserving principal component analysis. In *Symposium on Theory of Computing, STOC 2014, New York, NY, USA, May 31 June 03, 2014*, pages 11–20, 2014.
- Carl-Gustav Esseen. On the liapunov limit error in the theory of probability. *Ark. Mat. Astr. Fys.*, 28: 1–19, 1942.
- Georgina Evans and Gary King. Statistically valid inferences from differentially private data releases, with application to the facebook urls dataset. *Political Analysis*, 2021 2021.
- C. Ferrando, Shu-Fan Wang, and D. Sheldon. General-purpose differentially-private confidence intervals. *ArXiv*, abs/2006.07749, 2020.
- Gonzalo M Garrido, Xiaoyuan Liu, Florian Matthes, and Dawn Song. Lessons learned: Surveying the practicality of differential privacy in the industry. *Proceedings on Privacy Enhancing Technologies*, 2:151–170, 2023.
- Wassily Hoeffding. A class of statistics with asymptotically normal distribution. Annals of Mathematical Statistics, 19:293–325, 1948.
- Samuel B Hopkins, Gautam Kamath, Mahbod Majid, and Shyam Narayanan. Robustness implies privacy in statistical estimation. In *Proceedings of the 55th Annual ACM Symposium on Theory of Computing*, pages 497–506, 2023.
- Maurice George Kendall. Rank correlation methods. 1948.
- Alexander Knop and Thomas Steinke. Differentially private linear regression via medians. 2022.
- Xiyang Liu, Weihao Kong, and Sewoong Oh. Differential privacy and robust statistics in high dimensions. In *Conference on Learning Theory*, pages 1167–1246. PMLR, 2022.
- Xiyang Liu, Prateek Jain, Weihao Kong, Sewoong Oh, and Arun Suggala. Label robust and differentially private linear regression: Computational and statistical efficiency. In *Thirty-seventh Conference on Neural Information Processing Systems*, 2023.
- Frank McSherry and Kunal Talwar. Mechanism design via differential privacy. In 48th Annual IEEE Symposium on Foundations of Computer Science (FOCS 2007), October 20-23, 2007, Providence, RI, USA, Proceedings, pages 94–103, 2007.
- Kelly Ramsay and Shoja'eddin Chenouri. Differentially private depth functions and their associated medians. *arXiv preprint arXiv:2101.02800*, 2021.

- Jayshree Sarathy, Sophia Song, Audrey Haque, Tania Schlatter, and Salil Vadhan. Don't look at the data! how differential privacy reconfigures the practices of data science. In *Proceedings of the 2023 CHI Conference on Human Factors in Computing Systems*, pages 1–19, 2023.
- Pranab Kumar Sen. Estimates of the regression coefficient based on kendall's tau. *Journal of the American statistical association*, 63(324):1379–1389, 1968.
- Or Sheffet. Differentially private ordinary least squares. In *Proceedings of the 34th International Conference on Machine Learning, ICML 2017, Sydney, NSW, Australia, 6-11 August 2017*, pages 3105–3114, 2017. URL http://proceedings.mlr.press/v70/sheffet17a.html.
- Adam Smith. Privacy-preserving statistical estimation with optimal convergence rates. In *Proceedings* of the forty-third annual ACM symposium on Theory of computing, pages 813–822, 2011.
- Henri Theil. A rank-invariant method of linear and polynomial regression analysis, 3; confidence regions for the parameters of polynomial regression equations. *Indagationes Mathematicae*, 1(2): 467–482, 1950.
- Yu-Xiang Wang. Revisiting differentially private linear regression: optimal and adaptive prediction & estimation in unbounded domain. In *Proceedings of the Thirty-Fourth Conference on Uncertainty* in Artificial Intelligence, UAI 2018, Monterey, California, USA, August 6-10, 2018, pages 93–103, 2018.
- Jun Zhang, Zhenjie Zhang, Xiaokui Xiao, Yin Yang, and Marianne Winslett. Functional mechanism: Regression analysis under differential privacy. *Proc. VLDB Endow.*, 5(11):1364–1375, 2012.

Appendix A. Related Work, continued

We begin by describing related work in more detail. Several recent works have leveraged the connection between robustness and differential privacy to design privacy-preserving algorithms for specific statistical tasks, affirming Dwork and Lei (2009)'s initial findings that robust estimators are a valuable starting point for accurate DP algorithms. For example, Couch et al. (2019) find that robust estimators perform better than parametric estimators under differential privacy, even when the data come from a parametric model, but they focus on hypothesis testing and do not provide a theoretical utility analysis. Avella-Medina (2021) designs DP estimators based on bounded influence M-estimators, including DP tests for Wald, score, and likelihood ratio tests; they provide asymptotic analysis of the statistical guarantees, while our work focuses on finite-sample analysis.

More generally, recent work has significantly advanced our understanding of the connection between robust and private algorithms. Work by Asi et al. (2023) establishes a tight connection between privacy and robustness, providing a black-box transformation from optimal robust to optimal differentially private algorithms. They also design and analyze estimators for DP linear regression for high-dimensional tasks under assumptions of Gaussianity. Work by Hopkins et al. (2023) shows how to implement this black-box transformation in a computationally efficient manner via the sum-of-squares method. While these methods provide important general toolkits for DP algorithm design, they do not provide finite-sample bounds for specific estimators such as DPTheilSen.

Many recent works in the DP literature have considered linear regression, but most focus on empirical or asymptotic analysis rather than finite-sample guarantees. Sheffet (2017) considered differentially private ordinary least squares (OLS) methods and corresponding DP confidence intervals, but unlike our work, these methods assume normality of errors, require input data bounds, and satisfy approximate, rather than pure, DP. Wang (2018) studied private ridge regression and considered DP confidence intervals, but these methods require consuming additional privacy budget for estimating Hessians. Barrientos et al. (2017) and Evans and King (2021) use the subsample-and-aggregate framework, but their approaches rely on normality assumptions or normal approximations that only hold for large n. Bernstein and Sheldon (2018) consider a Bayesian approach, but unlike our work, they require a prior on the distribution of both the regression coefficients and the independent variables.

Finally, recent studies have further highlighted the need for more usable DP algorithms for basic statistical tasks such as simple linear regression. A study of data practitioners' use of DP by Sarathy et al. (2023) calls for algorithms with minimal hyperparameters and assumptions, as well as useful uncertainty measures, that will enable data analysts to navigate the constraints of the DP analysis process. Furthermore, a study by Barrientos et al. (2021) evaluates a range of DP linear regression algorithms in terms of feasibility for real-world use. They develop criteria around evaluating algorithms, including: assumptions that align with practical applications, ease of implementation, computational efficiency, minimal tuning parameters, and accompanied by uncertainty estimates. This study finds that suprisingly few algorithms in the DP literature satisfy these criteria. Our work on developing the design and analysis of DPTheilSen makes this family of algorithms measure up across all of these categories, suggesting that it will be useful for practical deployments of DP.

Appendix B. Widened Exponential Mechanism (DPWide)

In this section, we describe the *widened exponential mechanism* for quantile estimation designed by Alabi et al. (2022), which is a variant of the standard exponential mechanism (McSherry and Talwar, 2007) described below.

Definition B.0.1 (Exponential Mechanism (McSherry and Talwar, 2007)) The exponential mechanism is defined with respect to a utility function u, which maps (data set, output) pairs to real values. Given dataset s and range of possible outputs, [-R, R], the exponential mechanism outputs $r \in [-R, R]$ with probability density proportional to $\exp\left(\frac{\varepsilon u(s,r)}{2\Delta u}\right)$, where $\Delta(u) = \sup_{r \in [-R,R]} \sup_{s \sim s'} |u(s,r) - u(s',r)|$. The exponential mechanism is ε -DP.

The standard instantiation of the exponential mechanism for quantile estimation (used, e.g., in Smith (2011)) uses a utility function that assigns a score to output r based on how far r is *in rank* from the desired quantile of **s**. For all outputs r within the range [-R, R], and for a target quantile $q \in (0, 1)$, the standard utility function is:

$$u(\mathbf{s}, r) = -|| \# \text{below } r - n \cdot q||$$

where #below r denotes the number of datapoints in s that are less than or equal to r in value. Note that this utility function assigns the same utility score to every output r in the interval between two data points.

One issue with this mechanism is that when the output space is the real line and the data is highly concentrated, the mechanism may not place enough probability density near the target quantile. To mitigate this issue, Alabi et al. (2022) design a variation on the standard utility function. For widening parameter $\theta > 0$, and target quantile $q \in (0, 1)$, the widened utility function is:

$$u(\mathbf{s},r) = -\lfloor \min_{w \in [r-\theta,r+\theta]} | \# \text{below } w - n \cdot q | \rfloor$$

This utility function provides a lower bound on the probability density the mechanism assigns around the target quantile. Following Alabi et al. (2022), an efficient implementation of the θ -widened exponential mechanism for the median is given in Algorithm B.1.

Below, we state the standard utility theorem for DPWide with widening hyperparameter θ on a fixed dataset s. Let F_s^{-1} be the inverse empirical distribution function for the set of slopes, s. We use the following assumption on the range and widening hyperparameters $R, \theta \in \mathbb{R}_{>0}$.

Assumption B.0.2 For a target $q \in (0,1)$ and fixed dataset $\mathbf{s} \in \mathbb{R}^N$, the true quantile $F_{\mathbf{s}}^{-1}(q) \in [-R + \theta, R - \theta]$.

Theorem B.0.3 (Utility of DPWide) Let $s \in \mathbb{R}^N$ be a fixed sample of N points, and let F_s^{-1} be the inverse empirical distribution function. For $q \in (0, 1)$, let [-R, R], the range hyperparameter, satisfy Assumption B.0.2, and let $\theta, \varepsilon > 0$. Let $\tilde{q}^{\text{DPWide}} = \text{DPWide}(s, \varepsilon, (q, -R, R, \theta))$. Then, for $0 < c < \min(q, 1/2)$,

$$\Pr_{\substack{\text{DPWide}(\boldsymbol{s},\varepsilon, \\ (q,-R,R,\theta))}} \left(\widetilde{q}^{\text{DPWide}} \notin [F_{\boldsymbol{s}}^{-1}(q-c) - \theta, F_{\boldsymbol{s}}^{-1}(q+c) + \theta] \right) \leq \frac{R}{\theta} \exp\left(-\frac{\varepsilon cN}{2}\right)$$

Algorithm B.1 Widened Exponential Mechanism for Quantile (DPWide): ε -DP Algorithm

Data: $\mathbf{s} = (s_1, \ldots, s_N) \in \mathbb{R}^N$ **Privacy params:** $\varepsilon \in \mathbb{R}_{\geq 0}$ **Hyperparams:** $q \in (0, 1), -R, R \in \mathbb{R}^2; \theta \in \mathbb{R}_{>0}$ Sort s /* Clip **s** to the range [-R,R] and insert space θ around the qth quantile: */ for $i \in [1, \lfloor Nq \mid]$ do $| \mathbf{s}[i] = \min(\max(-R, \mathbf{s}[i] - \theta), R)$ end for $i \in [|Nq| + 1, N]$ do $|\mathbf{s}[i] = \max(\min(R, \mathbf{s}[i] + \theta), -R)$ end Insert -R and R into **s** and set N = N + 2Set maxNoisyScore = $-\infty$ Set argMaxNoisyScore = -1for $i \in [2, N)$ do score = $\log(\mathbf{s}[i] - \mathbf{s}[i-1]) - \frac{\varepsilon}{2} \cdot \lfloor |i - Nq||$ $Z \sim \text{Gumbel}(0,1)$ noisyScore = score + Zif noisyScore > maxNoisyScore then maxNoisyScore = noisyScore argMaxNoisyScore = iend end left = s[argMaxNoisyScore - 1]right = s[argMaxNoisyScore]Sample $\tilde{q}^{\text{DPWide}} \sim \text{Unif}[\text{left}, \text{right}]$ return $\widetilde{q}^{\text{DPWide}}$

Proof Let F_s be the empirical distribution function for s. The utility score of an output $r \in [-R, R]$ is $-\lfloor \min_{w \in [r-\theta, r+\theta]} N \cdot |F_s(w) - q| \rfloor$. Let $r^* \in [\max(-R, r-\theta), \min(R, r+\theta)]$ be the value that maximizes the utility function for potential output value r. Therefore, we can rewrite the utility score of r as $-\lfloor N |F_s(r^*) - q \rfloor \rfloor$.

Next, let us upper bound the probability that the mechanism selects an output with score $\leq -cN$. The exponential mechanism assigns un-normalized probability density of at most $\exp(-\varepsilon cN/2)$ to each of these outputs, and they span at most the interval [-R, R]. On the other hand, the exponential mechanism assigns un-normalized probability density of 1 to output values with score of zero, which we know exist in the range [-R, R] by Assumption B.0.2. In particular, all outputs within θ of $F_s^{-1}(q)$ have score of zero. Therefore, we have that

$$\begin{aligned} \Pr_{\text{DPWide}(\mathbf{s},\varepsilon,(q,-R,R,\theta))} & \left(\widetilde{q}^{\text{DPWide}*} \notin [F_{\mathbf{s}}^{-1}(q-c), F_{\mathbf{s}}^{-1}(q+c)] \right) \\ &= \Pr_{\text{DPWide}(\mathbf{s},\varepsilon,(q,-R,R,\theta))} \left(\left| F_{\mathbf{s}}(\widetilde{q}^{\text{DPWide}*}) - q \right| \ge c \right) \\ &= \Pr_{\text{DPWide}(\mathbf{s},\varepsilon,(q,-R,R,\theta))} \left(N \left| F_{\mathbf{s}}(\widetilde{q}^{\text{DPWide}*}) - F_{\mathbf{s}}(\widehat{q}) \right| \ge c N \right) \\ &\le \frac{2R \exp(-\varepsilon c N/2)}{2\theta \exp(-\varepsilon c N/2)} \\ &= \frac{R \exp(-\varepsilon c N/2)}{\theta} \end{aligned}$$

Since $\tilde{q}^{\text{DPWide}}$ is at most θ away from $\tilde{q}^{\text{DPWide*}}$, we can expand the interval by θ on each side and obtain the desired bound for $\tilde{q}^{\text{DPWide}}$.

The widening parameter θ needs to be carefully chosen. All outputs within θ of the target quantile are given the same utility score, so a large θ represents a lower bound on the performance. Conversely, choosing θ too small may result in the area around the target quantile not being given sufficient weight in the sampled distribution. We describe how to set θ for DPWideTS in Section E.1.

Appendix C. Finite-Sample Convergence of U-Statistic

In this section, we develop a finite-sample convergence bound for U-statistics (Definition 4.0.1) that will be a key component towards proving the finite-sample convergence bound for DPWideTS. The bound is stated in Theorem 4.2.1. To develop this bound, we adapt the convergence bounds for linear statistics of i.i.d. mean-0 random variables (Chen et al., 2011) to work for U-statistics of independent, *non-identical* random variables.

We will start by pointing out another fact about U-statistics.

Fact C.0.1 (Sen (1968)) For all $z \in \mathbb{R}$, U_z is a non-increasing function.

Next, we analyze the expectation and variance of the U-statistic in question.

C.1. Expectation and Variance of U-statistic

Here, we consider the expectation and variance of U_z (Equation 2) in a very general form. These expressions are not fully evaluated in order to be as general as possible. They can be further

characterized (as shown in Section E) based on features of the x values (number of ties, spacings, etc.) and the paired datapoints that are included within the set S.

Lemma C.1.1 Let x_1, \ldots, x_n satisfy Assumption 1.1.1 and have variance σ_x^2 , and let y_1, \ldots, y_n be the corresponding response variables under the model $y_i = \beta_0 + \beta_1 x_i + e_i$, where $\beta_0, \beta_1 \in \mathbb{R}$ and each e_i is sampled i.i.d from a continuous, symmetric, mean-0 distribution F_e with variance σ_e^2 . Let S be a set of unordered pairs of datapoints, let N = |S|, and let U_z be defined accordingly as in Definition 4.0.1, Equation (2). Let F_{diff} refer to the CDF of the difference in any two i.i.d. noise variables e_j, e_i , and let $\Delta_{ij} = x_j - x_i$. Then, we have that

$$\mathbf{E}_{e_1,\dots,e_n}[U_z] = \frac{1}{N} \sum_{\{i,j\}\in S, i < j} \operatorname{sign}(x_j - x_i) \cdot (1 - 2 \cdot F_{\operatorname{diff}}(z \cdot \Delta_{ij}))$$

Proof Using the definition of F_{diff} , we can expand the expectation of U_z as follows.

$$\begin{split} \mathbf{E}_{e_1,\dots,e_n}[U_z] &= \frac{1}{N} \sum_{\{i,j\} \in S, i < j} \mathbf{E}_{e_i,e_j} \left[\operatorname{sign}(x_j - x_i) \cdot \operatorname{sign}\left(e_j - e_i - z \cdot \Delta_{ij}\right) \right] \\ &= \frac{1}{N} \sum_{\{i,j\} \in S, i < j} \operatorname{sign}(x_j - x_i) \cdot \left(1 - 2 \cdot P_{e_j,e_i}\left(e_j - e_i < z \cdot \Delta_{ij}\right)\right) \\ &= \frac{1}{N} \sum_{\{i,j\} \in S, i < j} \operatorname{sign}(x_j - x_i) \cdot \left(1 - 2 \cdot F_{\operatorname{diff}}(z \cdot \Delta_{ij})\right) \end{split}$$

The next lemma bounds the variance of the U-statistic, U_z .

Lemma C.1.2 Let x_1, \ldots, x_n satisfy Assumption 1.1.1 and have variance σ_x^2 , and let y_1, \ldots, y_n be the corresponding response variables under the model $y_i = \beta_0 + \beta_1 x_i + e_i$, where $\beta_0, \beta_1 \in \mathbb{R}$ and each e_i is sampled i.i.d from a continuous, symmetric, mean-0 distribution F_e with variance σ_e^2 . Let S be a set of unordered pairs of datapoints, let N = |S|, and let U_z be defined accordingly as in Definition 4.0.1, Equation (2). Let $B_{ij}(z) = sign(x_j - x_i) \cdot sign(e_j - e_i - z \cdot (x_j - x_i))$. Then, we have that

$$\frac{1}{N^2} \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}|=1}} \operatorname{Cov} \left(B_{ij}(z), B_{st}(z)\right) \le \operatorname{Var}_{e_1, \dots, e_n} \left[U_z\right] \le \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}|=1}} \operatorname{Cov} \left(B_{ij}(z), B_{st}(z)\right) \le \operatorname{Var}_{e_1, \dots, e_n} \left[U_z\right] \le \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}|=1}} \operatorname{Cov} \left(B_{ij}(z), B_{st}(z)\right) \le \operatorname{Var}_{e_1, \dots, e_n} \left[U_z\right] \le \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}|=1}} \operatorname{Cov} \left(B_{ij}(z), B_{st}(z)\right) \le \operatorname{Var}_{e_1, \dots, e_n} \left[U_z\right] \le \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}|=1}} \operatorname{Cov} \left(B_{ij}(z), B_{st}(z)\right) \le \operatorname{Var}_{e_1, \dots, e_n} \left[U_z\right] \le \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}|=1}} \operatorname{Cov} \left(B_{ij}(z), B_{st}(z)\right) \le \operatorname{Var}_{e_1, \dots, e_n} \left[U_z\right] \le \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{i,j\} \in S, s < t \\ \{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}|=1}} \operatorname{Cov} \left(B_{ij}(z), B_{st}(z)\right) \le \operatorname{Var}_{e_1, \dots, e_n} \left[U_z\right] \le \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{i,j\} \in S, s < t \\ \{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}|=1}} \operatorname{Var}_{e_1, \dots, e_n} \left[U_z\right] \le \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{i,j\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}|=1}} \operatorname{Var}_{e_1, \dots, e_n} \left[U_z\right] \le \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{i,j\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}|=1}}} \operatorname{Var}_{e_1, \dots, e_n} \left[U_z\right] \le \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{i,j\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}|=1}}} \operatorname{Var}_{e_1, \dots, e_n} \left[U_z\right] \le \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{i,j\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}|=1}}} \operatorname{Var}_{e_1, \dots, e_n} \left[U_z\right] \le \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{i,j\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}|=1}}} \operatorname{Var}_{e_1, \dots, e_n} \left[U_z\right] = \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{i,j\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}|=1}}} \operatorname{Var}_{e_1, \dots, e_n} \left[U_z\right] = \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{i,j\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}|=1}}} \operatorname{Var}_{e_1, \dots, e_n} \left[U_z\right] = \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{i,j\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}|=1}}} \operatorname{Var}_{e_1, \dots, e_n} \left[U_z\right] = \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{i,j\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}|=1}}} \operatorname{Var}_{e_1, \dots, e_n} \left[U_z\right] = \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{i,j\} \in S, s < t$$

Proof We begin by rewriting the variance of U_z as a sum of covariances. Then, we can split up the sum based on the number of overlaps of datapoints i, j, s, t where $\{i, j\}, \{s, t\} \in S$.

$$\operatorname{Var}_{e_1,\ldots,e_n} \left[U_z \right]$$

= $\frac{1}{N^2} \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S, s < t}} \operatorname{Cov} \left(B_{ij}(z) \cdot B_{st}(z) \right)$

$$= \frac{1}{N^2} \left(\sum_{\substack{\{i,j\}\in S, i$$

The first sum in the second line can be evaluated by noting that |S| = N and since $\Pr_{e_i,e_j}[B_{ij} \in \{-1,1\}] = 1$, $\operatorname{Var}(B_{ij}(z)) \leq 1$. For the third sum, note that for $|\{i,j\} \cap \{s,t\}| = 0$, $B_{ij}(z)$ and $B_{st}(z)$ are independent, so $\operatorname{Cov}(B_{ij}(z), B_{st}(z)) = 0$. This leaves the second sum, which can be evaluated based on additional assumptions about the number of ties and spacings between x values, as in Lemma E.0.3. Note that since $\operatorname{Var}(B_{ij}(z)) \geq 0$, we also have that

$$Var_{e_1,...,e_n} [U_z] \ge \frac{1}{N^2} \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}| = 1}} \text{Cov} \left(B_{ij}(z), B_{st}(z)\right)$$

which gives the desired result.

Using Lemma C.1.2, we will evaluate the variance of the null U-statistic, U_0 .

Corollary C.1.3 Let x_1, \ldots, x_n satisfy Assumption 1.1.1 and have variance σ_x^2 , and let y_1, \ldots, y_n be the corresponding response variables under the model $y_i = \beta_0 + \beta_1 x_i + e_i$, where $\beta_0, \beta_1 \in \mathbb{R}$ and each e_i is sampled i.i.d from a continuous, symmetric, mean-0 distribution F_e with variance σ_e^2 . Let S be a set of unordered pairs of datapoints, let N = |S|, and let U_0 be defined accordingly as in Definition 4.0.1, Equation (3). Then, we have that

$$\operatorname{Var}_{e_1,\dots,e_n}[U_0] = \frac{1}{N} + \left(\frac{1}{3 \cdot N^2} \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}| = 1}} \operatorname{sign}(x_j - x_i) \cdot \operatorname{sign}(x_t - x_s) \right)$$

Proof For any $\{i, j\} \in S, i < j$, let $B_{ij}(0) = \operatorname{sign}(x_j - x_i) \cdot \operatorname{sign}(e_j - e_i)$. First, note that $\operatorname{Var}[B_{ij}(0)] = 1$ and $\operatorname{Cov}(B_{ij}(0), B_{st}(0))$ can be expanded as follows.

 $\begin{aligned} \operatorname{Cov} \left(B_{ij}(0), B_{st}(0)\right) \\ &= \operatorname{E}_{e_i, e_j, e_s, e_t}[B_{ij}(0) \cdot B_{st}(0)] - \operatorname{E}_{e_i, e_j}[B_{ij}(0)] \cdot \operatorname{E}_{e_s, e_t}[B_{st}(0)] \\ &= \operatorname{sign}(x_j - x_i) \cdot \operatorname{sign}(x_t - x_s) \cdot \left(\operatorname{E}\left[\operatorname{sign}(e_j - e_i) \cdot \operatorname{sign}(e_t - e_s)\right] - \operatorname{E}\left[\operatorname{sign}(e_j - e_i)\right] \cdot \operatorname{E}\left[\operatorname{sign}(e_t - e_s)\right]\right) \end{aligned}$

It can be shown that for all $\{i, j\} \in S, i < j$ and $\{s, t\} \in S, s < t$ such that $|\{i, j\} \cap \{s, t\}| = 1$, we have that $E[sign(e_j - e_i) \cdot sign(e_t - e_s)] = 1/3$ (Kendall, 1948). In addition, by the symmetric,

mean-0 nature of F_e , we have that $E[sign(e_i - e_i)] = E[sign(e_t - e_s)] = 0$. Plugging these into the characterization of $Var[U_z]$ from Lemma C.1.2 gives the desired result.

1

$$\begin{aligned} \operatorname{Var}_{e_1,\dots,e_n}[U_0] &= \frac{1}{N^2} \left(\sum_{\{i,j\} \in S, i < j} \operatorname{Var}\left(B_{ij}(0)\right) \right) + \left(\frac{1}{N^2} \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}| = 1}} \operatorname{Cov}\left(B_{ij}(0), B_{st}(0)\right) \right) \\ &= \frac{1}{N} + \left(\frac{1}{3 \cdot N^2} \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}| = 1}} \operatorname{sign}(x_j - x_i) \cdot \operatorname{sign}(x_t - x_s) \right) \end{aligned}$$

C.2. Berry-Esseen-type bound for U-statistic

Now, we move onto developing the finite-sample convergence bound for independent, non-identical U-statistics, shown in Theorem 4.2.1. First, we state a standard Berry-Esseen theorem for independent, mean-0, (not necessarily identical) statistics.

Theorem C.2.1 (Berry-Esseen bound for independent mean-0 r.v.s (Berry, 1941; Esseen, 1942)) Let ζ_1, \ldots, ζ_n be independent random variables with $E[\zeta_i] = 0$, $E[\zeta_i^2] > 0$, and $E[|\zeta_i|^3] = \rho_i^3 < \infty$, for $i \in [n]$. Let $\sum_{i=1}^n \operatorname{Var}(\zeta_i) = 1$ and $\rho^3 = \sum_{i=1}^n \rho_i^3$. Let $W = \sum_{i=1}^n \zeta_n$ and Φ be the standard normal cdf. Then,

$$\sup_{t \in \mathbb{R}} |\Pr(W \le t) - \Phi(t)| \le C \cdot \rho^3$$

where C > 0 is a universal constant.

Now, to prove Theorem 4.2.1, we begin by stating an inequality presented by Chen et al. (2011) that bounds the difference between the cumulative distributions of a non-linear function T and a linear approximation function W. Note that this inequality, stated below in Lemma C.2.2, does not require i.i.d random variables, but the subsequent theorems in Chen et al. (2011) that rely on this bound do require identical marginals.

Lemma C.2.2 (Chen et al. (2011)) Let ζ_1, \ldots, ζ_n be independent random variables satisfying $E[\zeta_i] = 0$, $E[\zeta_i^2] > 0$, and $E[|\zeta_i|^3] < \infty$ for $i \in [n]$. Let $\sum_{i=1}^n Var(\zeta_i) = 1$, $W = \sum_{i=1}^n \zeta_i$, and $T = W + \Delta$ for some $\Delta := \Delta(\zeta_1, \ldots, \zeta_n)$. For each $i \in [n]$, let Δ_i be a random variable such that ζ_i and $(W - \zeta_i, \Delta_i)$ are independent. Then, for all $t \in \mathbb{R}$,

$$\left| \Pr(T \le t) - P(W \le t) \right| \le 2\sum_{i=1}^{n} \mathbb{E}[\zeta_{i}^{2}] \mathbb{1}_{|\zeta_{i}| > 1} + 2\sum_{i=1}^{n} \mathbb{E}[|\zeta_{i}|^{3}] \mathbb{1}_{|\zeta_{i}| \le 1} + \mathbb{E}[|W\Delta|] + \sum_{i=1}^{n} \mathbb{E}[|\zeta_{i}(\Delta - \Delta_{i})|]$$

Using the inequality above in Lemma C.2.2, along with a standard Berry-Esseen theorem (Theorem C.2.1) to replace $\Pr[W \leq t]$ with $\Phi(t)$, Chen et al. (2011) develop a Berry-Esseen bound for non-linear statistics of i.i.d. mean-0 random variables. We will adapt this bound for non-identical yet independent, mean-0 random variables to match the setting of our U-statistic, U_z .

To do so, we first state some definitions that allow us to rewrite U_z as the sum of a linear approximation function W and a remainder Δ . These include defining the variables Ψ_1, \ldots, Ψ_n and ζ_1, \ldots, ζ_n . To help parse the notation in the following lemma, note that each Ψ_i is conditioned on the corresponding noise variable e_i , so Ψ_1, \ldots, Ψ_n are independent. In addition, note that the expectation μ_{Ψ} and variance σ_{Ψ} of the sum of the Ψ_i 's can be related to the expectation and variance of U_z (as shown in Lemma C.2.4). The linear approximation function W is the sum of $\zeta_i, i \in [n]$, which are normalized versions of the Ψ_i random variables.

Lemma C.2.3 Let S be a set of unordered pairs of datapoints from $(x_i, y_i)_{i=1}^n$, where |S| = N = nk/2. For $\{i, j\} \in S, i < j$, let $a_{ij} = sign(x_j - x_i)$, $b_{ij} = sign(e_j - e_i)$, and $c_{ij}(z) = sign(e_j - e_i - z \cdot (x_j - x_i))$, and $B'_{ij} = E_{e'}[sign(e' - e_i - z \cdot (x_j - x_i)) | e_i]$ where e' is a fresh draw from the distribution F_e . Then, for $i \in [n]$, let

$$\Psi_i = \frac{1}{k} \sum_{j:\{i,j\} \in S} a_{ij} \cdot B'_{ij} \quad and \quad \zeta_i = \frac{\Psi_i - \mu_i}{\sigma_{\Psi} \cdot n}$$

where $\mu_i = E_{e_1,\dots,e_n}[\Psi_i]$ and $\sigma_i^2 = \operatorname{Var}_{e_1,\dots,e_n}[\Psi_i]$. In addition, let $\mu_{\Psi} = \sum_{i=1}^n \mu_i/n$ and $\sigma_{\Psi}^2 = \sum_{i=1}^n \sigma_i^2/n^2$. Now, we define the following:

$$\bar{h}(i,j) = a_{ij} \cdot c_{ij} - \frac{1}{2} \cdot \Psi_i - \frac{1}{2} \cdot \Psi_j$$

$$\Delta = \frac{1}{2 \cdot \sigma_{\Psi} \cdot N} \sum_{\{i,j\} \in S} \bar{h}(i,j) \quad and \quad \Delta_l = \frac{1}{2 \cdot \sigma_{\Psi} \cdot N} \sum_{\substack{\{i,j\} \in S \\ i,j \neq l}} \bar{h}(i,j)$$

Finally, let $W = \sum_{i=1}^{n} \zeta_i$. Then, we claim the following.

- 1. ζ_1, \ldots, ζ_n are independent
- 2. $W + \Delta = \left(U_z \mu_{\Psi}\right) / (\sigma_{\Psi})$
- *3. For each* $l \in [n]$ *,* ζ_l *is independent of* $(W \zeta_l, \Delta_l)$ *.*

Proof First, note that each ζ_i conditions on the corresponding noise variable e_i , and the only randomness in each ζ_i is a fresh draw e' from the distribution F_e , so ζ_1, \ldots, ζ_n are independent.

Next, using the definitions of Ψ_i and h(i, j, z), we can expand $W + \Delta$ as follows. In particular, note that the Ψ_i and Ψ_j variables cancel out, so we are left with sums of $a_{ij}c_{ij}$ and μ_i .

$$W + \Delta = \sum_{i=1}^{n} \zeta_i + \frac{1}{\sigma_{\Psi} \cdot N} \sum_{\{i,j\} \in S} \bar{h}(i,j)$$
$$= \frac{1}{\sigma_{\Psi} \cdot n} \sum_{i=1}^{n} (\Psi_i - \mu_i) + \frac{1}{2 \cdot \sigma_{\Psi} \cdot N} \sum_{\{i,j\} \in S} \left(a_{ij} c_{ij} - \frac{1}{2} \cdot \Psi_i - \frac{1}{2} \cdot \Psi_j \right)$$

$$= \frac{1}{2 \cdot \sigma_{\Psi} \cdot N} \sum_{\{i,j\} \in S} a_{ij} c_{ij} - \frac{1}{\sigma_{\Psi} \cdot n} \sum_{i=1}^{n} \mu_i$$
$$+ \frac{1}{\sigma_{\Psi} \cdot n} \sum_{i=1}^{n} \Psi_i - \frac{1}{4 \cdot \sigma_{\Psi} \cdot N} \sum_{\{i,j\} \in S} (\Psi_i + \Psi_j)$$
$$= \frac{1}{\sigma_{\Psi} \cdot N} \sum_{\{i,j\} \in S, i < j} a_{ij} c_{ij} - \frac{1}{\sigma_{\Psi} \cdot n} \sum_{i=1}^{n} \mu_i$$
$$= \frac{U_z - \mu_{\Psi}}{\sigma_{\Psi}}$$

Finally, note that for each $l \in [n]$, the random variables $W - \zeta_l$ and Δ_l are functions of ζ_j , $j \neq l$. As ζ_1, \ldots, ζ_n are independent of each other, this means that ζ_l is independent of $(W - \zeta_l, \Delta_l)$.

Below, we relate the quantities μ_{Ψ} and σ_{Ψ}^2 to $\mu(z)$ and $\sigma^2(z)$, the expectation and variance of U_z .

Lemma C.2.4 Let S be the set of unordered pairs of datapoints from $(x_i, y_i)_{i=1}^n$, and let N = |S|. Let $\mu_{\Psi}, \sigma_{\Psi}$ be defined as in Lemma C.2.2. In addition, let U_z be defined as in (2), namely $\mu(z) = E_{e_1,\dots,e_n}[U_z]$, and $\sigma^2(z) = \operatorname{Var}_{e_1,\dots,e_n}[U_z]$. Then,

$$\mu_{\Psi} = \mu(z)$$
 and $\sigma_{\Psi}^2 = \sigma^2(z) + \Theta(1/N)$

Proof First, we show that $\mu_{\Psi}(z)$ can be rewritten as $\mu(z)$. Let $B_{ij} = E_{e'}[e_j - e_i - z(x_j - x_i) | e_i]$ and $B'_{ij} = E_{e'}[e' - e_i - z(x_j - x_i) | e_i]$, where e' is a fresh draw of the random variable e. Note that as e_j and e' are identical random variables, both of which are independent from e_i , $B_{ij} = B'_{ij}$. Then, we have that

$$\mu_{\Psi} = \frac{1}{n} \sum_{i=1}^{n} \mathbf{E}_{e_1,\dots,e_n} \left[\Psi_i \right]$$
$$= \frac{1}{n} \sum_{i=1}^{n} \mathbf{E}_{e_1,\dots,e_n} \left[\frac{1}{k} \sum_{j:\{i,j\} \in S} a_{ij} \cdot B'_{ij} \right]$$
$$= \frac{2}{nk} \sum_{\{i,j\} \in S, i < j} a_{ij} \cdot B_{ij}$$
$$= E_{e_1,\dots,e_n} \left[U_z \right]$$
$$= \mu(z)$$

Next, for $\sigma(z)$, we know from Lemma C.1.2 that

$$\frac{1}{N^2} \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}|=1}} \operatorname{Cov} \left(B_{ij}(z), B_{st}(z)\right) \le \sigma^2(z) \le \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}|=1}} \operatorname{Cov} \left(B_{ij}(z), B_{st}(z)\right) \le \sigma^2(z) \le \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}|=1}} \operatorname{Cov} \left(B_{ij}(z), B_{st}(z)\right) \le \sigma^2(z) \le \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}|=1}} \operatorname{Cov} \left(B_{ij}(z), B_{st}(z)\right) \le \sigma^2(z) \le \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}|=1}} \operatorname{Cov} \left(B_{ij}(z), B_{st}(z)\right) \le \sigma^2(z) \le \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}\}|=1}} \operatorname{Cov} \left(B_{ij}(z), B_{st}(z)\right) \le \sigma^2(z) \le \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}\}|=1}} \operatorname{Cov} \left(B_{ij}(z), B_{st}(z)\right) \le \sigma^2(z) \le \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}\}|=1}} \operatorname{Cov} \left(B_{ij}(z), B_{st}(z)\right) \le \sigma^2(z) \le \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}\}|=1}} \operatorname{Cov} \left(B_{ij}(z), B_{st}(z)\right) \le \sigma^2(z) \le \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}\}|=1}} \operatorname{Cov} \left(B_{ij}(z), B_{st}(z)\right) \le \sigma^2(z) \le \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}\}|=1}} \operatorname{Cov} \left(B_{ij}(z), B_{st}(z)\right) \le \sigma^2(z) \le \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}\}|=1}} \operatorname{Cov} \left(B_{ij}(z), B_{st}(z)\right) \le \sigma^2(z) \le \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}\}|=1}} \operatorname{Cov} \left(B_{ij}(z), B_{st}(z)\right) \le \sigma^2(z) \le \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}\}|=1}} \operatorname{Cov} \left(B_{ij}(z), B_{st}(z)\right) \le \sigma^2(z) \le \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}\}|=1}} \operatorname{Cov} \left(B_{ij}(z), B_{st}(z)\right) \le \sigma^2(z) \le \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}\}|=1}} \operatorname{Cov} \left(B_{ij}(z), B_{st}(z)\right) \le \sigma^2(z) \ge \frac{1}{N} + \frac{1}{N^2} \sum_{\substack{\{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}\}|=1}} \operatorname{Cov} \left(B_{ij}(z), B_{st}(z)\right)$$

In addition, note that $Cov(B'_{ij}, B''_{st}) = Cov(B_{ij}, B_{st})$. To see this, notice that e' is identical to e_j , and both are independent from the other random variables e_i, e_s, e_t, e'' , so e', e_j are interchangeable

in this expression. Similarly, e'' is identical to e_t , and both are independent from e_i, e_j, e', e_s , so e'' and e_t are interchangeable in this expression.

Then, note that we can rewrite σ_{Ψ} as follows.

$$\begin{split} \sigma_{\Psi}^{2} &= \frac{1}{n^{2}} \sum_{i=1}^{n} \operatorname{Var}[\Psi_{i}] \\ &= \frac{1}{n^{2}} \sum_{i=1}^{n} \left(\frac{1}{k^{2}} \sum_{\substack{j:\{i,j\} \in S \\ t:\{i,t\} \in S}} \operatorname{Cov}(B'_{ij}, B''_{il}) \right) \\ &= \frac{1}{n^{2}k^{2}} \sum_{i=1}^{n} \left(\sum_{\substack{j:\{i,j\} \in S \\ t:\{i,t\} \in S}} \operatorname{Cov}(B'_{ij}, B''_{ij}) + \sum_{\substack{j:\{i,j\} \in S \\ t:\{i,t\} \in S, t \neq j}} \operatorname{Cov}(B'_{ij}, B''_{ij}) + \frac{4}{n^{2}k^{2}} \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S, s < t \\ |\{i,j\} \cap \{s,t\}| = 1}} \operatorname{Cov}(B'_{ij}, B''_{ij}) + \frac{1}{N^{2}} \cdot \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S \\ |\{i,j\} \cap \{s,t\}| = 1}} \operatorname{Cov}(B_{ij}, B''_{ij}) + \frac{1}{N^{2}} \cdot \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S \\ |\{i,j\} \cap \{s,t\}| = 1}} \operatorname{Cov}(B_{ij}, B''_{ij}) + \frac{1}{N^{2}} \cdot \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S \\ |\{i,j\} \cap \{s,t\}| = 1}} \operatorname{Cov}(B_{ij}, B''_{ij}) + \frac{1}{N^{2}} \cdot \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S \\ |\{i,j\} \cap \{s,t\}| = 1}} \operatorname{Cov}(B_{ij}, B''_{ij}) + \frac{1}{N^{2}} \cdot \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S \\ |\{i,j\} \cap \{s,t\}| = 1}} \operatorname{Cov}(B_{ij}, B''_{ij}) + \frac{1}{N^{2}} \cdot \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S \\ |\{i,j\} \cap \{s,t\}| = 1}} \operatorname{Cov}(B_{ij}, B''_{ij}) + \frac{1}{N^{2}} \cdot \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S \\ |\{i,j\} \cap \{s,t\}| = 1}} \operatorname{Cov}(B_{ij}, B''_{ij}) + \frac{1}{N^{2}} \cdot \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S \\ |\{i,j\} \cap \{s,t\}| = 1}}} \operatorname{Cov}(B_{ij}, B''_{ij}) + \frac{1}{N^{2}} \cdot \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S \\ |\{i,j\} \cap \{s,t\}| = 1}}} \operatorname{Cov}(B_{ij}, B''_{ij}) + \frac{1}{N^{2}} \cdot \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S \\ |\{i,j\} \cap \{s,t\}| = 1}}} \operatorname{Cov}(B_{ij}, B''_{ij}) + \frac{1}{N^{2}} \cdot \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S \\ |\{i,j\} \cap \{s,t\}| = 1}}} \operatorname{Cov}(B_{ij}, B''_{ij}) + \frac{1}{N^{2}} \cdot \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S \\ |\{i,j\} \cap \{s,t\}| = 1}}} \operatorname{Cov}(B_{ij}, B''_{ij}) + \frac{1}{N^{2}} \cdot \sum_{\substack{\{i,j\} \in S, i < j \\ \{s,t\} \in S \\ |\{i,j\} \cap \{s,t\}| = 1}}} \operatorname{Cov}(B_{ij}, B''_{ij}) + \frac{1}{N^{2}} \cdot \sum_{\substack{\{i,j\} \in S \\ \{s,t\} \in S \\ |\{i,j\} \cap \{s,t\}| = 1}}} \operatorname{Cov}(B_{ij}, B''_{ij}) + \frac{1}{N^{2}} \cdot \sum_{\substack{\{i,j\} \in S \\ \{s,t\} \in S \\ |\{i,j\} \cap \{s,t\}| = 1}}} \operatorname{Cov}(B_{ij}, B''_{ij}) + \frac{1}{N^{2}} \cdot \sum_{\substack{\{i,j\} \in S \\ \{s,t\} \in S \\ |\{i,j\} \cap \{s,t\}| = 1, n \in S }}} \operatorname{Cov}(B_{ij}, B''_{ij}) + \frac{1}{N^{2}} \cdot \sum_{\substack{\{i,j\} \in S$$

As we can see, this gives the desired result.

Next, we use bounds for $E[\Delta^2]$ and $E[(\Delta - \Delta_l)^2]$ shown by Chen et al. (2011). The proof is lengthy so it is omitted here.

Lemma C.2.5 (Chen et al. (2011)) Let Δ , Δ_l and ζ_l (for $l \in [n]$) be defined as in Lemma C.2.3. In addition, for U_z as defined in (2), let $\sigma^2(z) = \operatorname{Var}_{e_1,\ldots,e_n}[U_z]$, and let $\sigma_{\Psi}^2 = \sum_{l=1}^n \operatorname{Var}[\Psi_l]/n^2$. Then,

$$\mathbf{E}\left[\Delta^{2}\right] \leq \frac{\sigma^{2}(z)}{2(n-1) \cdot \sigma_{\Psi}^{2}} \quad and \quad \mathbf{E}\left[(\Delta - \Delta_{l})^{2}\right] \leq \frac{2\sigma^{2}(z)}{2n(n-1) \cdot \sigma_{\Psi}^{2}}$$

The above lemma allows us to bound the right side of the inequality in Lemma C.2.2.

Lemma C.2.6 Let W, Δ, Δ_l , and ζ_l (for $l \in [n]$) be defined as in Lemma C.2.3, where $\rho^3 = \sum_{l=1}^{n} E[|\zeta_l|^3]$. Let γ be defined as follows.

$$\gamma = C \cdot \rho^3 + 2\sum_{i=1}^n \mathbf{E}[\zeta_i^2] \mathbb{1}_{|\zeta_i| > 1} + 2\sum_{i=1}^n \mathbf{E}[|\zeta_i|^3] \mathbb{1}_{|\zeta_i| \le 1} + \mathbf{E}[|W\Delta|] + \sum_{l=1}^n \mathbf{E}[|\zeta_l(\Delta - \Delta_l)|]$$

where C > 0 is a universal constant. In addition, let $\sigma^2(z) = \operatorname{Var}_{e_1,\ldots,e_n}[U_z]$, where U_z is defined in (2). Then, for sufficiently large $n, \gamma = O(1/(n^2\sigma^3(z)))$.

Proof First, note that $|\Psi_i - \mu_i| \le 2$ for all $i \in [n]$, because this term consists of a product and expectation of signs. Therefore, we have that

$$|\zeta_i| = \left|\frac{\Psi_i - \mu_i}{2 \cdot \sigma_{\Psi} \cdot n}\right| \le \frac{1}{\sigma_{\Psi} \cdot n}, \quad \rho^3 = \sum_{i=1}^n \operatorname{E}\left[|\zeta_i|^3\right] \le \frac{1}{\sigma_{\Psi}^3 \cdot n^2}$$

Note that $\sigma_{\Psi} = \Theta(1/\sqrt{n})$, so $\zeta_i = \Theta(1/\sqrt{n})$. Thus, for sufficiently large n, we have that

$$\sum_{i=1}^{n} \mathbf{E}[\zeta_{i}^{2}] \cdot \mathbb{1}_{|\zeta_{i}| > 1} + 2\sum_{i=1}^{n} \mathbf{E}[|\zeta_{i}|^{3}] \cdot \mathbb{1}_{|\zeta_{i}| \le 1} = 0 + \rho^{3} \le \frac{1}{\sigma_{\Psi}^{3} \cdot n^{2}}$$

Next, note that $E[W] = E[\sum_{i=1}^{n} \zeta_i] = 0$. Therefore,

$$E[W^{2}] = Var[W] = \frac{1}{4 \cdot \sigma_{\Psi}^{2} \cdot n^{2}} \sum_{i \in [n]} \sigma_{i}^{2} = O\left(\frac{n^{2} \sigma_{\Psi}^{2}}{\sigma_{\Psi}^{2} n^{2}}\right) = O(1)$$

Using this and the fact from Lemma C.2.5 that $E[\Delta^2] = O\left(\sigma^2(z)/n \cdot \sigma_{\Psi}^2\right)$, we have that

$$\begin{split} \mathbf{E}[|W\Delta|] &\leq \mathbf{E}[|W||\Delta|] \\ &\leq \sqrt{\mathbf{E}\left[|W|^2\right] \cdot \mathbf{E}\left[|\Delta|^2\right]} \\ &= \sqrt{\mathbf{E}\left[W^2\right] \cdot \mathbf{E}\left[\Delta^2\right]} \\ &= \sqrt{\mathbf{E}\left[W^2\right] \cdot \mathbf{E}\left[\Delta^2\right]} \\ &= O\left(\frac{\sigma(z)}{\sqrt{n} \cdot \sigma_{\Psi}}\right) \end{split}$$

Similarly, using the fact from Lemma C.2.5 that $E[(\Delta - \Delta_l)^2] = O(\sigma^2(z)/n^2 \cdot \sigma_{\Psi}^2)$, we have that

$$\begin{split} \sum_{l \in [n]} \mathrm{E}\left[|\zeta_l \cdot (\Delta - \Delta_l)|\right] &\leq \sum_{l \in [n]} \sqrt{\mathrm{E}\left[\zeta_l^2\right] \cdot \mathrm{E}\left[(\Delta - \Delta_l)^2\right]} \\ &= \sum_{l \in [n]} \sqrt{\frac{\sigma_l^2}{n^2 \cdot \sigma_\Psi^2} \cdot O\left(\frac{\sigma^2(z)}{n^2 \cdot \sigma_\Psi^2}\right)} \\ &= O\left(\frac{\sigma(z)}{n^2 \cdot \sigma_\Psi^2}\right) \cdot \sum_{l \in [n]} \sigma_l \\ &= O\left(\frac{\sigma(z)}{n^2 \cdot \sigma_\Psi^2}\right) \cdot O(n) \\ &= O\left(\frac{\sigma(z)}{n \cdot \sigma_\Psi^2}\right) \end{split}$$

where the second to last line follows from noting that $\sigma_l \in (0, 1]$. This gives us that

$$\gamma = O\left(\frac{1}{n^2 \cdot \sigma_{\Psi}^3}\right) + O\left(\frac{\sigma(z)}{\sqrt{n} \cdot \sigma_{\Psi}}\right) + O\left(\frac{\sigma(z)}{n \cdot \sigma_{\Psi}^2}\right) = O\left(\frac{1}{n^2 \cdot \sigma^3(z)}\right)$$

where the second step follows from noting that $\sigma_{\Psi} < 1$ for sufficiently large *n*, and by replacing σ_{Ψ} with $\sigma(z)$ via Lemma C.2.4.

Below, we put all the above lemmas together to state and prove the finite-sample convergence of the distribution of the U-statistic to the standard normal distribution.

Theorem [4.2.1] Let x_1, \ldots, x_n satisfy Assumption 1.1.1 (ie. not all equal) and let y_1, \ldots, y_n be the corresponding response variables under the model $y_i = \beta_0 + \beta_1 x_i + e_i$, where $\beta_0, \beta_1 \in \mathbb{R}$ and each e_i is sampled i.i.d from a continuous, symmetric, mean-0 distribution. Let S be a set of unordered pairs of datapoints, and let U_z be defined with respect to S as in (2). Let $\mu(z) = \mathbb{E}_{e_1,\ldots,e_n}[U_z]$, and $\sigma^2(z) = \operatorname{Var}_{e_1,\ldots,e_n}[U_z]$. Then, for sufficiently large n,

$$\sup_{t \in \mathbb{R}} \left| \Pr\left[\frac{U_z - \mu(z)}{\sigma(z)} \le t \right] - \Phi(t) \right| = O\left(\frac{1}{n^2 \cdot \sigma^3(z)} \right)$$

where Φ is the cdf of a standard normal distribution.

Proof Recall from Lemma C.2.3 the following definitions: for $a_{ij} = \text{sign}(x_j - x_i)$ and $B'_{ij} = E_{e'} [\text{sign} (e' - e_i - z \cdot (x_j - x_i)) | e_i]$, where e' is a fresh draw of the random variable e, we let

$$\Psi_i = \frac{1}{k} \sum_{j:\{i,j\} \in S} a_{ij} \cdot B'_{ij} \text{ and } \zeta_i = \frac{\Psi_i - \mu_i}{\sigma_{\Psi} \cdot n}$$

where $\mu_i = E_{e_1,\dots,e_n}[\Psi_i]$, $\sigma_i = \operatorname{Var}_{e_1,\dots,e_n}[\Psi_i]$, and $\sigma_{\Psi}^2 = \sum_{i=1}^n \sigma_i^2/n^2$. From these definitions, it can be shown that $E[\zeta_i] = 0$, $E[\zeta_i^2] > 0$, $E[|\zeta_i|^3] < \infty$, and $\sum_{i=1}^n \operatorname{Var}[\zeta_i] = 1$. In addition, Lemma C.2.4 shows that for $\mu_{\Psi} = \sum_{i=1}^n \mu_i/n$,

$$\mu(z) = \mu_{\Psi}, \quad \sigma^2(z) = \sigma_{\Psi}^2 + \Theta(1/N),$$

where N = |S|. By Lemma C.2.3, we therefore have that

$$\frac{U_z - \mu(z)}{\sigma(z)} = W + \Delta,$$

and for each $i \in [n]$, ζ_i is independent of $(W - \zeta_i, \Delta_i)$. Then, putting together Lemma C.2.2 and Lemma C.2.6 directly gives the desired result.

Appendix D. Finite-Sample Convergence Bound for DPWideTS

Here, we prove our main result: the finite-sample convergence bound for $\tilde{\beta}_1^{\text{DPWideTS}}$. As described earlier in Section 4.1, the proof of Theorem 4.0.2 follows in two main steps; first, we argue that $U(\tilde{\beta}_1^{\text{DPWideTS}})$ is close to 0 with high probability by utility theorem of the exponential mechanism (Theorem B.0.3). Second, we use the convergence of the U-statistic distribution (Theorem 4.2.1) to show that $\tilde{\beta}_1^{\text{DPWideTS}}$ is close to β_1 .

Theorem [4.0.2] Let x_1, \ldots, x_n satisfy Assumption 1.1.1 and have empirical variance σ_x^2 . Let y_1, \ldots, y_n be the corresponding response variables under the model $y_i = \beta_0 + \beta_1 x_i + e_i$, where $\beta_0, \beta_1 \in \mathbb{R}$ and each e_i is sampled i.i.d from a continuous, symmetric, mean-0 distribution F_e . Let $\tilde{\beta}_1^{DPWideTS} = DPTheilSen(\{x_i, y_i\}_{i=1}^n, \varepsilon, (S, DPWide, \theta, -R, R))$, where $\varepsilon, R, \theta > 0, \beta_1 \in [-R + \theta, R - \theta]$, and $S \in {[n] \choose 2}$ is a set of unordered pairs with distinct x-values. Let U_z be the U-statistic defined with respect to S.

Then, there exists a constant c > 0 such that for sufficiently large n, with probability at least 1 - p, $\beta_1^{\text{DPWideTS}} \in [\beta_1 - z - \theta, \beta_1 + z + \theta]$ for every z that satisfies the following:

$$\frac{-\mu(z) - \frac{4\ln(4R/p\theta)}{\varepsilon \cdot n}}{\sigma(z)} \ge \Phi^{-1} \left(1 - \frac{p}{4} + \frac{c}{n^2 \cdot \sigma^3(z)}\right)$$

where $\mu(z) = E[U_z], \sigma^2(z) = Var[U_z]$, and Φ is the standard normal cdf.

Proof First, we consider the event that $\widetilde{\beta_1}^{\text{DPWideTS}} > \beta_1 + z + \theta$. We take a union bound over two possibilities that could lead to this event: the first is that $U(\widetilde{\beta_1}^{\text{DPWideTS}} - \theta)$ is less than an arbitrary value $-c \in (-1,0)$ (ie. DPWide returned an output that is more than cN/2 away in rank from the median of the N slopes), which implies that $\widetilde{\beta_1}^{\text{DPWideTS}}$ is not within z of β_1 with high probability. The second that $U(\widetilde{\beta_1}^{\text{DPWideTS}} - \theta)$ is within c of 0, yet $\widetilde{\beta_1}^{\text{DPWideTS}} - \theta$ is still more than z greater than β_1 . We can simplify these expressions by noting that U is a non-increasing function (Fact C.0.1). Then, for all $z \in \mathbb{R}$ and sufficiently large n, we have that

where F_s is the empirical distribution function for the set of slopes computed by DPWideTS. Then, the first term in the last line comes from Theorem B.0.3 and the second term follows from Lemma 4.2.1. We can run through a similar analysis for the event that $\tilde{\beta}_1^{\text{DPWideTS}} < \beta_1 - z - \theta$ to get the following:

$$\Pr_{Y, \text{DPWide}} \left[\widetilde{\beta_1}^{\text{DPWideTS}} < \beta_1 - z - \theta \right] \le \frac{R}{\theta} \exp\left(-\varepsilon \cdot c \cdot N/4 \right) + \Phi\left(\frac{c - \mu(-z)}{\sigma(-z)} \right) + O\left(\frac{1}{n^2 \cdot \sigma^3(-z)} \right)$$

Setting the first terms less than or equal to p/4 and then solving for c and z gives the desired result. Finally, the expressions for $\mu(z)$ and $\sigma(z)$ follow directly from Lemmas C.1.1 and C.1.2. This result is not very easily interpretable in the general case. In the next section, we apply this bound to a special setting, and in Section 5, we provide intuition and comparison with convergence bounds for other non-DP and DP algorithms.

Appendix E. Evaluating DPWideTS Bound for Special Case

We consider the special setting where the x-values are evenly split between the endpoints of an interval of length Δ_x . The DPWideTS algorithm computes $N = \lfloor n/2 \rfloor \cdot \lceil n/2 \rceil$ slopes. This section evaluates the bound from Theorem 4.0.2 for this special setting.

To do so, we first solve for the expectation and variance of the U-statistic for this case so that we can plug these values into the general convergence bound. We rely on the following approximation of the normal CDF.

Lemma E.0.1 (Normal CDF approximation) Let $F(\cdot)$ be the cumulative distribution function for a Gaussian with mean β_1 and variance σ^2 . For all y such that $|y - \beta_1|/\sigma$ is sufficiently small, we have that

$$F(y) = \frac{1}{2} + \frac{y - \beta_1}{\sigma\sqrt{2\pi}} + O\left(\frac{(y - \beta_1)^3}{\sigma^3}\right)$$

Proof For any $y \in \mathbb{R}$, the distribution function F(y) is defined as follows.

$$F(y) = \frac{1}{2} \left(1 + \operatorname{erf}\left(\frac{y - \beta_1}{\sigma\sqrt{2}}\right) \right)$$

The error function erf(z) can be rewritten using a power series expansion. For z close to 0, we have:

$$\operatorname{erf}(z) = \frac{2}{\sqrt{\pi}} \int_0^z e^{-t^2} dt$$

= $\frac{2}{\sqrt{\pi}} \int_0^z \left(1 - t^2 + \frac{t^4}{2!} + \dots\right) dt$
= $\frac{2}{\sqrt{\pi}} \left(z - \frac{z^3}{3} + \dots\right)$
= $\frac{2}{\sqrt{\pi}} z + O(z^3)$

Setting $z = (y - \beta_1)/(\sigma\sqrt{2})$, we have that when z is close to 0,

$$F(y) = \frac{1}{2} \left(1 + \frac{\sqrt{2}(y - \beta_1)}{\sqrt{\pi}\sigma} + O\left(\frac{(y - \beta_1)^3}{\sigma^3}\right) \right)$$
$$= \frac{1}{2} + \frac{y - \beta_1}{\sigma\sqrt{2\pi}} + O\left(\frac{(y - \beta_1)^3}{\sigma^3}\right)$$

which gives us the desired result.

Now, we can evaluate the expectation and variance of U_z for this special setting.

Lemma E.0.2 Suppose that x_1, \ldots, x_n be at two endpoints of an interval of size Δ_x such that they satisfy Assumption 5.0.1, and let e_1, \ldots, e_n be drawn i.i.d from $F_e = \mathcal{N}(0, \sigma_e^2)$ according to Assumption 5.0.2. Let y_1, \ldots, y_n be the corresponding response variables under the model $y_i = \beta_0 + \beta_1 x_i + e_i$, where $\beta_0, \beta_1 \in \mathbb{R}$. Let S be the set of unordered pairs of datapoints, and let U_z be defined accordingly as in (2). Then, we have that for sufficiently small $z \cdot \Delta_x / \sigma_e$,

$$\mathbf{E}_{e_1,\dots,e_n}\left[U_z\right] = \frac{-z \cdot \Delta_x}{\sqrt{\pi}\sigma_e} + O\left(\frac{z^3 \cdot \Delta_x^3}{\sigma_e^3}\right)$$

Proof First, let F_{diff} refer to the CDF of the difference in any two i.i.d. noise variables e_j, e_i . In addition, let N = |S|. Then, we have from Lemma C.1.1 that

$$\mathbf{E}_{e_1,\dots,e_n}[U_z] = \frac{1}{N} \sum_{\{i,j\} \in S, i < j} \operatorname{sign}(\Delta_x) \cdot (1 - 2 \cdot F_{\operatorname{diff}}(z \cdot \Delta_x))$$

Next, note that for our special case (assumptions 5.0.1 and 5.0.2), $sign(\Delta_x) = 1$ and F_{diff} is the CDF of the difference between two i.i.d noise variables from a $\mathcal{N}(0, \sigma_e^2)$ distribution. Therefore, using a Taylor approximation of the normal cdf from Lemma E.0.1, we have that for sufficiently small $z \cdot \Delta_x / \sigma_e$,

$$\begin{aligned} \mathbf{E}_{e_1,\dots,e_n} \left[U_z \right] &= \frac{1}{N} \sum_{\{i,j\} \in S, i < j} \left(1 - 2 \cdot F_{\text{diff}}(z \cdot \Delta_x) \right) \\ &= \frac{1}{N} \sum_{\{i,j\} \in S, i < j} \left(1 - 2 \left(\frac{1}{2} + \frac{z \Delta_x}{2\sqrt{\pi}\sigma_e} + O\left(\frac{z^3 \Delta_x^3}{\sigma_e^3} \right) \right) \right) \\ &= \frac{-z \cdot \Delta_x}{\sqrt{\pi}\sigma_e} + O\left(\frac{z^3 \cdot \Delta_x^3}{\sigma_e^3} \right) \end{aligned}$$

which gives the desired result.

Lemma E.0.3 Suppose that x_1, \ldots, x_n be at two endpoints of an interval of size Δ_x such that they satisfy Assumption 5.0.1, and let e_1, \ldots, e_n be drawn i.i.d from $F_e = \mathcal{N}(0, \sigma_e^2)$ according to Assumption 5.0.2. Let y_1, \ldots, y_n be the corresponding response variables under the model $y_i = \beta_0 + \beta_1 x_i + e_i$, where $\beta_0, \beta_1 \in \mathbb{R}$. Let S be the set of unordered pairs of datapoints, and let U_z be defined accordingly as in (2). Then, for sufficiently small $z \cdot \Delta_x / \sigma_e$,

$$\operatorname{Var}_{e_1,\dots,e_n}\left[U_z\right] = \frac{4}{3n} + O\left(\frac{z \cdot \Delta_x}{n \cdot \sigma_e}\right) + O\left(\frac{1}{n^2}\right)$$

Proof Let $B_{ij}(z) = \text{sign}(\Delta_x) \cdot c_{ij}(z)$, where $c_{ij} = \text{sign}(e_j - e_i - z \cdot \Delta_x)$. Starting with the result from Lemma C.1.2, we have that

$$\operatorname{Var}_{e_1, \dots, e_n}[U_z] \le \left(\frac{1}{N^2} \sum_{\substack{\{i, j\} \in S \\ \{s, t\} \in S \\ |\{i, j\} \cap \{s, t\}| = 1}} \operatorname{Cov} \left(B_{ij}(z), B_{st}(z) \right) \right) + \frac{1}{N}$$

Note that under our additional assumptions 5.0.1 and 5.0.2, $sign(\Delta_x) = 1$ and $N = \lfloor n/2 \rfloor \cdot \lceil n/2 \rceil$. Next, for $|\{i, j\} \cap \{s, t\}| = 1$ (and supposing i = s), we have that

$$\operatorname{Cov} (B_{ij}(z), B_{st}(z)) \leq \operatorname{E}_{e_1, \dots, e_n} [B_{ij}(z) \cdot B_{st}(z)]$$

= $\operatorname{E}_{e_i, e_j, e_t} [c_{ij}(z) \cdot c_{it}(z)]$
= $\operatorname{E}_{e_i} [E_{e_j, e_t} [c_{ij}(z) \cdot c_{it}(z) \mid e_i]]$
= $\operatorname{E}_{e_i} [E_{e_j} [c_{ij}(z) \mid e_i] \cdot E_{e_t} [c_{it}(z) \mid e_i]]$
= $\operatorname{E}_{e_i} [(1 - 2 \cdot F_e(e_i + z \cdot \Delta_x))^2]$

The same expression follows for the case where j = t instead of i = s. Note that F_e is the cumulative distribution function for $e_i, i \in [n]$, which by Assumption 5.0.2 is $\mathcal{N}(0, \sigma_e^2)$.

Setting $w = z\Delta_x$, let $g(w) = \mathbb{E}_{e_i} \left[(1 - 2 \cdot F_e(e_i + w))^2 \right]$. Now, we can rewrite e_i as $F_e^{-1}(U)$, where U is a uniform [0, 1] random variable, and then compute g(0) as follows.

$$g(w) = \mathcal{E}_U \left[\left(1 - 2 \cdot F_e \left(F_e^{-1}(U) + w \right) \right)^2 \right]$$

$$g(0) = E_U \left[\left(1 - 2 \cdot F_e (F_e^{-1}(U)) \right)^2 \right] = E_U \left[(1 - 2 \cdot U)^2 \right] = 1/3$$

Next, we know that for any constant c > 0, if $|g'(w)| \le c$ for all $w \in \mathbb{R}$, then

$$g(w) \in (g(0) - cw, g(0) + cw)$$

To bound g'(w), we can first express it as follows.

$$g'(w) = \mathcal{E}_{U} \left[2 \cdot \left(1 - 2 \cdot F_{e} \left(F_{e}^{-1}(U) + w \right) \right) \cdot \left(-2 \cdot F_{e}' \left(F_{e}^{-1}(U) + w \right) \right) \right]$$
$$= \mathcal{E}_{U} \left[-4 \cdot \left(1 - 2 \cdot F_{e} \left(F_{e}^{-1}(U) + w \right) \right) \cdot \frac{1}{\sigma_{e} \cdot \sqrt{2\pi}} \cdot \exp \left(\frac{-\left(F_{e}^{-1}(U) + w \right)^{2}}{2\sigma_{e}^{2}} \right) \right]$$

where the second line follows by noting that F'_e is a pdf of a $\mathcal{N}(0, \sigma_e^2)$ random variable. As a cumulative distribution function, $F_e(\cdot)$ is between 0 and 1. In addition, we can see that the exp factor is between 0 and 1. Therefore, we have that

$$|g'(w)| \le 4 \cdot \frac{1}{\sigma_e \cdot \sqrt{2\pi}} = O\left(\frac{1}{\sigma_e}\right), \text{ so } g(w) = g(0) + O\left(\frac{w}{\sigma_e}\right)$$

Finally, the number of terms that share one datapoint, ie. $\#\{i, j\} \in S, \{s, t\} \in S$ s.t. $|\{i, j\} \cap \{s, t\}| = 1$, divided by the N^2 factor, is less than or equal to

$$\frac{2 \cdot (n/2)^2 \cdot (n/2 - 1)}{(n/2)^4} = \frac{4(n-2)}{n^2}$$

Putting this all together, we have that for $w = z\Delta_x$, the following holds.

$$\operatorname{Var}_{e_1,\dots,e_n} \left[U(\delta_n, (x_i, y_i)_{i=1}^n) \right] \le \left(\frac{1}{N^2} \sum_{\substack{\{i,j\} \in S \\ \{s,t\} \in S \\ |\{i,j\} \cap \{s,t\}|=1}} \operatorname{Cov} \left(B_{ij}(z), B_{st}(z) \right) \right) + \frac{1}{N}$$

$$\leq \left(\frac{1}{N^2} \sum_{\substack{\{i,j\} \in S \\ \{s,t\} \in S \\ \{s,t\} \in S \\ |\{i,j\} \cap \{s,t\}| = 1}} \operatorname{E}_{e_i} \left[(1 - 2 \cdot F_e(e_i + z \cdot \Delta_x))^2 \right] \right) + \frac{1}{N}$$

$$= \left(\frac{1}{N^2} \sum_{\substack{\{i,j\} \in S \\ \{s,t\} \in S \\ |\{i,j\} \cap \{s,t\}| = 1}} g(w) \right) + \frac{1}{N}$$

$$\leq \frac{1}{(n/2)^4} \cdot 2 \cdot (n/2)^2 \cdot (n/2 - 1) \cdot \left(\frac{1}{3} + O\left(\frac{w}{\sigma_e}\right) \right) + \frac{1}{(n/2)^2}$$

$$= \frac{4(n-2)}{n^2} \left(\frac{1}{3} + O\left(\frac{w}{\sigma_e}\right) \right) + O\left(\frac{1}{n^2}\right)$$

$$\leq \frac{4}{3n} + O\left(\frac{z \cdot \Delta_x}{n \cdot \sigma_e}\right) + O\left(\frac{1}{n^2}\right)$$

Using the above characterizations of the expectation and variance, we can now restate and prove the convergence bound for this special case.

Theorem [1.2.1] Let x_1, \ldots, x_n be at two endpoints of an interval of size Δ_x such that they satisfy Assumption 5.0.1, and let e_1, \ldots, e_n be drawn i.i.d from $F_e = \mathcal{N}(0, \sigma_e^2)$ according to Assumption 5.0.2. Let y_1, \ldots, y_n be the corresponding response variables under the model $y_i = \beta_0 + \beta_1 x_i + e_i$, where $\beta_0, \beta_1 \in \mathbb{R}$. Let $\tilde{\beta}_1^{\text{DPWideTS}} = \text{DPTheilSen}(\{x_i, y_i\}_{i=1}^n, \varepsilon, (\text{DPWide}, \theta, -R, R))$, where $\varepsilon, R, \theta > 0$, and $\beta_1 \in [-R + \theta, R - \theta]$ as in Assumption B.0.2. Let τ be defined as follows.

$$\tau = \Phi^{-1} \left(1 - \frac{p}{8} \right) \cdot \sqrt{\frac{4}{3n}} + \frac{8 \ln(4R/p\theta)}{\varepsilon n}$$

where Φ^{-1} is the inverse standard normal distribution function. Then, we have that for sufficiently large n, where for some constant c, $n \ge c \left(\log(R/p\theta) / \varepsilon \sqrt{\log(1/p)} \right)^2$, and sufficiently small τ , we have that with probability at least 1 - p,

$$\widetilde{\beta_1}^{\text{DPWideTS}} \in [\beta_1 - z - \theta, \beta_1 + z + \theta]$$

for

$$z = \frac{\sqrt{\pi} \cdot \sigma_e}{\Delta_x} \cdot \left(\tau + O\left(\tau^{3/2}\right)\right).$$

Proof We start with the bound from Theorem 4.0.2 and substitute the expressions for $\mu(z)$ and $\sigma^2(z)$ using Lemmas E.0.2 and E.0.3. This gives the following: for sufficiently large n and sufficiently

small $z \cdot \Delta_x / \sigma_e$, $\widetilde{\beta_1}^{\text{DPWideTS}} \in [\beta_1 - z - \theta, \beta_1 + z + \theta]$ provided that

$$z \ge \frac{\sqrt{\pi} \cdot \sigma_e}{\Delta_x} \cdot \left(\Phi^{-1} \left(1 - \frac{p}{4} + \frac{c_1}{n^2 \cdot \sigma^3(z)} \right) \cdot \sqrt{\left(\frac{4}{3n} + e_{\sigma^2}(z) \right) + \frac{8\ln(4R/p\theta)}{\varepsilon n} + e_\mu(z)} \right)$$
(5)

where

$$e_{\sigma^2}(z) \leq \frac{c_2 \cdot \Delta_x \cdot z}{n \cdot \sigma_e} + \frac{c_3}{n^2} \quad \text{and} \quad e_{\mu}(z) \leq \frac{c_4 \cdot \Delta_x^3 \cdot z^3}{\sigma_e^3}$$

and where $c_1, c_2, c_3, c_4 > 0$ are constants. Now, we will show that z can actually be set so as to satisfy condition (5). In particular, it can be shown that setting z as

$$z := \frac{2\sqrt{\pi} \cdot \sigma_e}{\Delta_x} \cdot \left(\tau + c_2 \cdot \tau^{3/2}\right) \quad \text{for} \quad \tau = \Phi^{-1} \left(1 - \frac{p}{8}\right) \cdot \sqrt{\frac{4}{3n}} + \frac{8\ln(4R/p\theta)}{\varepsilon n}$$

satisfies condition (5). To see this, first note that for any $z \in \mathbb{R}$, $\sigma^2(z) \ge 4/3n$ by Lemma E.0.3. Then, for $n \ge (6c_1/p)^2$, we have that

$$\Phi^{-1}\left(1 - \frac{p}{4} + \frac{c_1}{n^2 \cdot \sigma^3(z)}\right) \le \Phi^{-1}\left(1 - \frac{p}{8}\right).$$

Using this, we can see that for $n \ge \max(3c_3/4, 8c_2^2(\Phi^{-1}(1-p/8))^2, (6c_1/p)^2)$, we can bound the highest-order term on the right-hand side of (5) as follows.

$$\begin{split} \frac{\sqrt{\pi} \cdot \sigma_e}{\Delta_x} \cdot \Phi^{-1} \left(1 - \frac{p}{4} + \frac{c_1}{n^2 \cdot \sigma^3(z)} \right) \cdot \sqrt{\left(\frac{4}{3n} + e_{\sigma^2}(z)\right)} \\ &\leq \frac{\sqrt{\pi} \cdot \sigma_e}{\Delta_x} \cdot \Phi^{-1} \left(1 - \frac{p}{8} \right) \cdot \left(\sqrt{\frac{4}{3n}} + \sqrt{\frac{2 \cdot c_2 \cdot \tau}{n}} + \sqrt{\frac{2 \cdot c_2^2 \cdot \tau^{3/2}}{n}} + \sqrt{\frac{c_3}{n^2}} \right) \\ &\leq \frac{\sqrt{\pi} \cdot \sigma_e}{\Delta_x} \cdot \tau \cdot \left(1 + \sqrt{3 \cdot c_2 \cdot \tau} + \sqrt{3} \cdot c_2 \cdot \tau^4 \right) \\ &\leq \frac{2\sqrt{\pi} \cdot \sigma_e}{\Delta_x} \cdot \left(\tau + c_2 \cdot \tau^{3/2} \right) \\ &= z \end{split}$$

where the first inequality follows from the condition that $n > (6c_1/p)^2$, the second inequality follows for $n > 3c_3/4$ and the definition of τ , and the third inequality follows from $n > 8(c_2 \cdot \Phi^{-1}(1-p/8))^2$. From this, we can see that there exists a setting of z that satisfies condition (5), which completes the proof.

E.1. Choosing the widening parameter, θ

Theorem 1.2.1 offers insight on how to set θ in the DPWideTS algorithm, which was an open question raised by Alabi et al. (2022). For fixed $p, \varepsilon, R, \sigma_e, \Delta_x$, and n that satisfy the conditions stated in the Theorem, we can set θ to minimize the bound as follows:

$$\theta \approx \max\left(\frac{\sigma_e \cdot \ln(R\varepsilon n\Delta_x/p\sigma_e)}{\varepsilon n\Delta_x}, \frac{R}{p}\exp(-\varepsilon n)\right)$$

The first term in the max comes from allowing the two terms involving θ in Thm. 1.2.1 to be approximately equal, whereas the second term in the max comes from upper bounding $O(\ln(R/p\theta)/\varepsilon n)$ by some constant. The reason for having both terms is to account for both widespread and concentrated slopes.¹⁴ The factor $\sigma_e/(n\Delta_x)$ in the first term corresponds to the standard deviation of the slopes computed by DPWideTS; when the slopes are highly concentrated, the first term becomes small. The second term, however, is independent of σ_e and Δ_x , which allows θ to remain bounded away from 0 and prevents a blowup in the convergence bound. Note that R, ε, n and p are known in practice; if the experimental design suggests that the slopes may be concentrated (e.g. if the x-values are located at one of two endpoints of an interval as in this special case), our analysis suggests setting θ to scale with the second term.

Appendix F. Confidence Intervals

In this section, we consider DP confidence intervals for the linear regression slope, β_1 . We begin by defining the type of confidence intervals we are interested in.

Definition F.0.1 (Confidence intervals for simple linear regression.) Let $d = (x_i, y_i)_{i=1}^n$ be a dataset real-valued pairs, where x_1, \ldots, x_n are fixed values that are not all equal, and y_1, \ldots, y_n are the corresponding response variables under the model $y_i = \beta_0 + \beta_1 x_i + e_i$, where $\beta_0, \beta_1 \in \mathbb{R}$ and each e_i is sampled i.i.d from a continuous, symmetric, mean-0 distribution F_e . Let $I_{\mathbb{R}}$ be the set of all intervals in \mathbb{R} , and let $M_{nonpriv}$: Multisets ($\mathbb{R} \times \mathbb{R}, n$) $\times \mathcal{H} \to I_{\mathbb{R}}$ be a deterministic mechanism that outputs an interval. For any $\alpha_1 \in (0, 1)$, $M_{nonpriv}$ outputs a $(1 - \alpha_1)$ -confidence interval for the slope β_1 if for all hyperparams $\in \mathcal{H}$, and sufficiently large n,

$$\Pr_{e_1,\ldots,e_n \sim F_e} \left[\beta_1 \in M_{nonpriv}(\boldsymbol{d}, hyperparams) \right] \ge 1 - \alpha_1$$

where the probability is over the randomness of the dataset d.

Next, let M_{DP} : $Multisets(\mathbb{R} \times \mathbb{R}, n) \times \mathbb{R}_{>0} \times \mathcal{H} \to I_{\mathbb{R}}$ be a randomized ε -DP mechanism that outputs an interval. For any $\alpha \in (0, 1)$, M_{DP} outputs a $(1 - \alpha)$ -DP confidence interval for the slope β_1 if for all privacy loss parameters $\varepsilon \in \mathbb{R}_{>0}$, hyperparams $\in \mathcal{H}$, and sufficiently large n,

$$\Pr_{\substack{e_1,\ldots,e_n\sim F_e\\M_{DP}}} [\beta_1 \in M_{DP}(\boldsymbol{d},\varepsilon,hyperparams)] \ge 1 - \alpha$$

where the probability is over the randomness of both the dataset **d** and the mechanism M_{DP} .

F.1. Nonprivate Confidence Intervals for Theil-Sen

Now, we will define the non-private Theil-Sen confidence interval (Sen, 1968) and prove its coverage validity.

Definition F.1.1 Let S be a set of N unordered pairs of points, and let U_0 be the corresponding null U-statistic (defined in)Definition 4.0.1, Equation 3). For the corresponding set of slopes s, let F_s be

^{14.} Handling the case of concentrated slopes was Alabi et al. (2022)'s original motivation for designing the widened exponential mechanism.

the empirical distribution function for s. Let $\sigma^2(0) = \operatorname{Var}_{e_1,\ldots,e_n}[U_0]$ as evaluated in Corollary C.1.3. For any $\alpha_1 \in (0,1)$, let b be defined as follows:

$$b := \frac{1}{2} \cdot \Phi^{-1} \left(1 - \frac{\alpha_1}{8} \right) \cdot \sigma(0)$$

where Φ is the standard normal cdf. Let $\hat{\beta}_{1L} = F_s^{-1}(1/2 - b)$ and $\hat{\beta}_{1U} = F_s^{-1}(1/2 + b)$. Then, the non-private $1 - \alpha_1$ -Theil-Sen confidence interval is $[\hat{\beta}_{1L}, \hat{\beta}_{1U}]$.

The proof of this interval's validity can be found in Lemma F.1.2.

Lemma F.1.2 Let x_1, \ldots, x_n satisfy Assumption 1.1.1 and let y_1, \ldots, y_n be the corresponding response variables under the model $y_i = \beta_0 + \beta_1 x_i + e_i$, where $\beta_0, \beta_1 \in \mathbb{R}$ and each e_i is sampled *i.i.d* from a continuous, symmetric, mean-0 distribution F_e . Let S be a set of N unordered pairs of datapoints, and let U_0 be the corresponding null U-statistic defined in Definition 4.0.1 Equation (3). Let $\sigma^2(0) = \operatorname{Var}_{e_1,\ldots,e_n}[U_0]$ as evaluated in Corollary C.1.3. For any $\alpha_1 \in (0,1)$, let $[\hat{\beta}_{1L}, \hat{\beta}_{1U}]$ be the non-private Theil-Sen confidence interval as in Definition F.1.1. Then, there exists a constant c > 0 such that for all $n \ge c/\alpha_1^2$, $[\hat{\beta}_{1L}, \hat{\beta}_{1U}]$ is a $(1 - \alpha_1)$ -confidence interval for the true slope β_1 .

Proof We will consider the upper limit of the interval, $\hat{\beta}_{1U}$, for simplicity. First, note that

$$U(\hat{\beta}_{1U}) = 1 - 2F_{\mathbf{s}}(\hat{\beta}_{1U}) = 1 - 2F_{\mathbf{s}}(F_{\mathbf{s}}^{-1}(1/2 + b))$$

= $-2b = \Phi^{-1}(\alpha_1/8) \cdot \sigma(0)$

Next, we rely on the fact that if $\beta_1 > \hat{\beta}_{1U}$, $U(\beta_1) < U(\hat{\beta}_{1U})$. Therefore, we can use the distribution of $U(\beta_1)$ and its convergence to a standard normal via Theorem 4.2.1 to evaluate the probability that $\beta_1 > \hat{\beta}_{1U}$.

$$\begin{aligned} \Pr_{\mathbf{s}} \left[\beta_1 > \hat{\beta}_{1U} \right] &= \Pr_{\mathbf{s}} \left[U(\beta_1) < U\left(\hat{\beta}_{1U} \right) \right] \\ &\leq \Phi \left(\frac{U\left(\hat{\beta}_{1U} \right)}{\sigma(0)} \right) + O\left(\frac{1}{n^2 \cdot \sigma^3(0)} \right) \\ &= \Phi \left(\frac{\Phi^{-1}(\alpha_1/8) \cdot \sigma(0)}{\sigma(0)} \right) + O\left(\frac{1}{n^2 \cdot \sigma^3(0)} \right) \\ &= \alpha_1/8 + O\left(\frac{1}{n^2 \cdot \sigma^3(0)} \right) \\ &= \alpha_1/8 + O\left(\frac{1}{\sqrt{n}} \right) \\ &\leq \alpha_1/2 \end{aligned}$$

where the second to last line holds follows from Corollary C.1.3 and the last line holds for $n \ge c/\alpha_1^2$. We can go through a similar argument for $\hat{\beta}_{1L}$, which gives the desired result.

Algorithm F.1 DPTSCI: ε-DP Algorithm

 $\begin{array}{l} \hline \mathbf{Data:} \ \boldsymbol{d} = (x_i, y_i)_{i=1}^n \in \mathrm{Multisets}(\mathbb{R} \times \mathbb{R}, n) \\ \mathbf{Privacy params:} \ \boldsymbol{\varepsilon} \in \mathbb{R}_{\geq 0} \\ \mathbf{Hyperparams:} \ \boldsymbol{S} \in {[n] \choose 2}, \ \mathrm{DPMedCI}, \ \mathrm{hyperparams} \in \mathcal{H} \\ \mathbf{s} = \{\} \\ \mathbf{for} \ each \ \{i, j\} \in S \ (such \ that \ x_i \neq x_j) \ \mathbf{do} \\ \mid \ s = (y_j - y_i)/(x_j - x_i) \\ \mathrm{Add} \ s \ \mathrm{to} \ \mathbf{s} \\ \mathbf{end} \\ \mathrm{Let} \ k = \max_{i \in [n]} \{ \# j \in [n] : \{i, j\} \in S \} \\ [\widetilde{\beta}_{1L}^{\mathrm{TS}}, \widetilde{\beta}_{1U}^{\mathrm{TS}}] = \mathrm{DPMedCI} \ (\mathbf{s}, \boldsymbol{\varepsilon}/k, \ \mathrm{hyperparams}) \\ \mathbf{return} \ [\widetilde{\beta}_{1L}^{\mathrm{TS}}, \widetilde{\beta}_{1U}^{\mathrm{TS}}] \end{array}$

F.2. DPTSCI Algorithm

Below, we provide pseudocode for the general algorithm for confidence intervals via DPTS. Note that this algorithm is similar to the DPTS point estimator (Alg 3.1), except it replaces the call to DPMed with a call to DPMedCI, a DP algorithm for confidence intervals for the median.

Lemma F.2.1 Algorithm F.1 is ε -DP.

Proof The call to DPMedCI is ε/k -DP by definition, so by simple composition and post-processing, DPTSCI is ε -DP.

F.3. Union-bound confidence interval

Now, we consider the coverage validity and width of the DP confidence interval when we use DPWideCIUnion (Alg 6.1) as the DPMedCI algorithm.

Theorem F.3.1 (Validity and width of DPWideTSCIUnion) Let x_1, \ldots, x_n satisfy Assumption 1.1.1, and let y_1, \ldots, y_n be the corresponding response variables under the model $y_i = \beta_0 + \beta_1 x_i + e_i$, where $\beta_0, \beta_1 \in \mathbb{R}$ and each e_i is sampled i.i.d from a continuous, symmetric, mean-0 distribution F_e . Let S be a set of N unordered datapoints. For $\alpha \in (0,1)$, $r_\alpha \in (0,1)$, $\varepsilon, R, \theta > 0$, and $\beta_1 \in [-R + \theta, R - \theta]$ as in Assumption B.0.2, let $[\widetilde{\beta}_{1L}^{DPWideTS}, \widetilde{\beta}_{1U}^{DPWideTS}] =$ DPTSCI($\{x_i, y_i\}_{i=1}^n, \varepsilon, S, DPWideCIUnion, (\alpha, r_\alpha, \theta, -R, R, Union = 1)$). Then, the interval $[\widetilde{\beta}_{1L}^{DPWideTS}, \widetilde{\beta}_{1U}^{DPWideTS}]$ is a $1 - \alpha$ -confidence interval for the true slope β_1 over the randomness in $(x_i, y_i)_{i=1}^n$ and DPWideTSCIUnion.

In addition, let U_z be defined as in Definition 4.0.1, Equation (2) according to set S. Then, there exists a constant v > 0 such that for sufficiently large n, we have that with probability at least $1 - \alpha$, $\tilde{\beta}_{1U}^{\text{DPWideTS}} - \tilde{\beta}_{1L}^{\text{DPWideTS}} < 2z + 2\theta$ for all z that satisfies the following.

$$\frac{-\mu(z) - \frac{8\ln(2R/\alpha\theta)}{\varepsilon \cdot n}}{\sigma(z)} \ge 2 \cdot \Phi^{-1} \left(1 - \frac{\alpha}{2} + \frac{v}{n^2 \cdot \sigma^3(z)} \right)$$

where $\mu(z) = \mathbb{E}[U_z], \sigma^2(z) = \operatorname{Var}[U(z)], and \Phi$ is the standard normal cdf.

Proof First, let us prove the validity of the confidence interval

 $[\tilde{\beta}_{1L}^{\text{DPWideTS}}, \tilde{\beta}_{1U}^{\text{DPWideTS}}]$. We consider the upper limit of the interval, $\tilde{\beta}_{1U}^{\text{DPWideTS}}$, for simplicity. First, we will bound the probability that the upper limit of the private interval fails ($\tilde{\beta}_{1U}^{\text{DPWideTS}} < \beta_1$) conditioned on the success of the upper limit of the non-private interval ($\beta_1 < \hat{\beta}_{1U}$). This analysis uses a translation between the U-statistic and the distribution function F_s , as well as the utility of the exponential mechanism (Theorem B.0.3) with target quantile $q_U = 1/2 + b + c$.

$$\begin{aligned} &\Pr_{\boldsymbol{d}, \mathsf{DPWideCI}} \left(\widetilde{\beta}_{1U}^{\text{DPWideIS}} < \beta_1 \land \beta_1 \leq \widehat{\beta}_{1U} \right) \\ &\leq \mathsf{E}_{\boldsymbol{d}} \left[\Pr_{\mathsf{DPWideCI}} \left(\widetilde{\beta}_{1U}^{\text{DPWideTS}} < \widehat{\beta}_{1U} \mid \boldsymbol{d} \right) \right] \\ &\leq \mathsf{E}_{\boldsymbol{d}} \left[\Pr_{\mathsf{DPWideCI}} \left(U(\widetilde{\beta}_{1U}^{\text{DPWideTS}}) > U(\widehat{\beta}_{1U}) \mid \boldsymbol{d} \right) \right] \\ &= \mathsf{E}_{\boldsymbol{d}} \left[\Pr_{\mathsf{DPWideCI}} \left(1 - 2F_{\mathbf{s}}(\widetilde{\beta}_{1U}^{\text{DPWideTS}}) > -2b \mid \boldsymbol{d} \right) \right] \\ &= \mathsf{E}_{\boldsymbol{d}} \left[\Pr_{\mathsf{DPWideCI}} \left(F_{\mathbf{s}}(\widetilde{\beta}_{1U}^{\text{DPWideTS}}) < 1/2 + b \mid \boldsymbol{d} \right) \right] \\ &= \mathsf{E}_{\boldsymbol{d}} \left[\Pr_{\mathsf{DPWideCI}} \left(F_{\mathbf{s}}(\widetilde{\beta}_{1U}^{\text{DPWideTS}}) - q_U < -c \mid \boldsymbol{d} \right) \right] \\ &\leq \mathsf{E}_{\boldsymbol{d}} \left[\alpha_2/2 \right] \\ &= \alpha_2/2 \end{aligned}$$

We use this bound to evaluate the total failure probability of the private upper limit. We also use the failure probability of the upper limit of the *non-private* confidence interval, $\hat{\beta}_{1U}$, as characterized in Lemma F.1.2.

$$\begin{aligned} &\Pr_{\boldsymbol{d}, \text{DPWideCI}} \left[\widehat{\beta}_{1U}^{\text{DPWideTS}} < \beta_1 \right] \\ &= \Pr_{\boldsymbol{d}, \text{DPWideCI}} \left[\widehat{\beta}_{1U}^{\text{DPWideTS}} < \beta_1 \land \beta_1 > \widehat{\beta}_{1U} \right] + \Pr_{\boldsymbol{d}, \text{DPWideCI}} \left[\widehat{\beta}_{1U}^{\text{DPWideTS}} < \beta_1 \land \beta_1 \le \widehat{\beta}_{1U} \right] \\ &\leq \Pr_{\boldsymbol{d}} \left[\beta_1 > \widehat{\beta}_{1U} \right] + \Pr_{\text{DPWideCI}} \left[\widehat{\beta}_{1U}^{\text{DPWideTS}} < \widehat{\beta}_{1U} \right] \\ &\leq \alpha_1/2 + \alpha_2/2 \\ &= \alpha/2 \end{aligned}$$

The same analysis holds for the other end of the interval, so this proves the validity of the confidence interval. Next, let us consider the width of the interval. For simplicity, let us consider the event that $\tilde{\beta}_{1U}^{\text{DPWideTS}} > \beta_1 + z + \theta$. The analysis is similar to the proof of Theorem 4.0.2. For all $z \in \mathbb{R}$ and sufficiently large n, we have that

$$\begin{split} &\Pr_{Y,\text{DPWideCI}} \left[\widetilde{\beta}_{1U}^{\text{DPWideTS}} > \beta_1 + z + \theta \right] \\ & \leq \Pr_{Y,\text{DPWideCI}} \left[U(\widetilde{\beta}_{1U}^{\text{DPWideTS}} - \theta) < -d \right] + \\ & \Pr_{Y} \left[U(\widetilde{\beta}_{1U}^{\text{DPWideTS}} - \theta) \geq -d \wedge \widetilde{\beta}_{1U}^{\text{DPWideTS}} - \theta > \beta_1 + z \right] \\ & \text{DPWideCI} \end{split}$$

$$\begin{split} &\leq \Pr_{Y, \text{DPWideCI}} \left[U(\widetilde{\beta_1}_{U}^{\text{DPWideTS}}) < -d \right] + \Pr_{Y} \left[U(\beta_1 + z) \geq -d \right] \\ &\leq \Pr_{Y, \text{DPWideCI}} \left[F_{\mathbf{s}}(\widetilde{\beta_1}_{U}^{\text{DPWideTS}}) - q_U > d/2 - b - c \right] + \Pr_{Y} \left[U(\beta_1 + z) \geq -d \right] \\ &\leq \frac{R}{2\theta} \exp\left(-\varepsilon \cdot (d/2 - b - c) \cdot N/2 \right) + \left(1 - \Phi\left(\frac{-d - \mu(z)}{\sigma(z)}\right) \right) + O\left(\frac{1}{n^2 \cdot \sigma^3(z)}\right) \end{split}$$

where b, c are defined as in Algorithm 6.1. The first term in the last line follows from Theorem B.0.3 and the second term follows from Lemma 4.2.1. Setting this probability to $\leq \alpha$ and solving for d and z gives us the desired result. We can run through a similar analysis for the event that $\tilde{\beta}_{1L}^{\text{DPWideTS}} < \beta_1 - z - \theta$.

F.4. Tighter confidence interval

return q_L, q_U

Now, we consider the coverage validity of the tighter confidence interval, DPWideTSCI, obtained via the following subroutine.

 $\begin{array}{l} \hline \textbf{Algorithm F.2 ComputeExpMechCITargets} \\ \hline \textbf{Data: } s = (s_1, \ldots, s_N) \\ \hline \textbf{Input: } \varepsilon \in \mathbb{R}_{>0}, \alpha \in (0, 1), [-R, R] \subset \mathbb{R}, \theta \in \mathbb{R}_{>0} \\ \textbf{for } t_L \in \mathbb{N}, 1 \leq t_L \leq N/2 \textbf{ do} \\ \mid \gamma_{t_L} = 1 - F_U \left(1 - 2(t_L - 1)/N\right) + \sum_{m=t_L}^n F'_U \left(1 - 2m/N\right) \cdot \frac{R}{\theta} \exp(-(m - t_L) \cdot \varepsilon/2) \\ \textbf{end} \\ // \quad F_U, F'_U \text{ are the CDF and PDF of the null U-statistic corresponding to } s \\ q_L = \max_{t_L \in \mathbb{N}, 1 \leq t_L < \lceil N/2 \rceil} \{t_L : \gamma_{t_L} \leq \alpha/2\}/N \\ \textbf{for } t_U \in \mathbb{N}, N/2 \leq t_U < N \textbf{ do} \\ \mid \gamma_{t_U} = F_U \left(1 - 2t_U/N\right) + \sum_{m=1}^{t_U} F'_U \left(1 - 2m/N\right) \cdot \frac{R}{\theta} \exp(-(t_U - m) \cdot \varepsilon/2) \\ \textbf{end} \\ q_U = \min_{t_U \in \mathbb{N}, \lceil N/2 \rceil < t_U < N} \{j : \gamma_{t_U} \leq \alpha/2\}/N \end{array}$

To analyze this algorithm, we will first state a lemma that translates between the empirical distribution F_s to the distribution of the null U-statistic, F_U .

Lemma F.4.1 Let $s = (s_1, ..., s_N) \in \mathbb{R}^N$ be a multiset of (not necessarily independent) slopes computed via the DPTSCI algorithm, where each s_j is drawn from a marginal distribution function that has true median β_1 . Let F_s be the empirical distribution of the slopes, let U_0 be the corresponding U-statistic (as defined in (3), and let F_U, F'_U be the cdf and pdf, respectively, of $U_0(\beta_1, (x_i, y_i)_{i=1}^n)$. Then, for any $m \in [N]$,

$$\Pr\left[F_{s}(\beta_{1}) = \frac{m}{N}\right] = F'_{U}\left(1 - \frac{2m}{N}\right)$$
$$\Pr\left[F_{s}(\beta_{1}) \leq \frac{m}{N}\right] = 1 - F_{U}\left(1 - \frac{2m}{N}\right).$$

Proof We can rewrite $F_{\mathbf{s}}(\beta_1)$ as $1/2 - U_0(\beta_1)/2$, which directly gives that

$$\Pr\left[F_{\mathbf{s}}(\beta_{1}) = \frac{m}{N}\right] = \Pr\left[U_{0}(\beta_{1}) = 1 - \frac{2m}{N}\right]$$
$$\Pr\left[F_{\mathbf{s}}(\beta_{1}) \le \frac{m}{N}\right] = \Pr\left[U_{0}(\beta_{1}) \ge 1 - \frac{2m}{N}\right].$$

Using the above lemma, we show the validity of the DPWideTSCI using ComputeExpMechCITargets.

Theorem F.4.2 Let x_1, \ldots, x_n satisfy Assumption 1.1.1, and let y_1, \ldots, y_n be the corresponding response variables under the model $y_i = \beta_0 + \beta_1 x_i + e_i$, where $\beta_0, \beta_1 \in \mathbb{R}$ and each e_i is sampled i.i.d from a continuous, symmetric, mean-0 distribution F_e . Let S be a set of unordered pairs of datapoints. For $\alpha \in (0, 1)$, $r_{\alpha} = 1/2$ (as a default, since it won't be used), $\varepsilon, R, \theta > 0$, and $\beta_1 \in [-R + \theta, R - \theta]$ as in Assumption B.0.2, let $[\widetilde{\beta}_{1L}^{DPWideTS}, \widetilde{\beta}_{1U}^{DPWideTS}] = DPTSCI(\{x_i, y_i\}_{i=1}^n, \varepsilon, S, DPWideCI, (\alpha, r_{\alpha}, \theta, -R, R, Union = 0))$. Then, the interval $[\widetilde{\beta}_{1L}^{DPWideTS}, \widetilde{\beta}_{1U}^{DPWideTS}]$ is a $1 - \alpha$ -confidence interval for the true slope β_1 over the randomness in $(x_i, y_i)_{i=1}^n$ and DPWideTSCI.

Proof For simplicity, we consider just the upper endpoint of the interval. Recall that for a given $t_U \in [N]$, $\tilde{U}_{\alpha}(\mathbf{s}, t_U) = \text{DPWide}(\mathbf{s}, \varepsilon/2k, (t_U/N, \theta, -R, R))$, and $F_{\mathbf{s}}$ is the empirical distribution of the set of slopes \mathbf{s} . We will consider the probability that $\tilde{U}_{\alpha}(\mathbf{s}, t_U) + \theta < \beta_1$ by splitting this event into two cases: first, when $F_{\mathbf{s}}(\beta_1) \leq t_U/N$, and second, when $F_{\mathbf{s}}(\beta_1) > t_U/N$.

$$\begin{split} & \Pr_{\mathbf{s}, \mathsf{DPWideCI}} \left[\tilde{U}_{\alpha}(\mathbf{s}, t_{U}) + \theta < \beta_{1} \right] \\ &= \sum_{m=1}^{t_{U}} \Pr_{\mathsf{DPWideCI}} \left[\tilde{U}_{\alpha}(\mathbf{s}, t_{U}) + \theta < \beta_{1} \mid F_{\mathbf{s}}(\beta_{1}) = \frac{m}{N} \right] \cdot \Pr_{\mathbf{s}} \left[F_{\mathbf{s}}(\beta_{1}) = \frac{m}{N} \right] \\ &+ \sum_{m=t_{U}+1}^{N} \Pr_{\mathsf{DPWideCI}} \left[\tilde{U}_{\alpha}(\mathbf{s}, t_{U}) + \theta < \beta_{1} \mid F_{\mathbf{s}}(\beta_{1}) = \frac{m}{N} \right] \cdot \Pr_{\mathbf{s}} \left[F_{\mathbf{s}}(\beta_{1}) = \frac{m}{N} \right] \end{split}$$

We can simplify the first sum using Lemma F.4.1 and the utility of the widened exponential mechanism, Theorem B.0.3:

$$\begin{split} \sum_{m=1}^{m=t_U} \Pr_{\mathbf{DPWideCI}} \left[\tilde{U_{\alpha}}(\mathbf{s}, t_U) + \theta < \beta_1 \mid F_{\mathbf{s}}(\beta_1) = \frac{m}{N} \right] \cdot \Pr_{\mathbf{s}} \left[F_{\mathbf{s}}(\beta_1) = m/N \right] \\ \leq \sum_{m=1}^{m=t_U} \Pr_{\mathbf{DPWideCI}} \left[\tilde{U_{\alpha}}(\mathbf{s}, t_U) < \beta_1 \mid F_{\mathbf{s}}(\beta_1) = \frac{m}{N} \right] \cdot \Pr_{\mathbf{s}} \left[F_{\mathbf{s}}(\beta_1) = m/N \right] \\ = \sum_{m=1}^{m=t_U} \Pr_{\mathbf{DPWideCI}} \left[F_{\mathbf{s}} \left(\tilde{U_{\alpha}}(\mathbf{s}, t_U) \right) < m/N \right] \cdot F'_U \left(1 - \frac{2m}{N} \right) \\ \leq \sum_{m=1}^{m=t_U} \Pr_{\mathbf{DPWideCI}} \left[\frac{t_U}{N} - F_{\mathbf{s}} \left(\tilde{U_{\alpha}}(\mathbf{s}, t_U) \right) > \frac{t_U - m}{N} \right] \cdot F'_U \left(1 - \frac{2m}{N} \right) \end{split}$$

$$\leq \sum_{m=1}^{m=t_U} \frac{R}{\theta} \exp\left(-(t_U - m) \cdot \varepsilon/2\right) \cdot F'_U\left(1 - \frac{2m}{N}\right)$$

For the second sum, we simply upper bound the first probability in the summation by 1 and use Lemma F.4.1 to characterize the distribution of $\mathbb{1}_{F_s(\beta_1)=m/N}$.

$$\sum_{m=t_U+1}^{m=N} \Pr_{\substack{\mathbf{D} \mathsf{P} \mathsf{w} \mathsf{ideCI}}} \left[\tilde{U}_{\alpha}(\mathbf{s}, t_U) + \theta < \beta_1 \mid F_{\mathbf{s}}(\beta_1) = \frac{m}{N} \right] \cdot \Pr_{\mathbf{s}} \left[F_{\mathbf{s}}(\beta_1) = \frac{m}{N} \right]$$
$$\leq \sum_{m=t_U+1}^{m=N} \Pr_{\mathbf{s}} \left[F_{\mathbf{s}}(\beta_1) = \frac{m}{N} \right]$$
$$\leq F_U \left(1 - \frac{2 \cdot t_U}{N} \right)$$

Therefore, we have that

$$\Pr_{\mathbf{s}, \mathsf{DPWideCI}}\left(\tilde{U}_{\alpha}(\mathbf{s}, t_{U}) + \theta < \beta_{1}\right) \leq F_{U}\left(1 - \frac{2 \cdot t_{U}}{N}\right) + \sum_{m=1}^{t_{U}} F_{U}'\left(1 - \frac{2m}{N}\right) \cdot \frac{R}{\theta} \exp\left(-(t_{U} - m) \cdot \varepsilon/2\right)$$

A similar result holds for the other side of the interval. Then, validity of the interval follows directly from the selection of the target quantiles t_L, t_U (viaComputeExpMechCITargets) such that the interval has at least $1 - \alpha$ coverage.