CONNECTED REDUCED PRODUCTS

Miloš S. Kurilić¹

Abstract

A structure $\mathbb{X} = \langle X, \rho \rangle$, where ρ is a binary relation on the set X, is connected iff the minimal equivalence relation containing ρ is the full relation on X. Connectivity is not a first order property but it is expressible by a $L_{\omega_1\omega}$ -sentence. We show that, for a set I the following conditions are equivalent

- (a) |I| is less than the first measurable cardinal,
- (b) For each filter Φ ⊂ P(I) and each family {X_i : i ∈ I} of binary structures, the reduced product Π_Φ X_i is connected, iff there are a finite set K ⊂ I and n ∈ ω such that X_i is connected, for each i ∈ K and {i ∈ I : X_i is of diameter ≤ n} ∪ K ∈ Φ,
- (c) The ultraproduct $\prod_{\mathcal{U}} \mathbb{G}_{\omega}$ is a disconnected graph for each non-principal ultrafilter $\mathcal{U} \subset P(I)$, where \mathbb{G}_{ω} is the linear graph on ω .

Moreover, the implication " \Leftarrow " in (b) holds in ZFC. For a disconnected $\prod_{\Phi} X_i$ we describe the components of $\prod_{\Phi} X_i$.

2010 Mathematics Subject Classification: 03C20

Key words and phrases: Reduced product, connected binary relation, components.

1 Preliminaries

Reduced products Let $L = \{R\}$ be the language with one binary relational symbol. Let I be a non-empty set, $\Phi \subset P(I)$ a filter and $\mathbb{X}_i = \langle X_i, \rho_i \rangle$, for $i \in I$, L-structures. Let \sim_{Φ} be the binary relation on the product $\prod_{i \in I} X_i$ defined in the following way: for $x = \langle x_i : i \in I \rangle$, $y = \langle y_i : i \in I \rangle \in \prod_{i \in I} X_i$

$$x \sim_{\Phi} y \Leftrightarrow \{i \in I : x_i = y_i\} \in \Phi.$$

It is evident that the relation \sim_{Φ} is an equivalence relation on the set $\prod_{i \in I} X_i$. Let $[x]_{\Phi} = \{y \in \prod_{i \in I} X_i : x \sim_{\Phi} y\} \in (\prod_{i \in I} X_i) / \sim_{\Phi}$ denote the equivalence class of x. The corresponding *reduced product* $(\prod_{i \in I} X_i) / \Phi$ (shortly $\prod_{\Phi} X_i$) is the structure

$$\left\langle (\prod_{i\in I} X_i)/\sim_{\Phi}, \rho \right\rangle$$

where the binary relation ρ on the quotient $(\prod_{i \in I} X_i) / \sim_{\Phi}$ is defined by

$$[x]_{\Phi} \ \rho \ [y]_{\Phi} \Leftrightarrow \{i \in I : x_i \ \rho_i \ y_i\} \in \Phi.$$

We will use the following evident fact (see [1], p. 177).

¹Department of Mathematics and Informatics, Faculty of Sciences, University of Novi Sad, Trg Dositeja Obradovića 4, 21000 Novi Sad, Serbia. e-mail: milos@dmi.uns.ac.rs

Fact 1.1 If Φ is a filter on I and $J \in \Phi$, then $\Phi \upharpoonright J := \Phi \cap P(J)$ is a filter on Jand $f : \prod_{\Phi} X_i \cong \prod_{\Phi \upharpoonright J} X_j := (\prod_{i \in J} X_i) / \sim_{\Phi \upharpoonright J}$, where $f([x]_{\Phi}) = [x \upharpoonright J]_{\Phi \upharpoonright J}$.

First-order properties of reduced products are well understood. If L is any language, an L-formula is a *basic Horn formula* iff it is of the form $\eta_0 \vee \ldots \vee \eta_{m-1}$, where η_i 's are literals and at most one of them is atomic. The set of *Horn formulas* is the closure of the set of basic Horn formulas under \land , \exists and \forall . It is known that each Horn formula $\varphi(v_0, \ldots, v_{n-1})$ is *preserved in reduced products*; i.e., for each set I, filter $\Phi \subset P(I)$, family of L-structures $\{\mathbb{X}_i : i \in I\}$ and n-tuple $\langle x^0, \ldots, x^{n-1} \rangle \in (\prod_{i \in I} X_i)^n$ we have

$$\left\{i \in I : \mathbb{X}_i \models \varphi[x_i^0, \dots, x_i^{n-1}]\right\} \in \Phi \implies \prod_{\Phi} \mathbb{X}_i \models \varphi\Big[[x^0]_{\Phi}, \dots, [x^{n-1}]_{\Phi}\Big].$$
(1)

By the well known result of Keisler (under CH) and Galvin (in ZFC) we have: an *L*-sentence φ is *preserved in reduced products*, i.e., for each *I*, Φ and $\{X_i : i \in I\}$ as above

$$\{i \in I : \mathbb{X}_i \models \varphi\} \in \Phi \implies \prod_{\Phi} \mathbb{X}_i \models \varphi, \tag{2}$$

iff φ is logically equivalent to a Horn *L*-sentence (see [1], p. 365).

Putting the implication " \Leftarrow " in (2) we obtain a definition of a sentence *preserved under reduced factors*. It is known that the positive sentences (without \neg) are of that kind (see [1], p. 340).

Connected binary structures Let $L = \{R\}$ be the language with one binary relational symbol. Let $\mathbb{X} = \langle X, \rho \rangle$ be an *L*-structure, where $\rho \subset X^2$ is a reflexive binary relation and let ρ^{-1} be the inverse of ρ . For $\varepsilon \in 2 := \{0, 1\}$ let

$$\rho^{\varepsilon} = \begin{cases} \rho & \text{if } \varepsilon = 0, \\ \rho^{-1} & \text{if } \varepsilon = 1. \end{cases}$$
(3)

If $x, y \in X$ and $n \in \omega$, a *path of length* n + 1 *from* x *to* y is a pair $\langle \overline{z}, \varepsilon \rangle$, where $\overline{z} = \langle z_0, \ldots, z_{n-1} \rangle \in X^n$ and $\varepsilon = \langle \varepsilon_0, \ldots, \varepsilon_n \rangle \in 2^{n+1}$ and such that

$$x \rho^{\varepsilon_0} z_0 \rho^{\varepsilon_1} z_1 \rho^{\varepsilon_2} \dots \rho^{\varepsilon_{n-1}} z_{n-1} \rho^{\varepsilon_n} y_1$$

(So, for n = 0, $x \rho^{\varepsilon_0} y$.) Then we will write $d(x, y) \le n + 1$ and $d(x, y) = \infty$ will denote that there is no (finite) path from x to y.

The binary relation \sim on the set X defined by $x \sim y$ iff there is finite path from x to y, is an equivalence relation, the corresponding equivalence classes are the *(connected) components of* X and the structure X is *connected* iff it has only one component. (It is easy to see that \sim is the transitive closure $(\rho_s)_t$ of the symmetrization $\rho_s := \rho \cup \rho^{-1}$ of the relation ρ .) We extend these definitions for arbitrary *L*-structure $\mathbb{X} = \langle X, \rho \rangle$ in the following way: instead of ρ we take its reflexivization $\rho_R := \rho \cup \Delta_X$ and call \mathbb{X} connected iff $\mathbb{X}_R := \langle X, \rho_R \rangle$ is connected, components of \mathbb{X} are components of \mathbb{X}_R , etc.

The statement $d(x, y) \le n + 1$ is expressible by the first order formula

$$\varphi_{d \le n+1}(x,y) := \exists z_0, \dots, z_{n-1} \bigvee_{\varepsilon \in 2^{n+1}} x \ \rho^{\varepsilon_0} \ z_0 \ \rho^{\varepsilon_1} \dots \rho^{\varepsilon_{n-1}} \ z_{n-1} \ \rho^{\varepsilon_n} \ y.$$

It is evident that $\varphi_{d \le n+1}$ is positive and $\neg \varphi_{d \le n+1}$ is Horn. The property of an *L*-structure that each two points are connected by a path of length $\le n+1$ is expressible by the first order sentence

$$\varphi_n^{\operatorname{conn}} := \forall x, y \; \varphi_{d \le n+1}(x, y).$$

If such $n \in \omega$ exists we will say that \mathbb{X} is *of finite diameter*. The connectedness of an *L*-structure is expressible by the $L_{\omega_1\omega}$ -sentence

$$\varphi^{\operatorname{conn}} := \forall x, y \quad \bigvee_{n \in \omega} \varphi_{d \le n+1}(x, y).$$

2 Connected reduced products. Components

In the sequel we assume that

(*) $I \neq \emptyset$ is a set, Φ is a filter in P(I), $\mathbb{X}_i = \langle X_i, \rho_i \rangle$, $i \in I$, are *L*-structures and $\prod \mathbb{X}_i := \langle \prod_{i \in I} X_i, \sigma \rangle$ and $\prod_{\Phi} \mathbb{X}_i := \langle \prod_{i \in I} X_i / \sim_{\Phi}, \rho \rangle$ are the corresponding direct and reduced product respectively.

Fact 2.1 (*) If $x, y \in \prod_{i \in I} X_i$, $A \subset I$, $x \upharpoonright (I \setminus A) = y \upharpoonright (I \setminus A)$, $n \in \omega$, and there is $\varepsilon \in 2^{n+1}$ such that for each $i \in A$ there is $\overline{z} \in X_i^n$ such that $\langle \overline{z}, \varepsilon \rangle$ is a path from x_i to y_i , then in $\prod_{\Phi} X_i$ there is a path from $[x]_{\Phi}$ to $[y]_{\Phi}$.

Proof. Let $\varepsilon = \langle \varepsilon_0, \ldots, \varepsilon_n \rangle$, and for $i \in A$ let $\langle \langle z_i^0, \ldots, z_i^{n-1} \rangle, \varepsilon \rangle$ be a path from x_i to y_i

$$x_i \,\rho_i^{\varepsilon_0} \, z_i^0 \,\rho_i^{\varepsilon_1} \, z_i^1 \,\rho_i^{\varepsilon_2} \, \dots \,\rho_i^{\varepsilon_{n-1}} \, z_i^{n-1} \,\rho_i^{\varepsilon_n} \, y_i. \tag{4}$$

Let $t^m \in \prod_{i \in I} X_i$, for m < n, be defined by

$$t_i^m = \begin{cases} x_i & \text{if } i \in I \setminus A, \\ z_i^m & \text{if } i \in A. \end{cases}$$
(5)

By (4) in the direct product we have $x \sigma^{\varepsilon_0} t^0 \sigma^{\varepsilon_1} t^1 \sigma^{\varepsilon_2} \dots \sigma^{\varepsilon_{n-1}} t^{n-1} \sigma^{\varepsilon_n} y$ and, hence, in the reduced product $\prod_{\Phi} X_i$ we have

$$[x]_{\Phi} \rho^{\varepsilon_0} [t^0]_{\Phi} \rho^{\varepsilon_1} [t^1]_{\Phi} \rho^{\varepsilon_2} \dots \rho^{\varepsilon_{n-1}} [t^{n-1}]_{\Phi} \rho^{\varepsilon_n} [y]_{\Phi}.$$
(6)

So, this is a path from $[x]_{\Phi}$ to $[y]_{\Phi}$.

Fact 2.2 If $\kappa \geq \omega$ is a non-measurable cardinal and $\Phi \subset P(\kappa)$ a filter consisting of infinite sets, then there is a partition $\kappa = \bigcup_{n \in \omega} I_n$ such that for each $F \in \Phi$ we have $F \cap I_n \neq \emptyset$, for infinitely many $n \in \omega$.

Proof. First let $\bigcap \Phi = \emptyset$. Let $\mathcal{U} \subset P(\kappa)$ be an ultrafilter such that $\Phi \subset \mathcal{U}$. Then the ultrafilter \mathcal{U} is non-principal and, since the cardinal κ is non-measurable, \mathcal{U} is not ω -complete. Thus there is a partition $\kappa = \bigcup_{n \in \omega} I_n$ such that $I_n \notin \mathcal{U}$ and, hence, $\kappa \setminus I_n \in \mathcal{U}$, for each $n \in \omega$. Assuming that there are $F \in \Phi$ and $m \in \omega$ such that $F \subset \bigcup_{n \leq m} I_n$, we would have $F \cap \bigcap_{n \leq m} \kappa \setminus I_n = F \setminus \bigcup_{n \leq m} I_n = \emptyset \in \mathcal{U}$, which is false. Thus, $|\{n \in \omega : F \cap I_n \neq \emptyset\}| = \omega$, for each $F \in \overline{\Phi}$.

Second let $A := \bigcap \Phi \neq \emptyset$. If $|A| < \omega$, then, since $\Phi \subset P(\kappa) \setminus [\kappa]^{<\omega}$, $\Phi' := \Phi \upharpoonright (\kappa \setminus A)$ is a non-principal filter on $\kappa \setminus A$ and we work as above.

If $|A| \ge \omega$, $I_0 = \kappa \setminus A$ and $A = \bigcup_{n \in \mathbb{N}} I_n$ is any partition of A, then for each $F \in \Phi$ we have $F \cap I_n \neq \emptyset$, for all $n \in \mathbb{N}$.

Theorem 2.3 (*) If the size of the index set I is less than the first measurable cardinal, then the following conditions are equivalent

- (a) The reduced product $\prod_{\Phi} X_i$ is connected,
- (b) There are a finite set K ⊂ I and n ∈ ω such that X_i is connected, for each i ∈ K, and {i ∈ I : X_i ⊨ φ_n^{conn}} ∪ K ∈ Φ.

The implication $(b) \Rightarrow (a)$ is true in ZFC (without any assumptions on |I|).

Proof. W.l.o.g. we can assume that $\rho_i \subset X_i$, $i \in I$, are reflexive relations. (a) \Rightarrow (b). We prove the contrapositive: assuming that

$$\forall K \in [I]^{<\omega} \left(\forall i \in K \ (\mathbb{X}_i \models \varphi^{\mathrm{conn}}) \Rightarrow \forall n \in \omega \ \{i \in I : \mathbb{X}_i \models \varphi^{\mathrm{conn}}_n\} \cup K \notin \Phi \right)$$
(7)

we show that $\prod_{\Phi} X_i$ is disconnected.

If the filter Φ contains a finite set, then Φ is a principal filter generated by some $K \in [I]^{<\omega}$ and, by (7), there is $i_0 \in K$ such that \mathbb{X}_{i_0} is disconnected. Since $\Phi \upharpoonright K = \{K\}$, by Fact 1.1 we have $\prod_{\Phi} \mathbb{X}_i \cong \prod_{\Phi \upharpoonright K} \mathbb{X}_i \cong \prod_{i \in K} \mathbb{X}_i$. We take $x_{i_0}, y_{i_0} \in X_{i_0}$ such that there is no path from x_{i_0} to y_{i_0} in \mathbb{X}_{i_0} and choose arbitrary $x_i = y_i \in X_i$, for $i \in K \setminus \{i_0\}$. Assuming that there is a path from $x := \langle x_i : i \in K \rangle$ to $y := \langle y_i : i \in K \rangle$ in $\prod_{i \in K} \mathbb{X}_i$ we would obtain a path from x_{i_0} to y_{i_0} in \mathbb{X}_{i_0} , which is impossible. Thus the reduced product $\prod_{\Phi} \mathbb{X}_i$ is disconnected.

So, the remaining case is when Φ is a filter consisting of infinite subsets of I. Let $A_n := \{i \in I : \mathbb{X}_i \models \varphi_n^{\text{conn}}\}$, for $n \in \omega$. Then $A_0 \subset A_1 \subset A_2 \subset \ldots$ and, defining $I_0 := A_0, I_n := A_n \setminus A_{n-1}$, for $n \in \mathbb{N}$, and $I_\infty = I \setminus \bigcup_{n \in \omega} A_n$, we have

$$I = \bigcup_{n \in \omega} I_n \cup I_{\infty}.$$

We continue with a case analysis.

1. $I_{\infty} \in \Phi$. Then $|I_{\infty}| \ge \omega$ is a non-measurable cardinal, $\Phi \upharpoonright I_{\infty}$ is a filter on I_{∞} consisting of infinite sets, and, by Fact 2.2, there is a partition $I_{\infty} = \bigcup_{n \in \omega} I'_n$ such that for each $F \in \Phi \upharpoonright I_{\infty}$ we have $F \cap I'_n \ne \emptyset$, for infinitely many $n \in \omega$. If $n \in \omega$, then for $i \in I'_n$ we have $i \notin A_n$ and we choose $x_i, y_i \in X_i$ such that $d(x_i, y_i) > n + 1$. For $i \in I \setminus I_{\infty}$ we choose arbitrary $x_i = y_i \in X_i$. Let $x := \langle x_i : i \in I \rangle, y := \langle y_i : i \in I \rangle$ and suppose that in $\prod_{\Phi} X_i$ there is a path from $[x]_{\Phi}$ to $[y]_{\Phi}$

$$[x]_{\Phi} \rho^{\varepsilon_0} [z^0]_{\Phi} \rho^{\varepsilon_1} [z^1]_{\Phi} \rho^{\varepsilon_2} \dots \rho^{\varepsilon_{n-1}} [z^{n-1}]_{\Phi} \rho^{\varepsilon_n} [y]_{\Phi}.$$
(8)

Then there would be $F \in \Phi$ which witnesses that; in other words,

$$\forall i \in F \ x_i \ \rho_i^{\varepsilon_0} \ z_i^0 \ \rho_i^{\varepsilon_1} \ z_i^1 \ \rho_i^{\varepsilon_2} \ \dots \ \rho_i^{\varepsilon_{n-1}} \ z_i^{n-1} \ \rho_i^{\varepsilon_n} \ y_i. \tag{9}$$

Now let m > n be such that there is $i \in F \cap I'_m$. By (9) there is an (n + 1)-path from x_i to y_i , which is impossible by the choice of x_i and y_i $(d(x_i, y_i) > n + 1)$. So, there is no path from $[x]_{\Phi}$ to $[y]_{\Phi}$ in $\prod_{\Phi} X_i$ and this structure is disconnected.

2. $I_{\infty} \notin \Phi$. Then we have $F \cap \bigcup_{n \in \omega} I_n \neq \emptyset$, for all $F \in \Phi$.

2.1. Each $F \in \Phi$ intersects infinitely many sets I_n . If $n \in \mathbb{N}$, then for $i \in I_n$ we have $i \notin A_{n-1}$ and we choose $x_i, y_i \in X_i$ such that $d(x_i, y_i) > n$. For $i \in I \setminus \bigcup_{n \in \mathbb{N}} I_n$ we choose arbitrary $x_i = y_i \in X_i$. Let $x := \langle x_i : i \in I \rangle$, $y := \langle y_i : i \in I \rangle$ and suppose that in $\prod_{\Phi} X_i$ there is a path from $[x]_{\Phi}$ to $[y]_{\Phi}$

$$[x]_{\Phi} \rho^{\varepsilon_0} [z^0]_{\Phi} \rho^{\varepsilon_1} [z^1]_{\Phi} \rho^{\varepsilon_2} \dots \rho^{\varepsilon_{n-1}} [z^{n-1}]_{\Phi} \rho^{\varepsilon_n} [y]_{\Phi}.$$
(10)

Then there would be $F \in \Phi$ which witnesses that; in other words,

$$\forall i \in F \ x_i \ \rho_i^{\varepsilon_0} \ z_i^0 \ \rho_i^{\varepsilon_1} \ z_i^1 \ \rho_i^{\varepsilon_2} \ \dots \ \rho_i^{\varepsilon_{n-1}} \ z_i^{n-1} \ \rho_i^{\varepsilon_n} \ y_i.$$
(11)

By the assumption of 2.1 there is m > n + 1 such that $F \cap I_m \neq \emptyset$. So, for $i \in F \cap I_m$, by (11) there is an (n + 1)-path from x_i to y_i , which is impossible by

the choice of x_i and y_i ($d(x_i, y_i) > m$). So, there is no path from $[x]_{\Phi}$ to $[y]_{\Phi}$ in $\prod_{\Phi} X_i$ and this structure is disconnected.

2.2. There are $n_0 \in \omega$ and $F_0 \in \Phi$ such that $F_0 \subset A_{n_0} \cup I_\infty$. By (7) taking $K = \emptyset$ we obtain $A_n \notin \Phi$, for all $n \in \omega$. So, assuming that $F \cap I_\infty = \emptyset$, for some $F \in \Phi$, we would have $A_{n_0} \in \Phi$, which is false. Thus

$$\forall F \in \Phi \ F \cap I_{\infty} \neq \emptyset, \tag{12}$$

and $\Phi \upharpoonright I_{\infty}$ is a proper filter on I_{∞} .

2.2.1. $|F \cap I_{\infty}| \geq \omega$, for all $F \in \Phi$. Then $|I_{\infty}| \geq \omega$ is a non-measurable cardinal, $\Phi \upharpoonright I_{\infty}$ is a filter on I_{∞} consisting of infinite sets, and, by Fact 2.2, there is a partition $I_{\infty} = \bigcup_{n \in \omega} I'_n$ such that for each $F \in \Phi \upharpoonright I_{\infty}$ we have $F \cap I'_n \neq \emptyset$, for infinitely many $n \in \omega$. Now we continue exactly as in Case 1.

2.2.2. The set $F \cap I_{\infty}$ is finite, for some $F \in \Phi$. Then $\Phi \upharpoonright I_{\infty}$ is a principal filter generated by some finite set $J \subset I_{\infty}$ and, hence, $A_{n_0} \cup J \in \Phi$. So, by (7), there is $i_0 \in J$ such that \mathbb{X}_{i_0} is disconnected. We take $x_{i_0}, y_{i_0} \in X_{i_0}$ such that there is no path from x_{i_0} to y_{i_0} in \mathbb{X}_{i_0} and choose arbitrary $x_i = y_i \in X_i$, for $i \in I \setminus \{i_0\}$. Let $x := \langle x_i : i \in I \rangle$, $y := \langle y_i : i \in I \rangle$ and suppose that there is a path from $[x]_{\Phi}$ to $[y]_{\Phi}$ in $\prod_{\Phi} \mathbb{X}_i$,

$$[x]_{\Phi} \rho^{\varepsilon_0} [z^0]_{\Phi} \rho^{\varepsilon_1} [z^1]_{\Phi} \rho^{\varepsilon_2} \dots \rho^{\varepsilon_{n-1}} [z^{n-1}]_{\Phi} \rho^{\varepsilon_n} [y]_{\Phi}.$$
(13)

Then there would be $F \in \Phi$ which witnesses that; in other words,

$$\forall i \in F \ x_i \ \rho_i^{\varepsilon_0} \ z_i^0 \ \rho_i^{\varepsilon_1} \ z_i^1 \ \rho_i^{\varepsilon_2} \ \dots \ \rho_i^{\varepsilon_{n-1}} \ z_i^{n-1} \ \rho_i^{\varepsilon_n} \ y_i.$$
(14)

But $i_0 \in J \subset F$ and we obtain a path from x_{i_0} to y_{i_0} in \mathbb{X}_{i_0} , which is impossible. Thus the reduced product $\prod_{\Phi} \mathbb{X}_i$ is disconnected.

(b) \Rightarrow (a). Let $K \in [I]^{<\omega}$, $n \in \omega$ and, defining $A := \{i \in I : \mathbb{X}_i \models \varphi_n^{\text{conn}}\}$, suppose that

(i) \mathbb{X}_i is connected, for each $i \in K$, and

(ii) $A \cup K \in \Phi$.

Since $\varphi_n^{\text{conn}} \Rightarrow \varphi^{\text{conn}}$ we can assume that $A \cap K = \emptyset$. For $x = \langle x_i : i \in I \rangle, y = \langle y_i : i \in I \rangle \in \prod_{i \in I} X_i$ we construct a path from $[x]_{\Phi}$ to $[y]_{\Phi}$ in $\prod_{\Phi} X_i$.

If $i \in A$, then $\mathbb{X}_i \models \varphi_n^{\text{conn}}$ and, since $x_i, y_i \in X_i$, there is a path $\langle \bar{z}, \varepsilon \rangle$ of length n + 1 from x_i to y_i ; thus we can choose one such ε . Let $\varepsilon^k \in 2^{n+1}$, k < l, be an enumeration of the n + 1-tuples chosen in this way for $i \in A$. Then defining $A_{\varepsilon^k} = \{i \in A : \varepsilon^k \text{ is chosen for } x_i \text{ and } y_i\}$, for k < l, we have

$$A = \bigcup_{k < l} A_{\varepsilon^k}.$$

Let $y' \in \prod_{i \in I} X_i$ be defined by

$$y'_{i} = \begin{cases} x_{i} & \text{if } i \in I \setminus A, \\ y_{i} & \text{if } i \in A. \end{cases}$$
(15)

We define sequences $s^m = \langle s_i^m : i \in I \rangle \in \prod_{i \in I} X_i$, for $m \leq l-2$, such that there are *l*-many paths of length (n+1) witnessing that $[x]_{\Phi} \sim [s^0]_{\Phi} \sim [s^1]_{\Phi} \sim \ldots \sim [s^{l-2}]_{\Phi} \sim [y']_{\Phi}$, in the following way

$$s_i^m = \begin{cases} x_i & \text{if } i \in I \setminus \bigcup_{k \le m} A_{\varepsilon^k}, \\ y_i & \text{if } i \in \bigcup_{k \le m} A_{\varepsilon^k}. \end{cases}$$
(16)

Now, x and s^0 satisfy the assumptions of Fact 2.1 over A_{ε^0} ; so, $[x]_{\Phi} \sim [s^0]_{\Phi}$. s^0 and s^1 satisfy the assumptions of Fact 2.1 over A_{ε^1} ; so, $[s^0]_{\Phi} \sim [s^1]_{\Phi}$. Continuing in this way we have that s^{l-2} and y' satisfy the assumptions of Fact 2.1 over $A_{\varepsilon^{l-1}}$; so, $[s^{l-2}]_{\Phi} \sim [y']_{\Phi}$. Thus we have $[x]_{\Phi} \sim [y']_{\Phi}$.

Let $K = \{i_j : j < k\}$ be an enumeration. For j < k the structure \mathbb{X}_{i_j} is connected so we can choose a path $\langle \bar{z}_j, \varepsilon_j \rangle$ of length n_j from x_{i_j} to y_{i_j} . By (ii) we have $F := A \cup K \in \Phi$. Let $y'' \in \prod_{i \in I} X_i$ be defined by

$$y_i'' = \begin{cases} x_i & \text{if } i \in I \setminus F, \\ y_i & \text{if } i \in F. \end{cases}$$
(17)

We define sequences $t^j = \langle t_i^j : i \in I \rangle \in \prod_{i \in I} X_i$, for $j \leq k - 2$, such that $[y']_{\Phi} \sim [t^0]_{\Phi} \sim [t^1]_{\Phi} \sim \ldots \sim [t^{k-2}]_{\Phi} \sim [y'']_{\Phi}$, by

$$t_i^j = \begin{cases} x_i & \text{if } i \in I \setminus (A \cup \{i_0, \dots, i_j\}), \\ y_i & \text{if } i \in A \cup \{i_0, \dots, i_j\}. \end{cases}$$
(18)

Namely, y' and t^0 satisfy the assumptions of Fact 2.1 over $\{i_0\}$; so, $[y']_{\Phi} \sim [t^0]_{\Phi}$. t^0 and t^1 satisfy the assumptions of Fact 2.1 over $\{i_1\}$; so, $[t^0]_{\Phi} \sim [t^1]_{\Phi}$. Continuing in this way we have that t^{k-2} and y'' satisfy the assumptions of Fact 2.1 over $\{i_{k-1}\}$; so $[t^{k-2}]_{\Phi} \sim [y'']_{\Phi}$. Thus we have proved that $[y']_{\Phi} \sim [y'']_{\Phi}$ and, since $[x]_{\Phi} \sim [y']_{\Phi}$ and $[y'']_{\Phi} = [y]_{\Phi}$, we finally have $[x]_{\Phi} \sim [y]_{\Phi}$.

Remark 2.4 Regarding Theorem 2.3, if $|I| \ge \omega$ and the filter Φ contains all cofinite sets, then condition (b) can be replaced by the following simpler condition

(b') There is $n \in \omega$ such that $\{i \in I : \mathbb{X}_i \models \varphi_n^{\operatorname{conn}}\} \in \Phi$.

In particular cases, when the filter Φ is minimal (i.e. $\Phi = \{I\}$) and when the filter Φ is maximal (an ultrafilter in P(I)) by Theorem 2.3 we have the following two corollaries.

Corollary 2.5 (Direct products and powers) If (*) holds and $\Phi = \{I\}$, then

(a) $\prod_{i \in I} X_i$ is connected iff all the structures X_i are connected and there is

 $n \in \omega$ such that \mathbb{X}_i is of diameter $\leq n$, for all except finitely many $i \in I$;

(b) The direct power $\prod_{i \in I} \mathbb{X}$ is connected iff \mathbb{X} is of finite diameter; (c) If $|I| < \omega$, then $\prod_{i \in I} \mathbb{X}_i$ is connected iff all the structures \mathbb{X}_i are connected.

Corollary 2.6 (Ultraproducts and ultrapowers) If (*) holds, $\omega \leq |I|$ is less than the first measurable cardinal and $U \subset P(I)$ is a non-principal ultrafilter, then

(a) $\prod_{\mathcal{U}} \mathbb{X}_i$ is connected, iff $\{i \in I : \mathbb{X}_i \models \varphi_n^{\text{conn}}\} \in \mathcal{U}$, for some $n \in \omega$;

(b) The ultrapower $\prod_{\mathcal{U}} \mathbb{X}$ is connected iff \mathbb{X} is of finite diameter.

The implications \leftarrow *are true in ZFC (without any assumptions on* |I|).

Theorem 2.7 Let |I| be a measurable cardinal, \mathcal{U} a non-principal ω -complete ultrafilter on I and $\mathbb{X}_i = \langle X_i, \rho_i \rangle$, $i \in I$, L-structures. Then

 $\prod_{\mathcal{U}} \mathbb{X}_i \text{ is connected} \Leftrightarrow \{i \in I : \mathbb{X}_i \text{ is connected}\} \in \mathcal{U}.$

Proof. (\Leftarrow) Let $U := \{i \in I : \mathbb{X}_i \text{ is connected}\} \in \mathcal{U}$ and let $x = \langle x_i : i \in I \rangle, y = \langle y_i : i \in I \rangle \in \prod_{i \in I} X_i$. We construct a path from $[x]_{\mathcal{U}}$ to $[y]_{\mathcal{U}}$ in $\prod_{\mathcal{U}} \mathbb{X}_i$. For $i \in U$ there is a path $\langle \overline{z}^i, \varepsilon^i \rangle$ of length $n_i + 1$ from x_i to y_i and we choose such $\varepsilon_i \in 2^{n_i+1}$. Since $|\bigcup_{n \in \omega} 2^{n+1}| = \omega$, there are $\leq \omega$ such choices; let $\{\varepsilon^k : k < \lambda\}$ be an enumeration of them; so, $\lambda \leq \omega$. Then we have $U = \bigcup_{k < \lambda} A_k$, where $A_k := \{i \in U : \varepsilon^i = \varepsilon^k\}$ and, since \mathcal{U} is ω -complete, there is $k < \lambda$ such that $A_k \in \mathcal{U}$. Now, using Fact 2.1 we construct a path from $[x]_{\mathcal{U}}$ to $[y]_{\mathcal{U}}$.

 (\Rightarrow) Let $U := \{i \in I : X_i \text{ is not connected}\} \in \mathcal{U}$. For each $i \in U$ we pick $x_i, y_i \in X_i$ such that $d(x_i, y_i) = \infty$, for $i \in I \setminus U$ we take arbitrary $x_i = y_i \in X_i$ and define $x = \langle x_i : i \in I \rangle, y = \langle y_i : i \in I \rangle \in \prod_{i \in I} X_i$. Assuming that there is a path from $[x]_{\mathcal{U}}$ to $[y]_{\mathcal{U}}$ in $\prod_{\mathcal{U}} X_i$,

$$[x]_{\mathcal{U}} \rho^{\varepsilon_0} [z^0]_{\mathcal{U}} \rho^{\varepsilon_1} [z^1]_{\mathcal{U}} \rho^{\varepsilon_2} \dots \rho^{\varepsilon_{n-1}} [z^{n-1}]_{\mathcal{U}} \rho^{\varepsilon_n} [y]_{\mathcal{U}},$$
(19)

there would be $U_1 \in \mathcal{U}$ which witnesses that and, hence,

$$\forall i \in U \cap U_1 \ x_i \ \rho_i^{\varepsilon_0} \ z_i^0 \ \rho_i^{\varepsilon_1} \ z_i^1 \ \rho_i^{\varepsilon_2} \ \dots \ \rho_i^{\varepsilon_{n-1}} \ z_i^{n-1} \ \rho_i^{\varepsilon_n} \ y_i.$$
(20)

which is impossible because $d(x_i, y_i) = \infty$, for all $i \in U$. So, there is no path from $[x]_{\mathcal{U}}$ to $[y]_{\mathcal{U}}$ in $\prod_{\mathcal{U}} X_i$ and this structure is disconnected.

A simple example of a connected *L*-structure of infinite diameter is the linear graph $\mathbb{G}_{\omega} = \langle \omega, \rho \rangle$, where $\rho = \{\langle m, n \rangle : |m - n| = 1\}$. In the absence of measurable cardinals Theorem 2.3 gives a characterization of connected reduced products of *L*-structures. In the presence of measurable cardinals we have the following equivalence.

Theorem 2.8 For each infinite set I the following conditions are equivalent (i) The cardinal |I| is less than the first measurable cardinal,

(ii) The implication (a) \Rightarrow (b) of Theorem 2.3 is true for each filter $\Phi \subset P(I)$ and each family of L-structures $\{X_i : i \in I\}$,

(iii) The ultraproduct $\prod_{\mathcal{U}} \mathbb{G}_{\omega}$ is a disconnected graph for each non-principal ultrafilter $\mathcal{U} \subset P(I)$.

Proof. The implication (i) \Rightarrow (ii) is proved in Theorem 2.3.

(ii) \Rightarrow (iii). Let (ii) hold and let $\mathcal{U} \subset P(I)$ be a non-principal ultrafilter. Assuming that the ultraproduct $\prod_{\mathcal{U}} \mathbb{G}_{\omega}$ is connected we would have a finite set $K \subset I$ and $n \in \omega$ such that $\{i \in I : \mathbb{G}_{\omega} \models \varphi_n^{\text{conn}}\} \cup K = K \in \Phi$, which is false.

(iii) \Rightarrow (i). Let (iii) hold. Suppose that $|I| \ge \kappa$, where κ is the first measurable cardinal and w.l.o.g. suppose that $\kappa \subset I$. Let $\mathcal{U}_{\kappa} \subset P(\kappa)$ be a non-principal ω -complete ultrafilter and let $\mathcal{U} \subset P(I)$ be its extension to I. Then by Fact 1.1 we have $\prod_{\mathcal{U}} \mathbb{G}_{\omega} := (\prod_{i \in I} \mathbb{G}_{\omega})/\mathcal{U} \cong (\prod_{i \in \kappa} \mathbb{G}_{\omega})/\mathcal{U}_{\kappa}$ and by Theorem 2.7 the ultraproduct $(\prod_{i \in \kappa} \mathbb{G}_{\omega})/\mathcal{U}_{\kappa}$ is connected, which is false by (iii).

The connectivity relation (and, hence, the components) of a reduced product are described by the following theorem.

Theorem 2.9 (*) If $x = \langle x_i : i \in I \rangle$, $y = \langle y_i : i \in I \rangle \in \prod_{i \in I} X_i$, then

$$[x]_{\Phi} \sim [y]_{\Phi} \quad \Leftrightarrow \quad \exists n \in \omega \ \{i \in I : \mathbb{X}_i \models \varphi_{d \le n+1}[x_i, y_i]\} \in \Phi.$$

Proof. (\Rightarrow) If $[x]_{\Phi} \sim [y]_{\Phi}$, then in $\prod_{\Phi} X_i$ there is a path from $[x]_{\Phi}$ to $[y]_{\Phi}$; namely, there are $n \in \omega, \varepsilon \in 2^{n+1}$ and $z^0, \ldots, z^{n-1} \in \prod_{i \in I} X_i$ such that we have $[x]_{\Phi} \rho^{\varepsilon_0} [z^0]_{\Phi} \rho^{\varepsilon_1} [z^1]_{\Phi} \rho^{\varepsilon_2} \ldots \rho^{\varepsilon_{n-1}} [z^{n-1}]_{\Phi} \rho^{\varepsilon_n} [y]_{\Phi}$. Since Φ is closed under finite intersections there is $F \in \Phi$ such that for each $i \in F$ we have $x_i \rho_i^{\varepsilon_0} z_i^0 \rho_i^{\varepsilon_1} z_i^{1} \rho_i^{\varepsilon_2} \ldots \rho^{\varepsilon_{n-1}} z_i^{n-1} \rho_i^{\varepsilon_n} y_i$. So $F \subset \{i \in I : X_i \models \varphi_{d \leq n+1}[x_i, y_i]\} \in \Phi$.

 $(\Leftarrow) \text{ Let } n \in \omega \text{ and } A := \{i \in I : \mathbb{X}_i \models \varphi_{d \leq n+1}[x_i, y_i]\} \in \Phi. \text{ If } i \in A, \text{ then there is a path } \langle \bar{z}, \varepsilon \rangle \text{ of length } n+1 \text{ from } x_i \text{ to } y_i; \text{ thus we can choose one such } \varepsilon. \text{ Let } \varepsilon^k \in 2^{n+1}, k < l, \text{ be an enumeration of the } n+1\text{-tuples chosen in this way for } i \in A. \text{ Then defining } A_{\varepsilon^k} = \{i \in A : \varepsilon^k \text{ is chosen for } x_i \text{ and } y_i\}, \text{ for } k < l, \text{ we have } A = \bigcup_{k < l} A_{\varepsilon^k}. \text{ Let } y' \in \prod_{i \in I} X_i, \text{ where } y' \upharpoonright A = y \upharpoonright A \text{ and } y' \upharpoonright (I \setminus A) = x \upharpoonright (I \setminus A). \text{ Then } [y']_{\Phi} = [y]_{\Phi}. \text{ We define sequences } s^m = \langle s_i^m : i \in I \rangle \in \prod_{i \in I} X_i, \text{ for } m \leq l-2, \text{ such that there are } l\text{-many paths of length } (n+1) \text{ witnessing that } [x]_{\Phi} \sim [s^0]_{\Phi} \sim [s^1]_{\Phi} \sim \ldots \sim [s^{l-2}]_{\Phi} \sim [y']_{\Phi}, \text{ by } t \leq l \leq l \}$

$$s_i^m = \begin{cases} x_i & \text{if } i \in I \setminus \bigcup_{k \le m} A_{\varepsilon^k}, \\ y_i & \text{if } i \in \bigcup_{k < m} A_{\varepsilon^k}. \end{cases}$$
(21)

Now, x and s^0 satisfy the assumptions of Fact 2.1 over A_{ε^0} ; so, $[x]_{\Phi} \sim [s^0]_{\Phi}$. s^0 and s^1 satisfy the assumptions of Fact 2.1 over A_{ε^1} ; so, $[s^0]_{\Phi} \sim [s^1]_{\Phi}$. Continuing

in this way we have that s^{l-2} and y' satisfy the assumptions of Fact 2.1 over $A_{\varepsilon^{l-1}}$; so, $[s^{l-2}]_{\Phi} \sim [y']_{\Phi}$. Thus we have $[x]_{\Phi} \sim [y']_{\Phi} = [y]_{\Phi}$. \Box

Acknowledgement This research was supported by the Science Fund of the Republic of Serbia, Program PROMIS, Grant No. 6062228, *Classification of Large Objects–Ultrafilters and Directed Sets–*CLOUDS.

References

[1] C. C. Chang, H. J. Keisler, Model Theory, North Holland, Amsterdam, 1973.