

CROSS-CONNECTIONS IN CLIFFORD SEMIGROUPS

P.A. AZEEF MUHAMMED AND C.S. PREENU

ABSTRACT. An inverse Clifford semigroup (often referred to as just a Clifford semigroup) is a semilattice of groups. It is an inverse semigroup and in fact, one of the earliest studied classes of semigroups [6]. In this short note, we discuss various structural aspects of a Clifford semigroup from a cross-connection perspective. In particular, given a Clifford semigroup S , we show that the semigroup $T\mathbb{L}(S)$ of normal cones is isomorphic to the original semigroup S , even when S is not a monoid. Hence, we see that cross-connection description degenerates in Clifford semigroups. Further, we specialise the discussion to provide the description of the cross-connection structure in an arbitrary semilattice, also.

1. INTRODUCTION

Grillet introduced cross-connections as a pair of functions to describe the inter-relationship between the posets of principal left and right ideals of a regular semigroup. This construction involved building two intermediary semigroups and further identifying a fundamental image of the semigroup as a subdirect product, using the cross-connection functions. But isomorphic posets give rise to isomorphic cross-connections. So, Nambooripad replaced posets with certain small categories to overcome this limitation. Hence, using the categorical theory of cross-connections, Nambooripad constructed arbitrary semigroups from their ideal structure.

Starting from a regular semigroup S , Nambooripad identified two small categories: $\mathbb{L}(S)$ and $\mathbb{R}(S)$ which abstract the principal left and right ideal structures of the semigroup, respectively. He showed that these categories are interconnected using a pair of functors. It can be seen that the object maps of these functors coincides with Grillet's cross-connection functions and so Nambooripad called his functors also cross-connections. Further, Nambooripad showed that this correspondence can be extended to an explicit category equivalence between the category of regular semigroups and the category of cross-connections. Hence the ideal structure of a regular semigroup can be completely captured using these 'cross-connected' categories.

Being a rather technical construction, it is instructive to work out the simplifications that arise in various special classes of semigroups. There has been several works in this direction [1–5, 10] and in this short article, we propose to outline how the construction simplifies in a couple of very natural classes of regular semigroups: namely Clifford semigroups and semilattices. As the reader may see, this exercise also provides some useful illustrations to several cross-connection related subtleties.

In fact, Clifford semigroups are one of the first classes of inverse semigroups whose structure was studied. It was originally defined [6] as a union of groups in which idempotents commute. It may be noted that some authors refer to general union of groups also as Clifford semigroups but we shall follow [8] and refer to a

semilattice of groups as a Clifford semigroup. The following characterizations of Clifford semigroups will be useful in the sequel.

Theorem 1.1. [8, Theorem 4.2.1] [7, Theorem 1.3.11] Let S be a semigroup. Then the following statements are equivalent.

- (1) S is a Clifford semigroup;
- (2) S is a semilattice of groups;
- (3) S is a strong semilattice of groups;
- (4) S is regular and the idempotents of S are central;
- (5) Every \mathcal{D} class of S has a unique idempotent.

In Nambooripad's cross-connection description, starting from a regular semigroup S , two small categories of principal left and right ideals (denoted by $\mathbb{L}(S)$ and $\mathbb{R}(S)$, respectively in the sequel) are defined. Then their inter-relationship is abstracted as a pair of functors called cross-connections. Conversely, given an abstractly defined pair of cross-connected categories (with some special properties), one can construct a regular semigroup. This correspondence between regular semigroups and cross-connections is proved to be a category equivalence.

The construction of the regular semigroup from the category happens in several layers and the crucial object here is the intermediary regular semigroup $T\mathbb{L}(S)$ of normal cones from the given category $\mathbb{L}(S)$. We shall see that when the semigroup S is Clifford, then the semigroup $T\mathbb{L}(S)$ is isomorphic to S . It is known that this is not true in general [5] even for an inverse semigroup S . This in turn, highly degenerates the cross-connection structure in Clifford semigroups. This is discussed in the next section. In the last section, we specialise our discussion to an arbitrary semilattice and describe the cross-connections therein.

As mentioned above, since the article can be seen as a part of a continuing project of studying the various classes of regular semigroups within the cross-connection framework, we refer the reader to [1, 5, 10] for the preliminary notions and formal definitions in Nambooripad's cross-connection theory. We also refer the reader to [9] for the original treatise on cross-connections.

2. NORMAL CATEGORIES AND CROSS-CONNECTIONS IN CLIFFORD SEMIGROUPS

Recall from [9] that given a regular semigroup S , the normal category $\mathbb{L}(S)$ of principal left ideals are defined as follows. The object set

$$v\mathbb{L}(S) := \{Se : e \in E(S)\}$$

and the morphisms in $\mathbb{L}(S)$ are partial right translations. In fact, the set of all morphisms between two objects Se and Sf may be characterised as the set $\{\rho(e, u, f) : u \in eSf\}$, where the map $\rho(e, u, f)$ sends $x \in Se$ to $xu \in Sf$.

First, we proceed to discuss some special properties of the category $\mathbb{L}(S)$ when S is a Clifford semigroup. This will lead us to the characterisation of the semigroup $T\mathbb{L}(S)$ of all normal cones in S .

Proposition 2.1. Let S be a Clifford semigroup. Two objects in $\mathbb{L}(S)$ are isomorphic if and only if they are identical.

Proof. Clearly, identical objects are always isomorphic. Conversely, suppose Se and Sf are two isomorphic objects in $\mathbb{L}(S)$. Then by [9, Proposition III.13(c)], we have $e \mathcal{D} f$. Recall that in a Clifford semigroup, the Green's relations \mathcal{L} , \mathcal{R} and \mathcal{D} are identical. Therefore $e\mathcal{L}f$ and hence $Se = Sf$. \square

Given the normal category $\mathbb{L}(S)$ of principal left ideals of a regular semigroup S , it is known that two morphisms are equal, i.e. $\rho(e, u, f) = \rho(g, v, h)$ if and only if $e \mathcal{L} g$, $f \mathcal{L} h$, and $v = gu$. Also, given two morphisms $\rho(e, u, f)$ and $\rho(g, v, h)$, they are composable if and only $Sf = Sg$ so that $\rho(e, u, f)\rho(g, v, h) = \rho(e, uv, h)$. Now, we see that the equality of morphisms simplify when S is a Clifford semigroup.

Proposition 2.2. Let S be a Clifford semigroup, then $\rho(e, u, f) = \rho(g, v, h)$ in the category $\mathbb{L}(S)$ if and only if $e = g$, $u = v$ and $f = h$.

Proof. Suppose that two morphisms $\rho(e, u, f) = \rho(g, v, h)$ are equal. Recall that by [9, Lemma II.12], this implies that $Se = Sg$, $Sf = Sh$ and $v = gu$. That is, the elements e and g are two \mathcal{L} related idempotents. But since \mathcal{L} and \mathcal{H} are identical in a Clifford semigroup S and since a \mathcal{H} -class can contain at most one idempotent, we have $e = g$. Similarly we get $f = h$. So, the sets $eSf = gSh$ are equal and also $v = gu = eu = u$. \square

Now, we proceed to characterise the building blocks of the cross-connection construction, namely the normal cones in the category $\mathbb{L}(S)$. An ‘order-respecting’ collection of morphisms in a normal category is defined as a normal cone.

Definition 2.1. Let $\mathbb{L}(S)$ be the normal category of principal left ideals in a regular semigroup S and $Sd \in v\mathbb{L}(S)$. A *normal cone with apex Sd* is a function $\gamma: v\mathbb{L}(S) \rightarrow \mathbb{L}(S)$ such that:

- (1) for each $Se \in v\mathbb{L}(S)$, one has $\gamma(Se) \in \mathbb{L}(S)(Se, Sd)$;
- (2) $\iota(Sf, Sg)\gamma(Sg) = \gamma(Sf)$ whenever $Sf \subseteq Sg$;
- (3) $\gamma(Sm)$ is an isomorphism for some $Sm \in v\mathbb{L}(S)$.

Now, for each $a \in S$, we can define a function $\rho^a: v\mathbb{L}(S) \rightarrow \mathbb{L}(S)$ as follows:

$$(1) \quad \rho^a(Se) := \rho(e, ea, f) \text{ where } f \mathcal{L} a.$$

It is easy to verify that the map ρ^a is a well-defined normal cone with apex Sf in the sense of Definition 2.1, see [9, Lemma III.15].

In the sequel, the normal cone ρ^a is called the *principal cone* determined by the element a . In particular, observe that, for an idempotent $e \in E(S)$, we have a principal cone ρ^e such that $\rho^e(Se) = \rho(e, e, e) = 1_{Se}$. This leads us to the most crucial proposition of this article.

Proposition 2.3. In a Clifford semigroup S , every normal cone in $\mathbb{L}(S)$ is a principal cone.

Proof. Suppose γ is a normal cone in $\mathbb{L}(S)$ with vertex Se so that $\gamma(Se) = \rho(e, u, e)$ for some $u \in S$. Then for any $Sf \in v\mathbb{L}(S)$, we shall show that $\gamma(Sf) = \rho(f, fu, e) = \rho^u$ for some $u \in S$. To this end, first observe that since idempotents commute in S , we have $Se f = S f e \subseteq Se$. So, by (2) of Definition 2.1, we have $\gamma(Se f) = \rho(e f, e f, e)\gamma(Se)$. Then,

$$\begin{aligned} \gamma(Se f) &= \rho(e f, e f, e)\gamma(Se) && \text{since } Se f \subseteq Se \\ &= \rho(e f, e f, e)\rho(e, u, e) \\ &= \rho(e f, e f u, e) \\ &= \rho(e f, f u, e) && \text{since } u \in e S e \text{ and } e f = f e. \end{aligned}$$

Now for any $Sf \in v\mathbb{L}(S)$, let $\gamma(Sf) = \rho(f, v, e)$ for some $v \in fSe$. Then since $Se f \subseteq Sf$ also, we have

$$\begin{aligned}
\gamma(Se f) &= \rho(ef, ef, f)\gamma(Sf) \\
&= \rho(ef, ef, f)\rho(f, v, e) \\
&= \rho(ef, efv, e) \\
&= \rho(ef, ev, e) && \text{since } fv = v \\
&= \rho(ef, ve, e) && \text{since idempotents are central in } S \\
&= \rho(ef, v, e) && \text{since } v \in fSe.
\end{aligned}$$

Now, from the discussion above, we see that $\rho(ef, fu, e) = \rho(ef, v, e)$. Then, using Proposition 2.2, this implies that $v = fu$. Hence for all $Sf \in v\mathbb{L}(S)$, we have $\gamma(Sf) = \rho(f, fu, e)$ and so $\gamma = \rho^u$. \square

In general, for an arbitrary regular semigroup S , two distinct principal cones ρ^a and ρ^b may be equal in $\mathbb{L}(S)$ even when $a \neq b$. But when S is Clifford, we proceed to show that it is not the case.

Proposition 2.4. Let S be a Clifford semigroup. Given two principal cones ρ^a and ρ^b , we have $\rho^a = \rho^b$ if and only if $a = b$.

Proof. Clearly when $a = b$, then $\rho^a = \rho^b$. Conversely suppose $\rho^a = \rho^b$. Then their vertices coincide and so, we have $Sa = Sb$, then since S is Clifford and Green's relations coincide, we have $a\mathcal{H}b$. Now let e be the idempotent in $H_a = H_b$, the Green's \mathcal{H} class containing a and b . Then $\rho^a(Se) = \rho(e, ea, e) = \rho(e, a, e)$. Similarly we get $\rho^b(Se) = \rho(e, b, e)$. Since the cones are equal, the corresponding morphism components at each vertex coincide. Hence using Proposition 2.2, we have $a = b$. \square

Recall from [9, Section III.1] that the set of all normal cones in a normal category forms a regular semigroup, under a natural binary operation. So, in particular, given the normal category $\mathbb{L}(S)$, the set $T\mathbb{L}(S)$ of all normal cones in the category $\mathbb{L}(S)$ is a regular semigroup. Now, we proceed to characterise this semigroup when S is Clifford.

Theorem 2.5. Let S be a Clifford semigroup. Then the semigroup $T\mathbb{L}(S)$ of all normal cones in $\mathbb{L}(S)$ is isomorphic to the semigroup S .

Proof. Recall from [9, Section III.3.2] that the map $\bar{\rho}: a \mapsto \rho^a$ from a regular semigroup S to the semigroup $T\mathbb{L}(S)$ is a homomorphism. Now, when S is Clifford, by 2.3, we have seen that every normal cone in $T\mathbb{L}(S)$ is principal and hence the map $\bar{\rho}$ is surjective. Also, by Proposition 2.4, we see that the map $\bar{\rho}$ is injective. Hence the map $\bar{\rho}$ is an isomorphism from the Clifford semigroup S to the semigroup $T\mathbb{L}(S)$. \square

The above theorem characterises the semigroup of $T\mathbb{L}(S)$ of all normal cones in $\mathbb{L}(S)$; this naturally leads us to the complete description of the cross-connection structure of the semigroup S as follows.

Recall from [9, Section III.4] that given a normal category, it has an associated dual category whose objects are certain set-valued functors and morphisms are natural transformations. Now we proceed to characterise the normal dual $N^*\mathbb{L}(S)$ of the normal category $\mathbb{L}(S)$ of principal left ideals of a Clifford semigroup S .

Theorem 2.6. Let S be a Clifford semigroup. Then the normal dual $N^*\mathbb{L}(S)$ of the normal category $\mathbb{L}(S)$ of principal left ideals in S is isomorphic to the normal category $\mathbb{R}(S)$ of principal right ideals in S .

Proof. It is known that [9, Theorem III.25] the normal dual $N^*\mathbb{L}(S)$ of the normal category $\mathbb{L}(S)$ is isomorphic to the normal category $\mathbb{R}(T\mathbb{L}(S))$ of principal right ideals of the regular semigroup $T\mathbb{L}(S)$. When S is a Clifford semigroup, by Theorem 2.5, we see that the semigroup $T\mathbb{L}(S)$ is isomorphic to S and hence $N^*\mathbb{L}(S)$ is isomorphic to $\mathbb{R}(S)$ as normal categories. \square

Dually, we can easily prove the following results:

Theorem 2.7. Let S be a Clifford semigroup. The semigroup $T\mathbb{R}(S)$ of all normal cones in the category $\mathbb{R}(S)$ of all principal right ideals in S is anti-isomorphic to the semigroup S . The normal dual $N^*\mathbb{R}(S)$ of the normal category $\mathbb{R}(S)$ is isomorphic to the normal category $\mathbb{L}(S)$.

Recall from [9, Theorem IV.1] that the cross-connection of a regular semigroup S is defined as a quadruplet $(\mathbb{L}(S), \mathbb{R}(S), \Gamma, \Delta)$ such that $\Gamma: \mathbb{R}(S) \rightarrow N^*\mathbb{L}(S)$ and $\Delta: \mathbb{L}(S) \rightarrow N^*\mathbb{R}(S)$ are functors satisfying certain properties. Now, using Theorems 2.6 and 2.7, we see that both the functors Γ and Δ are in fact isomorphisms. And hence the cross-connection structure degenerates to isomorphisms of the associated normal categories in a Clifford semigroup S .

3. CROSS-CONNECTIONS OF A SEMILATTICE

Clearly, a semilattice is a Clifford semigroup. So, we specialise our discussion to a semilattice using the results in the previous section. The following theorem follows from Theorem 2.5.

Theorem 3.1. Let S be a semilattice. Then the semigroup $T\mathbb{L}(S)$ of all normal cones in $\mathbb{L}(S)$ is isomorphic to the semilattice S .

Theorems 2.6 and 2.7 when applied to a semilattice can be unified as follows:

Theorem 3.2. Let S be a semilattice. Then the normal dual $N^*\mathbb{L}(S)$ of the normal category $\mathbb{L}(S)$ of principal left ideals in S is isomorphic to the normal category $\mathbb{R}(S)$ of principal right ideals in S . The normal dual $N^*\mathbb{R}(S)$ of the normal category $\mathbb{R}(S)$ is isomorphic to the normal category $\mathbb{L}(S)$.

Hence, we see that both the cross-connections functors Γ and Δ are isomorphisms, when we have a semilattice also. So, the cross-connection structure degenerates to isomorphisms of the associated normal categories in a semilattice, too.

REFERENCES

- [1] P. A. Azeef Muhammed. Cross-connections and variants of the full transformation semigroup. *Acta Sci. Math. (Szeged)*, (84):377–399, 2018.
- [2] P. A. Azeef Muhammed. Cross-connections of linear transformation semigroups. *Semigroup Forum*, 97(3):457–470, 2018.
- [3] P. A. Azeef Muhammed and A. R. Rajan. Cross-connections of completely simple semigroups. *Asian-European J. Math.*, 09(03):1650053, 2016.
- [4] P. A. Azeef Muhammed and A. R. Rajan. Cross-connections of the singular transformation semigroup. *J. Algebra Appl.*, 17(3):1850047, 2018.
- [5] P. A. Azeef Muhammed, M. V. Volkov, and K. Auinger. Cross-connection structure of locally inverse semigroups. 2022. arXiv:1912.00214 (submitted).

- [6] A. H. Clifford. Semigroups admitting relative inverses. *Annals of Mathematics*, 42(4):1037–1049, 1941.
- [7] P. M. Higgins. *Techniques of semigroup theory*. Clarendon Press, 1992.
- [8] J. M. Howie. *Fundamentals of semigroup theory*. Number 12. Oxford University Press, 1995.
- [9] K. S. S. Nambooripad. *Theory of Cross-connections*. Centre for Mathematical Sciences, 1994.
- [10] A. R. Rajan, C. S. Preenu, and K. S. Zeenath. Cross connections for normal bands and partial order on morphisms. *Semigroup Forum*, (104):125–147, 2022.

CENTRE FOR RESEARCH IN MATHEMATICS AND DATA SCIENCE, WESTERN SYDNEY UNIVERSITY,
PENRITH, NSW 2751, AUSTRALIA.

DEPARTMENT OF MATHEMATICS, UNIVERSITY COLLEGE, THIRUVANANTHAPURAM, KERALA 695034,
INDIA.