

A point to set principle for finite-state dimension

Elvira Mayordomo*

August 2, 2022

Abstract

Effective dimension has proven very useful in geometric measure theory through the point-to-set principle [7] that characterizes Hausdorff dimension by relativized effective dimension. Finite-state dimension is the least demanding effectivization in this context [2] that among other results can be used to characterize Borel normality [1].

In this paper we prove a characterization of finite-state dimension in terms of information content of a real number at a certain precision. We then use this characterization to prove a finite-state dimension point-to-set principle.

1 Introduction

Effective dimension was introduced in [6, 5] as an effectivization of Hausdorff dimension. One of its generalizations is finite-state dimension [2] that is a robust notion that interacts with compression and characterizes Borel normality [1].

In [7] Lutz and Lutz proved a point-to-set principle that characterizes Hausdorff dimension in terms of relativized effective dimension. This principle has already produced a number of interesting results in geometric fractal theory through computability based proofs (for an already outdated list see [4, 8] and their references).

In this paper we provide a characterization of finite-state dimension on the Euclidean space based on the finite-state information content of a real number at a certain precision, which also provides an alternative characterization of Borel normality. Then we use this dimension characterization to prove a finite-state dimension point-to-set principle.

*Departamento de Informática e Ingeniería de Sistemas, Instituto de Investigación en Ingeniería de Aragón, Universidad de Zaragoza, Spain. Research supported in part by Spanish Ministry of Science and Innovation grant PID2019-104358RB-I00 and by the Science dept. of Aragon Government: Group Reference T64_20R (COSMOS research group).

2 Preliminaries

Let Σ be a finite alphabet. We write Σ^* for the set of all (finite) *strings* over Σ and Σ^∞ for the set of all (infinite) *sequences* over Σ . We write $|x|$ for the length of a string or sequence x , and we write λ for the *empty string*, the string of length 0. For $x \in \Sigma^* \cup \Sigma^\infty$ and $0 \leq n < |x|$, we write $x \upharpoonright n = x[0..n-1]$. For $w \in \Sigma^*$ and $x \in \Sigma^* \cup \Sigma^\infty$, we say that w is a *prefix* of x , and we write $w \sqsubseteq x$, if $x \upharpoonright |w| = w$.

A Σ *finite-state transducer* (Σ -FST) is a 4-tuple $T = (Q, \delta, \nu, q_0)$, where

- Q is a nonempty, finite set of *states*,
- $\delta : Q \times \Sigma \rightarrow Q$ is the *transition function*,
- $\nu : Q \times \Sigma \rightarrow \Sigma^*$ is the *output function*, and
- $q_0 \in Q$ is the *initial state*.

For $q \in Q$ and $w \in \Sigma^*$, we define the *output* from state q on input w to be the string $\nu(q, w)$ defined by the recursion

$$\begin{aligned}\nu(q, \lambda) &= \lambda, \\ \nu(q, wa) &= \nu(q, w)\nu(\delta(q, w), a)\end{aligned}$$

for all $w \in \Sigma^*$ and $a \in \Sigma$. We then define the *output* of T on input $w \in \Sigma^*$ to be the string $T(w) = \nu(q_0, w)$.

For each integer $b \geq 1$ we let $\Sigma_b = \{0, 1, \dots, b-1\}$ be the alphabet of *base- b digits*. We use infinite sequences over Σ_b to represent real numbers in $[0, 1)$.

Each $S \in \Sigma_b^\infty$ is associated the real number $\text{real}_b(S) = \sum_{i=1}^{\infty} S[i-1]b^{-i}$ and for each $x \in [0, 1)$, $\text{seq}_b(x)$ is the infinite sequence S that does not finish with infinitely many $b-1$ and such that $x = \text{real}_b(S)$.

A set of real numbers $A \subseteq [0, 1)$ is represented by the set

$$\text{seq}_b(A) = \{\text{seq}_b(\alpha) \mid \alpha \in A\}$$

of sequences. If $X \subseteq \Sigma_b^\infty$ then

$$\text{real}_b(X) = \{\text{real}_b(x) \mid x \in X\}.$$

We will denote with \mathcal{D}_b the set of rational numbers that have finite representation in base b , that is,

$$\mathcal{D}_b = \{q \mid \text{seq}_b(q) = w0^\infty, w \in \Sigma_b^*\}.$$

We will write $\text{real}_b(w)$ for $\text{real}_b(w0^\infty)$ when $w \in \Sigma_b^*$.

3 Finite state dimension

Finite-state dimension was introduced in [2] in terms of gambling and characterized using information lossless compression.

Here we present an alternative definition on the Euclidean space and then prove its equivalence with [2].

Definition. Let T be a Σ -FST and let $w \in \Sigma^*$. The T -information content of w is

$$K^T(w) = \min \{ |\pi| \mid T(\pi) = w \}.$$

Definition. Let T be a Σ_b -FST, $\delta > 0$ and $x \in [0, 1)$. The base- b T -information content of x at precision δ is

$$K_\delta^T(x) = \min \{ K^T(w) \mid |\text{real}_b(w) - x| < \delta \}.$$

We next define the finite-state dimension of points and sets.

Definition. Let $b \geq 1$. Let $x \in [0, 1)$ and $A \subseteq [0, 1)$. The base- b finite-state dimension of x is

$$\dim_{\text{FS}}^b(x) = \inf_{T \Sigma_b\text{-FST}} \liminf_{\delta > 0} \frac{K_\delta^T(x)}{\log_b(1/\delta)},$$

the base- b finite-state dimension of A is

$$\dim_{\text{FS}}^b(A) = \inf_{T \Sigma_b\text{-FST}} \sup_{x \in A} \liminf_{\delta > 0} \frac{K_\delta^T(x)}{\log_b(1/\delta)}.$$

Observation 3.1 $\dim_{\text{FS}}^b(x) = \inf_{T \Sigma_b\text{-FST}} \liminf_n \frac{K_{b^{-n}}^T(x)}{n}.$

The definition of finite-state dimension from [2] is usually done in a space of infinite sequences, while identifying $[0, 1)$ and Σ_b^∞ through seq_b or base- b representation.

Doty and Moser [3] proved that finite dimension on sequences can be characterized in terms of finite-state transducers.

Theorem 3.2 ([3]) Let $S \in \Sigma^\infty$,

$$\dim_{\text{FS}}(S) = \inf_{T \Sigma\text{-FST}} \liminf_n \frac{K^T(S \upharpoonright n)}{n}.$$

We next show that the notion of information content at a certain precision characterizes finite-state dimension.

Theorem 3.3 For each $b \geq 1$, $x \in [0, 1)$, and $A \subseteq [0, 1)$

$$\dim_{\text{FS}}^b(x) = \dim_{\text{FS}}(\text{seq}_b(x)),$$

$$\dim_{\text{FS}}^b(A) = \dim_{\text{FS}}(\text{seq}_b(A)).$$

Proof. Let $x \in [0, 1)$, let $S = \text{seq}_b(x)$. Then for every $n \in \mathbb{N}$ and $T \Sigma_b$ -FST, $K_{b^{-n}}^T(x) \leq K^T(S \upharpoonright (n+1))$ and therefore $\dim_{\text{FS}}^b(x) \leq \dim_{\text{FS}}(S)$.

For each $w \in \Sigma_b^* \cup \Sigma_b^\infty$, let $\text{comp}(w)$ be the complementary of w , that is, $\text{comp}(w)[i] = b - 1 - w[i]$ for $0 \leq i < |w|$.

Claim 3.4 $\dim_{\text{FS}}(S) = \dim_{\text{FS}}(\text{comp}(S))$. $\dim_{\text{FS}}^b(x) = \dim_{\text{FS}}^b(\text{real}_b(\text{comp}(\text{seq}_b(x))))$.

Claim 3.5 $\dim_{\text{FS}}(S) \leq \dim_{\text{FS}}^b(x)$.

To prove this claim, notice that $\dim_{\text{FS}}^b(x)$ needs to be witnessed either by approximations from above or for approximations from below and that tighter approximations can be delayed.

That is, for every FST T there exist and infinitely many n_i such that

$$\lim_i \frac{K_{b^{-n_i}}^T(x)}{n_i} = \liminf_n \frac{K_{b^{-n}}^T(x)}{n}.$$

For each i let w_i be such that $K^T(w_i) = K_{b^{-n_i}}^T(x)$ and $|x - \text{real}_b(w_i)| < b^{-n_i}$. Let m_i be such that $b^{-m_i-1} \leq |x - \text{real}_b(w_i)| < b^{-m_i} \leq b^{-n_i}$. Then $\frac{K_{b^{-m_i}}^T(x)}{m_i} \leq \frac{K_{b^{-n_i}}^T(x)}{n_i}$, and for $T'(\pi) = \text{comp}(T(\pi))$, either

$$\liminf_n \frac{K^T(S \upharpoonright n)}{n} \leq \lim_i \frac{K_{b^{-m_i}}^T(x)}{m_i}$$

or

$$\liminf_n \frac{K^{T'}(\text{comp}(S) \upharpoonright n)}{n} \leq \lim_i \frac{K_{b^{-m_i}}^T(x)}{m_i}.$$

Therefore either $\dim_{\text{FS}}(S) \leq \dim_{\text{FS}}^b(x)$ or $\dim_{\text{FS}}(\text{comp}(S)) \leq \dim_{\text{FS}}^b(x)$ and the claim follows. \square

Corollary 3.6 Let $b \geq 1$, $x \in [0, 1)$. x is b -normal if and only if $\dim_{\text{FS}}^b(x) = 1$, that is,

$$\inf_{T \Sigma_b\text{-FST}} \liminf_{\delta > 0} \frac{K_\delta^T(x)}{\log_b(1/\delta)} = 1.$$

4 Point to set principle for finite-state dimension

We denote as *separator* a set $S \subseteq [0, 1)$ such that S is countable and dense in $[0, 1]$.

Definition. A *separator enumerator (SE)* is a function $f : \Sigma^* \rightarrow [0, 1)$ such that $\text{Im}(f)$ is a separator.

Given a separator enumerator we can define information content in $[0, 1)$.

Definition. Let $f : \Sigma^* \rightarrow [0, 1)$ be a SE. Let T be a Σ -FST, $\delta > 0$ and $x \in [0, 1)$. The *f-T-information content of x at precision δ* is

$$K_\delta^{T,f}(x) = \min \{K^T(w) \mid |f(w) - x| < \delta\}.$$

Definition. Let $f : \Sigma^* \rightarrow [0, 1)$ be a SE. Let $x \in [0, 1)$ and $A \subseteq [0, 1)$. The *f-enumerator finite-state dimension of x* is

$$\dim_{\text{FS}}^f(x) = \inf_{T \in \Sigma\text{-FST}} \liminf_{\delta > 0} \frac{K_\delta^{T,f}(x)}{\log_{|\Sigma|}(1/\delta)},$$

the *f-enumerator finite-state dimension of A* is

$$\dim_{\text{FS}}^f(A) = \inf_{T \in \Sigma\text{-FST}} \sup_{x \in A} \liminf_{\delta > 0} \frac{K_\delta^{T,f}(x)}{\log_{|\Sigma|}(1/\delta)}.$$

Theorem 4.1 Let $A \subseteq [0, 1)$.

$$\dim_{\text{H}}(A) = \min_{f \in \text{SE}} \dim_{\text{FS}}^f(A),$$

$$\dim_{\text{H}}(A) = \min_{f : \{0,1\}^* \rightarrow \mathcal{D}_2} \dim_{\text{FS}}^f(A).$$

Proof. Let C be such that $\dim_{\text{H}}(A) = \dim^C(A)$. Let $h : \{0, 1\}^* \rightarrow \{0, 1\}^*$ be such that $h(w) = U^C(w)$ when $U^C(w)$ is defined, and $U^C(w) = 0$ otherwise. Then $f(w) = \text{real}_2(h(w))$ is the required SE. \square

Notice that the previous theorem holds even when fixing a particular countable dense set. In terms of Borel normality, it shows that reordering the set \mathcal{D}_b of base- b finite representation numbers is enough to obtain normality for any other base.

Definition. Let $f : \Sigma^* \rightarrow [0, 1)$ be a SE, let $x \in [0, 1)$. x is *f-normal* if $\dim_{\text{FS}}^f(x)$.

5 Conclusions

We expect that our main theorem will be useful to produce lower bounds on Hausdorff dimension in different settings. Notice that the result can be directly translated to any separable metric space and any reasonable gauge family.

Our result helps clarify the oracle role in the point to set principles. The next step would be to classify the different enumerations of a countable dense set.

We believe the notion of f -normal sequence can be of independent interest.

References

- [1] C. Bourke, J. M. Hitchcock, and N. V. Vinodchandran. Entropy rates and finite-state dimension. *Theoretical Computer Science*, 349(3):392–406, 2005.
- [2] J. J. Dai, J. I. Lathrop, J. H. Lutz, and E. Mayordomo. Finite-state dimension. *Theoretical Computer Science*, 310:1–33, 2004.
- [3] D. Doty and P. Moser. Finite-state dimension and lossy decompressors. Technical Report CoRR abs/cs/0609096, Arxiv, 2006.
- [4] J. Lutz and N. Lutz. Who asked us? how the theory of computing answers questions about analysis. In *Complexity and Approximation: In Memory of Ker-I Ko*. Springer, ding-zhu du and jie wang (eds.) edition, 2020.
- [5] J. H. Lutz. Dimension in complexity classes. *SIAM Journal on Computing*, 32(5):1236–1259, 2003.
- [6] J. H. Lutz. The dimensions of individual strings and sequences. *Information and Computation*, 187(1):49–79, 2003.
- [7] J. H. Lutz and N. Lutz. Algorithmic information, plane Kakeya sets, and conditional dimension. *ACM Transactions on Computation Theory*, 10, 2018. Article 7.
- [8] J. H. Lutz and E. Mayordomo. Algorithmic fractal dimensions in geometric measure theory. In V. Brattka and P. Hertling, editors, *Handbook of Computability and Complexity in Analysis*. Springer-Verlag, 2021.