ON (STRONGLY) (Θ -)DISCRETE HOMOGENEOUS SPACES

VITALIJ A. CHATYRKO AND ALEXANDRE KARASSEV

ABSTRACT. We introduce the classes of (strongly) (Θ -)discrete homogeneous spaces. We discuss the relationships of these classes to other classes of spaces possessing homogeneity-related properties, such as (strongly) (*n*-)homogeneous spaces. Many examples are given distinguishing discrete homogeneity and other types of homogeneity.

1. INTRODUCTION

A family of subsets \mathcal{D} of a topological space X is *discrete* if each point $x \in X$ has a neighborhood Ox which intersects at most one set from \mathcal{D} . If each element of \mathcal{D} is a singleton, we obtain the definition of a discrete subset of X. Note that if X is T_1 then each discrete subset of X is closed.

Recall also [BBHS] that a subset D of a topological space X is Θ discrete if each point $x \in X$ has a neighbourhood $Ox \subset X$ whose closure ClOx contains at most one point of the set D.

It is clear that each Θ -discrete subset of a topological space is discrete. There exist Hausdorff spaces in which some discrete subsets are not Θ -discrete ([BBHS, Example 1]). However, in regular Hausdorff spaces Θ -discrete subsets and discrete subsets are the same.

In [BBHS] the authors proved the following interesting result.

Theorem 1.1. ([BBHS, Theorem 1 and Corollary 2]) Let X be the connected countable Hausdorff space B due to Bing ([Bi], see also the appendix for the definition) or the connected, locally connected, countable Hausdorff space R due to Ritter ([R], see Section 3 for the definition). Then any bijection $f: A \to B$ between Θ -discrete subsets A and B of X extends to a homeomorphism $f: X \to X$.

²⁰²⁰ Mathematics Subject Classification. Primary 57S05; Secondary 54F99, 54A10.

Key words and phrases. (strong) discrete homogeneity, (strong) n-homogeneity, Bing space, Ritter space.

This result and the definition of (strongly) n-homogeneous spaces (see further in the text) motivates the following definition (for a set S by |S| we denote its cardinality).

Definition 1.2. A Hausdorff topological space X is called

- discrete homogeneous (DH), if for any two discrete subsets A and B of X with |A| = |B| there exists a homeomorphism f of X onto itself such that f(A) = B;
- strongly discrete homogeneous (sDH), if for any two discrete subsets A and B of X and any bijection f: A → B this bijection extends to a homeomorphism of the whole X onto itself.

Replacing in the above definition discrete subsets with Θ -discrete, we obtain the definitions of Θ -discrete homogeneous (Θ -DH), and strongly Θ -discrete homogeneous (s Θ -DH) spaces, respectively.

Note that the Bing space B and the Ritter space R are examples of s Θ -DH spaces. It is also easy to see that each s(Θ -)DH space is (Θ -)DH, and each (s)DH space is (s) Θ -DH. Note, however, that the Bing space B is not a DH space ([BBHS, Example 2]). It is natural to ask the following question.

Question 1.3. Is the space R an (s)DH-space?

Recall ([Bu]) that a topological space X is called *n*-homogeneous for a positive integer n if for any two subsets A and B of X of cardinality n there exists a homeomorphism $f: X \to X$, such that f(A) = B. The 1-homogeneous spaces are called homogeneous. Further, X is called strongly n-homogeneous if any bijection $f: A \to B$ between subsets A and B of X of cardinality n extends to a homeomorphism $f: X \to X$.

Since any finite subset of a Hausdorff space is Θ -discrete we have the following fact.

Proposition 1.4. ([BBHS, Corollary 1 for the space B and Corollary 3 for the space R]) Each $(s)\Theta$ -DH space (resp. (s)DH space) is (strongly) n-homogeneous for every positive integer n.

In this article we will discuss the introduced classes of topological spaces, and their relationships to other classes of spaces with homogeneity-type properties. We prove that several classical spaces possess the (strong) discrete homogeneity.

The reader can find all standard notions mentioned in the text in [E] or [M]. Everywhere below, \mathbb{N} , \mathbb{Q} , \mathbb{P} , \mathbb{C} , \mathbb{R} , S^1 , \mathbb{S} denote the sets of natural, rational, and irrational numbers, the Cantor set, real line, the circle, and the Sorgenfrey line, respectively.

2. The case of regular Hausdorff spaces

The following statement is evident.

Proposition 2.1. Let X be a regular Hausdorff space. Then X is an $(s)\Theta$ -DH space iff it is an (s)DH space.

It is easy to see that the real line \mathbb{R} is not strongly 3-homogeneous (although it is strongly 2-homogeneous) and the circle S^1 is not strongly 4homogeneous (although it is strongly 3-homogeneous). Therefore both these spaces are not sDH spaces. Note also that S^1 is an example of a DH space that is not sDH.

Proposition 2.2. Any infinite discrete space X is strongly n-homogeneous for every positive integer n, but not a DH space.

Proof. It is clear that X is strongly n-homogeneous for every positive integer n. We will show that X is not a DH space. Let a, b be two different points of X. Put $A = X \setminus \{a\}$ and $B = X \setminus \{a, b\}$. Clearly, |A| = |B|, and A, B are discrete subsets of X. However, there is no bijection $f: X \to X$ such that f(A) = B.

A Hausdorff space is called *strongly homogeneous* (see [vM]) provided that all nonempty clopen subspaces are homeomorphic.

The spaces \mathbb{C} , \mathbb{Q} , \mathbb{P} , $\mathbb{C} \times \mathbb{Q}$, $\mathbb{C} \times \mathbb{P}$ and \mathbb{S} are strongly homogeneous (for the product $\mathbb{C} \times \mathbb{Q}$ see [AU], and for the product $\mathbb{C} \times \mathbb{P}$ see [vM]).

It is easy to see that each zero-dimensional in the sense of dimension ind first countable strongly homogeneous space is homogeneous. See [E, p. 383] for the definition of the small inductive dimension ind.

The following interesting result was obtained in [vM].

Theorem 2.3. ([vM, Theorem 3.1]) Let X be a strongly homogeneous separable metrizable zero-dimensional space X and let F and G be closed nowhere dense subsets of X. If $f: F \to G$ is a homeomorphism then f can be extended to an automorphism \overline{f} of X.

From Theorem 2.3 we immediately obtain the following.

Proposition 2.4. Every strongly homogeneous separable metrizable zero-dimensional space is sDH. In particular, the spaces \mathbb{C} , \mathbb{Q} , \mathbb{P} , $\mathbb{C} \times \mathbb{Q}$, $\mathbb{C} \times \mathbb{P}$ are sDH.

In [vM] it was observed that the zero-dimensional separable metrizable topological group H from [vD] (in fact, H is a subgroup of the circle group) is not strongly homogeneous. The necessary property of H for the above fact is the following: if F and G are homeomorphic closed subspaces of H then $H \setminus F$ and $H \setminus G$ are homeomorphic (note that we do not require that F and G are proper closed subsets of H).

We will prove that H is sDH.

Let us say that a Hausdorff space X possesses a property \mathcal{D} if for any proper clopen homeomorphic subspaces F and G of X the subspaces $X \setminus F$ and $X \setminus G$ are also homeomorphic.

It is easy to see that the group H as well as any strongly homogeneous space possesses property \mathcal{D} .

Recall further that a space X is called *collectionwise normal* if X is T_1 and for every discrete family $\{F_s\}_{s\in S}$ of closed subsets of X there exists a discrete family $\{V_s\}_{s\in S}$ of open subsets of X such that $F_s \subset V_s$ for every $s \in S$. It is a well-known fact that every paracompact space is collectionwise normal ([E, p. 305]).

Lemma 2.5. Let X be a collectionwise normal space with $\operatorname{ind} X = 0$, and A be a discrete subset of X. Then there exists disjoint family $\{V_a: a \in A\}$ of clopen subsets of X such that $a \in V_a$ for each $a \in A$, and the set $X \setminus \bigcup_{a \in A} V_a$ is also clopen.

Proof. Since the set A is discrete and the space X is T_1 , the family $F_a = \{\{a\}: a \in A\}$, is discrete and consists of closed subsets of X. Therefore there exists a discrete family $\{V_a\}_{a\in A}$ of open subsets of X such that $a \in V_a$ for every $a \in A$. We may assume that all $V_a, a \in A$, are clopen because ind X = 0. Since the family $\{V_a\}_{a\in A}$ is discrete, the set $\bigcup_{a\in A}V_a$ is closed. Hence the set $X \setminus \bigcup_{a\in A}V_a$ is also clopen. \Box

Theorem 2.6. Let X be a collectionwise normal homogeneous space with ind X = 0 and with property \mathcal{D} . Then X is sDH.

Proof. Consider two discrete subsets A, B of X and any bijection $f: A \rightarrow B$. Since the space X is collectionwise normal and ind X = 0 we can apply Lemma 2.5 to each of the sets A and B. So

- (a) there exists a disjoint family $\{V_a : a \in A\}$ of clopen subsets of X such that $a \in V_a$ for each $a \in A$, and the set $X \setminus \bigcup_{a \in A} V_a$ is proper clopen, and
- (b) there exists a disjoint family $\{W_b : b \in B\}$ of clopen subsets of X such that $b \in W_b$ for each $b \in B$, and the set $X \setminus \bigcup_{b \in B} W_b$ is proper clopen.

Furthermore, since X is homogeneous we can assume that for each $a \in A$ the subspaces V_a and $W_{f(a)}$ are homeomorphic and for a homeomorphism $f_a: V_a \to W_{f(a)}$ we have $f_a(a) = f(a) = b$. Set now $F = \bigcup_{a \in A} V_a$ and $G = \bigcup_{b \in B} W_b$. Note that F and G are clopen, proper and homeomorphic. So by the property \mathcal{D} the subspaces $X \setminus F$ and $X \setminus G$ are also homeomorphic. This easily implies that there exists an automorphism \overline{f} of X which extends f.

Corollary 2.7. Let X be a collectionwise normal first countable strongly homogeneous topological space with ind X = 0. Then X is sDH.

Corollary 2.8. The Sorgenfrey line S is sDH.

Corollary 2.9. The topological group H from [vD] is sDH.

Question 2.10. (a) Is H^n sDH for each positive integer n? (b) Is \mathbb{S}^n sDH for each positive integer n?

Theorem 2.3 motivates the following definition.

Definition 2.11. A separable metrizable zero-dimensional topological space X is called nowhere dense homogeneous (NDH), if for any two closed nowhere dense subsets F and G of X and any homeomorphism $f: F \to G$ there exists an automorphism \bar{f} of X which extends f.

The discussion above indicates that each separable metrizable zerodimensional strongly homogeneous space X is NDH, and each NDH space is sDH.

Question 2.12. (a) Is there NDH space which is not strongly homogeneous?

(b) Is there separable metrizable zero-dimensional sDH space which is not NDH?

Since the group H is sDH but not strongly homogeneous, H provides a positive answer to one of the questions.

Let us continue with the case ind ≥ 1 .

For x in a metric space X and r > 0 let B(x, r) be the open ball of radius r centred at x. We will also let $B(x, 0) = \{x\}$.

Next, we will prove that \mathbb{R}^n is sDH for all $n \ge 2$. Since the proofs differ for n = 2 and n > 2, we divide the proofs into two propositions below. Note that these results show that the product of two DH spaces (in this case, \mathbb{R} , which is not sDH) can be sDH.

Proposition 2.13. \mathbb{R}^2 is an sDH space.

Proof. First, we will show that \mathbb{R}^2 is DH. Let A be a countable discrete subset of \mathbb{R}^2 . We will construct a homeomorphism $h: \mathbb{R}^2 \to \mathbb{R}^2$ such that $f(A) = \mathbb{N}$, where \mathbb{N} is viewed as a subset of \mathbb{R}^2 . Using general position argument, we may assume that each circle centred at the origin contains at most one point from A. Hence, $A = \{a_k \mid k \ge 1\}$ and we may assume that $0 < ||a_1|| < ||a_2|| < ||a_3|| < \ldots$ Let $\epsilon_k = \min\{||a_{k+1}|| - ||a_k||, ||a_k|| - ||a_{k-1}||\}, k \ge 1$ (here, we let $a_0 = (0, 0)$). Let U_k be $\epsilon_k/3$ - neighbourhood of a circle of radius $||a_k||$ centred at the origin. For each k, there exists a homeomorphism f_k of \mathbb{R}^2 moving a_k to $(||a_k||, 0)$ and being identity outside U_k . We can now define the limit f of $f_k \circ$ $f_{k-1}... \circ f_1$. Clearly, f is a homeomorphism. Note that A' = f(A)is a discrete subset of the positive part of first coordinate axis in \mathbb{R}^2 . Clearly, there exists a homeomorphism g of \mathbb{R}^2 such that $g(A') = \mathbb{N}$. Now we let $h = g \circ f$.

Further, let A and B be two discrete countable subsets of \mathbb{R}^2 . Again, using general position argument, we may assume that $A \cap B = \emptyset$. We want to extend a bijection between A and B to a homeomorphism of \mathbb{R}^2 . For this, we first consider a homeomorphism of \mathbb{R}^2 that maps $A \cup B$ onto N (it exists because of the first part of the proof). Thus, we reduce the task of extending bijection between A and B to the following: for two disjoint infinite subsets C and D of N such that $C \cup D = \mathbb{N}$ and a bijection of C onto D, find a homeomorphism of \mathbb{R}^2 extending this bijection. So let $C = \{c_k \mid k \geq 1\}, D = \{d_k \mid k \geq 1\}$ $k \geq 1$, and we want to extend a bijection sending c_k to d_k for each k. Fix a countable collection of rays $R_k, k \geq 1$, in \mathbb{R}^2 , such that $R_i \cap R_j = (0,0)$ for all $i \neq j$. Further, for each k, let U_k and V_k be 1/3neighbourhoods of circles of radii $||c_k||$ and $||d_k||$, respectively, centred at the origin. There exists a homeomorphism $f_k, k = 1, 2, ...$, such that $c'_k = f_k(c_k) \in R_k, \, d'_k = f_k(d_k) \in R_k, \, \|c'_k\| = \|c_k\|, \, \|d'_k\| = \|d_k\| \text{ and } f_k$ is the identity outside of $U_k \cup V_k$. Let f be the limit of $f_k \circ f_{k-1} \dots \circ f_1$. Clearly, f is a homeomorphism. Note that the collection of straight line segments $[c'_k, d'_k]$, joining c'_k and d'_k , is discrete. For each k, let g_k be a homeomorphism of \mathbb{R}^2 such that $g_k(c'_k) = d'_k$, which is the identity outside of some "narrow" neighbourhood of $[c'_k, d'_k]$. We can define a homeomorphism g which is the limit of $g_k \circ g_{k-1} \dots \circ g_1$. The homeomorphism $f^{-1} \circ q \circ f$ is the one we are looking for.

Proposition 2.14. Every finite-dimensional Euclidean space \mathbb{R}^n , n > 2, is an sDH space.

Proof. Indeed, let A and B be two discrete subsets of \mathbb{R}^n of the same cardinality. Hence both A and B are either finite or countable. We will assume A and B are countable (the finite case is similar). Therefore $A = \{a_k : k \ge 1\}$ and $B = \{b_k : k \ge 1\}$. We want to extend a bijection sending a_k to b_k for each $k \ge 1$. For any k there exists a path P_k joining a_k and b_k , that avoids the ball of radius $\frac{1}{2}\min\{||a_k||, ||b_k||\}$ centred at the origin, and also avoids all other a's and b's. Using general position argument in \mathbb{R}^n , n > 2, we may also assume that $P_i \cap P_j = \emptyset$ for $i \ne j$. There exists a homeomorphism $f_k : \mathbb{R}^n \to \mathbb{R}^n$ that maps a_k to b_k and is the identity outside some "narrow" neighbourhood of the path P_k . We can now define the limit of $f_k \circ f_{k-1} \dots \circ f_1$, which is the homeomorphism extending the bijection between A and B.

Proposition 2.15. The separable Hilbert space l_2 is sDH.

Proof. As in the proof of the previous propositon, consider two countable subsets $A = \{a_k : k \ge 1\}$ and $B = \{b_k : k \ge 1\}$ of l_2 . We want to extend a bijection sending a_k to b_k for each $k \ge 1$.

For each $x \in l_2$ there exists $r_x > 0$ such that $B(x, r_x) \cap (A \cup B) = \{x\} \cap (A \cup B)$. Additionally, we may assume that $r_{a_k} < 1/k$ and $r_{b_k} < 1/k$ for all k. Let $\mathcal{U} = \{B(x, r_x) \mid x \in l_2\}$.

Join each a_k to b_k by a straight line segment. This collection of segments can be viewed as a map $g: \bigoplus_{k=1}^{\infty} ([0,1] \times \{k\}) \to l_2$. Therefore there exists a map $g': \bigoplus_{k=1}^{\infty} ([0,1] \times \{k\}) \to l_2$ such that the family $\{g'([0,1] \times \{k\}) \mid k \in \mathbb{N}\}$ is discrete. Each $g'([0,1] \times \{k\})$ is a path such that $g'(0) \in B(a_k, r_{a_k})$ and $g'(1) \in B(b_k, r_{b_k})$. By adding straight line segments from a_k to g'(0) and from b_k to g'(1) we obtain a discrete family of paths P_k in l_2 , joining a_k and b_k . Moreover, we may even assume that each P_k is piecewise linear arc. There exists a homeomorphism $f_k: l_2 \to l_2$ that maps a_k to b_k and is identity outside some "narrow" neighbourhood V_k of the path P_k , such that the family $\{V_k: k \ge 1\}$ is discrete. As in the proof of the previous proposition, the limit of $f_k \circ f_{k-1} \dots \circ f_1$ is the desired homeomorphism.

Recall that a space X is called strongly locally homogeneous (SLH) if it has a basis of open sets \mathcal{B} such that for every $U \in \mathcal{B}$ and any two points x and y in U there exists a homeomorphism $f: X \to X$ such that f(x) = y and f is identity on $X \setminus U$. Note that any n-manifold for n > 1 and any Hilbert space manifold are examples of SLH spaces. It was shown in [Ba] that a connected SLH space, no two-point subset of which disconnects it, is strongly n-homogeneous for any n.

Example 2.16. Let $M = \mathbb{R} \times S^1$. Then M is a 2-manifold which is strongly n-homogeneous for any n and SLH but not DH

Proof. Being a 2-manifold, M is both strongly n-homogeneous and SLH. Fix a point $s \in S^1$ and let $A = \{(n, s) : n \in \mathbb{N}\}$ and $B = \{(\pm n, s) : n \in \mathbb{N}\}$. We claim that there is no autohomeomorphism of M that maps A onto B. Indeed, suppose $f \colon M \to M$ is such homeomorphism. Let $N = [0, \infty) \times S^1$. Note that f(N) is a submanifold of M, containing B, with the boundary being simple closed curve $C = f(\{0\} \times S^1)$. If C is nullhomotopic, then C is a common boundary of a topological disk and a manifold, homeomorphic to a sphere with three holes. Thus, in this case f(N) cannot be homeomorphic to N. Otherwise, C divides M into two submonifolds, none of which can contain the whole B, contradiction.

Question 2.17. Which metrizable manifolds are DH or sDH?

Problem 2.18. Find conditions for SLH space to be (s)DH.

In this question, one can impose various additional conditions such as being completely metrizable and separable, connected, arc-wise connected, locally compact, etc.

Since every discrete subspace of a compact Hausdorff space is finite we have the following statement.

Proposition 2.19. Let X be a compact Hausdorff space. Then X is a (s)DH space iff it is (strongly) n-homogeneous for every positive integer n.

Besides the Cantor set \mathbb{C} (see Proposition 2.4), an obvious example of a zero-dimensional sDH compact Hausdorff space is the Alexandroff double arrow space [AU2]. Note also, that since \mathbb{C} is the only homogeneous metrizable zero-dimensional infinite compactum, it is also the only metrizable zero-dimensional infinite compactum which is sDH. In general case, however, the following problem seems interesting.

Problem 2.20. Characterize all zero-dimensional sDH compacta.

Recall that a closed subset A of a metric space X is called a Z-set if for any $\epsilon > 0$ there exists a map $f: X \to X$ such that f is ϵ -close to the identity map and $f(X) \cap A = \emptyset$. Let μ^k denote the Menger cube for each $k = 1, 2, \ldots$, and $Q = [0, 1]^{\omega}$ be the Hilbert cube. It is well-known that any finite subset of μ^k or Q is a Z-set in the respective space, and any homeomorphism between two Z-sets of μ^k or Q can be extended to a homeomorphism of the whole space (see, e.g. [Ch] for the case of the Hilbert cube, and [A, Be] for the case of Menger compacta). Consequently, Q and μ^k are examples of sDH compacta.

Moreover, it is well-known [A] that μ^1 and S^1 are the only homogeneous 1-dimensional Peano continua (by a continuum we mean metrizable compact connected space). By the result of [U], any 2homogeneous continuum is locally connected. Therefore we have the following observation.

Proposition 2.21. μ^1 and S^1 are the only DH1-dimensional continua.

Since as was observed earlier S^1 is not sDH but μ^1 is sDH, we get immediately the following corollary.

Corollary 2.22. μ^1 is the only sDH 1-dimensional continuum

8

Note that according to [KKT], the product $\mu^1 \times \mu^1$ is not 2-homogeneous, and hence not DH. Moreover, it is easy to see that $\mathbb{R}^n \times \mathbb{Q}$, $n \ge 1$, is also not 2-homogeneous. Therefore the property (s)DH is not preserved by products.

In view of the above discussion about n-homogeneity, as well as Example 2.16, the following question appears to be interesting.

Question 2.23. Let X be a connected separable metrizable (complete, locally compact, locally connected etc.) space, that is (strongly) n-homogeneous for any n. What additional conditions on X imply that X is (s)DH?

3. The case of non-regular Hausdorff spaces

Since every countable regular Hausdorff space cannot be connected, the Bing space B and the Ritter spaces R are non-regular.

Let X be a topological space with the topology τ .

Recall (cf. [MRV]) that the topology $\hat{\tau} \subseteq \tau$ on the set X generated by the base consisting of all regular open subsets of the topological space X is called the semi-regularization of τ . (An open set O of a topological space X is called regular open if O coincides with the interior of its closure Cl(O) in X.)

We will denote the space $(X, \hat{\tau})$ by \hat{X} . Recall ([R]) that $\hat{B} = R$. Below we recall some facts about the semi-regularization topologies.

Proposition 3.1. (cf. [MRV]) Let X be a topological space with the topology τ . Then we have the following.

- (a) \hat{X} is Hausdorff iff X is Hausdorff;
- (b) X is connected iff X is connected;
- (c) if X is locally connected then \hat{X} is also locally connected but the converse is false.

Note that the Bing space B is not locally connected. The following lemma is evident.

Lemma 3.2. Let X be a Hausdorff topological space. If D is a Θ -discrete (resp. discrete) subset of \hat{X} then D is a Θ -discrete (resp. discrete) subset of X.

Theorem 3.3. (For the spaces B and R, [BBHS]) If X is an $s\Theta$ -DH space then \hat{X} is also an $s\Theta$ -DH space.

Proof. Let A and B be Θ -discrete subsets of \hat{X} and $f: A \to B$ be a bijection. By Lemma 3.2 A and B are Θ -discrete subsets of X. Hence f can be extended to a homeomorphism $f: X \to X$ of X which is also

a homeomorphism of \hat{X} since $f(\operatorname{Int}(\operatorname{Cl}(O))) = \operatorname{Int}(\operatorname{Cl}(f(O)))$ for any open subset O of X.

In the light of the examples of Bing and Ritter, the following question looks interesting.

Question 3.4. (a) Are there examples of countable connected $s\Theta$ -DH spaces, other than B and R?

(b) Are there examples of countable connected Hausdorff sDH spaces?

The following fact is relevant to this question.

Proposition 3.5. ([MRV, Lemma 5]) For every topological space X with the topology τ we have $\hat{\tau} = \hat{\tau}$, i.e. the process of semi-regularizing of a topological space is an idempotent operation.

In particular, we have $\hat{R} = R$.

4. Appendix

The Bing space B ([Bi]) is the rational half plane $\{(x, y) \in \mathbb{Q} \times \mathbb{Q} : y \geq 0\}$ endowed with the topology τ consisting of the sets $U \subset \mathbb{B}$ such that for every point $(a, b) \in U$ there exists $\epsilon > 0$ such that

 $\{(x,0) \in B \colon |x - (a - b/\sqrt{3})| < \epsilon\} \cup \{(x,0) \in B \colon |x - (a + b/\sqrt{3})| < \epsilon\} \subset U.$

References

- [AU] P. Alexandroff and P. Urysohn Uber nulldimensionalle Punktmengen, Math. Ann. 98 (1928) 89-106.
- [AU2] P. Alexandroff and P. Urysohn; "Mémoire sur les espaces topologiques compacts." Verh. Akad. Wetensch. Amsterdam 14 (1929).
- [A] R.D. Anderson, A characterization of the universal curve and a proof of its homogeneity, Ann. of Math. (2) 67 (1958) 313-324.
- [Ba] J.W. Bales, Representable and strongly locally homogeneous spaces and strongly n-homogeneous spaces, Houston J. of Math. 2, no. 3 (1976) 315-327
- [BBHS] I. Banakh, T. Banakh, O. Hryniv, Y. Stelmakh, The connected countable spaces of Bing and Ritter are topologically homogeneous, Topol. Proc. 57 (2021) 149-158
- [Be] M. Bestvina, Characterizing k-dimensional universal Menger compacta, Bull. Amer. Math. Soc. 11, no. 2 (1984) 369-370.
- [Bi] R. H. Bing, A connected countable Hausdorff space, Proc. Amer. Math. Soc. 4 (1953) 474
- [Bu] C. E. Burgess, Some theorems on n-homogeneous continua, Proc. Amer. Math. Soc. 1 (1954) 136-143
- [Ch] T. A. Chapman, Lectures on Hilbert Cube Manifolds, CBMS Regional Conference Series in Mathematics 28, 1976.
- [E] R. Engelking, General Topology, Warszawa, Heldermann Verlag, Berlin, 1989

- [vD] E. K. van Douwen, A measure that knows which sets are homeomorphic, Top. Structures 2, Mathematical Centre Tracts 115 (1979) 67-71
- [KKT] K.K. Kuperberg, W. Kuperberg, W.R.R. Transue, On the 2-homogeneity of Cartesian products, Fund. Math. 110 (1980) no. 2, 131-134
- [vM] J. van Mill, Characterization of some zero-dimensional separable metric spaces, Trans. of the AMS 264, no. 1 (1981) 205-215
- [MRV] M. Mrsevic, I. L. Reilly, M. K. Vamanamurthy, On semi-regularization topologies, J. Austral. Math. Soc. 38 (1985) 40-54
- [M] J. R. Munkres, *Topology*, Prentice Hall, Upper Saddle River, 2000
- [R] G. Ritter, A connected, locally connected, countable Hausdorff space, Amer. Math. Monthly 83:3 (1976) 185-186
- [U] G.S. Ungar, On all kinds of homogeneous spaces, Trans. of the AMS 212 (1975) 393-400.

(V.A. Chatyrko)

Department of Mathematics, Linkoping University, 581 83 Linkoping, Sweden.

vitalij.tjatyrko@liu.se

(A. Karassev)

Department of Mathematics, Nipissing University, North Bay, Canada alexandk@nipissingu.ca