# RIGIDITY OF ACUTE TRIANGULATIONS OF THE PLANE

#### TIANQI WU

ABSTRACT. We show that a uniformly acute triangulation of the plane is rigid under Luo's discrete conformal change, extending previous results on hexagonal triangulations. Our result is a discrete analogue of the conformal rigidity of the plane. We followed He's analytical approach in his work on the rigidity of disk patterns. The main tools include maximum principles, a discrete Liouville theorem, smooth and discrete extremal lengths on networks. The key step is relating the Euclidean discrete conformality to the hyperbolic discrete conformality, to obtain an  $L^{\infty}$  bound on the discrete conformal factor.

## Contents

1. Introduction	1
1.1. Other Related Works	2
1.2. Notations and Conventions	3
1.3. Organization of the Paper	3
1.4. Acknowledgement	3
2. Preparations for the Proof	4
2.1. Extremal Length and Modulus of Annuli	4
2.2. Discrete Harmonic Functions	5
2.3. Differential of the Curvature Map	5
2.4. Maximum Principles	5
2.5. Key Estimates on the Conformal Factors for Geodesic Embeddings	6
3. Proof of Theorem 1.2	6
3.1. Proof of Theorem 1.2 Assuming the Boundedness of $\bar{u}$	6
3.2. Boundedness of the Conformal Factor	8
4. Discrete Extremal Length and the Discrete Liouville Theorem	9
4.1. Electrical Networks and Discrete Extremal Length	9
4.2. Proof of the Discrete Liouville Theorem	10
5. Hyperbolic Maximum Principles and Proof of Lemma 2.9	11
5.1. Proof of the Hyperbolic Maximum Principle	13
Appendix A. Proof of Lemma 5.1	15
References	16

### 1. INTRODUCTION

A fundamental property in conformal geometry is that a conformal embedding of the plane  $\mathbb{R}^2$  to itself must be a similar transformation. In this paper we discretize the plane by triangulations and prove a similar rigidity result under the notion of discrete conformal change introduced by Luo [Luo04].

Let T = (V, E, F) be an (infinite) simplicial topological triangulation of the Euclidean plane  $\mathbb{R}^2$ , where V is the set of vertices, E is the set of edges and F is the set of faces. Given a subcomplex  $T_0 = (V_0, E_0, F_0)$  of T, denote  $|T_0| \to \mathbb{R}^2$  is called *geodesic* if  $\phi$  maps each edge of  $T_0$  to a geodesic arc, i.e., a straight closed line segment. A *piecewise linear metric* (*PL metric* for short) on  $T_0$  is represented by an edge length function  $l \in \mathbb{R}_{>0}^{E_0}$  satisfying the triangle inequalities. A geodesic embedding  $\phi$  of  $T_0$  naturally induces a PL metric  $l = l(\phi)$  on  $T_0$  by letting  $l_{ij} = |\phi(i) - \phi(j)|_2$ . Luo [Luo04] introduced the following notion of discrete conformality.

**Definition 1.1** (Luo [Luo04]). Two PL metrics l, l' on  $T_0 = (V_0, E_0, F_0)$  are discretely conformal if there exists some  $u \in \mathbb{R}^{V_0}$  such that for any edge  $ij \in E_0$ 

$$l'_{ij} = e^{\frac{1}{2}(u_i + u_j)} l_{ij}$$

In this case, u is called a discrete conformal factor, and we denote l' = u \* l.

Given a PL metric l on  $T_0$ , let  $\theta^i_{jk}$  denote the inner angle at the vertex i in the triangle  $\Delta i j k$  under the metric l. Then l is called

- (a) uniformly nondegenerate if there exists a constant  $\epsilon > 0$  such that  $\theta_{jk}^i \ge \epsilon$  for all  $\triangle ijk$  in  $T_0$ , and
- (b) uniformly acute if there exists a constant  $\epsilon > 0$  such that  $\theta_{jk}^i \leq \pi/2 \epsilon$  for all  $\triangle ijk$  in  $T_0$ , and
- (c) Delaunay if  $\theta_{ij}^k + \theta_{ij}^{k'} \leq \pi$  for any pair of adjacent triangles  $\triangle ijk$  and  $\triangle ijk'$  in  $T_0$ .

A uniformly acute PL metric is clearly uniformly nondegenerate and Delaunay. The main result of the paper is the following.

**Theorem 1.2.** Suppose  $\phi$  is a geodesic homeomorphism of T and  $\psi$  is a geodesic embedding of T. If  $l(\phi), l(\psi)$  are discretely conformal and both uniformly acute, then they differ by a constant scaling.

Wu-Gu-Sun [WGS15] first proved Theorem 1.2 for the special case where  $\phi(T)$  is a regular hexagonal triangulation. Dai-Ge-Ma [DGM22] and Luo-Sun-Wu [LSW20] generalized Wu-Gu-Sun's result by allowing  $l(\psi)$  to be only Delaunay rather than uniformly acute. All these works essentially rely on the lattice structure of the embedded vertices  $\phi(V)$ , and apparently cannot be generalized to triangulations without translational invariance. To prove Theorem 1.2, we adopted a different approach, which is developed by He [He99] in his state-of-art work on the rigidity of disk patterns.

1.1. Other Related Works. After Luo introducing the Definition 1.1, various properties regarding the rigidity and convergence of the discrete conformality were discussed in [BPS15][WGS15][GLW19][WZ20][LSW20][LWZ21a][DGM22]. To solve the problem of singularity in the discrete Yamabe flow, Gu et al. [GLSW18][GGL<sup>+</sup>18] proposed a revised notion of discrete conformality for piecewise Euclidean (or hyperbolic) metrics on closed surfaces with marked points, and perfectly solved the prescribed curvature problem. This major improvement in the theory of discrete conformality inspired new advanced numerical methods in computing conformal maps [SWGL15][GSC21][CCS<sup>+</sup>21], as well as further theoretical investigations [Spr19][LW19]. Gu et al. [GLSW18][GGL<sup>+</sup>18] proposed to use the discrete Yamabe flow to numerically compute the target metric in the prescribed curvature problem.

 $\mathbf{2}$ 

Since the discrete Yamabe flow may pass through different combinatorial triangulations, diagonal switches might be needed along the flow. In [Wu14] it is proved that only finitely many diagonal switches are needed in a Yamabe flow. Other works on discrete geometric flows or deformations of triangle meshes could be found in [ZGZ<sup>+</sup>14][GH18] [ZX19][FLZ20][WX21][LWZ21b][LWZ21c][LWZ22][Luo22].

1.2. Notations and Conventions. In the remaining of the paper, we will identify the plane  $\mathbb{R}^2$  as the complex plane  $\mathbb{C}$ . Given 0 < r < r', denote  $D_r = \{z \in \mathbb{C} : |z| < r\}$  and  $A_{r,r'} = \{z \in \mathbb{C} : r < |z| < r'\}$ . We also denote  $D = D_1$  as the unit open disk. Given a subset A of  $\mathbb{C}$ ,  $A^c$  denote the complement  $\mathbb{C} \setminus A$  and  $\partial A$  denotes the boundary of A in  $\mathbb{C}$ . Given two subsets A, B of  $\mathbb{C}$ , the diameter of A is denoted by

$$\operatorname{diam}(A) = \sup\{|z - z'| : z, z' \in A\},\$$

and the distance between A, B is denoted by

$$d(A, B) = \inf\{|z - z'| : z \in A, z' \in B\}.$$

Given a subset  $V_0$  of V, we use the following notations and conventions.

- (a) The complement of  $V_0$  is denoted as  $V_0^c = V \setminus V_0$ .
- (b) The *boundary* of  $V_0$  is denoted as

 $\partial V_0 = \{i \in V_0 : \text{there exists } j \in V_0^c \text{ such that } ij \in E\}.$ 

(c) The *interior* of  $V_0$  is denoted as

$$int(V_0) = V_0 \setminus \partial V_0 = \{i \in V_0 : j \in V_0 \text{ if } ij \in E\}.$$

(d) The *closure* of  $V_0$  is denoted as

$$\overline{V_0} = V_0 \cup \partial(V_0^c) = (int(V_0^c))^c.$$

- (e) The subcomplex generated by  $V_0$  is denoted as  $T(V_0)$ .
- (f) Denote  $E(V_0) = \{ij \in E : i \in int(V_0) \text{ or } j \in int(V_0)\}$ . Notice that  $E(V_0)$  generally is not the set of edges in  $T(V_0)$ .
- (g) A real-valued function on  $V_0$  is often identifies as a vector in  $\mathbb{R}^{V_0}$ .

Given  $i \in V$ , the 1-ring neighborhood of i is the subcomplex generated by i and its neighbors. In other words, the 1-ring neighborhood of i is

$$T(\{i\} \cup \{j \in V : ij \in E\}).$$

Furthermore, we denote  $R_i$  as the underlying space of the 1-ring neighborhood of i. Given a subcomplex  $T_0 = (V_0, E_0, F_0)$  of T and  $l \in \mathbb{R}^{E_0}$  and  $u \in \mathbb{R}^{V_0}$ , if u \* l is a PL metric then

(a)  $\theta_{jk}^i(u) = \theta_{jk}^i(u, l)$  denotes the inner angle of  $\triangle ijk$  at *i* under u \* l, and (b)  $K_i(u) = K_i(u * l)$  denotes the discrete curvature

$$K_i(u) = 2\pi - \sum_{jk: \triangle ijk \in F} \theta_{jk}^i(u).$$

1.3. Organization of the Paper. In Section 2 we introduce necessary properties and tools for the proof of the main theorem. The proof of main Theorem 1.2 is given in Section 3. Section 4 gives a proof of a discrete Liouville theorem, which is used in proving Theorem 1.2. Section 5 proves a key estimate for the discrete conformal factor by relating to the hyperbolic discrete conformality.

1.4. Acknowledgement. The work is supported in part by NSF 1760471.

#### 2. Preparations for the Proof

2.1. Extremal Length and Modulus of Annuli. We briefly review the notions of extremal length and conformal modulus. The definitions and properties discussed here are mostly well-known. One may refer [Ahl10] and [LV73] for more comprehensive introductions.

A closed annulus is a subset of  $\mathbb{C}$  that is homeomorphic to  $\{z \in \mathbb{C} : 1 \leq |z| \leq 2\}$ . An *(open) annulus* is the interior of a closed annulus. Given an annulus A, denote  $\Gamma = \Gamma(A)$  as the set of smooth simple closed curves in A separating the two boundary components of A. A real-valued Borel measurable function f on A is called *admissible* if  $\int_{\gamma} fds \geq 1$  for all  $\gamma \in \Gamma$ . Here ds denotes the element of arc length. The *(conformal) modulus* of A is defined as

$$\operatorname{Mod}(A) = \inf\{\int_A f^2 : f \text{ is admissible}\},\$$

where  $\int_A f^2$  denotes the integral of  $f(z)^2$  against the 2-dim Lebesgue measure on A. From the definition it is straightforward to verify that Mod(A) is conformally invariant. Furthermore, if  $f : A \to A'$  is a K-quasiconformal homeomorphism between two annuli, then

$$\frac{1}{K} \cdot \operatorname{Mod}(A) \le \operatorname{Mod}(A') \le K \cdot \operatorname{Mod}(A).$$

Given 0 < r < r', denote  $A_{r,r'}$  as the annulus  $\{z \in \mathbb{C} : r < |z| < r'\}$ . It is well-known that

$$\operatorname{Mod}(A_{r,r'}) = \frac{1}{2\pi} \log \frac{r'}{r}$$

Intuitively, the conformal modulus measures the relative thickness of an annulus. If an annulus A in  $\mathbb{C}\setminus\{0\}$  contains  $A_{r,r'}$ , then it is "thicker" than  $A_{r,r'}$  and one can show that

$$\operatorname{Mod}(A) \ge \operatorname{Mod}(A_{r,r'}) = \frac{1}{2\pi} \log \frac{r'}{r}.$$

On the other hand, we have that

**Lemma 2.1.** Suppose  $A \subseteq \mathbb{C} \setminus \{0\}$  is an annulus separating 0 from the infinity. If  $Mod(A) \ge 100$ , then  $A \supseteq A_{r,2r}$  for some r > 0.

*Proof.* Deonte B as the bounded component of  $\mathbb{C} - A$ , and  $r = \max\{|z| : z \in B\}$  and  $R = \min\{|z| : z \in (B \cup A)^c\}$ . If  $R \ge 2r$  we are done. So we may assume R < 2r.

Then  $D_{2r} \cap \gamma \neq \emptyset$  for all  $\gamma \in \Gamma(A)$ . Let f be a function on A such that f(z) = 1/ron  $A \cap D_{3r}$  and f(z) = 0 on  $A \setminus D_{3r}$ . If  $\gamma \in \Gamma$  and  $\gamma \subseteq D_{3r}$ ,

$$\int_{\gamma} f ds = s(\gamma) \cdot \frac{1}{r} \ge 2 \cdot \operatorname{diam}(B) \cdot \frac{1}{r} = 2r \cdot \frac{1}{r} > 1.$$

If  $\gamma \in \Gamma$  and  $\gamma \subsetneq D_{3r}$ , then  $\gamma$  is a connected curve connecting  $D_{2r}$  and  $D_{3r}^c$  and

$$\int_{\gamma} f ds \ge d(D_{2r}, D_{3r}^c) \cdot \frac{1}{r} = r \cdot \frac{1}{r} = 1$$

So f is admissible and

$$\operatorname{Mod}(A) \le \int_{A} f^{2} = \frac{1}{r^{2}} \cdot \operatorname{Area}(A \cap D_{3r}) \le \frac{\pi (3r)^{2}}{r^{2}} = 9\pi < 100.$$

This contradicts with our assumption.

**Remark 2.2.** To some extend, Lemma 2.1 is a consequence of Teichmüller's result on extremal annuli (see Theorem 4-7 in [Ahl10]). The constant 100 is chosen for convenience and should not be optimal.

2.2. Discrete Harmonic Functions. Given  $V_0 \subset V$  and  $\eta \in \mathbb{R}_{>0}^{E(V_0)}$ , a discrete function  $f : V_0 \to \mathbb{R}$ , or equivalently a vector  $f \in \mathbb{R}^{V_0}$ , is called *harmonic at*  $i \in int(V_0)$  if

$$\sum_{j:ij\in E}\eta_{ij}(f_j-f_i)=0.$$

The following result is well-known and easy to prove.

**Proposition 2.3.** Suppose  $V_0$  is a finite subset of V and  $\eta_{ij} \in \mathbb{R}_{>0}^{E(V_0)}$ .

(a) If  $f \in \mathbb{R}^{V_0}$  is harmonic at i for all  $i \in int(V_0)$ , then for all  $i \in V_0$ 

$$|f_i| \le \max_{j \in \partial V_0} |f_j|.$$

(b) Given  $g: \partial V_0 \to \mathbb{R}$ , there exists a unique function  $f: V_0 \to \mathbb{R}$  such that (i)  $f_i = g_i$  on  $\partial V_0$ , and (ii) f is harmonic at any  $i \in int(V_0)$ . Furthermore, such a map  $(\eta, g) \mapsto f$  is smooth from  $\mathbb{R}_{>0}^{E(V_0)} \times \mathbb{R}^{\partial V_0}$  to  $\mathbb{R}^{V_0}$ .

Given  $\eta \in \mathbb{R}^{E}_{>0}$ ,  $f \in \mathbb{R}^{V}$  is called *harmonic* if it is harmonic at all points in V. It is well-known by Liouville's Theorem that any bounded smooth harmonic function on the plane is constant. Here we have a discrete version of Liouville's Theorem.

**Theorem 2.4.** Suppose  $\phi$  is a geodesic embedding of T and  $l(\phi)$  is uniformly nondegenerate. Given  $\eta \in \mathbb{R}^{E}_{\geq 0}$  with  $|\eta|_{\infty} < \infty$ , then any bounded harmonic function on  $(T, \eta)$  is constant.

The proof of Theorem 2.4 is postponed to Section 4.

2.3. Differential of the Curvature Map. The differential of  $K_i(u)$  has the following elegant formula, first proposed by Luo [Luo04].

**Proposition 2.5** (Adapted from Theorem 2.1 in [Lu04]). Suppose  $T_0 = (V_0, E_0, F_0)$  is a 1-ring neighborhood of  $i \in V$  and  $l \in \mathbb{R}^{E_0}$ . Then  $K_i = K_i(u)$  is a smooth function on an open set in  $\mathbb{R}^{V_0}$ , and

$$dK_i = \sum_{j:ij\in E} \eta_{ij} (du_i - du_j).$$

where  $\eta_{ij} = \eta_{ij}(u)$  is defined to be

(2.1) 
$$\eta_{ij}(u) = \frac{1}{2} \left( \cot \theta_{ij}^k(u) + \cot \theta_{ij}^{k'}(u) \right),$$

where  $\triangle ijk, \triangle ijk'$  are the two triangles in F containing edge ij.

2.4. Maximum Principles. We need the following maximum principle.

**Lemma 2.6.** Suppose  $V_0$  is a finite subset of V, and u \* l, u' \* l are Delaunay PL metrics on  $T(V_0)$ . If  $K_i(u) = K_i(u') = 0$  for all  $i \in int(V_0)$ , then for all  $i \in V_0$  $|u'_i - u_i| \leq \max_{j \in \partial V_0} |u'_j - u_j|.$ 

Lemma 2.6 is a standard consequence the following local maximum principle, which is adapted from Lemma 2.12 in [DGM22] (or Theorem 3.1 in [LSW20]).

**Lemma 2.7.** Suppose  $i \in V$  and  $T_0 = (V_0, E_0, F_0)$  is the 1-ring neighborhood of i in V. Given  $l \in \mathbb{R}^{E_0}$ , if u \* l, u' \* l are two Delaunay PL metrics on  $T_0$  and  $K_i(u) = K_i(u') = 0$ , then

$$u_i' - u_i \le \max_{j: ij \in E} (u_j' - u_j)$$

and the equality holds if  $(u'_i - u_i) = (u'_j - u_j)$  for any neighbor j of i.

**Remark 2.8.** Lemma 2.12 in [DGM22] is a special case of our Lemma 2.7, where  $u_i = u'_i = 0$  is further assumed. However, by the scaling invariance these two Lemmas are really equivalent.

### 2.5. Key Estimates on the Conformal Factors for Geodesic Embeddings.

**Lemma 2.9.** Suppose  $\epsilon > 0$  and  $\phi, \psi$  are two geodesic embeddings of a subcomplex  $T_0 = (V_0, E_0, F_0)$  of T, such that

(i)  $l(\psi) = u * l(\phi)$  for some  $u \in \mathbb{R}^{V_0}$ , and

(ii) the inner angles in both PL metrics  $l(\phi)$  and  $l(\psi)$  are at most  $\pi/2 - \epsilon$ .

Given r, r' > 0 and  $i \in V$ , if

 $\phi(|T_0|) \subseteq D_r$ 

and

$$\psi(i) \in D_{r'/2} \subseteq D_{r'} \subseteq \psi(|T_0|)$$

then

$$u_i \ge \log(r'/r) - M$$

for some constant  $M = M(\epsilon) > 0$ .

### 3. Proof of Theorem 1.2

Assume  $l(\psi) = \bar{u} * l(\phi)$ , and all the inner angles in  $l(\phi), l(\psi)$  are at most  $\pi/2 - \epsilon$  for a constant  $\epsilon > 0$ . We will first prove Theorem 1.2 assuming  $\bar{u} : V \to \mathbb{R}$  is bounded in Section 3.1, and then prove  $\bar{u}$  is bounded in Section 3.2.

3.1. Proof of Theorem 1.2 Assuming the Boundedness of  $\bar{u}$ . Let us prove by contradiction and assume that  $\bar{u}$  is not constant. Without loss of generality, we can do a scaling and assume

$$\inf_{i \in V} \bar{u}_i < 0 < \sup_{i \in V} \bar{u}_i$$

and

$$-\inf_{i\in V}\bar{u}_i = \sup_{i\in V}\bar{u}_i = |\bar{u}|_{\infty}.$$

By a standard compactness argument, it is not difficult to see that there exists a small constant  $\delta = \delta(\epsilon, \bar{u}) \in (0, |\bar{u}|_{\infty})$  such that if  $|u|_{\infty} < 2\delta$ ,

$$\theta^i_{jk}(u) = \theta^i_{jk}(u, l(\phi)) \ge \pi/2 - \epsilon/2$$

for all  $\triangle ijk \in F$ . Pick a sequence of increasing subsets  $V_n$  of V such that  $\bigcup_{n=1}^{\infty} V_n = V$ . For each  $n \in \mathbb{Z}_{>0}$ , we will construct a smooth  $\mathbb{R}^{V_n}$ -valued function  $u^{(n)}(t) = [u_i^{(n)}(t)]_{i \in V_n}$  on  $(-2\delta, 2\delta)$  such that

- (a)  $u^{(n)}(0) = 0$ , and
- (b)  $\dot{u}_i^{(n)}(t) = \bar{u}_i / |\bar{u}|_{\infty}$  if  $i \in \partial V_n$ , and

(3.1) 
$$\sum_{j:ij\in E} \eta_{ij}(u^{(n)}(t))(\dot{u}_i^{(n)}(t) - \dot{u}_j^{(n)}(t)) = 0$$

(c) if  $i \in int(V_n)$  then

where  $\eta_{ij}(u)$  is defined for all  $ij \in E(V_n)$  as in equation (2.1).

The conditions (b) and (c) give an autonomous ODE system on

$$\mathcal{U}_n = \{ u \in \mathbb{R}^{V_n} : |u|_\infty < 2\delta \}.$$

Notice that  $\eta_{ij}(u) > 0$  if  $u \in \mathcal{U}_n$ . Then by part (b) of Lemma 2.3,  $\dot{u}^{(n)}(t)$  is smoothly determined by  $u^{(n)}(t)$  on  $\mathcal{U}_n$ . Given the initial condition  $u^{(n)}(0) = 0$ , assume the maximum existence interval for this ODE system on  $\mathcal{U}_n$  is  $(t_{\min}, t_{\max})$ where  $t_{\min} \in [-\infty, 0)$  and  $t_{\max} \in (0, \infty]$ . By the maximum principle (part (a) in Lemma 2.3), for all  $i \in V_n$ 

$$|\dot{u}^{(n)}|_{\infty} \le \max_{j \in \partial V_n} |\dot{u}_j^{(n)}| = \max_{j \in \partial V_n} |\bar{u}_j| / |\bar{u}|_{\infty} \le 1.$$

So  $|u^{(n)}(t)|_{\infty} \leq t \leq t_{\max}$  for all  $t \in [0, t_{\max})$ . By the maximality of  $t_{\max}$ ,  $t_{\max} = \infty$  or

$$|u^{(n)}(t)|_{\infty} \to 2\delta$$
 as  $t \to t_{\max}$ .

So  $t_{\max} \ge 2\delta$  and by a similar reason  $t_{\min} \le -2\delta$ .  $u^{(n)}(t)$  is indeed well-defined on  $(-2\delta, 2\delta)$ . By Proposition 2.5 and equation (3.1),  $K_i(u^{(n)}(t)) = 0$  for all  $i \in int(V_n)$ . Then by Lemma 2.6, for all  $i \in V_n$ 

(3.2) 
$$|\bar{u}_i - u_i^{(n)}(\delta)| \leq \max_{j \in \partial V_n} |\bar{u}_j - u_j^{(n)}(\delta)| = \max_{j \in \partial V_n} \left( \bar{u}_j - \delta \cdot \frac{\bar{u}_j}{|\bar{u}|_{\infty}} \right)$$
$$\leq (1 - \frac{\delta}{|\bar{u}|_{\infty}}) |\bar{u}|_{\infty} = |\bar{u}|_{\infty} - \delta.$$

By picking a subsequence, we may assume that  $u_i^{(n)}$  converge to  $u_i^*$  on  $[0, \delta]$  uniformly for all  $i \in V$ . Then  $u^* = [u_i^*]_{i \in V}$  satisfies the following.

(a)  $u_i^*(t)$  is 1-Lipschitz for all  $i \in V$ . As a consequence, for all  $i \in V$ ,  $u_i^*(t)$  is differentiable at a.e.  $t \in [0, \delta]$ .

(b) For all  $\Delta ijk \in F$ ,  $\theta^i_{jk}(u^*(t)) \leq \frac{\pi}{2} - \frac{\epsilon}{2}$ . As a consequence  $\theta^i_{jk}(u^*(t)) \geq \epsilon$  for all  $\Delta ijk \in F$  and  $\eta_{ij}(u^*(t)) \leq 2 \cot \epsilon$  for all  $ij \in E$ .

(c) For all  $i \in V$ ,  $K_i(u^*(t)) = 0$ . As a consequence for a.e.  $t \in [0, \delta]$ ,

$$0 = \frac{d}{dt} K_i(u^*(t)) = \sum_{j:ij \in E} \eta_{ij}(u^*(t))(\dot{u}_i^*(t) - \dot{u}_j^*(t)),$$

for all  $i \in V$ .

(d) By Theorem 2.4,  $\dot{u}^*(t)$  is constant on V for a.e.  $t \in [0, \delta]$ . As a consequence  $u_i^*(\delta)$  equals to a constant c independent on  $i \in V$ .

(f) By equation (3.2),

$$|\bar{u}_i - c| = |\bar{u}_i - u_i^*(\delta)| \le |\bar{u}|_{\infty} - \delta$$

for all  $i \in V$ . As a consequence we get the following contradiction

$$2|\bar{u}|_{\infty} = |\sup_{i \in V} \bar{u}_i - \inf_{i \in V} \bar{u}_i| \le |\sup_{i \in V} \bar{u}_i - c| + |\inf_{i \in V} \bar{u}_i - c| \le 2|\bar{u}|_{\infty} - 2\delta$$

3.2. Boundedness of the Conformal Factor. Without loss of generality, we may assume that  $\psi \circ \phi^{-1}$  is linear on each triangle  $\phi(\triangle ijk)$ . Then  $\psi \circ \phi^{-1}$  is K-quasiconformal for some constant  $K = K(\epsilon) > 0$ . We will prove the boundedness of  $\bar{u}$  by showing that for any  $j, j' \in V$ ,

$$|\bar{u}_j - \bar{u}_{j'}| \le 2M + 2\log C + \log C' - \log 2,$$

where  $M = M(\epsilon)$  is the constant given in Lemma 2.9 and  $C = C(\epsilon)$  is the constant given in Lemma 4.3 and  $C' = C'(\epsilon) = e^{200\pi K}$ .

Assume  $j, j' \in V$ . For convenience, let us assume  $\phi(j) = \psi(j) = 0$  by translations. Pick r > 0 sufficiently large such that  $|\phi(j')| < r/(2C)$  and  $\phi(R_j) \subseteq D_r$ . Let  $V_1 = \{i \in V : \phi(i) \in D_r\}$  and  $V_2 = \{i \in V : \phi(i) \in D_{CC'r}\}$  and  $T_1 = T(V_1)$  and  $T_2 = T(V_2)$ . Then by Lemma 4.3 we have

(3.3) 
$$\{\phi(j), \phi(j')\} \subseteq D_{r/(2C)} \subseteq D_{r/C} \subseteq \phi(|T_1|),$$

and

$$\phi(|T_1|) \subseteq D_r \subseteq D_{C'r} \subseteq \phi(|T_2|)$$

and

(3.4) 
$$\phi(|T_2|) \subseteq D_{CC'r}$$

So  $A = A_{r,C'r}$  separates  $\phi(|T_1|)$  and  $\phi(|T_2|)^c$ , and then  $A' = \psi \circ \phi^{-1}(A) \ni \psi(j) = 0$ separates  $\psi(T_1)$  and  $\psi(T_2)^c$ . Furthermore

$$\operatorname{Mod}(A') \ge \frac{1}{K} \cdot \operatorname{Mod}(A) = \frac{1}{K} \cdot \frac{1}{2\pi} \log \frac{C'r}{r} = 100.$$

Then by Lemma 2.1 there exists r' > 0 such that  $A_{r',2r'} \subseteq A'$ . So  $A_{r',2r'}$  separates  $\psi(T_1)$  and  $\psi(T_2)^c$  and then

(3.5) 
$$\psi(|T_1|) \subseteq D_{r'}$$

and

(3.6) 
$$\{\psi(j), \psi(j')\} \subseteq D_{r'} \subseteq D_{2r'} \subseteq \psi(|T_2|).$$

By Lemma 2.9 and equations (3.4) and (3.6), both  $\bar{u}_j, \bar{u}_{j'}$  are at least

$$\log \frac{2r'}{CC'r} - M = \log \frac{r'}{r} + \log \frac{2}{CC'} - M.$$

Again by Lemma 2.9 and equations (3.5) and (3.3), both  $-\bar{u}_j$  and  $-\bar{u}_{j'}$  are at least

$$\log \frac{r/C}{r'} - M = \log \frac{r}{r'} - \log C - M.$$

So both  $\bar{u}_i$  and  $\bar{u}_{i'}$  are in the interval

$$\left[\log\frac{r'}{r} + \log\frac{2}{CC'} - M, \log\frac{r'}{r} + \log C + M\right],$$

and  $|\bar{u}_j - \bar{u}_{j'}|$  is bounded by the length of this interval

$$2M + \log C - \log \frac{2}{CC'} = 2M + 2\log C + \log C' - \log 2.$$

4.1. Electrical Networks and Discrete Extremal Length. Discrete harmonic functions are closely related to the theory of electrical networks. Here the 1-skeleton (V, E) of the triangulation T could be viewed as an electrical network, and  $\eta_{ij}$  denotes the conductance of the edge ij, and the function f denotes the electric potentials at the vertices. Then f is harmonic at i if and only if the outward electric flux at i is 0. The theory of electrical networks is closely related to discrete (edge) extremal length, originally introduced by Duffin [Duf62]. Here we briefly review the theory of discrete (edge) extremal length, adapted to our setting. All the definitions and properties here are well-known and one may read [Duf62][He99] for references.

Assume  $V_1, V_2$  are two nonempty disjoint subsets of V such that  $V_0 = (V_1 \cup V_2)^c$ is finite. A *path* p between  $V_1$  and  $V_2$  is a finite set of edges in

$$E_0 = E_0(V_1, V_2) = \{ ij \in E : i \in V_0 \text{ or } j \in V_0 \}$$

such that  $\gamma_p = \bigcup \{e : e \in p\}$  is a simple curve connecting  $V_1$  and  $V_2$ . Denote  $P = P(V_1, V_2)$  as the set of paths between  $V_1$  and  $V_2$ . A *cut* q between  $V_1$  and  $V_2$  is a finite set of edges in  $E_0$  such that q separates  $V_1$  and  $V_2$ , i.e., for any path  $p \in P$ ,  $p \cap q \neq \emptyset$ . Denote  $Q = Q(V_1, V_2)$  as the set of cuts between  $V_1$  and  $V_2$ .

Given  $\mu \in \mathbb{R}^{E}_{>0}$ , the discrete (edge) extremal length  $EL = EL(V_1, V_2, \mu)$  is defined as

(4.1) 
$$EL = \min\{\sum_{e \in E_0} \mu_e w_e^2 : w \in \mathbb{R}^{E_0}, \sum_{e \in q} w_e \ge 1 \text{ for all } q \in Q\},\$$

and the discrete (edge) extremal width  $EW = EW(V_1, V_2, \mu)$  is defined as

$$EW = \min\{\sum_{e \in E_0} \mu_e w_e^2 : w \in \mathbb{R}^{E_0}, \sum_{e \in p} \mu_e w_e \ge 1 \text{ for all } p \in P\}.$$

Here  $\mu_e$  should be viewed as the resistance of edge  $e \in E$ . Then the conductance of edge  $e \in E$  should be  $\eta_e = 1/\mu_e$ . If  $f: V \to \mathbb{R}$  is harmonic on  $V_0$  with respect to  $\eta$  and  $f|_{V_1} = 0$  and  $f|_{V_2} = 1$ , then  $w_{ij} = |f_j - f_i|/\mu_{ij}$  gives the unique minimizer in the quadratic minimization problem in equation (4.1). If we view such f as an electric potential, then  $w_e$  represents the current on edge  $e \in E_0$  and  $EL = \sum_{e \in E_0} \mu_e w_e^2$  is the electrical power in the network, which is equal to the *(equivalent) resistance* between  $V_1$  and  $V_2$ . The discrete extremal length and width satisfy the following reciprocal theorem.

**Theorem 4.1** (Adapted from Corollary 1 in [Duf62]).  $EL(V_1, V_2, \mu) \cdot EW(V_1, V_2, \mu) = 1.$ 

Now assume  $\emptyset \neq V_0, V_1, V_2...$  is an increase sequence of subsets of V and  $\bigcup_{k=0}^{\infty} V_k = V$ . Then the electric network  $(T, \mu)$  is called *recurrent* if

$$EL(V_0, V_n^c, \mu) \to \infty$$

as  $n \to \infty$ . The recurrency of the network does not depend on the choice of  $V_n$ 's. Intuitively the recurrency means that the equivalent resistance between a finite set and the infinity is infinite. Discrete extremal length is a useful tool to prove the discrete Liouville theorem, since the recurrency implies the discrete Liouville property. **Lemma 4.2** (Lemma 5.5 in [He99]). Assume  $(T, \mu)$  is recurrent, and let  $\eta_e = 1/\mu_e$ for all  $e \in E$ . Then any bounded harmonic function on  $(T, \eta)$  is bounded.

4.2. Proof of the Discrete Liouville Theorem. We need the following lemma for the proof.

**Lemma 4.3.** Suppose  $\phi: |T| \to \mathbb{R}^2$  is a geodesic homeomorphism and any inner angle in  $l(\phi)$  is at least  $\epsilon > 0$ . Let  $a \in V$  be a vertex and assume  $\phi(a) = 0$ . Given r > 0, denote  $V_r = \{i \in V : |\phi(i)| < r\}$  and  $T_r = T(V_r)$ . Then there exists a constant  $C = C(\epsilon) > 0$  such that if  $\phi(R_a) \subseteq D_r$ ,

- (a)  $D_{r/C} \subseteq \phi(|T_r|)$ , and (b) as a consequence  $|\phi(i)| \ge r/C$  for all  $i \in \partial V_r$ .

*Proof.* By a standard compactness argument, it is not difficult to show that there exists a constant  $\delta = \delta(\epsilon) > 0$  such that for all  $\triangle ijk \in F$ ,

 $d(U_{ijk}^c, \phi(\triangle ijk)) \ge \delta \cdot \operatorname{diam}(\phi(\triangle ijk))$ 

where

$$U_{ijk} = int(\phi(R_i)) \cup int(\phi(R_j)) \cup int(\phi(R_k)) \supseteq \phi(\triangle ijk)$$

We claim that  $C = 1 + 2/\delta$  is a desired constant. Let us prove by contradiction. Suppose  $r > \max\{|\phi(i)| : ai \in E\}$  and  $D_{r/C} \not\subseteq \phi(|T_r|)$ . Then there exists  $z \in$  $D_{r/C} \setminus \phi(|T_r|)$ . Since  $\phi$  is a geodesic homeomorphism, there exists a triangle  $\triangle ijk \in$ F such that  $z \in \phi(\triangle ijk)$ . Then  $\triangle ijk$  is not a triangle in  $T_r$  and we may assume  $i \notin V_r$ . So  $|\phi(i)| \ge r$  and  $ai \notin E$  and  $0 = \phi(a) \notin U_{ijk}$ . Then

$$\begin{aligned} r/C &\ge |0-z| \ge d(U_{ijk}^c, \phi(\triangle ijk)) \ge \delta \cdot \operatorname{diam}(\phi(\triangle ijk)) \\ &\ge \delta \cdot |\phi(i)-z| \ge \delta \cdot (r-r/C) = (r/C) \cdot \delta(C-1) = 2r/C \\ &\text{a contradiction.} \end{aligned}$$

and we get a contradiction.

Proof of Theorem 2.4. By replacing  $\eta$  by  $\eta/|\eta|_{\infty}$  we may assume that  $|\eta|_{\infty} = 1$ . Assume  $\mu \in \mathbb{R}^E$  is defined as  $\mu_e = 1/\eta_e \ge 1$  for all  $e \in E$ . Then by Lemma 4.2 we only need to show that  $(T, \mu)$  is recurrent. Let  $\mathbf{1} = (1, 1, ..., 1) \in \mathbb{R}^{E}$ . Then by the definition (equation (4.1))  $EL(V_1, V_2, \mu) \ge EL(V_1, V_2, \mathbf{1})$  whenever well-defined. So we only need to show that (T, 1) is recurrent.

Suppose  $a \in V$  is a vertex and without loss of generality we may assume that  $\phi(a) = 0$ . Let  $\epsilon > 0$  be the infimum of the inner angles in the PL metric  $l(\phi)$ , and  $C = C(\epsilon) > 1$  be the constant given in Lemma 4.3.

Let  $r_0 = \max\{|\phi(i)| : ai \in E\}$  and  $r_n = (2C)^n r_0$  and  $V_n = \{i \in V : \phi(i) \in D_{r_n}\}$ for all  $n \in \mathbb{Z}_{\geq 0}$ . Clearly  $V_n$  is an increasing sequence of subsets of V and  $\bigcup_{n=1}^{\infty} V_n =$ V. We will prove the recurrency of  $(T, \mathbf{1})$  by showing that  $EL(V_0, V_n^c, \mathbf{1}) \to \infty$  as  $n \to \infty$ .

By Lemma 4.3 (b),  $|\phi(i)| \ge r_n/C = 2r_{n-1}$  if  $i \in \partial V_n \cup V_n^c = \overline{V_n^c}$ . So  $V_{n-1} \cap \overline{V_n^c} =$  $\emptyset$ , i.e.,  $V_{n-1} \subseteq (\overline{V_n^c})^c = int(V_n)$ . It is easy to see

(4.2) 
$$E_0(V_{n-1}, \overline{V_n^c}) \subseteq E(V_n) \setminus E(V_{n-1}).$$

From the definition of extremal length, we have

$$EL(V_0, V_n^c) \ge EL(V_0, \overline{V_1^c}) + EL(V_1, \overline{V_2^c}) + \dots + EL(V_{n-1}, \overline{V_n^c})$$

since

(1)  $E_0(V_0, \overline{V_1^c}), E_0(V_1, \overline{V_2^c}), \dots, E_0(V_{n-1}, \overline{V_n^c})$  are disjoint by equation (4.2), and (2)  $Q(V_0, \overline{V_1^c}), Q(V_1, \overline{V_2^c}), \dots, Q(V_{n-1}, \overline{V_n^c})$  are all subsets of  $Q(V_0, V_n^c)$ .

So it suffices to show that for all n,

$$EL(V_{n-1}, \overline{V_n^c}, \mathbf{1}) \ge \frac{\sin^2 \epsilon}{12\pi C^2},$$

which by Theorem 4.1 is equivalent to

$$EW(V_{n-1}, \overline{V_n^c}, \mathbf{1}) \le \frac{12\pi C^2}{\sin^2 \epsilon}.$$

In the remaining of the proof we denote  $E_0 = E_0(V_{n-1}, \overline{V_n^c})$ . Pick  $w_e = l_e/r_{n-1}$ , and then for any  $p \in P = P(V_{n-1}, \overline{V_n^c})$ ,

$$\sum_{e \in P} w_e = \frac{1}{r_{n-1}} \sum_{e \in P} l_e \ge \frac{1}{r_{n-1}} \cdot d(\phi(V_{n-1}), \phi(\overline{V_n^c})) \ge \frac{1}{r_{n-1}} \cdot (2r_{n-1} - r_{n-1}) = 1.$$

 $\operatorname{So}$ 

$$EW(V_{n-1}, \overline{V_n^c}, \mathbf{1}) \le \sum_{e \in E_0} w_e^2 = \frac{1}{r_{n-1}^2} \sum_{e \in E_0} l_e^2$$

and it remains to show

$$\sum_{e \in E_0} l_e^2 \le \frac{12\pi C^2}{\sin^2 \epsilon} \cdot r_{n-1}^2$$

Given  $e \in E$ , denote  $\triangle_e, \triangle'_e$  as the two triangles in T containing e. If  $e \in E_0$ , then e contains at least 1 vertex in  $(\overline{V_n^c})^c = int(V_n)$  and  $\triangle_e, \Delta'_e$  are both triangles in  $T_n$ , i.e.,  $\phi(\triangle_e), \phi(\triangle'_e)$  are both in  $D_{r_n}$ . Given a triangle  $\triangle \in F$ , we denote  $|\triangle|$  as the area of  $\phi(\triangle)$ . Then by the sine law

$$|\triangle ijk| = \frac{1}{2} l_{ij} l_{jk} \sin \theta_{ik}^j \ge \frac{1}{2} l_{ij}^2 \cdot \frac{\sin \theta_{jk}^i}{\sin \theta_{ij}^k} \cdot \sin \theta_{ik}^j \ge l_{ij}^2 \cdot \frac{\sin^2 \epsilon}{2}.$$

Notice that a triangle  $\triangle \in F$  is counted for at most 3 times in  $\sum_{e \in E_0} (|\triangle_e| + |\triangle'_e|)$  and then

$$\sum_{e \in E_0} l_e^2 \le \frac{1}{\sin^2 \epsilon} \sum_{e \in E_0} (|\triangle_e| + |\triangle'_e|) \le \frac{1}{\sin^2 \epsilon} \sum_{\triangle: \phi(\triangle) \subseteq D_{r_n}} 3|\triangle| = \frac{3\pi r_n^2}{\sin^2 \epsilon} = \frac{12\pi C^2}{\sin^2 \epsilon} \cdot r_{n-1}^2.$$

### 5. Hyperbolic Maximum Principles and Proof of Lemma 2.9

Given  $z_1, z_2 \in D$ , we denote  $d_h(z_1, z_2)$  as the hyperbolic distance between  $z_1, z_2$  in the Poincaré disk model. The (Euclidean) discrete conformal change is related with the hyperbolic discrete conformal change as follows.

**Lemma 5.1.** Suppose  $z_1, z_2, z'_1, z'_2 \in D$  and  $u_1, u_2, u^h_1, u^h_2 \in \mathbb{R}$  are such that

$$u_i^h = u_i + \log \frac{1 - |z_i|^2}{1 - |z_i'|^2}$$

for i = 1, 2. Then

$$|z_1' - z_2'| = e^{\frac{1}{2}(u_1 + u_2)} |z_1 - z_2|$$

if and only if

(5.1) 
$$\sinh \frac{d_h(z'_i, z'_j)}{2} = e^{\frac{1}{2}(u^h_i + u^h_j)} \sinh \frac{d_h(z_i, z_j)}{2}$$

**Remark 5.2.** Equation (5.1) is indeed the formula of the discrete conformal change for piecewise hyperbolic metric. This formula was first proposed by Bobenko-Pinkall-Springborn [BPS15], and  $u_i^h$  in the formula is called the hyperbolic discrete conformal factor at i.

Lemma 5.1 could be verified by elementary computations. The proof is given in Appendix. The hyperbolic discrete conformal factor  $u^h$  also satisfies a maximum principle.

**Lemma 5.3.** Suppose  $V_0$  is a subset of V and  $u \in \mathbb{R}^{V_0}$  and  $\phi, \psi$  are Euclidean geodesic embeddings of  $T(V_0)$ , such that  $\phi(|T(V_0)|), \psi(|T(V_0)|) \subseteq D$  and  $l(\phi), l(\psi)$  are both uniformly acute and  $l(\psi) = u * l(\phi)$ . For all  $i \in V_0$ , denote  $z_i = \phi(i)$  and  $z'_i = \psi(i)$  and

$$u_i^h = u_i + \log \frac{1 - |z_i|^2}{1 - |z_i'|^2}.$$

(a) If  $i \in int(V_0)$  and  $u_i^h < 0$ , then there exists a neighbor j of i such that

$$u_j^h < u_i^h.$$

(b) If  $u_i^h \ge 0$  for all  $i \in \partial V_0$ , then  $u_i^h \ge 0$  for all  $i \in V_0$ .

We first prove Lemma 2.9 using the hyperbolic maximum princple and then prove Lemma 5.3.

Proof of Lemma 2.9. For any  $\triangle ijk \in F$ ,

$$e^{\frac{1}{2}(u_j - u_i)} = \frac{e^{(u_j + u_k)/2}}{e^{(u_i + u_k)/2}} = \frac{l_{jk}(\psi)/l_{jk}(\phi)}{l_{ik}(\psi)/l_{ik}(\phi)} = \frac{l_{jk}(\psi)}{l_{ik}(\psi)} \cdot \frac{l_{ik}(\phi)}{l_{jk}(\phi)} \ge \sin^2 \epsilon.$$

So there exists a constant  $C = C(\epsilon) > 0$  such that  $|u_j - u_i| \leq 2C$  for all  $ij \in E$ . We will show that  $M(\epsilon) = C(\epsilon) + 3$  is a satisfactory constant. By a scaling, we only need to prove for the special case where r' = 1 and  $r = e^{-C-2}$ .

Denote  $V_1 = \{i \in V : \psi(i) \in D\}$  and  $z_i = \phi(i)$  and  $z'_i = \psi(i)$ . Define  $u^h \in \mathbb{R}^{V_1}$  as

$$u_i^h = u_i + \log \frac{1 - |z_i|^2}{1 - |z_i'|^2}$$

for all  $i \in V_1$ . Assume  $i \in \partial V_1$ , then there exists  $j \in V_0 - V_1$  such that  $ij \in E$ . We claim that  $u_i^h \ge 0$ , i.e.,

$$e^{u_i} \cdot \frac{1 - |z_i|^2}{1 - |z_i'|^2} \ge 1.$$

Notice that

$$1 - |z_i'| \le |z_i' - z_j'| = e^{\frac{1}{2}(u_i + u_j)} |z_i - z_j| \le e^{u_i + C} \cdot 2r = 2e^{-2}e^{u_i}.$$

So

$$e^{u_i} \cdot \frac{1 - |z_i|^2}{1 - |z_i'|^2} \ge \frac{e^2}{2} \cdot \frac{1 - |z_i|^2}{1 + |z_i'|} \ge \frac{e^2}{2} \cdot \frac{1 - r^2}{2} \ge \frac{e^2}{2} \cdot \frac{1 - (e^{-2})^2}{2} > 1$$

By the hyperbolic maximum principle Lemma 5.3 (b),  $u_i^h \ge 0$  for all  $i \in V_1$ . Then for all  $i \in V_0$  with  $|z_i'| < 1/2$ ,

$$u_i = u_i^h - \log \frac{1 - |z_i|^2}{1 - |z_i'|^2} \ge -\log \frac{1 - |z_i|^2}{1 - |z_i'|^2} \ge \log(1 - |z_i'|^2) \ge -1 = \log(r'/r) - M.$$

12

5.1. **Proof of the Hyperbolic Maximum Principle.** For the proof of Lemma 5.3, we need to briefly review the notion of hyperbolic Delaunay. Given a subcomplex  $T_0$  of T, an embedding  $\phi : |T_0| \to D$  is called a *hyperbolic geodesic embedding* if  $\phi_h$  maps each edge of  $T_1$  to a hyperbolic geodesic arc in  $(D, d_h)$ . Given a triangle  $\triangle ijk$  in  $T_0$  and a Euclidean or hyperbolic geodesic embedding  $\phi$  of  $T_0$ , denote  $C_{ijk} = C_{ijk}(\phi)$  as the circumcircle of  $\phi(\triangle ijk)$ , i.e., a round circle in the Riemann sphere  $\hat{\mathbb{C}}$  passing through the three vertices of  $\phi(\triangle ijk)$ . Furthermore, we denote  $D_{ijk} = D_{ijk}(\phi)$  as the circumdisk of  $\phi(\triangle ijk)$ , i.e., the closed round disk in  $\hat{\mathbb{C}}$  such that  $\partial D_{ijk} = C_{ijk}$  and  $\phi(\triangle ijk) \subseteq D_{ijk}$ . For a Euclidean geodesic embedding  $\phi$  of  $T_0$ , it is well-known that  $l(\phi)$  is Delaunay if and only if that for any pair of adjacent triangles  $\triangle ijk, \triangle ijk'$  in  $T_0$ ,

$$\phi(k') \notin int(D_{ijk}).$$

So here we naturally call a Euclidean or hyperbolic geodesic embedding  $\phi~Delaunay$  if

$$\phi(k') \notin int(D_{ijk})$$

for any pair of adjacent triangles  $\triangle ijk$ ,  $\triangle ijk'$  in  $T_0$ .

Proof of Lemma 5.3 (a). Assume  $i \in int(V_0)$  and  $T_1 = (V_1, E_1, F_1)$  is the 1-ring neighborhood of i. Then by Lemma 5.4 below there exists a hyperbolic Delaunay geodesic embedding  $\phi_h$  (resp.  $\psi_h$ ) of  $T_1$  such that  $\phi_h(j) = z_j$  (resp.  $\psi_h(j) = z'_j$ ) for all  $j \in V_1$ . By Lemma 5.1,

$$\sinh \frac{d_h(z'_j, z'_k)}{2} = e^{\frac{1}{2}(u^h_j + u^h_k)} \sinh \frac{d_h(z_j, z_k)}{2}$$

for all  $jk \in E_1$ . Suppose  $f_1, f_2 : D \to D$  are hyperbolic isometries such that  $f_1(z_i) = 0$  and  $f_2(z'_i) = 0$ . Then  $\tilde{\phi}_h = f_1 \circ \phi_h$  (resp.  $\tilde{\psi}_h = f_2 \circ \psi_h$ ) is a hyperbolic Delaunay geodesic embedding. Denote  $\tilde{z}_j = \tilde{\phi}_h(j)$  (resp.  $\tilde{z}'_j = \tilde{\psi}_h(j)$ ) for all  $j \in V_1$ . Then  $z_i = z'_i = 0$  and

$$\sinh \frac{d_h(\tilde{z}'_j, \tilde{z}'_k)}{2} = e^{\frac{1}{2}(u^h_j + u^h_k)} \sinh \frac{d_h(\tilde{z}_j, \tilde{z}_k)}{2}$$

for all  $jk \in E_1$ . It is not hard to see that there exists a Euclidean Delaunay geodesic embedding  $\tilde{\phi}$  (resp.  $\tilde{\psi}$ ) of  $T_1$  such that  $\tilde{\phi}(j) = \tilde{z}_j$  (resp.  $\tilde{\psi}(j) = \tilde{z}'_j$ ). By Lemma 5.1  $l(\tilde{\psi}) = \tilde{u} * l(\tilde{\phi})$  where

$$\tilde{u}_j = u_j^h - \log \frac{1 - |\tilde{z}_j|^2}{1 - |\tilde{z}'_j|^2}.$$

By the Euclidean maximum principle Lemma 2.7,  $\tilde{u}_j \leq \tilde{u}_i < 0$  for some neighbor j of i. Then

$$|\tilde{z}'_j| = l_{ij}(\tilde{\psi}) = e^{\frac{1}{2}(\tilde{u}_i + \tilde{u}_j)} l_{ij}(\tilde{\phi}) = e^{\frac{1}{2}(\tilde{u}_i + \tilde{u}_j)} |\tilde{z}_j| < |\tilde{z}_j|$$

and

$$u_j^h = \tilde{u}_j + \log \frac{1 - |\tilde{z}_j|^2}{1 - |\tilde{z}_j'|^2} < \tilde{u}_j \le \tilde{u}_i = u_i^h - \frac{1 - |\tilde{z}_i|^2}{1 - |\tilde{z}_i'|^2} = u_i^h.$$

Proof of Lemma 5.3 (b). If not, assume  $u_i^h = \min_{j:j \in V_0} u_j^h < 0$  and then  $i \in int(V_0)$ . By the minimality of  $u_i^h$ ,  $u_j^h \ge u_i^h$  for any neighbor j of i. This contradicts with part (a).

**Lemma 5.4.** Suppose  $i \in V$  and  $T_1 = (V_1, E_1, F_1)$  is a 1-ring neighborhood of i. If  $\phi$  is a geodesic embedding of  $T_1$  such that  $\phi(|T_1|) \subseteq D$  and  $l(\phi)$  is uniformly acute, then there exists a hyperbolic geodesic embedding  $\phi_h$  of  $T_1$  such that  $\phi_h(j) = \phi(j)$ for all  $j \in V_1$ . Furthermore, such  $\phi_h$  is Delaunay.

*Proof.* Let  $j_1, j_2, ..., j_m$  be the neighbors of *i* listed counterclockwise in  $\phi(|T_1|)$ . Denote  $z_0 = \phi(i)$  and  $z_k = \phi(j_k)$  for k = 1, ..., m. If  $\gamma(t) : [0, 1] \to D$  is a smooth curve such that  $\gamma(0) = z_0$ , then  $\dot{\gamma}(t)$  could be viewed as not only a complex number but also a vector in the tangent space  $T_{z_0}D$  of  $(D, d_h)$  at  $z_0$ . By this way we naturally identify  $T_{z_0}D$  with  $\mathbb{C}$ .

Given  $z \in D$ , let  $v(z) = \exp_{z_0}^{-1} z \in T_{z_0} D = \mathbb{C}$  where  $\exp_{z_0} : T_{z_0} D \to D$  is the exponential map at  $z_0$  on the hyperbolic plane D. We first show that  $v(z_1), ..., v(z_m)$ are counterclockwise around 0 and wrap around 0 once. More specifically, we will show that

(5.2) 
$$\arg\left(\frac{v(z_{k+1})}{v(z_k)}\right) \in (0,\pi)$$

and

(5.3) 
$$\sum_{k=1}^{m} \arg\left(\frac{v(z_{k+1})}{v(z_k)}\right) = 2\pi$$

where  $z_{m+1} = z_1$  and  $\arg(z)$  denotes the argument of z.

Assume  $k \in \{1, ..., m\}$ . Denote  $\gamma$  (resp.  $\gamma_h$ ) as the Euclidean straight line in  $\mathbb{C}$ (hyperbolic geodesic in D) containing  $z_0, z_k$ . Then  $\gamma$  (resp.  $\gamma_h$ ) cuts  $\mathbb{C}$  (resp. D) into two open subsets P, P' (resp.  $P_h, P'_h$ ). We may assume

$$P = \{z \in \mathbb{C} : \arg\left(\frac{z - z_0}{z_k - z_0}\right) \in (0, \pi)\}$$

and

$$P_h = \{z \in D : \arg\left(\frac{v(z)}{v(z_k)}\right) \in (0,\pi)\}.$$

Then  $z_{k+1} \in P$ . If  $\gamma_h$  is a straight line,  $P_h = P \ni z_{k+1}$  and we have proved equation (5.2). If  $\gamma_h$  is a round circular arc orthogonal to  $\{|z|=1\}$ , there are two different cases.

Case 1: assume  $z_0, z_k$  are counterclockwise on  $\gamma_h$  (see Figure 1 (A)). If  $z_{k+1} \in$  $P \setminus P_h$ ,  $\angle z_0 z_k z_{k+1} > \pi/2$  or  $\angle z_k z_0 z_{k+1} > \pi/2$  and it is contradictory to the acuteness assumption. So  $z_{k+1} \in P_h$ .

Case 2: assume  $z_0, z_k$  are clockwise on  $\gamma_h$  (see Figure 1 (B)). If  $z_{k+1} \in P \setminus P_h$ ,  $\angle z_0 z_{k+1} z_k > \pi/2$  and it is contradictory to the acuteness assumption. So  $z_{k+1} \in$  $P_h$ .

So we proved equation (5.2) and now prove equation (5.3). It is easy to see that

$$\operatorname{arg}\left(\frac{v(z_k)}{z_k-z_0}\right) \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right).$$

for all k = 1, ..., m. We claim that

(5.4) 
$$\arg\left(\frac{v(z_{k+1})}{v(z_k)}\right) + \arg\left(\frac{v(z_k)}{z_k - z_0}\right) = \arg\left(\frac{z_{k+1} - z_0}{z_k - z_0}\right) + \arg\left(\frac{v(z_{k+1})}{z_{k+1} - z_0}\right).$$
Since

ç

$$\exp(\sqrt{-1} \cdot LHS) = \exp(\sqrt{-1} \cdot RHS) = \frac{v(z_{k+1})}{z_k - z_0},$$

RIGIDITY OF ACUTE TRIANGULATIONS OF THE PLANE



Figure 1

we have that

$$LHS = RHS + 2n\tau$$

for some integer n. On the other hand LHS and RHS are both bounded in

$$(0-\frac{\pi}{2},\pi+\frac{\pi}{2})=(-\frac{\pi}{2},\frac{3\pi}{2})$$

so LHS = RHS. Now by adding up equation (5.4) for k = 1, ..., m we have that

$$\sum_{k=1}^{m} \arg\left(\frac{v(z_{k+1})}{v(z_k)}\right) = \sum_{k=1}^{m} \arg\left(\frac{z_{k+1} - z_0}{z_k - z_0}\right) = 2\pi$$

since  $\phi$  is a geodesic embedding. So we proved equations (5.2) and (5.3), and as a consequence there exists a hyperbolic embedding  $\phi_h$  of  $|T_1|$  such that  $\phi_h(j) = z_j$  for all  $j \in V_1$ .

By equation (5.2) it is not difficult to see that the two circumdisks  $D_{ij_kj_{k+1}}(\phi)$ and  $D_{ij_kj_{k+1}}(\phi_h)$  are the same for k = 1, ..., m. So  $\phi_h$  is Delaunay since  $\phi$  is Delaunay.

### Appendix A. Proof of Lemma 5.1

Proof of Lemma 5.1. It suffices to show that for all  $z_1, z_2 \in D$ ,

$$\sinh \frac{d_h(z_1, z_2)}{2} = \frac{|z_1 - z_2|}{\sqrt{(1 - |z_1|^2)(1 - |z_2|^2)}}.$$

We first consider a special case where  $z_1 = 0$  and  $z_2 = r \in (0, 1)$  is real. Then

$$d_h(z_1, z_2) = \ln \frac{1 \cdot (1+r)}{1 \cdot (1-r)} = \ln \frac{1+r}{1-r}$$

and

$$\sinh \frac{d_h(z_1, z_2)}{2} = \frac{1}{2}\sqrt{\frac{1+r}{1-r}} - \frac{1}{2}\sqrt{\frac{1-r}{1+r}} = \frac{r}{\sqrt{1-r^2}} = \frac{|z_1 - z_2|}{\sqrt{(1-|z_1|^2)(1-|z_2|^2)}}.$$

For general  $z_1, z_2 \in D$ , we can find a hyperbolic isometric map  $f(z) = \frac{z-a}{1-\bar{a}z}$  such that  $f(z_1) = 0$  and  $f(z_2)$  is a positive real number. We only need to verify that

$$\frac{|f(z_1) - f(z_2)|^2}{(1 - |f(z_1)|^2)(1 - |f(z_2)|^2)} = \frac{|z_1 - z_2|^2}{(1 - |z_1|^2)(1 - |z_2|^2)}$$

This equality can be derived from

$$\frac{|f(z_1) - f(z_2)|^2}{(1 - |f(z_1)|^2)(1 - |f(z_2)|^2)} = \frac{|(z_1 - a)(1 - \bar{a}z_2) - (1 - \bar{a}z_1)(z_2 - a)|^2}{(|1 - \bar{a}z_1|^2 - |z_1 - a|^2) \cdot (|1 - \bar{a}z_2|^2 - |z_2 - a|^2)} = \frac{|(1 - a\bar{a})(z_1 - z_2)|^2}{(|1 - \bar{a}z_1|^2 - |z_1 - a|^2) \cdot (|1 - \bar{a}z_2|^2 - |z_2 - a|^2)}$$

and

$$|1 - \bar{a}z_1|^2 - |z_1 - a|^2 = (1 - \bar{a}z_1)(1 - a\bar{z}_1) - (z_1 - a)(\bar{z}_1 - \bar{a})$$
  
=  $(1 - a\bar{a})(1 - z_1\bar{z}_1) = (1 - a\bar{a})(1 - |z_1|^2).$ 

and similarly

$$|1 - \bar{a}z_2|^2 - |z_2 - a|^2 = (1 - a\bar{a})(1 - |z_2|^2).$$

### References

- [Ahl10] Lars Valerian Ahlfors. Conformal invariants: topics in geometric function theory, volume 371. American Mathematical Soc., 2010.
- [BPS15] Alexander I Bobenko, Ulrich Pinkall, and Boris A Springborn. Discrete conformal maps and ideal hyperbolic polyhedra. Geometry & Topology, 19(4):2155–2215, 2015.
- [CCS<sup>+</sup>21] Marcel Campen, Ryan Capouellez, Hanxiao Shen, Leyi Zhu, Daniele Panozzo, and Denis Zorin. Efficient and robust discrete conformal equivalence with boundary. ACM Transactions on Graphics (TOG), 40(6):1–16, 2021.
- [DGM22] Song Dai, Huabin Ge, and Shiguang Ma. Rigidity of the hexagonal delaunay triangulated plane. *Peking Mathematical Journal*, 5(1):1–20, 2022.
- [Duf62] RJ Duffin. The extremal length of a network. Journal of Mathematical Analysis and Applications, 5(2):200–215, 1962.
- [FLZ20] Ke Feng, Aijin Lin, and Xiaoxiao Zhang. Combinatorial p-th calabi flows for discrete conformal factors on surfaces. *The Journal of Geometric Analysis*, 30(4):3979–3994, 2020.
- [GGL<sup>+</sup>18] Xianfeng Gu, Ren Guo, Feng Luo, Jian Sun, Tianqi Wu, et al. A discrete uniformization theorem for polyhedral surfaces ii. Journal of differential geometry, 109(3):431– 466, 2018.
- [GH18] Huabin Ge and Bobo Hua. On combinatorial calabi flow with hyperbolic circle patterns. Advances in Mathematics, 333:523–538, 2018.
- [GLSW18] Xianfeng David Gu, Feng Luo, Jian Sun, and Tianqi Wu. A discrete uniformization theorem for polyhedral surfaces. Journal of differential geometry, 109(2):223–256, 2018.
- [GLW19] David Gu, Feng Luo, and Tianqi Wu. Convergence of discrete conformal geometry and computation of uniformization maps. Asian Journal of Mathematics, 23(1):21– 34, 2019.
- [GSC21] Mark Gillespie, Boris Springborn, and Keenan Crane. Discrete conformal equivalence of polyhedral surfaces. ACM Transactions on Graphics (TOG), 40(4):1–20, 2021.
- [He99] Zheng-Xu He. Rigidity of infinite disk patterns. Annals of Mathematics, pages 1–33, 1999.
- [LSW20] Feng Luo, Jian Sun, and Tianqi Wu. Discrete conformal geometry of polyhedral surfaces and its convergence. arXiv preprint arXiv:2009.12706, 2020.

16

- [Lu004] Feng Luo. Combinatorial yamabe flow on surfaces. Communications in Contemporary Mathematics, 6(05):765–780, 2004.
- [Luo22] Yanwen Luo. Spaces of geodesic triangulations of surfaces. Discrete & Computational Geometry, pages 1–19, 2022.
- [LV73] Olli Lehto and Kaarlo Ilmari Virtanen. Quasiconformal mappings in the plane, volume 126. Citeseer, 1973.
- [LW19] Feng Luo and Tianqi Wu. Koebe conjecture and the weyl problem for convex surfaces in hyperbolic 3-space. arXiv preprint arXiv:1910.08001, 2019.
- [LWZ21a] Yanwen Luo, Tianqi Wu, and Xiaoping Zhu. The convergence of discrete uniformizations for genus zero surfaces. arXiv preprint arXiv:2110.08208, 2021.
- [LWZ21b] Yanwen Luo, Tianqi Wu, and Xiaoping Zhu. The deformation space of geodesic triangulations and generalized tutte's embedding theorem. arXiv preprint arXiv:2105.00612, 2021.
- [LWZ21c] Yanwen Luo, Tianqi Wu, and Xiaoping Zhu. The deformation spaces of geodesic triangulations of flat tori. arXiv preprint arXiv:2107.05159, 2021.
- [LWZ22] Yanwen Luo, Tianqi Wu, and Xiaoping Zhu. The deformation space of delaunay triangulations of the sphere. arXiv preprint arXiv:2202.06402, 2022.
- [Spr19] Boris Springborn. Ideal hyperbolic polyhedra and discrete uniformization. Discrete & Computational Geometry, pages 1–46, 2019.
- [SWGL15] Jian Sun, Tianqi Wu, Xianfeng Gu, and Feng Luo. Discrete conformal deformation: algorithm and experiments. SIAM Journal on Imaging Sciences, 8(3):1421–1456, 2015.
- [WGS15] Tianqi Wu, Xianfeng Gu, and Jian Sun. Rigidity of infinite hexagonal triangulation of the plane. Transactions of the American Mathematical Society, 367(9):6539–6555, 2015.
- [Wu14] Tianqi Wu. Finiteness of switches in discrete Yamabe flow. PhD thesis, Master Thesis, Tsinghua University, Beijing, 2014.
- [WX21] Tianqi Wu and Xu Xu. Fractional combinatorial calabi flow on surfaces. *arXiv preprint* arXiv:2107.14102, 2021.
- [WZ20] Tianqi Wu and Xiaoping Zhu. The convergence of discrete uniformizations for closed surfaces, 2020.
- [ZGZ<sup>+</sup>14] Min Zhang, Ren Guo, Wei Zeng, Feng Luo, Shing-Tung Yau, and Xianfeng Gu. The unified discrete surface ricci flow. *Graphical Models*, 76(5):321–339, 2014.
- [ZX19] Xiang Zhu and Xu Xu. Combinatorial calabi flow with surgery on surfaces. Calculus of Variations and Partial Differential Equations, 58(6):1–20, 2019.

Department of Mathematics, Clark University, 950 Main St, Worcester, MA 01610, USA

Email address: mike890505@gmail.com