

Non-degenerate Hypergraphs with Exponentially Many Extremal Constructions

József Balogh* Felix Christian Clemen† Haoran Luo‡

Abstract

For every integer $t \geq 0$, denote by F_5^t the hypergraph on vertex set $\{1, 2, \dots, 5 + t\}$ with hyperedges $\{123, 124\} \cup \{34k : 5 \leq k \leq 5 + t\}$. We determine $\text{ex}(n, F_5^t)$ for every $t \geq 0$ and sufficiently large n and characterize the extremal F_5^t -free hypergraphs. In particular, if n satisfies certain divisibility conditions, then the extremal F_5^t -free hypergraphs are exactly the balanced complete tripartite hypergraphs with additional hyperedges inside each of the three parts (V_1, V_2, V_3) in the partition; each part V_i spans a $(|V_i|, 3, 2, t)$ -design. This generalizes earlier work of Frankl and Füredi on the Turán number of $F_5 := F_5^0$.

Our results extend a theory of Erdős and Simonovits about the extremal constructions for certain fixed graphs. In particular, the hypergraphs F_5^{6t} , for $t \geq 1$, are the first examples of hypergraphs with exponentially many extremal constructions and positive Turán density.

1 Introduction

For a family of r -uniform hypergraphs (r -graphs) \mathcal{F} , a hypergraph H is \mathcal{F} -free if H contains no copy of any $F \in \mathcal{F}$ as a subhypergraph. The *Turán number* $\text{ex}(n, \mathcal{F})$ is the maximum possible number of hyperedges in an n -vertex \mathcal{F} -free r -graph. Let $\pi(\mathcal{F}) := \lim_{n \rightarrow \infty} \text{ex}(n, \mathcal{F}) / \binom{n}{r}$ be the *Turán density* of \mathcal{F} . A family \mathcal{F} is *non-degenerate* if $\pi(\mathcal{F}) > 0$. When $\mathcal{F} = \{F\}$, we will simply use F -free, $\text{ex}(n, F)$, and $\pi(F)$, respectively. We say an n -vertex \mathcal{F} -free r -graph H is *extremal* (or an *extremal construction*) for \mathcal{F} if H has $\text{ex}(n, \mathcal{F})$ hyperedges. Determining

*Department of Mathematics, University of Illinois at Urbana-Champaign, Urbana, Illinois 61801, USA. E-mail: jobal@illinois.edu. Research is partially supported by NSF Grant DMS-1764123 and RTG DMS-1937241, Arnold O. Beckman Research Award (UIUC Campus Research Board RB 22000), the Langan Scholar Fund (UIUC), and the Simons Fellowship.

†Department of Mathematics, Karlsruhe Institute of Technology, 76131 Karlsruhe, Germany, Email: felix.clemen@kit.edu. Research was partially performed while the second author was at the University of Illinois at Urbana-Champaign.

‡Department of Mathematics, University of Illinois at Urbana-Champaign, Urbana, Illinois 61801, USA. E-mail: haoranl8@illinois.edu. Research is partially supported by UIUC Campus Research Board RB 22000.

Turán numbers and the corresponding extremal constructions is one of the central problems in Extremal Combinatorics and has received extensive attention.

This problem is reasonably well-understood for graphs (the case where $r = 2$). The Erdős-Stone theorem [7], as pointed out by Erdős and Simonovits [5], determines the Turán density of all graphs. However, for hypergraphs, the Turán problem is notoriously difficult, and there are only very few exact results. The *tetrahedron*, denoted by K_4^3 , is the complete 3-graph on four vertices. It is a famous problem of Turán [23] to determine the Turán density of the tetrahedron. There are exponentially many conjectured extremal constructions by Brown [2], Kostochka [17], Fon-der-Flaass [8], and Frohmader [10], see also [20] by Razborov.

In this paper, we give the first examples of hypergraphs which share this (conjectured) property with the tetrahedron in the following sense: each of them has exponentially many non-isomorphic extremal constructions and positive Turán density. We note that our extremal constructions are close (i.e. $O(n^2)$) to each other in edit-distance, while for the tetrahedron, the (conjectured) extremal constructions differ more from each other (some of them have edit-distance $\Omega(n^3)$). Following Liu, Mubayi [18] and Liu, Mubayi, Reiher [19], Hou, Li, Liu, Mubayi, and Zhang [13] recently constructed finite families of 3-graphs such that each family has $\Omega(n)$ non-isomorphic extremal constructions far (i.e. $\Omega(n^3)$) from each other in edit-distance, which is a new phenomenon compared to graphs. It remains open to find a single hypergraph with $\omega(1)$ extremal hypergraphs far from each other in edit-distance. We reiterate that the tetrahedron is conjectured to be such a hypergraph.

A classical theorem in hypergraph Turán theory is the following result of Bollobás [1]. The *generalized triangle*, denoted by F_5 , is the 3-graph on vertex set $\{1, 2, 3, 4, 5\}$ with hyperedges $\{123, 124, 345\}$, see Figure 1. Let K_4^- be the 3-graph on vertex set $\{1, 2, 3, 4\}$ with hyperedges $\{123, 124, 134\}$, and denote by $S(n)$ the complete tripartite 3-graph on vertex set $[n] := \{1, 2, \dots, n\}$ with parts of sizes as equal as possible, see Figure 2. The number of hyperedges in $S(n)$ is $s(n) := \lfloor n/3 \rfloor \cdot \lfloor (n+1)/3 \rfloor \cdot \lfloor (n+2)/3 \rfloor$. Bollobás [1] proved that $\text{ex}(n, \{K_4^-, F_5\}) = s(n)$. This result was extended by Frankl and Füredi [9], who proved that $\text{ex}(n, F_5) = s(n)$ for $n \geq 3000$, and it was improved again to $n \geq 33$ by Keevash and Mubayi [16].




Figure 1: Hypergraph F_5 .




Figure 2: Hypergraph $S(n)$.

For every integer $t \geq 0$, denote by F_5^t the hypergraph on vertex set $\{1, 2, \dots, 5 + t\}$ with hyperedges $\{123, 124\} \cup \{34k : 5 \leq k \leq 5 + t\}$, see Figure 3. Hence $F_5^0 = F_5$. Furthermore, denote by F_2^t the hypergraph on vertex set $\{1, 2, \dots, 3 + t\}$ with hyperedges $\{12k : 3 \leq k \leq 3 + t\}$. Note that F_2^t -free hypergraphs are exactly the hypergraphs with maximum codegree

at most t , i.e., every pair of vertices is in at most t hyperedges. Let $\mathcal{H}(n, t)$ be the family of extremal n -vertex F_2^t -free 3-graphs $H(n, t)$. Define $\mathcal{S}_t^\circ(n)$ as the family of 3-graphs $S_t^\circ(n)$ on vertex set $[n]$, where $S_t^\circ(n)$ is obtained from $S(n)$ by adding a copy of $H(|V_i|, t) \in \mathcal{H}(|V_i|, t)$ to every part V_i in the partition of $S(n)$, see Figure 4. Let

$$s_t^\circ(n) := \left\lfloor \frac{n}{3} \right\rfloor \left\lfloor \frac{n+1}{3} \right\rfloor \left\lfloor \frac{n+2}{3} \right\rfloor + \text{ex} \left(\left\lfloor \frac{n}{3} \right\rfloor, F_2^t \right) + \text{ex} \left(\left\lfloor \frac{n+1}{3} \right\rfloor, F_2^t \right) + \text{ex} \left(\left\lfloor \frac{n+2}{3} \right\rfloor, F_2^t \right)$$

be the number of hyperedges in every $S_t^\circ(n) \in \mathcal{S}_t^\circ(n)$. Our main result is the following theorem.

Theorem 1.1. *For every integer $t \geq 0$ and sufficiently large n , the F_5^t -free 3-graphs on vertex set $[n]$ with maximum number of hyperedges are exactly the hypergraphs $S_t^\circ(n) \in \mathcal{S}_t^\circ(n)$, and hence $\text{ex}(n, F_5^t) = s_t^\circ(n)$.*

Thus, we solve the Turán problem for each hypergraph in this infinite family of forbidden hypergraphs for sufficiently large n and characterize their extremal constructions. Our extremal constructions $\mathcal{S}_t^\circ(n)$ are similar in soul to the following result of Simonovits [22]. For $r, d \geq 2$, let $Q(r, d)$ be the complete $(d+1)$ -partite graph on $rd+1$ vertices with one part of size 1 and d parts of size r . Let $\mathcal{U}(n, r, d)$ be the family of graphs, each of which is obtained from the n -vertex complete d -partite graph with parts of sizes as equal as possible by adding maximum number of edges inside each part such that every vertex is adjacent to at most $r-1$ other vertices in the same part. Let $\mathcal{U}^*(n, r, d)$ be the family of graphs in $\mathcal{U}(n, r, d)$ with the extra requirement that there is no triangle inside any part. Simonovits [22], generalizing an unpublished result of Erdős, proved that all the graphs in $\mathcal{U}^*(n, r, d)$ are extremal constructions for $Q(r, d)$ and all the extremal constructions for $Q(r, d)$ are in $\mathcal{U}(n, r, d)$, for sufficiently large n .

For our proof of Theorem 1.1, we derive a stability result, followed up by a cleaning method, which reveals the structure of the extremal F_5^t -free hypergraphs step by step. One




Figure 3: Hypergraph F_5^t .




Figure 4: Hypergraph $S_t^\circ(n)$.

of the roadblocks in our proof is that there exists another family of almost extremal F_5^t -free 3-graphs, whose number of hyperedges is smaller than $s_t^\circ(n)$ only by $O(n)$. Hence for the proof of Theorem 1.1, we need to have a good understanding of $\text{ex}(n, F_2^t)$. The value of $\text{ex}(n, F_2^t)$ is closely related to designs. An (n, k, r, t) -design is a family X of distinct k -subsets of an n -set

V , such that every r -subset of V belongs to exactly t elements of X . Hence, if every pair of vertices in hypergraph $H(n, t) \in \mathcal{H}(n, t)$ is in exactly t hyperedges, then $H(n, t)$ is just an $(n, 3, 2, t)$ -design and thus $H(n, t)$ has $\frac{t}{3}\binom{n}{2}$ hyperedges. Dehon [3] proved the following necessary and sufficient condition for the existence of $(n, 3, 2, t)$ -designs.

Theorem 1.2 (Dehon [3]). *Let t and n be integers such that $0 \leq t \leq n - 2$. Then, there exists an $(n, 3, 2, t)$ -design iff $tn(n - 1) \equiv 0 \pmod{6}$ and $t(n - 1) \equiv 0 \pmod{2}$.*

If t is a multiple of 6, then there always exists an $(n, 3, 2, t)$ -design for sufficiently large n . Keevash proved that the number of non-isomorphic $(n, 3, 2, t)$ -designs (see Theorem 6.1 in [15]) is

$$t!^{-\binom{n}{2}} \left(\left(\frac{t}{e} \right)^2 (n - 2) + o(n - 2) \right)^{\frac{t}{3}\binom{n}{2}} = n^{\Omega(n^2)}.$$

Thus, by Theorem 1.1, there are exponentially many extremal constructions for F_5^{6t} whenever $t \geq 1$.

Note that if $n = 6k + 1$ or $n = 6k + 3$ for some integer $k \geq (t + 1)/6$, then there always exists an $(n, 3, 2, t)$ -design. Thus, we have the following corollary of Theorem 1.1.

Corollary 1.3. *For every $t \geq 0$, if n is sufficiently large and $n \equiv 3$ or $9 \pmod{18}$, then*

$$\text{ex}(n, F_5^t) = \frac{1}{27}n^3 + \frac{1}{18}tn^2 - \frac{1}{6}tn.$$

There does not exist an $(n, 3, 2, t)$ -design for every n and t , and therefore it is non-trivial to give a reasonably good lower bounds on $\text{ex}(n, F_2^t)$ in general. In Section 2.2, we derive a lower bound on $\text{ex}(n, F_2^t)$ which is good enough for our purpose. We cannot give an explicit formula for $\text{ex}(n, F_2^t)$. Similar phenomena often occur in Extremal Combinatorics. An interesting classical example is by Erdős and Simonovits [6]. Denote by $K_{2,2,2}$ the complete tripartite graph with each part of size 2. They [6] proved that for sufficiently large n , the extremal $K_{2,2,2}$ -free graph is realized by the graphs whose vertex set can be partitioned into two subsets A and B such that $G[A]$ is C_4 -free and $G[B]$ is P_3 -free, where C_4 is the cycle with 4 vertices and P_3 is the path with 3 vertices. However, they did not determine the exact sizes of A and B , as $\text{ex}(n, C_4)$ is not known precisely for every n . Similarly to $\text{ex}(n, C_4)$, in our case $\text{ex}(n, F_2^t)$ is not known precisely for every n . We are still able to confirm the above-mentioned unified description of the extremal F_5^t -free hypergraphs and in particular, prove that the three parts in the partition have to be balanced, i.e., they have sizes $\lfloor n/3 \rfloor$, $\lfloor (n + 1)/3 \rfloor$ and $\lfloor (n + 2)/3 \rfloor$.

Our paper is organized as follows. In Section 2, we introduce our notation, give an estimate on $\text{ex}(n, F_2^t)$ and some useful results about $S_t^\circ(n)$, and deduce a stability result for F_5^t . In Section 3, we prove Theorem 1.1, our main result. We discuss some open problems in Section 4.

2 Preliminaries

2.1 Notation

For a positive integer m , we use $[m]$ for the set $\{1, 2, \dots, m\}$. We write xy for the set $\{x, y\}$ and xyz for the set $\{x, y, z\}$. For a set S and a non-negative integer k , we use $\binom{S}{k}$ to denote the family of all k -subsets of S . For an n -vertex r -graph H , we will assume its vertex set $V(H)$ is $[n]$. We often use H for the hyperedge set of H and denote by $|H|$ the number of hyperedges in H . Let H be an n -vertex 3-graph. For vertices $x, y \in [n]$ and (not necessarily disjoint) subsets of vertices $S, T \subseteq [n]$, let

- $L_{S,T}(x) := \{yz : y \in S, z \in T, xyz \in H\}$ be the *link graph* of x between S and T ,
- $d_{S,T}(x) := |L_{S,T}(x)|$ be the *degree* of x between S and T ,
- $L_S(x, y) := \{z : z \in S, xyz \in H\}$ be the set of *neighbors* of x and y in S , and
- $d_S(x, y) := |L_S(x, y)|$ be the *codegree* of x and y in S .

For a given partition $\pi = (V_1, V_2, V_3)$ of a subset of $[n]$, we say that π is *balanced* if the sizes of its parts differ by at most 1. We denote by K_π the set of triples with exactly one vertex in each part of π . Let $H_\pi := H \cap K_\pi$ and $\bar{H}_\pi := K_\pi \setminus H_\pi$. We call hyperedges in H_π the *crossing hyperedges* (of π) and call hyperedges in \bar{H}_π the *missing crossing hyperedges* (of π). For $i, j \in \{1, 2, 3\}$, we simply write $L_{i,j}(x)$ for $L_{V_i, V_j}(x)$ and $L_i(x, y)$ for $L_{V_i}(x, y)$. Similarly, we use $d_{i,j}(x)$ for $d_{V_i, V_j}(x)$ and $d_i(x, y)$ for $d_{V_i}(x, y)$. We also write $L(x, y)$ for $L_{[n]}(x, y)$, $L(x)$ for $L_{[n], [n]}(x)$, $d(x, y)$ for $d_{[n]}(x, y)$, and $d(x)$ for $d_{[n], [n]}(x)$.

2.2 Results on $s_t^\circ(\mathbf{n})$

Proposition 2.1. *For every integer $t \geq 0$ and sufficiently large n , we have*

$$\frac{t}{3} \binom{n}{2} \geq \text{ex}(n, F_2^t) \geq \begin{cases} \frac{t}{3} \binom{n}{2} - \frac{n}{3} - \frac{t^2}{6} - 3t & \text{if } t \text{ is odd and } n \text{ is even,} \\ \frac{t}{3} \binom{n}{2} - \frac{8t}{3} & \text{if } t \text{ is odd and } n \text{ is odd,} \\ \frac{t}{3} \binom{n}{2} - \frac{2t}{3} & \text{if } t \text{ is even.} \end{cases}$$

We give the proof of Proposition 2.1 in the Appendix.

Lemma 2.2. *For every integer $t \geq 0$ and sufficiently large n , we have*

$$s_t^\circ(n) \in \left[\frac{1}{27}n^3 + \frac{1}{18}tn^2 - \left(\frac{1}{6}t + \frac{4}{9} \right)n - \left(\frac{1}{2}t^2 + \frac{80}{9}t + \frac{2}{27} \right), \frac{1}{27}n^3 + \frac{1}{18}tn^2 - \frac{1}{6}tn \right].$$

Proof. By the definition of $s_t^\circ(n)$ and Proposition 2.1, we have the following statements.

Case: $n \equiv 0 \pmod{3}$

$$\begin{aligned} s_t^\circ(n) &= \left(\frac{n}{3} \right)^3 + 3 \cdot \text{ex} \left(\frac{n}{3}, F_2^t \right) \\ &\in \left[\frac{1}{27}n^3 + \frac{1}{18}tn^2 - \left(\frac{1}{6}t + \frac{1}{3} \right)n - \left(\frac{1}{2}t^2 + 9t \right), \frac{1}{27}n^3 + \frac{1}{18}tn^2 - \frac{1}{6}tn \right]. \end{aligned}$$

Case: $n \equiv 1 \pmod{3}$

$$\begin{aligned} s_t^\circ(n) &= \left(\frac{n-1}{3}\right)^2 \binom{n+2}{3} + 2 \cdot \text{ex}\left(\frac{n-1}{3}, F_2^t\right) + \text{ex}\left(\frac{n+2}{3}, F_2^t\right) \\ &\in \left[\frac{1}{27}n^3 + \frac{1}{18}tn^2 - \left(\frac{1}{6}t + \frac{4}{9}\right)n - \left(\frac{1}{2}t^2 + \frac{80}{9}t - \frac{2}{27}\right), \right. \\ &\quad \left. \frac{1}{27}n^3 + \frac{1}{18}tn^2 - \left(\frac{1}{6}t + \frac{1}{9}\right)n + \left(\frac{1}{9}t + \frac{2}{27}\right) \right]. \end{aligned}$$

Case: $n \equiv 2 \pmod{3}$

$$\begin{aligned} s_t^\circ(n) &= \left(\frac{n-2}{3}\right) \binom{n+1}{3} + \text{ex}\left(\frac{n-2}{3}, F_2^t\right) + 2 \cdot \text{ex}\left(\frac{n+1}{3}, F_2^t\right) \\ &\in \left[\frac{1}{27}n^3 + \frac{1}{18}tn^2 - \left(\frac{1}{6}t + \frac{4}{9}\right)n - \left(\frac{1}{2}t^2 + \frac{80}{9}t + \frac{2}{27}\right), \right. \\ &\quad \left. \frac{1}{27}n^3 + \frac{1}{18}tn^2 - \left(\frac{1}{6}t + \frac{1}{9}\right)n + \left(\frac{1}{9}t - \frac{2}{27}\right) \right]. \quad \square \end{aligned}$$

For a partition $\pi = (V_1, V_2, V_3)$ of $[n]$, let $\mathcal{S}_{t,\pi}^\circ(n)$ be the family of hypergraphs on vertex set $[n]$ whose hyperedges contain K_π , and each part V_i spans a copy of $H(|V_i|, t) \in \mathcal{H}(|V_i|, t)$. Hence, $\mathcal{S}_{t,\pi}^\circ(n)$ is just $\mathcal{S}_t^\circ(n)$ when π is balanced. Let

$$s_{t,\pi}^\circ(n) := |V_1||V_2||V_3| + \sum_{i=1}^3 \text{ex}(|V_i|, F_2^t)$$

be the number of hyperedges in every $S_{t,\pi}^\circ(n) \in \mathcal{S}_{t,\pi}^\circ(n)$.

Lemma 2.3. *Let $t \geq 0$. For every sufficiently large integer n , we have $s_t^\circ(n) \geq s_{t,\pi}^\circ(n)$ for every partition π of $[n]$, where equality holds only if π is balanced.*

Proof. Suppose that there is a partition $\pi = (V_1, V_2, V_3)$ of $[n]$ that achieves the maximum value of $s_{t,\pi}^\circ(n)$, but π is not balanced. Assume that the three parts of π have sizes $a_1 \leq a_2 \leq a_3$, where $a_1 + a_2 + a_3 = n$ and $a_1 + 2 \leq a_3$. We first prove $a_2 \geq (\frac{1}{3} - 0.01)n$. Otherwise, by Proposition 2.1 and Lemma 2.2, we have

$$\begin{aligned} s_{t,\pi}^\circ(n) &\leq a_1 a_2 a_3 + \frac{t}{3} \left(\binom{a_1}{2} + \binom{a_2}{2} + \binom{a_3}{2} \right) \leq \frac{a_1 + a_3}{2} a_2 \frac{a_1 + a_3}{2} + t \binom{n}{2} \\ &= a_2 \left(\frac{n - a_2}{2} \right)^2 + t \binom{n}{2} \leq \left(\frac{1}{3} - 0.01 \right) n \left(\frac{n - (\frac{1}{3} - 0.01)n}{2} \right)^2 + t \binom{n}{2} \\ &< \left(\frac{1}{27} - 10^{-10} \right) n^3 < s_t^\circ(n), \end{aligned}$$

where the third-to-last inequality holds because the function $x \rightarrow x((n-x)/2)^2$ is monotone increasing for $x \leq n/3$. This contradicts our choice of π . Now, let π' be the partition of $[n]$ obtained from π by moving a vertex from V_3 to V_1 . Then,

$$s_{t,\pi'}^\circ(n) - s_{t,\pi}^\circ(n) = (a_3 - a_1 - 1)a_2 + \text{ex}(a_1 + 1, F_2^t) + \text{ex}(a_3 - 1, F_2^t) - \text{ex}(a_1, F_2^t) - \text{ex}(a_3, F_2^t).$$

By Proposition 2.1,

$$\text{ex}(a_1 + 1, F_2^t) - \text{ex}(a_1, F_2^t) \geq \frac{t}{3} \binom{a_1 + 1}{2} - \frac{a_1 + 1}{3} - \frac{t^2}{6} - 3t - \frac{t}{3} \binom{a_1}{2} \geq \frac{t}{3} a_1 - \frac{a_1 + 1}{3} - 4t^2$$

and

$$\text{ex}(a_3 - 1, F_2^t) - \text{ex}(a_3, F_2^t) \geq \frac{t}{3} \binom{a_3 - 1}{2} - \frac{a_3 - 1}{3} - 4t^2 - \frac{t}{3} \binom{a_3}{2} = -\frac{t+1}{3}(a_3 - 1) - 4t^2.$$

We conclude

$$\begin{aligned} s_{t,\pi'}^\circ(n) - s_{t,\pi}^\circ(n) &\geq (a_3 - a_1 - 1) \left(a_2 - \frac{t}{3} \right) - \frac{a_1 + a_3}{3} - 8t^2 \geq \left(a_2 - \frac{t}{3} \right) - \frac{a_1 + a_3}{3} - 8t^2 \\ &= \frac{4}{3}a_2 - \frac{n}{3} - \frac{t}{3} - 8t^2 > 0, \end{aligned}$$

contradicting our choice of π . □

2.3 An almost extremal F_5^t -free hypergraph

For $T = \{v_1, \dots, v_t\} \subseteq [n]$ and a partition $\varpi = (W_1, W_2, W_3)$ of $[n] \setminus T$, define $S_{t,\varpi}^\Delta(n)$ to be the 3-graph on vertex set $[n]$ with hyperedges

$$K_\varpi \cup \binom{T}{3} \cup \bigcup_{i=1}^3 \{vx_1x_2 : v \in T, x_1, x_2 \in W_i\}.$$

For an illustration of $S_{t,\varpi}^\Delta(n)$, see Figure 5. Let

$$s_{t,\varpi}^\Delta(n) := |W_1||W_2||W_3| + t \sum_{i=1}^3 \binom{|W_i|}{2} + \binom{t}{3}$$

be the number of hyperedges in $S_{t,\varpi}^\Delta(n)$. We remark that similar constructions were proved by Simonovits to be extremal constructions for certain graphs (see Theorem 2 in [22]).

Lemma 2.4. *Let $t \geq 1$. For every sufficiently large integer n , we have $s_t^\circ(n) > s_{t,\varpi}^\Delta(n) + \frac{n}{10}$ for every set $\{v_1, \dots, v_t\} \subseteq [n]$ and partition ϖ of $[n] \setminus \{v_1, \dots, v_t\}$.*




Figure 5: Hypergraph $S_{t, \varpi}^{\Delta}(n)$.

Proof. Define function

$$f_{n,t}(x_1, x_2, x_3) := x_1 x_2 x_3 + t \left(\frac{x_1(x_1 - 1)}{2} + \frac{x_2(x_2 - 1)}{2} + \frac{x_3(x_3 - 1)}{2} \right) + \binom{t}{3}$$

on the domain $D := \{(x_1, x_2, x_3) \in \mathbb{R}^3 : x_1, x_2, x_3 \geq 0, x_1 + x_2 + x_3 = n - t\}$. Note that for every set $T = \{v_1, \dots, v_t\} \subseteq [n]$ and partition $\varpi = (W_1, W_2, W_3)$ of $[n] \setminus T$, we have $s_{t, \varpi}^{\Delta}(n) = f(|W_1|, |W_2|, |W_3|)$. Also note that

$$f_{n,t} \left(\frac{n-t}{3}, \frac{n-t}{3}, \frac{n-t}{3} \right) = \frac{1}{27} n^3 + \frac{1}{18} t n^2 - \left(\frac{2}{9} t^2 + \frac{1}{2} t \right) n + \frac{8}{27} t^3 + \frac{1}{3} t.$$

Since D is compact, $f_{n,t}$ achieves its maximum value at some point $(y_1, y_2, y_3) \in D$. If $y_1 \leq t$, then

$$f_{n,t}(y_1, y_2, y_3) \leq t n^2 + 3t \frac{n^2}{2} + t^3 < \frac{n^3}{27} < f_{n,t} \left(\frac{n-t}{3}, \frac{n-t}{3}, \frac{n-t}{3} \right),$$

a contradiction. Thus, $y_1 > t$. Similarly, $y_2 > t$ and $y_3 > t$. We have

$$f_{n,t} \left(y_1, \frac{y_2 + y_3}{2}, \frac{y_2 + y_3}{2} \right) - f_{n,t}(y_1, y_2, y_3) = \frac{1}{4} (y_1 - t)(y_2 - y_3)^2 \geq 0,$$

with equality iff $y_2 = y_3$. Using the symmetry of f , we conclude that the maximum value of $f_{n,t}$ is achieved when $x_1 = x_2 = x_3 = \frac{n-t}{3}$. For $t \geq 1$, by Lemma 2.2, we have

$$s_t^{\circ}(n) - f_{n,t} \left(\frac{n-t}{3}, \frac{n-t}{3}, \frac{n-t}{3} \right) \geq \left(\frac{2}{9} t^2 + \frac{1}{3} t - \frac{4}{9} \right) n - \left(\frac{8}{27} t^3 + \frac{1}{2} t^2 + \frac{83}{9} t + \frac{2}{27} \right) > \frac{n}{10}.$$

Therefore, $s_t^{\circ}(n) > s_{t, \varpi}^{\Delta}(n) + \frac{1}{10} n$. \square

2.4 Stability result

In this section, we prove the following stability theorem for F_5^t .

Theorem 2.5 (Stability theorem for F_5^t). *For every $t \geq 0$ and $\varepsilon > 0$, there exist $\delta \in (0, \varepsilon)$ and $N > 0$ such that for every $n > N$, the following statement holds. Let H be an F_5^t -free 3-graph H on vertex set $[n]$ with at least $(\frac{1}{27} - \delta)n^3$ hyperedges, and let $\pi = (V_1, V_2, V_3)$ be a partition of $[n]$ that maximizes $|H_{\pi}|$. Then $|H \setminus H_{\pi}| \leq \varepsilon n^3$.*

Claim 2.6. *The partition $\pi = (V_1, V_2, V_3)$ in Theorem 2.5 has the following additional properties.*

(i) *For $i \in [3]$, we have $(\frac{1}{3} - 3\varepsilon^{1/2})n < |V_i| < (\frac{1}{3} + 6\varepsilon^{1/2})n$.*

(ii) *For $\{i, j, k\} = [3]$, we have $|\{xy : x \in V_i, y \in V_j, d_k(x, y) \leq |V_k| - \varepsilon^{1/2}n\}| \leq 2\varepsilon^{1/2}n^2$.*

(iii) *For $\{i, j, k\} = [3]$ and every $x \in V_i$, we have $d_{i,j}(x), d_{i,k}(x) \leq 3\varepsilon^{1/2}n^2$.*

We need the following theorems and definitions for proving Theorem 2.5. For an r -graph H , a *blow-up* of H is an r -graph obtained from H by replacing every vertex v by a set S_v , where $S_{v_1} \cap S_{v_2} = \emptyset$ for distinct vertices v_1, v_2 and replacing every hyperedge $\{v_1, \dots, v_r\}$ by the complete r -partite r -graph on $S_{v_1} \cup \dots \cup S_{v_r}$. When $|S_v| = t$ for every vertex v , the blow-up is called the t -*blow-up* of H and denoted by $H[t]$. Note that F_5^t is a blow-up of F_5 and $F_5^t \subseteq F_5[t + 1]$ for every $t \geq 0$. We will see that Theorem 2.5 follows easily from the stability result for F_5 and the fact that blow-ups inherit the stability of the original hypergraph.

Theorem 2.7 (Stability result for F_5 , Keevash and Mubayi [16]). *For every $a > 0$, there exist $b > 0$ and $N > 0$ such that if $n > N$ and H is an F_5 -free 3-graph on vertex set $[n]$ with at least $(\frac{1}{27} - b)n^3$ hyperedges, then there exists a partition $\pi = (V_1, V_2, V_3)$ of $[n]$ such that $|H \setminus H_\pi| \leq an^3$.*

Lemma 2.8 (e.g. [14]). *For fixed integers $l \geq r \geq 2$, $t \geq 1$ and every $\alpha > 0$, there exists $n_0 = n_0(l, r, t, \alpha)$ such that the following holds. Let F be an r -graph on l vertices and H be an r -graph on $n \geq n_0$ vertices. If H contains at least αn^l copies of F , then H contains a copy of $F[t]$.*

We also use the hypergraph removal lemma.

Theorem 2.9 (Rödl, Nagle, Skokan, Schacht, and Kohayakawa [21], Gower [12]). *For fixed integers $l \geq r \geq 2$ and every $\mu > 0$, there exists $\zeta = \zeta(l, r, \mu) > 0$ and $n_0 = n_0(l, r, \mu)$ such that the following holds. Let F be an r -graph on l vertices and H be an r -graph on $n \geq n_0$ vertices. If H contains at most ζn^l copies of F , then one can delete μn^r hyperedges of H to make it F -free.*

Proof of Theorem 2.5. Assume that n is sufficiently large. Let b be given by Theorem 2.7 when applying it for $a = \varepsilon/2$. Let $\delta := \min(b/2, \varepsilon/2)$. Let H be an F_5^t -free 3-graph on vertex set $[n]$ with at least $(\frac{1}{27} - \delta)n^3$ hyperedges and let π be a 3-partition of $[n]$ maximizing $|H_\pi|$.

Setting $\mu = \delta$, $l = 5$, and $r = 3$ in Theorem 2.9, we get that there exists $\zeta > 0$ such that if an n -vertex 3-graph F contains at most ζn^5 copies of F_5 , then we can delete at most δn^3 hyperedges to make it F_5 -free. Setting $\alpha = \zeta$ in Theorem 2.8, we get that if H contains at least ζn^5 copies of F_5 , then H contains a copy of $F[t + 1] \supseteq F_5^t$, a contradiction to that H is F_5^t -free, so H indeed contains at most ζn^5 copies of F_5 . Therefore, H contains a spanning F_5 -free subhypergraph H' with at least $|H| - \delta n^3 \geq (\frac{1}{27} - 2\delta)n^3 \geq (\frac{1}{27} - b)n^3$ hyperedges. By Theorem 2.7, there exists a 3-partition π' of $[n]$ such that $|H' \setminus H'_{\pi'}| \leq \varepsilon n^3/2$. Thus,

$$|H \setminus H_\pi| \leq |H \setminus H_{\pi'}| \leq |H \setminus H'| + |H' \setminus H_{\pi'}| = |H \setminus H'| + |H' \setminus H'_{\pi'}| \leq \varepsilon n^3. \quad \square$$

Proof of Claim 2.6. For (i), assume $|V_i| \leq (\frac{1}{3} - 3\varepsilon^{1/2})n$ for some $i \in [3]$. The function $x \rightarrow x(n-x)/2$ is monotone increasing when $x < n/3$, and therefore

$$|H| \leq |V_1||V_2||V_3| + |H \setminus H_\pi| \leq |V_i| \left(\frac{n - |V_i|}{2} \right)^2 + \varepsilon n^3 \leq \left(\frac{1}{27} - \frac{5}{4}\varepsilon - \frac{27}{4}\varepsilon^{3/2} \right) n^3 < |H|,$$

a contradiction. Thus $|V_i| > (\frac{1}{3} - 3\varepsilon^{1/2})n$ for every $i \in [3]$. Since $|V_1| + |V_2| + |V_3| = n$, we have $|V_i| < (\frac{1}{3} + 6\varepsilon^{1/2})n$ for $i \in [3]$.

Assume that (ii) does not hold. Then $|\bar{H}_\pi| > \varepsilon^{1/2}n \cdot 2\varepsilon^{1/2}n^2 = 2\varepsilon n^3$, and thus

$$|H| = |V_1||V_2||V_3| - |\bar{H}_\pi| + |H \setminus H_\pi| \leq \left(\frac{n}{3} \right)^3 - 2\varepsilon n^3 + \varepsilon n^3 < |H|,$$

a contradiction.

For (iii), assume for contradiction that $d_{i,j}(x) > 3\varepsilon^{1/2}n^2$ for some $x \in V_i$. Then, by (ii), there exist $x' \in V_i$ and $y \in V_j$ such that $xx'y \in H$ and $d_k(x', y) > |V_k| - \varepsilon^{1/2}n$. By our assumption that π maximizes $|H_\pi|$, we have $d_{j,k}(x) \geq d_{i,j}(x) > 3\varepsilon^{1/2}n^2$; otherwise moving x to V_k will strictly increase $|H_\pi|$. There exists a vertex $z \in V_k$ such that $x'yz \in H$ and $d_j(x, z) \geq t + 2$; otherwise for every z with $x'yz \in H$ we have $d_j(x, z) \leq t + 1$ and then

$$d_{j,k}(x) \leq (t + 1)d_k(x', y) + |V_j| (|V_k| - d_k(x', y)) \leq (t + 1)n + n \cdot \varepsilon^{1/2}n < 3\varepsilon^{1/2}n^2,$$

a contradiction. Now choose distinct vertices $y_1, \dots, y_{t+1} \in L_j(x, z) \setminus \{y\}$. The hyperedges $\{x'yx, x'yz, xzy_1, \dots, xzy_{t+1}\}$ form a copy of F_5^t in H , a contradiction. \square

3 Proof of Theorem 1.1

In Section 3.1, using the stability result for F_5^t , Theorem 2.5, we show that if H is an extremal F_5^t -free hypergraph, then $|\bar{H}_\pi| = O(n^2)$ for some partition π . For a hypergraph H on vertex set $[n]$, denote by $\delta(H) := \min\{d(v) : v \in [n]\}$ the *minimum degree* of H . In Sections 3.2, 3.3, and 3.4, we prove Theorem 1.1 with an extra assumption on the minimum degree, by distinguishing “typical” and “non-typical” vertices and using their respective properties. Finally, in Section 3.5, we derive Theorem 1.1.

3.1 Upper bound on $|\bar{H}_\pi|$

Lemma 3.1. *Let H be an F_5^t -free 3-graph on vertex set $[n]$. For every pair of vertices $x, x' \in [n]$ with $d(x, x') \geq t + 1$ and disjoint subsets of vertices $S, T \subseteq [n]$, we have*

$$d_{S,T}(x) + d_{S,T}(x') \leq |S||T| + (t + 3)n$$

and

$$d_{S,S}(x) + d_{S,S}(x') \leq \frac{1}{2}|S|^2 + (t + 4)n.$$

Proof. Fix $t + 1$ distinct vertices $v_1, \dots, v_{t+1} \in L(x, x')$. There are at least $(|S| - t - 3)(|T| - t - 3)$ pairs yz such that $y \in S$, $z \in T$, and $\{y, z\} \cap \{x, x', v_1, \dots, v_{t+1}\} = \emptyset$. For each such pair, $yzx \notin H$ or $yzx' \notin H$; otherwise the hyperedges $\{yzx, yzx', xx'v_1, \dots, xx'v_{t+1}\}$ form a copy of F_5^t in H . Then,

$$d_{S,T}(x) + d_{S,T}(x') \leq 2|S||T| - (|S| - t - 3)(|T| - t - 3) \leq |S||T| + (t + 3)n.$$

Similarly, there are at least $\binom{|S| - t - 3}{2}$ pairs yz such $y, z \in S$ and $\{y, z\} \cap \{x, x', v_1, \dots, v_{t+1}\} = \emptyset$. Again, for each such pair, $yzx \notin H$ or $yzx' \notin H$. We conclude

$$d_{S,S}(x) + d_{S,S}(x') \leq 2\binom{|S|}{2} - \binom{|S| - t - 3}{2} \leq \frac{1}{2}|S|^2 + (t + 4)n. \quad \square$$

Theorem 3.2. *For every $t \geq 0$ and $\varepsilon > 0$, there exists $N > 0$ such that if $n > N$ the following holds. Let H be an F_5^t -free 3-graph on vertex set $[n]$ with $|H| \geq s_t^\circ(n)$ and π be a partition of $[n]$ maximizing $|H_\pi|$. Then, $|\bar{H}_\pi| \leq (\frac{1}{9} + \varepsilon)tn^2$.*

Proof. We can assume that ε is sufficiently small and n is sufficiently large. Let $\beta > 0$ be the real number such that $\varepsilon = (2t + 25)\beta^{1/2}$. For $i \in [3]$, we define

$$P_i := \{xx' : x, x' \in V_i, d(x, x') \leq t\} \quad \text{and} \quad Q_i := \{xx' : x, x' \in V_i, d(x, x') \geq t + 1\}.$$

Let M_i be a maximum matching of Q_i , and let E_i be the set of endpoints of the pairs in M_i , i.e., the vertices $x \in V_i$ such that there exists $x' \in V_i$ with $xx' \in M_i$. Note that $|E_i| = 2|M_i|$ and every pair in Q_i contains at least one point in E_i , since M_i is a maximum matching of Q_i .

Claim 3.3. *We have $|Q_i| \leq (t + 2)\beta^{1/2}n^2$ and $|M_i| \leq (t + 2)\beta^{1/2}n$ for $i \in [3]$.*

Proof. For $\{i, j, k\} = [3]$, define $S_{j,k}$ to be the set of pairs yz such that $y \in V_j, z \in V_k$, and $d_i(y, z) \geq |V_i| - \beta^{1/2}n$. By Claim 2.6 (i) and (ii),

$$|S_{j,k}| \geq |V_j||V_k| - 2\beta^{1/2}n^2 \geq \left(\left(\frac{1}{3} - 3\beta^{1/2} \right) n \right)^2 - 2\beta^{1/2}n^2 \geq \frac{n^2}{10}.$$

Thus, we can choose $t + 2$ pairwise disjoint pairs $y_1z_1, \dots, y_{t+2}z_{t+2}$ from $S_{j,k}$. Define U_i to be the set of vertices $x \in V_i$ such that $xy_a z_a \notin H$ for some $a \in [t + 2]$. Then, $|U_i| \leq (t + 2)\beta^{1/2}n$.

For $x_1, x_2 \in V_i \setminus U_i$, we have $x_1x_2 \notin Q_i$; otherwise there are $t + 1$ distinct vertices $v_1, \dots, v_{t+1} \in L(x_1, x_2)$ and there is some $a \in [t + 2]$ such that $\{v_1, \dots, v_{t+1}\} \cap \{y_a, z_a\} = \emptyset$, and hence, the hyperedges $\{y_a z_a x_1, y_a z_a x_2, x_1 x_2 v_1, \dots, x_1 x_2 v_{t+1}\}$ form a copy of F_5^t in H , a contradiction. Therefore, every pair in Q_i contains at least one vertex in U_i . We conclude $|Q_i| \leq |U_i||V_i| \leq (t + 2)\beta^{1/2}n^2$ and $|M_i| \leq |U_i| \leq (t + 2)\beta^{1/2}n$. \square

Claim 3.4. *We have*

$$|H \setminus H_\pi| \leq \left(\frac{1}{6} + 6\beta^{1/2} + 54\beta \right) tn^2 + |M|(2t + 16)\beta^{1/2}n^2.$$

Proof. For $i \in [3]$, let H_i be the set of hyperedges in H containing at least two vertices in V_i . The number of hyperedges in H_i that contain a pair in P_i is most $t|P_i|$, because every pair in P_i has codegree at most t by definition. The number of hyperedges in H_i that contain no pair in P_i and also no vertex in $[n] \setminus V_i$ is at most $|E_i||Q_i|$. The number of hyperedges in H_i that contain no pair in P_i but a vertex in $[n] \setminus V_i$ is at most $|E_i| \cdot 2 \cdot 3\beta^{1/2}n^2$ by Claim 2.6 (iii). Adding up the number of hyperedges of these three types, we get

$$\begin{aligned} |H_i| &\leq t \binom{\left(\frac{1}{3} + 6\beta^{1/2}\right)n}{2} + 2(t+2)\beta^{1/2}n^2|M_i| + 12\beta^{1/2}n^2|M_i| \\ &\leq \left(\frac{1}{18} + 2\beta^{1/2} + 18\beta\right)tn^2 + (2t+16)\beta^{1/2}n^2|M_i|, \end{aligned}$$

where the first inequality follows from Claims 2.6 (i) and 3.3. Then,

$$|H \setminus H_\pi| = \sum_{i=1}^3 |H_i| \leq \left(\frac{1}{6} + 6\beta^{1/2} + 54\beta\right)tn^2 + |M|(2t+16)\beta^{1/2}n^2. \quad \square$$

Claim 3.5. *We have*

$$|\bar{H}_\pi| + \sum_{i=1}^3 \text{ex}(|V_i|, F_2^t) \geq |M| \left(\frac{1}{9} - (2t+7)\beta^{1/2}\right)n^2 + \left(\frac{1}{18} - 2\beta^{1/2}\right)tn^2.$$

Proof. For $\{i, j, k\} = [3]$ and $x_1x_2 \in M_i$, by Lemma 3.1, we have

$$d_{V_j \setminus E_j, V_k \setminus E_k}(x_1) + d_{V_j \setminus E_j, V_k \setminus E_k}(x_2) \leq |V_j \setminus E_j||V_k \setminus E_k| + (t+3)n,$$

so there are at least

$$|M_i| (2|V_j \setminus E_j||V_k \setminus E_k| - (|V_j \setminus E_j||V_k \setminus E_k| + (t+3)n))$$

missing crossing hyperedges containing exactly one vertex in each of E_i , $V_j \setminus E_j$, and $V_k \setminus E_k$. By Claims 2.6 (i) and 3.3, we have

$$|\bar{H}_\pi| \geq |M| \left(\left(\left(\frac{1}{3} - 3\beta^{1/2} - 2(t+2)\beta^{1/2} \right) n \right)^2 - (t+3)n \right) \geq |M| \left(\frac{1}{9} - (2t+7)\beta^{1/2} \right) n^2.$$

By Claim 2.6 (i) and the lower bounds on $\text{ex}(n, F_2^t)$ in Proposition 2.1, we have

$$\sum_{i=1}^3 \text{ex}(|V_i|, F_2^t) \geq t \binom{\left\lfloor \left(\frac{1}{3} - 3\beta^{1/2} \right) n \right\rfloor}{2} - \frac{n}{3} - \frac{t^2}{2} - 9t \geq \left(\frac{1}{18} - 2\beta^{1/2} \right) tn^2. \quad (1)$$

Summing them up, we get the lower bound desired. \square

Claim 3.6. *We have $|M| \leq t$.*

Proof. Assume $|M| \geq t + 1$. We get $|H \setminus H_\pi| < |\bar{H}_\pi| + \sum_{i=1}^3 \text{ex}(|V_i|, F_2^t)$ by Claims 3.4 and 3.5. This implies $|H| < s_{t,\pi}^\circ(n) < s_t^\circ(n)$ by Lemma 2.3, a contradiction. \square

Finally, Theorem 3.2 follows from a short calculation. Since $|H| \geq s_{t,\pi}^\circ(n)$, we have $|\bar{H}_\pi| \leq |H \setminus H_\pi| - \sum_{i=1}^3 \text{ex}(|V_i|, F_2^t)$ and thus by Claims 3.4, 3.6, and (1), we get

$$\begin{aligned} |\bar{H}_\pi| &\leq \left(\frac{1}{6} + 6\beta^{1/2} + 54\beta \right) tn^2 + (2t + 16)\beta^{1/2}tn^2 - \left(\frac{1}{18} - 2\beta^{1/2} \right) tn^2 \\ &\leq \left(\frac{1}{9} + (2t + 25)\beta^{1/2} \right) tn^2 = \left(\frac{1}{9} + \varepsilon \right) tn^2. \end{aligned} \quad \square$$

Remark 3.7. For $t = 0$, Theorem 3.2 states that an extremal F_5 -free n -vertex 3-graph is complete tripartite for sufficiently large n . This proves Theorem 1.1 in the case $F_5^0 = F_5$.

3.2 Typical and non-typical vertices

For every integer $t \geq 0$, let $c = c(t) := (t + 100)^{-1000}$. For a 3-graph H on vertex set $[n]$ and a partition $\pi = (V_1, V_2, V_3)$ of $[n]$, define

$$\begin{aligned} B_i &:= \{x \in V_i : d_{j,k}(x) \leq cn^2\}, \quad A_i := V_i \setminus B_i, \quad \text{for } \{i, j, k\} = [3], \\ B &:= B_1 \cup B_2 \cup B_3, \quad \text{and} \quad A := A_1 \cup A_2 \cup A_3. \end{aligned}$$

We call vertices in A *typical* and vertices in B *non-typical*. In Sections 3.2.1 and 3.2.2, we prove properties of non-typical and typical vertices, respectively.

3.2.1 Non-typical vertices

Lemma 3.8. Let $t \geq 0$, n be sufficiently large, and H be an F_5^t -free 3-graph on vertex set $[n]$ with $|H| \geq s_t^\circ(n)$ and $\delta(H) \geq \frac{1}{9}n^2 - n$. Further, let $\pi = (V_1, V_2, V_3)$ be a partition of $[n]$ maximizing $|H_\pi|$ and c, B be defined as above. The following statements hold.

(i) $|B| \leq t$.

(ii) For every vertex $v \in B$, we have $d_{1,2}(v), d_{1,3}(v), d_{2,3}(v) \leq cn^2$.

(iii) For every pair of distinct vertices $v, v' \in B$, we have $d(v, v') \leq t$.

Proof of (i). By Claim 2.6 (i), Theorem 3.2, and the definition of B_i , we first have $|B_i| \leq t$ for $i \in [3]$. Let $\{i, j, k\} = [3]$. For every $x \in B_i$, the number of missing crossing hyperedges that contain x and do not contain any vertex from $B_j \cup B_k$ is at least

$$(|V_j| - |B_j|)(|V_k| - |B_k|) - cn^2 \geq \left(\frac{1}{3} - 4c \right)^2 n^2 - cn^2 \geq \left(\frac{1}{9} - 4c \right) n^2.$$

Thus, by Theorem 3.2, we have $|B| \left(\frac{1}{9} - 4c \right) n^2 \leq |\bar{H}_\pi| \leq \left(\frac{1}{9} + c \right) tn^2$, implying $|B| \leq t$. \square

Proof of (ii). Let $\{i, j, k\} = [3]$. By the definition of B_i , we have $d_{j,k}(x) \leq cn^2$ for every $x \in B_i$. As π maximizes $|H_\pi|$, we have $d_{i,j}(x), d_{i,k}(x) \leq d_{j,k}(x) \leq cn^2$. \square

Proof of (iii). Assume for contradiction that there exist vertices $v, v' \in B$ such that $d(v, v') \geq t + 1$. By Claim 2.6 (i) and Lemma 3.1, we have

$$\sum_{i=1}^3 (d_{i,i}(v) + d_{i,i}(v')) \leq \sum_{i=1}^3 \left(\frac{1}{2}|V_i|^2 + (t+4)n \right) \leq \frac{n^2}{5}.$$

Thus, without loss of generality, we can assume that $\sum_{i=1}^3 d_{i,i}(v) \leq \frac{1}{10}n^2$. By (ii), we have

$$d(v) = \sum_{i=1}^3 d_{i,i}(v) + \sum_{1 \leq i < j \leq 3} d_{i,j}(v) \leq \frac{1}{10}n^2 + 3cn^2 < \frac{1}{9}n^2 - n,$$

contradicting the minimum degree assumption. \square

3.2.2 Typical vertices

Lemma 3.9. *Let $t \geq 0$, n be sufficiently large, and H be an F_5^t -free 3-graph on vertex set $[n]$ with $|H| \geq s_t^\circ(n)$ and $\delta(H) \geq \frac{1}{9}n^2 - n$. Further, let $\pi = (V_1, V_2, V_3)$ be a partition of $[n]$ maximizing $|H_\pi|$ and c, A_1, A_2, A_3 be defined as above. The following statements hold.*

(i) *Let $\{i, j, k\} = [3]$. For every vertex $x \in A_i$, we have*

- (a) $d_{i,j}(x), d_{i,k}(x) \leq 3cn^2$,
- (b) $d_{i,i}(x) \leq n^{3/2}$,
- (c) $d_{j,j}(x), d_{k,k}(x) \leq cn^2$, and
- (d) $d_{j,k}(x) \geq (\frac{1}{9} - 10c)n^2$.

(ii) *For $i \in [3]$ and every pair of distinct vertices $x, x' \in A_i$, we have $d(x, x') \leq t$.*

(iii) *For $i \neq j \in [3]$ and vertices $x \in A_i, y \in A_j$, if $d_{A_i}(x, y) + d_{A_j}(x, y) > 0$, then we have $d_{A_k}(x, y) \leq 50cn$.*

Proof of (i). (a) follows from Claim 2.6 (iii). For (b), assume for contradiction that there exists vertex $x \in A_i$ such that $d_{i,i}(x) > n^{3/2}$. Let $S_x := \{x' \in V_i : d_i(x, x') \geq t + 1\}$. Then

$$n^{3/2} < d_{i,i}(x) \leq \frac{1}{2}(|S_x||V_i| + |V_i \setminus S_x| \cdot t) \leq \frac{1}{2}(|S_x|n + tn),$$

implying $|S_x| \geq n^{1/2}$. For every $x' \in S_x$, by Lemma 3.1, we have $d_{j,k}(x) + d_{j,k}(x') \leq |V_j||V_k| + (t+3)n$. By the definition of A_i , we have $d_{j,k}(x) \geq cn^2$ and hence $d_{j,k}(x') \leq |V_j||V_k| + (t+3)n - cn^2 \leq |V_j||V_k| - \frac{1}{2}cn^2$. Thus,

$$|\bar{H}_\pi| = \sum_{x' \in V_i} (|V_j||V_k| - d_{j,k}(x')) \geq \sum_{x' \in S_x} (|V_j||V_k| - d_{j,k}(x')) \geq |S_x| \cdot \frac{1}{2}cn^2 \geq \frac{1}{2}cn^{\frac{5}{2}},$$

contradicting Theorem 3.2. Next, we prove (c). For every $x \in V_i$, define $W_j := \{y \in V_j : d_j(x, y) > n^{3/4}\}$, $W_k := \{z \in V_k : d_k(x, z) > n^{3/4}\}$, $\bar{W}_j := V_j \setminus W_j$, and $\bar{W}_k := V_k \setminus W_k$. By the definitions of W_j and W_k , we have

$$d_{W_j, \bar{W}_j}(x), d_{\bar{W}_j, W_j}(x), d_{W_k, \bar{W}_k}(x), d_{\bar{W}_k, W_k}(x) \leq n^{3/4} \cdot n \leq n^{7/4}.$$

Further, for every $y \in W_j$, we have $d_k(x, y) \leq n^{3/4}$; otherwise either there exist $y' \in L_j(x, y)$ and $z \in L_k(x, y)$ satisfying $d_i(y', z) \geq t + 2$, and we find a copy of F_5^t in H ; or for every $y' \in L_j(x, y)$ and $z \in L_k(x, y)$, it holds that $d_i(y', z) < t + 2$, in which case

$$|\bar{H}_\pi| \geq d_j(x, y)d_k(x, y)(|V_i| - t - 2) \geq n^{3/2}(|V_i| - t - 2),$$

contradicting Theorem 3.2 by Claim 2.6 (i). We conclude $d_{W_j, W_k}(x), d_{W_j, \bar{W}_k}(x) \leq n^{7/4}$, and by a similar argument also $d_{\bar{W}_j, W_k}(x) \leq n^{7/4}$. Therefore,

$$d_{j,j}(x) + d_{j,k}(x) + d_{k,k}(x) \leq \binom{|W_j|}{2} + (|V_j| - |W_j|)(|V_k| - |W_k|) + \binom{|W_k|}{2} + 7n^{7/4}. \quad (2)$$

On the other hand, by (a), (b), and the minimum degree assumption, we have

$$d_{j,j}(x) + d_{j,k}(x) + d_{k,k}(x) \geq d(x) - d_{i,i}(x) - d_{i,j}(x) - d_{i,k}(x) \geq \left(\frac{1}{9} - 7c\right)n^2. \quad (3)$$

Combining (2) and (3) and using Claim 2.6 (i), we get

$$\frac{1}{2}|W_j|^2 + \left(\left(\frac{1}{3} + 6c\right)n - |W_j|\right) \left(\left(\frac{1}{3} + 6c\right)n - |W_k|\right) + \frac{1}{2}|W_k|^2 \geq \left(\frac{1}{9} - 8c\right)n^2,$$

which implies $|W_j| + |W_k| \leq 50cn$ or $|W_j| + |W_k| \geq \left(\frac{2}{3} - 38c\right)n$. If $|W_j| + |W_k| \geq \left(\frac{2}{3} - 38c\right)n$, then by the definition of A_i , we have

$$\begin{aligned} cn^2 &< d_{j,k}(x) = d_{\bar{W}_j, \bar{W}_k}(x) + d_{W_j, \bar{W}_k}(x) + d_{\bar{W}_j, W_k}(x) + d_{W_j, W_k}(x) \\ &\leq \left(\left(\frac{1}{3} + 6c\right)n - |W_j|\right) \left(\left(\frac{1}{3} + 6c\right)n - |W_k|\right) + 3n^{7/4} \\ &\leq (25cn)^2 + 3n^{7/4} < cn^2, \end{aligned}$$

a contradiction. We conclude $|W_j|, |W_k| \leq 50cn$ and thus

$$d_{j,j}(x) = d_{W_j, W_j}(x) + d_{W_j, \bar{W}_j}(x) + d_{\bar{W}_j, W_j}(x) \leq \binom{50cn}{2} + 2n^{7/4} \leq cn^2.$$

Similarly, $d_{k,k}(x) \leq cn^2$. Finally, (d) follows by the minimum degree assumption, (a), (b), and (c). \square

Proof of (ii). If $x, x' \in A_i$ are distinct vertices such that $d(x, x') \geq t + 1$, then by Lemma 3.1, $d_{j,k}(x) + d_{j,k}(x') \leq |V_j||V_k| + (t + 3)n \leq \frac{1}{8}n^2$. This contradicts (d) in (i). \square

Proof of (iii). Fix vertex $v \in A_i \cup A_j$ such that $xyv \in H$. Without loss of generality, we assume $v \in A_i$. By (i) and Lemma 3.8 (i), we have

$$d_{A_j, A_k}(v) \geq d_{j,k}(v) - d_{B_j, V_k}(v) - d_{V_j, B_k}(x) \geq d_{j,k}(v) - t|V_i| - t|V_j| \geq \left(\frac{1}{9} - 11c\right)n^2. \quad (4)$$

For every $z \in L_{A_k}(x, y)$, we have $d_{A_j}(v, z) \leq t + 1$; otherwise there exist $y_1, \dots, y_{t+1} \in A_j \setminus \{y\}$ and then hyperedges $\{xyv, xyz, vzy_1, \dots, vzy_{t+1}\}$ form a copy of F_5^t in H . Therefore,

$$d_{A_j, A_k}(v) \leq (t + 1)d_{A_k}(x, y) + |A_j|(|A_k| - d_{A_k}(x, y)). \quad (5)$$

Combining (4) with (5) and Claim 2.6 (i), we get $d_{A_k}(x, y) \leq 50cn$. \square

3.3 All vertices are typical

Theorem 3.10. *Let $t \geq 0$, n be sufficiently large, and H be an F_5^t -free 3-graph on vertex set $[n]$ with $|H| \geq s_t^c(n)$ and $\delta(H) \geq \frac{1}{9}n^2 - n$. Further, let $\pi = (V_1, V_2, V_3)$ be a partition of $[n]$ maximizing $|H_\pi|$, and let B be defined as in Section 3.2. Then, $|B| = 0$.*

Let $c, A_1, A_2, A_3, A, B_1, B_2$, and B_3 be defined as in Section 3.2. We write ϖ for the 3-partition with parts (A_1, A_2, A_3) , and let $A_4 := B$. For $i, j, k \in [4]$, define H_{ijk} to be the set of hyperedges $xyz \in H$ where $x \in A_i, y \in A_j$, and $z \in A_k$. Further, define $H_{bad} := \bigcup_{i, j \in [3], i \neq j} H_{iij}$. Then,

$$|H| = |H_{123}| + \sum_{i=1}^3 |H_{iii}| + |H_{bad}| + \sum_{1 \leq i \leq j \leq 3} |H_{ij4}| + \sum_{i=1}^4 |H_{i44}|. \quad (6)$$

Recall that we defined $H_\varpi = H_{123}$ and $\bar{H}_\varpi = K_\varpi \setminus H_\varpi$ in Section 2.1. Note that $H_\varpi \subseteq H_\pi$ and $\bar{H}_\varpi \subseteq \bar{H}_\pi$. For $i \in [3]$, by Claim 2.6 (i) and Lemma 3.8 (i), we have

$$\left(\frac{1}{3} - 3c\right)n - t \leq \left(\frac{1}{3} - 3c\right)n - |B| \leq |A_i| \leq |V_i| \leq \left(\frac{1}{3} + 6c\right)n. \quad (7)$$

For $i \neq j \in [3]$, let G_{ij} be an auxiliary bipartite graph on $A_i \cup A_j$, where there is an edge between $x \in A_i$ and $y \in A_j$ iff there exists a vertex $v \in (A_i \cup A_j)$ such that $xyv \in H$. Denote by $d_{G_{ij}}(x)$ the degree of vertex x in G_{ij} .

Proof of Theorem 3.10. The proof is split into several claims.

Claim 3.11. $|H_{bad}| \leq 12tc^{1/2}n^2$.

Proof. We bound $|H_{iij}|$ for every $i \neq j \in [3]$. For every $xy \in G_{ij}$, by Lemma 3.9 (iii), we have $d_{A_k}(x, y) \leq 50cn$. By Theorem 3.2, we have

$$|G_{ij}|(|A_k| - 50c) \leq |\bar{H}_\varpi| \leq |\bar{H}_\pi| \leq \left(\frac{1}{9} + c\right)tn^2,$$

implying $|G_{ij}| \leq tn$ by (7). We partition $A_i = S \cup T$, where S is the set of vertices $x \in A_i$ satisfying $d_{G_{ij}}(x) > c^{1/2}n$ and $T := A_i \setminus S$. Since $|G_{ij}| \leq tn$, we have $|S| \leq tc^{-1/2}$. By Lemma 3.9 (i), we have $d_{A_j, A_j}(x) \leq d_{j,j}(x) \leq cn^2$ for every $x \in S$. Therefore, the number of hyperedges in H_{ijj} containing a vertex in S is at most $tc^{1/2}n^2$. The number of hyperedges in H_{ijj} containing a vertex in T is at most

$$\sum_{x \in T} \binom{d_{G_{ij}}(x)}{2} \leq \sum_{x \in T} d_{G_{ij}}^2(x) \leq c^{1/2}n \sum_{x \in T} d_{G_{ij}}(x) \leq c^{1/2}n|G_{ij}| \leq tc^{1/2}n^2.$$

We conclude $|H_{ijj}| \leq 2tc^{1/2}n^2$ and thus $|H_{bad}| \leq 12tc^{1/2}n^2$. \square

Claim 3.12. *For every vertex $v \in B$, we have $d_{A_i, A_i}(v) \geq \binom{|A_i|}{2} - c^{1/3}n^2$ for $i \in [3]$.*

Proof. Assume that there exist vertex $v \in B$ and $k \in [3]$ such that $d_{A_k, A_k}(v) < \binom{|A_k|}{2} - c^{1/3}n^2$. Then, for every $i \in [3]$,

$$|H_{ii4}| \leq |B| \binom{|A_i|}{2} - \mathbb{1}_{\{i=k\}} c^{1/3}n^2 \quad \text{and} \quad 3|H_{iii}| + |H_{ii4}| \leq t \binom{|A_i|}{2},$$

where the second inequality holds by Lemma 3.9 (ii). Combining these two inequalities, we get

$$\sum_{i=1}^3 (|H_{iii}| + |H_{ii4}|) \leq \left(\frac{t}{3} + \frac{2|B|}{3} \right) \left(\sum_{i=1}^3 \binom{|A_i|}{2} \right) - \frac{2}{3} c^{1/3}n^2 \leq \frac{t}{18}n^2 + \frac{|B|}{9}n^2 - \frac{1}{2}c^{1/3}n^2, \quad (8)$$

where the last inequality holds by (7). Now we have

$$|H_{123}| \leq |A_1||A_2||A_3| \leq \left(\frac{n-|B|}{3} \right)^3 = \frac{1}{27}n^3 - \frac{1}{9}|B|n^2 + \frac{1}{9}|B|^2n - \frac{1}{27}|B|^3, \quad (9)$$

$$\sum_{1 \leq i < j \leq 3} |H_{ij4}| \leq |B| \cdot 3cn^2 \leq 3tcn^2 \quad \text{by Lemma 3.8 (i) (ii), and} \quad (10)$$

$$\sum_{i=1}^4 |H_{i44}| \leq \binom{|B|}{2} t \leq t^3 \quad \text{by Lemma 3.8 (i) (iii).} \quad (11)$$

Combining (6), (8), (9), (10), (11), and Claim 3.11, we get

$$|H| \leq \frac{1}{27}n^3 + \left(\frac{1}{18}t - \frac{1}{3}c^{1/3} \right) n^2 < s_t^\circ(n),$$

a contradiction. \square

Claim 3.13. *For vertex $v \in B$ and $i \in [3]$, let $S_{v,i} := \{x \in A_i : d_{A_i}(v, x) \leq t + 2\}$. Then $|S_{v,i}| \leq 10c^{1/3}n$.*

Proof. By the definition of $S_{v,i}$ and Claim 3.12,

$$\binom{|A_i| - |S_{v,i}|}{2} + |S_{v,i}|(t+2) \geq d_{A_i, A_i}(v) \geq \binom{|A_i|}{2} - c^{1/3}n^2,$$

so $|S_{v,i}| \leq 10c^{1/3}n$, where we also use (7). \square

Claim 3.14. *If $|B| \geq 1$, then $|H_{iii}| \leq (t+1)c^{1/3}n^2$ for $i \in [3]$.*

Proof. Let $v \in B$ and $i \in [3]$. By Claim 3.13, $|S_{v,i}| \leq 10c^{1/3}n$. For every $xx'x'' \in H_{iii}$ where $x'' \in A_i \setminus S_{v,i}$, we have $xx' \notin L_{A_i, A_i}(v)$; otherwise there exist distinct vertices $x_1, \dots, x_{t+1} \in L_{A_i}(v, x'') \setminus \{x, x'\}$, and the hyperedges $\{xx'v, xx'x'', vx''x_1, \dots, vx''x_{t+1}\}$ form an F_5^t in H , a contradiction. By Lemma 3.9 (ii), every pair in A_i has codegree at most t . By Claim 3.12, there are at most $\left(\binom{|A_i|}{2} - d_{A_i, A_i}(v)\right)t \leq tc^{1/3}n^2$ hyperedges in H_{iii} containing at least one vertex in $A_i \setminus S_{v,i}$. There are at most $\binom{|S_{v,i}|}{2}t \leq c^{1/3}n^2$ hyperedges in H_{iii} containing three vertices in $S_{v,i}$. Therefore, $|H_{iii}| \leq (t+1)c^{1/3}n^2$. \square

Claim 3.15. $|B| \notin [1, t-1]$.

Proof. Assume $|B| \in [1, t-1]$. By Claim 3.14, $\sum_{i=1}^3 |H_{iii}| \leq 3(t+1)c^{1/3}n^2$. By (7), $\sum_{i=1}^3 |H_{ii4}| \leq |B| \sum_{i=1}^3 \binom{|A_i|}{2} \leq |B|(\frac{1}{6} + 7c)n^2$. Then, by (6), (9), (10), (11), and Claim 3.11, we get

$$|H| \leq \frac{n^3}{27} + \frac{|B|n^2}{18} + 0.01n^2 < s_t^\circ(n),$$

a contradiction. \square

Claim 3.16. $|B| \neq t$.

Proof. Assume $|B| = t$. By Lemma 3.9 (ii), we have

$$\sum_{i=1}^3 |H_{iii}| + |H_{bad}| + \sum_{i=1}^3 |H_{ii4}| \leq t \sum_{i=1}^3 \binom{|A_i|}{2}. \quad (12)$$

If $\sum_{1 \leq i < j \leq 3} |H_{ij4}| \leq \frac{n}{11}$, then by (6), (11), (12), and Lemma 2.4,

$$|H| \leq |A_1||A_2||A_3| + t \sum_{i=1}^3 \binom{|A_i|}{2} + \frac{n}{11} + t^3 \leq s_{t, \varpi}^\Delta(n) + \frac{n}{10} < s_t^\circ(n),$$

a contradiction. Hence, $\sum_{1 \leq i < j \leq 3} |H_{ij4}| > \frac{n}{11}$, so there exist $v \in B$ and $i \neq j \in \{1, 2, 3\}$ such that $d_{A_i, A_j}(v) \geq \frac{n}{33t}$. For every $x \in A_i, y \in A_j$ with $xyv \in H$ and every $z \in L_{A_k}(x, y)$, we have $d_{A_k}(v, z) < t+1$; otherwise there exist distinct vertices $z_1, \dots, z_{t+1} \in L_{A_k}(v, z)$, and then hyperedges $\{xyv, xyz, vzz_1, \dots, vzz_{t+1}\}$ form a copy of F_5^t in H , a contradiction. By Claim 3.13, $d_{A_k}(x, y) \leq |S_{v,k}| \leq 10c^{1/3}n$. Thus, $|\bar{H}_\varpi| \geq \frac{n}{33t}(|A_k| - 10c^{1/3}n)$.

Let $\{i, j, k\} = [3]$. For every vertex $v \in B_i$, by Lemma 3.8 (ii), there are at least $|A_j||A_k| - cn^2$ missing crossing hyperedges in \bar{H}_π containing v and two vertices in A . By (7), we conclude

$$|\bar{H}_\pi| \geq |\bar{H}_\varpi| + t \left(\left(\frac{1}{3}n - 3cn - t \right)^2 - cn^2 \right) > \left(\frac{t}{9} + \frac{1}{100t} \right) n^2 > \left(\frac{1}{9} + c \right) tn^2,$$

contradicting Theorem 3.2. □

By Lemma 3.8 (i) and Claims 3.15 and 3.16, $|B| = 0$. □

3.4 Proof of Theorem 1.1 with a minimum degree assumption

Theorem 3.17. *For every $t \geq 0$ and sufficiently large n , let H be a 3-graph on vertex set $[n]$ with maximum number of hyperedges among the n -vertex F_5^t -free 3-graphs satisfying $\delta(H) \geq \frac{1}{9}n^2 - n$. Then, $H \in \mathcal{S}_t^\circ(n)$.*

Proof. Let $\pi = (V_1, V_2, V_3)$ be a partition of $[n]$ maximizing $|H_\pi|$, and let c, A_1, A_2, A_3, B be defined as in Section 3.2. Note that $\delta(\mathcal{S}_t^\circ(n)) \geq \delta(\mathcal{S}(n)) \geq \frac{1}{9}n^2 - n$, so $|H| \geq s_t^\circ(n)$. By Theorem 3.10, $|B| = 0$, and hence $V_i = A_i$ for $i \in [3]$. We will prove $H_{bad} = \emptyset$. Without loss of generality, we assume $|H_{112}| + |H_{221}| \geq |H_{113}| + |H_{331}| \geq |H_{223}| + |H_{332}|$.

Claim 3.18. *If $|H_{bad}| \geq 1$, then $|G_{12}| > \frac{1}{4}n^{1/2}$.*

Proof. By Lemma 2.3, we have $|H| \geq s_t^\circ(n) \geq s_{t,\pi}^\circ(n)$, and hence,

$$\sum_{i=1}^3 |H_{iii}| + |H_{bad}| = |H \setminus H_\pi| \geq |\bar{H}_\pi| + \sum_{i=1}^3 \text{ex}(|V_i|, F_2^t). \quad (13)$$

For $i \in [3]$, by Lemma 3.9 (ii), every pair of distinct vertices in A_i has codegree at most t , so $|H_{iii}| \leq \text{ex}(|V_i|, F_2^t)$. Hence, we have $|H_{bad}| \geq |\bar{H}_\pi|$. By (7) and Lemma 3.9 (iii), we have

$$|\bar{H}_\pi| \geq |G_{12}| \cdot \left(\frac{1}{3} - 3c - 50c \right) n. \quad (14)$$

If $|H_{bad}| \geq 1$, then $|H_{112}| + |H_{221}| \geq 1$ and hence $|G_{12}| \geq 1$ by the definition of G_{12} . we have

$$2 \binom{|G_{12}|}{2} \geq |H_{112}| + |H_{221}| \geq \frac{1}{3}|H_{bad}| \geq \frac{1}{3}|\bar{H}_\pi| \geq \frac{1}{3} \left(\frac{1}{3} - 53c \right) n,$$

implying $|G_{12}| > \frac{1}{4}n^{1/2}$. □

Claim 3.19. *If $|H_{bad}| \geq 1$, then there exist $x_0 \in V_1, y_0 \in V_2$ such that $d_{G_{12}}(x_0) + d_{G_{12}}(y_0) \geq \left(\frac{1}{3} - 54c \right) n$.*

Proof. By Lemma 3.9 (ii),

$$3|H_{iii}| + |H_{ijj}| + |H_{ikk}| \leq t \binom{|V_i|}{2} \quad \text{for } \{i, j, k\} = [3]. \quad (15)$$

By the definition of G_{12} ,

$$|H_{ijj}| + |H_{jjj}| \leq |H_{112}| + |H_{221}| \leq \sum_{x \in V_1} \binom{d_{G_{12}}(x)}{2} + \sum_{y \in V_2} \binom{d_{G_{12}}(y)}{2} \quad \text{for } i \neq j \in [3]. \quad (16)$$

Combining (15) and (16), we have

$$\begin{aligned} \sum_{i=1}^3 |H_{iii}| + |H_{bad}| &\leq \frac{1}{3}t \sum_{i=1}^3 \binom{|V_i|}{2} + \frac{2}{3} \cdot 3 \left(\sum_{x \in V_1} \binom{d_{G_{12}}(x)}{2} + \sum_{y \in V_2} \binom{d_{G_{12}}(y)}{2} \right) \\ &\leq \sum_{i=1}^3 \text{ex}(|V_i|, F_2^t) + \frac{n}{3} + \frac{t^2}{2} + 9t + \sum_{x \in V_1} d_{G_{12}}^2(x) + \sum_{y \in V_2} d_{G_{12}}^2(y), \end{aligned} \quad (17)$$

where for the last inequality, we use the lower bound on $\text{ex}(m, F_2^t)$ from Proposition 2.1. By (13) and (17),

$$\sum_{\{x,y\} \in G_{12}} (d_{G_{12}}(x) + d_{G_{12}}(y)) = \sum_{x \in V_1} d_{G_{12}}^2(x) + \sum_{y \in V_2} d_{G_{12}}^2(y) \geq |\bar{H}_\pi| - \frac{n}{3} - \frac{t^2}{2} - 9t. \quad (18)$$

Using (14) and (18), we get

$$\sum_{\{x,y\} \in G_{12}} (d_{G_{12}}(x) + d_{G_{12}}(y)) \geq |G_{12}| \left(\frac{1}{3} - 53c \right) n - \frac{n}{3} - \frac{t^2}{2} - 9t \geq |G_{12}| \left(\frac{1}{3} - 54c \right) n,$$

where the last inequality holds by Claim 3.18. By the pigeon-hole principle, there exist $x_0 \in V_1$, $y_0 \in V_2$ such that $d_{G_{12}}(x_0) + d_{G_{12}}(y_0) \geq (\frac{1}{3} - 54c)n$. \square

Claim 3.20. $H_{bad} = \emptyset$.

Proof. Assume for contradiction that $|H_{bad}| \geq 1$. By Claim 3.19, we can assume that there exists $x_0 \in V_1$ such that $d_{G_{12}}(x_0) \geq (\frac{1}{6} - 27c)n$. By (7) and Lemma 3.9 (iii), x_0 is contained in at least $(\frac{1}{6} - 27c)n \cdot (\frac{1}{3} - 53c)n > \frac{1}{20}n^2$ missing crossing hyperedges, contradicting Lemma 3.9 (i). Hence, $H_{bad} = \emptyset$. \square

By Claim 3.20 and Lemmas 3.9 (ii) and 2.3, we get $|H| \leq s_{t,\pi}^\circ(n) \leq s_t^\circ(n)$, where equality holds iff $H \in \mathcal{S}_t^\circ(n)$. \square

3.5 Proof of Theorem 1.1

Theorem 3.17 \Rightarrow *Theorem 1.1*. Assume that the statement of Theorem 1.1 does not hold. Then, for some $t \geq 0$, there exist infinitely many positive integers n such that there exists an n -vertex F_5^t -free hypergraph $H_n \notin \mathcal{S}_t^\circ(n)$ with $|H_n| \geq s_t^\circ(n)$. Choose a sufficiently large n among them. If H_n contains a vertex v with degree less than $\frac{1}{9}n^2 - n$, let $H_{n-1} = H_n - v$. Repeat this process until there is no such low-degree vertex. We obtain a sequence of hypergraphs H_n, H_{n-1}, \dots , where H_m is obtained from H_{m+1} by deleting a vertex in H_{m+1} with degree less than $\frac{1}{9}(m+1)^2 - (m+1)$. Note that we have $|H_{n-l}| \geq s_t^\circ(n-l) + l$, since $s_t^\circ(m) - s_t^\circ(m-1) > \frac{1}{9}m^2 - m$ for sufficiently large m by Lemma 2.2. This sequence is finite and ends with more than $n_0 = \frac{1}{100}n^{1/3}$ vertices remaining; otherwise there would be at least $n - n_0 > \binom{n_0}{3}$ hyperedges in H_{n_0} . When the process ends, we reach an \hat{n} -vertex F_5^t -free 3-graph $H_{\hat{n}}$, where $\hat{n} > n_0$, $|H_{\hat{n}}| > s_t^\circ(\hat{n})$, and the minimum degree of $H_{\hat{n}}$ is at least $\frac{1}{9}\hat{n}^2 - \hat{n}$.

Now taking n to infinity, we have that there are infinitely many integers m such that there exists an m -vertex F_5^t -free 3-graph H_m with $|H_m| > s_t^\circ(m)$ and $\delta(H_m) \geq \frac{1}{9}m^2 - m$, contradicting Theorem 3.17. \square

4 Concluding Remarks

Recall that $S(n)$ denotes the complete tripartite 3-graph on vertex set $[n]$ and $s(n) = |S(n)|$. An interesting question is to determine all 3-graphs H with $\text{ex}(n, H) = s(n)$ for sufficiently large n . However, this seems to be rather difficult. A more approachable question is the following one. Recall that we write $F_5[t]$ for the t -blow-up of F_5 . For $t \geq 1$, let \mathcal{H}^t be the family of 3-graphs F such that $F_5 \subseteq F \subseteq F_5[t]$.

Question 4.1. *For which $F \in \mathcal{H}^t$ does $\text{ex}(n, F) = s(n)$ hold for sufficiently large n ?*

Let $S'(m)$ be the 3-graph obtained from $S(m)$ by adding a hyperedge containing three vertices in V_1 , see Figure 6. Let $S''(m)$ be the 3-graph obtained from $S(m)$ by adding a hyperedge containing two vertices in V_1 and one vertex in V_2 , see Figure 7. Inspired by Simonovits's result [22] about critical graphs, one may have the natural guess that $\text{ex}(n, F) = s(n)$ for sufficiently large n iff $F \subseteq S'(m)$ and $F \subseteq S''(m)$ for some $m \geq 1$. This assumption is necessary, since $|S'(n)| = |S''(n)| = s(n) + 1$. However, it is not sufficient, as shown by the following example.

Let \hat{F} be the 3-graph with vertex set

$$\{1, 2, 3, 4, 5\} \cup \{c_{12}, c_{23}, c_{31}\} \cup \bigcup_{\substack{l=1,2 \\ i=1,2,3}} \{a_i^l, b_i^l\}$$

and hyperedge set

$$\{123, 124, 345\} \cup \{c_{12}b_1^2a_2^2, c_{23}b_2^2a_3^2, c_{31}b_3^2a_1^2\} \cup \bigcup_{i=1,2,3} \{5a_i^1b_i^1, 5a_i^1b_i^2, 5a_i^2b_i^1, 5a_i^2b_i^2\},$$




Figure 6: Hypergraph $S'(m)$.




Figure 7: Hypergraph $S''(m)$.

see Figure 8. Hence, the 3-graph \hat{F} contains F_5 and other hyperedges such that the link graph of vertex 5 contains three extra disjoint complete bipartite graphs on $\{a_i^1, a_i^2\} \cup \{b_i^1, b_i^2\}$. Besides, there are three vertices c_{12}, c_{23}, c_{31} connecting these three complete bipartite graphs.




Figure 8: Hypergraph \hat{F} .

Claim 4.2. *We have $F_5 \subseteq \hat{F} \subseteq F_5[7]$, $\hat{F} \subseteq S'(30), S''(30)$, and $\text{ex}(n, \hat{F}) > s(n)$ for sufficiently large n .*

Proof. The first two claims hold by the definition of \hat{F} . For the third claim, let $\hat{S}(n)$ be the 3-graph obtained from $S(n-1)$ by adding a new vertex v and adding hyperedges such that $L_{1,1}(v)$ and $L_{2,2}(v)$ are complete graphs $K_{|V_1|}$ and $K_{|V_2|}$ respectively, and $L_{3,3}(v)$ is an extremal $K_{2,2}$ -free graph. We have $|\hat{S}(n)| > s(n)$ for sufficiently large n , since $\text{ex}(n, K_{2,2}) \geq cn^{3/2}$ for some constant $c > 0$ by Erdős, Rényi, and Sós [4]. If there exists a copy of \hat{F} in $\hat{S}(n)$, then the vertex in $\hat{S}(n)$ corresponding to vertex 5 in \hat{F} can only be the new vertex v , and then $\{a_i^1, a_i^2, b_i^1, b_i^2\}$ is in different parts of the partition of $\hat{S}(n)$ for different i . However, $L_{3,3}(v)$ is $K_{2,2}$ -free, a contradiction. \square

We have the following sufficient condition for Question 4.1. Let $S^+(m)$ be the 3-graph obtained from $S(m-1)$ by adding a new vertex v and a hyperedge containing v and two vertices in V_1 , see Figure 9. Note that $S^+(m) \subseteq S'(m+3), S''(m+3)$. Using a cleaning method similar to the one in Section 3, we can prove the following claim; we omit its proof.

Claim 4.3. For 3-graph $F \in \mathcal{H}^t$, if $F \subseteq S^+(m)$ for some $m \geq 1$, then $\text{ex}(n, F) = s(n)$ for sufficiently large n .

However, this is not a complete answer to Question 4.1, due to the following example. Let F'_5 be the 3-graph on vertex set $\{1, 2, 3, 3', 4, 4', 5\}$ with hyperedges $\{123, 124, 345, 3'4'5\}$, see Figure 10.




Figure 9: Hypergraph $S^+(m+1)$.




Figure 10: Hypergraph F'_5 .

Claim 4.4. We have $F_5 \subseteq F'_5 \subseteq F_5[2]$, F'_5 is not a subhypergraph of $S^+(m)$ for any $m \geq 1$, and $\text{ex}(n, F'_5) = s(n)$ for every sufficiently large n .

Proof. The first claim holds by the definition of F'_5 . Assume that $S^+(m)$ contains a copy of F'_5 . Then, only the new vertex v in $S^+(m)$ can be the vertex 5 in F'_5 . However, vertex 5 has degree two in F'_5 while v has degree one in $S^+(m)$, a contradiction.

Trivially, we have $\text{ex}(n, F'_5) \geq \text{ex}(n, F_5) \geq s(n)$. For the other direction, using a similar argument as in Section 3.5, we can assume that the extremal F'_5 -free 3-graph has minimum degree at least $\frac{1}{9}n^2 - n$. For every n -vertex 3-graph H with $|H| > s(n)$, by Theorem 1.1, H contains a copy of F_5 . Let this copy be $\{v_1v_2v_3, v_1v_2v_4, v_3v_4v_5\}$. By the minimum degree assumption, v_5 is contained in a hyperedge e that does contain any v_i for $i \in [4]$. Now, $\{v_1v_2v_3, v_1v_2v_4, v_3v_4v_5, e\}$ forms a copy of F'_5 . Therefore, $\text{ex}(n, F'_5) = s(n)$ for every sufficiently large n . \square

Acknowledgments. We are very grateful to Xizhi Liu, Dhruv Mubayi, and Oleg Pikhurko for many helpful comments on our work. This project started as an IGL 2021 Summer program. The undergraduate students Junsheng Liu, Alexander Roe, Yuzhou Wang, and Zihan Zhou took part in the initial discussions. During Summer 2021, Alexander Roe was partially supported by RTG NSF grant DMS-1937241.

References

- [1] B. Bollobás. Three-graphs without two triples whose symmetric difference is contained in a third. *Discrete Math.*, 8:21–24, 1974.

- [2] W. G. Brown. On an open problem of Paul Turán concerning 3-graphs. In *Studies in pure mathematics*, pages 91–93. Birkhäuser, Basel, 1983.
- [3] M. Dehon. On the existence of 2-designs $S_\lambda(2, 3, v)$ without repeated blocks. *Discrete Math.*, 43(2–3):155–171, 1983.
- [4] P. Erdős, A. Rényi, and V. T. Sós. On a problem of graph theory. *Studia Sci. Math. Hungar.*, 1:215–235, 1966.
- [5] P. Erdős and M. Simonovits. A limit theorem in graph theory. *Studia Sci. Math. Hungar.*, 1:51–57, 1966.
- [6] P. Erdős and M. Simonovits. An extremal graph problem. *Acta Math. Acad. Sci. Hungar.*, 22:275–282, 1971/72.
- [7] P. Erdős and A. H. Stone. On the structure of linear graphs. *Bull. Amer. Math. Soc.*, 52:1087–1091, 1946.
- [8] D. G. Fon-Der-Flaass. A method for constructing $(3, 4)$ -graphs. *Mat. Zametki*, 44(4):546–550, 559, 1988.
- [9] P. Frankl and Z. Füredi. A new generalization of the Erdős-Ko-Rado theorem. *Combinatorica*, 3(3-4):341–349, 1983.
- [10] A. Frohmader. More constructions for Turán’s $(3, 4)$ -conjecture. *Electron. J. Combin.*, 15(1):Research Paper 137, 23, 2008.
- [11] S. Glock, D. Kühn, A. Lo, and D. Osthus. The existence of designs via iterative absorption: hypergraph F -designs for arbitrary F . *arXiv:1611.06827*, 2016.
- [12] W. T. Gowers. Hypergraph regularity and the multidimensional Szemerédi theorem. *Ann. of Math. (2)*, 166(3):897–946, 2007.
- [13] J. Hou, H. Li, X. Liu, D. Mubayi, and Y. Zhang. Hypergraphs with infinitely many extremal constructions. *arXiv:2206.03948*, 2022.
- [14] P. Keevash. Hypergraph Turán problems. In *Surveys in combinatorics 2011*, volume 392 of *London Math. Soc. Lecture Note Ser.*, pages 83–139. Cambridge Univ. Press, Cambridge, 2011.
- [15] P. Keevash. Counting designs. *J. Eur. Math. Soc. (JEMS)*, 20(4):903–927, 2018.
- [16] P. Keevash and D. Mubayi. Stability theorems for cancellative hypergraphs. *J. Combin. Theory Ser. B*, 92(1):163–175, 2004.
- [17] A. V. Kostochka. A class of constructions for Turán’s $(3, 4)$ -problem. *Combinatorica*, 2(2):187–192, 1982.

- [18] X. Liu and D. Mubayi. A hypergraph Turán problem with no stability. *Combinatorica*, Feb 2022.
- [19] X. Liu, D. Mubayi, and C. Reiher. Hypergraphs with many extremal configurations. *arXiv:2102.02103*, 2021.
- [20] A. A. Razborov. On the Fon-der-Flaass interpretation of extremal examples for Turán’s $(3, 4)$ -problem. *Tr. Mat. Inst. Steklova*, 274(Algoritmicheskie Voprosy Algebry i Logiki):269–290, 2011.
- [21] V. Rödl, B. Nagle, J. Skokan, M. Schacht, and Y. Kohayakawa. The hypergraph regularity method and its applications. *Proc. Natl. Acad. Sci. USA*, 102(23):8109–8113, 2005.
- [22] M. Simonovits. A method for solving extremal problems in graph theory, stability problems. In *Theory of Graphs (Proc. Colloq., Tihany, 1966)*, pages 279–319. Academic Press, New York, 1968.
- [23] P. Turán. Eine Extremalaufgabe aus der Graphentheorie. *Mat. Fiz. Lapok*, 48:436–452, 1941.

Appendix

In this appendix, we prove Proposition 2.1. The upper bound follows directly from the fact that an F_2^t -free hypergraph has maximum codegree at most t and the equality is achieved only for $(n, 3, 2, t)$ -designs. Next, we prove the lower bounds.

Let G and F be r -graphs. An (F, t) -design of G is a family \mathcal{F} of distinct copies of F in G such that every hyperedge of G is contained in exactly t of these copies. The main tool for proving Proposition 2.1 is a result by Glock, Kühn, Lo, and Osthus [11], which gives sufficient conditions on G to contain an (F, t) -design. Denote by K_n the complete graph on n vertices. Here, we will only state their result for the case we are interested in: $r = 2$ and $F = K_3$. An n -vertex graph G is (c, h, p) -typical if for every set $A \subseteq V(G)$ with $|A| \leq h$, we have $(1 - c)np^{|A|} \leq |\bigcap_{v \in A} N(v)| \leq (1 + c)np^{|A|}$, where $N(v)$ is the set of neighbors of v in G . A graph G is (K_3, t) -divisible if $t|E(G)|$ is divisible by 3 and $t|N(v)|$ is divisible by 2 for every $v \in V(G)$.

Theorem 5.1 (Glock, Kühn, Lo, and Osthus [11]). *For $h = 2812$ and every $c, p \in (0, 1]$ with $c \leq 0.9(p/2)^h / (36^2 \cdot 4^{36})$, there exist n_0 and $\gamma > 0$ such that the following holds for all $n \geq n_0$. Let t be a positive integer with $t \leq \gamma n$. Suppose that G is a (c, h, p) -typical graph on n vertices. Then, G has a (K_3, t) -design if it is (K_3, t) -divisible.*

We are ready to prove the following lemmas corresponding to the lower bounds in Proposition 2.1.

Lemma 5.2. *For every even $t \geq 1$ and sufficiently large n , we have $\text{ex}(n, F_2^t) \geq \frac{t}{3} \binom{n}{2} - \frac{2t}{3}$.*

Proof. Let $G_1 = K_n$, G_2 be the graph obtained from K_n by removing an edge and G_3 be the graph obtained from K_n by removing a matching containing two edges. Note that G_1, G_2, G_3 are all $(c, 2812, 1)$ -typical for every $c \leq 0.9(1/2)^{2812}/(36^2 \cdot 4^{36})$. We have $|E(G_1)| = |E(G_2)| + 1 = |E(G_3)| + 2$, and hence there exists $i \in \{1, 2, 3\}$ such that 3 divides $|E(G_i)|$. Therefore, G_i is (K_3, t) -divisible. By Theorem 5.1, there exists a (K_3, t) -design of G_i . The corresponding hypergraph H , i.e., the n -vertex 3-graph where a triple is a hyperedge in H if it is a K_3 in the (K_3, t) -design, satisfies that every pair of vertices has codegree at most t and $|H| \geq \frac{t}{3} \left(\binom{n}{2} - 2 \right) = \frac{t}{3} \binom{n}{2} - \frac{2t}{3}$. \square

Lemma 5.3. *For every odd $t \geq 1$ and sufficiently large odd n , we have $\text{ex}(n, F_2^t) \geq \frac{t}{3} \binom{n}{2} - \frac{8t}{3}$.*

Proof. Denote by C_4 the cycle with 4 vertices. Let $G_1 = K_n$, G_2 be the graph obtained from K_n by removing 4 edges that form a copy of C_4 and G_3 be the graph obtained from K_n by removing 8 edges that form two vertex-disjoint copies of C_4 . Note that G_1, G_2, G_3 are all $(c, 2812, 1)$ -typical for every $c \leq 0.9(1/2)^{2812}/(36^2 \cdot 4^{36})$ and every vertex in each of these three graphs has an even degree. We have $|E(G_1)| = |E(G_2)| + 4 = |E(G_3)| + 8$, and hence there exists $i \in \{1, 2, 3\}$ such that 3 divides $|E(G_i)|$. Therefore, G_i is (K_3, t) -divisible. By Theorem 5.1, there exists a (K_3, t) -design of G_i . The corresponding hypergraph H satisfies that every pair of vertices has codegree at most t and $|H| \geq \frac{t}{3} \left(\binom{n}{2} - 8 \right) = \frac{t}{3} \binom{n}{2} - \frac{8t}{3}$. \square

Claim 5.4. *For every even $n \geq 4$, there exists an n -vertex 3-graph H with $d(x, y) \geq 1$ for every $x, y \in V(H)$ and $|H| \leq \frac{1}{3} \binom{n}{2} + \frac{n}{3}$.*

Proof. Since n is even, by Theorem 1.2, there exists an $(n-1, 3, 2, 1)$ -design or an $(n+1, 3, 2, 1)$ -design. In the first case, take an $(n-1, 3, 2, 1)$ -design on $[n-1]$, and add one vertex n and hyperedges $\{2i-1, 2i, n\}$ for $i = 1, \dots, n/2 - 1$ and hyperedge $\{1, n-1, n\}$. The resulting hypergraph H has the desired properties: $d(x, y) \geq 1$ for every $x, y \in [n]$ and

$$|H| = \frac{1}{3} \binom{n-1}{2} + \frac{n}{2} = \frac{1}{3} \binom{n}{2} + \frac{n}{6} + \frac{1}{3} \leq \frac{1}{3} \binom{n}{2} + \frac{n}{3}.$$

Now, suppose there exists an $(n+1, 3, 2, 1)$ -design on $[n+1]$. Let $\{v_i u_i(n+1)\}$ be the hyperedges containing vertex $n+1$. Remove vertex $n+1$, and then add hyperedge $v_i u_i z_i$, where z_i is an arbitrary vertex, for every i . The resulting hypergraph H has the desired properties: $d(x, y) \geq 1$ for every $x, y \in [n]$ and

$$|H| \leq \frac{1}{3} \binom{n+1}{2} = \frac{1}{3} \binom{n}{2} + \frac{n}{3}. \quad \square$$

Claim 5.5. *For every odd $t \geq 1$, there exists a $(t+3)$ -vertex 3-graph H with $|H| \geq \frac{t}{3} \binom{t+3}{2} - \frac{t+3}{3}$ such that each pair of vertices has codegree at most t .*

Proof. Take a copy of the complete 3-graph on $t+3$ vertices and remove the hyperedge set of a $(t+3)$ -vertex 3-graph H' , where $|H'| \leq \frac{1}{3} \binom{t+3}{2} + \frac{t+3}{3}$ and $d(x, y) \geq 1$ for every pair of vertices x, y in H' . Such a hypergraph exists by Claim 5.4. The resulting hypergraph H has the desired properties: every pair of vertices has codegree at most t and

$$|H| \geq \binom{t+3}{3} - \left(\frac{1}{3} \binom{t+3}{2} + \frac{t+3}{3} \right) = \frac{t}{3} \binom{t+3}{2} - \frac{t+3}{3}. \quad \square$$

Lemma 5.6. *For odd $t \geq 1$ and sufficiently large even n , we have $\text{ex}(n, F_2^t) \geq \frac{t}{3} \binom{n}{2} - \frac{n}{3} - \frac{t^2}{6} - 3t$.*

Proof. Let $s = s(n) := \lfloor \frac{n}{t+3} \rfloor$. Partition $[n]$ into disjoint sets $B_1 \cup \dots \cup B_s \cup J$, where $|B_k| = t + 3$ for each $k \in [s]$ and $0 \leq |J| < t + 3$. Note that $|B_k|$ and $|J|$ are both even. Assume $J = \{w_1, \dots, w_{|J|}\}$. Define G_1 to be the n -vertex graph where $uv \notin E(G_1)$ iff $u, v \in B_k$ for some $k \in [s]$ or $\{u, v\} = \{w_{2j-1}, w_{2j}\}$ for some $j \in [|J|/2]$. Let G_2 be the graph obtained from G_1 by removing 4 edges that form a copy of C_4 , every vertex of which comes from a different set B_k . Let G_3 be the graph obtained from G_1 by removing 8 edges that form two vertex-disjoint copies of C_4 , where every vertex in these two C_4 's comes from a different set B_k . Note that G_1, G_2, G_3 are all $(c, 2812, 1)$ -typical for every $c \leq 0.9(1/2)^{2812}/(36^2 \cdot 4^{36})$ and every vertex in the graphs G_1, G_2 and G_3 has an even degree. Further, we have that $|E(G_1)| = |E(G_2)| + 4 = |E(G_3)| + 8$, and hence there exists $i \in \{1, 2, 3\}$ such that 3 divides $|E(G_i)|$. Therefore, G_i is (K_3, t) -divisible. By Theorem 5.1, there exists a (K_3, t) -design of G_i . Let H be the corresponding 3-graph. Note that if $u, v \in B_k$ for some $k \in [s]$, then $uv \notin E(G_i)$, and hence $d(u, v) = 0$ in H . We have

$$|H| \geq \frac{t}{3} \left(\binom{n}{2} - s \binom{t+3}{2} - \frac{|J|}{2} - 8 \right).$$

Now, we construct an n -vertex hypergraph H' from H by adding to every set B_k a copy of a $(t+3)$ -vertex 3-graph with at least $\frac{t}{3} \binom{t+3}{2} - \frac{t+3}{3}$ hyperedges, each pair of vertices in which has codegree at most t . Such a hypergraph exists by Claim 5.5. Then, H' satisfies that every pair of vertices has codegree at most t and

$$\begin{aligned} |H'| &\geq \frac{t}{3} \left(\binom{n}{2} - s \binom{t+3}{2} - \frac{|J|}{2} - 8 \right) + s \left(\frac{t}{3} \binom{t+3}{2} - \frac{t+3}{3} \right) \\ &= \frac{t}{3} \binom{n}{2} - \frac{t|J|}{6} - \frac{8t}{3} - \frac{s(t+3)}{3} \geq \frac{t}{3} \binom{n}{2} - \frac{t^2}{6} - 3t - \frac{n}{3}. \quad \square \end{aligned}$$