Characteristic quasi-polynomials for deformations of Coxeter arrangements of types A, B, C, and D

Yusuke Mori^{*} and Norihiro Nakashima[†]

Abstract

Kamiya, Takemura, and Terao introduced a characteristic quasi-polynomial which enumerates the numbers of elements in the complement of hyperplane arrangements modulo positive integers. In this paper, we compute the characteristic quasi-polynomials for arrangements which contain the Coxeter arrangement of types A, B, C, and D described by the orthonormal basis.

Key Words: hyperplane arrangement, characteristic quasi-polynomial, Coxeter arrangement

2020 Mathematics Subject Classification: Primary 32S22, Secondary 52C35.

1 Introduction

Let $S \in \operatorname{Mat}_{m \times n}(\mathbb{Z})$ be an $m \times n$ matrix with integral entries and $\mathbf{b} \in \mathbb{Z}^n$ an integral vector. Let $q \in \mathbb{Z}_{>0}$ and define $\mathbb{Z}_q := \mathbb{Z}/q\mathbb{Z}$. For $a \in \mathbb{Z}$, let $[a]_q := a + q\mathbb{Z} \in \mathbb{Z}_q$ be the q reduction of a. For a matrix or vector A with integral entries, let $[A]_q$ stand for the entry-wise q reduction of A. The matrix S and the vector \mathbf{b} define an arrangement \mathcal{A} of n hyperplanes in \mathbb{R}^m , while $[S]_q$ and $[\mathbf{b}]_q$ define an arrangement $\mathcal{A}_q = \mathcal{A}_q(S, b)$ in \mathbb{Z}_q^m . Indeed if $S = (\mathbf{s}_1 \dots \mathbf{s}_n)$ and $\mathbf{b} = (b_1, \dots, b_n)$ where $\mathbf{s}_1, \dots, \mathbf{s}_n$ are the columns of S, then the arrangement \mathcal{A} is the collection of hyperplanes $H_i = \{\mathbf{x} = (x_1, \dots, x_m) \in \mathbb{R}^m \mid \mathbf{x}\mathbf{s}_i = b_i\}$ for $i = 1, \dots, n$, while \mathcal{A}_q is the collection of $H_{i,q} = \{\mathbf{x} = (x_1, \dots, x_m) \in \mathbb{Z}_q^m \mid \mathbf{x}[\mathbf{s}_i]_q = [b_i]_q\}$ for $i = 1, \dots, n$. We call the case $\mathbf{b} = \mathbf{0}$ the central case, and call the case $\mathbf{b} \neq \mathbf{0}$ the non-central case. Kamiya, Takemura, and Terao [4, 7] showed that the complement $\mathcal{M}(\mathcal{A}_q)$ of \mathcal{A}_q in \mathbb{Z}_q^m is a quasi-polynomial, which is called the characteristic quasi-polynomials, in two ways. The first proof is via the theory

^{*}Department of Mathematics, Nagoya Institute of Technology, Aichi, 466-8555, Japan. Email: y.mori.404@stn.nitech.ac.jp

[†]Department of Mathematics, Nagoya Institute of Technology, Aichi, 466-8555, Japan. Email: nakashima@nitech.ac.jp (Corresponding author)

of elementary divisors. The second is via the theory of the Ehrhart quasi-polynomials which is studied by Beck and Zaslavsky [2] as an "inside-out" polytope. Kamiya, Takemura, and Terao [4, 7] also gave a period ρ_S , which is called the lcm period, of this quasi-polynomial. The lcm period ρ_S depends only on the matrix S (does not depend on the vector \boldsymbol{b} in the non-central case). In general the lcm period is not the minimum period. We say that a period collapse occurs if the minimum period is strictly less than the lcm period. There is a remarkable study about a period collapse. Higashitani, Tran, and Yoshinaga [8] constructed non-central arrangements so that the characteristic quasi-polynomial has lcm period p and has minimum period s for all nonnegative integers p and s with s|p. (This means that there exist an arrangement such that a period collapse occurs if s < p.) In the same paper, they proved that the lcm period always coincides with the minimum period in the central case.

Coxeter arrangements and their deformations (e.g., the extended Shi arrangements and the extended Linial arrangements) are interesting subjects of research on characteristic quasipolynomials. Kamiya, Takemura, and Terao [6] presented a formula for the generating function of the characteristic quasi-polynomial of a Coxeter arrangement described by the basis using root systems. This formula is essentially obtained by Athanasiadis [1] and by Blass and Sagan [3]. In the first version [5] of the paper [6], Kamiya, Takemura, and Terao explicitly determined the characteristic quasi-polynomials of Coxeter arrangements of types B, C, and D described by the orthonormal basis. Yoshinaga [11] determined the characteristic quasi-polynomial of the extended Shi arrangement and showed that it is a polynomial. The conjecture by Postnikov and Stanley [9] for the roots of the characteristic polynomials of the extended Linial arrangements was partially shown by Yoshinaga [12], using representations of the characteristic quasi-polynomials in terms of Eulerian polynomials. Later, Tamura [10] has completely resolved the conjecture by Postnikov and Stanley.

In this paper, we define two classes of matrices (and hence define two classes of arrangements). The first class of arrangements contains the Coxeter arrangement of type A described by the orthonormal basis, and the second class of arrangements contains the Coxeter arrangements of types B, C, and D described by the orthonormal basis. The characteristic quasi-polynomials of these arrangements are explicitly computed by counting the number of elements in the complement of arrangements. On the other hand, the method of computing the characteristic quasi-polynomials of the Coxeter arrangements of types B, C, and D in the paper [5] is to construct some equalities of characteristic quasi-polynomials and solve the equalities. Therefore the method of computation in this paper differs from that in the paper [5], and our main result gives another proof of the explicit formula of characteristic quasi-polynomials of types B, C, and D described by the orthonormal basis.

The organization of this paper is as follows. In Section 2, we define the characteristic quasipolynomial, while we introduce the explicit formulas of the characteristic quasi-polynomials of the Coxeter arrangements of types B, C, and D described by the orthonormal basis. In Section 3, we define the two classes of arrangements, and compute the characteristic quasi-polynomials of the arrangements.

The authors have no conflicts of interest directly relevant to the content of this article.

2 Preliminaries

In this section, we fix notations and introduce results for characteristic quasi-polynomials. A function $\chi : \mathbb{Z}_{>0} \to \mathbb{Z}$ is called a quasi-polynomial if there exist a positive integer $\rho \in \mathbb{Z}_{>0}$ and polynomials $\chi^1(t), \chi^2(t), \ldots, \chi^{\rho}(t) \in \mathbb{Z}[t]$ such that for $q \in \mathbb{Z}_{>0}$,

$$\chi(q) = \begin{cases} \chi^1(q) & q \equiv 1 \mod \rho, \\ \chi^2(q) & q \equiv 2 \mod \rho, \\ \vdots & \vdots \\ \chi^{\rho}(q) & q \equiv \rho \mod \rho. \end{cases}$$

In addition, we call χ a monic quasi-polynomial if $\chi^1(t), \chi^2(t), \ldots, \chi^{\rho}(t)$ are monic polynomials. The number ρ is called a period and the polynomials $\chi^1(t), \chi^2(t), \ldots, \chi^{\rho}(t)$ are called the *k*-constituent of the quasi-polynomial χ . A quasi-polynomial χ with a period ρ is said to have the gcd property with respect to ρ if the *k*-constituents $\chi^1(t), \chi^2(t), \ldots, \chi^{\rho}(t)$ depend on *k* only through $gcd(\rho, k)$, i.e., $\chi^a(t) = \chi^b(t)$ if $gcd(\rho, a) = gcd(\rho, b)$.

We next define the lcm period and the characteristic quasi-polynomial. In this paper, we consider only the central case. Let $S \in \operatorname{Mat}_{m \times n}(\mathbb{Z})$ be an $m \times n$ matrix with integral entries. For $\emptyset \neq J \subseteq \{1, \ldots, n\}$, the matrix $S_J \in \operatorname{Mat}_{m \times |J|}(\mathbb{Z})$ is defined by the submatrix of S consisting of the columns indexed by J. Set $\ell(J) := \operatorname{rank}(S_J)$. Let $e_{J,1}|e_{J,2}|\cdots|e_{J,\ell(J)}$ be the elementary divisors of S_J . Then the lcm period of S is defined by

$$\rho_S := \operatorname{lcm}\left(e_{J,\ell(J)} \mid \emptyset \neq J \subseteq \{1,\ldots,n\}\right).$$

For $q \in \mathbb{Z}_{>0}$, let $\mathbb{Z}_q^{\times} := \mathbb{Z}_q \setminus \{0\}$ and we define

$$M_S(q) := \left\{ \boldsymbol{x} = (x_1, \dots, x_m) \in \mathbb{Z}_q^m \, \big| \, \boldsymbol{x}[S]_q \in \left(\mathbb{Z}_q^{\times}\right)^n \right\}.$$

Then $M_S(q)$ is the complement of the arrangement defined by $[S]_q$.

Theorem 2.1 (Kamiya, Takemura, and Terao [4, 7]). The function $|M_S(q)|$ is a monic quasipolynomial with a period ρ_S having the gcd property with respect to ρ_S .

The quasi-polynomial in Theorem 2.1 is called the characteristic quasi-polynomial of S (or of the arrangement defined by S), and we write $\chi_S^{quasi}(q) = |M_S(q)|$. The next theorem states that we can obtain the minimum period by computing the lcm period in the central case¹.

¹There exist examples of arrangements such that the lcm period is not the minimum period in the noncentral case (see Theorem 1.1 in [8])

Theorem 2.2 (Higashitani, Tran, and Yoshinaga [8]). The lcm period of $\chi_S^{quasi}(q)$ coincides with the minimum period of $\chi_S^{quasi}(q)$ in the central case.

Let $e_i \in \mathbb{Z}^m$ be the column vector with 1 in the *i*th entry and 0's elsewhere.

Definition 2.3. The following matrices A_m , B_m , C_m , and D_m define Coxeter arrangements of types A, B, C, and D described by the orthonormal basis.

- Let A_m be the $m \times \binom{m}{2}$ matrix consisting of the columns $e_i e_j$ $(1 \le i < j \le m)$.
- Let B_m be the $m \times m^2$ matrix consisting of the columns e_i $(1 \le i \le m)$ and $e_i e_j$, $e_i + e_j$ $(1 \le i < j \le m)$.
- Let C_m be the $m \times m^2$ matrix consisting of the columns $2\mathbf{e}_i$ $(1 \le i \le m)$ and $\mathbf{e}_i \mathbf{e}_j$, $\mathbf{e}_i + \mathbf{e}_j$ $(1 \le i < j \le m)$.
- Let D_m be the $m \times (m^2 m)$ matrix consisting of the columns $e_i e_j$, $e_i + e_j$ $(1 \le i < j \le m)$.

Kamiya, Takemura, and Terao computed the characteristic quasi-polynomials for A_m , B_m , C_m , and D_m in section 4.5 in the paper [5] (the first version of the published paper [6]).

Theorem 2.4 (Kamiya, Takemura, and Terao [5]). The minimum periods and the characteristic quasi-polynomials for A_m , B_m , C_m , and D_m are as follows.

$$\rho_{A_m} = 1,
\chi_{A_m}^{quasi}(q) = \chi_{A_m}^1(q) = \prod_{i=1}^m (q - i + 1), \quad \text{for } q \in \mathbb{Z}_{>0}.$$

$$\rho_{B_m} = 2,$$
(2.1)

$$\chi_{B_m}^{quasi}(q) = \begin{cases} \chi_{B_m}^1(q) = \prod_{i=1}^m (q-2i+1) & \text{if } q \text{ is odd,} \\ \chi_{B_m}^2(q) = (q-m) \prod_{i=1}^{m-1} (q-2i) & \text{if } q \text{ is even.} \end{cases}$$
(2.2)

$$\rho_{C_m} = 2,
\chi_{C_m}^{quasi}(q) = \begin{cases} \chi_{C_m}^1(q) = \prod_{i=1}^m (q - 2i + 1) & \text{if } q \text{ is odd,} \\ \chi_{C_m}^2(q) = \prod_{i=1}^m (q - 2i) & \text{if } q \text{ is even.} \end{cases}$$

$$\rho_{D_m} = 2,$$
(2.3)

$$\chi_{D_m}^{quasi}(q) = \begin{cases} \chi_{D_m}^1(q) = (q - m + 1) \prod_{i=1}^{m-1} (q - 2i + 1) & \text{if } q \text{ is odd,} \\ \chi_{D_m}^2(q) = (q^2 - 2(m - 1)q + m(m - 1)) \prod_{i=1}^{m-2} (q - 2i) & \text{if } q \text{ is even.} \end{cases}$$
(2.4)

3 Characteristic quasi polynomials for deformation of Coxeter arrangements

3.1 Deformation of type A

Let $t \in \mathbb{Z}_{\geq 0}$ with $0 \leq t \leq m$, and let s_1, \ldots, s_t be positive integers with $s_t | \ldots | s_1$. Then we define a tuple s by $s := (s_1, \ldots, s_t)$. We define an $m \times \frac{m^2 - m + 2t}{2}$ matrix $A_m(s)$ by adding the columns $s_1 e_1, \ldots, s_t e_t$ to A_m , that is,

$$A_m(\boldsymbol{s}) := egin{pmatrix} s_1 & \cdots & 0 & \ dots & \ddots & dots & \ 0 & \cdots & s_t & A_m \ dots & & dots & \ 0 & \cdots & 0 & \end{pmatrix}$$

We note that if t = 0, then $A_m(\mathbf{s}) = A_m$ and $\rho_{A_m} = 1$. If $t \neq 0$, then $\rho_{A_m(\mathbf{s})} = s_1$. By Theorem 2.2, the minimum period of $\chi^{quasi}_{A_m(\mathbf{s})}(q)$ is $\begin{cases} s_1 & \text{if } t \neq 0, \\ 1 & \text{if } t = 0. \end{cases}$

Let k be a positive integer with $k|\rho_{A_m(s)}$. When $t \ge 1$, we define $d_i := \gcd(k, s_i)$ for $1 \le i \le t$. For $q \in k + \rho_{A_m(s)}\mathbb{Z}_{\ge 0}$, we have $d_i = \gcd(k, s_i) = \gcd(q, \rho_{A_m(s)}, s_i) = \gcd(q, s_i)$ since $s_i|\rho_{A_m(s)}$.

Lemma 3.1. Let $t \ge 1$. Let $q \in k + \rho_{A_m(s)} \mathbb{Z}_{\ge 0}$, and let h be an integer with $1 \le h \le t$. We assume that $x_1, \ldots, x_{h-1} \in \mathbb{Z}_q$ satisfy

$$s_i x_i \neq 0 \quad \text{for} \quad 1 \le i \le h - 1, \tag{3.1}$$

$$x_i - x_j \neq 0$$
 for $1 \le i < j \le h - 1$. (3.2)

Then we have

$$\#\left(\{x \in \mathbb{Z}_q \mid s_h x = 0\} \bigcup \{x_1, \dots, x_{h-1}\}\right) = d_h + h - 1.$$

Proof. By the inclusion–exclusion principle,

$$\#\left(\{x \in \mathbb{Z}_q \mid s_h x = 0\} \bigcup \{x_1, \dots, x_{h-1}\}\right)$$

=#\{x \in \mathbb{Z}_q \mathbb{Z}_q \mathbf{S}_h x = 0\} + \#\{x_1, \ldots, x_{h-1}\} - \#\left(\{x \in \mathbb{Z}_q \mathbf{S}_n x = 0\}\left(\{x_1, \ldots, x_{h-1}\}\right).

Since $gcd(q, s_h) = d_h$ and $\{x \in \mathbb{Z}_q \mid s_h x = 0\} = \left\{0, \frac{q}{d_h}, \frac{2q}{d_h}, \dots, \frac{(d_h - 1)q}{d_h}\right\}$, we have $\#\{x \in \mathbb{Z}_q \mid s_h x = 0\} = d_h$. By the condition (3.2), we have $\#\{x_1, \dots, x_{h-1}\} = h - 1$. To prove the assertion, it remains to prove that $\#(\{x \in \mathbb{Z}_q \mid s_h x = 0\} \cap \{x_1, \dots, x_{h-1}\}) = 0$. Assume that $\#(\{x \in \mathbb{Z}_q \mid s_h x = 0\} \cap \{x_1, \dots, x_{h-1}\}) \neq 0$. Then there exists i < h such that $s_h x_i = 0$. Since $s_h \mid s_i$, we can write $s_i = ls_h$ for some $l \in \mathbb{Z}$. Thus $s_i x_i = ls_h x_i = 0$. This is in contradiction to the condition (3.1). Therefore we have $\#(\{x \in \mathbb{Z}_q \mid s_h x = 0\} \cap \{x_1, \dots, x_{h-1}\}) = 0$, while $\#(\{x \in \mathbb{Z}_q \mid s_h x = 0\} \cup \{x_1, \dots, x_{h-1}\}) = d_h + h - 1$.

We write $\prod_{i=a}^{b} f_i = 1$ for integers $f_i \ (i \in \mathbb{Z})$ if a > b.

Theorem 3.2. Let $k | \rho_{A_m(s)}$. For $q \in k + \rho_{A_m(s)} \mathbb{Z}_{\geq 0}$, we have

$$\chi_{A_m(s)}^k(q) = \prod_{i=1}^t (q - d_i - i + 1) \prod_{i=t+1}^m (q - i + 1).$$
(3.3)

Proof. If t = 0, then k = 1 and $q \in \mathbb{Z}_{>0}$ since $\rho_{A_m(s)} = 1$. The assertion follows from Theorem 2.4.

Let $t \ge 1$. The value $\chi^k_{A_m(s)}(q) = |M_{A_m(s)}(q)|$ is the number of elements $\boldsymbol{x} = (x_1, \ldots, x_m) \in \mathbb{Z}_q^m$ with

$$\begin{cases} s_i x_i \neq 0, & \text{for } 1 \leq i \leq t, \\ x_i - x_j \neq 0, & \text{for } 1 \leq i < j \leq m. \end{cases}$$
(3.4)

We enumerate the elements $x_i \in \mathbb{Z}_q$ $(1 \le i \le m)$ from the first entry x_1 to the last entry x_m so that x_1, \ldots, x_m satisfy the condition (3.4). We first consider x_1 . Since $\#\{x_1 \in \mathbb{Z}_q \mid s_1x_1 = 0\} = \gcd(q, s_1) = d_1$, the number of elements $x_1 \in \mathbb{Z}_q$ with $s_1x_1 \neq 0$ is $q - d_1$. Next let $2 \le h \le t$, and we take $x_1, \ldots, x_{h-1} \in \mathbb{Z}_q$ with

$$\begin{cases} s_i x_i \neq 0, & \text{for } 1 \leq i \leq h-1, \\ x_i - x_j \neq 0 & \text{for } 1 \leq i < j \leq h-1 \end{cases}$$

Then x_1, \ldots, x_h satisfy the condition (3.4) if and only if $s_h x_h \neq 0$ and $x_h \neq x_l$ for $l = 1, \ldots, h-1$. By Lemma 3.1,

$$\# \left\{ \{ x_h \in \mathbb{Z}_p \mid s_h x_h \neq 0 \} \bigcap \{ x_h \in \mathbb{Z}_q \mid x_h \neq x_l, \text{ for } l < h \} \right\}$$
$$= q - \# \left\{ \{ x_h \in \mathbb{Z}_p \mid s_h x_h = 0 \} \bigcup \{ x_1, \dots, x_{h-1} \} \right\}$$
$$= q - d_h - h + 1.$$

Therefore the number of elements $(x_1, \ldots, x_t) \in \mathbb{Z}_q^t$ with the condition (3.4) is

$$\prod_{i=1}^{t} (q - d_i - i + 1) = (q - d_1)(q - d_2 - 1) \cdots (q - d_t - t + 1).$$

Finally let $t < h \le m$. After taking x_1, \ldots, x_{h-1} , we can take x_h from $\mathbb{Z}_q \setminus \{x_1, \ldots, x_{h-1}\}$. Hence the number of elements $x_h \in \mathbb{Z}_q$ so that x_1, \ldots, x_h satisfy the condition (3.4) is $q - \# \{x_l \mid l < h\} = q - h + 1$. Hence we have $\chi^k_{A_m(s)}(q) = (q - d_1)(q - d_2 - 1) \cdots (q - d_t - t + 1)(q - t) \cdots (q - m + 1) = \prod_{i=1}^t (q - d_i - i + 1) \prod_{i=t+1}^m (q - i + 1)$.

Example 3.3. Let t = 0. Then $A_m(s) = A_m$ and $\rho_{A_m} = 1$. If $k | \rho_{A_m}$, then k = 1. By substituting t = 0 to the equality (3.3), we have

$$\chi^{1}_{A_{m}}(q) = \prod_{i=1}^{m} (q-i+1)$$

and it coincides with the equality (2.1).

3.2 Deformation of type D

Let $r, t \in \mathbb{Z}_{\geq 0}$ with $0 \leq r \leq t \leq m$. We take positive integers s_1, \ldots, s_t , where s_1, \ldots, s_r are even, s_{r+1}, \ldots, s_t are odd, and $s_t | \ldots | s_r | \ldots | s_1$. Let $\mathbf{s} := (s_1, \ldots, s_t)$. Then we define an $m \times (m^2 - m + t)$ matrix $D_m(\mathbf{s})$ by adding the columns $s_1 \mathbf{e}_1, \ldots, s_t \mathbf{e}_t$ to D_m , that is,

$$D_m(\boldsymbol{s}) = \begin{pmatrix} s_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & s_t & D_m \\ \vdots & & \vdots \\ 0 & \cdots & 0 \end{pmatrix}.$$

We note that the matrices B_m , C_m , and D_m are described as follows:

$$r = 0, \ t = m, \text{ and } s_i = 1 \ (i = 1, \dots, t) \quad \Rightarrow \quad D_m(s) = B_m,$$

$$r = t = m \text{ and } s_i = 2 \ (i = 1, \dots, t) \quad \Rightarrow \quad D_m(s) = C_m,$$

$$r = t = 0 \quad \Rightarrow \quad D_m(s) = D_m$$

By Theorem 2.2, the minimum period of $\chi_{D_m(s)}^{quasi}(q)$ is $\rho_{D_m(s)} = \begin{cases} \operatorname{lcm}(s_1, 2) & \text{if } t \neq 0, \\ 2 & \text{if } t = 0. \end{cases}$

Let k be a positive integer with $k|\rho_{D_m(s)}$ and let $q \in k + \rho_{D_m(s)}\mathbb{Z}_{\geq 0}$. When $t \geq 1$, we define $d_i := gcd(k, s_i)$ for $1 \leq t \leq t$. Then $d_i = gcd(k, s_i) = gcd(q, \rho_{D_m(s)}, s_i) = gcd(q, s_i)$ since $s_i|\rho_{D_m(s)}$. In addition, since $\rho_{D_m(s)}$ is even, k is even if and only if q is even.

Lemma 3.4. Let $t \ge 1$. Let $q \in k + \rho_{D_m(s)} \mathbb{Z}_{\ge 0}$ and let h be an integer with $1 \le h \le t$. We assume that $x_1, \ldots, x_{h-1} \in \mathbb{Z}_q$ satisfy

$$s_i x_i \neq 0 \quad \text{for} \quad 1 \le i \le h - 1, \tag{3.5}$$

$$x_i - x_j \neq 0$$
 for $1 \le i < j \le h - 1$, (3.6)

$$x_i + x_j \neq 0$$
 for $1 \le i < j \le h - 1$. (3.7)

In addition, if k is even and $r+1 \le h-1$, then we assume that $x_i \ne \frac{q}{2}$ for $r+1 \le i \le h-1$. Then we have

$$\#\left(\{x \in \mathbb{Z}_q \mid s_h x = 0\} \bigcup \{x_i, -x_i \mid 1 \le i \le h - 1\}\right) = d_h + 2h - 2.$$

Proof. By the inclusion–exclusion principle,

$$\# \left(\{ x \in \mathbb{Z}_q \mid s_h x = 0 \} \bigcup \{ x_i, -x_i \mid 1 \le i \le h - 1 \} \right) \\
= \# \{ x \in \mathbb{Z}_q \mid s_h x = 0 \} + \# \{ x_i, -x_i \mid 1 \le i \le h - 1 \} \\
- \# \left(\{ x \in \mathbb{Z}_q \mid s_h x = 0 \} \bigcap \{ x_i, -x_i \mid 1 \le i \le h - 1 \} \right).$$

Here we have $\#\{x \in \mathbb{Z}_q \mid s_h x = 0\} = d_h$ and $\#(\{x \in \mathbb{Z}_q \mid s_h x = 0\} \bigcap \{x_i, -x_i \mid 1 \le i \le h-1\}) = 0$ similarly to Lemma 3.1. It remains to prove the assertion that $\#\{x_i, -x_i \mid 1 \le i \le h-1\} = 2h-2$. By the conditions (3.6) and (3.7), we have $x_i \ne \pm x_1, \ldots, x_i \ne \pm x_{i-1}$ for $1 \le i \le h-1$. To prove $\#\{x_i, -x_i \mid 1 \le i \le h-1\} = 2h-2$, we prove that $x_i \ne -x_i$ for $1 \le i \le h-1$ by a case study. Let $1 \le i \le h-1$.

- When k is odd ($\Leftrightarrow q$ is odd), it holds that $\{x \in \mathbb{Z}_q \mid x = -x\} = \{0\}$. By the condition (3.5), we have $x_i \neq 0$, and hence $x_i \neq -x_i$.
- When k is even ($\Leftrightarrow q$ is even) and $i \leq r$, it holds that $\{x \in \mathbb{Z}_q \mid x = -x\} = \{0, \frac{q}{2}\}$ while s_i is even. If $x_i = 0$, then $s_i x_i = 0$. If $x_i = \frac{q}{2}$, then $s_i x_i = q \frac{s_i}{2} = 0$. These are in contradiction to the condition (3.5). Therefore $x_i \neq -x_i$.
- When k is even and $r+1 \le i \le h-1$, we have $x_i \ne -x_i$ by the assumption $x_i \ne \frac{q}{2}$ for $r+1 \le i \le h-1$.

Thus we have $\#\{x_i, -x_i \mid 1 \le i \le h-1\} = 2h-2$, and hence $\#(\{x_h \in \mathbb{Z}_q \mid s_h x = 0\} \bigcup \{x_i, -x_i \mid 1 \le i \le h-1\}) = d_h + 2h - 2$.

We write
$$\sum_{i=a}^{b} f_i = 0$$
 and $\prod_{i=a}^{b} f_i = 1$ for integers f_i $(i \in \mathbb{Z})$ if $a > b$.

Theorem 3.5. Let $k | \rho_{D_m(s)}$. If k is odd, then

$$\chi_{D_m(s)}^k(q) = \prod_{i=1}^t (q - d_i - 2i + 2) \left(\prod_{i=t+1}^m (q - 2i + 1) + (m - t) \prod_{i=t+1}^{m-1} (q - 2i + 1) \right)$$
(3.8)

for $q \in k + \rho_{D_m(s)} \mathbb{Z}_{\geq 0}$. If k is even, then

$$\chi_{D_m(s)}^k(q) = \prod_{i=1}^r (q - d_i - 2i + 2)(P_1 + P_2), \qquad (3.9)$$

where

$$P_{1} := \prod_{i=r+1}^{t} (q - d_{i} - 2i + 1) \left(\prod_{i=t+1}^{m} (q - 2i) + 2(m - t) \prod_{i=t+1}^{m-1} (q - 2i) + (m - t)(m - t - 1) \prod_{i=t+1}^{m-2} (q - 2i) \right),$$

$$P_{2} := \left(\sum_{i=r+1}^{t} \prod_{j=1}^{i-1} (q - d_{j} - 2j + 1) \prod_{j=i+1}^{t} (q - d_{j} - 2j + 3) \right) \left(\prod_{i=t+1}^{m} (q - 2i + 2) + (m - t) \prod_{i=t+1}^{m-1} (q - 2i + 2) \right),$$

for $q \in k + \rho_{D_m(s)} \mathbb{Z}_{\geq 0}$.

In particular, let t = m. If k is odd, then

$$\chi^k_{D_m(s)}(q) = \prod_{i=1}^m (q - d_i - 2i + 2)$$

for $q \in k + \rho_{D_m(s)} \mathbb{Z}_{\geq 0}$. If k is even, then

$$\chi_{D_m(s)}^k(q) = \prod_{i=1}^r (q - d_i - 2i + 2) \left(\prod_{i=r+1}^t (q - d_i - 2i + 1) + \sum_{i=r+1}^m \prod_{j=r+1}^{i-1} (q - d_j - 2j + 1) \prod_{j=i+1}^m (q - d_j - 2j + 3) \right)$$

for $q \in k + \rho_{D_m(s)} \mathbb{Z}_{\geq 0}$.

Proof. The value $\chi^k_{D_m(s)}(q)$ is the number of elements $\boldsymbol{x} = (x_1, \ldots, x_m) \in \mathbb{Z}_q^m$ with

$$\begin{cases} s_i x_i \neq 0 & \text{for } 1 \leq i \leq t, \\ x_i - x_j \neq 0 & \text{for } 1 \leq i < j \leq m, \\ x_i + x_j \neq 0 & \text{for } 1 \leq i < j \leq m. \end{cases}$$
(3.10)

To prove the assertion, we enumerate the elements $x_i \in \mathbb{Z}_q$ $(1 \le i \le m)$ from the first entry x_1 to the last entry x_m so that x_1, \ldots, x_m satisfy the condition (3.10).

First let k be odd. By Lemma 3.4, the number of elements $(x_1, \ldots, x_t) \in \mathbb{Z}_q^t$ with the condition (3.10) is

$$(q-d_1)(q-d_2-2)\cdots(q-d_t-2t+2) = \prod_{i=1}^t (q-d_i-2i+2).$$

Here we fix (x_1, \ldots, x_t) , and determine the number of elements $(x_{t+1}, \ldots, x_m) \in \mathbb{Z}_q^{m-t}$ with the condition (3.10) by the following case study.

- When $x_i \neq 0$ for all $t+1 \leq i \leq m$, the number of $(x_{t+1}, \dots, x_m) \in \mathbb{Z}_q^{m-t}$ with the condition (3.10) is $(q-1-2t)(q-1-2t+2)\cdots(q-1-2m+2) = \prod_{i=t+1}^m (q-2i+1)$.
- When there exists $t+1 \leq i \leq m$ such that $x_i = 0$, such i is unique by the condition (3.10). We fix an index i with $x_i = 0$. Then the number of elements $(x_{t+1}, \ldots, x_{i-1}, x_{i+1}, \ldots, x_m) \in \mathbb{Z}_q^{m-t-1}$ with the condition (3.10) is $(q-1-2t)(q-1-2t+2)\cdots(q-1-2m+4)$. Since there is (m-t) ways to select such index i, the number of $(x_{t+1}, \ldots, x_m) \in \mathbb{Z}_q^{m-t}$ with the condition (3.10) is $(m-t)(q-1-2t)(q-1-2t+2)\cdots(q-1-2m+4) = (m-t)\prod_{i=1}^{m-1} (q-2i+1)$.

From the argument above, we have

$$\chi_{D_m(s)}^k(q) = \prod_{i=1}^t (q - d_i - 2i + 2) \left(\prod_{i=t+1}^m (q - 2i + 1) + (m - t) \prod_{i=t+1}^{m-1} (q - 2i + 1) \right).$$

Next let k be even. We recall that q is even in this case. By Lemma 3.4, the number of elements $(x_1, \ldots, x_r) \in \mathbb{Z}_q^t$ with the condition (3.10) is

$$(q-d_1)(q-d_2-2)\cdots(q-d_t-2r+2) = \prod_{i=1}^r (q-d_i-2i+2).$$

We fix (x_1, \ldots, x_r) , and determine the number of elements $(x_{r+1}, \ldots, x_m) \in \mathbb{Z}_q^{m-t}$ with the condition (3.10) by a case study. We note that $x_1 \notin \{0, \frac{q}{2}\}, \ldots, x_r \notin \{0, \frac{q}{2}\}, x_{r+1} \neq 0, \ldots, x_t \neq 0$ since s_1, \ldots, s_r are even.

since s_1, \ldots, s_r are even. (i) When $\frac{q}{2}$ is not contained in $\{x_{r+1}, \ldots, x_t\}$, the number of (x_{r+1}, \ldots, x_t) 's is

$$(q-1-d_{r+1}-2r)(q-1-d_{r+2}-2r-2)\cdots(q-1-d_t-2t+2) = \prod_{i=r+1}^t (q-d_i-2i+1)$$

by Lemma 3.4. Furthermore we fix (x_{r+1}, \ldots, x_t) in this setting, and determine the number of (x_{t+1}, \ldots, x_m) 's.

- (i-a) When both 0 and $\frac{q}{2}$ are not contained in $\{x_{t+1},\ldots,x_m\}$, we can take x_i from $\mathbb{Z}_q \setminus$ $\left\{0, \frac{q}{2}, \pm x_1, \dots, \pm x_{i-1}\right\}$ for $t+1 \le i \le m$. Therefore the number of (x_{t+1}, \dots, x_m) 's is $\prod_{i=t+1} (q-2-2i+2) = \prod_{i=t+1}^{m} (q-2i).$
- (i-b) When either 0 or $\frac{q}{2}$ is contained in $\{x_{t+1}, \ldots, x_m\}$, we select an index from $\{t+1, \ldots, m\}$ so that $x_i \in \left\{0, \frac{q}{2}\right\}$, and there is (m-t) ways to select such index *i*. We may assume that $x_m \in \left\{0, \frac{q}{2}\right\}$ after selecting such index. The other elements x_j can be taken from $\mathbb{Z}_q \setminus \left\{0, \frac{q}{2}, \pm x_1, \dots, \pm x_{j-1}\right\}$. Hence the number of (x_{t+1}, \dots, x_m) 's is $2(m-t) \prod_{i=t+1} (q-t)$ $2 - 2i + 2) = 2(m - t) \prod_{i=t+1}^{m-1} (q - 2i).$

(i-c) When both 0 and
$$\frac{q}{2}$$
 are contained in $\{x_{t+1}, \ldots, x_m\}$, the number of indices j, l with $x_j = 0, x_l = \frac{q}{2}$ is $(m-t)(m-t-1)$. Similarly to the case (i-b), the number of (x_{t+1}, \ldots, x_m) 's is $(m-t)(m-t-1) \prod_{i=t+1}^{m-2} (q-2-2i+2) = (m-t)(m-t-1) \prod_{i=t+1}^{m-2} (q-2i)$.

Hence in the case (i), the number of (x_{r+1}, \ldots, x_m) 's is P_1 . (ii) When $\frac{q}{2}$ is contained in $\{x_{r+1}, \ldots, x_t\}$, the index *i* with $x_i = \frac{q}{2}$ is unique by the condition (3.10). If $r+1 \leq j \leq i-1$, then we can take $x_j \in \mathbb{Z}_q \setminus \left\{\frac{q}{2}\right\}$ with $s_j x_j \neq 0$, $x_j \neq \pm x_1, \ldots, x_j \neq \pm x_{j-1}$. If $i+1 \le j \le t$, then we can take $x_j \in \mathbb{Z}_q \setminus \left\{\frac{q}{2}\right\}$ with $s_j x_j \neq 0$, $x_j \neq \pm x_1, \ldots, x_j \neq \pm x_{i-1}, x_j \neq \pm x_{i+1}, \ldots, x_j \neq \pm x_{j-1}$. Therefore by Lemma 3.4, the number of (x_{r+1}, \ldots, x_t) 's is

$$\sum_{i=r+1}^{t} \prod_{j=r+1}^{i-1} (q-1-d_j-2j+2) \prod_{j=i+1}^{t} (q-1-d_j-2j+4)$$
$$= \sum_{i=r+1}^{t} \prod_{j=r+1}^{i-1} (q-d_j-2j+1) \prod_{j=i+1}^{t} (q-d_j-2j+3).$$

Furthermore we fix (x_{r+1}, \ldots, x_t) in this setting, and determine the number of (x_{t+1}, \ldots, x_m) 's. We note that $x_{t+1} \neq \frac{q}{2}, \ldots, x_m \neq \frac{q}{2}$ since $\frac{q}{2} \in \{x_{r+1}, \ldots, x_t\}$.

(ii-a) When $0 \notin \{x_{t+1}, \dots, x_m\}$, we fix an index $r+1 \le i \le t$ with $x_i = \frac{q}{2}$. Then we can take x_j from $\mathbb{Z}_q \setminus \{0, \frac{q}{2}, \pm x_1, \dots, \pm x_{i-1}, \pm x_{i+1}, \dots, \pm x_{j-1}\}$ for $t+1 \le j \le m$. Therefore the number of (x_{t+1}, \dots, x_m) 's is $(q-2-2t+2)(q-2-2t)\cdots(q-2-2m+4) = \prod_{i=t+1}^m (q-2i+2)$.

(ii-b) When $0 \in \{x_{t+1}, \dots, x_m\}$, the number of (x_{t+1}, \dots, x_m) 's is $(m-t)(q-2-2t+2)(q-2-2t)\cdots(q-2-2m+6) = (m-t)\prod_{i=t+1}^{m-1}(q-2i+2).$

In the case (ii), the number of (x_{r+1}, \ldots, x_m) 's is P_2 . Hence we have

$$\chi_{D_m(s)}^k(q) = \prod_{i=1}^r (q - d_i - 2i + 2)(P_1 + P_2).$$

We can compute the characteristic quasi-polynomials of B_m , C_m , and D_m from Theorem 3.5.

Example 3.6. $[B_m]$ Let r = 0, t = m, and $s_1 = \cdots = s_m = 1$. Then we have that $D_m(\mathbf{s}) = B_m$, $\rho_{B_m} = \operatorname{lcm}(s_1, 2) = 2$, and $d_1 = \cdots = d_m = 1$. By the equalities (3.8) and (3.9), we have

$$\chi_{B_m}^1(q) = \prod_{i=1}^m (q-1-2i+2) = \prod_{i=1}^m (q-2i+1),$$

$$\chi_{B_m}^2(q) = \prod_{i=1}^m (q-1-2i+1) + \sum_{i=1}^m \prod_{j=1}^{i-1} (q-1-2j+1) \prod_{j=i+1}^m (q-1-2j+3)$$

$$= \prod_{i=1}^m (q-2i) + \sum_{i=1}^m \prod_{j=1}^{i-1} (q-2j) \prod_{j=i+1}^m (q-2(j-1)) = \prod_{i=1}^m (q-2i) + \sum_{i=1}^m \prod_{j=1}^{m-1} (q-2j)$$

$$= \prod_{i=1}^m (q-2i) + m \prod_{j=1}^{m-1} (q-2j) = (q-m) \prod_{i=1}^{m-1} (q-2i).$$

 $[C_m]$ Let r = t = m, and $s_1 = \cdots = s_m = 2$. Then we have $D_m(s) = C_m$ and $\rho_{C_m} = \operatorname{lcm}(s_1, 2) = 2$. In the case when k = 1, we have $d_1 = \cdots = d_m = 1$. By the equality (3.8), we have

$$\chi^{1}_{C_{m}}(q) = \prod_{i=1}^{m} (q-1-2i+2) = \prod_{i=1}^{m} (q-2i+1).$$

In the case when k = 2, we have $d_1 = \cdots = d_m = 2$. By the equality (3.9), we have

$$\chi^2_{C_m}(q) = \prod_{i=1}^m (q-2-2i+2) = \prod_{i=1}^m (q-2i).$$

 $[D_m]$ Let r = t = 0. Then we have $D_m(s) = D_m$ and $\rho_{D_m} = 2$. By the equality (3.8) and (3.9), we have

$$\chi^{1}_{D_{m}}(q) = \prod_{i=1}^{m} (q-2i+1) + m \prod_{i=1}^{m-1} (q-2i+1) = (q-m+1) \prod_{i=1}^{m-1} (q-2i+1),$$

$$\chi^{2}_{D_{m}}(q) = \prod_{i=1}^{m} (q-2i) + 2m \prod_{i=1}^{m-1} (q-2i) + m(m-1) \prod_{i=1}^{m-2} (q-2i)$$

$$= ((q-2m)(q-2m+2) + 2m(q-2m+2) + m(m-1)) \prod_{i=1}^{m-2} (q-2i)$$

$$= (q^{2} - 2(m-1)q + m(m-1)) \prod_{i=1}^{m-2} (q-2i).$$

These quasi-polynomials coincide with that of (2.2), (2.3), and (2.4).

Acknowledgments

This work was supported by JSPS KAKENHI Grant Number JP20K14282.

References

- C. A. Athanasiadis. Characteristic polynomials of subspace arrangements and finite fields. Adv. Math., 122, 193–233, (1996).
- [2] M. Beck and T. Zaslavsky. Inside-out polytopes. Adv. Math., 205 (1), 134–162, (2006).
- [3] A. Blass and B. Sagan. Characteristic and Ehrhart polynomials. J. Algebraic Combin., 7, 115–126, (1998).
- [4] H. Kamiya, A. Takemura, and H. Terao. Periodicity of hyperplane arrangements with integral coefficients modulo positive integers. J. Algebr. Comb., 27 (3), 317–330, (2008).
- [5] H. Kamiya, A. Takemura, and H. Terao. The characteristic quasi-polynomials of the arrangements of root systems. arXiv:0707.1381v1.

- [6] H. Kamiya, A. Takemura, and H. Terao. The characteristic quasi-polynomials of the arrangements of root systems and mid-hyperplane arrangements. Arrangements, local systems and singularities, Progr. Math., 283,177–190 Birkhäuser Verlag, Basel, (2010).
- [7] H. Kamiya, A. Takemura, and H. Terao. Periodicity of non-Central integral arrangements modulo positive integers. Ann. Comb., 15 (3), 449–464, (2011).
- [8] A. Higashitani, T. N. Tran, and M. Yoshinaga. Period collapse in characteristic quasipolynomials of hyperplane arrangements. Preprint. arXiv:2105.02367
- [9] A. Postnikov and R. Stanley, Deformations of Coxeter hyperplane arrangements. J. Combin. Theory Ser. A, 91, 544–597, (2000).
- [10] S. Tamura, Deformations of Coxeter hyperplane arrangements. Preprint. arXiv:2012.05634
- [11] M. Yoshinaga. Worpitzky partitions for root systems and characteristic quasi-polynomials. Tohoku Math. J., 70 (1), 39–63, (2018).
- [12] M. Yoshinaga. Characteristic polynomials of Linial arrangements for exceptional root systems. J. Combin. Theory Ser. A, 157, 267–286, (2018).