# $\epsilon$ -Expansion of Multivariable Hypergeometric Functions Appearing in Feynman Integral Calculus

Souvik Bera

Centre for High Energy Physics, Indian Institute of Science, Bangalore-560012, Karnataka, India

Email : souvikbera@IISc.ac.in

#### Abstract

We present a new methodology to perform the  $\epsilon$ -expansion of hypergeometric functions with linear  $\epsilon$  dependent Pochhammer parameters in any number of variables. Our approach allows one to perform Taylor as well as Laurent series expansion of multivariable hypergeometric functions. Each of the coefficients of  $\epsilon$  in the series expansion is expressed as a linear combination of multivariable hypergeometric functions with the same domain of convergence as that of the original hypergeometric function thereby providing a closed system of expressions. We present illustrative examples of hypergeometric functions in one, two and three variables which are typical of Feynman integral calculus.

Keywords :  $\epsilon$ -expansion, Feynman integrals, multivariable hypergeometric functions

## 1 Introduction

The evaluation of Feynman integrals originating in the perturbative calculations of quantum field theory is a challenging task. The difficulty of the calculation increases as the loops and legs and the scales involved in Feynman integral increases. These integrals can be represented by multivariable hypergeometric functions. These are well studied in the mathematics literature [1–4]. Methods based on Mellin-Barnes [5, 6], the method of negative dimension [7, 8], the method of bracket [9–11] evaluate Feynman integrals in terms of multivariable hypergeometric functions. Generalized hypergeometric functions are common in Feynman integral calculus. To give a few examples, two loop massive sunset diagram [12, 13], three loop vacuum diagram [14], one loop two and three point scalar functions [15, 16] are evaluated in terms of multivariable hypergeometric functions. Functional relations are used to find one loop Feynman integrals (see [17] and references within) in terms of multivariable hypergeometric functions. Feynman integrals can also be realized as GKZ hypergeometric system [18–20].

In dimensional regularization [21], the space-time dimension D (more precisely, integer multiples of D/2) of the Feynman integral appears in the Pochhammer parameters of the corresponding multivariable hypergeometric functions and the ratios of the scales involved in the integral take its place in the arguments of multivariable hypergeometric function. These functions have certain domain of convergence in its argument space. To find the numerical value of these functions beyond their defining domain of convergence one has to perform analytic continuations. The hypergeometric functions appearing in the Feynman integral calculus are usually expanded in Laurent series of  $\epsilon = (4 - D)/2$ . As a result, special functions appear in each order of  $\epsilon$ . These functions are analytically continued beyond their defining domain of convergences.

There are many results of analytic continuations of hypergeometric functions with more than one variable available in the mathematics literature [1-4, 22-26]. The analytic continuations of the double variable Appell  $F_2$  are found which are the backbone of the numerical package AppellF2.wl [28]. This package allows one to find the numerical value of the Appell  $F_2$  function for general values of Pochhammer parameters and real values of x

and y. Analytic continuations of three variable Srivastava  $H_C$  functions are obtained in [29]. Study of analytic continuations and numerical evaluations of other hypergeometric functions are ongoing [30, 31]. Recently, the process of finding analytic continuations of general hypergeometric functions is automated with Mathematica based Olsson.wl package [27]. The package allows one to obtain analytic continuations of hypergeometric functions with any number of variables and the region of convergences of double variable hypergeometric functions only. This package eases the calculation of finding analytic continuations of multivariable hypergeometric functions associated with a Feynman integral.

It is customary to express the result of Feynman integral in dimensional regularization scheme as a series in the parameter  $\epsilon = (4 - D)/2$ . This motivates to determine the  $\epsilon$ -expansion of the multivariable hypergeometric functions. The series expansion of the one variable Gauss hypergeometric function  $_2F_1$  [32] and  $_pF_{p-1}$  [33–35] are well studied. There are publicly available packages that can perform the expansion of  $_pF_{p-1}$  around integer and half-integer Pochhammer parameters analytically [36, 37] and numerically [38]. The  $\epsilon$ -expansion of double variable Appell and Kampé de Fériet functions are discussed in [39, 40]. In [41–43], differential equation approach is used to find the series expansion of multivariable hypergeometric function related to Feynman integrals (see also [44, 45]). Interested readers can find a comprehensive review of multivariable hypergeometric functions and their  $\epsilon$ -expansion related to the Feynman integral calculus in [43].

In this article we present a new approach to find the series expansion of multivariable hypergeometric functions analytically where the Pochhammer parameters are linearly dependent on  $\epsilon$ . There is no restriction on other parameters of the function and these could be complex numbers. Our approach is applicable for hypergeometric functions with any number of variables. Each of the coefficients of the  $\epsilon$  expansion is expressed as linear combination of hypergeometric functions. The domain of convergence of the hypergeometric functions appearing as coefficient of  $\epsilon$  is same as that of the starting hypergeometric function. Hence in the context of Feynman integrals, the hypergeometric structure remains intact thought out our method. Inside the associated domain of convergence, these analytic expressions can be used to find numerical values. We have applied our methodology to one, two and three variable hypergeometric functions and results are found to be consistent.

The article is organized as follows. In Section 2 we explain our methodology. A number of illustrative examples of hypergeometric functions in Section 3 are presented, followed by the conclusions. The ancillary notebook contains the first few terms of  $\epsilon$ -expansions of the hypergeometric functions including those presented in the article.

## 2 The methodology

The parameter  $\epsilon = (4 - D)/2$  appears linearly in the Pochhammer parameters of a multivariable hypergeometric function associated with a Feynman integral. The expansion of a multivariable hypergeometric function around  $\epsilon = 0$  can either be or Taylor type of Laurent type. If the lower Pochhammer parameters (i.e. Pochhammer parameters in the denominator) of a multivariable hypergeometric function has the form

$$(B_0 + B_1 \epsilon)_* \tag{1}$$

where  $B_0$  is zero or negative integer, then it is called singular. The "\*" denotes linear combination of summation indices which does not play any role in what follows. If one or more lower Pochhammer parameters of multivariable hypergeometric function are singular, then  $\epsilon$ -expansion of that function may be of Laurent type. If the lower Pochhammer parameters are non-singular, the  $\epsilon$ -expansion of the corresponding function is of Taylor type. Our methodology can be divided into two parts. In the first part we discuss the Taylor expansion of a multivariable hypergeometric function and in the second part the Laurent expansion (whenever applicable) is studied.

#### 2.1 Taylor expansion of multivariable hypergeometric functions

The Taylor series expansion around  $\epsilon = 0$  of a multivariable hypergeometric function which we denote by  $F(\epsilon)$ , suppressing dependencies of all the other parameters but  $\epsilon$ , can be found by taking successive derivatives with respect to  $\epsilon$ 

$$F(\epsilon) = \sum_{i=0}^{\infty} \frac{\epsilon^i}{i!} \frac{d^i}{d\epsilon^i} F(\epsilon) \Big|_{\epsilon=0}$$
<sup>(2)</sup>

In practice we only need first few terms.

Since the parameter  $\epsilon$  takes its place in the Pochhammer parameters of a multivariable hypergeometric function, taking derivative of Pochhammer parameters with respect to its argument plays crucial role in finding the Taylor

series expansion. Derivatives of Gauss hypergeometric function  ${}_2F_1$  with respect to the Pochhammer parameters are studied in [46]. Similar study for general  ${}_pF_q$  function can be found in [35, 47–49]. Discussions on the Pochhammer derivative of hypergeometric functions of more than one variable can be found in [39, 40, 50–52]. In this section we show how the Pochhammer derivative of a hypergeometric function with n fold summations can be represented as hypergeometric function with n + 1 fold summations and the same set of variables. This fact is used to find the Taylor series expansion of multivariable hypergeometric function in  $\epsilon$ . The coefficients of  $\epsilon$  themselves are multivariable hypergeometric functions.

#### 2.1.1 Derivative of Pochhammer parameters

We start by taking derivative of Pochhammer parameter with respect to its argument. In what follows, we assume that m, n, k takes non-negative integer values.

$$\frac{d(a)_n}{da} = (a)_n \left( \psi^{(0)}(a+n) - \psi^{(0)}(a) \right) = (a)_n \sum_{k=0}^{n-1} \frac{1}{a+k} = (a)_n \frac{1}{a} \sum_{k=0}^{n-1} \frac{(a)_k}{(a+1)_k}$$
(3)

where  $\psi^0$  is polygamma function. If a is  $\epsilon$  dependent, the chain rule can be used to find the derivative with respect to  $\epsilon$ ,

$$\frac{d(a)_m}{d\epsilon} = \frac{da}{d\epsilon} \frac{d(a)_m}{da} \tag{4}$$

Derivative of Pochhammer in other circumstances can be obtained by Pochhammer identities (listed in the Appendix A), chain rule and Eq. (3). We list few cases below.

• Derivative of powers of Pochhammer can be found by applying chain rule first and then Eq. (3). For example,

$$\frac{d}{da} [(a)_m]^p = p [(a)_m]^{p-1} \frac{d}{da} (a)_m$$
(5)

• To find the derivative of  $(a)_{pm}$ , where p is a positive integer, Gauss's multiplication formula can be used to simplify  $(a)_{pm}$  and then chain rule and Eq. (3) can be applied. For example,

$$\frac{d(a)_{2m}}{da} = \frac{d}{da} \left[ 2^{2m} \left( \frac{a}{2} + \frac{1}{2} \right)_m \left( \frac{a}{2} \right)_m \right] = 2^{2m} \frac{d}{da} \left[ \left( \frac{a}{2} + \frac{1}{2} \right)_m \right] \left( \frac{a}{2} \right)_m + 2^{2m} \left( \frac{a}{2} + \frac{1}{2} \right)_m \frac{d}{da} \left[ \left( \frac{a}{2} \right)_m \right]$$
(6)

•  $(a)_{-m}$  can be simplified using the Pochhammer identities and then the derivative of the result can be found using Eq. (3)

$$\frac{d}{da}(a)_{-m} = \frac{d}{da} \left[ \frac{(-1)^m}{(1-a)_m} \right] = (-1)^m \frac{d}{da} \left[ \frac{1}{(1-a)_m} \right]$$
(7)

• Using the Pochhammer identities, Pochhammer like  $(a)_{m+n}$  can be written as products of several Pochhammers. Then chain rule and Eq. (3) can be used to find the derivative of  $(a)_{m+n}$ .

$$\frac{d}{da}(a)_{m+n} = \frac{d}{da}\left[(a)_m(a+m)_n\right] = \frac{d}{da}\left[(a)_m\right](a+m)_n + (a)_m\frac{d}{da}\left[(a+m)_n\right]$$
(8)

Equipped with all the techniques of finding derivative of Pochhammer parameters, we are ready to consider Pochhammer derivative of multivariable hypergeometric function.

#### 2.1.2 Pochhammer derivative of multivariable hypergeometric function

We assume the parameter with respect to which the derivative of a multivariable hypergeometric function is to be taken is a. First the Pochhammer containing a can be simplified using Pochhammer identities. Then the chain rule of derivative can be applied. Essentially, the task of finding the Pochhammer derivative of a multivariable hypergeometric function boils down to evaluating a number of expressions that look like the following

$$\sum_{n=0}^{\infty} B(n) \frac{x^n}{n!} \frac{d}{da}(a)_n \tag{9}$$

Following [52], the summation of the index n is explicitly shown as the Pochhammer that is being differentiated contains n. Here B(n) collectively denotes the summation of indices other than n of the multivariable hypergeometric function. Using Eq. (3) we write

$$\sum_{n=0}^{\infty} B(n) \frac{x^n}{n!} \frac{d}{da}(a)_n = \sum_{n=0}^{\infty} B(n) \frac{x^n}{n!} (a)_n \frac{1}{a} \sum_{k=0}^{n-1} \frac{(a)_k}{(a+1)_k}$$
(10)

Now shifting  $n \to n+1$ ,

$$\sum_{n=0}^{\infty} B(n) \frac{x^n}{n!} \frac{d}{da}(a)_n = \sum_{n=0}^{\infty} B(n+1) \frac{x^{n+1}}{(n+1)!} (a)_{n+1} \frac{1}{a} \sum_{k=0}^n \frac{(a)_k}{(a+1)_k}$$
(11)

Reshuffling the summation of indices n, k as,

$$\sum_{n=0}^{\infty} \sum_{k=0}^{n} A(n,k) \to \sum_{n,k=0}^{\infty} A(n+k,k)$$
(12)

we find

$$\sum_{n=0}^{\infty} B(n) \frac{x^n}{n!} \frac{d}{da}(a)_n = \sum_{n,k=0}^{\infty} B(n+k+1) \frac{x^{n+k+1}}{(n+k+1)!} (a)_{n+k+1} \frac{1}{a} \frac{(a)_k}{(a+1)_k}$$
$$= x \sum_{n,k=0}^{\infty} B(n+k+1) \frac{(1)_k (1)_n (a)_k (a+1)_{k+n}}{(a+1)_k (2)_{k+n}} \frac{x^n x^k}{k! n!}$$
(13)

Taking the Pochhammer derivative of multivariable hypergeometric function introduces a new summation index. Thus the summation fold is increased by one. The region of convergence of the resultant hypergeometric function is same as the initial one [52].

#### 2.2 Laurent Expansion

As discussed above, if one or more lower Pochhammer parameters of multivariable hypergeometric function (primary function) are singular, then  $\epsilon$  expansion of that function may be of Laurent type. In such cases, one can find a suitable step down operator that relates the hypergeometric function in hand with the another one (secondary function) with lower Pochhammer parameter

$$(B_1\epsilon+1)_*$$

The hypergeometric function with non-singular lower Pochhammer parameters (i.e. secondary function) can be expanded in  $\epsilon$  as Taylor series (see Section 2.1). Once the Taylor series is found, the step down operator can be made to act on it to obtain the Laurent series of the hypergeometric function (primary function) we started with. All the  $\epsilon^{-p}$  (with p a positive integer) terms in the resultant Laurent series comes from the step down operator.

#### 2.2.1 The step down operator

There exists contiguous relations that relate multivariable hypergeometric functions whose Pochhammer parameters differ by integers. An algorithm based on Gröbner bases calculation is proposed by Takayama [53] to find the operators that generates such relations. These operators play a crucial role in the differential reduction technique of multivariable hypergeometric functions [54, 55]. For our purpose we only need one kind of operator, called the **step down** operator, to decrease the argument of the lower Pochhammer parameters by unity.

Let us define a multivariable hypergeometric function as

$$F(\mathbf{a}, \mathbf{b}, \mathbf{x}) = \sum_{\mathbf{m} \in \mathbb{N}_0^r} \frac{\Gamma(\mathbf{a} + \mu.\mathbf{m}) / \Gamma(\mathbf{a})}{\Gamma(\mathbf{b} + \nu.\mathbf{m}) / \Gamma(\mathbf{b})} \frac{\mathbf{x}^{\mathbf{m}}}{\mathbf{m}!} = \sum_{\mathbf{m} \in \mathbb{N}_0^r} \frac{(\mathbf{a})_{\mu.\mathbf{m}}}{(\mathbf{b})_{\nu.\mathbf{m}}} \frac{\mathbf{x}^{\mathbf{m}}}{\mathbf{m}!} = \sum_{\mathbf{m} \in \mathbb{N}_0^r} A(\mathbf{m}) \mathbf{x}^{\mathbf{m}}$$
(14)

Here **a**, **b** and **m** are vectors of length p, q and r respectively.  $\mu$  and  $\nu$  are matrices of size  $p \times r$  and  $q \times r$  respectively with integers as their elements. The expression is the definition of the r variable hypergeometric function with parameters **a** and **b**.  $\mathbb{N}_0$  denotes natural numbers including zero. We have used the vector notation here.

- $\mathbf{a} := \{a_1, a_2, \dots, a_n\}$
- $\mathbf{x}^{\mathbf{m}} := \prod_{i=1}^{n} x_i^{m_i}$
- $\mathbf{m}! := \prod_{i=1}^{n} (m_i!)$
- $\Gamma(\mathbf{a}) := \prod_{i=1}^{n} \Gamma(a_i)$
- $(\mathbf{a})_{\mathbf{m}} := \prod_{i=1}^{n} (a_i)_{m_i} = \prod_{i=1}^{n} \Gamma(a_i + m_i) / \Gamma(a_i)$
- $\sum_{\mathbf{m}\in\mathbb{N}_0^n}:=\sum_{m_1=0}^\infty\cdots\sum_{m_n=0}^\infty$

These multivariable hypergeometric functions are known to satisfy partial differential equations [1]. Let,

$$P_i = \frac{A(\mathbf{m} + \mathbf{e}_i)}{A(\mathbf{m})} = \frac{g_i(\mathbf{m})}{h_i(\mathbf{m})} \qquad , i = 1, \dots, r$$
(15)

where  $\mathbf{e}_i$  is unit vector with 1 in its i-th entry. The annihilators  $L_i$  of  $F(\mathbf{a}, \mathbf{b}, \mathbf{x})$  (Eq. (14)) are given by

$$L_{i} = \left[h_{i}(\boldsymbol{\theta})\frac{1}{x_{i}} - g_{i}(\boldsymbol{\theta})\right]$$
(16)

where  $\boldsymbol{\theta} = \{\theta_1, \dots, \theta_r\}$  is a vector containing Euler operators  $\theta_i = x_i \partial_{x_i}$ 

Following [53, 54], we define the unit step down operator for the lower Pochhammer parameter

$$F(\mathbf{a}, \mathbf{b}, \mathbf{x}) = \frac{1}{b_i} \left( \sum_{j=1}^r \nu_{ij} \theta_{x_j} + b_i \right) \bullet F(\mathbf{a}, \mathbf{b} + \mathbf{e_i}, \mathbf{x}) = H(b_i) \bullet F(\mathbf{a}, \mathbf{b} + \mathbf{e_i}, \mathbf{x})$$
(17)

Here the bullet meaning the action of the operator H on a function. The step down operator can be applied multiple times to decrease any lower Pochhammer parameter by suitable amount. In such cases, the product of the unit step down operators modulo the ideal generated by  $L_i$ 's can be taken as the step down operator.

$$H = \left[\prod_{i=1} H(b_i)\right] / \langle L_1, \dots, L_r \rangle$$
(18)

The calculation of Gröbner bases of the annihilators and the reduction of the step down operator with respect to the ideal generated by the annihilators can be found using computer programs [56] in Maple as well as [57, 58] in Mathematica. We present an example of step down operator for the one variable Gauss hypergeometric function below.

#### Example of Gauss hypergeometric function

The unit step down operator for the Gauss hypergeometric function is given by  $H(c) = \frac{1}{c}(\theta + c)$ , ie.

$${}_{2}F_{1}(a,b;c;x) = H(c) \bullet {}_{2}F_{1}(a,b;c+1,x)$$
(19)

Here the Euler operator  $\theta = x \partial_x$ . The above identity can be proved from the definition of the Gauss hypergeometric function.

$$c_{2}F_{1}(a,b;c;x) = \sum_{m=0}^{\infty} \frac{c(a)_{m}(b)_{m}}{(c)_{m}} \frac{x^{m}}{m!}$$
$$= \sum_{m=0}^{\infty} \frac{(c+m)(a)_{m}(b)_{m}}{(c+1)_{m}} \frac{x^{m}}{m!}$$
$$= (\theta+c) \bullet_{2}F_{1}(a,b;c+1;x)$$
(20)

In the second step we have used the Pochhammer identity  $\frac{c}{(c)_m} = \frac{c(c+m)\Gamma(c)}{(c+m)\Gamma(c+m)} = \frac{(c+m)}{(c+1)_m}$  and the last equality is obtained using  $\theta \bullet x^m = mx^m$ .

The unit step down operator can be used repetitively to find a relation between  ${}_{2}F_{1}(a,b;-n+c\epsilon;x)$  to  ${}_{2}F_{1}(a,b;c\epsilon+1;x)$ , where n is any positive integer. Note that the later hypergeometric function can be expanded in Taylor series. For example

$${}_{2}F_{1}(a,b;c-1;x) = H(c-1)H(c) \bullet {}_{2}F_{1}(a,b;c+1,x)$$
(21)

In the case, the step down operator H is H(c-1)H(c) modulo L, where L the annihilator of the hypergeometric function  ${}_{2}F_{1}(a,b;c+1,x)$ .

$$L \bullet _{2}F_{1}(a,b;c+1,x) = \left[-ab + (-x(a+b+1)+c+1)\partial_{x} - (x-1)x\partial_{x}^{2}\right] \bullet _{2}F_{1}(a,b;c+1,x) = 0$$
(22)

Hence we find the step down operator

$$H = \left(1 - \frac{abx}{(c-1)c(x-1)}\right) - \frac{x(a+b+1) - 2cx + c - 1}{(c-1)c(x-1)}\theta$$
(23)

which matches with the result obtained from the package pfq.m [54]. The above expression simplifies to

$${}_{2}F_{1}(a,b;c-1;x) = \left(1 - \frac{abx}{(c-1)c(x-1)}\right) {}_{2}F_{1}(a,b;c+1;x) + \frac{abx(x(-(a+b+1)) + c(2x-1)+1)}{(c-1)c(c+1)(x-1)} {}_{2}F_{1}(a+1,b+1;c+2;x)$$
(24)

#### 2.2.2 Argument derivative of multivariable hypergeometric function

Usually, the step down operators contain derivative operators with respect to the arguments of the corresponding multivariable hypergeometric function (see the examples in Section 3). We call the action of these derivative operators on multivariable hypergeometric function as 'argument derivative' in order to distinguish it from Pochhammer derivative discussed above.

Similar to the discussion in Section 2.1.2, we explicitly write the summation of the index n related to the argument with respect to which the differential is being taken.

$$\frac{d}{dx}\sum_{n=0}^{\infty}B(n)\frac{x^n}{n!}\tag{25}$$

Here, as before, B(n) collectively denotes the dependence of all the other summation indices in a multivariable hypergeometric function. So we have

$$\frac{d}{dx}\sum_{n=0}^{\infty}B(n)\frac{x^n}{n!} = \sum_{n=1}^{\infty}B(n)\frac{x^{n-1}}{(n-1)!} = \sum_{n=0}^{\infty}B(n+1)\frac{x^n}{n!}$$
(26)

We have shifted  $n \rightarrow n+1$  to get the last equality. Hence the argument derivative of a multivariable hypergeometric function is also another multivariable hypergeometric function. It is straight forward to realise that the domain of convergence of the resultant multivariable hypergeometric functions remain same as the initial one.

We evaluate  $\epsilon$ -expansion of one, two and three variables hypergeometric functions to demonstrate the methodology below.

#### 3 Examples

#### 3.1 Gauss $_2F_1$ example with half integer Pochhammer

We begin this section by providing the  $\epsilon$ -expansion of the following Gauss hypergeometric function

$$G := {}_{2}F_{1}(3, 2, \epsilon - 3/2, x) = \sum_{m=0}^{\infty} \frac{(2)_{m}(3)_{m}}{(\epsilon - \frac{3}{2})_{m}} \frac{x^{m}}{m!} \qquad \text{where } |x| < 1$$
(27)

There is only one lower Pochhammer parameter and it is not singular. Hence this Gauss function has Taylor expansion around  $\epsilon = 0$ . So we do not need any step down operators in this example. The coefficients of  $\epsilon^i$  are found by taking successive derivative with respect to  $\epsilon$  (Eq. (2)).

$$\begin{aligned} G &= {}_{2}F_{1}\left(2,3;-\frac{3}{2};x\right) \\ &- \frac{8x}{3}\epsilon F\left[\begin{array}{c} \{4,\{1,1\}\},\{3,\{1,1\}\},\{1,\{1,0\}\},\{1,\{0,1\}\},\{-\frac{3}{2},\{0,1\}\}\\ \{2,\{1,1\}\},\{-\frac{1}{2},\{1,1\}\},\{-\frac{1}{2},\{0,1\}\}\\ &- \frac{16x}{9}\epsilon^{2} F\left[\begin{array}{c} \{4,\{1,1\}\},\{3,\{1,1\}\},\{1,\{1,0\}\},\{1,\{0,1\}\},\{-\frac{3}{2},\{0,1\}\}\\ \{2,\{1,1\}\},\{-\frac{1}{2},\{1,1\}\},\{-\frac{1}{2},\{0,1\}\}\\ \{2,\{1,1\}\},\{-\frac{1}{2},\{1,1\}\},\{-\frac{1}{2},\{0,1\}\}\\ &\{3,\{1,1,1\}\},\{1,\{1,0,0\}\},\{-\frac{1}{2},\{0,1,1\}\},\{1,\{0,0,1\}\},\{1,\{0,0,1\}\},\{-\frac{1}{2},\{0,0,1\}\}\\ &\{3,\{1,1,1\}\},\{\frac{1}{2},\{1,1\}\},\{\frac{1}{2},\{0,1,1\}\},\{\frac{1}{2},\{0,0,1\}\}\\ &+ O(\epsilon^{3}) \end{aligned} \end{aligned}$$

The coefficients of  $\epsilon^3$  and  $\epsilon^4$  can be found in the ancillary file. We show the Taylor expansion obtained using HypExp [36, 37] below for comparison.

$$G = \frac{-315i\sqrt{-\frac{x}{x-1}}(4x+7)x^{2} \text{ HPL}\left(\{\text{plus}\}, i\sqrt{-\frac{x}{x-1}}\right) + 140x^{4} + 4294x^{3} + 2784x^{2} - 320x + 32}{32(x-1)^{6}} \\ + \frac{\epsilon x}{48(x-1)^{6}} \left[12i(143x+329)\sqrt{-\frac{x}{x-1}}x\text{HPL}\left(\{\text{plus}\}, i\sqrt{-\frac{x}{x-1}}\right) \\ + 945i(4x+7)\sqrt{-\frac{x}{x-1}}x\text{HPL}\left(\{0, \text{plus}\}, i\sqrt{-\frac{x}{x-1}}\right) - 144x^{3} + 2822x^{2} + 6912x - 128\right] \\ + \frac{\epsilon^{2}x}{72(x-1)^{6}} \left[-11340i\sqrt{-\frac{x}{x-1}}x^{2}\text{HPL}\left(\{0, 0, \text{plus}\}, i\sqrt{-\frac{x}{x-1}}\right) \\ - 18i\sqrt{-\frac{x}{x-1}}(42x+145)x\text{HPL}\left(\{\text{plus}\}, i\sqrt{-\frac{x}{x-1}}\right) - 36i\sqrt{-\frac{x}{x-1}}(143x+329)x\text{HPL}\left(\{0, \text{plus}\}, i\sqrt{-\frac{x}{x-1}}\right) \\ - 19845i\sqrt{-\frac{x}{x-1}}x\text{HPL}\left(\{0, 0, \text{plus}\}, i\sqrt{-\frac{x}{x-1}}\right) + 36x^{3} + 14474x^{2} + 20736x - 128\right]$$
(29)

Here HPL's are harmonic polylogarithms [59]. We evaluate Eq. (28) and Eq. (29) at a certain x(< 1) for comparison. The results are shown in Appendix C.1 and found to be consistent.

#### **3.2** Gauss $_2F_1$ example

Let us consider the another Gauss hypergeometric function

$$G := {}_{2}F_{1}(\epsilon, -\epsilon, \epsilon - 1, x) = \sum_{m=0}^{\infty} \frac{(\epsilon)_{m}(-\epsilon)_{m}}{(\epsilon - 1)_{m}} \frac{x^{m}}{m!} \qquad \text{where } |x| < 1$$
(30)

Here the lower Pochhammer parameter is singular so we find the step down operator first. This will allow us to relate the above function with another hypergeometric function which can be expanded in Taylor expansion.

$$G = H \bullet {}_{2}F_{1}(\epsilon, -\epsilon, \epsilon + 1, x) \tag{31}$$

We find the step down operator H to be,

$$H = \frac{\left(x(\epsilon(2x-1)-x+1)\right)}{(\epsilon-1)\epsilon(x-1)}\partial_x + \frac{\epsilon(2x-1)-x+1}{(\epsilon-1)(x-1)}$$
$$= \frac{1}{\epsilon}x\partial_x + \epsilon^0\left(1-\frac{x^2\partial_x}{x-1}\right) + \epsilon\left(-\frac{x(x\partial_x+1)}{x-1}\right) + \epsilon^2\left(-\frac{x(x\partial_x+1)}{x-1}\right) + \epsilon^3\left(-\frac{x(x\partial_x+1)}{x-1}\right) + O(\epsilon^4) \quad (32)$$

In the last step, we expanded the step down operator in series of  $\epsilon$ .

Next we find the Taylor expansion of  $_2F_1$  appearing in the right hand side of Eq. (31).

$${}_{2}F_{1}(\epsilon, -\epsilon, \epsilon+1, x) = 1 - \epsilon^{2} \left( x \ {}_{3}F_{2}(1, 1, 1; 2, 2; x) \right) + \epsilon^{3} \left( \frac{2}{3} x \ {}_{3}F_{2}(1, 1, 1; 2, 2; x) \right) \\ + \frac{x}{3} F \left[ \begin{array}{c} \{1, \{1, 1\}\}, \{1, \{1, 1\}\}, \{1, \{1, 0\}\}, \{1, \{0, 1\}\}, \{1, \{0, 1\}\} \\ \{2, \{1, 1\}\}, \{2, \{1, 1\}\}, \{2, \{1, 1\}\}, \{2, \{0, 1\}\} \end{array} \right| \left\{ x, x \right\} \right] \\ + \frac{x^{2}}{12} F \left[ \begin{array}{c} \{2, \{1, 1\}\}, \{2, \{1, 1\}\}, \{1, \{1, 0\}\}, \{2, \{0, 1\}\}, \{1, \{0, 1\}\} \\ \{3, \{1, 1\}\}, \{3, \{1, 1\}\}, \{3, \{0, 1\}\} \end{array} \right| \left\{ x, x \right\} \right] \right) + O(\epsilon^{4})$$
(33)

The coefficient of  $\epsilon^4$  can be found in the ancillary file.

Now applying the step down operator (i.e. Eq. (32)) on the Taylor series (Eq. (33)), we find the Laurent expansion of the Gauss  $_2F_1$  we started with (i.e. Eq. (30)),

$$\begin{split} G &= 1 + \epsilon \left( -\frac{x}{x-1} - x_{3}F_{2}(1,1,1;2,2;x) - \frac{x^{2}}{4} {}_{3}F_{2}(2,2,2;3,3;x) \right) \\ &+ \epsilon^{2} \left[ -\frac{x}{x-1} + \frac{(x(2x+1))}{3(x-1)} {}_{3}F_{2}(1,1,1;2,2;x) + \frac{(x^{2}(5x-2))}{12(x-1)} {}_{3}F_{2}(2,2,2;3,3;x) \right. \\ &+ \frac{x}{3} F \left[ \left. \{ 1, \{1,1\}\}, \{1, \{1,1\}\}, \{1, \{1,0\}\}, \{1, \{0,1\}\}, \{1, \{0,1\}\}\} \right| \left\{ x, x \} \right] \\ &+ \frac{x^{2}}{24} F \left[ \left. \{ 2, \{1,1\}\}, \{2, \{1,1\}\}, \{2, \{1,1\}\}, \{2, \{0,1\}\}\}, \{2, \{0,1\}\}\} \right| \left\{ x, x \} \right] \\ &+ \frac{x^{2}}{12} F \left[ \left. \{ 2, \{1,1\}\}, \{2, \{1,1\}\}, \{2, \{1,0\}\}, \{2, \{0,1\}\}, \{2, \{0,1\}\}\} \right| \left\{ x, x \} \right] \\ &+ \frac{x^{2}}{12} F \left[ \left. \{ 2, \{1,1\}\}, \{2, \{1,1\}\}, \{2, \{1,0\}\}, \{2, \{0,1\}\}, \{1, \{0,1\}\}\} \right| \left\{ x, x \} \right] \\ &+ \frac{x^{2}}{6} F \left[ \left. \{ 2, \{1,1\}\}, \{2, \{1,1\}\}, \{3, \{1,1\}\}, \{3, \{1,1\}\}, \{3, \{0,1\}\}, \{2, \{0,1\}\}\} \right| \left\{ x, x \} \right] \\ &+ \frac{2x^{3}}{81} F \left[ \left. \{ 3, \{1,1\}\}, \{3, \{1,1\}\}, \{3, \{1,1\}\}, \{2, \{1,0\}\}, \{2, \{0,1\}\}, \{1, \{0,1\}\}\} \right| \left\{ x, x \} \right] \\ &+ \frac{x^{3}}{27} F \left[ \left. \{ 3, \{1,1\}\}, \{3, \{1,1\}\}, \{2, \{1,0\}\}, \{2, \{0,1\}\}, \{1, \{0,1\}\}\} \right| \left\{ x, x \} \right] \right] + O(\epsilon^{3}) \end{split}$$
(34)

We present the Laurent expansion obtained from HypExp package [36] [37] for comparison,

$$G = 1 + \epsilon \left( \log(1-x) - \frac{x}{x-1} \right) + \epsilon^2 \left( -\frac{x}{x-1} + \frac{1}{2} \log^2(1-x) - \frac{x \log(1-x)}{x-1} \right) + O(\epsilon^3)$$
(35)

We found that our result Eq. (34) is in agreement with the result from HypExp Eq. (35) numerically at given x (< 1) evaluated with sufficient number of terms. It is also consistent with NumExp package [38]. The numerical comparison can be found in Appendix C.2.

#### **3.3** $_2F_2$ example

In this example we apply the methodology  $_2F_2$ . Consider the following function

$$G := {}_2F_2(1,2;\epsilon,\epsilon-1,x) = \sum_{m=0}^{\infty} \frac{(1)_m(2)_m}{(\epsilon)_m(\epsilon-1)_m} \frac{x^m}{m!} \qquad \text{where } |x| < \infty$$

$$(36)$$

Note that the lower Pochhammer parameters are  $\epsilon$  and  $\epsilon - 1$ . So, the function G may have Laurent series expansion. On the other hand, the function  $_2F_2(1,2;\epsilon+1,\epsilon+1,x)$  has Taylor series expansion. We find the step down operator that relates to the former i.e. Eq. (36) with the later

$$G = H \bullet_{2}F_{2}(1, 2; \epsilon + 1, \epsilon + 1, x)$$
(37)

The step down operator is found to be

$$H = \frac{\epsilon^3 - \epsilon^2 + 2x}{(\epsilon - 1)\epsilon^2} + \frac{x\left(2\epsilon^2 - \epsilon + 4x - 1\right)}{(\epsilon - 1)\epsilon^2}\partial_x + \frac{x^2(\epsilon + x - 1)}{(\epsilon - 1)\epsilon^2}\partial_x^2 \tag{38}$$

Expanding the operator H in series, and collecting the coefficients of  $\epsilon$  we find

$$H = \left[ -(x-1)x^2 \partial_x^2 + x(1-4x)\partial_x - 2x \right] \frac{1}{\epsilon^2} + \left[ -x^3 \partial_x^2 + (2-4x)x \partial_x - 2x \right] \frac{1}{\epsilon} + \left[ -x^3 \partial_x^2 - 4x^2 \partial_x - 2x + 1 \right] + \left[ -x^3 \partial_x^2 - 4x^2 \partial_x - 2x \right] \epsilon + O(\epsilon^2)$$
(39)

The Taylor expansion of the  $_2F_2$  function on the right and side of Eq. (37) is found to be

$${}_{2}F_{2}(1,2;\epsilon+1,\epsilon+1,x) = {}_{1}F_{1}(2;1,x) + -4\epsilon x F \begin{bmatrix} \{3,\{1,1\}\},\{1,\{1,0\}\},\{1,\{0,1\}\},\{1,\{0,1\}\} \\ \{2,\{1,1\}\},\{2,\{1,1\}\},\{2,\{0,1\}\} \end{bmatrix} | \{x,x\} \end{bmatrix} \\ + 6x\epsilon^{2} F \begin{bmatrix} \{3,\{1,1\}\},\{1,\{1,0\}\},\{1,\{0,1\}\},\{2,\{0,1\}\} \\ \{2,\{1,1\}\},\{2,\{1,1\}\},\{2,\{0,1\}\} \end{bmatrix} | \{x,x\} \end{bmatrix} \\ + \frac{3x^{2}}{4}\epsilon^{2} F \begin{bmatrix} \{4,\{1,1,1\}\},\{1,\{1,0,0\}\},\{2,\{0,1,1\}\},\{1,\{0,1,0\}\},\{1,\{0,1,0\}\},\{1,\{0,0,1\}\} \\ \{3,\{1,1,1\}\},\{3,\{1,1,1\}\},\{3,\{0,1,1\}\},\{2,\{0,1,0\}\} \end{bmatrix} | \{x,x,x\} \end{bmatrix} \\ + \frac{9x^{2}}{8}\epsilon^{2} F \begin{bmatrix} \{4,\{1,1,1\}\},\{1,\{1,0,0\}\},\{2,\{0,1,1\}\},\{1,\{0,1,0\}\},\{2,\{0,0,1\}\},\{1,\{0,0,1\}\} \\ \{3,\{1,1,1\}\},\{3,\{1,1,1\}\},\{3,\{0,1,1\}\},\{3,\{0,0,1\}\} \end{bmatrix} | \{x,x,x\} \end{bmatrix} \\ + O(e^{3}) \tag{40}$$

To find the Laurent series expansion of Eq. (36), we apply the step down operator Eq. (39) on the Taylor series expansion of  ${}_{2}F_{2}(1,2;\epsilon+1,\epsilon+1,x)$ , i.e. Eq. (40). The Laurent expansion reads,

$$\begin{split} G &= \left[ -2x_{1}F_{1}(2;1;x) + 2(1-4x)x_{1}F_{1}(3;2;x) - 3(x-1)x^{2} {}_{1}F_{1}(4;3;x) \right] \frac{1}{\epsilon^{2}} \\ &+ \left[ -2x_{1}F_{1}(2;1;x) + 2(2-4x)x_{1}F_{1}(3;2;x) - 3x^{3} {}_{1}F_{1}(4;3;x) \right] \\ &+ 4x(6x-1) F \left[ \begin{cases} \{3, \{1,1\}\}, \{1, \{1,0\}\}, \{1, \{0,1\}\}, \{1, \{0,1\}\} \\ \{2, \{1,1\}\}, \{2, \{1,1\}\}, \{2, \{0,1\}\} \\ \{2, \{1,1\}\}, \{2, \{1,1\}\}, \{2, \{0,1\}\} \\ \{3, \{1,1\}\}, \{3, \{1,1\}\}, \{2, \{0,1\}\} \\ \{3, \{1,1\}\}, \{3, \{1,1\}\}, \{2, \{0,1\}\} \\ \{3, \{1,1\}\}, \{3, \{1,1\}\}, \{2, \{0,1\}\} \\ \{3, \{1,1\}\}, \{3, \{1,1\}\}, \{2, \{0,1\}\} \\ \{3, \{1,1\}\}, \{3, \{1,1\}\}, \{3, \{0,1\}\} \\ + \frac{9x^{2}(2x-1)}{2} F \left[ \begin{cases} \{4, \{1,1\}\}, \{1, \{1,0\}\}, \{2, \{0,1\}\}, \{2, \{0,1\}\} \\ \{3, \{1,1\}\}, \{3, \{1,1\}\}, \{3, \{0,1\}\} \\ \{4, \{1,1\}\}, \{4, \{1,1\}\}, \{2, \{0,1\}\} \\ \{4, \{1,1\}\}, \{4, \{1,1\}\}, \{2, \{0,1\}\} \\ \{4, \{1,1\}\}, \{4, \{1,1\}\}, \{2, \{0,1\}\} \\ \{4, \{1,1\}\}, \{4, \{1,1\}\}, \{3, \{0,1\}\} \\ \{4, x,x\} \right] \\ &+ \frac{8(x-1)x^{3}}{9} F \left[ \begin{cases} \{5, \{1,1\}\}, \{1, \{1,0\}\}, \{3, \{0,1\}\}, \{3, \{0,1\}\} \\ \{4, \{1,1\}\}, \{4, \{1,1\}\}, \{4, \{0,1\}\} \\ \{4, \{1,1\}\}, \{4, \{1,1\}\}, \{4, \{0,1\}\} \\ \{4, x,x\} \right] \\ &+ \frac{8(x-1)x^{3}}{9} F \left[ \begin{cases} \{5, \{1,1\}\}, \{1, \{1,0\}\}, \{3, \{0,1\}\}, \{3, \{0,1\}\} \\ \{4, \{1,1\}\}, \{4, \{1,1\}\}, \{4, \{0,1\}\} \\ \{4, \{1,1\}\}, \{4, \{1,1\}\}, \{4, \{0,1\}\} \\ \{4, x,x\} \right] \\ &+ \frac{8(x-1)x^{3}}{9} F \left[ \begin{cases} \{5, \{1,1\}\}, \{1, \{1,0\}\}, \{3, \{0,1\}\}, \{3, \{0,1\}\} \\ \{4, \{1,1\}\}, \{4, \{1,1\}\}, \{4, \{0,1\}\} \\ \{4, \{1,1\}\}, \{4, \{1,1\}\}, \{4, \{0,1\}\} \\ \{4, \{1,1\}\}, \{4, \{1,1\}\}, \{4, \{0,1\}\} \\ \{4, 5, 4\} \\ \{4, 5, 5\} \\ \{4, 5, 5, 5\} \\ \{4, 5, 5, 5\} \\ \{4, 5, 5, 5\} \\ \{4, 5, 5, 5\} \\ \{4, 5, 5, 5\} \\ \{4, 5, 5, 5\} \\ \{4, 5, 5\} \\ \{4, 5, 5\} \\ \{4, 5, 5\} \\ \{4, 5, 5\} \\ \{4, 5, 5\} \\ \{4, 5, 5\} \\ \{4, 5, 5\} \\ \{4, 5, 5\} \\ \{4, 5, 5\} \\ \{4, 5, 5\} \\ \{4, 5, 5\} \\ \{4, 5, 5\} \\ \{4, 5, 5\} \\ \{4, 5, 5\} \\ \{4, 5, 5\} \\ \{4, 5, 5\} \\ \{4, 5, 5\} \\ \{4, 5\}$$

The next two higher order terms are also evaluated and can be found in the ancillary files. A numerical test of the above result can be found in Appendix C.3.

## **3.4** Appell $F_1$ example

Let us consider the following double variable Appell  $F_1$  function which we wish to find the  $\epsilon$  expansion of,

$$F_1(3/2, 2\epsilon + 1, 4 - \epsilon, \epsilon - 2, x, y) = \sum_{m,n=0}^{\infty} \frac{\left(\frac{3}{2}\right)_{m+n} (2\epsilon + 1)_m (4-\epsilon)_n}{(\epsilon - 2)_{m+n}} \frac{x^m}{m!} \frac{y^n}{n!} \qquad \text{where} \quad |x| < 1 \land |y| < 1 \qquad (42)$$

The corresponding Appell  $F_1$  function that can be expanded in Taylor series is given by

$$F_1(3/2, 2\epsilon + 1, 4 - \epsilon, \epsilon + 1, x, y) = \sum_{m,n=0}^{\infty} \frac{\left(\frac{3}{2}\right)_{m+n} (2\epsilon + 1)_m (4 - \epsilon)_n}{(\epsilon + 1)_{m+n}} \frac{x^m}{m!} \frac{y^n}{n!} \qquad \text{where} \quad |x| < 1 \land |y| < 1 \qquad (43)$$

We find the Taylor expansion of the Appell  $F_1$  function in Eq. (43) and act the step down operator on it to find Laurent series expansion of the Appell  $F_1$  function Eq. (42).

The step down operator that relates the above Appell  $F_1$  function in Eq. (42) with the Appell  $F_1$  function in Eq. (43) is found to be,

$$H = C_0 + C_1 \partial_x + C_2 \partial_y \tag{44}$$

where are rational functions  $C_i$ 's are

$$\begin{split} C_{0} &= \frac{1}{4\epsilon \left(\epsilon^{2} - 3\epsilon + 2\right) \left(x - 1\right)^{2} \left(y - 1\right)^{2}} \left[ 4\epsilon^{3} \left(x - 1\right)^{2} \left(y - 1\right)^{2} - 12\epsilon^{2} \left(x^{2} \left(2y^{2} - 5y + 2\right) - 2x \left(y^{2} - 4y + 2\right) - y^{2} - y + 1\right) \right. \\ &+ \epsilon \left(x^{2} \left(5y^{2} - 160y + 68\right) + 2x \left(91y^{2} + 10y - 14\right) - 139y^{2} + 44y + 8\right) \\ &+ 3 \left(x^{2} \left(95y^{2} - 54y + 11\right) - 4x \left(39y^{2} - 14y + 1\right) + 4y(17y - 4)\right) \right] \end{split}$$
(45)  
$$\begin{aligned} C_{1} &= \frac{1}{4(\epsilon - 2)(\epsilon - 1)\epsilon(x - 1)^{2}(y - 1)^{2}} \left[ x \left( 4\epsilon^{2} \left(x^{2} \left(y^{2} + 1\right) + x \left(-4y^{2} + y - 1\right) + 4y^{2} - 3y + 1\right) \right. \\ &- 2\epsilon \left(x^{2} \left(22y^{2} + 6y + 1\right) + x \left(-78y^{2} + 39y - 19\right) + 44y^{2} - 21y + 6\right) + x^{2} \left(225y^{2} - 178y + 57\right) \\ &- 2x \left(155y^{2} - 66y + 15\right) + 8 \left(15y^{2} - 3y + 1\right) \right) \right] \end{aligned}$$
(46)  
$$\begin{aligned} C_{2} &= \frac{1}{4(\epsilon - 2)(\epsilon - 1)\epsilon(x - 1)^{2}(y - 1)^{2}} \left[ y \left( 4\epsilon^{2} \left(x^{2}(y + 1)^{2} - 2x(3y + 1)y + 7y^{2} - 4y + 1\right) \right. \\ &- 4\epsilon \left(x^{2} \left(11y^{2} + 18y - 5\right) - 6x \left(9y^{2} - 2y + 1\right) + 35y^{2} - 14y + 3\right) \\ &+ x^{2} \left(225y^{2} - 106y + 24\right) - 2x \left(191y^{2} - 57y + 9\right) + 171y^{2} - 36y + 8 \right) \right] \end{aligned}$$
(47)

Now expanding all the three coefficients  $C_i$ , in  $\epsilon$ , we club coefficients of different powers of  $\epsilon$  together,

$$H = A_1 \frac{1}{\epsilon} + A_2 \epsilon^0 + A_3 \epsilon + A_4 \epsilon^2 + O(\epsilon^3)$$

$$\tag{48}$$

where,

$$\begin{split} A_{1} &= \frac{x \left(x^{2} \left(225 y^{2} - 178 y + 57\right) - 2x \left(155 y^{2} - 66 y + 15\right) + 8 \left(15 y^{2} - 3 y + 1\right)\right) \partial_{x}}{8 (x - 1)^{2} (y - 1)^{2}} \\ &+ \frac{y \left(x^{2} \left(225 y^{2} - 106 y + 24\right) - 2x \left(191 y^{2} - 57 y + 9\right) + 171 y^{2} - 36 y + 8\right) \partial_{y}}{8 (x - 1)^{2} (y - 1)^{2}} \\ &+ \frac{3 \left(x^{2} \left(95 y^{2} - 54 y + 11\right) - 4x \left(39 y^{2} - 14 y + 1\right) + 4y (17 y - 4)\right)\right)}{8 (x - 1)^{2} (y - 1)^{2}} \end{split}$$
(49)  
$$A_{2} &= \frac{x \left(587 x^{2} y^{2} - 558 x^{2} y + 167 x^{2} - 618 x y^{2} + 240 x y - 14 x + 184 y^{2} + 12 y\right) \partial_{x}}{16 (x - 1)^{2} (y - 1)^{2}} \\ &+ \frac{y \left(587 x^{2} y^{2} - 462 x^{2} y + 112 x^{2} - 714 x y^{2} + 246 x y - 6x + 233 y^{2} + 4y\right) \partial_{y}}{16 (x - 1)^{2} (y - 1)^{2}} \\ &+ \frac{865 x^{2} y^{2} - 806 x^{2} y + 235 x^{2} - 1040 x y^{2} + 544 x y - 92 x + 334 y^{2} - 56 y + 16}{16 (x - 1)^{2} (y - 1)^{2}} \end{aligned}$$
(50)  
$$A_{3} &= \frac{x \left(1327 x^{2} y^{2} - 1318 x^{2} y + 403 x^{2} - 1298 x y^{2} + 472 x y + 2x + 376 y^{2} + 36 y\right) \partial_{x}}{32 (x - 1)^{2} (y - 1)^{2}} \\ &+ \frac{y \left(1327 x^{2} y^{2} - 1142 x^{2} y + 304 x^{2} - 1474 x y^{2} + 478 x y + 18 x + 469 y^{2} + 20 y\right) \partial_{y}}{32 (x - 1)^{2} (y - 1)^{2}} \\ &+ \frac{3 \left(643 x^{2} y^{2} - 618 x^{2} y + 181 x^{2} - 696 x y^{2} + 304 x y - 20 x + 214 y^{2} - 8y\right)}{32 (x - 1)^{2} (y - 1)^{2}} \end{aligned}$$
(51)

The Taylor Expansion of the Appell  $F_1$  in Eq. (43) is found taking successive derivative with respect to  $\epsilon$  i.e.,

$$F_{1}(\epsilon, x, y) = F_{1}(0, x, y) + \epsilon F_{1}'(\epsilon, x, y)\Big|_{\epsilon=0} + \frac{1}{2!}\epsilon^{2}F_{1}''(\epsilon, x, y)\Big|_{\epsilon=0} + \frac{1}{3!}\epsilon^{3}F_{1}'''(\epsilon, x, y)\Big|_{\epsilon=0} + O(e^{4})$$
(52)  
=  $f_{0} + \epsilon f_{1} + \epsilon^{2}f_{2} + \epsilon^{3}f_{3} + O(\epsilon^{4})$ (53)

All the Pochhammer parameters of the Appell  $F_1$  in Eq. (43) are collectively denoted as  $\epsilon$  in the argument of  $F_1$  in Eq. (52) to give emphasis on the fact that the parameters are  $\epsilon$  dependent. The primes in the right hand side refers to the derivative with respect to  $\epsilon$ . We write the first few  $f_i$  below.

$$\begin{aligned} f_{0} &= F_{1}\left(\frac{3}{2}, 1, 4, 1, x, y\right) \\ f_{1} &= 3x \ F\left[\begin{array}{c} \left\{\frac{5}{2}, \{1, 1, 1\}\right\}, \{1, \{1, 0, 0\}\}, \{1, \{1, 0, 0\}\}, \{1, \{0, 1, 0\}\}, \{4, \{0, 0, 1\}\} \\ \{2, \{1, 1, 1\}\}, \{2, \{1, 0, 0\}\} \\ \left\{\frac{5}{2}, \{1, 1, 1\}\right\}, \{1, \{1, 0, 1\}\}, \{1, \{1, 0, 0\}\}, \{1, \{0, 1, 0\}\}, \{4, \{0, 0, 1\}\} \\ \{2, \{1, 1, 1\}\}, \{2, \{1, 0, 1\}\} \\ \left\{\frac{5}{2}, \{1, 1, 1\}\right\}, \{5, \{1, 0, 1\}\}, \{1, \{1, 0, 0\}\}, \{1, \{1, 0, 0\}\}, \{1, \{0, 1, 0\}\}, \{1, \{0, 0, 1\}\} \\ \{2, \{1, 1, 1\}\}, \{2, \{1, 0, 1\}\}, \{2, \{1, 0, 0\}\} \\ \left\{\frac{5}{2}, \{1, 1, 1\}\right\}, \{5, \{1, 0, 1\}\}, \{4, \{1, 0, 0\}\}, \{1, \{1, 0, 0\}\}, \{1, \{0, 1, 0\}\}, \{1, \{0, 0, 1\}\} \\ \left\{\frac{3y}{2}, F\left[\begin{array}{c} \left\{\frac{5}{2}, \{1, 1, 1\}\right\}, \{5, \{1, 0, 1\}\}, \{4, \{1, 0, 0\}\}, \{1, \{1, 0, 0\}\}, \{1, \{0, 1, 0\}\}, \{1, \{0, 0, 1\}\} \\ \{2, \{1, 1, 1\}\}, \{2, \{1, 0, 1\}\}, \{5, \{1, 0, 1\}\}, \{4, \{1, 0, 0\}\}, \{1, \{0, 1, 0\}\}, \{1, \{0, 0, 1\}\} \\ \{2, \{1, 1, 1\}\}, \{2, \{1, 0, 1\}\}, \{5, \{1, 0, 1\}\}, \{4, \{1, 0, 0\}\}, \{1, \{0, 1, 0\}\}, \{1, \{0, 0, 1\}\} \\ \{2, \{1, 1, 1\}\}, \{2, \{1, 0, 1\}\}, \{5, \{1, 0, 1\}\}, \{4, \{1, 0, 0\}\}, \{1, \{0, 1, 0\}\}, \{1, \{0, 0, 1\}\} \\ \{2, \{1, 1, 1\}\}, \{2, \{1, 0, 1\}\}, \{5, \{1, 0, 1\}\}, \{4, \{1, 0, 0\}\}, \{1, \{0, 1, 0\}\}, \{1, \{0, 0, 1\}\} \\ \{4, x, y\}\right\end{bmatrix} \end{aligned}$$
(55)

We do not show the functions  $f_2$ ,  $f_3$  here due to its long expression. It is provided in the ancillary notebook. Now applying the step down operator on the Appell  $F_1$  in Eq. (43),

$$F_1(3/2, 2\epsilon + 1, 4 - \epsilon, \epsilon - 2, x, y) = H \bullet F_1(3/2, 2\epsilon + 1, 4 - \epsilon, \epsilon + 1, x, y)$$
(56)

Using Eq. (48) and Eq. (53) on the right hand side of the above equation,

$$F_{1}(3/2, 2\epsilon + 1, 4 - \epsilon, \epsilon - 2, x, y) = \left(A_{1}\frac{1}{\epsilon} + A_{2}\epsilon^{0} + A_{3}\epsilon + A_{4}\epsilon^{2} + O(\epsilon^{3})\right) \bullet \left(f_{0} + \epsilon f_{1} + e^{2}f_{2} + e^{3}f_{3} + O(e^{4})\right)$$
$$= \frac{1}{\epsilon}\left[A_{1} \bullet f_{0}\right] + \left[A_{1} \bullet f_{1} + A_{2} \bullet f_{0}\right] + \epsilon\left[A_{3} \bullet f_{0} + A_{2} \bullet f_{1} + A_{1} \bullet f_{2}\right]$$
$$+ \epsilon^{2}\left[A_{4} \bullet f_{0} + A_{3} \bullet f_{1} + A_{2} \bullet f_{2} + A_{1} \bullet f_{3}\right] + O(\epsilon^{4})$$
(57)

We apply  $A_1$  (Eq. (49)) on  $f_0$  (Eq. (54)) to find the coefficient of  $\epsilon^{-1}$  of the Laurent expansion,

$$A_{1} \bullet f_{0} = \frac{3\left(x^{2}\left(95y^{2} - 54y + 11\right) - 4x\left(39y^{2} - 14y + 1\right) + 4y(17y - 4)\right)}{8(x - 1)^{2}(y - 1)^{2}} F_{1}\left(\frac{3}{2}, 1, 4, 1, x, y\right) \\ + \frac{3x\left(x^{2}\left(225y^{2} - 178y + 57\right) - 2x\left(155y^{2} - 66y + 15\right) + 8\left(15y^{2} - 3y + 1\right)\right)}{16(x - 1)^{2}(y - 1)^{2}} F_{1}\left(\frac{5}{2}, 2, 4, 2, x, y\right) \\ + \frac{3y\left(x^{2}\left(225y^{2} - 106y + 24\right) - 2x\left(191y^{2} - 57y + 9\right) + 171y^{2} - 36y + 8\right)}{4(x - 1)^{2}(y - 1)^{2}} F_{1}\left(\frac{5}{2}, 1, 5, 2, x, y\right)$$
(58)

Similarly one find the higher order term of the Laurent series. The coefficient of  $\epsilon^0, \epsilon^1$  in the Laurent series can be found in the ancillary notebook. The coefficients are numerically tested and found to be consistent. The results of the numerical tests can be found in the Appendix C.4.

## **3.5** Lauricella $F_C^{(3)}$ Example

To provide an example of three variable hypergeometric function we chose Lauricella  $F_C^{(3)}$  function which is linked with two loop different mass sunset integral [12, 13]. In particular, we will find the  $\epsilon$ -expansion of one of the four Lauricella function in Eq. (16) of [12]. Let us take,

$$G := F_C^{(3)} (1, 1 - \nu, 1 + \nu, 1 - \nu, 1 + \nu, z_1, z_2, z_3)$$
  
=  $\sum_{n_1, n_2, n_3 = 0}^{\infty} \frac{(1)_{n_1 + n_2 + n_3}(e)_{n_1 + n_2 + n_3}}{(2 - e)_{n_1}(2 - e)_{n_3}(e)_{n_2}} \frac{z_1^{n_1} z_2^{n_2} z_3^{n_3}}{n_1! n_2! n_3!}$  (59)

which is valid for  $\sqrt{|z_1|} + \sqrt{|z_2|} + \sqrt{|z_3|} < 1$ . We have used  $\nu = 1 - \epsilon$ .

Since one of the lower Pochhammer parameters is singular, we find the step down operator first.

$$H = \frac{z_2 \partial_{z_2}}{\epsilon} + 1 \tag{60}$$

Hence,

$$F_{C}^{(3)}(1,\epsilon,2-\epsilon,2-\epsilon,\epsilon,z_{1},z_{2},z_{3}) = H \bullet F_{C}^{(3)}(1,\epsilon,2-\epsilon,2-\epsilon,\epsilon+1,z_{1},z_{2},z_{3})$$
(61)

The Taylor series expansion of the Lauricella function with non-singular lower Pochhammer parameter is found to be

$$F_{C}^{(3)}(1,\epsilon,2-\epsilon,2-\epsilon,\epsilon+1,z_{1},z_{2},z_{3}) = 1 + \left[\frac{1}{2}z_{1\,2}F_{1}(1,1;3;z_{1}) + z_{2}F\left[\begin{array}{c}\{2,\{1,1\}\},\{1,\{1,1\}\},\{1,\{0,1\}\}\\\{2,\{1,0\}\},\{2,\{0,1\}\},\{2,\{0,1\}\}\end{array}\right| \{z_{1},z_{2}\}\right] \\ + \frac{z_{3}}{2}F\left[\begin{array}{c}\{2,\{1,1,1\}\},\{1,\{1,1,1\}\},\{1,\{0,0,1\}\}\\\{2,\{1,0,0\}\},\{1,\{0,1,0\}\},\{3,\{0,0,1\}\},\{2,\{0,0,1\}\}\end{vmatrix}\right| \{z_{1},z_{2},z_{3}\}\right]\right]\epsilon + O(\epsilon^{2})$$

$$(62)$$

Now applying the step down operator (Eq. (60)) on the Taylor expansion (Eq. (62)), we find

$$F_{C}^{(3)}(1,\epsilon,2-\epsilon,2-\epsilon,\epsilon,z_{1},z_{2},z_{3}) = 1 + z_{2} F \begin{bmatrix} \{2,\{1,1\}\},\{1,\{1,1\}\},\{1,\{0,1\}\}\\ \{2,\{1,0\}\},\{2,\{0,1\}\},\{2,\{0,1\}\} \end{bmatrix} | \{z_{1},z_{2}\} \end{bmatrix} + \frac{z_{2}^{2}}{2} F \begin{bmatrix} \{3,\{1,1\}\},\{2,\{1,1\}\},\{2,\{0,1\}\}\\ \{2,\{1,0\}\},\{3,\{0,1\}\},\{3,\{0,1\}\} \end{bmatrix} | \{z_{1},z_{2}\} \end{bmatrix} + z_{2}z_{3} F \begin{bmatrix} \{3,\{1,1,1\}\},\{2,\{1,1,1\}\},\{2,\{1,1,1\}\},\{1,\{0,0,1\}\}\\ \{2,\{1,0,0\}\},\{2,\{0,1,0\}\},\{3,\{0,0,1\}\},\{2,\{0,0,1\}\} \end{bmatrix} | \{z_{1},z_{2},z_{3}\} \end{bmatrix} + O(\epsilon)$$

$$(63)$$

The  $\epsilon$  order term can be found in the ancillary file. A numerical consistency check of Eq. (59) and Eq. (63) can be found in Appendix C.5.

### 4 Conclusions

We have presented a new method to obtain the  $\epsilon$ -expansion of hypergeometric functions in one or more variables. In our approach, when the expansion is of Taylor type, successive derivative with respect to  $\epsilon$  has to evaluated. We have demonstrated the process of Pochhammer derivative of a multivariable hypergeometric functions which plays important role in finding the Taylor expansion. When lower Pochhammer parameters of a hypergeometric function are singular, the series expansion may be of Laurent type. In this situation, an appropriate step down operator can be found, which relates the starting hypergeometric function to another one with shifted lower Pochhammer parameters. This hypergeometric function with non-singular lower Pochhammer parameters can be expanded in Taylor series expansion. Finally the obtained step down operator can be made to act on the Taylor series of the hypergeometric function with non-singular lower Pochhammer parameters to find the  $\epsilon$ -expansion of the starting function. In our method, each of the coefficient of  $\epsilon$  in the series expansion of a multivariable hypergeometric function is linear combination of hypergeometric functions with same domain of convergence as the original one. These analytic expressions can be used to find the numerical value at given point in its argument space within the domain of convergence. The summation of such multi-fold series has to be performed with sufficient number of terms for the required precision.

This methodology allows one to find series expansion of any multivariable hypergeometric function related to the Feynman integrals. To illustrate the approach we have presented examples containing one variable Gauss  ${}_{2}F_{1}$  and  ${}_{2}F_{2}$ , two variable Appell  $F_{1}$  and three variable Lauricella function, which is linked to two loop sunset Feynman integral with different masses. The examples given above are performed using Mathematica. In the present work many of the steps are done manually. An automated package for handling more complicated cases is under construction and will be published soon.

In this approach, higher order terms in the series expansion contain multivariable hypergeometric functions where all the variables are not independent. In such cases, it is possible to reduce the fold of the hypergeometric functions. Some reduction formulae of Kampé de Fériet series are discuss in [4]. Reduction formulae for a general multivariable hypergeometric functions are unknown and beyond the scope of the paper.

## Acknowledgments

The author would like to thank Prof. B. Ananthanarayan and Tanay Pathak for the discussions and comments on the draft. This is a part of author's doctoral work at CHEP, IISc.

# Appendices

## A Pochhammer identities

Pochhammer parameter is ratio of Euler's gamma functions,

$$(a)_m = \frac{\Gamma(a+m)}{\Gamma(a)} \tag{64}$$

Some useful identities of the Pochhammer parameters are listed below.

- $(a)_{m_1+\dots+m_n} = (a)_{m_1}(a+m_1)_{m_2}(a+m_1+m_2)_{m_3}\dots(a+m_1+\dots+m_{n-1})_{m_n}$
- $(a)_{-n} \to \frac{(-1)^n}{(1-a)_n}, \quad a \notin \mathbb{Z} \text{ and } n \in \mathbb{N}$
- Gauss's Multiplication theorem :

$$(a)_{km} = k^{km} \left(\frac{a}{k}\right)_m \left(\frac{a+1}{k}\right)_m \dots \left(\frac{a+k-1}{k}\right)_m \qquad \text{where } k \in \mathbb{N}^*$$

## **B** Notation used for multivariable hypergeometric functions

In this article we have used a common notation of hypergeometric functions for more than one variable. We denote a hypergeometric function with r fold summations with the set of variables  $\mathbf{x}$  in vector notation as,

$$F\begin{bmatrix} \{\mathbf{a},\mathbf{m}\} \\ \{\mathbf{b},\mathbf{n}\} \end{bmatrix} = \sum_{s_1=0}^{\infty} \cdots \sum_{s_r=0}^{\infty} \frac{(\mathbf{a})_{\mathbf{m}\cdot\mathbf{s}}}{(\mathbf{b})_{\mathbf{n}\cdot\mathbf{s}}} \frac{\mathbf{x}^{\mathbf{s}}}{\mathbf{s}!}$$
(65)

where  $\mathbf{m}, \mathbf{n}$  are vectors of length p and q respectively. Each of the elements of  $\mathbf{m}$  and  $\mathbf{n}$  themselves are vectors of length r with integers in their elements.

$$\mathbf{s} = \{s_1, \dots, s_r\} \qquad \{\mathbf{x}\} = \{x_1, \dots, x_r\} \\ \{\mathbf{a}, \mathbf{m}\} = \{a_1, \mathbf{m}_1\}, \dots, \{a_p, \mathbf{m}_p\} \qquad \{\mathbf{b}, \mathbf{n}\} = \{b_1, \mathbf{n}_1\}, \dots, \{b_q, \mathbf{n}_q\} \\ (\mathbf{a})_{\mathbf{m} \cdot \mathbf{s}} = (a_1)_{\mathbf{m}_1 \cdot \mathbf{s}} \dots (a_p)_{\mathbf{m}_p \cdot \mathbf{s}} \qquad (\mathbf{b})_{\mathbf{n} \cdot \mathbf{s}} = (b_1)_{\mathbf{n}_1 \cdot \mathbf{s}} \dots (b_q)_{\mathbf{n}_q \cdot \mathbf{s}} \\ \mathbf{m}_j \cdot \mathbf{s} = \sum_{i=0}^r (m_j)_i s_i \qquad j = 1, \dots, p \qquad \mathbf{n}_j \cdot \mathbf{s} = \sum_{i=0}^r (n_j)_i s_i \qquad j = 1, \dots, q \\ \mathbf{x}^{\mathbf{s}} = \prod_{i=1}^r x_i^{s_i} \qquad \mathbf{x}! = \prod_{i=1}^r (x_i!)$$

In this notation the Appell  $F_1$  can be written as

$$F_1(a, b_1, b_2, c, x, y) = \sum_{m,n=0}^{\infty} \frac{(a)_{m+n} (b_1)_m (b_2)_n}{(c)_{m+n}} \frac{x^m y^n}{m! n!}$$
$$= F \begin{bmatrix} \{a, \{1,1\}\}, \{b_1, \{1,0\}\}, \{b_2, \{0,1\}\} \\ \{c, \{1,1\}\} \end{bmatrix} \{x, y\} \end{bmatrix}$$

## C Numerical tests

In this Appendix we test the results obtained different examples in Section 3 numerically.

#### C.1 Gauss $_2F_1$ example with half integer Pochhammer

We compare our result from Section 3.1, Eq. (28) numerically with HypExp result Eq. (29) and the result obtained from NumExp below. The coefficients of the series expansion of  $\epsilon$  are evaluated at x = 0.3 and tabulate in Table 1. We observe that our results are numerically consistent with others.

	$\epsilon^0$	$\epsilon^1$	$\epsilon^2$	$\epsilon^3$
Eq. $(28)$	127.514	-38.7907	610.309	-189.456
result from HypExp	127.514	-38.7907	610.309	-189.456
result from NumExp	127.514	-38.7907	610.309	-189.528

Table 1: The numerical value of coefficients of  $\epsilon$  at x = 0.3 from Eq. (28), HypExp and NumExp are compared. The NumExp results are obtained with  $e_h = 10^{-4}$ .

#### C.2 Gauss $_2F_1$ example

We compare our result from Section 3.2, Eq. (34) numerically with HypExp result Eq. (35) and the result obtained from NumExp below. The coefficients of the series expansion of  $\epsilon$  are evaluated at x = 0.1 and tabulate in Table 2. We observe that our results are numerically consistent with others.

	$\epsilon^0$	$\epsilon^1$	$\epsilon^2$	$\epsilon^3$
Eq. (34)	1	0.0057506	0.104955	0.0998262
result from HypExp	1	0.0057506	0.104955	0.0998262
result from NumExp	1	0.0057506	0.104955	0.099735

Table 2: The numerical value of coefficients of  $\epsilon$  at x = 0.1 from Eq. (34), HypExp and NumExp are compared. The NumExp results are obtained with  $e_h = 10^{-4}$ .

#### C.3 The $_2F_2$ example

To check the result (i.e. Eq. (41)) of the  ${}_{2}F_{2}$  example, we multiply both sides of Eq. (41) by  $\epsilon^{2}$ . The right hand side becomes a series with non-negative powers of  $\epsilon$ .

$$\epsilon^2 {}_2F_2(1,2;\epsilon,\epsilon-1,x) = C_0 + C_1\epsilon + C_2\epsilon^2 + C_3\epsilon^3 + O(\epsilon^4)$$

Each of the coefficients with appropriate powers of  $\epsilon$  are evaluated at x = 0.1 with  $\epsilon = 10^{-4}$  and tabulated in Table 3.

$C_0$	-0.07305179768480031690
$C_1\epsilon$	$-1.795147633233522925^{*}10^{-5}$
$C_2 \epsilon^2$	$7.021511837197325735^{*}10^{-9}$
$C_3\epsilon^3$	$-1.52211510291782210*10^{-13}$

Table 3: The values of  $C_i \epsilon^i$ , for i = 0, 1, 2, 3 at x = 0.1 with  $\epsilon = 10^{-4}$  are shown.

The left hand side i.e.  $\epsilon^2 {}_2F_2(1,2;\epsilon,\epsilon-1,x)$  is calculated at the same x and  $\epsilon$  with sufficient number of terms and compared with the sum of  $C_0, C_1\epsilon, C_2\epsilon^2$  and  $C_3\epsilon^3$  in the Table 4.

$\epsilon^2 {}_2F_2(1,2;\epsilon,\epsilon-1,x)$	$C_0 + C_1\epsilon + C_2\epsilon^2 + C_3\epsilon^3$
-0.07306974213977305910	-0.0730697421397730264

Table 4: Comparison of the values of both sides of the series of  $\epsilon^2 {}_2F_2(1,2;\epsilon,\epsilon-1,x)$  upto  $\epsilon^3$  terms at x = 0.1 with  $\epsilon = 10^{-4}$ . The result from the series expansion upto  $\epsilon^3$  terms is in agreement with the value of  $\epsilon^2 {}_2F_2(\ldots,x)$ 

#### C.4 Appell $F_1$ example

We perform a test similar to the test done for  ${}_{2}F_{2}$  example (Appendix C.3). Here  $\epsilon$  is multiplied to the both sides of the Laurent expansion of the Appell function considered in Section 3.4. The right hand side becomes a series with non-negative powers of  $\epsilon$ .

$$\epsilon F_1(3/2, 2\epsilon + 1, 4 - \epsilon, \epsilon - 2, x, y) = C_0 + C_1\epsilon + C_2\epsilon^2 + O(\epsilon^3)$$

Each of the coefficients with appropriate powers of  $\epsilon$  is evaluated at (x, y) = (0.1, 0.2) with  $\epsilon = 10^{-3}$  with sufficient numbers of terms and tabulated in Table 5

$C_0$	$C_1\epsilon$	$C_2\epsilon^2$	
8.511583950834586989	0.00180071895404555128	0.00001320237872566561435	

Table 5: The values of  $C_i \epsilon^i$ , for i = 0, 1, 2 at x = 0.1 and y = 0.2 with  $\epsilon = 10^{-3}$  are shown.

The numerical value of  $\epsilon F_1(3/2, 2\epsilon + 1, 4 - \epsilon, \epsilon - 2, x, y)$  is calculated with same x, y and  $\epsilon$  and compared with sum the first three terms of its series expansion in Table 6.

$\epsilon F_1(3/2, 2\epsilon + 1, 4 - \epsilon, \epsilon - 2, x, y)$	$C_0 + C_1\epsilon + C_2\epsilon^2$
8.513397873773039908	8.51339787216735821

Table 6: Comparison of the values of both sides of the series of  $\epsilon F_1(3/2, 2\epsilon + 1, 4 - \epsilon, \epsilon - 2, x, y)$  upto  $\epsilon^2$  terms at (x, y) = (0.1, 0.2) with  $\epsilon = 10^{-3}$ . The value of the series expansion upto  $\epsilon^2$  terms is in agreement upto 8 decimal places with value of  $\epsilon F_1(\ldots, x, y)$ .

## C.5 Lauricella $F_C^{(3)}$ example

We compare the numerical value of the Taylor series expansion of the Lauricella function Eq. (63).

$$F_C^{(3)}(1,\epsilon,2-\epsilon,2-\epsilon,\epsilon,z_1,z_2,z_3) = C_0 + C_1\epsilon + O(\epsilon^2)$$
(66)

The first two terms in the series are calculated at  $(z_1, z_2, z_3) = (.1, .02, .03)$  with  $\epsilon = 10^{-3}$  and are tabulated in Table 7.

$C_0$	$C_1\epsilon$
1.023722811318515905790	0.00007405624680453594333

Table 7: The values of  $C_i \epsilon^i$ , for i = 0, 1 at  $(z_1, z_2, z_3) = (.1, .02, .03)$  with  $\epsilon = 10^{-3}$  are shown.

We compare the numerical value of  $F_C^{(3)}(1, \epsilon, 2 - \epsilon, 2 - \epsilon, \epsilon, z_1, z_2, z_3)$  and the sum of first two terms of its series expansion in Table 8.

$F_C^{(3)}\left(1,\epsilon,2-\epsilon,2-\epsilon,\epsilon,z_1,z_2,z_3\right)$	$C_0 + C_1 \epsilon$
1.023796910877940231962	1.023796867565320441419

Table 8: Comparison of the values of both sides of the series of  $F_C^{(3)}(1, \epsilon, 2 - \epsilon, 2 - \epsilon, \epsilon, z_1, z_2, z_3)$  upto  $\epsilon$  terms at  $(z_1, z_2, z_3) = (0.1, 0.02, 0.03)$  with  $\epsilon = 10^{-3}$ . The value of the series expansion upto  $\epsilon^2$  terms is in agreement upto 6 decimal places with value of the Lauricella function.

## References

- [1] H. Bateman. Higher Transcendental Functions. 1953.
- [2] L. Slater. Generalized Hypergeometric Functions. 1966.
- [3] H. Exton. Multiple hypergeometric functions and applications. 1976.
- [4] H. M. Srivastava and P. W. Karlsson. Multiple gaussian hypergeometric series. 1985.
- [5] V. A. Smirnov. Analytic tools for Feynman integrals. Vol. 250. 2012.
- [6] B. Ananthanarayan, S. Banik, S. Friot, and S. Ghosh. "Multiple Series Representations of N-fold Mellin-Barnes Integrals". Phys. Rev. Lett. 127.15 (2021), p. 151601. arXiv: 2012.15108 [hep-th].
- [7] "Negative dimensional integrals. I. Feynman graphs". Physics Letters B 193.2 (1987), pp. 241–246.
- [8] A. T. Suzuki. Massless two-loop "master" and three-loop two point function in NDIM. 2014. arXiv: 1408.4064 [math-ph].
- [9] I. Gonzalez and V. H. Moll. "Definite integrals by the method of brackets. Part 1". Advances in Applied Mathematics 45.1 (2010), pp. 50–73.
- [10] I. Gonzalez, V. H. Moll, and A. Straub. The method of brackets. Part 2: examples and applications. 2010. arXiv: 1004.2062 [math-ph].
- [11] B. Ananthanarayan, S. Banik, S. Friot, and T. Pathak. "On the Method of Brackets" (2021). arXiv: 2112.09679 [hep-th].
- [12] F. A. Berends, M. Buza, M. Bohm, and R. Scharf. "Closed expressions for specific massive multiloop selfenergy integrals". Z. Phys. C 63 (1994), pp. 227–234.
- [13] B. Ananthanarayan, S. Friot, and S. Ghosh. "New Series Representations for the Two-Loop Massive Sunset Diagram". Eur. Phys. J. C 80.7 (2020), p. 606. arXiv: 1911.10096 [hep-ph].
- [14] Z.-H. Gu, H.-B. Zhang, and T.-F. Feng. "Hypergeometric expression for a three-loop vacuum integral". Int. J. Mod. Phys. A 35.19 (2020), p. 2050089.
- [15] T.-F. Feng, C.-H. Chang, J.-B. Chen, Z.-H. Gu, and H.-B. Zhang. "Evaluating Feynman integrals by the hypergeometry". Nucl. Phys. B 927 (2018), pp. 516–549. arXiv: 1706.08201 [hep-ph].
- [16] T.-F. Feng, C.-H. Chang, J.-B. Chen, and H.-B. Zhang. "The system of partial differential equations for the C<sub>0</sub> function". Nucl. Phys. B 940 (2019), pp. 130–189. arXiv: 1809.00295 [hep-th].
- [17] O. V. Tarasov. "Functional reduction of one-loop Feynman integrals with arbitrary masses". JHEP 06 (2022), p. 155. arXiv: 2203.00143 [hep-ph].
- [18] L. de la Cruz. "Feynman integrals as A-hypergeometric functions". JHEP 12 (2019), p. 123. arXiv: 1907.00507 [math-ph].
- [19] R. P. Klausen. "Hypergeometric Series Representations of Feynman Integrals by GKZ Hypergeometric Systems". JHEP 04 (2020), p. 121. arXiv: 1910.08651 [hep-th].
- [20] T.-F. Feng, C.-H. Chang, J.-B. Chen, and H.-B. Zhang. "GKZ-hypergeometric systems for Feynman integrals". Nucl. Phys. B 953 (2020), p. 114952. arXiv: 1912.01726 [hep-th].
- [21] G. 't Hooft and M. J. G. Veltman. "Regularization and Renormalization of Gauge Fields". Nucl. Phys. B 44 (1972), pp. 189–213.
- [22] S. I. Bezrodnykh. "Horn's hypergeometric functions with three variables". Integral Transforms and Special Functions 32.3 (2021), 207–223. eprint: {https://doi.org/10.1080/10652469.2020.1814770}.
- [23] S. I. Bezrodnykh. "Analytic continuation of the Horn hypergeometric series with an arbitrary number of variables". Integral Transforms and Special Functions 31.10 (2020), pp. 788–803. eprint: https://doi.org/10.1080/10652469.
- [24] S. I. Bezrodnykh. "Analytic continuation of the Lauricella function F<sub>D</sub><sup>(N)</sup> with arbitrary number of variables". Integral Transforms and Special Functions 29.1 (2018), pp. 21–42. eprint: https://doi.org/10.1080/10652469.2017.1402
- [25] S. I. Bezrodnykh. "Analytic continuation of Lauricella's function  $F_D^{(N)}$  for variables close to unit near hyperplanes  $\{z_j = z_l\}$ ". Integral Transforms and Special Functions 0.0 (2021), pp. 1–15. eprint: https://doi.org/10.1080/1065
- [26] S. I. Bezrodnykh. "Analytic continuation of Lauricella's functions  $F_A^{(N)}$ ,  $F_B^{(N)}$ , and  $F_D^{(N)}$ ". Integral Transforms and Special Functions 31.11 (2020), 921–940. eprint: {https://doi.org/10.1080/10652469.2020.1762081}.
- [27] B. Ananthanarayan, S. Bera, S. Friot, and T. Pathak. "Olsson.wl : a *Mathematica* package for the computation of linear transformations of multivariable hypergeometric functions" (2021). arXiv: 2201.01189 [cs.MS].

- [28] B. Ananthanarayan, S. Bera, S. Friot, O. Marichev, and T. Pathak. "On the evaluation of the Appell F<sub>2</sub> double hypergeometric function" (2021). arXiv: 2111.05798 [math.CA].
- [29] S. Friot and G. Suchet-Bernard. "Linear transformations of Srivastava's  $H_C$  triple hypergeometric function" (2022). arXiv: 2205.06247 [math-ph].
- [30] B. Ananthanarayan, S. Bera, S. Friot, and T. Pathak. "work in progress".
- [31] B. Ananthanarayan, S. Bera, and S. Friot. "work in progress".
- [32] M. Y. Kalmykov. "Gauss hypergeometric function: Reduction, epsilon-expansion for integer/half-integer parameters and Feynman diagrams". JHEP 04 (2006), p. 056. arXiv: hep-th/0602028.
- [33] M. Y. Kalmykov, B. F. L. Ward, and S. A. Yost. "On the all-order epsilon-expansion of generalized hypergeometric functions with integer values of parameters". *JHEP* 11 (2007), p. 009. arXiv: 0708.0803 [hep-th].
- [34] M. Y. Kalmykov and B. A. Kniehl. "Towards all-order Laurent expansion of generalized hypergeometric functions around rational values of parameters". *Nucl. Phys. B* 809 (2009), pp. 365–405. arXiv: 0807.0567 [hep-th].
- [35] D. Greynat and J. Sesma. "A new approach to the epsilon expansion of generalized hypergeometric functions". Comput. Phys. Commun. 185 (2014), pp. 472–478. arXiv: 1302.2423 [math-ph].
- [36] T. Huber and D. Maitre. "HypExp: A Mathematica package for expanding hypergeometric functions around integer-valued parameters". Comput. Phys. Commun. 175 (2006), pp. 122–144. arXiv: hep-ph/0507094.
- [37] T. Huber and D. Maitre. "HypExp 2, Expanding Hypergeometric Functions about Half-Integer Parameters". Comput. Phys. Commun. 178 (2008), pp. 755–776. arXiv: 0708.2443 [hep-ph].
- [38] Z.-W. Huang and J. Liu. "NumExp: Numerical epsilon expansion of hypergeometric functions". Comput. Phys. Commun. 184 (2013), pp. 1973–1980. arXiv: 1209.3971 [physics.comp-ph].
- [39] D. Greynat, J. Sesma, and G. Vulvert. "Epsilon expansion of Appell and Kampé de Fériet functions" (2013). arXiv: 1310.7700 [math-ph].
- [40] D. Greynat, J. Sesma, and G. Vulvert. "Derivatives of the Pochhammer and reciprocal Pochhammer symbols and their use in epsilon-expansions of Appell and Kampé de Fériet functions". J. Math. Phys. 55 (2014), p. 043501.
- [41] S. A. Yost, V. V. Bytev, M. Y. Kalmykov, B. A. Kniehl, and B. F. L. Ward. "The Epsilon Expansion of Feynman Diagrams via Hypergeometric Functions and Differential Reduction" (2011). arXiv: 1110.0210 [math-ph].
- [42] V. V. Bytev, M. Y. Kalmykov, and B. A. Kniehl. "When epsilon-expansion of hypergeometric functions is expressible in terms of multiple polylogarithms: the two-variables examples". *PoS* LL2012 (2012). Ed. by T. Riemann, p. 029. arXiv: 1212.4719 [hep-th].
- [43] M. Kalmykov, V. Bytev, B. A. Kniehl, S.-O. Moch, B. F. L. Ward, and S. A. Yost. "Hypergeometric Functions and Feynman Diagrams". Antidifferentiation and the Calculation of Feynman Amplitudes. 2020. arXiv: 2012.14492 [hep-th].
- [44] J. Blümlein, M. Saragnese, and C. Schneider. "Hypergeometric Structures in Feynman Integrals" (2021). arXiv: 2111.15501 [math-ph].
- [45] J. Bluemlein, M. Saragnese, and C. Schneider. "Computer Algebra and Hypergeometric Structures for Feynman Integrals" (2022). arXiv: 2207.08524 [math-ph].
- [46] L. U. Ancarani and G. Gasaneo. "Derivatives of any order of the Gaussian hypergeometric function  ${}_2F_1(a, b, c; z)$  with respect to the parameters a, b and c". Journal of Physics A: Mathematical and Theoretical 42.39 (2009), p. 395208.
- [47] L. U. Ancarani and G. Gasaneo. "Derivatives of any order of the hypergeometric function  ${}_{p}F_{q}(a_{1},...,a_{p};b_{1}...,b_{q};z)$  with respect to the parameters  $a_{i}$  and  $b_{i}$ ". Journal of Physics A: Mathematical and Theoretical 43.8 (2010).
- [48] H. Kang and C. An. "Differentiation formulas of some hypergeometric functions with respect to all parameters". Applied Mathematics and Computation 258 (2015), pp. 454–464.
- [49] B. X. Fejzullahu. "Parameter derivatives of the generalized hypergeometric function". Integral Transforms and Special Functions 28.11 (2017), pp. 781–788.
- [50] V. Sahai and A. Verma. "DERIVATIVES OF APPELL FUNCTIONS WITH RESPECT TO PARAME-TERS". Journal of Inequalities and Special Functions 6 (3 2015).
- [51] L. U. Ancarani, J. A. Del Punta, and G. Gasaneo. "Derivatives of Horn hypergeometric functions with respect to their parameters". Journal of Mathematical Physics 58.7 (2017), p. 073504. eprint: https://doi.org/10.1063/1.499405

- [52] V. V. Bytev and B. A. Kniehl. "Derivatives of any Horn-type hypergeometric functions with respect to their parameters". Nuclear Physics B 952 (2020), p. 114911.
- [53] N. Takayama. "Groebner basis and the problem of contiguous relation". Japan Journal of Applied Mathematics 6 (1989), pp. 147–160.
- [54] V. V. Bytev, M. Y. Kalmykov, and B. A. Kniehl. "HYPERDIRE, HYPERgeometric functions DIfferential REduction: MATHEMATICA-based packages for differential reduction of generalized hypergeometric functions *pFp-1*, *F*<sub>1</sub>,*F*<sub>2</sub>,*F*<sub>3</sub>,*F*<sub>4</sub>". Comput. Phys. Commun. 184 (2013), pp. 2332–2342. arXiv: 1105.3565 [math-ph].
- [55] V. V. Bytev, M. Y. Kalmykov, and S.-O. Moch. "HYPERgeometric functions DIfferential REduction (HY-PERDIRE): MATHEMATICA based packages for differential reduction of generalized hypergeometric functions: F<sub>D</sub> and F<sub>S</sub> Horn-type hypergeometric functions of three variables". Comput. Phys. Commun. 185 (2014), pp. 3041–3058. arXiv: 1312.5777 [math-ph].
- [56] F. Chyzak. "Gröbner Bases, Symbolic Summation and Symbolic Integration". Gröbner Bases and Applications. Cambridge University Press, 1998, pp. 32–60.
- [57] C. Koutschan. "A fast approach to creative telescoping". Mathematics in Computer Science 4.2 (2010), pp. 259–266.
- [58] C. Koutschan. "Advanced Applications of the Holonomic Systems Approach". ACM Commun. Comput. Algebra 43.3/4 (2010), p. 119.
- [59] D. Maître. "HPL, a Mathematica implementation of the harmonic polylogarithms". Computer Physics Communications 174.3 (2006), pp. 222–240. arXiv: 0507152v2 [arXiv:hep-ph].