SOME NEW CLASSES OF (ALMOST) PERFECT *c*-NONLINEAR PERMUTATIONS

MOHIT PAL

ABSTRACT. The concept of differential uniformity was recently extended to the c-differential uniformity. An interesting problem in this area is the construction of functions with low c-differential uniformity and a lot of research has been done in this direction in the recent past. Here, we present three classes of (almost) perfect c-nonlinear permutations over finite fields of even characteristic.

1. INTRODUCTION

Let \mathbb{F}_{q^n} be the finite field with q^n elements, where q is a prime power and n is a positive integer. We denote by $\mathbb{F}_{q^n}^*$, the multiplicative cyclic group of non-zero elements of \mathbb{F}_{q^n} . Let f be a function from \mathbb{F}_{q^n} to itself. Lagrange's interpolation formula allows us to write f as a polynomial in $\mathbb{F}_{q^n}[X]/(X^{q^n}-X)$ uniquely. Thus, we shall use the term function and polynomial for f interchangeably. A polynomial $f \in \mathbb{F}_{q^n}[X]$ is called a permutation polynomial if the associated mapping $c \mapsto f(c)$ permutes the elements of \mathbb{F}_{q^n} . Here we consider permutation polynomials of the shape $f(X) = \gamma X + \operatorname{Tr}_{q^n/q}(X^k)$, where $\operatorname{Tr}_{q^n/q} : \mathbb{F}_{q^n} \to \mathbb{F}_q$ is the trace map given by $\operatorname{Tr}_{q^n/q}(X) = \sum_{i=0}^{n-1} X^{q^i}$ and $\gamma \in \mathbb{F}_{q^n}^*$. These permutation polynomials are interesting because of their simple algebraic structure. Charpin and Kyureghyan [2] first studied the polynomials of the form f and gave a complete characterization of permutation polynomials for q = 2. In 2016, Kyureghyan and Zieve [10] studied permutation polynomials of the shape $X + \gamma \operatorname{Tr}_{q^n/q}(X^k)$ over \mathbb{F}_{q^n} for odd q and constructed several classes of permutation polynomials. Motivated by the results of [2, 10], Li et al. [12] studied polynomials of the form $\gamma X + \operatorname{Tr}_{q^n/q}(X^k)$ over finite fields of even characteristic. To avoid repetitive work from [2], the authors in [12] restricted themselves to the case q > 2 and constructed fifteen new classes of permutation polynomials. For more details on permutation polynomials of the shape $\gamma X + \text{Tr}_{a^n/a}(X^k)$, reader may refer to [2, 6, 9, 10, 12, 16] and references therein.

Permutation polynomials have wide applications in the areas of mathematics and engineering such as coding theory [3, 11], cryptography [15, 19] and combinatorial designs [4]. In symmetric key cryptography, permutation polynomials are widely applied to the design of block ciphers, where they are used in the design of so-called substitution boxes. In fact, permutation polynomials must be choosen in a way so as S-box becomes immune to some of the known attacks against block ciphers. For instance, the differential attack introduced by Biham and Shamir [1] is one of the most powerful attacks on block ciphers. In 1993, Nyberg [18] introduced the notion of differential uniformity to quantify the immunity of a function against differential attacks. Recently, Ellingsen et al. [5] generalised the notion of differential uniformity and defined the c-differential uniformity of a function f for any $c \in \mathbb{F}_{q^n}$. For any $a, c \in \mathbb{F}_{q^n}$, the c-derivative of f in the direction of a is given by ${}_cD_f(a, X) := f(X+a) - cf(X)$ for all $X \in \mathbb{F}_{q^n}$. For any $a, b \in \mathbb{F}_{q^n}$, the c-Difference Distribution Table (c-DDT) entry at

²⁰¹⁰ Mathematics Subject Classification. 11T06, 94A60.

Key words and phrases. Finite fields, permutation polynomials, c-differential uniformity.

M. PAL

point (a, b) of f is given by ${}_{c}\Delta_{f}(a, b) := |\{X \in \mathbb{F}_{q^{n}} | {}_{c}D_{f}(a, X) = b\}|$. The c-differential uniformity of f is defined as ${}_{c}\Delta_{f} := \max\{{}_{c}\Delta_{f}(a, b) | a, b \in \mathbb{F}_{q^{n}} \text{ and } a \neq 0 \text{ when } c = 1\}$. When ${}_{c}\Delta_{f} = 1, 2$, then the function f is called perfect c-nonlinear (PcN) function and almost perfect c-nonlinear (APcN) function, respectively. When c = 1, then the notions of c-DDT and c-differential uniformity coincide with the classical notions of DDT and differential uniformity, respectively.

Since the appearence of the notion of the *c*-differential uniformity, several functions having low *c*-differential uniformity have been constructed in the recent past. The most recent list of power maps along with their *c*-differential uniformity can be found in [13, Table 1], and the list of non-power maps with their *c*-differential uniformity is given in [13, Table 2] for $c \neq 1$. However, over finite fields of even characteristic, there are only few PcN and APcN functions, see, e.g., [5, 7, 8, 17, 20]. In this paper, we shall restrict ourselves to finite fields of even characteristic and study the *c*-differential uniformity of three classes of permutation polynomials of the shape $\gamma X + \text{Tr}_{q^n/q}(X^k)$ over \mathbb{F}_{q^n} for $c \in \mathbb{F}_{q^n} \setminus \{1\}$. The rationale behind excluding the case c = 1 is the fact that in this case linear term outside the trace function vanishes and the varible X remains only in the trace term. As a consequence, polynomials of the form $\gamma X + \text{Tr}_{q^n/q}(X^k)$ will have very high differential uniformity.

The rest of the paper is organised as follows. In Section 2, we consider the *c*-differential uniformity of the permutation polynomial $f(X) := \gamma X + \operatorname{Tr}_{q^{2n}/q}(X^{2^{i}(q+1)})$, where $\gamma \in \mathbb{F}_{q^{2}}^{*}$, $q = 2^{t}$ and i, n, t > 0 are integers. We show that f is PcN for all $c \in \mathbb{F}_{q^{2}} \setminus \mathbb{F}_{2}$ and if $\operatorname{gcd}(i,t) = 1$ then $_{c}\Delta_{f} \leq 2$ for all $c \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_{q^{2}}$. Since we are considering only permutation polynomials, and for c = 0 every permutation polynomial is PcN, we have excluded the trivial case c = 0 in rest of the paper. In Section 3, we shall show that the permutation polynomial $g(X) := \gamma X + \operatorname{Tr}_{q^{2n}/q}(X^{q^{2}+1})$, where $\gamma \in \mathbb{F}_{q}^{*}$, $q = 2^{t}$ and n, t > 0 are integers, is PcN for all $c \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_{2}$. Section 4 is devoted to the *c*-differential uniformity of the permutation polynomial $h(X) := \gamma X + \operatorname{Tr}_{q^{2n}/q}(X^{2^{i}(q^{2}+1)})$, where $\gamma \in \mathbb{F}_{q^{2}}^{*}$, $q = 2^{t}$ and i, n, t > 0 are integers and either *n* is even; or *n* is odd and $\operatorname{Tr}_{q^{2}/q}\left(\left(\frac{1}{\gamma}\right)^{2^{i+1}}\right)^{\frac{q^{-1}}{\gcd(2^{i+1}-1,2^{t}-1)}} \neq 1$. We show that *h* is

PcN for all $c \in \mathbb{F}_{q^2} \setminus \mathbb{F}_2$, and if gcd(i, t) = 1 then the *c*-differential uniformity of *h* is ≤ 2 for all $c \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_{q^2}$. Finally, we summarize the paper with some concluding remarks in Section 5.

2. The *c*-differential uniformity of the function f

In this section, we shall consider the *c*-differential uniformity of the function $f(X) = \gamma X + \operatorname{Tr}_{q^{2n}/q}(X^{2^i(q+1)})$, where $\gamma \in \mathbb{F}_{q^2}^*$, $q = 2^t$ and i, n, t > 0 are integers. It has been proved in [12, Theorem 3.1] that f is a permutation polynomial over $\mathbb{F}_{q^{2n}}$. We first recall a lemma that will be used throughout this paper.

Lemma 2.1. [14] Let $q = p^t$, *i* be a positive integer and $f(X) = X^{p^i} - \alpha X - \beta$ where $\alpha \in \mathbb{F}_q^*$ and $\beta \in \mathbb{F}_q$. Then, in the field \mathbb{F}_q , *f* has either zero, one or p^d roots where d = gcd(i, t).

The following theorem gives the *c*-differential uniformity of the function f for all $c \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_2$.

Theorem 2.2. Let $q = 2^t$ and $f(X) = \gamma X + \operatorname{Tr}_{q^{2n}/q}(X^{2^i(q+1)})$, where $\gamma \in \mathbb{F}_{q^2}^*$ and i, n, t > 0 are integers. Then

(i) f is PcN over $\mathbb{F}_{q^{2n}}$ for all $c \in \mathbb{F}_{q^2} \setminus \mathbb{F}_2$, (ii) ${}_c\Delta_f \leq 2$ for all $c \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_{q^2}$ when gcd(i, t) = 1. *Proof.* Let $c \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_2$. Recall that, for any $(a, b) \in \mathbb{F}_{q^{2n}} \times \mathbb{F}_{q^{2n}}$, the c-DDT entry $c\Delta_f(a, b)$ is given by the number of solutions $X \in \mathbb{F}_{q^{2n}}$ of the following equation.

$$(2.1) f(X+a) + cf(X) = b.$$

It is easy to see that when a = 0 then for each $b \in \mathbb{F}_{q^{2n}}$, Equation (2.1) has a unique solution $X \in \mathbb{F}_{q^{2n}}$. For $a \neq 0$, Equation (2.1) can be written as

(2.2)
$$\gamma X + \operatorname{Tr}_{q^{2n}/q}(X^{2^{i}(q+1)}) + C \cdot \operatorname{Tr}_{q^{2n}/q}((X^{q}a + a^{q}X)^{2^{i}}) = B,$$

where $B = C(b + \gamma a + \operatorname{Tr}_{q^{2n}/q}(a^{2^{i}(q+1)}))$ and $C = \frac{1}{1+c}$. Notice that, when $a \in \mathbb{F}_q^*$, then the above equation reduces to

$$B = \gamma X + \operatorname{Tr}_{q^{2n}/q}(X^{2^{i}(q+1)}) + a^{2^{i}}C \cdot \operatorname{Tr}_{q^{2n}/q}((X^{2^{i}})^{q} + X^{2^{i}}) = \gamma X + \operatorname{Tr}_{q^{2n}/q}(X^{2^{i}(q+1)}),$$

which has a unique solution for each $B \in \mathbb{F}_{q^{2n}}$ (and hence for each $b \in \mathbb{F}_{q^{2n}}$). Now, let $a \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_q$. Then, from Equation (2.2) we have

(2.3)
$$\gamma X + \operatorname{Tr}_{q^{2n}/q}(X^{2^{i}(q+1)}) + C \cdot \operatorname{Tr}_{q^{2n}/q}\left(((X^{q^{2}} + X)a^{q})^{2^{i}}\right) = B.$$

We shall now study the solutions of Equation (2.3), in three different cases, namely, $c \in \mathbb{F}_q \setminus \mathbb{F}_2$, $c \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$ and $c \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_{q^2}$, respectively.

Case 1 Let $c \in \mathbb{F}_q \setminus \mathbb{F}_2$. Now, let $u := \operatorname{Tr}_{q^{2n}/q}(X^{2^i(q+1)}) + C \cdot \operatorname{Tr}_{q^{2n}/q}\left(((X^{q^2} + X)a^q)^{2^i}\right)$. Then, from Equation (2.3), we have $X = \frac{u+B}{\gamma}$. Putting the value of X into Equation (2.3), we have

$$u = \frac{1}{\gamma^{2^{i}(q+1)}} \operatorname{Tr}_{q^{2n}/q} \left(\left(u^{2} + (B^{q} + B)u + B^{q+1} \right)^{2^{i}} \right) + CB_{1}$$
$$= \frac{1}{\gamma^{2^{i}(q+1)}} \operatorname{Tr}_{q^{2n}/q} (B^{2^{i}(q+1)}) + CB_{1},$$

where $B_1 = \operatorname{Tr}_{q^{2n}/q}\left(\left(\frac{(B^{q^2}+B)a^q}{\gamma}\right)^{2^i}\right)$ and the first equality holds as $u, \gamma^{q+1} \in \mathbb{F}_q$ and the second equality holds as $\operatorname{Tr}_{q^{2n}/q}(1) = 0$ and $\operatorname{Tr}_{q^{2n}/q}(B^q+B) = 0$. Thus, for $c \in \mathbb{F}_q \setminus \mathbb{F}_2$, Equation (2.3) has a unique solution for each $a \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_q$ and $b \in \mathbb{F}_{q^{2n}}$.

Case 2 Let $c \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$. Let $u := \operatorname{Tr}_{q^{2n}/q}(X^{2^i(q+1)})$ and $v := \operatorname{Tr}_{q^{2n}/q}\left(((X^{q^2} + X)a^q)^{2^i}\right)$. Then, from Equation (2.3), we have $X = \frac{u + Cv + B}{\gamma}$. Using the properties of the trace function, we shall now simplify u and v. Consider

$$\begin{split} u &= \frac{1}{\gamma^{2^{i}(q+1)}} \operatorname{Tr}_{q^{2n}/q} \left(\left(u^{2} + ((B+Cv)^{q} + B+Cv)u + C^{q+1}v^{2} + (B^{q}C+BC^{q})v + B^{q+1} \right)^{2^{i}} \right) \\ &= \frac{1}{\gamma^{2^{i}(q+1)}} \operatorname{Tr}_{q^{2n}/q} \left(\left(((BC^{q})^{q} + BC^{q})v + B^{q+1} \right)^{2^{i}} \right) \\ &= \frac{1}{\gamma^{2^{i}(q+1)}} \operatorname{Tr}_{q^{2n}/q} \left(B^{2^{i}(q+1)} \right), \end{split}$$

where the first equality holds because $u, v \in \mathbb{F}_q$ and $C \in \mathbb{F}_{q^2}$ and the second equality holds because $C^{q+1} \in \mathbb{F}_q$, $\operatorname{Tr}_{q^{2n}/q}((B+Cv)^{q \cdot 2^i}) = \operatorname{Tr}_{q^{2n}/q}((B+Cv)^{2^i})$ and the third equality holds as $\operatorname{Tr}_{q^{2n}/q}((BC^q)^{q\cdot 2^i}) = \operatorname{Tr}_{q^{2n}/q}((BC^q)^{2^i})$. Now consider v which is given by

$$v = \operatorname{Tr}_{q^{2n}/q} \left(\left(\left(\frac{u + Cv + B}{\gamma} \right)^{q^2} + \left(\frac{u + Cv + B}{\gamma} \right) \right)^{2^i} a^{q \cdot 2^i} \right) = B_1,$$

where the last equality holds as $u, v \in \mathbb{F}_q$ and $C, \gamma \in \mathbb{F}_{q^2}$. Thus, $c \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$, Equation (2.3) has a unique solution for each $a \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_q$ and $b \in \mathbb{F}_{q^{2n}}$.

Case 3 Let $c \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_{q^2}$. Again, let $u := \operatorname{Tr}_{q^{2n}/q}(X^{2^i(q+1)}), v := \operatorname{Tr}_{q^{2n}/q}\left(((X^{q^2} + X)a^q)^{2^i}\right)$. Then, from Equation (2.3), we have

(2.4)
$$X = \frac{u + Cv + B}{\gamma}.$$

We shall now simplify u and v. Consider u which is given by

$$u = \frac{1}{\gamma^{2^{i}(q+1)}} \operatorname{Tr}_{q^{2n}/q} \left((u^{2} + ((B+Cv)^{q} + B+Cv)u + C^{q+1}v^{2} + (BC^{q} + B^{q}C)v + B^{q+1})^{2^{i}} \right)$$

= $\frac{1}{\gamma^{2^{i}(q+1)}} \left(v^{2^{i+1}}\mathcal{C} + v^{2^{i}}\mathcal{B} + \operatorname{Tr}_{q^{2n}/q}(B^{2^{i}(q+1)}) \right),$

where $\mathcal{B} = \operatorname{Tr}_{q^{2n}/q}((BC^q + B^qC)^{2^i}), \mathcal{C} = \operatorname{Tr}_{q^{2n}/q}(C^{2^i(q+1)})$ and the first equality holds because $u, v \in \mathbb{F}_q$, and last equality holds as $\operatorname{Tr}_{q^{2n}/q}((B+Cv)^{q\cdot 2^i}) = \operatorname{Tr}_{q^{2n}/q}((B+Cv)^{2^i})$. Now consider v which is given by

$$v = \operatorname{Tr}_{q^{2n}/q} \left(\left(\left(\frac{u + Cv + B}{\gamma} \right)^{q^2} + \left(\frac{u + Cv + B}{\gamma} \right) \right)^{2^i} a^{q \cdot 2^i} \right) = v^{2^i} C_1 + B_1,$$

$$C_1 = \operatorname{Tr}_{q^{2n}/q} \left(\left(\frac{(C^{q^2} + C)a^q}{\gamma} \right)^{2^i} \right) \text{ and the last equality holds as } u, v \in \mathbb{F}_q \text{ and } \gamma \in \mathbb{F}_{q^2}.$$

When $C_1 = 0$, then $v = B_1$ and by putting the values of u and v in Equation (2.4), we get a unique solution

$$X = \frac{1}{\gamma^{2^{i}(q+1)+1}} \left((B_{1})^{2^{i+1}} \mathcal{C} + (B_{1})^{2^{i}} \mathcal{B} + \operatorname{Tr}_{q^{2n}/q}(B^{2^{i}(q+1)}) \right) + \frac{(CB_{1} + B)}{\gamma}$$

of Equation (2.3). In the event when $C_1 \neq 0$, consider the equation

$$v^{2^{i}} + C_{1}^{-1}v + B_{1}C_{1}^{-1} = 0.$$

Since gcd(i,t) = 1, from Lemma 2.1 the above equation can have at most two solutions $v \in \mathbb{F}_q$. Thus, for $c \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_{q^2}$, Equation (2.3) can have at most two solutions for each $a \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_q$ and $b \in \mathbb{F}_{q^{2n}}$. This completes the proof.

3. The *c*-differential uniformity of the function g

In this section, we shall consider the *c*-differential uniformity of the function $g(X) := \gamma X + \operatorname{Tr}_{q^{2n}/q}(X^{q^2+1})$, where $\gamma \in \mathbb{F}_q^*$, $q = 2^t$ and n, t > 0 are integers. In [12, Theorem 3.2], Li et al. proved that g is a permutation polynomial over $\mathbb{F}_{q^{2n}}$. The following theorem gives the *c*-differential uniformity of the function g for all $c \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_2$.

Theorem 3.1. Let $q = 2^t$ and $g(X) = \gamma X + \operatorname{Tr}_{q^{2n}/q}(X^{q^2+1})$, where $\gamma \in \mathbb{F}_q^*$ and n, t > 0 are integers. Then g is PcN over $\mathbb{F}_{q^{2n}}$ for all $c \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_2$

where

Proof. Let $c \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_2$. For any $(a, b) \in \mathbb{F}_{q^{2n}} \times \mathbb{F}_{q^{2n}}$ the c-DDT entry $c\Delta_g(a, b)$ is given by the number of solutions $X \in \mathbb{F}_{q^{2n}}$ of the following equation.

(3.1)
$$\gamma X + \operatorname{Tr}_{q^{2n}/q}(X^{q^2+1}) + C \cdot \operatorname{Tr}_{q^{2n}/q}((X^{q^2}a + a^{q^2}X)) = B,$$

where $B = C\left(b + \gamma a + \operatorname{Tr}_{q^{2n}/q}(a^{q^2+1})\right)$ and $C = \frac{1}{1+c}$. When $a \in \mathbb{F}_q$, then the above equation reduces to $\gamma X + \operatorname{Tr}_{q^{2n}/q}(X^{q^2+1}) = B$, which has a unique solution for each $B \in \mathbb{F}_{q^{2n}}$. Now, let $a \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_q$ then Equation (3.1) can be rewritten as

(3.2)
$$\gamma X + \operatorname{Tr}_{q^{2n}/q}(X^{q^2+1}) + C \cdot \operatorname{Tr}_{q^{2n}/q}((X^{q^4} + X)a^{q^2}) = B.$$

We shall split the analysis of the solutions of Equation (3.2) into three different cases, namely, $c \in \mathbb{F}_q \setminus \mathbb{F}_2$, $c \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$ and $c \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_{q^2}$, respectively.

Case 1 Let $c \in \mathbb{F}_q \setminus \mathbb{F}_2$. Now, let $u := \operatorname{Tr}_{q^{2n}/q}(X^{q^2+1}) + C \cdot \operatorname{Tr}_{q^{2n}/q}((X^{q^4}+X)a^{q^2})$. Then, from Equation (3.2), we have $X = \frac{u+B}{\gamma}$. Putting the value of X into Equation (3.2), we have

$$u = \frac{1}{\gamma^2} \operatorname{Tr}_{q^{2n}/q} (u^2 + (B^{q^2} + B)u + B^{q^2 + 1}) + C \cdot \operatorname{Tr}_{q^{2n}/q} \left(\left(\frac{B^{q^4} + B}{\gamma} \right) a^{q^2} \right)$$
$$= \frac{1}{\gamma^2} \operatorname{Tr}_{q^{2n}/q} (B^{q^2 + 1}) + CB_1,$$

where $B_1 := \frac{1}{\gamma} \operatorname{Tr}_{q^{2n}/q} \left((B^{q^4} + B)a^{q^2} \right)$ and equalities hold because $u, \gamma \in \mathbb{F}_q$. Thus, for $c \in \mathbb{F}_q \setminus \mathbb{F}_2$, Equation (3.2) has a unique solution for each $a \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_q$ and $b \in \mathbb{F}_{q^{2n}}$. **Case 2** Let $c \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$. In this case, let $u := \operatorname{Tr}_{q^{2n}/q}(X^{(q^2+1)})$ and $v := \operatorname{Tr}_{q^{2n}/q}((X^{q^4} + X)a^{q^2})$. Then, from Equation (3.2), we have $X = \frac{u + Cv + B}{\gamma}$. Using properties of the trace function, we shall now simplify u and v. Consider u which is given by

$$u = \frac{1}{\gamma^2} \operatorname{Tr}_{q^{2n}/q} (u^2 + (Cv)^2 + (B^{q^2} + B)u + ((BC)^{q^2} + BC)v + B^{q^2+1})$$

= $\frac{1}{\gamma^2} \left(v^2 \operatorname{Tr}_{q^{2n}/q} (C^2) + \operatorname{Tr}_{q^{2n}/q} (B^{q^2+1}) \right),$

where equalities hold because $u, v, \gamma \in \mathbb{F}_q$, $\operatorname{Tr}_{q^{2n}/q}(B^{q^2}+B) = 0$ and $\operatorname{Tr}_{q^{2n}/q}((BC)^{q^2}+BC) = 0$. Now consider v which is given by

$$v = \operatorname{Tr}_{q^{2n}/q}\left(\left(\left(\frac{u+Cv+B}{\gamma}\right)^{q^4} + \left(\frac{u+Cv+B}{\gamma}\right)\right)a^{q^2}\right) = B_1$$

where last equality holds as $u, v, \gamma \in \mathbb{F}_q$ and $C \in \mathbb{F}_{q^2}$. Thus, for $c \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$, Equation (3.2) has a unique solution for each $a \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_q$ and $b \in \mathbb{F}_{q^{2n}}$. **Case 3** Let $c \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_{q^2}$. Again, let $u := \operatorname{Tr}_{q^{2n}/q}(X^{(q^2+1)})$ and $v := \operatorname{Tr}_{q^{2n}/q}((X^{q^4}+X)a^{q^2})$. Then, from Equation (3.2), we have

(3.3)
$$X = \frac{u + Cv + B}{\gamma}$$

Now, consider u which is given by

$$u = \frac{1}{\gamma^2} \operatorname{Tr}_{q^{2n}/q} \left(u^2 + ((B + Cv)^{q^2} + B + Cv)u + C^{q^2 + 1}v^2 + (BC^{q^2} + B^{q^2}C)v + B^{q^2 + 1} \right)$$

= $\frac{1}{\gamma^2} \left(v^2 \mathcal{C} + v\mathcal{B} + \operatorname{Tr}_{q^{2n}/q}(B^{q^2 + 1}) \right),$

where $\mathcal{B} = \operatorname{Tr}_{q^{2n}/q}(BC^{q^2} + B^{q^2}C), C = \operatorname{Tr}_{q^{2n}/q}(C^{q^2+1})$ and equalities hold because $u, v \in \mathbb{F}_q$ and $\operatorname{Tr}_{q^{2n}/q}((B+Cv)^{q^2} + B + Cv) = 0$. Now consider v which is given by

$$v = \operatorname{Tr}_{q^{2n}/q}\left(\left(\left(\frac{u+Cv+B}{\gamma}\right)^{q^4} + \left(\frac{u+Cv+B}{\gamma}\right)\right)a^{q^2}\right) = vC_1 + B_1,$$

where $C_1 = \frac{1}{\gamma} \operatorname{Tr}_{q^{2n}/q} \left((C^{q^4} + C)a^{q^2} \right)$ and the last equality holds as $u, v, \gamma \in \mathbb{F}_q$. From above equation we get a unique solution for v. Thus, for $c \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_{q^2}$, Equation (3.2) has a unique solution for each $a \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_q$ and $b \in \mathbb{F}_{q^{2n}}$. This completes the proof. \Box

4. The *c*-differential uniformity of the function h

In this section, we shall consider the *c*-differential uniformity of a more general class of permutations than *g*. Let $h(X) = \gamma X + \operatorname{Tr}_{q^{2n}/q}(X^{2^i(q^2+1)})$, where $\gamma \in \mathbb{F}_{q^2}^*$, $q = 2^t$ and i, n, t > 0 are integers. It has been proved in [12, Theorem 3.3] that *h* is a permutation polynomial over $\mathbb{F}_{q^{2n}}$ if either *n* is even; or *n* is odd and $\operatorname{Tr}_{q^2/q}\left(\frac{1}{\gamma^{2^i+1}}\right)^{\frac{q-1}{\gcd(2^{i+1}-1,2^t-1)}} \neq 1$. The following theorem gives the *c*-differential uniformity of the function *h* for all $c \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_2$.

Theorem 4.1. Let $q = 2^t$ and $h(X) = \gamma X + \operatorname{Tr}_{q^{2n}/q}(X^{2^i(q^2+1)})$, where $\gamma \in \mathbb{F}_{q^2}^*$, i, n, t > 0 are integers and either n is even; or n is odd and $\operatorname{Tr}_{q^2/q}\left(\frac{1}{\gamma^{2^{i+1}}}\right)^{\frac{q-1}{\gcd(2^{i+1}-1,2^{t-1})}} \neq 1$. Then

- (i) h is PcN over $\mathbb{F}_{q^{2n}}$ for all $c \in \mathbb{F}_{q^2} \setminus \mathbb{F}_2$,
- (ii) $_{c}\Delta_{h} \leq 2$ for all $c \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_{q^{2}}$ when gcd(i, t) = 1.

Proof. Let $c \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_2$. For any $(a, b) \in \mathbb{F}_{q^{2n}} \times \mathbb{F}_{q^{2n}}$ the c-DDT entry $c\Delta_h(a, b)$ is given by the number of solutions $X \in \mathbb{F}_{q^{2n}}$ of the following equation.

(4.1)
$$\gamma X + \operatorname{Tr}_{q^{2n}/q}(X^{2^{i}(q^{2}+1)}) + C \cdot \operatorname{Tr}_{q^{2n}/q}((X^{q^{2}}a + a^{q^{2}}X)^{2^{i}}) = B,$$

where $B = C(b + \gamma a + \operatorname{Tr}_{q^{2n}/q}(a^{2^{i}(q^{2}+1)}))$ and $C = \frac{1}{1+c}$. Notice that when $a \in \mathbb{F}_{q}$, Equation (4.1) reduces to $B = \gamma X + \operatorname{Tr}_{q^{2n}/q}(X^{2^{i}(q^{2}+1)})$, which has a unique solution for each $B \in \mathbb{F}_{q^{2n}}$. Now, let $a \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_{q}$. Then, from Equation (4.1) we have

(4.2)
$$\gamma X + \operatorname{Tr}_{q^{2n}/q}(X^{2^{i}(q^{2}+1)}) + C \cdot \operatorname{Tr}_{q^{2n}/q}\left(\left((X^{q^{4}} + X)a^{q^{2}}\right)^{2^{i}}\right) = B.$$

Now we shall split the analysis of the solutions of Equation (4.2) into three different cases, namely, $c \in \mathbb{F}_q \setminus \mathbb{F}_2$, $c \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$ and $c \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_{q^2}$, respectively.

Case 1 Let
$$c \in \mathbb{F}_q \setminus \mathbb{F}_2$$
 and $u := \operatorname{Tr}_{q^{2n}/q}(X^{2^i(q^2+1)}) + C \cdot \operatorname{Tr}_{q^{2n}/q}\left(\left((X^{q^4} + X)a^{q^2}\right)^{2^i}\right)$. Then, from Equation (4.2), we have $X = \frac{u+B}{\gamma}$. Putting the value of X into Equation (4.2), we

have

$$\begin{split} u &= \mathrm{Tr}_{q^{2n}/q} \left(\left(\frac{u^2 + (B^{q^2} + B)u + B^{q^2 + 1}}{\gamma^2} \right)^{2^i} \right) + C_1 \\ &= \mathrm{Tr}_{q^2/q} \left(\mathrm{Tr}_{q^{2n}/q^2} \left(\left(\frac{u^2 + (B^{q^2} + B)u + B^{q^2 + 1}}{\gamma^2} \right)^{2^i} \right) \right) + C_1 \\ &= \mathrm{Tr}_{q^2/q} \left(\left(\frac{u}{\gamma} \right)^{2^{i+1}} \mathrm{Tr}_{q^{2n}/q^2}(1) \right) + \mathrm{Tr}_{q^{2n}/q} \left(\left(\frac{B}{\gamma} \right)^{2^i(q^2 + 1)} \right) + C_1 \\ &= \begin{cases} B_1 + C_1 & \text{when } n \text{ is even} \\ u^{2^{i+1}}T_1 + B_1 + C_1 & \text{when } n \text{ is odd,} \end{cases} \end{split}$$

where
$$B_1 := \operatorname{Tr}_{q^{2n}/q}\left(\left(\frac{B}{\gamma}\right)^{2^i(q^2+1)}\right)$$
, $C_1 := C \cdot \operatorname{Tr}_{q^{2n}/q}\left(\left(\frac{(B^{q^4}+B)a^{q^2}}{\gamma}\right)^{2^i}\right)$ and $T_1 := \operatorname{Tr}_{q^2/q}\left(\frac{1}{\gamma^{2^{i+1}}}\right)$. Notice that when n is odd, $T_1^{\overline{\operatorname{gcd}(2^{i+1}-1,2^{t-1})}} \neq 1$ and hence we get only one solution $u \in \mathbb{F}_q$. Thus, for $c \in \mathbb{F}_q \setminus \mathbb{F}_2$, Equation (4.2) has a unique solution for each $a \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_q$ and $b \in \mathbb{F}_{q^{2n}}$.
Case 2 Let $c \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$. Let $u := \operatorname{Tr}_{q^{2n}/q}\left(X^{2^i(q^2+1)}\right)$ and $v := \operatorname{Tr}_{q^{2n}/q}\left(((X^{q^4}+X)a^{q^2})^{2^i}\right)$.
Then, from Equation (4.2), we have

(4.3)
$$X = \frac{u + Cv + B}{\gamma}.$$

Using the properties of the trace function, we shall now simplify u and v. Consider u which is given by

$$\begin{split} u &= \mathrm{Tr}_{q^{2n}/q} \left(\left(\frac{u^2 + (B^{q^2} + B)u + C^2 v^2 + ((BC)^{q^2} + BC)v + B^{q^2 + 1}}{\gamma^2} \right)^{2^i} \right) \\ &= \mathrm{Tr}_{q^2/q} \left(\frac{1}{\gamma^{2^{i+1}}} \mathrm{Tr}_{q^{2n}/q^2} \left(\left(u^2 + (B^{q^2} + B)u + C^2 v^2 + ((BC)^{q^2} + BC)v + B^{q^2 + 1} \right)^{2^i} \right) \right) \\ &= \mathrm{Tr}_{q^2/q} \left(\frac{1}{\gamma^{2^{i+1}}} \mathrm{Tr}_{q^{2n}/q^2} \left(\left(u^2 + C^2 v^2 + B^{q^2 + 1} \right)^{2^i} \right) \right) \\ &= \mathrm{Tr}_{q^2/q} \left(\frac{u^{2^{i+1}}}{\gamma^{2^{i+1}}} \mathrm{Tr}_{q^{2n}/q^2} \left(1 \right) + \frac{v^{2^{i+1}}}{\gamma^{2^{i+1}}} \mathrm{Tr}_{q^{2n}/q^2} \left(C^{2^{i+1}} \right) + \frac{1}{\gamma^{2^{i+1}}} \mathrm{Tr}_{q^{2n}/q^2} \left(B^{2^i(q^2 + 1)} \right) \right) \\ &= \begin{cases} v^{2^{i+1}}C_2 + B_1 & \text{when } n \text{ is even} \\ u^{2^{i+1}}T_1 + v^{2^{i+1}}C_2 + B_1 & \text{when } n \text{ is odd,} \end{cases}$$

where $C_2 := \operatorname{Tr}_{q^{2n}/q}\left(\left(\frac{C}{\gamma}\right)^{2^{i+1}}\right)$. Again, since $T_1^{\frac{q-1}{\gcd(2^{i+1}-1,2^{t}-1)}} \neq 1$ when n is odd, for any fixed v, we get a unique solution $u \in \mathbb{F}_q$ of the above equation. Now consider v which is given

by

$$v = \operatorname{Tr}_{q^{2n}/q}\left(\left(\left(\frac{u+Cv+B}{\gamma}\right)^{q^4} + \left(\frac{u+Cv+B}{\gamma}\right)\right)^{2^i} a^{q^22^i}\right) = \frac{C_1}{C}$$

Now putting the values of u, v into Equation (4.3), we get a unique solution $X \in \mathbb{F}_{q^{2n}}$ of Equation (4.2). Thus, for $c \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$, Equation (4.2) has a unique solution for each $a \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_q$ and $b \in \mathbb{F}_{q^{2n}}$.

Case 3 Let $c \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_{q^2}$. Again, in this case too, let $u := \operatorname{Tr}_{q^{2n}/q} \left(X^{2^i(q^2+1)} \right)$ and $v := \operatorname{Tr}_{q^{2n}/q} \left(\left((X^{q^4} + X)a^{q^2} \right)^{2^i} \right)$. Then, from Equation (4.2), we have $X = \frac{u + Cv + B}{\gamma}$. Now consider u which is given by

$$\begin{split} u &= \mathrm{Tr}_{q^{2n}/q} \left(\left(\frac{u^2 + ((B+Cv)^{q^2} + B + Cv)u + C^{q^2+1}v^2 + (C^{q^2}B + CB^{q^2})v + B^{q^2+1}}{\gamma^2} \right)^{2^i} \right) \\ &= \mathrm{Tr}_{q^2/q} \left(\frac{1}{\gamma^{2^{i+1}}} \mathrm{Tr}_{q^{2n}/q^2} \left((u^2 + C^{q^2+1}v^2 + (C^{q^2}B + CB^{q^2})v + B^{q^2+1})^{2^i} \right) \right) \\ &= u^{2^{i+1}} \mathrm{Tr}_{q^2/q} \left(\frac{1}{\gamma^{2^{i+1}}} \mathrm{Tr}_{q^{2n}/q^2}(1) \right) + v^{2^{i+1}} \mathrm{Tr}_{q^{2n}/q} \left(\left(\frac{C}{\gamma} \right)^{2^i(q^2+1)} \right) \\ &+ v^{2^i} \mathrm{Tr}_{q^{2n}/q} \left(\left(\frac{C^{q^2}B + CB^{q^2}}{\gamma^2} \right)^{2^i} \right) + \mathrm{Tr}_{q^{2n}/q} \left(\left(\frac{B}{\gamma} \right)^{2^i(q^2+1)} \right) \\ &= \begin{cases} v^{2^{i+1}}\mathcal{C} + v^{2^i}\mathcal{B} + B_1 & \text{when } n \text{ is even} \\ u^{2^{i+1}}T_1 + v^{2^{i+1}}\mathcal{C} + v^{2^i}\mathcal{B} + B_1 & \text{when } n \text{ is odd,} \end{cases} \end{split}$$

where
$$\mathcal{B} = \operatorname{Tr}_{q^{2n}/q}\left(\left(\frac{BC^{q^2} + B^{q^2}C}{\gamma^2}\right)^{2^i}\right), C = \operatorname{Tr}_{q^{2n}/q}\left(\left(\frac{C}{\gamma}\right)^{2^i(q^2+1)}\right)$$
 and these equalities

hold because $u, v \in \mathbb{F}_q$, $\gamma \in \mathbb{F}_{q^2}^*$ and $\operatorname{Tr}_{q^{2n}/q}((B+Cv)^{q^2}) = \operatorname{Tr}_{q^{2n}/q}(B+Cv)$. Now consider v which is given by

(4.5)
$$v = \operatorname{Tr}_{q^{2n}/q}\left(\left(\left(\frac{u+Cv+B}{\gamma}\right)^{q^4} + \left(\frac{u+Cv+B}{\gamma}\right)\right)a^{q^2}\right) = v^{2^i}C_3 + \frac{C_1}{C},$$

where $C_3 = \text{Tr}_{q^{2n}/q} \left(\left(\frac{(C^{q^4} + C)a^{q^2}}{\gamma} \right)^{2^*} \right)$ and the second equality holds as $u, v \in \mathbb{F}_q$ and $\gamma \in \mathbb{F}_{q^2}^*$. Now, if $C_3 = 0$, then we have a unique solution $v \in \mathbb{F}_q$ of Equation (4.5). Thus,

 $\gamma \in \mathbb{F}_{q^2}^*$. Now, if $C_3 = 0$, then we have a unique solution $v \in \mathbb{F}_q$ of Equation (4.5). Thus, Equation (4.2) has a unique solution in this case. If $C_3 \neq 0$, then Equation (4.5) can be rewritten as

$$v^{2^{i}} + \frac{v}{C_3} + \frac{C_1}{CC_3} = 0.$$

From Lemma 2.1, we know that if gcd(i, t) = 1, then the above equation can have at most two solutions $v \in \mathbb{F}_q$. Also, corresponding to each solution v of Equation (4.5), we get a unique solution u of Equation (4.4). Thus, for $c \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_{q^2}$, Equation (4.2) has at most two solutions for each $a \in \mathbb{F}_{q^{2n}} \setminus \mathbb{F}_q$ and $b \in \mathbb{F}_{q^{2n}}$. This completes the proof.

8

5. Conclusion

We have constructed three new classes of PcN permutations and two new classes of APcN permutations for some specific values of c. Since there are only few PcN and APcN permutations over finite fields of even characteristic, it would be an interesting problem to construct more such permutations.

References

- [1] E. Biham, A. Shamir, Differential cryptanalysis of DES-like cryptosystems, J. Cryptol. 4(1) (1991) 3–72.
- [2] P. Charpin, G. M. Kyureghyan, On a class of permutation polynomials over F_{2ⁿ}. In: Sequences and Their Applications - SETA 2008. Ed. by Solomon W. Golomb, Matthew G. Parker, Alexander Pott, and Arne Winterhof. Lect. Notes Comput. Sci. 5203. Berlin, Heidelberg: Springer Berlin Heidelberg, 2008, pp. 368–376.
- [3] C. Ding, T. Helleseth, Optimal ternary cyclic codes from monomials, IEEE Trans. Inf. Theory 59 (2013) 5898-5904.
- [4] C. Ding, J. Yuan, A family of skew Hadamard difference sets, J. Comb. Theory, Ser. A 113 (2006) 1526–1535.
- [5] P. Ellingsen, P. Felke, C. Riera, P. Stănică, A. Tkachenko, C-differentials, multiplicative uniformity, and (almost) perfect c-nonlinearity, IEEE Trans. Inf. Theory, 66(9) (2020) 5781–5789.
- [6] D. Gerike, G. M. Kyureghyan, Results on permutation polynomials of shape X^t + γTr_{qⁿ/q}(X^d). In: Combinatorics and Finite Fields. Ed. by Kai-Uwe Schmidt and Arne Winterhof. Radon Ser. Comput. Appl. Math. 23. Berlin, Boston: De Gruyter, 2019, pp. 67–78.
- [7] S. U. Hasan, M. Pal, P. Stănică, On the c-differential uniformity and boomerang uniformity of two classes of permutation polynomials, IEEE Trans. Inf. Theory, 68 (2022) 679–691.
- [8] J. Jeong, N. Koo, S. Kwon, On non-monomial APcN permutations over finite fields of even characteristic, ArXiv 2022. https://arxiv.org/abs/2205.11418.
- [9] G. M. Kyureghyan, Constructing permutations of finite fields via linear translators, J. Comb. Theory. Ser. A 118(3) (2011) 1052–1061.
- [10] G. M. Kyureghyan, M. E. Zieve, Permutation polynomials of the form $X + \gamma \text{Tr}(X^k)$. In: Contemporary Developments in Finite Fields and Applications. Ed. by Anne Canteaut, Gove Effinger, Sophie Huczynska, Daniel Panario, and Leo Storme. Singapore: World Scientific, 2016, pp. 178–194.
- [11] Y. Laigle-Chapuy, Permutation polynomials and applications to coding theory, Finite Fields Appl. 13 (2007) 58-70.
- [12] K. Li, L. Qu, X. Chen, C. Li, Permutation polynomials of the form $cx + \text{Tr}_{q^{\ell}/q}(x^a)$ and permutation trinomials over finite fields with even characteristic, Cryptogr. Commun. 10(3) (2018) 531–554.
- [13] C. Li, C. Riera, P. Stănică, Low c-differentially uniform functions via an extension of Dillon's switching method, ArXiv (2022). https://arxiv.org/abs/2204.08760.
- [14] J. Liang, On the solutions of trinomial equations over finite fields, Bull. Calcutta Math. Soc. 70 (1978) 379–382.
- [15] R. Lidl, W.B. Mullen, Permutation polynomials in RSA-cryptosystems, in: Advances in Cryptology, Plenum, New York (1984) 293–301.
- [16] J. Ma, G. Ge, A note on permutation polynomials over finite fields, Finite Fields Appl. 48 (2017) 261–270.
- [17] S. Mesnager, C. Riera, P. Stănică, H. Yan, Z. Zhou, Investigations on c-(almost) perfect nonlinear functions, IEEE Trans. Inf. Theory, 67 (2021) 6916–6925.
- [18] K. Nyberg, Differentially uniform mappings for cryptography. In: Advances in Cryptology-Eurocrypt'93.
 Ed. by Tor Helleseth. Lect. Notes Comput. Sci. 765 Springer, Berlin 1993, pp. 55–64.
- [19] J. Schwenk, K. Huber, Public key encryption and digital signatures based on permutation polynomials, Electron. Lett. 34 (1998) 759–760.
- [20] Z. Tu, X. Zeng, Y. Jiang, X. Tang, A class of APcN power functions over finite fields of even characteristic, arXiv (2021). https://arxiv.org/abs/2107.06464v1

DEPARTMENT OF INFORMATICS, UNIVERSITY OF BERGEN, PB 7803, 5020, BERGEN, NORWAY *Email address*: mohit.pal@uib.no