# Disproof of a conjecture on the minimum Wiener index of signed trees

Songlin Guo Wei Wang\* Chuanming Wang

School of Mathematics, Physics and Finance, Anhui Polytechnic University, Wuhu 241000, P. R. China

#### Abstract

The Wiener index of a connected graph is the sum of distances between all unordered pairs of vertices. Sam Spiro [The Wiener index of signed graphs, Appl. Math. Comput., 416(2022)126755] recently introduced the Wiener index for a signed graph and conjectured that the path  $P_n$  with alternating signs has the minimum Wiener index among all signed trees with n vertices. By constructing an infinite family of counterexamples, we prove that the conjecture is false whenever n is at least 30.

Keywords: Wiener index; signed tree; signed graph

AMS Classification: 05C09; 05C22

## 1 Introduction

A signed graph is a graph where each edge has a positive or negative sign. We usually write a signed graph as a pair  $(G, \sigma)$ , where G is the underlying graph and  $\sigma \colon E(G) \mapsto \{+1, -1\}$  describes the sign of each edge. For a path P in  $(G, \sigma)$ , the length of P (under the signing  $\sigma$ ) is  $\ell_{\sigma}(P) = |\Sigma_{e \in E(P)} \sigma(e)|$ . A path P in  $(G, \sigma)$  is called a uv-path if it has u and v as its endvertices. For two distinct vertices  $u, v \in V(G)$ , the signed distance [3] of u, v in  $(G, \sigma)$ , is

$$d_{\sigma}(u,v) = \min\{\ell_{\sigma}(P): P \text{ is a } uv\text{-path in } (G,\sigma)\}.$$

**Definition 1** ([3]). Let  $(G, \sigma)$  be a signed graph. The Wiener index of  $(G, \sigma)$ , denoted by  $W_{\sigma}(G)$ , is  $\sum d_{\sigma}(u, v)$ , where the summation is taken over all unordered pairs  $\{u, v\}$  of distinct vertices in G.

Let (G, +) denote a signed graph where each edge is positive. It is easy to see that the Wiener index  $W_+(G)$  coincides with the classic Wiener index W(G) of the ordinary graph G, introduced by Harry Wiener [5] in 1947. As the oldest topological index of a molecule, Wiener index has many applications in molecular chemistry, see the monograph [4].

<sup>\*</sup>Corresponding author. Email: wangwei.math@gmail.com

A tree is a connected graph with no cycles. There are numerous studies of properties of the Wiener indices of trees, see the survey paper [1]. Entringer, Jackson and Snyder [2] proved that, among all trees of any fixed order n, the path  $P_n$  (resp. the star  $K_{1,n}$ ) has the maximum (resp. minimum) Wiener index. Note that for any connected graph G together with any signing  $\sigma$ , we have  $W_{\sigma}(G) \leq W_{+}(G) = W(G)$ . Consequently, the above result of Entringer et al. indicates that  $W_{\sigma}(T) \leq W(P_n)$  for any signed n-vertex tree  $(T, \sigma)$ .

Let  $\sigma$  be a signing of the path  $P_n$ . We call  $\sigma$  (or  $(P_n, \sigma)$ ) alternating if any two adjacent edges have opposite signs. We usually use  $\alpha$  to denote an alternating signing of a path. The following interesting conjecture was proposed recently by Spiro [3].

Conjecture 1 ([3]). Among all signed trees of order n, the alternating path  $(P_n, \alpha)$  has the minimum Wiener index.

In this short note, we disprove Conjecture 1 by constructing infinite counterexamples.

**Theorem 1.** Conjecture 1 fails for every  $n \geq 30$ .

The proof of Theorem 1 is given at the end of the next section.

## 2 An infinite family of counterexamples

Let  $k \ge 0$  and  $a_1, a_2, \ldots, a_k$  be k nonnegative integers. Let  $T(a_1, a_2, \ldots, a_k)$  denote a rooted tree with  $1 + k + \sum_{i=1}^k a_i$  vertices constructing by the following two rules:

- (i) The root vertex has k neighbors  $u_1, u_2, \ldots, u_k$ ; such k vertices will be called branch vertices.
- (ii) For each  $i \in \{1, 2, ..., k\}$ , the branch vertex  $u_i$  has  $a_i$  neighbors other than the root vertex; such  $a_i$  neighbors will be called *leaf* vertices.

**Definition 2.** Let  $\sigma$  be a signing of a rooted tree  $T(a_1, a_2, \ldots, a_k)$ . We call  $\sigma$  *nice* if it satisfies the following two conditions:

- (i) Among k edges incident to the root vertex, the numbers of positive edges and negative edges differ by at most one.
- (ii) For each branch vertex u, all edges connecting u and leaf vertices have the same sign which is opposite to the sign of the edge connecting u and the root vertex.

Figure 1 illustrates a nice signing for the rooted tree T(3, 4, 4, 4, 4, 4), where we use dashed (resp. solid) lines to represent negative (resp. positive) edges.

**Theorem 2.** If  $\sigma$  is a nice then

$$W_{\sigma}(T(a_1, a_2, \dots, a_k)) = 2\sum_{i=1}^k \binom{a_i}{2} + 2\binom{\lfloor \frac{k}{2} \rfloor}{2} + 2\binom{\lceil \frac{k}{2} \rceil}{2} + k\left(1 + \sum_{i=1}^k a_i\right).$$

*Proof.* Write  $T = T(a_1, a_2, ..., a_k)$  and let P be any path in  $(T, \sigma)$ . Clearly, P contains at most four edges. Since  $\sigma$  is nice, one easily sees from Definition 2(ii) that any path in  $(T, \sigma)$  with 4 edges have exactly 2 positive edges and hence satisfies  $\ell_{\sigma}(P) = 0$ . Similarly, if P has




Figure 1: T(3,4,4,4,4,4) with a nice signing.

exactly 2 edges and  $\ell_{\sigma}(P) > 0$  then the two endvertices of P must be either two leaf vertices adjacent to a common branch vertex, or two branch vertices adjacent to the root vertex by two edges sharing the same sign. Note that the numbers of positive edges and negative edges are  $\lfloor \frac{k}{2} \rfloor$  and  $\lceil \frac{k}{2} \rceil$  (or in reverse order) by Definition 2(i). Thus, the contribution of such paths to  $W_{\sigma}(T)$  is

$$2\sum_{i=1}^{k} \binom{a_i}{2} + 2\binom{\lfloor \frac{k}{2} \rfloor}{2} + 2\binom{\lceil \frac{k}{2} \rceil}{2}.$$

Furthermore, noting that each path P with exactly one or three edges satisfies  $\ell_{\sigma}(P) = 1$  and there exists such a path between branch vertices and the remaining vertices, we see that the contribution of path with one or three edges is exactly

$$k\left(1+\sum_{i=1}^{k}a_{i}\right).$$

Adding the above two expressions completes the proof.

**Lemma 1.** Let  $\alpha$  be an alternating signing of  $P_n$ . Then  $W_{\alpha}(P_n) = \lfloor \frac{n}{2} \rfloor \lceil \frac{n}{2} \rceil$ .

Proof. Let (U, V) be the bipartition of  $P_n$  as a bipartite graph, where we assume  $|U| \leq |V|$ . Then  $|U| = \lfloor \frac{n}{2} \rfloor$  and  $|V| = \lceil \frac{n}{2} \rceil$ . Let u, v be any two vertices of  $P_n$ . It is easy to see that  $d_{\alpha}(u, v) = 0$  if u and v are in the same part, and  $d_{\alpha}(u, v) = 1$  otherwise. Thus,  $W_{\alpha}(P_n) = |U||V| = \lfloor \frac{n}{2} \rfloor \lceil \frac{n}{2} \rceil$ , as desired.

Noting that T(3, 4, 4, 4, 4, 4) has exactly 30 vertices, the following proposition gives a counterexample to Conjecture 1.

**Proposition 1.** Let  $\alpha$  be an alternating signing of  $P_{30}$  and  $\sigma$  be a nice signing of T = T(3, 4, 4, 4, 4, 4). Then  $W_{\sigma}(T) < W_{\alpha}(P_{30})$ .

*Proof.* Using Theorem 2 and Lemma 1, we find that  $W_{\sigma}(T) = 222$  while  $W_{\alpha}(P_{30}) = 225$ . Thus  $W_{\sigma}(T) < W_{\alpha}(P_{30})$ , as desired.

We shall show that for any  $n \geq 30$ , there exists a counterexample to Conjecture 1.

## Definition 3.

$$\mathcal{T}_k = \bigcup_{0 \le s \le k} \left\{ T(\underbrace{k-1, \dots, k-1}_{k-s}, \underbrace{k, \dots, k}_{s}), T(\underbrace{k, \dots, k}_{k-s}, \underbrace{k+1, \dots, k+1}_{s}) \right\}.$$

Note that  $\mathcal{T}_k$  contains exactly 2k+1 rooted trees of consecutive orders from  $k^2+1$  to  $(k+1)^2$ , see Figure 2 for the five rooted trees in  $\mathcal{T}_2$ .




Figure 2: The family  $\mathcal{T}_2$ .

**Lemma 2.** Let  $k \geq 10$  and T be any rooted tree in  $\mathcal{T}_k$ . Let n = |V(T)|. Then  $W_{\sigma}(T) < W_{\alpha}(P_n)$  where  $\sigma$  is nice while  $\alpha$  is alternating.

*Proof.* Write  $m = k^2 + 1$  and  $M = (k+1)^2$ . By Theorem 2 and Lemma 1, it is not difficult to see that both  $W_{\sigma}(T)$  and  $W_{\alpha}(P_n)$  are increasing as a function of n = |V(T)|. Thus we are done if we can show that  $W_{\sigma}(T_M) < W_{\alpha}(P_m)$  where  $T_M = T(\underbrace{k+1,\ldots,k+1})$ .

By Theorem 2 we have

$$W_{\sigma}(T_{M}) = 2k \binom{k+1}{2} + 2\binom{\lfloor \frac{k}{2} \rfloor}{2} + 2\binom{\lceil \frac{k}{2} \rceil}{2} + k(1+k(k+1))$$

$$< 2k \binom{k+1}{2} + 2\binom{\frac{k}{2}}{2} + 2\binom{\frac{k+1}{2}}{2} + k(1+k(k+1))$$

$$= 2k^{3} + \frac{5}{2}k^{2} + \frac{1}{2}k - \frac{1}{4}.$$
(1)

On the other hand, by Lemma 1, we have

$$W_{\alpha}(P_m) = \left\lfloor \frac{m}{2} \right\rfloor \left\lceil \frac{m}{2} \right\rceil = \left\lfloor \frac{k^2 + 1}{2} \right\rfloor \left\lceil \frac{k^2 + 1}{2} \right\rceil > \frac{1}{4}k^4.$$

It follows that

$$\frac{W_{\sigma}(T_M)}{W_{\alpha}(P_m)} < \frac{8}{k} + \frac{10}{k^2} + \frac{2}{k^3} - \frac{1}{k^4} < \frac{8}{k} + \frac{10}{k^2} + \frac{2}{k^3} \le \frac{8}{10} + \frac{10}{10^2} + \frac{2}{10^3} < 1.$$

Thus  $W_{\sigma}(T_M) < W_{\alpha}(P_m)$ , as desired. The proof is complete.

Proof of Theorem 1. Let  $\mathcal{T} = \bigcup_{k=0}^{\infty} \mathcal{T}_k$ . It is clear that  $\mathcal{T}$  contains exactly one *n*-vertex (rooted) tree for every positive integer n. We use  $T_n$  to denote the unique n-vertex tree in the family  $\mathcal{T}$ . Let  $\sigma$  be a nice signing of  $T_n$  and  $\alpha$  be an alternating signing of  $P_n$ . By Lemma 2, we see that  $W_{\sigma}(T_n) < W_{\alpha}(P_n)$  whenever  $n \geq 10^2 + 1$ . On the other hand, we

know from Proposition 1 that there does exist a 30-vertex tree T (with a nice signing  $\sigma$ ) such that  $W_{\sigma}(T) < W_{\alpha}(P_{30})$ . It remains to consider the case that  $n \in \{31, 32, \ldots, 100\}$ .

We claim that  $W_{\sigma}(T_n) < W_{\alpha}(P_n)$  for each  $n \in \{31, 32, \dots, 100\}$ . This can be checked directly using Theorem 2 and Lemma 1. Take n = 31 as an example. As  $31 \in [5^2+1, (5+1)^2]$ , we find that  $T_{31} \in \mathcal{T}_5$  and moreover  $T_{31} = T(5, 5, 5, 5, 5)$ . Using Theorem 2 for  $T_{31}$ , we obtain that  $W_{\sigma}(T_{31}) = 238$ . By Lemma 1, we have  $W_{\alpha}(P_{31}) = \lfloor \frac{31}{2} \rfloor \lceil \frac{31}{2} \rceil = 240$ . Thus  $W_{\sigma}(T_n) < W_{\alpha}(P_n)$  for n = 31. The proof is complete.

We remark that the counterexamples constructed in this note also disprove another conjecture of Spiro. For a graph G, the minimal signed Wiener index of G, denoted by  $W_*(G)$ , is the minimum of  $W_{\sigma}(G)$  for all possible signings  $\sigma$ . Spiro [3] conjectured that  $W_*(T) \geq W_*(P_n)$  for any n-vertex tree T. Let  $n \geq 30$  and  $T_n$  be the tree used in the proof of Theorem 1. Clearly,  $W_*(T_n) \leq W_{\sigma}(T_n)$ , where  $\sigma$  is a nice signing of  $T_n$ . On the other hand, it is easy to see that  $W_*(P_n) = W_{\alpha}(P_n)$ . Since  $W_{\sigma}(T_n) < W_{\alpha}(P_n)$ , we obtain  $W_*(T_n) < W_*(P_n)$ , disproving this conjecture.

# 3 Asymptotic property

It is still unknown which signed trees have the minimum Wiener index among all signed trees of a fixed order n. We use  $(\hat{T}_n, \hat{\sigma})$  to denote an n-vertex signed tree whose Wiener index is minimum among all signed trees of order n. And let  $(T_n, \sigma)$  be the n-vertex tree in  $\bigcup_{k=0}^{\infty} \mathcal{T}_k$  with a nice signing  $\sigma$ . One referee kindly points out that  $(T_n, \sigma)$  is optimal up to a constant factor. Precisely,

$$\limsup_{n \to \infty} \frac{W_{\sigma}(T_n)}{W_{\hat{\sigma}}(\hat{T}_n)} \le C,$$

for some constant C.

**Lemma 3.**  $W_{\sigma}(T_n) = (2 + o(1))n^{\frac{3}{2}}$ .

*Proof.* Let  $k = \lfloor \sqrt{n-1} \rfloor$ ,  $m = k^2 + 1$  and  $M = (k+1)^2$ . Then we have  $m \leq n \leq M$ . Note that  $T_m = T(\underbrace{k, \ldots, k}_k)$  and  $T_M = T(\underbrace{k+1, \ldots, k+1}_k)$ . Using Theorem 2, we have

$$W_{\sigma}(T_m) = 2k \binom{k}{2} + 2 \binom{\lfloor \frac{k}{2} \rfloor}{2} + 2 \binom{\lceil \frac{k}{2} \rceil}{2} + k(1+k^2) = (2+o(1))k^3$$
 (2)

and

$$W_{\sigma}(T_M) = 2k \binom{k+1}{2} + 2 \binom{\lfloor \frac{k}{2} \rfloor}{2} + 2 \binom{\lceil \frac{k}{2} \rceil}{2} + k(1+k(k+1)) = (2+o(1))k^3.$$
 (3)

Noting that  $k^3 \sim n^{\frac{3}{2}}$  and  $W_{\sigma}(T_m) \leq W_{\sigma}(T_n) \leq W_{\sigma}(T_M)$ , we have  $W_{\sigma}(T_n) = (2 + o(1))n^{\frac{3}{2}}$  by Squeeze Theorem.

The following lower bound is due to Sam Spiro.

**Lemma 4.**  $W_{\hat{\sigma}}(\hat{T}_n) \ge (\sqrt{2} + o(1))n^{\frac{3}{2}}$ .

*Proof.* Let U, V be the bipartition of  $\hat{T}_n$  with  $|U| \leq |V|$ . Label vertices in U as  $u_1, u_2, \ldots, u_k$ , where k = |U|. Let  $d_i^+$  (resp.  $d_i^-$ ) denote the number of positive (resp. negative) edges incident with  $u_i$  for each i. It is not too difficult to show that

$$W_{\hat{\sigma}}(\hat{T}_n) \ge |U||V| + 2\sum_{i=1}^k \left( \binom{d_i^+}{2} + \binom{d_i^-}{2} \right).$$
 (4)

Indeed, the first term comes from all paths of odd length and the term  $\binom{d_i^+}{2} + \binom{d_i^-}{2}$  comes from the paths of length 2 between two neighbors of  $u_i$  with the same sign. As the function  $\binom{x}{2} = \frac{1}{2}x(x-1)$  is convex, we have

$$\sum_{i=1}^{k} \left( \binom{d_i^+}{2} + \binom{d_i^-}{2} \right) \ge 2k \binom{\frac{1}{2k} \sum_{i=1}^{k} (d_i^+ + d_i^-)}{2}, \tag{5}$$

by Jensen's Inequality. As |U| = k, |V| = n - k and  $\sum_{i=1}^{k} (d_i^+ + d_i^-)$  equals n - 1, which is the number of edges in  $\hat{T}_n$ , we obtain from Eqs. (4) and (5) that

$$W_{\hat{\sigma}}(\hat{T}_n) \geq k(n-k) + 4k \binom{\frac{n-1}{2k}}{2}$$

$$= kn + \frac{n^2}{2k} - k^2 + \frac{1}{2k}((2k+1) - (2k+2)n)$$

$$\geq kn + \frac{n^2}{2k} - k^2 - 2n. \tag{6}$$

Using the basic inequality  $a + b \ge 2\sqrt{ab}$  for a, b > 0, we have

$$kn + \frac{n^2}{2k} \ge 2\sqrt{\frac{n^3}{2}} = \sqrt{2}n^{\frac{3}{2}}. (7)$$

Recall that  $k \leq n/2$ . Thus  $n - k \geq n/2$ . If  $k \geq 2\sqrt{2n}$  then from the trivial inequality  $W_{\hat{\sigma}}(\hat{T}_n) \geq k(n-k)$  we obtain

$$W_{\hat{\sigma}}(\hat{T}_n) \ge (2\sqrt{2n}) \cdot \frac{n}{2} = \sqrt{2n}^{\frac{3}{2}}.$$

Now assume  $k < 2\sqrt{2n}$ . Then by (6) and (7), we find

$$W_{\hat{\sigma}}(\hat{T}_n) > \sqrt{2}n^{\frac{3}{2}} - k^2 - 2n > \sqrt{2}n^{\frac{3}{2}} - 10n = (\sqrt{2} + o(1))n^{\frac{3}{2}}.$$

Thus we always have  $W_{\hat{\sigma}}(\hat{T}_n) \geq (\sqrt{2} + o(1))n^{\frac{3}{2}}$ , as desired.

The following theorem is a direct consequence of Lemmas 3 and 4.

### Theorem 3.

$$\limsup_{n \to \infty} \frac{W_{\sigma}(T_n)}{W_{\hat{\sigma}}(\hat{T}_n)} \le \sqrt{2}.$$

We end this note by leaving the following problem suggested by one referee.

**Problem 1.** Is it true that

$$\lim_{n \to \infty} \frac{W_{\sigma}(T_n)}{W_{\hat{\sigma}}(\hat{T}_n)} = 1?$$

## Acknowledgments

The authors would like to thank the anonymous reviewer for her/his instructive suggestions. In particular, the results in the final section are suggested by the reviewer. We thank Sam Spiro for pointing out a preliminary version of Lemma 4. The second author is supported by the National Natural Science Foundation of China under the grant number 12001006.

## References

- [1] A. A. Dobrynin, R. Entringer, I. Gutman, Wiener index of trees: Theory and applications, Acta Appl. Math. 66(2001) 211-249.
- [2] R. C. Entringer, D. E. Jackson, D. A. Snyder, Distance in graphs, Czechoslovak Math. J. 26(1976) 283-296.
- [3] S. Spiro, The Wiener index of signed graphs, Appl. Math. Comput., 416(2022)126755.
- [4] N. Trinajstić, Chemical Graph Theory, 2nd ed., CRC Press, 1992
- [5] H. Wiener, Structural determination of paraffin boiling points, J. Am. Chem. Soc. 69 (1947) 17-20.