

# Common Pairs of Graphs

Natalie Behague\*, Natasha Morrison† and Jonathan A. Noel‡

Department of Mathematics and Statistics, University of Victoria, Victoria, B.C., Canada.

{nbehague,nmorrison,noelj}@uvic.ca

December 5, 2022

## Abstract

A graph  $H$  is said to be *common* if the number of monochromatic labelled copies of  $H$  in a red/blue edge colouring of a large complete graph is asymptotically minimized by a random colouring with an equal proportion of each colour. We extend this notion to an asymmetric setting. That is, we define a pair  $(H_1, H_2)$  of graphs to be  $(p, 1 - p)$ -common if a particular linear combination of the density of  $H_1$  in red and  $H_2$  in blue is asymptotically minimized by a random colouring in which each edge is coloured red with probability  $p$  and blue with probability  $1 - p$ . We extend many of the results on common graphs to this asymmetric setting. In addition, we obtain several novel results for common pairs of graphs with no natural analogue in the symmetric setting. We also obtain new examples of common graphs in the classical sense and propose several open problems.

## 1 Introduction

Ramsey's Theorem [41] states that, for every graph  $H$ , there exists an integer  $N$  such that every colouring of the edges of the complete graph  $K_N$  with red and blue contains a monochromatic copy of  $H$ . The minimum such  $N$  is known as the *Ramsey number* of  $H$  and denoted  $R(H)$ . The problem of estimating Ramsey numbers is widely studied and notoriously difficult; see, e.g., the survey of Conlon, Fox and Sudakov [6].

The closely related *Ramsey multiplicity problem* asks for the asymptotics of the minimum possible number of monochromatic copies of  $H$  in a 2-edge colouring of  $K_N$  as  $N$  tends to

---

\*Supported by a PIMS Postdoctoral Fellowship.

†Research supported by NSERC Discovery Grant RGPIN-2021-02511, NSERC Early Career Supplement DGEER-2021-00047 and a Start-Up Grant from the University of Victoria.

‡Research supported by NSERC Discovery Grant RGPIN-2021-02460, NSERC Early Career Supplement DGEER-2021-00024 and a Start-Up Grant from the University of Victoria.

infinity [4,17,19]. While Ramsey multiplicity problems typically focus on counting monochromatic copies of a single graph, we will focus on the problem of minimizing the number of red copies of a graph  $H_1$  plus blue copies of another graph  $H_2$ , appropriately normalized and weighted. Asymmetric Ramsey multiplicity questions of this type were recently investigated for some specific weightings in the case that  $H_1 = K_3$  and  $H_2 \in \{K_4, K_5\}$  by Parczyk, Pokutta, Spiegel, and Szabó [39]. Our focus in this paper is more specifically on classifying pairs of graphs for which the minimum is attained by a random colouring in which each colour appears with a prescribed frequency. Such problems have a rich history starting with the beautiful result of Goodman [20, Theorem 1] which says that the number of monochromatic triangles in a 2-edge colouring of a large complete graph is at least  $(\frac{1}{4} - o(1)) \binom{n}{3}$ ; see Theorem 5.2. That is, the number of monochromatic triangles is asymptotically minimized by an unbiased random colouring.

In order to state our results, we require some notation and terminology. A *homomorphism* from a graph  $H$  to a graph  $G$  is a function  $f : V(H) \rightarrow V(G)$  such that  $f(u)f(v) \in E(G)$  whenever  $uv \in E(H)$ . We let  $\text{Hom}(H, G)$  denote the set of all homomorphisms from  $H$  to  $G$  and  $\text{hom}(H, G) := |\text{Hom}(H, G)|$ . The *homomorphism density* of  $H$  in  $G$  is defined by

$$t(H, G) := \frac{\text{hom}(H, G)}{v(G)^{v(H)}}$$

where  $v(F) = |V(F)|$  for any graph  $F$ . In other words,  $t(H, G)$  is the probability that a function from  $V(H)$  to  $V(G)$  is a homomorphism. Note that, if the number of vertices of  $G$  is large, then the deviation between  $\text{hom}(H, G)$  and the number of labelled copies of  $H$  in  $G$  is only  $O(v(G)^{v(H)-1})$ . Thus, the homomorphism density can be seen as an approximate measure of the number of copies of  $H$  in  $G$ .

Now, let  $H$  be a graph, suppose that  $N$  is large with respect to  $H$  and let  $G$  be an Erdős–Rényi graph on  $N$  vertices obtained by including each edge independently with probability  $1/2$ . Then  $t(H, G)$  and  $t(H, \overline{G})$  are both  $(1/2)^{e(H)} \pm o(1)$  with high probability, where  $\overline{G}$  is the complement of  $G$  and  $e(H) = |E(H)|$ . If we think of the edges of  $G$  as being red and the edges of  $\overline{G}$  as being blue, then this implies that the density of monochromatic copies of  $H$  is approximately  $(1/2)^{e(H)-1}$ . Following [25, p. 123], a graph  $H$  is said to be *common* if

$$t(H, G) + t(H, \overline{G}) \geq (1/2)^{e(H)-1} - o(1)$$

for any graph  $G$  on  $N$  vertices, where the  $o(1)$  term tends to zero as  $N$  tends to infinity. For example, the aforementioned result of Goodman [20] implies that  $K_3$  is common. In 1962, Erdős [13, Equation (3)] conjectured that all complete graphs are common. This conjecture fits nicely with one of the guiding philosophies of Ramsey theory, originating in a 1947 paper of Erdős [15], that colouring the edges of  $K_N$  randomly is often a good strategy for avoiding monochromatic structures. In 1980, Burr and Rosta [3] boldly extended Erdős’ conjecture to the statement that every graph  $H$  is common. Both of these conjectures turn out to be false; Sidorenko [48, p. 881] showed that a triangle with a pendant edge (the *paw* graph) is uncommon and Thomason [50] proved that  $K_4$  is uncommon. The latter result was extended by Jagger, Šťovíček and Thomason [25, Theorem 12] to the very general statement that any graph containing  $K_4$  is uncommon; see Theorem 3.17 for a further generalization.

The problem of classifying common graphs remains interesting and elusive. It is intimately linked to Sidorenko’s Conjecture [45], which says that, if  $H$  is bipartite, then

$$t(H, G) \geq t(K_2, G)^{e(H)} \tag{1.1}$$

for every graph  $G$ . We remark that a seemingly weaker, but technically equivalent, conjecture was made earlier by Erdős and Simonovits [14]. If  $H$  satisfies (1.1) for every  $G$ , then we say that  $H$  is *Sidorenko*. A simple convexity argument shows that, if  $H$  is Sidorenko, then  $H$  is common; see Corollary 4.10 for a generalization of this statement. Therefore, all of the positive results on Sidorenko’s Conjecture automatically yield examples of common graphs. Despite a great deal of recent progress [5, 7–9, 23, 27, 49], Sidorenko’s Conjecture remains open. Interestingly, there seem to be no examples of bipartite graphs which are currently known to be common and are not known to be Sidorenko.

Examples of non-bipartite common graphs are harder to come by. Sidorenko [48, Corollary 1] proved that all odd cycles are common, and Jagger, Šťovíček and Thomason [25] proved that even wheels are common. Sidorenko [46] showed that any graph obtained from gluing together triangles on vertices in a tree-like way is common, and an analogous result for gluing triangles along edges was proved in [25]. Recently, these two results were unified by Grzesik, Lee, Lidický and Volec [21, Theorem 1.1] to allow a mixture of gluing along vertices and edges; see Section 6 for several strengthenings of this statement. In addition, they proved that the so called “beachball graphs” obtained from even cycles by adding two vertices which are adjacent to every vertex of the cycle, but not to one another, are common. The first example of a common graph of chromatic number four, namely the 5-wheel, was obtained by Hatami, Hladký, Král’, Norine and Razborov in [24] using the powerful flag algebra method of [42]. Perhaps the most intriguing open problem in the area has been to determine whether there are common graphs of arbitrary large chromatic number; see [25, p. 124], [24, p. 739] and [6, Problem 2.28]. This was recently settled in the affirmative by Král’, Volec and Wei [32]; see also [29]. Analogous problems for more colours are also interesting [12, 25, 31]; see Section 8 for discussion of the multicolour case.

There has been some significant recent interest in analogues of Sidorenko’s Conjecture and the notion of common graphs in other areas, such as additive combinatorics [18, 26, 26, 43, 52, 53]. Here, the aim is to classify equations, or systems of equations, over  $\mathbb{F}_q$  with the property that the number of solutions to the system in a subset  $A \subseteq \mathbb{F}_q^n$  of a given size (in the Sidorenko problem) or a set  $A$  and its complement (in the commonness problem) is asymptotically minimized when  $A$  is chosen randomly.

Given the vast literature on off-diagonal Ramsey numbers (see [6, 40] for references), it is natural to wonder whether there is a sensible way to extend the notion of common graphs to an asymmetric setting. This question first came to our attention when it was asked by Hladký after a webinar talk given by the third author [38]. Specifically, Hladký’s question focused on pairs of graphs with the same number of edges and unbiased random colourings. In this paper, we consider the more general situation in which the two graphs can have different numbers of edges and the optimal colouring is random with different proportions of red and blue edges.

**Definition 1.2.** Let  $p_1, p_2 \in (0, 1)$  such that  $p_1 + p_2 = 1$  and let  $H_1$  and  $H_2$  be non-empty graphs. We say that  $(H_1, H_2)$  is  $(p_1, p_2)$ -common if, for any positive integer  $n$  and graphs  $G_1$  and  $G_2$  such that  $E(G_1) \cup E(G_2) = E(K_n)$ ,

$$\frac{t(H_1, G_1)}{e(H_1)p_1^{e(H_1)-1}} + \frac{t(H_2, G_2)}{e(H_2)p_2^{e(H_2)-1}} \geq \frac{p_1}{e(H_1)} + \frac{p_2}{e(H_2)} - o(1)$$

where the  $o(1)$  term tends to zero as  $n$  tends to infinity.

At first glance the coefficients in this expression may seem strange or arbitrary but this is in fact the only natural definition. Let  $c_1, c_2$  be real numbers and suppose that we wish to show that  $c_1 \cdot t(H_1, G) + c_2 \cdot t(H_2, \overline{G})$  is asymptotically minimized when  $G$  is a random graph with density  $p_1$ . By considering a random graph with a different density, say  $p_1 + x$  for  $-p_1 \leq x \leq p_2$ , we get a necessary condition on  $c_1$  and  $c_2$  that the function  $c_1(p_1 + x)^{e(H_1)} + c_2(p_2 - x)^{e(H_2)}$  must be minimized at  $x = 0$ . Setting the derivative of this function equal to zero at  $x = 0$  gives us that  $c_1 e(H_1) p_1^{e(H_1)-1} = c_2 e(H_2) p_2^{e(H_2)-1}$ . Thus, the coefficients in Definition 1.2 are the only possible choice, up to multiplying both of them by the same positive scalar.

Note also that by plugging  $p_1 = p_2 = 1/2$  and  $H_1 = H_2 = H$  into Definition 1.2, one immediately sees that a graph  $H$  is common if and only if the pair  $(H, H)$  is  $(1/2, 1/2)$ -common, and thus this definition does generalize the usual definition of common graphs.

Our goal in this paper is to initiate the study of  $(p_1, p_2)$ -common pairs of graphs. As with any new variation of an existing concept, there are several questions which naturally spring to mind, and we aim to address as many of these questions as we can. In Section 2, we translate this new notion into the natural setting of graph limits and build up some standard preliminaries. We prove some necessary conditions in Section 3. For instance, we show that, if  $(H_1, H_2)$  is  $(p, 1-p)$ -common for some  $p \in (0, 1)$ , then both of  $H_1$  and  $H_2$  must be  $K_4$ -free. We also show that, if  $H_1$  and  $H_2$  have odd girth, then there is at most one  $p$  for which  $(H_1, H_2)$  can be  $(p, 1-p)$ -common.

In Sections 4-7, we apply several different methods, all based on convexity (in one form or another), to obtain a wide range of examples of  $(p_1, p_2)$ -common pairs of graphs. In particular, in Section 6, we show that graphs obtained by gluing together cycles and paths in a tree-like way are  $(p_1, p_2)$ -common, under certain conditions. The proof involves two main ingredients: a binomial inequality comparing homomorphism densities of such graphs to homomorphism densities of odd cycles, proved in that section using entropy methods, and a strong form of the commonness property for  $K_3$  and  $C_5$  that is established in Section 5. These ingredients are fed into two general lemmas for propagating  $(p_1, p_2)$ -commonness from one pair of graphs to another proved in Sections 4 and 5. In Section 7, we analyse the set of  $p$  for which the pair  $(C_4, C_5)$  is  $(p, 1-p)$ -common. In particular, this set contains  $p = 1/2$  and  $p = 1/3$ , but does not contain any  $p \geq 0.518$ . We conclude the paper in Sections 8 and 9 by discussing a multicolour generalization and proposing several open problems.

## 2 Graph Limits

Many of the central problems and results in asymptotic extremal graph theory—especially those involving homomorphism densities—are most naturally understood in the language of graph limits. When writing about graph limits, we mainly follow the notation and terminology from the book of Lovász [35].

A *kernel* is a measurable function  $U : [0, 1]^2 \rightarrow \mathbb{R}$  which is bounded and *symmetric* in the sense that  $U(x, y) = U(y, x)$  for all  $x, y \in [0, 1]$ . A *graphon*  $W$  is a kernel such that  $0 \leq W(x, y) \leq 1$  for all  $x, y \in [0, 1]$ . Informally, a graphon can be thought of as a graph with vertex set  $[0, 1]$  such that, for  $x, y \in [0, 1]$ , the edge  $xy$  is assigned a weight of  $W(x, y)$ . A kernel can be thought of in similar terms, the only difference being that the edge weights of a kernel are not constrained to  $[0, 1]$ .

The notion of homomorphism density naturally extends from graphs to kernels by setting

$$t(H, U) = \int_{[0,1]^{V(H)}} \prod_{uv \in E(H)} U(x_u, x_v) dx_{V(H)}$$

for any graph  $H$  and kernel  $U$ , where  $x_{V(H)}$  is a vector  $(x_u : u \in V(H))$  of variables indexed by  $V(H)$ . Given a graph  $G$ , there is a natural graphon  $W_G$  corresponding to  $G$ ; simply take  $(A_v : v \in V(G))$  to be a partition of  $[0, 1]$  into  $v(G)$  sets of equal measure and, for  $(x, y) \in A_u \times A_v$ , we set  $W(x, y) = 1$  if  $uv \in E(G)$  and  $W(x, y) = 0$  otherwise. It is easily observed that  $t(H, G) = t(H, W_G)$  for any graphs  $H$  and  $G$ .

A sequence  $G_1, G_2, \dots$  of graphs is said to *converge* if the sequence  $t(H, G_1), t(H, G_2), \dots$  converges for every graph  $H$ . A graphon  $W$  is said to be the *limit* of the sequence if  $\lim_{n \rightarrow \infty} t(H, G_n) = t(H, W)$  for every graph  $H$ . Perhaps the most fundamentally important result in graph limits is the theorem of Lovász and Szegedy [36] which says that every convergent sequence of graphs has a limit. On the other hand, for every graphon  $W$ , there is a sequence of graphs whose limit is  $W$ . For this, it suffices to take a sequence  $G_1, G_2, \dots$  of graphs such that  $G_n$  is a graph on  $n$  vertices obtained by sampling  $n$  points  $x_1, x_2, \dots, x_n$  uniformly at random from  $[0, 1]$  and putting an edge from the  $i$ th vertex to the  $j$ th vertex with probability  $W(x_i, x_j)$ ; this construction is known as a *W-random graph*.

One benefit of dealing with graph limits as opposed to finite graphs is that smaller order asymptotic terms that are necessary in the setting of finite graphs tend to vanish in the limit. This often yields slightly more elegant expressions. For example, in the language of graph limits, a pair  $(H_1, H_2)$  is  $(p_1, p_2)$ -common if and only if

$$\frac{t(H_1, W_1)}{e(H_1)p_1^{e(H_1)-1}} + \frac{t(H_2, W_2)}{e(H_2)p_2^{e(H_2)-1}} \geq \frac{p_1}{e(H_1)} + \frac{p_2}{e(H_2)} \quad (2.1)$$

for any graphons  $W_1$  and  $W_2$  such that  $W_1 + W_2 = 1$ . Note that the  $o(1)$  term in Definition 1.2 is necessary for graphs, but not for graphons. For example, there exists a triangle-free graph  $G$  on five vertices such that  $\overline{G}$  is also triangle-free, and so  $t(K_3, G) + t(K_3, \overline{G}) = 0$  despite the fact that  $K_3$  is common.

We close this section with a simple fact about homomorphism densities of disjoint unions of graphs. Given two graphs  $F_1$  and  $F_2$ , let  $F_1 \sqcup F_2$  denote the vertex-disjoint union of  $F_1$  and  $F_2$ . Given a graph  $F$  and a positive integer  $m$ , let  $m \cdot F$  denote the graph  $\bigsqcup_{i=1}^m F$ .

**Lemma 2.2.** *For any graphs  $F_1$  and  $F_2$  and kernel  $U$ ,*

$$t(F_1 \sqcup F_2, U) = t(F_1, U)t(F_2, U).$$

*Proof.* We have

$$t(F_1 \sqcup F_2, U) = \int_{[0,1]^{V(F_1) \cup V(F_2)}} \prod_{uv \in E(F_1) \cup E(F_2)} U(x_u, x_v) dx_{V(F_1) \cup V(F_2)}.$$

This integral is clearly separable and, therefore, evaluates to  $t(F_1, U)t(F_2, U)$ .  $\square$

## 2.1 Algebraic Expansion

One approach to analyzing the homomorphism density of a graph  $H$  in a graphon  $W$  is to re-parameterize  $W$  as a constant function plus a perturbation and to expand the product inside of the integral in the definition of  $t(H, W)$ . This is a standard trick that is used in several papers on common graphs [11, 21, 22, 25, 46, 51] and Sidorenko's Conjecture [34].

Given  $p \in (0, 1)$  and a graphon  $W$ , consider the kernel  $U : [0, 1]^2 \rightarrow \mathbb{R}$  defined by  $U := W - p$ . Then

$$t(H, W) = \int_{[0,1]^{V(H)}} \prod_{uv \in E(H)} W(x_u, x_v) dx_{V(H)} = \int_{[0,1]^{V(H)}} \prod_{uv \in E(H)} (p + U(x_u, x_v)) dx_{V(H)}.$$

Given a graph  $H$  and a set  $E \subseteq E(H)$ , let  $H[E]$  be the graph with vertex set  $V(H)$  and edge set  $E$ . Expanding the product inside of the above integral yields

$$\int_{[0,1]^{V(H)}} \left( \sum_{E \subseteq E(H)} p^{e(H)-|E|} \prod_{uv \in E} U(x_u, x_v) \right) dx_{V(H)} = \sum_{E \subseteq E(H)} p^{e(H)-|E|} t(H[E], U).$$

So, we have derived the following nice formula that is valid for any graph  $H$ , graphon  $W$  and  $p \in (0, 1)$ .

**Observation 2.3.** Let  $W$  be a graphon and  $p \in (0, 1)$  and let  $U$  be the kernel defined by  $U := W - p$ . Then

$$t(H, W) = \sum_{E \subseteq E(H)} p^{e(H)-|E|} t(H[E], U). \quad (2.4)$$

Observation 2.3 leads us to an alternative definition of a  $(p_1, p_2)$ -common pair of graphs.

**Lemma 2.5.** *Let  $p_1, p_2 \in (0, 1)$  such that  $p_1 + p_2 = 1$  and let  $H_1$  and  $H_2$  be non-empty graphs. Then  $(H_1, H_2)$  is  $(p_1, p_2)$ -common if and only if*

$$\sum_{\substack{E \subseteq E(H_1) \\ |E| \geq 2}} \frac{t(H_1[E], U)}{p_1^{|E|-1} e(H_1)} + \sum_{\substack{E \subseteq E(H_2) \\ |E| \geq 2}} (-1)^{|E|} \frac{t(H_2[E], U)}{p_2^{|E|-1} e(H_2)} \quad (2.6)$$

is non-negative for any kernel  $U$  such that  $-p_1 \leq U(x, y) \leq p_2$  for all  $x, y \in [0, 1]$ .

*Proof.* Let  $W_1$  and  $W_2$  be graphons such that  $W_1 + W_2 = 1$  and let  $U = W_1 - p_1$ . Then  $W_1 = p_1 + U$  and  $W_2 = p_2 - U$ . Moreover, it holds that  $-p_1 \leq U(x, y) \leq 1 - p_1 = p_2$  for all  $x, y \in [0, 1]$  since  $W$  is a graphon. By Observation 2.3 and the fact that  $t(F, cU) = c^{e(F)} t(F, U)$  every kernel  $U$  and  $c \in \mathbb{R}$ , the expression

$$\frac{t(H_1, W_1)}{e(H_1) p_1^{e(H_1)-1}} + \frac{t(H_2, W_2)}{e(H_2) p_2^{e(H_2)-1}}$$

can be rewritten as

$$\sum_{E \subseteq E(H_1)} \frac{p_1^{e(H_1)-|E|} t(H_1[E], U)}{e(H_1) p_1^{e(H_1)-1}} + \sum_{E \subseteq E(H_2)} (-1)^{|E|} \frac{p_2^{e(H_2)-|E|} t(H_2[E], U)}{e(H_2) p_2^{e(H_2)-1}}$$

or, equivalently,

$$\sum_{E \subseteq E(H_1)} \frac{t(H_1[E], U)}{e(H_1) p_1^{|E|-1}} + \sum_{E \subseteq E(H_2)} (-1)^{|E|} \frac{t(H_2[E], U)}{e(H_2) p_2^{|E|-1}}.$$

The contribution of the term  $E = \emptyset$  to this expression is precisely  $\frac{p_1}{e(H_1)} + \frac{p_2}{e(H_2)}$ . Thus,  $(H_1, H_2)$  is  $(p_1, p_2)$ -common if and only if the other terms sum to a non-negative value for every possible choice of  $U$ . The edge sets of cardinality one contribute a total of 1 to the first summation and  $-1$  to the second, and so their contributions cancel. This completes the proof.  $\square$

It is well-known that some additional cancellations occur in the symmetric case  $p_1 = p_2 = 1/2$  and  $H_1 = H_2$  [11, 21, 22, 25, 46, 51]. The proof is similar to that of Lemma 2.5, and so we omit it.

**Lemma 2.7.** *A graph  $H$  is common if and only if the following holds for any kernel  $U$  such that  $\|U\|_\infty \leq 1$*

$$\sum_{\substack{\emptyset \neq E \subseteq E(H) \\ |E| \text{ is even}}} t(H[E], U) \geq 0.$$

### 3 Necessary Conditions

Our next goal is to use the algebraic expansion described in the previous section to derive some necessary conditions for a pair  $(H_1, H_2)$  to be  $(p_1, p_2)$ -common. For this, we will use several different operations on kernels. The first can be thought of as “scaling down” a kernel onto a  $\delta$ -proportion of the vertex set.

**Definition 3.1.** Given a kernel  $U$  and a real number  $\delta \in (0, 1]$ , let  $U^\delta$  be the kernel defined by

$$U^\delta(x, y) = \begin{cases} U(x/\delta, y/\delta) & \text{if } x, y \in [0, \delta], \\ 0 & \text{otherwise.} \end{cases}$$

Given a graph  $F$ , let  $i(F)$  be the number of isolated vertices of  $F$ .

**Lemma 3.2.** For any graph  $H$ , kernel  $U$  and  $\delta \in (0, 1]$ ,

$$t(F, U^\delta) = \delta^{v(F)-i(F)} t(F, U).$$

*Proof.* We may assume that  $F$  has no isolated vertices. Indeed, for any isolated vertex  $w$ , the variable corresponding to  $w$  does not appear in the product  $\prod_{uv \in E(F)} U(x_u, x_v)$ . So, by integrating out that variable over  $[0, 1]$ , we see that  $t(F, U) = t(F - w, U)$  for any kernel  $U$ , and so we can conclude by induction on  $v(F)$ .

So, assume that  $F$  has no isolated vertices. By definition,  $U^\delta(x, y) = 0$  whenever  $x$  or  $y$  is not in  $[0, \delta]$ . Therefore, if  $x_{V(F)} \in [0, 1]^{V(F)}$  such that there is a vertex  $v$  with  $x_v \notin [0, \delta]$ , then the product  $\prod_{uv \in E(F)} U^\delta(x_u, x_v)$  is automatically zero. Therefore,

$$\begin{aligned} t(F, U^\delta) &= \int_{[0, \delta]^{V(F)}} \prod_{uv \in E(F)} U^\delta(x_u, x_v) dx_{V(F)} = \int_{[0, \delta]^{V(F)}} \prod_{uv \in E(F)} U(x_u/\delta, x_v/\delta) dx_{V(F)} \\ &= \delta^{v(F)} t(F, U). \end{aligned}$$

This completes the proof. □

In order to simplify the expression in Lemma 2.5, it is often useful to choose  $U$  to be “regular” in a certain sense.

**Definition 3.3.** Given a kernel  $U$  and  $x \in [0, 1]$ , the *degree* of  $x$  is defined to be

$$d_U(x) := \int_0^1 U(x, y) dy.$$

**Definition 3.4.** Given a kernel  $U$  and  $d \in \mathbb{R}$ , we say that  $U$  is *d-regular* if  $d_U(x) = d$  for almost every  $x \in [0, 1]$ .

**Lemma 3.5.** If  $U$  is a  $d$ -regular kernel and  $F$  is a graph containing a vertex  $w$  of degree one, then

$$t(F, U) = d \cdot t(F - w, U).$$



*Proof.* Let  $z$  be the unique neighbour of  $w$  in  $F$ . We have

$$\begin{aligned} t(F, U) &= \int_{[0,1]^{V(F)}} \prod_{uv \in E(F)} U(x_u, x_v) dx_{V(F)} \\ &= \int_{[0,1]^{V(F) \setminus \{w\}}} \left( \int_0^1 U(x_z, x_w) dx_w \right) \prod_{uv \in E(F) \setminus \{wz\}} U(x_u, x_v) dx_{V(F) \setminus \{w\}} = d \cdot t(F, U) \end{aligned}$$

as desired.  $\square$

Using the above ideas, we obtain a simple constraint on the shortest cycles in  $(p_1, p_2)$ -common pairs. Let  $C_k$  denote the cycle of length  $k$ . Given a graph  $H$  and integer  $k \geq 3$ , we let  $c_k(H)$  be the number of cycles of length  $k$  in  $H$ . The *girth* of  $H$  is  $g(H) := \min\{k : c_k(H) \geq 1\}$ . In particular,  $g(H) = \infty$  if  $H$  is a forest.

**Theorem 3.6.** *Suppose that  $(H_1, H_2)$  is  $(p_1, p_2)$ -common. If  $g(H_1)$  is odd, then either  $g(H_2) < g(H_1)$  or  $g(H_1) = g(H_2) = k$  and*

$$\frac{c_k(H_1)}{p_1^{k-1}e(H_1)} = \frac{c_k(H_2)}{p_2^{k-1}e(H_2)}.$$

*Proof.* We prove the contrapositive. Suppose that  $g(H_1) = k$  where  $k$  is an odd integer and that  $g(H_2) \geq k$ . Note that, if  $g(H_2) > k$ , then  $c_k(H_2) = 0$ ; so, in any case, we can assume, without loss of generality, that

$$\frac{c_k(H_1)}{p_1^{k-1}e(H_1)} > \frac{c_k(H_2)}{p_2^{k-1}e(H_2)}. \quad (3.7)$$

Let  $I_1$  and  $I_2$  be intervals of length  $1/2$  partitioning  $[0, 1]$  and let  $B$  be the kernel defined by

$$B(x, y) = \begin{cases} -1 & \text{if } (x, y) \in I_i^2 \text{ for some } i \in \{1, 2\}, \\ 1 & \text{otherwise.} \end{cases}$$

Observe that the homomorphism density of a cycle  $C_\ell$  in  $B$  is precisely  $2^{-\ell}$  times the sum of the trace of the  $\ell$ th power of the matrix  $J - 2I$ , where  $J$  and  $I$  are the  $2 \times 2$  all-ones and identity matrices, respectively. Since the eigenvalues of  $J - 2I$  are 0 and  $-2$ , we see that  $t(C_\ell, B) = (-1)^\ell$ .

Now, let  $p = \min\{p_1, p_2\}$ , let  $\delta \in (0, 1)$  be a small real number, to be specified later, and define  $U := p \cdot B^\delta$ . By construction,  $U$  is 0-regular. Therefore, by Lemma 3.5, we have  $t(F, U) = 0$  for any graph  $F$  which has a vertex of degree one. In particular, since  $g(H_2) \geq g(H_1) = k$ , this means that  $t(F, U) = 0$  for any subgraph  $F$  of  $H_1$  or  $H_2$  with fewer than  $k$  edges. Also, using Lemma 3.2, the fact that  $k$  is odd and the observation from the end of the previous paragraph, we see that  $t(C_k, U) = -p^k \delta^k$ .

For  $i \in \{1, 2\}$ , let  $b_i$  be the number of subgraphs of  $H_i$  with at least  $k + 1$  non-isolated vertices. By Lemma 3.2, we have  $t(F, U) \leq \delta^{v(F)-i(F)}$  for any graph  $F$ . Putting all of this together, the expression in (2.6) can be bounded above as follows:

$$\begin{aligned} & \sum_{\substack{E \subseteq E(H_1) \\ |E| \geq 2}} \frac{t(H_1[E], U)}{p_1^{|E|-1} e(H_1)} + \sum_{\substack{E \subseteq E(H_2) \\ |E| \geq 2}} (-1)^{|E|} \frac{t(H_2[E], U)}{p_2^{|E|-1} e(H_2)} \\ & \leq \frac{-c_k(H_1)p^k \delta^k}{p_1^{k-1} e(H_1)} + \frac{c_k(H_2)p^k \delta^k}{p_2^{k-1} e(H_2)} + \sum_{i=1}^2 \frac{b_i \delta^{k+1}}{p^{e(H_i)-1} e(H_i)}. \end{aligned}$$

Now, by (3.7), if  $\delta$  is chosen small enough with respect to  $H_1, H_2, p_1$  and  $p_2$ , then the above expression is negative. This completes the proof.  $\square$

The previous lemma yields a strong constraint on  $(p_1, p_2)$  in the case that  $H_1$  and  $H_2$  both have odd girth.

**Corollary 3.8.** *If  $H_1$  and  $H_2$  both have odd girth and  $(H_1, H_2)$  is  $(p_1, p_2)$ -common, then  $p_1 = \frac{1}{\alpha+1}$  where*

$$\alpha = \left( \frac{c_k(H_2)e(H_1)}{c_k(H_1)e(H_2)} \right)^{1/(k-1)}$$

and  $k = g(H_1) = g(H_2)$ .

*Proof.* Let  $k = g(H_1)$ . By Theorem 3.6, we must also have  $g(H_2) = k$ . Also, we must have

$$\frac{c_k(H_1)}{p_1^{k-1} e(H_1)} = \frac{c_k(H_2)}{p_2^{k-1} e(H_2)}.$$

The result follows by substituting  $p_2 = 1 - p_1$  and solving for  $p_1$ .  $\square$

Our next goal is to modify a construction from [25] to show that, if  $(H_1, H_2)$  is  $(p_1, p_2)$ -common, then both of  $H_1$  and  $H_2$  must be  $K_4$ -free. This is an asymmetric generalization of the result of [25] which says that every common graph is  $K_4$ -free. For this, we will use the notion of a tensor products of kernels, which features prominently in constructions showing that certain graphs are not common [11, 16, 51].

**Remark 3.9.** Note that  $[0, 1]$  and  $[0, 1]^2$  are atomless standard probability spaces, and so there is a measure preserving map  $\psi : [0, 1] \rightarrow [0, 1]^2$ . This function is fixed throughout the rest of this section. For  $x \in [0, 1]$ , let  $\psi_1(x)$  and  $\psi_2(x)$  be such that  $\psi(x) = (\psi_1(x), \psi_2(x))$ .

**Definition 3.10.** Let  $U_1$  and  $U_2$  be kernels. The *tensor product* of  $U_1$  and  $U_2$  is the function  $U_1 \otimes U_2 : [0, 1]^2 \rightarrow \mathbb{R}$  defined by

$$U_1 \otimes U_2(x, y) = U_1(\psi_1(x), \psi_1(y)) \cdot U_2(\psi_2(x), \psi_2(y))$$

for all  $(x, y) \in [0, 1]^2$ .

The following (standard) lemma describes a key property of the tensor product.

**Lemma 3.11** (E.g., [35, Equation (7.17)]). *Let  $F$  be a graph and let  $U_1$  and  $U_2$  be kernels. Then*

$$t(F, U_1 \otimes U_2) = t(F, U_1) \cdot t(F, U_2).$$

*Proof.* By definition,

$$\begin{aligned} t(F, U_1 \otimes U_2) &= \int_{[0,1]^{V(F)}} \prod_{uv \in E(F)} U_1 \otimes U_2(x_u, x_v) dx_{V(F)} \\ &= \int_{[0,1]^{V(F)}} \prod_{uv \in E(F)} U_1(\psi_1(x_u), \psi_1(x_v)) U_2(\psi_2(x_u), \psi_2(x_v)) dx_{V(F)}. \end{aligned}$$

Since  $\psi$  is measure-preserving, we can view this expression equivalently as an integral over  $2v(F)$  variables ( $y_v : v \in V(F)$ ) and ( $z_v : v \in V(F)$ ) where  $y_v = \psi_1(x_v)$  and  $z_v = \psi_2(x_v)$ .

We get

$$\int_{[0,1]^{V(F)}} \int_{[0,1]^{V(F)}} \prod_{uv \in E(F)} U_1(y_u, y_v) U_2(z_u, z_v) dy_{V(F)} dz_{V(F)}.$$

This integral is clearly separable and is equal to  $t(F, U_1) \cdot t(F, U_2)$ .  $\square$

It is often useful to take “tensor powers” of kernels. Given a kernel  $U$ , let  $U^{\otimes 0} := 1$  and, for  $k \geq 1$ , let  $U^{\otimes k} := U^{\otimes(k-1)} \otimes U$ .

**Corollary 3.12.** *For any graph  $H$ , kernel  $U$  and  $k \geq 1$ ,*

$$t(H, U^{\otimes k}) = t(H, U)^k.$$

The following lemma can be seen as a weak form of Lemma 10 from [25]; we have distilled it down to contain only the ingredients that we will need here. We note that the proof of this lemma is fairly involved and relies on Fourier analytic ideas; for this reason, we will describe the construction itself and establish a few of the properties, but will not repeat the full analysis here. An alternative argument showing that  $K_4$  is uncommon without Fourier analysis was recently provided by Cs3ka, Hubai and Lov3asz [11].

**Lemma 3.13** (Jagger, Št3v3ček and Thomason [25]). *There exists a kernel  $K$  such that*

$$(3.14) \quad t(K_4, K) = -1/2,$$

$$(3.15) \quad |t(F, K)| \leq 1/2 \text{ for every graph with at least one edge, and}$$

$$(3.16) \quad |t(F, K)| \leq 1/4 \text{ if } F \text{ is a connected graph on three or four vertices which is not isomorphic to } K_4.$$

*Proof Sketch.* Let  $I_1, I_2, I_3, I_4$  be a partition of  $[0, 1]$  into four intervals of length  $1/4$ . Let  $K$  be the kernel defined by

$$K(x, y) := \begin{cases} 1 & \text{if } (x, y) \in I_i^2 \text{ for some } i \in \{1, 2, 3, 4\}, \\ -1 & \text{otherwise.} \end{cases}$$

In other words,  $K$  is the kernel such that  $(1 - K)/2$  is precisely the graphon  $W_{K_4}$ .

By definition,  $K$  is  $(-1/2)$ -regular. Thus, by Lemma 3.5, for any graph  $F$  that has a vertex  $w$  of degree one, it holds that  $t(F, K) = -(1/2)t(F - w, K)$ . In particular,  $t(F, K) = (-1/2)^{e(F)}$  for any tree  $F$ . Moreover, in proving any of the three assertions of the lemma, it suffices to consider graphs that do not have any vertices of degree one.

Thus, to prove (3.14) and (3.16), we need only consider  $F \in \{K_4, K_3, C_4, D\}$ , where  $D$  is the graph obtained from  $K_4$  by deleting an edge, which we refer to as the *diamond*. It is tedious, but not difficult, to confirm that the homomorphism densities of the relevant graphs in  $K$  are the values given by the following table:

$K_4$	$K_3$	$C_4$	$D$
$-1/2$	$1/4$	$1/4$	$1/4$

The proof of property (3.15) is somewhat complicated and is therefore omitted from this proof sketch; see the proof of [25, Lemma 10] for details.  $\square$

Using the previous lemma, we prove that  $K_4$  subgraphs are a strong obstruction to being common in the asymmetric setting, analogous to the symmetric setting.

**Theorem 3.17.** *Let  $H_1$  and  $H_2$  be graphs and let  $p_1$  and  $p_2$  be positive real numbers such that  $p_1 + p_2 = 1$ . If  $H_1$  contains  $K_4$ , then  $(H_1, H_2)$  is not  $(p_1, p_2)$ -common.*

*Proof.* Our goal is to apply Lemma 2.5. To that end, we construct a kernel  $U$  such that the expression in (2.6) is negative. Let  $p := \min\{p_1, p_2\}$ .

Let  $k$  be an integer and let  $\delta$  be a real number which will be chosen later in the proof. We let  $U = p \cdot (K^{\otimes(2k+1)})^\delta$ , where  $K$  is a kernel satisfying the conclusion of Lemma 3.13. Then, by Lemmas 3.2 and 3.11, we have

$$|t(F, U)| = \delta^{v(F) - i(F)} p^{e(F)} t(F, K)^{2k+1}$$

for any graph  $F$ . In particular,  $t(K_4, U) = \delta^4 p^{e(F)} (-1/2)^{2k+1}$ . Given a graph  $F$ , let  $F^*$  be the graph obtained from  $F$  by deleting all isolated vertices. By applying Lemma 3.13 and combining it with Lemma 2.2 to deal with disconnected graphs, we get that

$$|t(F, U)| \leq \begin{cases} \delta^5 (1/2)^{2k+1} & \text{if } v(F) - i(F) \geq 5, \\ (1/4)^{2k+1} & \text{if } 3 \leq v(F) - i(F) \leq 4 \text{ and } F^* \notin \{K_2, K_4\}. \end{cases}$$

Now, for  $i \in \{1, 2\}$ , let  $b_i$  be the number of non-empty subsets  $E$  of  $E(H_i)$  such that  $H_i[E]$  has at least five vertices. Let  $s_i$  be the number of non-empty subsets  $E$  of  $E(H_i)$  such

that  $H_i[E]^*$  has at most four vertices and is not isomorphic to  $K_2$  nor  $K_4$ . Also, let  $n_i$  be the number of non-empty subsets  $E$  of  $E(H_i)$  such that  $H_i[E]$  is isomorphic to  $K_4$ . The expression in (2.6) can be bounded as follows:

$$\begin{aligned} & \sum_{\substack{E \subseteq E(H_1) \\ |E| \geq 2}} \frac{t(H_1[E], U)}{p_1^{|E|-1} e(H_1)} + \sum_{\substack{E \subseteq E(H_2) \\ |E| \geq 2}} (-1)^{|E|} \frac{t(H_2[E], U)}{p_2^{|E|-1} e(H_2)} \\ & \leq \sum_{i=1}^2 \left( \frac{b_i \delta^5 (1/2)^{2k+1}}{p_i^{e(H_i)-1} e(H_i)} + \frac{s_i (1/4)^{2k+1}}{p_i^5 e(H_i)} + \frac{n_i \delta^4 (-1/2)^{2k+1} p^6}{p_i^6 e(H_i)} \right). \end{aligned}$$

To complete the proof, first choose  $\delta$  small enough with respect to  $H_1, H_2, p_1$  and  $p_2$  so that

$$\sum_{i=1}^2 \frac{b_i \delta^5}{p_i^{e(H_i)-1} e(H_i)} < \frac{1}{2} \sum_{i=1}^2 \frac{n_i \delta^4 p^6}{p_i^6 e(H_i)}.$$

Now, let  $k$  be chosen large enough with respect to  $H_1, H_2, p_1, p_2$  and  $\delta$  so that

$$\sum_{i=1}^2 \frac{s_i (1/4)^{2k+1}}{p_i^5 e(H_i)} < \frac{1}{2} \sum_{i=1}^2 \frac{n_i \delta^4 (1/2)^{2k+1} p^6}{p_i^6 e(H_i)}.$$

Putting this together, we get that the expression in (2.6) is negative, and so  $(H_1, H_2)$  is not  $(p_1, p_2)$ -common.  $\square$

**Remark 3.18.** In [11], it is shown that any graph  $H$  containing  $K_4$  fails to be common in a certain “local” sense. Specifically, this corresponds to showing, in the case  $p_1 = p_2 = 1/2$  and  $H_1 = H_2 = H$ , that, for every  $\varepsilon > 0$ , there exists a kernel  $U$  with  $\|U\|_\infty \leq \varepsilon$  such that the expression in (2.6) is negative. The construction in the previous lemma easily yields a similar statement in the asymmetric setting. The only change is to start by letting  $U = \min\{\varepsilon, p_1, p_2\} \cdot (K^{\otimes(2k+1)})^\delta$  and then choosing  $\delta$  and  $k$  to additionally depend on  $\varepsilon$  at the end of the proof.

Given a graph  $F$ , let  $\mathcal{P}(F, 0) = \{F\}$  and, for  $\ell \geq 1$ , let  $\mathcal{P}(F, \ell)$  be the set of graphs which can be obtained from a graph in  $\mathcal{P}(F, \ell - 1)$  by adding a pendant vertex. Note that each graph in  $\mathcal{P}(F, \ell)$  has exactly  $v(F) + \ell$  vertices and  $e(F) + \ell$  edges. Next, we show that, if  $H_1$  is not Sidorenko, then, for any graph  $F$ , there exists  $p \in (0, 1)$  and  $\ell \in \mathbb{N}$  such that  $(H_1, H_2)$  is not  $(p, 1 - p)$ -common for any  $H_2 \in \mathcal{P}(F, \ell)$ . The proof is inspired by an argument used by Jagger, Stovíček and Thomason to show that non-bipartite graph can be made into an uncommon graph by gluing a sufficiently large tree onto it [25, Theorem 4]. Our proof applies the following result of Coregliano and Razborov [10].

**Theorem 3.19** (Coregliano and Razborov [10, Theorem 8.2]). *If  $H$  is not Sidorenko, then there exists  $d \in (0, 1)$  and a  $d$ -regular graphon  $W$  such that*

$$t(H, W) < t(K_2, W)^{e(H)}.$$

**Theorem 3.20.** *If  $H_1$  is not Sidorenko, then, for any graph  $F$ , there exists  $p \in (0, 1)$  and  $\ell \in \mathbb{N}$  such that  $(H_1, H_2)$  is not  $(p, 1 - p)$ -common for any  $H_2 \in \mathcal{P}(F, \ell)$ .*

*Proof.* Using Theorem 3.19, let  $W$  be a  $d$ -regular graphon such that  $t(H_1, W) < t(K_2, W)^{e(H_1)} = d^{e(H_1)}$ . For each  $\ell \in \mathbb{N}$ , let

$$\alpha(\ell) := \frac{e(F) \ln\left(\frac{1}{1-d}\right)}{\ell}.$$

We choose  $\ell$  sufficiently large so that  $\alpha := \alpha(\ell) < 2^{-e(H_1)} (d^{e(H_1)} - t(H_1, W))$ . Let

$$W_1(x, y) := W(x, y) + \alpha(1 - W(x, y)) = (1 - \alpha)W(x, y) + \alpha$$

for all  $(x, y) \in [0, 1]^2$  and let  $W_2 = 1 - W_1$ . Then  $W_2$  is  $(1 - \alpha)(1 - d)$ -regular. So, by Lemma 3.5, for any  $H_2 \in \mathcal{P}(F, \ell)$ , we have

$$t(H_2, W_2) = t(F, W_2)(1 - \alpha)^\ell (1 - d)^\ell \leq (1 - \alpha)^\ell (1 - d)^\ell \leq e^{-\alpha\ell} (1 - d)^\ell = (1 - d)^{e(H_2)}.$$

Now, an algebraic expansion similar to the one at the beginning of Subsection 2.1 gives us

$$\begin{aligned} t(H_1, W_1) &= \int_{[0,1]^{V(H)}} \prod_{uv \in E(H)} W_1(x_u, x_v) dx_{V(H)} \\ &= \sum_{E_1 \sqcup E_2 = E(H_1)} \int_{[0,1]^{V(H_1)}} \prod_{uv \in E_1} W(x_u, x_v) \cdot \alpha^{|E_2|} \cdot \prod_{uv \in E_2} (1 - W(x_u, x_v)) dx_{V(H)} \\ &\leq t(H_1, W) + 2^{e(H_1)} \alpha. \end{aligned}$$

By our choice of  $\ell$ , the above inequality implies that  $t(H_1, W_1) < d^{e(H_1)}$ . Therefore,

$$\frac{t(H_1, W_1)}{e(H_1)d^{e(H_1)-1}} + \frac{t(H_2, W_2)}{e(H_2)(1-d)^{e(H_2)-1}} < \frac{d}{e(H_1)} + \frac{1-d}{e(H_2)}$$

and so  $(H_1, H_2)$  is not  $(d, 1 - d)$ -common. This completes the proof.  $\square$

## 4 Simple Consequences of Convexity

Up to this point, we have focused mainly on negative results; i.e. showing that certain pairs  $(H_1, H_2)$  of graphs fail to be  $(p_1, p_2)$ -common. We now turn our attention to positive results.

A great deal of attention in modern extremal graph theory has been on proving the validity of binomial inequalities (as studied in depth in, e.g., [1, 2, 30]) of the following form:

$$\prod_{i=1}^m t(F_i, W)^{\alpha_i} \geq t(F, W)$$

where  $F_1, F_2, \dots, F_m$  and  $F$  are graphs,  $\alpha_1, \dots, \alpha_k$  are non-zero reals and the above inequality holds for all graphons  $W$  or, occasionally, for all kernels [34] or all graphons satisfying

certain constraints (e.g. having non-negative spectrum [44, 47]). A prominent example is Sidorenko's Conjecture which, by Lemma 2.2, is equivalent to the statement that  $t(H, W) \geq t(e(H) \cdot K_2, W)$  for every bipartite graph  $H$  and graphon  $W$ .

Many of the results on Sidorenko's Conjecture over the years have focused on establishing binomial inequalities of the form

$$t(H, W)t(K_2, W)^\alpha \geq t(F, W)^\beta \quad (4.1)$$

where  $F$  is a graph that is already known to be Sidorenko, either inductively or by applying existing results, and  $\alpha$  and  $\beta$  are positive integers such that  $e(H) = \beta e(F) - \alpha$ . Given such an inequality, the fact that  $H$  is Sidorenko follows trivially from the fact that  $F$  is Sidorenko. Inequalities of the form given by (4.1) are also valuable in the study of common graphs. An early application can be found in [25, Theorem 2]. A more recent example is the result of Grzesik et al. that triangle-trees are common, which follows directly from the result of Goodman [20] and an inequality of this type [21, Corollary 3.4].

Unsurprisingly, such inequalities are also valuable to us in studying the asymmetric variant of the problem. The following lemma is useful for translating bounds of the form given by (4.1) into results on common pairs of graphs. It will be applied mainly in Section 6.

**Lemma 4.2.** *Let  $F_1, F_2, H_1$  and  $H_2$  be non-empty graphs and let  $k_1, k_2, \ell_1, \ell_2$  be non-negative reals such that the following hold:*

$$(4.3) \quad \min\{k_1, k_2\} \geq \max\{\lceil \ell_1 \rceil, \lceil \ell_2 \rceil\} + 1,$$

$$(4.4) \quad k_1 e(F_1) \ell_2 = k_2 e(F_2) \ell_1,$$

$$(4.5) \quad e(H_i) = k_i e(F_i) - \ell_i \text{ for } i \in \{1, 2\},$$

$$(4.6) \quad t(H_i, W)t(K_2, W)^{\ell_i} \geq t(F_i, W)^{k_i} \text{ for every graphon } W \text{ and } i \in \{1, 2\}.$$

*If  $p_1, p_2 > 0$  such that  $p_1 + p_2 = 1$  and  $(F_1, F_2)$  is  $(p_1, p_2)$ -common, then  $(H_1, H_2)$  is  $(p_1, p_2)$ -common.*

*Proof.* Let  $W_1$  and  $W_2$  be graphons such that  $W_1 + W_2 = 1$ . If either  $W_1$  or  $W_2$  is equal to 1 almost everywhere, then (2.1) holds trivially. So, we may assume that  $t(K_2, W_1), t(K_2, W_2) > 0$ . By (4.6), we have

$$\frac{t(H_1, W_1)}{e(H_1)p_1^{e(H_1)-1}} + \frac{t(H_2, W_2)}{e(H_2)p_2^{e(H_2)-1}} \geq \frac{t(F_1, W_1)^{k_1}}{e(H_1)p_1^{e(H_1)-1}t(K_2, W_1)^{\ell_1}} + \frac{t(F_2, W_2)^{k_2}}{e(H_2)p_1^{e(H_2)-1}t(K_2, W_2)^{\ell_2}}.$$

Our goal is to bound the expression on the right side of this inequality below by  $\frac{p_1}{e(H_1)} + \frac{p_2}{e(H_2)}$ .

Since  $(F_1, F_2)$  is  $(p_1, p_2)$ -common, there exists a real number  $x$  such that both of the following hold:

$$\frac{t(F_1, W_1)}{e(F_1)p_1^{e(F_1)-1}} \geq \frac{p_1}{e(H_1)} + x \geq 0, \quad \frac{t(F_2, W_2)}{e(F_2)p_2^{e(F_2)-1}} \geq \frac{p_2}{e(H_2)} - x \geq 0.$$

In other words,

$$t(F_1, W_1) \geq p_1^{e(F_1)} (1 + xe(F_1)p_1^{-1}) \geq 0, \quad t(F_2, W_2) \geq p_2^{e(F_2)} (1 - xe(F_2)p_2^{-1}) \geq 0. \quad (4.7)$$

Suppose first that  $\ell_1 = 0$  which, by (4.3) and (4.4), implies that  $\ell_2 = 0$ . In this case, we get

$$\begin{aligned} \frac{t(F_1, W_1)^{k_1}}{e(H_1)p_1^{e(H_1)-1}} + \frac{t(F_2, W_2)^{k_2}}{e(H_2)p_1^{e(H_2)-1}} &\geq \frac{p_1^{k_1 e(F_1)} (1 + xe(F_1)p_1^{-1})^{k_1}}{e(H_1)p_1^{e(H_1)-1}} + \frac{p_2^{k_2 e(F_2)} (1 - xe(F_2)p_2^{-1})^{k_2}}{e(H_2)p_1^{e(H_2)-1}} \\ &= \frac{p_1 (1 + xe(F_1)p_1^{-1})^{k_1}}{e(H_1)} + \frac{p_2 (1 - xe(F_2)p_2^{-1})^{k_2}}{e(H_2)} \end{aligned}$$

where the last equality uses (4.5) and the fact that  $\ell_1 = \ell_2 = 0$ . By Bernoulli's Inequality (Theorem A.1), we can bound this below by

$$\frac{p_1}{e(H_1)} + \frac{p_2}{e(H_2)} + x \left( \frac{p_1}{e(H_1)} \left( \frac{e(F_1)k_1}{p_1} \right) - \frac{p_2}{e(H_2)} \left( \frac{e(F_2)k_2}{p_2} \right) \right) = \frac{p_1}{e(H_1)} + \frac{p_2}{e(H_2)}$$

where the last equality uses (4.5) and the fact that  $\ell_1 = \ell_2 = 0$ . This completes the proof in the case that  $\ell_1 = 0$ .

Now, suppose that  $\ell_1 \geq 1$  which, by (4.3) and (4.4), implies that  $\ell_2 \geq 1$ . Let  $-p_1 \leq y \leq p_2$  such that  $t(K_2, W_1) = p_1 + y$  and  $t(K_2, W_2) = p_2 - y$ . For  $i \in \{1, 2\}$ , define

$$b_i := \frac{t(F_i, W_i)^{k_i}}{e(H_i)p_i^{e(H_i)+\ell_i-1}} = \frac{t(F_i, W_i)^{k_i}}{e(H_i)p_i^{ke(F_i)-1}}$$

where the equality holds by (4.5). Then

$$\frac{t(F_1, W_1)^{k_1}}{e(H_1)p_1^{e(H_1)-1}t(K_2, W_1)^{\ell_1}} + \frac{t(F_2, W_2)^{k_2}}{e(H_2)p_1^{e(H_2)-1}t(K_2, W_2)^{\ell_2}} = \frac{b_1}{(1 + p_1^{-1}y)^{\ell_1}} + \frac{b_2}{(1 - p_2^{-1}y)^{\ell_2}}. \quad (4.8)$$

By (4.3), we can choose  $d$  to be an integer such that  $\min\{k_1, k_2\} \geq d \geq \max\{\ell_1, \ell_2\} + 1$ . By Lemma A.3, we get that the expression on the right side of equation (4.8) is at least

$$\frac{\left( p_1 \ell_1^{-1} (b_1 p_1^{-1} \ell_1)^{1/d} + p_2 \ell_2^{-1} (b_2 p_2^{-1} \ell_2)^{1/d} \right)^d}{(p_1 \ell_1^{-1} + p_2 \ell_2^{-1})^{d-1}}. \quad (4.9)$$

So, we will be done if we can bound the expression in (4.9) below by  $\frac{p_1}{e(H_1)} + \frac{p_2}{e(H_2)}$ . By (4.7), we have

$$b_1 \geq \frac{p_1 (1 + xe(F_1)p_1^{-1})^{k_1}}{e(H_1)}, \quad b_2 \geq \frac{p_2 (1 - xe(F_2)p_2^{-1})^{k_2}}{e(H_2)}$$

and so

$$p_1 \ell_1^{-1} (b_1 p_1^{-1} \ell_1)^{1/d} + p_2 \ell_2^{-1} (b_2 p_2^{-1} \ell_2)^{1/d}$$



$$\geq \frac{p_1}{\ell_1} \left( \frac{\ell_1}{e(H_1)} \right)^{1/d} \left( 1 + \frac{xe(F_1)}{p_1} \right)^{k_1/d} + \frac{p_2}{\ell_2} \left( \frac{\ell_2}{e(H_2)} \right)^{1/d} \left( 1 - \frac{xe(F_2)}{p_2} \right)^{k_2/d}.$$

By Theorem A.1 and the fact that  $\min\{k_1, k_2\} \geq d$ , this is at least

$$\begin{aligned} & \frac{p_1}{\ell_1} \left( \frac{\ell_1}{e(H_1)} \right)^{1/d} + \frac{p_2}{\ell_2} \left( \frac{\ell_2}{e(H_2)} \right)^{1/d} \\ & + x \left( \frac{p_1}{\ell_1} \left( \frac{\ell_1}{e(H_1)} \right)^{1/d} \left( \frac{e(F_1)k_1}{p_1 d} \right) - \frac{p_2}{\ell_2} \left( \frac{\ell_2}{e(H_2)} \right)^{1/d} \left( \frac{e(F_2)k_2}{p_2 d} \right) \right). \end{aligned}$$

By (4.4) and (4.5), we have that  $e(F_1)k_1/\ell_1 = e(F_2)k_2/\ell_2$  and  $e(H_1)/\ell_1 = e(H_2)/\ell_2$ . Therefore, all of the terms in the above expression involving  $x$  cancel out. Plugging the remaining terms into (4.9) and using the fact that  $e(F_1)k_1/\ell_1 = e(F_2)k_2/\ell_2$  twice, we get the following

$$\frac{\left( \frac{p_1}{\ell_1} \left( \frac{\ell_1}{e(H_1)} \right)^{1/d} + \frac{p_2}{\ell_2} \left( \frac{\ell_2}{e(H_2)} \right)^{1/d} \right)^d}{(p_1 \ell_1^{-1} + p_2 \ell_2^{-1})^{d-1}} = \frac{\left( \frac{\ell_1}{e(H_1)} \right) \left( \frac{p_1}{\ell_1} + \frac{p_2}{\ell_2} \right)^d}{(p_1 \ell_1^{-1} + p_2 \ell_2^{-1})^{d-1}} = \frac{p_1}{e(H_1)} + \frac{p_2}{e(H_2)}$$

which completes the proof.  $\square$

Next, we derive a couple of easy consequences of Lemma 4.2.

**Corollary 4.10.** *If  $H_1$  and  $H_2$  are Sidorenko, then  $(H_1, H_2)$  is  $(p_1, p_2)$ -common for any positive reals  $p_1$  and  $p_2$  such that  $p_1 + p_2 = 1$ .*

*Proof.* Apply Lemma 4.2 with  $F_1 = F_2 = K_2$ ,  $k_i = e(H_i)$  for  $i \in \{1, 2\}$  and  $\ell_1 = \ell_2 = 0$ . Hypothesis (4.6) of Lemma 4.2 is satisfied because  $H_1$  and  $H_2$  are Sidorenko.  $\square$

**Corollary 4.11.** *If  $(F_1, F_2)$  is  $(p_1, p_2)$ -common, then  $(k_1 \cdot F_1, k_2 \cdot F_2)$  is  $(p_1, p_2)$ -common for any positive integers  $k_1$  and  $k_2$ .*

*Proof.* Lemma 4.2 with  $\ell_1 = \ell_2 = 0$  and  $H_i = k_i \cdot F_i$  for  $i \in \{1, 2\}$ . Hypothesis (4.6) of Lemma 4.2 is satisfied because of Lemma 2.2.  $\square$

The following proposition generalizes Corollary 4.11.

**Proposition 4.12.** *Let  $F_1, F_2$  and  $F_3$  be non-empty graphs. If  $e(F_1) = e(F_2)$  and  $(F_1, F_3)$  and  $(F_2, F_3)$  are  $(p_1, p_2)$ -common, then  $(F_1 \sqcup F_2, F_3)$  is  $(p_1, p_2)$ -common.*

*Proof.* Let  $W_1$  and  $W_2$  be graphons such that  $W_1 + W_2 = 1$ . Let  $k = e(F_1) = e(F_2)$  and let

$$x := -\frac{p_1}{k} + \min \left\{ \frac{t(F_1, W_1)}{kp_2^{k-1}}, \frac{t(F_2, W_1)}{kp_2^{k-1}} \right\}$$

Then

$$\frac{t(F_i, W_1)}{kp_2^{k-1}} \geq \frac{p_1}{k} + x \geq 0 \text{ for } i \in \{1, 2\}.$$

Also, since  $(F_1, F_3)$  and  $(F_2, F_3)$  are both  $(p_1, p_2)$ -common, we have

$$\frac{t(F_3, W_2)}{e(F_3)p_2^{e(F_3)-1}} \geq \frac{p_2}{e(F_3)} - x.$$

Unfortunately, the right side of the above inequality may be negative, which forbids us from applying Bernoulli's Inequality. To remedy this, let  $y = \min\{x, p_2/e(F_3)\}$ . Note that, if  $x < 0$ , then we get  $y = x$  and, otherwise,  $y \leq x$ ; thus, the following still holds

$$\frac{t(F_i, W_1)}{kp_2^{k-1}} \geq \frac{p_1}{k} + y \geq 0 \text{ for } i \in \{1, 2\}.$$

Also,

$$\frac{t(F_3, W_2)}{e(F_3)p_2^{e(F_3)-1}} \geq \frac{p_2}{e(F_3)} - y \geq 0$$

now holds. Rearranging these last two inequalities yields

$$t(F_3, W_2) = p_2^{e(F_3)} (1 - ye(F_3)p_2^{-1}) \geq 0, \quad t(F_i, W_1) \geq p_2^k (1 + ykp_1^{-1}) \geq 0 \text{ for } i \in \{1, 2\}.$$

Therefore,

$$\begin{aligned} \frac{t(F_1 \sqcup F_2, W_1)}{2kp_1^{2k-1}} + \frac{t(F_3, W_2)}{e(F_3)p_2^{e(F_3)-1}} &\geq \frac{p_1(1 + ykp_1^{-1})(1 + ykp_1^{-1})}{2k} + \frac{p_2(1 - ye(F_3)p_2^{-1})}{e(F_3)} \\ &= \frac{p_1}{2k} + y + \frac{y^2k^2}{p_1(2k)^2} + \frac{p_2}{e(F_3)} - y \geq \frac{p_1}{2k} + \frac{p_2}{e(F_3)} \end{aligned}$$

as desired. □

## 5 Strongly Common Graphs

The results presented in the previous section are useful for transforming a  $(p_1, p_2)$ -common pair  $(F_1, F_2)$  of graphs into another  $(p_1, p_2)$ -common pair  $(H_1, H_2)$ , but they do not allow one to change the values of  $p_1$  and  $p_2$ . Thus, feeding the known examples of common graphs and binomial inequalities into these results only yields examples of  $(1/2, 1/2)$ -common pairs of graphs. Our two goals in this subsection are to show that stronger forms of the common graph property can be used to obtain examples of  $(p_1, p_2)$ -common pairs where  $(p_1, p_2) \neq (1/2, 1/2)$ , and to exhibit two examples, one old and one new, of graphs which possess this stronger property.

**Definition 5.1.** A graph  $H$  is said to be *strongly common* if, for any graphons  $W_1$  and  $W_2$  such that  $W_1 + W_2 = 1$ ,

$$t(H, W_1) + t(H, W_2) \geq t(K_2, W_1)^{e(H)} + t(K_2, W_2)^{e(H)}.$$

Since the function  $x^k + (1 - x)^k$  for  $k \geq 1$  is globally minimized at  $x = 1/2$ , we see that every strongly common graph is common. It is not clear whether or not the converse is true; see Problem 9.6. The classical example of a strongly common graph is the triangle; this follows from the following result, known as Goodman's Formula [20]. We include a proof for completeness, and because we will generalize it to  $C_5$  using a similar argument. For  $k \geq 1$ , let  $P_k$  denote the path on  $k$  vertices.

**Theorem 5.2** (Goodman's Formula [20]). *For any graphons  $W_1$  and  $W_2$  such that  $W_1 + W_2 = 1$ , we have that  $t(K_3, W_1) + t(K_3, W_2)$  is equal to*

$$t(K_2, W_1)^3 + t(K_2, W_2)^3 + \frac{3}{2}(t(P_3, W_1) + t(P_3, W_2) - t(K_2, W_1)^2 - t(K_2, W_2)^2).$$

*Proof.* Let  $U$  be the kernel such that  $W_1 = t(K_2, W_1) + U$  and  $W_2 = t(K_2, W_2) - U$ . Note that, by definition of  $t(K_2, W_1)$ , we must have  $t(K_2, U) = 0$ . Using Observation 2.3, we get that  $t(K_3, W_1) + t(K_3, W_2)$  is equal to

$$\sum_{E \subseteq E(K_3)} t(K_2, W_1)^{3-|E|} t(K_3[E], U) + \sum_{E \subseteq E(K_3)} t(K_2, W_2)^{3-|E|} t(K_3[E], -U).$$

Using the facts that  $t(K_2, U) = 0$  and  $t(F, -U) = (-1)^{e(F)} t(F, U)$  for any graph  $F$  and kernel  $U$ , we get that the above expression is equal to

$$\begin{aligned} & t(K_2, W_1)^3 + t(K_2, W_2)^3 + 3(t(K_2, W_1) + t(K_2, W_2))t(P_3, U) \\ &= t(K_2, W_1)^3 + t(K_2, W_2)^3 + 3t(P_3, U). \end{aligned}$$

Now, applying Observation 2.3 to  $t(P_3, W_1) + t(P_3, W_2)$ , and using the facts that  $t(K_2, U) = 0$  and  $t(P_3, -U) = t(P_3, U)$  again, we get that

$$t(P_3, W_1) + t(P_3, W_2) = t(K_2, W_1)^2 + t(K_2, W_2)^2 + 2t(P_3, U).$$

Solving for  $t(P_3, U)$  in this expression and substituting it into the expression for  $t(K_3, W_1) + t(K_3, W_2)$  derived above completes the proof.  $\square$

**Theorem 5.3** (Goodman [20]).  *$K_3$  is strongly common.*

*Proof.* This follows from Theorem 5.2 and the fact that  $P_3$  is Sidorenko.  $\square$

Our next result is an analogue of the Goodman Formula for  $C_5$ , which we will use to show that  $C_5$  is strongly common. While this is a very natural extension of Theorem 5.2, we have not seen it anywhere in the literature. It seems likely that this phenomenon extends to longer odd cycles, but we have been unable to prove this; see Conjecture 9.7.

**Theorem 5.4.** *For any graphons  $W_1$  and  $W_2$  such that  $W_1 + W_2 = 1$ , we have  $t(C_5, W_1) + t(C_5, W_2)$  is equal to*

$$\begin{aligned} & t(K_2, W_1)^5 + t(K_2, W_2)^5 + 5t(K_2, W_1)t(P_5, W_1) + 5t(K_2, W_2)t(P_5, W_2) \\ & - 5t(K_2, W_1)^2 t(P_4, W_1) - 5t(K_2, W_1)^2 t(P_4, W_1). \end{aligned}$$

*Proof.* For ease of notation, let  $t(K_2, W_1) = p_1$  and  $t(K_2, W_2) = p_2 = 1 - p_1$ . Let  $U$  be the kernel such that  $W_1 = p_1 + U$  and  $W_2 = p_2 - U$ . Note that, by definition of  $t(K_2, W_1)$ , we must have  $t(K_2, U) = 0$ . Using Observation 2.3, we get that  $t(C_5, W_1) + t(C_5, W_2)$  is equal to

$$\sum_{E \subseteq E(C_5)} p_1^{5-|E|} t(C_5[E], U) + \sum_{E \subseteq E(C_5)} p_2^{5-|E|} t(C_5[E], -U).$$

Using the facts that  $t(K_2, U) = 0$  and  $t(F, -U) = (-1)^{e(F)} t(F, U)$  for any graph  $F$  and kernel  $U$ , we get that the above expression is equal to

$$p_1^5 + p_2^5 + 5(p_1^3 + p_2^3)t(P_3, U) + 5(p_1^2 - p_2^2)t(P_4, U) + 5(p_1 + p_2)t(P_5, U).$$

In the same way, we can apply Observation 2.3 to deduce that  $t(K_2, W_1)t(P_5, W_1) + t(K_2, W_2)t(P_5, W_2)$  is equal to

$$\begin{aligned} & p_1(p_1^4 + 3p_1^2 t(P_3, U) + 2p_1 t(P_4, U) + t(P_5, U)) + p_2(p_2^4 + 3p_2^2 t(P_3, U) - 2p_2 t(P_4, U) + t(P_5, U)) \\ &= p_1^5 + p_2^5 + 3(p_1^3 + p_2^3)t(P_3, U) + 2(p_1^2 - p_2^2)t(P_4, U) + (p_1 + p_2)t(P_5, U), \end{aligned}$$

and that  $t(K_2, W_1)^2 t(P_4, W_1) - t(K_2, W_2)^2 t(P_4, W_2)$  is equal to

$$\begin{aligned} & p_1^2(p_1^3 + 2p_1 t(P_3, U) + t(P_4, U)) + p_2^2(p_2^3 + 2p_2 t(P_3, U) - t(P_4, U)) \\ &= p_1^5 + p_2^5 + 2(p_1^3 + p_2^3)t(P_3, U) + (p_1^2 - p_2^2)t(P_4, U) \end{aligned}$$

and the claimed equality follows.  $\square$

In order to use Theorem 5.4 to show that  $C_5$  is strongly common, we apply a correlation inequality for homomorphism densities of paths which follows from a recent result of Blekherman and Raymond [1].

**Lemma 5.5** (Blekherman and Raymond [1, Theorem 1.3 (1.1)]). *If  $0 \leq a \leq b \leq c$  are integers such that  $a$  and  $c$  are odd, then*

$$\text{hom}(P_a, G)^{c-b} \text{hom}(P_c, G)^{b-a} \geq \text{hom}(P_b, G)^{c-a}$$

for every graph  $G$ .

**Corollary 5.6.** *For any graphon  $W$ ,*

$$t(P_5, W) \geq t(K_2, W)t(P_4, W).$$

*Proof.* Note that  $K_2 = P_2$ . By Lemma 5.5, for any graph  $G$  we have

$$\begin{aligned} \text{hom}(P_1, G)^3 \text{hom}(P_5, G) &\geq \text{hom}(P_2, G)^4 \quad \text{and} \\ \text{hom}(P_1, G) \text{hom}(P_5, G)^3 &\geq \text{hom}(P_4, G)^4. \end{aligned}$$

Multiplying these and taking the fourth root we obtain

$$\text{hom}(P_1, G) \text{hom}(P_5, G) = v(G) \text{hom}(P_5, G) \geq \text{hom}(P_2, G) \text{hom}(P_4, G).$$

Thus

$$t(P_5, G) = \frac{\text{hom}(P_5, G)}{v(G)^5} \geq \frac{\text{hom}(P_2, G) \text{hom}(P_4, G)}{v(G)^6} = t(P_2, G)t(P_4, G).$$

Taking limits yields the same result for graphons.  $\square$

**Theorem 5.7.**  $C_5$  is strongly common.

*Proof.* This follows from Theorem 5.4 and Corollary 5.6.  $\square$

We close this section with a lemma which shows that, under certain conditions, strongly common graphs can give rise to examples of  $(p_1, p_2)$ -common pairs of graphs with  $(p_1, p_2) \neq (1/2, 1/2)$ . This will be applied in Section 6.

**Lemma 5.8.** Let  $F, H_1$  and  $H_2$  be non-empty graphs,  $k_1, k_2, \ell_1, \ell_2$  be non-negative integers and  $p_1, p_2 \in (0, 1)$  such that  $p_1 + p_2 = 1$  and the following hold:

$$(5.9) \quad k_i \geq \ell_i + 1 \text{ for } i \in \{1, 2\},$$

$$(5.10) \quad e(H_i) = k_i e(F) - \ell_i \text{ for } i \in \{1, 2\},$$

$$(5.11) \quad p_2^{e(F)-1} k_1 e(H_2) = p_1^{e(F)-1} k_2 e(H_1),$$

$$(5.12) \quad t(H_i, W)t(K_2, W)^{\ell_i} \geq t(F, W)^{k_i} \text{ for every graphon } W \text{ and } i \in \{1, 2\},$$

$$(5.13) \quad \text{either } k_1 = 1 \text{ or } k_1 = k_2.$$

If  $F$  is strongly common, then  $(H_1, H_2)$  is  $(p_1, p_2)$ -common.

Before presenting the somewhat technical proof of this lemma, let us illustrate its applicability with a concrete example. Recall that  $D$  is the *diamond* graph obtained from  $K_4$  by deleting one edge. The class of *triangle-vertex-trees* is defined recursively as follows:  $K_3$  is the smallest triangle-vertex-tree and any graph obtained from a triangle-vertex-tree by adding a pendant copy of  $K_3$  is also a triangle-vertex-tree.

**Proposition 5.14.** If  $T$  is a triangle-vertex-tree and  $p = \sqrt{30} - 5$ , then  $(T, D)$  is  $(p, 1 - p)$ -common.

*Proof.* First, consider the pair  $(K_3, D)$ . We apply Lemma 5.8 with  $F = K_3$ ,  $(H_1, H_2) = (K_3, D)$ ,  $(k_1, k_2, \ell_1, \ell_2) = (1, 2, 0, 1)$  and  $(p_1, p_2) = (\sqrt{30} - 5, 6 - \sqrt{30})$ . Then, clearly, (5.9), (5.10), (5.11) and (5.13) hold by definition. Also, (5.12) holds by [21, Corollary 3.4].

Now, to get the result for the pair  $(T, D)$ , we apply Lemma 4.2 with  $(F_1, F_2) = (K_3, D)$ ,  $(H_1, H_2) = (T, D)$ ,  $(k_1, k_2, \ell_1, \ell_2) = ((|V(T)| - 1)/2, 2, 0, 1)$  and  $(p_1, p_2) = (\sqrt{30} - 5, 6 - \sqrt{30})$ . Clearly, (4.3), (4.4) and (4.5) hold by definition. Also, (4.6) holds by [21, Corollary 3.4]. Thus, the proof is complete.  $\square$

*Proof of Lemma 5.8.* Let  $W_1$  and  $W_2$  be graphons such that  $W_1 + W_2 = 1$ . Define  $x$  so that  $t(K_2, W_1) = p_1 + x$  and  $t(K_2, W_2) = p_2 - x$ . Since  $F$  is strongly common, we know that

$$t(F, W_1) + t(F, W_2) \geq (p_1 + x)^{e(F)} + (p_2 - x)^{e(F)}.$$

Let  $y$  be such that

$$t(F, W_1) \geq (p_1 + x)^{e(F)} + y \geq 0, \quad t(F, W_2) \geq (p_2 - x)^{e(F)} - y \geq 0; \quad (5.15)$$

Then, by (5.12),

$$\frac{t(H_1, W_1)}{e(H_1)p_1^{e(H_1)-1}} + \frac{t(H_2, W_2)}{e(H_2)p_2^{e(H_2)-1}} \geq \frac{t(F, W_1)^{k_1}}{t(K_2, W_1)^{\ell_1}e(H_1)p_1^{e(H_1)-1}} + \frac{t(F, W_2)^{k_2}}{t(K_2, W_1)^{\ell_2}e(H_2)p_2^{e(H_2)-1}}$$

which is at least

$$\frac{((p_1 + x)^{e(F)} + y)^{k_1}}{(p_1 + x)^{\ell_1}e(H_1)p_1^{e(H_1)-1}} + \frac{((p_2 - x)^{e(F)} - y)^{k_2}}{(p_2 - x)^{\ell_2}e(H_2)p_2^{e(H_2)-1}}. \quad (5.16)$$

Let  $f(x, y)$  be the expression in (5.16), viewed as a function of  $x$  and  $y$  with domain given by  $-p_1 \leq x \leq p_2$  and  $-(p_1 + x)^{e(F)} \leq y \leq (p_2 - x)^{e(F)}$ . Our goal is to show that  $f(x, y)$  is at least  $\frac{p_1}{e(H_1)} + \frac{p_2}{e(H_2)}$  for all  $x$  and  $y$  in its domain. We now divide the proof into cases depending on the value of  $k_1$ .

*Case 1.*  $k_1 = 1$ .

In this case, we get  $\ell_1 = 0$  by (5.9). Thus,  $f(x, y)$  simplifies to

$$f(x, y) = \frac{(p_1 + x)^{e(F)} + y}{e(H_1)p_1^{e(H_1)-1}} + \frac{((p_2 - x)^{e(F)} - y)^{k_2}}{(p_2 - x)^{\ell_2}e(H_2)p_2^{e(H_2)-1}}.$$

If  $k_2 = 1$  as well, then we must have  $\ell_2 = 0$  and the whole thing simplifies to  $\frac{p_1}{e(H_1)} + \frac{p_2}{e(H_2)}$ , as desired. So, for the rest of the proof of this case, we assume that  $k_2 \geq 2$ .

The partial derivative of  $f$  with respect to  $y$  is

$$\frac{\partial f}{\partial y}(x, y) = \frac{1}{e(H_1)p_1^{e(H_1)-1}} - \frac{k_2((p_2 - x)^{e(F)} - y)^{k_2-1}}{(p_2 - x)^{\ell_2}e(H_2)p_2^{e(H_2)-1}}.$$

Thus,  $\frac{\partial f}{\partial y}(x, y)$  is negative at  $y = -(p_1 + x)^{e(F)}$  and positive at  $y = (p_2 - x)^{e(F)}$ . This implies that the global minimum of  $f$  on its domain is at a point  $(x, y)$  such that  $-(p_1 + x)^{e(F)} < y < (p_2 - x)^{e(F)}$  and  $\frac{\partial f}{\partial y}(x, y) = 0$ . Setting this partial derivative equal to zero and solving for  $y$  yields

$$y = (p_2 - x)^{e(F)} - \left( \frac{(p_2 - x)^{\ell_2}e(H_2)p_2^{e(H_2)-1}}{k_2e(H_1)p_1^{e(H_1)-1}} \right)^{\frac{1}{k_2-1}}.$$

By (5.11), (5.10) and the fact that  $k_1 = 1$  and  $e(H_1) = e(F)$ , this simplifies to

$$y = (p_2 - x)^{e(F)} - \left( \frac{p_2 - x}{p_2} \right)^{\frac{\ell_2}{k_2-1}} p_2^{e(F)}.$$

Substituting this back into  $f$  yields

$$\frac{(p_1 + x)^{e(F)}}{e(H_1)p_1^{e(H_1)-1}} + \frac{(p_2 - x)^{e(F)}}{e(H_1)p_1^{e(H_1)-1}} - \frac{\left( \frac{p_2 - x}{p_2} \right)^{\frac{\ell_2}{k_2-1}} p_2^{e(F)}}{e(H_1)p_1^{e(H_1)-1}} + \frac{\left( \frac{p_2 - x}{p_2} \right)^{\frac{\ell_2 k_2}{k_2-1}} p_2^{e(F)k_2}}{(p_2 - x)^{\ell_2} e(H_2) p_2^{e(H_2)-1}}$$

which, by (5.11) and the fact that  $e(H_1) = e(F)$ , simplifies to the following:

$$\left( \frac{p_1 + x}{p_1} \right)^{e(F)} \left( \frac{p_1}{e(H_1)} \right) + k_2 \left( \frac{p_2 - x}{p_2} \right)^{e(F)} \left( \frac{p_2}{e(H_2)} \right) - (k_2 - 1) \left( \frac{p_2 - x}{p_2} \right)^{\frac{\ell_2}{k_2-1}} \left( \frac{p_2}{e(H_2)} \right).$$

By three applications of Theorem A.1, this is at least

$$\begin{aligned} & \left( 1 + \frac{xe(F)}{p_1} \right) \left( \frac{p_1}{e(H_1)} \right) + k_2 \left( 1 - \frac{xe(F)}{p_2} \right) \left( \frac{p_2}{e(H_2)} \right) - (k_2 - 1) \left( 1 - \frac{x\ell_2}{p_2(k_2 - 1)} \right) \left( \frac{p_2}{e(H_2)} \right) \\ &= \frac{p_1}{e(H_1)} + \frac{p_2}{e(H_2)} + \frac{xe(F)}{e(H_1)} - \frac{xe(F)k_2}{e(H_2)} + \frac{x\ell_2}{e(H_2)} \\ &= \frac{p_1}{e(H_1)} + \frac{p_2}{e(H_2)} + \frac{xe(F)e(H_2)}{e(H_1)e(H_2)} - \frac{xe(F)k_2e(H_1)}{e(H_1)e(H_2)} + \frac{x\ell_2e(H_1)}{e(H_1)e(H_2)}. \end{aligned}$$

Using (5.10) and the fact that  $e(H_1) = e(F)$ , this expression simplifies to  $\frac{p_1}{e(H_1)} + \frac{p_2}{e(H_2)}$  and we are done in this case.

*Case 2.*  $k_1 = k_2$ .

In this case,  $f(x, y)$  simplifies to

$$f(x, y) = \frac{((p_1 + x)^{e(F)} + y)^{k_1}}{(p_1 + x)^{\ell_1} e(H_1) p_1^{e(H_1)-1}} + \frac{((p_2 - x)^{e(F)} - y)^{k_1}}{(p_2 - x)^{\ell_2} e(H_2) p_2^{e(H_2)-1}}.$$

We apply the Generalized Hölder Inequality (Theorem A.2) with  $n = 2$  and  $m = k_1$ . Define

$$a_{1,1}(x) := \frac{((p_1 + x)^{e(F)} + y)^{k_1}}{(p_1 + x)^{\ell_1} e(H_1) p_1^{e(H_1)-1}}, \quad a_{2,1}(x) := \frac{((p_2 - x)^{e(F)} - y)^{k_1}}{(p_2 - x)^{\ell_2} e(H_2) p_2^{e(H_2)-1}},$$

$$a_{1,j}(x) := a_1(x) := \left( \frac{p_1 + x}{p_1} \right)^{\frac{\ell_1}{k_1-1}} \left( \frac{p_1}{e(H_1)} \right) \text{ and}$$

$$a_{2,j}(x) := a_2(x) := \left( \frac{p_2 - x}{p_2} \right)^{\frac{\ell_2}{k_1-1}} \left( \frac{p_2}{e(H_2)} \right)$$

for  $2 \leq j \leq k_1$ . Note that  $f(x, y) = a_{1,1} + a_{2,1}$ . Also, let  $\alpha_1 = \alpha_2 = \dots = \alpha_{k_1} = \frac{1}{k_1}$ . Then, by Theorem A.2,

$$\prod_{j=1}^{k_1} (a_{1,j} + a_{2,j})^{1/k_1} \geq \prod_{j=1}^m a_{1,j}^{1/k_1} + \prod_{j=1}^m a_{2,j}^{1/k_1}$$

which implies that  $f(x, y)$  is at least

$$\begin{aligned} & \frac{\left( \left( \frac{((p_1+x)^{e(F)}+y)^{k_1}}{(p_1+x)^{\ell_1} e(H_1) p_1^{e(H_1)-1}} \right)^{\frac{1}{k_1}} a_1(x)^{\frac{k_1-1}{k_1}} + \left( \frac{((p_2-x)^{e(F)}-y)^{k_1}}{(p_2-x)^{\ell_2} e(H_2) p_2^{e(H_2)-1}} \right)^{\frac{1}{k_1}} a_2(x)^{\frac{k_1-1}{k_1}} \right)^{k_1}}{(a_1(x) + a_2(x))^{k_1-1}} \\ &= \frac{\left( \frac{((p_1+x)^{e(F)}+y) a_1(x)^{\frac{k_1-1}{k_1}}}{(p_1+x)^{\frac{\ell_1}{k_1}} e(H_1)^{\frac{1}{k_1}} p_1^{\frac{e(H_1)-1}{k_1}}} + \frac{((p_2-x)^{e(F)}-y) a_2(x)^{\frac{k_1-1}{k_1}}}{(p_2-x)^{\frac{\ell_2}{k_1}} e(H_2)^{\frac{1}{k_1}} p_2^{\frac{e(H_2)-1}{k_1}}} \right)^{k_1}}{(a_1(x) + a_2(x))^{k_1-1}}. \end{aligned} \quad (5.17)$$

Using the fact that  $k_2 = k_1$  (several times), we see that the expression in (5.17) is equal to  $(a_1(x) + a_2(x))^{-k_1+1}$  multiplied by

$$\begin{aligned} & \left( \frac{((p_1+x)^{e(F)}+y) a_1(x)^{\frac{k_1-1}{k_1}}}{(p_1+x)^{\frac{\ell_1}{k_1}} e(H_1)^{\frac{1}{k_1}} p_1^{\frac{e(H_1)-1}{k_1}}} + \frac{((p_2-x)^{e(F)}-y) a_2(x)^{\frac{k_1-1}{k_1}}}{(p_2-x)^{\frac{\ell_2}{k_1}} e(H_2)^{\frac{1}{k_1}} p_2^{\frac{e(H_2)-1}{k_1}}} \right)^{k_1} \\ &= \left( \frac{((p_1+x)^{e(F)}+y) \left( \frac{p_1+x}{p_1} \right)^{\frac{\ell_1}{k_1}} \left( \frac{p_1}{e(H_1)} \right)^{\frac{k_1-1}{k_1}}}{(p_1+x)^{\frac{\ell_1}{k_1}} e(H_1)^{\frac{1}{k_1}} p_1^{\frac{e(H_1)-1}{k_1}}} + \frac{((p_2-x)^{e(F)}-y) \left( \frac{p_2-x}{p_2} \right)^{\frac{\ell_2}{k_1}} \left( \frac{p_2}{e(H_2)} \right)^{\frac{k_1-1}{k_1}}}{(p_2-x)^{\frac{\ell_2}{k_1}} e(H_2)^{\frac{1}{k_1}} p_2^{\frac{e(H_2)-1}{k_1}}} \right)^{k_1} \\ &= \left( \frac{(p_1+x)^{e(F)}+y}{e(H_1) p_1^{e(F)-1}} + \frac{(p_2-x)^{e(F)}-y}{e(H_2) p_2^{e(F)-1}} \right)^{k_1}. \end{aligned}$$

By (5.11) and the fact that  $k_2 = k_1$ , we have  $e(H_1) p_1^{e(F)-1} = e(H_2) p_2^{e(F)-1}$ . Thus, the above expression is equal to

$$\left( \frac{(p_1+x)^{e(F)}}{e(H_1) p_1^{e(F)-1}} + \frac{(p_2-x)^{e(F)}}{e(H_2) p_2^{e(F)-1}} \right)^{k_1}.$$

Thus, the expression in (5.17) is at least

$$\frac{\left( \frac{(p_1+x)^{e(F)}}{e(H_1) p_1^{e(F)-1}} + \frac{(p_2-x)^{e(F)}}{e(H_2) p_2^{e(F)-1}} \right)^{k_1}}{\left( \left( \frac{p_1+x}{p_1} \right)^{\frac{\ell_1}{k_1-1}} \left( \frac{p_1}{e(H_1)} \right) + \left( \frac{p_2-x}{p_2} \right)^{\frac{\ell_2}{k_1-1}} \left( \frac{p_2}{e(H_2)} \right) \right)^{k_1-1}}$$



for all  $(x, y)$  in the domain of  $f$ . Applying Theorem A.1 to both the numerator and denominator, where the denominator also uses (5.9), we get a lower bound of

$$\frac{\left(\frac{p_1}{e(H_1)} + \frac{xe(F)}{e(H_1)} + \frac{p_2}{e(H_2)} - \frac{xe(F)}{e(H_2)}\right)^{k_1}}{\left(\frac{p_1}{e(H_1)} + \frac{x\ell_1}{(k_1-1)e(H_1)} + \frac{p_2}{e(H_2)} - \frac{x\ell_2}{(k_1-1)e(H_2)}\right)^{k_1-1}}.$$

A bit of rearranging yields

$$\frac{\left(\frac{p_1}{e(H_1)} + \frac{p_2}{e(H_2)} + \frac{xe(F)(\ell_1-\ell_2)}{e(H_1)e(H_2)}\right)^{k_1}}{\left(\frac{p_1}{e(H_1)} + \frac{p_2}{e(H_2)} + \frac{k_1xe(F)(\ell_1-\ell_2)}{(k_1-1)e(H_1)e(H_2)}\right)^{k_1-1}}. \quad (5.18)$$

If  $\ell_2 = \ell_1$ , then this expression simplifies to  $\frac{p_1}{e(H_1)} + \frac{p_2}{e(H_2)}$  and we are done. Otherwise, setting the derivative of this expression with respect to  $x$  equal to zero yields

$$\begin{aligned} & \frac{k_1 \left(\frac{p_1}{e(H_1)} + \frac{p_2}{e(H_2)} + \frac{xe(F)(\ell_1-\ell_2)}{e(H_1)e(H_2)}\right)^{k_1-1} \left(\frac{e(F)(\ell_1-\ell_2)}{e(H_1)e(H_2)}\right)}{\left(\frac{p_1}{e(H_1)} + \frac{p_2}{e(H_2)} + \frac{xe(F)(\ell_1-\ell_2)k_1}{(k_1-1)e(H_1)e(H_2)}\right)^{k_1-1}} \\ &= \frac{(k_1-1) \left(\frac{p_1}{e(H_1)} + \frac{p_2}{e(H_2)} + \frac{xe(F)(\ell_1-\ell_2)}{e(H_1)e(H_2)}\right)^{k_1} \left(\frac{e(F)(\ell_1-\ell_2)k_1}{(k_1-1)e(H_1)e(H_2)}\right)}{\left(\frac{p_1}{e(H_1)} + \frac{p_2}{e(H_2)} + \frac{xe(F)(\ell_1-\ell_2)k_1}{(k_1-1)e(H_1)e(H_2)}\right)^{k_1}} \end{aligned}$$

which implies that

$$\frac{p_1}{e(H_1)} + \frac{p_2}{e(H_2)} + \frac{xe(F)(\ell_1-\ell_2)k_1}{(k_1-1)e(H_1)e(H_2)} = \frac{p_1}{e(H_1)} + \frac{p_2}{e(H_2)} + \frac{xe(F)(\ell_1-\ell_2)}{e(H_1)e(H_2)}$$

and so  $x = 0$ . The second derivative evaluated at  $x = 0$  is

$$\frac{k_1(k_1-1)}{\frac{p_1}{e(H_1)} + \frac{p_2}{e(H_2)}} \left( \frac{e(F)(\ell_1-\ell_2)}{e(H_1)e(H_2)} - \frac{e(F)(\ell_1-\ell_2)k_1}{(k_1-1)e(H_1)e(H_2)} \right)^2$$

which is clearly positive, and so the expression in (5.18) is minimized at  $x = 0$ . Therefore,  $f(x, y) \geq \frac{p_1}{e(H_1)} + \frac{p_2}{e(H_2)}$  and so we are done in this case.  $\square$

## 6 Binomial Inequalities Via Entropy

As we have seen in Lemmas 4.2 and 5.8, certain types of binomial inequalities for homomorphism densities allow us to “propagate” the property of being  $(p_1, p_2)$ -common from one pair of graphs to another. The goal of this section is to describe an approach to proving such

inequalities that uses the information-theoretic notion of entropy and apply this approach to a certain family of graphs. This idea seems to have first appeared in a paper of Kopyarty and Rossman [30]. It has had a major impact on our understanding of Sidorenko's Conjecture [7, 8, 49] and related problems [21, 33]. In the applications of this method, it will be more convenient to deal with finite graphs as opposed to graphons, since it allows us to speak in terms of discrete random variables rather than continuous ones.

Suppose that  $H$  and  $F$  are graphs and that  $k$ ,  $\ell$  and  $m$  are positive rationals such that  $e(H) + \ell = ke(F)$  and  $v(H) + 2\ell + m = kv(F)$ . Let  $r$  be a positive integer so that  $rk$ ,  $r\ell$  and  $rm$  are all integers. Our aim will be to prove an inequality of the form

$$\text{hom}(H, G) \cdot \text{hom}(K_2, G)^\ell \cdot \text{hom}(K_1, G)^m \geq \text{hom}(F, G)^k \quad (6.1)$$

holds for all graphs  $G$ . If this holds, then, by dividing both sides by  $v(G)^{kv(F)}$ , we get that

$$t(H, G) \cdot t(K_2, G)^\ell \geq t(F, G)^k$$

which is precisely the type of inequality that we require in order to apply Lemma 4.2.

An elegant method for proving inequalities of the form given by (6.1) is to construct a distribution on homomorphisms from the disjoint union of  $r$  copies of  $H$ ,  $r\ell$  copies of  $K_2$  and  $rm$  copies of  $K_1$  into  $G$  which has entropy at least  $r$  times the entropy of a uniformly random homomorphism from  $F$  to  $G$ . As it turns out the existence of such a distribution implies a lower bound on the total number of homomorphisms from  $H$  to  $G$  which yields an inequality as in (6.1) (see Lemma 6.4 below). Let us formalize this now.

**Definition 6.2.** The *range* of a discrete random variable  $X$  is  $\text{rng}(X) = \{x : \mathbb{P}(X = x) > 0\}$ .

**Definition 6.3.** The *entropy* of a discrete random variable  $X$  is defined to be

$$\mathbb{H}(X) = \sum_{x \in \text{rng}(X)} \mathbb{P}(X = x) \log_2 \left( \frac{1}{\mathbb{P}(X = x)} \right).$$

We remark that the specific choice of the base of the logarithm will be unimportant in this paper. The following lemma follows from a simple application of Jensen's Inequality.

**Lemma 6.4.** *For any discrete random variable  $X$ ,*

$$\mathbb{H}(X) \leq \log_2(|\text{rng}(X)|)$$

*with equality if and only if  $X$  is uniformly distributed on  $\text{rng}(X)$ .*

Given a graph  $G$ , a tuple  $(X_v : v \in V(H))$  of random variables taking values in  $V(G)^{V(H)}$  is  $(H, G)$ -homomorphism supported if, for every  $(x_v : v \in V(H)) \in \text{rng}(X_v : v \in V(H))$ , the function  $f : V(H) \rightarrow V(G)$  defined by  $f(v) = x_v$  for all  $v \in V(H)$  is a homomorphism. Suppose that  $H$  and  $F$  are graphs such that, for every graph  $G$  and every  $(F, G)$ -homomorphism

supported random variable  $(X_v : v \in V(F))$ , there exists an  $(r \cdot H \sqcup r\ell \cdot K_2 \sqcup rm \cdot K_1, G)$ -homomorphism supported random variable  $(Y_v : v \in V(r \cdot H \sqcup r\ell \cdot K_2 \sqcup rm \cdot K_1))$  such that

$$\mathbb{H}(Y_v : v \in V(r \cdot H \sqcup r\ell \cdot K_2 \sqcup rm \cdot K_1)) \geq rk\mathbb{H}(X_v : v \in V(F)).$$

Then, in particular, this holds when  $(X_v : v \in V(F))$  is the  $(F, G)$ -homomorphism supported variable of maximum entropy. By applying Lemma 6.4 twice, we get

$$\begin{aligned} \log_2(\text{hom}(H, G)^r \text{hom}(K_2, G)^{r\ell} \text{hom}(K_1, G)^{rm}) &\geq \mathbb{H}(Y_v : v \in V(r \cdot H \sqcup r\ell \cdot K_2 \sqcup rm \cdot K_1)) \\ &\geq rk\mathbb{H}(X_v : v \in V(F)) = rk \log_2(\text{hom}(F, G)). \end{aligned}$$

Therefore, we get that (6.1) holds. Thus, the approach boils down to constructing distributions on homomorphisms of disjoint unions of copies of  $H$ ,  $K_2$  and  $K_1$  to an arbitrary graph  $G$  from distributions on homomorphisms from  $F$  to  $G$ . The notion of conditional entropy will be particularly valuable in describing and analyzing such distributions.

**Definition 6.5.** Let  $X$  and  $Y$  be discrete random variables and let  $y \in \text{rng}(Y)$ . The *range of  $X$  given  $Y = y$*  is the set  $\text{rng}(X | Y = y) = \{x : \mathbb{P}(X = x | Y = y) > 0\}$ .

**Definition 6.6.** Let  $X$  and  $Y$  be discrete random variables and let  $y \in \text{rng}(Y)$ . The *conditional entropy of  $X$  given  $Y = y$*  is

$$\mathbb{H}(X | Y = y) = \sum_{x \in \text{rng}(X|Y=y)} \mathbb{P}(X = x | Y = y) \log_2 \left( \frac{1}{\mathbb{P}(X = x | Y = y)} \right).$$

**Definition 6.7.** Let  $X$  and  $Y$  be discrete random variables. The *conditional entropy of  $X$  given  $Y$*  is

$$\mathbb{H}(X | Y) = \sum_{y \in \text{rng}(Y)} \mathbb{P}(Y = y) \mathbb{H}(X | Y = y).$$

**Definition 6.8.** Let  $X, Y$  and  $Z$  be discrete random variables. We say that  $X$  and  $Y$  are *conditionally independent given  $Z$*  if

$$\mathbb{P}(X = x | Z = z, Y = y) = \mathbb{P}(X = x | Z = z) \text{ and}$$

$$\mathbb{P}(Y = y | Z = z, X = x) = \mathbb{P}(Y = y | Z = z)$$

for every  $z \in \text{rng}(Z)$ ,  $x \in \text{rng}(X | Z = z)$  and  $y \in \text{rng}(Y | Z = z)$ .

The main properties of conditional entropy that we will require are summarized in the following lemma.

**Lemma 6.9.** *For any integers  $m \geq 1$  and  $k \geq 0$  and discrete random variables  $X_1, X_2, \dots, X_m$  and  $Y_1, Y_2, \dots, Y_k$ ,*

$$(6.10) \quad \mathbb{H}(X_1, \dots, X_m | Y_1, \dots, Y_k) = \sum_{i=1}^m \mathbb{H}(X_i | Y_1, \dots, Y_k, X_1, \dots, X_{i-1}),$$

$$(6.11) \quad \text{If } k \geq 1, \text{ then } \mathbb{H}(X_1, \dots, X_m \mid Y_1, \dots, Y_k) \leq \mathbb{H}(X_1, \dots, X_m \mid Y_1, \dots, Y_{k-1}),$$

$$(6.12) \quad \text{If } k \geq 1 \text{ and } (X_1, \dots, X_m) \text{ and } Y_k \text{ are conditionally independent given } (Y_1, \dots, Y_{k-1}), \\ \text{then } \mathbb{H}(X_1, \dots, X_m \mid Y_1, \dots, Y_k) = \mathbb{H}(X_1, \dots, X_m \mid Y_1, \dots, Y_{k-1}).$$

The following lemma is very useful in constructing distributions of high entropy; c.f. [33, Lemma 2.5].

**Lemma 6.13.** *Let  $Y_1, Y_2, Y'_2$  and  $Y_3$  be discrete random variables such that  $Y_2$  and  $Y'_2$  are identically distributed. Then there exist discrete random variables  $Z_1, Z_2$  and  $Z_3$  such that  $(Z_1, Z_2)$  and  $(Y_1, Y_2)$  are identically distributed,  $(Z_2, Z_3)$  and  $(Y'_2, Y_3)$  are identically distributed and  $Z_1$  and  $Z_3$  are conditionally independent given  $Z_2$ . In particular,*

$$\mathbb{H}(Z_1, Z_2, Z_3) = \mathbb{H}(Y_1, Y_2) + \mathbb{H}(Y'_2, Y_3) - \mathbb{H}(Y_2).$$

*Proof Sketch.* Sample  $(Z_1, Z_2)$  according to the distribution of  $(Y_1, Y_2)$ . Now, independently of  $(Z_1, Z_2)$ , sample  $(Z'_2, Z_3)$  according to the distribution of  $(Y'_2, Y_3)$ . Resample  $(Z'_2, Z_3)$  repeatedly until  $Z'_2 = Z_2$  (without resampling  $Z_1$  or  $Z_2$ ). The random variables  $Z_1, Z_2$  and  $Z_3$  have the desired properties.  $\square$

The way in which Lemma 6.13 tends to be used in the context of homomorphism densities is to combine a  $(F_1, G)$ -homomorphism supported random variable and a  $(F_2, G)$ -homomorphism supported random variable to get a  $(H, G)$ -homomorphism supported random variable, where  $H$  is obtained from  $F_1$  and  $F_2$  by a certain type of “gluing” operation, in such a way that the entropy of the combined distribution can be analyzed in terms of the other two. The next example, which is similar in spirit to [25, Theorem 2], illustrates a particularly simple gluing operation.

**Example 6.14.** Suppose that  $F$  is a graph and  $S \subseteq V(F)$  such that the subgraph of  $F$  induced by  $S$  has maximum degree one. Let  $H$  be the graph obtained from two disjoint copies of  $F$  by identifying each vertex in the first copy that is contained in  $S$  with its counterpart in the other copy. Then Lemma 6.13 tells us that, for any graph  $G$  such that  $\text{hom}(F, G) > 0$ , there is a distribution on  $\text{Hom}(H, G)$  with entropy  $2\mathbb{H}(X_v : v \in V(F)) - \mathbb{H}(X_v : v \in S)$  for any  $(F, G)$ -supported random variable  $(X_v : v \in V(F))$ . Thus, by Lemma 6.4,

$$\log_2(\text{hom}(H, G)) \geq 2\mathbb{H}(X_v : v \in V(F)) - \mathbb{H}(X_v : v \in S).$$

Let  $\ell$  be the number of edges of  $F$  contained in  $S$  and let  $m = |S| - 2\ell$ . Note that  $e(H) = 2e(F) - \ell$ . Then  $(X_v : v \in S)$  gives rise to a random homomorphism from  $\ell \cdot K_2 \sqcup m \cdot K_1$  to  $G$ . Thus, taking  $(X_v : v \in V(F))$  to be uniform on  $\text{Hom}(F, G)$ , we see, by two applications of Lemma 6.4, that

$$\begin{aligned} 2\mathbb{H}(X_v : v \in V(F)) - \mathbb{H}(X_v : v \in S) &= 2\log_2(\text{hom}(F, G)) - \mathbb{H}(X_v : v \in S) \\ &\geq 2\log_2(\text{hom}(F, G)) - \log_2(\text{hom}(\ell \cdot K_2 \sqcup m \cdot K_1, G)) = \log \left( \frac{\text{hom}(F, G)^2}{\text{hom}(K_2, G)^\ell \text{hom}(K_1, G)^m} \right). \end{aligned}$$

Putting all of this together, we get that (6.1) holds for this particular choice of  $H$  and  $F$ .

Let us now extend the idea in Example 6.14 to allow for more general gluing operations. The following two definitions describe the nature of these operations. Given a set  $X$ , let  $2^X := \{Y : Y \subseteq X\}$ .

**Definition 6.15.** Given a graph  $F$ , an  $F$ -tree is a pair  $(T, \varphi)$  such that  $T$  is a tree and  $\varphi : V(T) \cup E(T) \rightarrow 2^{V(F)}$  such that  $\varphi(st) \subseteq \varphi(s) \cap \varphi(t)$  for every edge  $st$  of  $T$ .

**Definition 6.16.** Given a graph  $F$  and an  $F$ -tree  $(T, \varphi)$ , let  $H(T, \varphi)$  be the graph obtained by taking a copy  $J_t$  of  $F[\varphi(t)]$  for each  $t \in V(T)$ , all of which are vertex disjoint, and then identifying each vertex of  $J_s$  corresponding to an element of  $\varphi(st)$  with the corresponding vertex of  $J_t$  for all  $st \in E(T)$ .




Figure 1: An example of an  $C_5$ -tree  $(T, \varphi)$  and the graph  $H(T, \varphi)$ .

The key result of this section is the following lemma. We remark that this statement is not new; it has appeared in different (and often more general) forms in, e.g., [7, 21, 30, 33, 49].

**Lemma 6.17.** *Let  $F$  be a graph, let  $(T, \varphi)$  be an  $F$ -tree and let  $H = H(T, \varphi)$ . Suppose that  $G$  is a graph such that  $\text{hom}(F, G) > 0$ . Then, for any  $(F, G)$ -supported random variable  $(X_v : v \in V(F))$ , there exists an  $(H, G)$ -homomorphism supported random variable  $(Y_v : v \in V(H))$  with entropy equal to*

$$\sum_{t \in V(T)} (X_v : v \in \varphi(t)) - \sum_{st \in E(T)} (X_v : v \in \varphi(st)). \quad (6.18)$$

*Proof.* What we will show is that there is an  $(H, G)$ -homomorphism supported random variable  $(Y_v : v \in V(H))$  such that, for every  $t \in V(T)$ , the restriction of this random variable to the vertices from the corresponding copy of  $F[\varphi(t)]$  is distributed according to  $(X_v : v \in \varphi(t))$  and, moreover, that the entropy of this distribution is the expression in the lemma statement. We proceed by induction on  $|V(T)|$ . In the case that  $|V(T)| = 1$ , we have

that  $H = F[\varphi(t)]$  for the unique vertex  $t$  of  $T$ . In this case, the desired random variable is simply  $(X_v : v \in \varphi(t))$ .

Now, suppose that  $|V(T)| \geq 2$ , let  $s$  be any leaf of  $T$  and let  $t$  be its unique neighbour. Let  $T' = T \setminus \{t\}$  and  $\varphi'$  be the restriction of  $\varphi$  to  $V(T')$ . Define  $H' = H(T', \varphi')$ . By induction, we let  $(Y_v : v \in V(H'))$  be an  $(H', G)$ -homomorphism supported random variable which satisfies the condition described at the beginning of the previous paragraph. Our goal is to apply Lemma 6.13. Let  $Y_1$  be the tuple consisting of  $Y_v$  for all vertices of  $H'$  that are not in the copy of  $F[\varphi(st)]$  contained within the copy of  $F[\varphi(s)]$  corresponding to  $s$ . Let  $Y_2$  be the tuple consisting of  $Y_v$  for all other vertices  $v$  of  $H'$ . Also, let  $Y_2' = (X_v : v \in \varphi(st))$  and  $Y_3 = (X_v : v \in \varphi(t) \setminus \varphi(st))$ . Then, by the inductive hypothesis, these four variables satisfy the conditions of Lemma 6.13. Let  $(Z_v : v \in V(H))$  be the random variable obtained by applying Lemma 6.13 to these four random variables. Now, to compute its entropy, we use the conditional independence guaranteed by Lemma 6.13 to get

$$\mathbb{H}(Z_v : v \in V(H)) = \mathbb{H}(Y_v : v \in V(H')) + \mathbb{H}(X_v : v \in \varphi(t)) - \mathbb{H}(X_v : v \in \varphi(st))$$

by Lemma 6.9 (6.12). Applying the induction hypothesis to  $(Y_v : v \in V(H'))$  completes the proof.  $\square$

## 6.1 Trees of Odd Cycles

Our next goal is to use the notion of entropy to prove binomial inequalities in homomorphism density for  $C_k$ -trees, where  $k$  is odd, under certain conditions that are described by the following definitions. In what follows, we always assume that the vertices of  $C_k$  are  $0, 1, \dots, k-1$  and two vertices are adjacent if they are consecutive modulo  $k$ .

**Definition 6.19.** For odd  $k$ , a  $C_k$ -tree  $(T, \varphi)$  is said to be *good* if

$$(6.20) \quad C_k[\varphi(t)] \text{ is a path for every } t \in V(T) \text{ and}$$

$$(6.21) \quad \text{either } \varphi(st) \text{ is a path or } |\varphi(st)| \leq 2 \text{ for all } st \in E(T).$$

**Definition 6.22.** Given a good  $C_k$ -tree  $(T, \varphi)$  and  $0 \leq j \leq k-2$ , define

$$(6.23) \quad \pi_j(T, \varphi) \text{ to be the number of } t \in V(T) \text{ such that } C_k[\varphi(t)] \simeq P_{j+1} \text{ minus the number of } st \in E(T) \text{ such that } C_k[\varphi(st)] \simeq P_{j+1},$$

$$(6.24) \quad \gamma_j(T, \varphi) \text{ to be the number of } st \in E(T) \text{ such that } \varphi(st) \text{ is an independent set of size two whose vertices are at distance } \min\{j, k-j\} \text{ in } C_k.$$

$$(6.25) \quad \vec{\sigma}(T, \varphi) = (\pi_j(T, \varphi) - \gamma_j(T, \varphi) : j = 0, \dots, k-2).$$

**Observation 6.26.** We have  $\gamma_j(T, \varphi) = 0$  for  $j \in \{0, 1\}$ . Also,  $\gamma_j(T, \varphi) = \gamma_{k-j}(T, \varphi)$  for all  $0 \leq j \leq k-2$ .

**Definition 6.27.** A  $C_k$ -tree  $(T, \varphi)$  is *very good* if it is good,  $H(T, \varphi)$  has exactly

$$k \cdot \sum_{j=0}^{\lfloor k/2 \rfloor} \gamma_j(T, \varphi)$$

vertices and edges and  $\vec{\sigma}(T, \varphi)$  is in the convex cone generated by  $\{-\vec{e}_j + 2\vec{e}_{j+1} - \vec{e}_{j+2} : j = 0, \dots, k-4\}$ , where  $\vec{e}_0, \dots, \vec{e}_k$  is the standard basis of  $\mathbb{R}^{k+1}$ .

The condition on  $\vec{\sigma}(T, \varphi)$  in Definition 6.27 will be used in conjunction with the following lemma.

**Lemma 6.28.** *Let  $G$  be a graph with  $\text{hom}(C_k, G) > 0$  and let  $(X_0, X_1, \dots, X_{k-1})$  be a  $(C_k, G)$ -homomorphism supported random variable of maximum entropy. Then, for any  $1 \leq j \leq k-2$ ,*

$$-\mathbb{H}(X_0, \dots, X_{j-1}) + 2\mathbb{H}(X_0, \dots, X_j) - \mathbb{H}(X_0, \dots, X_{j+1}) \geq 0.$$

*Proof.* By Lemma 6.4 and the fact that  $(X_0, X_1, \dots, X_{k-1})$  is a  $(C_k, G)$ -homomorphism supported random variable of maximum entropy, we see the function  $f : V(C_k) \rightarrow V(G)$  defined by  $f(v) = X_v$  for  $v \in V(C_k)$  is a uniformly random homomorphism from  $C_k$  to  $G$ . By symmetry,  $\mathbb{H}(X_0, \dots, X_{j-1}) = \mathbb{H}(X_1, \dots, X_j)$ . So, by Lemma 6.9 (6.10),

$$\mathbb{H}(X_0, \dots, X_j) - \mathbb{H}(X_0, \dots, X_{j-1}) = \mathbb{H}(X_0, \dots, X_j) - \mathbb{H}(X_1, \dots, X_j) = \mathbb{H}(X_0 \mid X_1, \dots, X_j).$$

By Lemma 6.9 (6.11), this is at least  $\mathbb{H}(X_0 \mid X_1, \dots, X_{j+1})$  which is equal to

$$\mathbb{H}(X_0, \dots, X_{j+1}) - \mathbb{H}(X_1, \dots, X_{j+1})$$

by Lemma 6.9 (6.10). By symmetry,  $\mathbb{H}(X_1, \dots, X_{j+1}) = \mathbb{H}(X_0, \dots, X_j)$ . Now, a bit of rearranging yields

$$-\mathbb{H}(X_0, \dots, X_{j-1}) + 2\mathbb{H}(X_0, \dots, X_j) - \mathbb{H}(X_0, \dots, X_{j+1}) \geq 0$$

as desired. □

Finally, let us put this all together to derive the key lemma of the section.

**Lemma 6.29.** *If  $k$  is odd,  $(T, \varphi)$  is a very good  $C_k$ -tree and  $H = H(T, \varphi)$ , then*

$$\text{hom}(H, G) \geq \text{hom}(C_k, G)^{e(H)/k}$$

*for every graph  $G$ .*

*Proof.* Let  $G$  be a graph with  $\text{hom}(C_k, G) > 0$  and let  $(X_0, \dots, X_{k-1})$  be an  $(F, G)$ -homomorphism supported random variable of maximum entropy. Then  $\mathbb{H}(X_0, \dots, X_{k-1}) = \log_2(\text{hom}(C_k, G))$ . By Lemma 6.17, there is an  $(H, G)$ -homomorphism supported random variable with entropy

$$\sum_{t \in V(T)} (X_v : v \in \varphi(t)) - \sum_{st \in E(T)} (X_v : v \in \varphi(st)). \quad (6.30)$$

By Lemma 6.4, this implies that  $\log_2(\text{hom}(H, G))$  is bounded below by this quantity. Thus, we will be done if we can bound the above expression below by  $(|E(H)|/k) \log_2(\text{hom}(C_k, G))$ .

For  $0 \leq j \leq k-1$ , define  $p_j := \mathbb{H}(X_0, \dots, X_j)$  and  $g_j := \mathbb{H}(X_0, X_j)$ . Note that  $p_j + p_{k-j} - g_j = \log_2(\text{hom}(C_k, G))$  by Lemma 6.9 (6.10). Using Definition 6.22, we can rewrite (6.30) as

$$\begin{aligned} & \sum_{j=0}^k \pi_j(T, \varphi) \cdot p_j - \sum_{j=0}^{\lfloor k/2 \rfloor} \gamma_j(T, \varphi) \cdot g_j \\ &= \sum_{j=0}^k (\pi_j(T, \varphi) - \gamma_j(T, \varphi)) \cdot p_j + \sum_{j=0}^{\lfloor k/2 \rfloor} \gamma_j(T, \varphi) \cdot (p_j + p_{k-j} - g_j) \\ &= \sum_{j=0}^k (\pi_j(T, \varphi) - \gamma_j(T, \varphi)) \cdot p_j + (|E(H)|/k) \log_2(\text{hom}(C_k, G)). \end{aligned}$$

So, all that remains is to prove that

$$\sum_{j=0}^k (\pi_j(T, \varphi) - \gamma_j(T, \varphi)) \cdot p_j \geq 0.$$

By the the assumption on  $\vec{\sigma}$  in Definition 6.27, the expression on the left side can be written as a linear combination of expressions of the form  $-p_j + 2p_{j+1} - p_{j+2}$  for  $0 \leq j \leq k-4$  with non-negative coefficients. By Lemma 6.28, all such expressions are non-negative. This completes the proof.  $\square$

**Definition 6.31.** Given a  $C_k$ -tree  $(T, \varphi)$ , let  $H'(T, \varphi)$  be the graph obtained from  $H(T, \varphi)$  by deleting all components on at most two vertices.

**Corollary 6.32.** For odd  $k \geq 3$ , let  $(T, \varphi)$  be a very good  $C_k$ -tree, let  $\ell$  be the number of 2-vertex components of  $H(T, \varphi)$  and let  $H = H'(T, \varphi)$ . Then, for every graph  $G$ ,

$$t(H, G) \geq \frac{t(C_k, G)^{(e(H)+\ell)/k}}{t(K_2, G)^\ell}.$$

*Proof.* Let  $m$  be the number of singleton components of  $H'(T, \varphi)$ . By Lemma 6.29,

$$\text{hom}(H, G) \text{hom}(K_1, G)^m \text{hom}(K_2, G)^\ell = \text{hom}(H(T, \varphi), G) \geq \text{hom}(C_k, G)^{(e(H)+\ell)/k}.$$

The result follows by rearranging and dividing both sides by  $v(G)^{v(H)}$ .  $\square$

The following result generalizes [21, Theorem 1.2] to the asymmetric setting and to graphs built up from odd cycles of arbitrary length, as opposed to just triangles. It also addresses, but does not completely settle, one of the questions in [21, Section 5].

**Theorem 6.33.** Let  $k \geq 3$  be odd. If  $(T_1, \varphi_1)$  and  $(T_2, \varphi_2)$  are very good  $C_k$ -trees,  $\ell_i$  is the number of 2-vertex components of  $H(T_i, \varphi_i)$  for  $i \in \{1, 2\}$ ,  $H_i = H'(T_i, \varphi_i)$  for  $i \in \{1, 2\}$  and the following hold:



$$(6.34) \quad \frac{1}{k} \min\{e(H_1) + \ell_1, e(H_2) + \ell_2\} \geq \max\{\ell_1, \ell_2\} + 1,$$

$$(6.35) \quad (e(H_1) + \ell_1)\ell_2 = (e(H_2) + \ell_2)\ell_1.$$

then  $(H_1, H_2)$  is  $(1/2, 1/2)$ -common.

*Proof.* We show that the conditions of Lemma 4.2 are satisfied where  $F_1 = F_2 = C_k$  and  $k_i = \frac{e(H_i) + \ell_i}{k}$  for  $i \in \{1, 2\}$ . By [48, Corollary 1],  $C_k$  is common; thus, the pair  $(C_k, C_k)$  is  $(1/2, 1/2)$ -common. Now, (4.3) and (4.4) hold by (6.34) and (6.35), respectively. For (4.5), note that  $k_i e(C_k) - \ell_i = e(H_i)$  by definition of  $k_i$ . Also, (4.6) holds by Corollary 6.32. This completes the proof.  $\square$

In much the same way, we can apply Lemma 5.8 and Theorems 5.2 and 5.4 to get interesting examples of  $(p_1, p_2)$ -common pairs with  $p_1 \neq p_2$ .

**Theorem 6.36.** *Let  $k \geq 3$  be odd such that  $C_k$  is strongly common and let  $p_1, p_2 \in (0, 1)$  such that  $p_1 + p_2 = 1$ . If  $(T_1, \varphi_1)$  and  $(T_2, \varphi_2)$  are very good  $C_k$ -trees,  $\ell_i$  is the number of 2-vertex components of  $H(T_i, \varphi_i)$ ,  $H_i = H'(T_i, \varphi_i)$  for  $i \in \{1, 2\}$  and the following hold:*

$$(6.37) \quad \frac{e(H_i) + \ell_i}{k} \geq \ell_i + 1 \text{ for } i \in \{1, 2\},$$

$$(6.38) \quad p_2^{k-1}(e(H_1) + \ell_1)e(H_2) = p_1^{k-1}(e(H_2) + \ell_2)e(H_1) \text{ and}$$

$$(6.39) \quad \text{either } e(H_1) + \ell_1 = k \text{ or } e(H_1) + \ell_1 = e(H_2) + \ell_2.$$

then  $(H_1, H_2)$  is  $(p_1, p_2)$ -common.

*Proof.* We show that the conditions of Lemma 5.8 are satisfied where  $F = C_k$  and  $k_i = \frac{e(H_i) + \ell_i}{k}$  for  $i \in \{1, 2\}$ . Observe that (5.9), (5.11) and (5.13) follow from (6.37), (6.38) and (6.39), respectively. Also, (5.10) holds by definition of  $k_i$  and the definition of a very good  $C_k$ -tree. Finally, (5.12) holds by Corollary 6.32. This completes the proof.  $\square$

To illustrate the applicability of Lemma 6.29, we close this section with a few examples of very good  $C_5$ -trees.

**Example 6.40.** Label the vertices of  $P_6$  by  $1, 2, \dots, 6$  in the order that they appear on the path. Consider the function  $\varphi : V(P_6) \cup E(P_6) \rightarrow 2^{V(C_5)}$  defined by

$$\begin{aligned} \varphi(1) = \varphi(4) = \varphi(5) &= \{0, 1, 2\}, & \varphi(2) = \varphi(3) &= \{2, 3, 4, 0\}, & \varphi(6) &= \{0\}, \\ \varphi(12) = \varphi(34) &= \{0, 2\}, & \varphi(23) &= \{2, 3, 4\}, & \varphi(45) &= \{0\}, & \varphi(56) &= \emptyset. \end{aligned}$$


Then we get that

$$\begin{aligned} \pi_0(P_6, \varphi) &= 1 - 1 = 0, & \pi_1(P_6, \varphi) &= 0 - 0 = 0, & \pi_2(P_6, \varphi) &= 3 - 1 = 2, \\ \pi_3(P_6, \varphi) &= 2 - 0 = 2, & \gamma_2(P_6, \varphi) &= \gamma_3(P_6, \varphi) = 2. \end{aligned}$$

Thus,

$$\vec{\sigma}(P_6, \varphi) = (0, 0, 2 - 2, 0) = (0, 0, 0, 0).$$

The graph  $H = H(P_6, \varphi)$  is depicted below.



Note that  $H$  has precisely 10 vertices and edges. Also,

$$5 \cdot \sum_{j=0}^{\lfloor 5/2 \rfloor} \gamma_j(T, \varphi) = 5 \cdot 2 = 10.$$

Therefore,  $(P_6, \varphi)$  is a very good tree. It follows from Lemma 6.29 that

$$t(H, G) \geq t(C_5, G)^2$$

for any graph  $G$ .

The graph  $H'$  in the following example is a special case of the construction used in the proof of [31, Theorem 1].

**Example 6.41.** Consider the function  $\varphi : V(P_5) \rightarrow 2^{V(C_5)}$  defined by

$$\begin{aligned} \varphi(1) = \varphi(3) = \varphi(4) &= \{0, 1, 2\}, & \varphi(2) &= \{2, 3, 4, 0\}, & \varphi(5) &= \{0, 1\}, \\ \varphi(12) = \varphi(34) &= \{0, 2\}, & \varphi(23) &= \{0\}, & \varphi(45) &= \emptyset. \end{aligned}$$


Then we get that

$$\begin{aligned} \pi_0(P_5, \varphi) &= 0 - 1 = -1, & \pi_1(P_5, \varphi) &= 1 - 0 = 1, & \pi_2(P_5, \varphi) &= 3 - 0 = 3, \\ \pi_3(P_5, \varphi) &= 1 - 0 = 1, & \gamma_2(P_5, \varphi) &= \gamma_3(P_5, \varphi) = 2. \end{aligned}$$

Thus,

$$\vec{\sigma}(P_5, \varphi) = (-1, 1, 3 - 2, 1 - 2) = (-1, 1, 1, -1) = (-1, 2, -1, 0) + (0, -1, 2, -1).$$

The graph  $H = H(P_5, \varphi)$  is depicted below.



Note that  $H$  has precisely 10 vertices and edges. Also,

$$5 \cdot \sum_{j=0}^{\lfloor 5/2 \rfloor} \gamma_j(T, \varphi) = 5 \cdot 2 = 10.$$

Therefore,  $(P_5, \varphi)$  is a very good tree. It follows from Lemma 6.29 that

$$t(H, G) \geq t(C_5, G)^2$$

for any graph  $G$ . Also, for  $H' = H'(P_5, \varphi)$ , we get that

$$t(H', G) \geq \frac{t(C_5, G)^2}{t(K_2, G)}$$

for any graph  $G$  by Corollary 6.32.

## 7 The Square and the Pentagon

In this section, we investigate the values of  $p$  such that the pair  $(C_4, C_5)$  is  $(p, 1 - p)$ -common. The main proofs in this section were discovered and verified via the flag algebra method. This method was introduced in a 2007 paper of Razborov [42] and led to an explosion of new results on classical problems in extremal combinatorics. The basic idea behind the approach is that, for certain problems in extremal combinatorics, it suffices to solve a certain type of semi-definite relaxation of the problem. The relaxed problem can often be solved approximately by computer using standard semi-definite programming packages. By “rounding” the approximate solution from the computer output, it is often possible to obtain an exact solution which completely settles the original extremal problem.

Let us now formally describe the nature of the semi-definite relaxations that typically arise in applications of the flag algebra method. First, we need to introduce the standard notions of “induced density” and “injective homomorphism density.”

**Definition 7.1.** For graphs  $J$  and  $G$ , let

$$\#(J, G) = |\{S \subseteq V(G) : G[S] \simeq J\}|.$$

**Definition 7.2.** Let  $J$  and  $G$  be graphs. The *induced density* of  $J$  in  $G$  is

$$d(J, G) := \begin{cases} \#(J, G) \cdot \binom{v(G)}{v(J)}^{-1} & \text{if } v(J) \leq v(G), \\ 0 & \text{otherwise.} \end{cases}$$

We pause for a rather simple observation.

**Observation 7.3.** Let  $G$  be a graph and  $0 \leq k \leq v(G)$ . Then

$$\sum_{J: v(J)=k} d(J, G) = 1$$

where the above sum ranges over all graphs  $J$  of order  $k$ , up to isomorphism.

Let  $H$  and  $J$  be graphs and define  $\text{Hom}_{\text{inj}}(H, J)$  to be the set of injective homomorphisms from  $H$  to  $J$  and  $\text{hom}_{\text{inj}}(H, J)$  to be its cardinality.

**Definition 7.4.** Let  $J$  and  $H$  be graphs with  $v(J) \leq v(H)$ . The *injective homomorphism density* of  $H$  in  $J$  is

$$t_{\text{inj}}(H, J) := \frac{\text{hom}_{\text{inj}}(H, J)(v(J) - v(H))!}{v(J)!}.$$

If  $v(H) > v(J)$ , then  $t_{\text{inj}}(H, J) := 0$ .

The following observation follows from a simple double-counting argument and the fact that, if  $H$  is fixed  $v(G)$  is large, then almost every function from  $V(H)$  to  $V(G)$  is injective.

**Observation 7.5.** Let  $H$  and  $G$  be graphs. Then

$$t(H, G) = \sum_{J:v(J)=k} t_{\text{inj}}(H, J)d(J, G) + o(1)$$

where the  $o(1)$  term goes to 0 as  $v(G) \rightarrow \infty$ .

The ideas that we have presented so far can already prove some lower bounds for certain homomorphism density problems, but they are usually very weak. For example, suppose that  $c_1, \dots, c_m$  are reals,  $H_1, \dots, H_m$  and  $G$  are graphs and  $G_i \in \{G, \overline{G}\}$  for all  $1 \leq i \leq m$ . Suppose that one is aiming for an asymptotic lower bound on  $\sum_{i=1}^m c_i t(H_i, G_i)$  that is valid for every graph  $G$ . For each graph  $J$ , let  $J_i = J$  if  $G_i = G$  and  $J_i = \overline{J}$  otherwise. Then, by Observation 7.5, for any fixed  $k$ ,

$$\begin{aligned} \sum_{i=1}^m c_i t(H_i, G_i) &= \sum_{J:v(J)=k} \left( \sum_{i=1}^m c_i t_{\text{inj}}(H_i, J_i) \right) d(J, G) + o(1) \\ &\geq \min_{J:v(J)=k} \left( \sum_{i=1}^m c_i t_{\text{inj}}(H_i, J_i) \right) + o(1) \end{aligned} \tag{7.6}$$

where the last inequality follows from Observation 7.3. This bound is rarely, if ever, useful on its own; the clever idea behind flag algebras is to introduce some additional terms into the sum in the second expression above to “boost” the final bound.

The two main ingredients in the flag algebra method are rather elementary: (i) double-counting, and (ii) the basic fact from linear algebra that, if  $A$  is a  $q \times q$  positive semi-definite matrix, then for any real-valued column vector  $\vec{x}$  of length  $q$ , it holds that  $\vec{x}^T A \vec{x} \geq 0$ . While both ideas are simple in isolation, Razborov’s insight [42] that these two ingredients can be blended together to create an automatized method for discovering and proving new combinatorial theorems was a profound breakthrough.

Let us first explain the types of double-counting arguments that are used. Fix a graph  $F$  and a partition  $S_1 \cup S_2$  of  $V(F)$  and let  $G$  be any graph; the vertices of  $S_1$  are viewed as the *roots* of  $F$ . Then one way to count induced copies of  $F$  in  $G$  is to sum, over all subsets  $T_1$  of

$V(G)$  such that  $G[T_1] \simeq F[S_1]$ , the number of sets  $T_2$  disjoint from  $T_1$  such that there is an isomorphism from  $F$  to  $G[T_1 \cup T_2]$  in which  $S_i$  maps to  $T_i$  for  $i \in \{1, 2\}$ . This procedure will count each induced copy of  $F$  in  $G$  the same number of times; specifically, each copy of  $F$  will be counted once for each partition  $S'_1 \cup S'_2$  of  $V(F)$  for which there is an automorphism of  $F$  mapping  $S_i$  to  $S'_i$  for  $i \in \{1, 2\}$ . This is nothing more than a generalization of the classical double-counting argument used in the proof of the Handshaking Lemma.

Now, let us consider a slightly more general version of this double-counting argument. Suppose that you have two graphs  $F_1$  and  $F_2$ , partitions  $S_{i,1} \cup S_{i,2}$  of  $F_i$  for  $i \in \{1, 2\}$  and bijections  $\psi_i : S_{i,1} \rightarrow \{1, \dots, |S_{i,1}|\}$  for  $i \in \{1, 2\}$  called *labellings*. Let  $G$  be a graph and suppose that we sum, over all pairs  $(T_1, \psi)$  where  $T_1 \subseteq V(G)$  and  $\psi : T_1 \rightarrow \{1, \dots, |T_1|\}$  is a bijection, the number of pairs of sets  $T_{1,2}$  and  $T_{2,2}$  which are disjoint from each other and from  $T_1$  such that there is an isomorphism  $\phi_i$  from  $F_i$  to  $G[T_1 \cup T_{i,2}]$  such that  $S_{i,1}$  maps to  $T_1$  and  $S_{i,2}$  maps to  $T_{i,2}$  for  $i \in \{1, 2\}$  and, additionally,  $\psi_i(v) = \psi(\phi_i(v))$  for all  $v \in S_{i,1}$ . Then this sum counts (several times) induced copies of any graph  $J$  obtained from  $F_1$  and  $F_2$  by gluing each vertex of  $S_{1,1}$  to the vertex of  $S_{2,1}$  with the same label and then adding some subset of the edges between  $S_{1,2}$  and  $S_{2,2}$ . Every such graph  $J$  is counted a specific number of times based on the number of triples  $(U_1, U_{2,1}, U_{2,2}, \varphi)$  such that  $U_1 \cup U_{1,2} \cup U_{2,2}$  partitions  $V(J)$ ,  $\varphi : U_1 \rightarrow \{1, \dots, |U_1|\}$  is a bijection and there are isomorphisms  $\theta_i$  for  $i \in \{1, 2\}$  from  $F_i$  to  $J[U_1 \cup U_{i,2}]$  such that  $\psi_i(v) = \varphi(\theta_i(v))$  for all  $v \in S_{i,1}$ .

Let us introduce some notation and terminology to formalize this. Let  $f_i := (F_i, S_{i,1}, \psi_i)$  for  $i \in \{1, 2\}$ . For each graph  $J$ , let  $c(f_1, f_2, J)$  be the number of times that copies of  $J$  are counted by the procedure in the previous paragraph. For example, if  $D$  is the diamond graph (two triangles glued along an edge),  $f_1 = f_2 = (K_3, S, \psi)$ , where  $S$  is a set of size two and  $\psi$  is an arbitrary function from  $S$  to  $\{1, 2\}$ , then  $c(f_1, f_2, D) = 4$  since there are two choices of the partition  $U_1 \cup U_{1,2} \cup U_{2,2}$  and, for every such partition, there are two ways to label  $U_1$ . The tuples  $f_1$  and  $f_2$  are referred to as *flags*; they are said to be *compatible* if  $F_1[S_{1,1}] \simeq F_2[S_{2,1}]$ .

Let  $f_1 = (F_1, S_{1,1}, \psi_1)$  and  $f_2 = (F_2, S_{2,1}, \psi_2)$  be compatible flags and let  $S_{i,2} = V(F_i) \setminus S_{i,1}$  for  $i \in \{1, 2\}$ . For each  $i \in \{1, 2\}$ , graph  $G$ , set  $T_1 \subseteq V(G)$  and bijection  $\psi : T_1 \rightarrow \{1, \dots, |T_1|\}$ , let  $d_G(f, T_1, \psi)$  be the number of pairs  $(\phi_i, T_{i,2})$  such that  $T_{i,2} \subseteq V(G)$  is disjoint from  $T_1$  and  $\phi_i$  is an isomorphism from  $F_i$  to  $G[T_1, T_{i,2}]$  such that  $S_{i,1}$  maps to  $T_1$ ,  $S_{i,2}$  maps to  $T_{i,2}$  and  $\psi_i(v) = \psi(\phi_i(v))$  for all  $v \in S_{i,1}$ . Then the above discussion implies that

$$\sum_{(T_1, \psi)} d_G(f_1, T_1, \psi) \cdot d_G(f_2, T_1, \psi) = \sum_J c(f_1, f_2, J) \cdot \#(J, G) + O(v(G)^{v(F_1)+v(F_2)-|S_1|-1}) \quad (7.7)$$

where the error term comes from the fact that  $d_G(f_1, T_1) \cdot d_G(f_2, T_1)$  counts cases in which the sets  $T_{1,2}$  and  $T_{2,2}$  have non-empty intersection while the expression on the right side does not.

Now, for the second main ingredient, suppose that  $A$  is a  $q \times q$  positive semidefinite matrix and let  $f_1, \dots, f_q$  be compatible flags. Let  $\vec{f}$  be the column vector of length  $q$  whose entries are  $f_1, \dots, f_q$ . Let  $A(i, j)$  be the entry on the  $i$ th row and  $j$ th column of  $A$ . Then,

by standard linear algebra, we have that, for any graph  $G$ , set  $T \subseteq V(G)$  and bijection  $\psi : T \rightarrow \{1, \dots, |T|\}$ ,

$$\sum_{i=1}^q \sum_{j=1}^q A(i, j) d_G(f_i, T, \psi) \cdot d_G(f_j, T, \psi) \geq 0. \quad (7.8)$$

Letting  $d_G(\vec{f}, T, \psi)$  be the column vector consisting of  $d_G(f_1, T, \psi), \dots, d_G(f_q, T, \psi)$  for each  $T \subseteq V(G)$  and bijection  $\psi : T \rightarrow \{1, \dots, |T|\}$ , and letting  $\ell = v(F_1) + v(F_2) - |S_1|$ , by (7.7) and (7.8), we get

$$\begin{aligned} & \sum_J \left( \sum_{i=1}^q \sum_{j=1}^q A(i, j) c(f_i, f_j, J) \right) d(J, G) \\ &= \binom{v(G)}{\ell}^{-1} \sum_{(T, \psi)} d_G(\vec{f}, T, \psi) \cdot A \cdot d_G(\vec{f}, T, \psi)^T - o(1) \geq -o(1). \end{aligned} \quad (7.9)$$

Let us come back to the problem of proving a lower bound on an expression of the form  $\sum_{i=1}^m c_i t(H_i, G_i)$ . We can now use (7.9) to “boost” the bound in (7.6) as follows. Let  $q_1, \dots, q_r$  be positive integers. For  $1 \leq t \leq r$ , let  $A_t$  be a  $q_t \times q_t$  positive semidefinite matrix and let  $f_{t,1}, \dots, f_{t,q_t}$  be compatible flags. We have

$$\begin{aligned} & \sum_{i=1}^m c_i t(H_i, G_i) = \sum_{J: v(J)=k} \left( \sum_{i=1}^m c_i t_{\text{inj}}(H_i, J_i) \right) d(J, G) - o(1) \\ & \geq \sum_{J: v(J)=k} \left( \sum_{i=1}^m c_i t_{\text{inj}}(H_i, J_i) - \sum_{t=1}^r \sum_{i=1}^{q_t} \sum_{j=1}^{q_t} A_t(i, j) c(f_{t,i}, f_{t,j}, J) \right) d(J, G) - o(1) \quad (7.10) \\ & \geq \min_{J: v(J)=k} \left( \sum_{i=1}^m c_i t_{\text{inj}}(H_i, J_i) - \sum_{t=1}^r \sum_{i=1}^{q_t} \sum_{j=1}^{q_t} A_t(i, j) c(f_{t,i}, f_{t,j}, J) \right) - o(1) \end{aligned}$$


The following lemma applies everything that we have observed so far in this section to yield a sufficient condition for a pair  $(H_1, H_2)$  to be  $(p_1, p_2)$ -common.

**Lemma 7.11.** *Let  $H_1$  and  $H_2$  be graphs and  $p_1, p_2 \in (0, 1)$  such that  $p_1 + p_2 = 1$  and let  $r, q_1, \dots, q_r$  and  $k$  be integers such that  $k \geq \max\{v(H_1), v(H_2)\}$ . For  $1 \leq t \leq r$ , let  $A_t$  be a  $q_t \times q_t$  positive semidefinite matrix and let  $f_{t,1}, \dots, f_{t,q_t}$  be compatible flags. If*

$$\frac{t_{\text{inj}}(H_1, J)}{e(H_1) \cdot p_1^{e(H_1)-1}} + \frac{t_{\text{inj}}(H_2, \bar{J})}{e(H_2) \cdot p_2^{e(H_2)-1}} - \sum_{t=1}^r \sum_{i=1}^{q_t} \sum_{j=1}^{q_t} A_t(i, j) c(f_{t,i}, f_{t,j}, J) \geq \frac{p_1}{e(H_1)} + \frac{p_2}{e(H_2)}$$

for every graph  $J$  on  $k$  vertices, then  $(H_1, H_2)$  is  $(p_1, p_2)$ -common.

Let us now use this lemma to obtain some additional examples of  $(p_1, p_2)$ -common graphs. In all of these applications of Lemma 7.11, we will have  $k = 5$ ,  $r = 4$  and  $q_1 = q_2 = q_3 = q_4 = 8$ . The flags  $f_{i,j}$  for  $1 \leq i \leq 4$  and  $1 \leq j \leq 8$  that we use are depicted below. In each of these depictions, the roots are represented by square vertices, non-roots by round vertices, edges by solid lines and non-edges by dotted lines. The roots are always labelled in increasing order from left to right.



Finally, let us prove the main results of this section.

**Theorem 7.12.**  $(C_4, C_5)$  is  $(1/2, 1/2)$ -common and  $(1/3, 2/3)$ -common.

*Proof.* First, let us show that  $(C_4, C_5)$  is  $(1/2, 1/2)$ -common. We define the following four matrices:

$$A_1 := \frac{1}{1036800} \begin{bmatrix} 25704 & 6806 & 6806 & 6806 & -8112 & -8112 & -8112 & -21786 \\ 6806 & 10512 & -1080 & -1080 & 481 & 481 & -9334 & -6786 \\ 6806 & -1080 & 10512 & -1080 & 481 & -9334 & 481 & -6786 \\ 6806 & -1080 & -1080 & 10512 & -9334 & 481 & 481 & -6786 \\ -8112 & 481 & 481 & -9334 & 11304 & -720 & -720 & 6620 \\ -8112 & 481 & -9334 & 481 & -720 & 11304 & -720 & 6620 \\ -8112 & -9334 & 481 & 481 & -720 & -720 & 11304 & 6620 \\ -21786 & -6786 & -6786 & -6786 & 6620 & 6620 & 6620 & 22284 \end{bmatrix}$$

$$A_2 := \frac{1}{1036800} \begin{bmatrix} 52260 & 4029 & 4029 & 31650 & -33264 & -7632 & -7632 & -43440 \\ 4029 & 19224 & -5400 & -864 & 864 & -5469 & -6120 & -6264 \\ 4029 & -5400 & 19224 & -864 & 864 & -6120 & -5469 & -6264 \\ 31650 & -864 & -864 & 35976 & -35484 & -111 & -111 & -30192 \\ -33264 & 864 & 864 & -35484 & 37140 & 216 & 216 & 29448 \\ -7632 & -5469 & -6120 & -111 & 216 & 14586 & -1230 & 5760 \\ -7632 & -6120 & -5469 & -111 & 216 & -1230 & 14586 & 5760 \\ -43440 & -6264 & -6264 & -30192 & 29448 & 5760 & 5760 & 45192 \end{bmatrix}$$

$$A_3 := \frac{1}{1036800} \begin{bmatrix} 40104 & 7458 & 11880 & 11880 & -10095 & -10095 & -17496 & -33636 \\ 7458 & 33852 & -3888 & -3888 & 978 & 978 & -15768 & -19722 \\ 11880 & -3888 & 16560 & 3390 & -3660 & -12075 & -2463 & -9744 \\ 11880 & -3888 & 3390 & 16560 & -12075 & -3660 & -2463 & -9744 \\ -10095 & 978 & -3660 & -12075 & 15366 & -144 & 105 & 9525 \\ -10095 & 978 & -12075 & -3660 & -144 & 15366 & 105 & 9525 \\ -17496 & -15768 & -2463 & -2463 & 105 & 105 & 24372 & 13608 \\ -33636 & -19722 & -9744 & -9744 & 9525 & 9525 & 13608 & 40188 \end{bmatrix}$$

$$A_4 := \frac{1}{1036800} \begin{bmatrix} 15552 & 4912 & 4912 & 4912 & -5166 & -5166 & -5166 & -14790 \\ 4912 & 10504 & -1728 & -1728 & 824 & 824 & -8640 & -4968 \\ 4912 & -1728 & 10504 & -1728 & 824 & -8640 & 824 & -4968 \\ 4912 & -1728 & -1728 & 10504 & -8640 & 824 & 824 & -4968 \\ -5166 & 824 & 824 & -8640 & 10760 & -1694 & -1694 & 4786 \\ -5166 & 824 & -8640 & 824 & -1694 & 10760 & -1694 & 4786 \\ -5166 & -8640 & 824 & 824 & -1694 & -1694 & 10760 & 4786 \\ -14790 & -4968 & -4968 & -4968 & 4786 & 4786 & 4786 & 15336 \end{bmatrix}$$

All four of these matrices are positive semidefinite. To complete the proof, all that is required is to show that the inequality in Lemma 7.11 is satisfied for every graph  $J$  on 5 vertices. In fact, this inequality turns out to be an exact equality for every such graph. It would be incredibly tedious to check this inequality for every such graph by hand; each of these calculations has been included in Appendix B in an ancillary file submitted with the arxiv version of this paper: <https://arxiv.org/src/2208.02045/anc/commonExtraAppendices.pdf>. For the purposes of illustration, let us just verify two simple examples by hand. Suppose that  $J = K_5$ . Then  $t_{\text{inj}}(C_4, K_5) = 1$  and  $t_{\text{inj}}(C_5, \overline{K_5}) = 0$ . The only way that  $K_5$  can be expressed as a product of two flags in our proof is in the product of  $f_{4,8}$  with itself. By definition,  $c(f_{4,8}, f_{4,8}, K_5) = \binom{5}{3} \cdot 2 \cdot 6 = 120$ . Therefore, in the case  $J = K_5$ , the left side of the inequality in Lemma 7.11 is

$$\frac{1}{4 \cdot (1/2)^3} + \frac{0}{5 \cdot (1/2)^4} - \frac{15336 \cdot 120}{1036800} = \frac{9}{40} = \frac{1/2}{4} + \frac{1/2}{5}$$

as desired. For a slightly more involved example, suppose that  $J = K_3 \sqcup K_2$ . Then  $t_{\text{inj}}(C_4, J) = t_{\text{inj}}(C_5, \overline{J}) = 0$ . We have

$$c(f_{2,4}, f_{2,4}, J) = 12,$$

$$c(f_{2,4}, f_{2,5}, J) = c(f_{2,5}, f_{2,4}, J) = 12,$$

$$c(f_{4,1}, f_{4,1}, J) = 12.$$

Therefore, the left side of the inequality in Lemma 7.11 is

$$0 - \frac{35976 \cdot 12}{1036800} + \frac{35484 \cdot 12}{1036800} + \frac{35484 \cdot 12}{1036800} - \frac{15552 \cdot 12}{1036800} = \frac{9}{40} = \frac{1/2}{4} + \frac{1/2}{5}$$



as desired.

The proof for  $(1/3, 2/3)$  is similar, except that the matrices  $A_1, A_2, A_3, A_4$  are replaced with  $\frac{1}{637009920}$  multiplied by the following four matrices, respectively:

$$\begin{bmatrix} 42246144 & 582094 & 582094 & 582094 & -42135552 & -42135552 & -42135552 & -92141034 \\ 582094 & 49243392 & -15842304 & -15842304 & 7796736 & 7796736 & -46514496 & -13049856 \\ 582094 & -15842304 & 49243392 & -15842304 & 7796736 & -46514496 & 7796736 & -13049856 \\ 582094 & -15842304 & -15842304 & 49243392 & -46514496 & 7796736 & 7796736 & -13049856 \\ -42135552 & 7796736 & 7796736 & -46514496 & 138599424 & 21676032 & 21676032 & 96865624 \\ -42135552 & 7796736 & -46514496 & 7796736 & 21676032 & 138599424 & 21676032 & 96865624 \\ -42135552 & -46514496 & 7796736 & 7796736 & 21676032 & 21676032 & 138599424 & 96865624 \\ -92141034 & -13049856 & -13049856 & -13049856 & 96865624 & 96865624 & 96865624 & 312532992 \end{bmatrix}$$

$$\begin{bmatrix} 107984172 & -18972513 & -18972513 & 41642346 & -163858752 & -76966047 & -76966047 & -243081216 \\ -18972513 & 98249718 & -4893696 & -36052992 & 56408379 & -75589632 & -25408512 & 11747760 \\ -18972513 & -4893696 & 98249718 & -36052992 & 56408379 & -25408512 & -75589632 & 11747760 \\ 41642346 & -36052992 & -36052992 & 162533628 & -171260478 & -38375424 & -38375424 & -198826518 \\ -163858752 & 56408379 & 56408379 & -171260478 & 487876608 & 83816640 & 83816640 & 233625600 \\ -76966047 & -75589632 & -25408512 & -38375424 & 83816640 & 359826702 & 20736000 & 244463616 \\ -76966047 & -25408512 & -75589632 & -38375424 & 83816640 & 20736000 & 359826702 & 244463616 \\ -243081216 & 11747760 & 11747760 & -198826518 & 233625600 & 244463616 & 244463616 & 1200867840 \end{bmatrix}$$

$$\begin{bmatrix} 101172144 & 5282292 & -6580224 & -6580224 & -85681152 & -85681152 & -110108160 & -214922952 \\ 5282292 & 126774228 & -8879739 & -8879739 & -55406592 & -55406592 & -39429504 & -177831936 \\ -6580224 & -8879739 & 151234560 & -62871552 & -3815424 & -96242688 & -30360144 & -4455024 \\ -6580224 & -8879739 & -62871552 & 151234560 & -96242688 & -3815424 & -30360144 & -4455024 \\ -85681152 & -55406592 & -3815424 & -96242688 & 466255872 & -44568576 & 109264896 & 245402496 \\ -85681152 & -55406592 & -96242688 & -3815424 & -44568576 & 466255872 & 109264896 & 245402496 \\ -110108160 & -39429504 & -30360144 & -30360144 & 109264896 & 109264896 & 367946496 & 108512640 \\ -214922952 & -177831936 & -4455024 & -4455024 & 245402496 & 245402496 & 108512640 & 1267716096 \end{bmatrix}$$

$$\begin{bmatrix} 64971648 & -8197220 & -8197220 & -8197220 & -54645562 & -54645562 & -54645562 & -93533184 \\ -8197220 & 104755556 & -34466080 & -34466080 & 15801488 & 15801488 & -69672960 & -1575936 \\ -8197220 & -34466080 & 104755556 & -34466080 & 15801488 & -69672960 & 15801488 & -1575936 \\ -8197220 & -34466080 & -34466080 & 104755556 & -69672960 & 15801488 & 15801488 & -1575936 \\ -54645562 & 15801488 & 15801488 & -69672960 & 214917120 & 22387584 & 22387584 & 70060032 \\ -54645562 & 15801488 & -69672960 & 15801488 & 22387584 & 214917120 & 22387584 & 70060032 \\ -54645562 & -69672960 & 15801488 & 15801488 & 22387584 & 22387584 & 214917120 & 70060032 \\ -93533184 & -1575936 & -1575936 & -1575936 & 70060032 & 70060032 & 70060032 & 346816512 \end{bmatrix}$$

Again, the verification of the hypotheses of Lemma 7.11 has been included in Appendix C in an ancillary file submitted with the arxiv version of this paper: <https://arxiv.org/src/2208.02045/anc/commonExtraAppendices.pdf>.  $\square$

Let us now exhibit an example to show that  $(C_4, C_5)$  is not  $(p, 1 - p)$ -common when  $p$  is larger than (roughly) 0.518.

**Proposition 7.13.** *If*

$$40p^4 + 32(1 - p)p^3 - 5 > 0,$$

*then  $(C_4, C_5)$  is not  $(p, 1 - p)$ -common.*

*Proof.* Let  $W_1$  be the graphon defined by

$$W_1(x, y) = \begin{cases} 1 & \text{if } x, y \in [0, 1/2] \text{ or } x, y \in (1/2, 1], \\ 0 & \text{otherwise} \end{cases}$$

and define  $W_2 = 1 - W_1$ . Then  $t(C_4, W_1) = 1/8$  and  $t(C_5, W_2) = 0$ . Thus,  $(C_4, C_5)$  fails to be  $(p, 1 - p)$ -common for any  $p$  such that

$$\frac{(1/8)}{4p^3} + \frac{0}{5(1-p)^4} < \frac{p}{4} + \frac{1-p}{5}$$

The result now follows by reorganizing this inequality.  $\square$

## 8 More Colours

Before closing the paper, let us propose a natural multicolour generalization of Definition 1.2.

**Definition 8.1.** Let  $k \geq 2$  and  $p_1, p_2, \dots, p_k \in (0, 1)$  such that  $\sum_{i=1}^k p_i = 1$  and let  $H_1, H_2, \dots, H_k$  be non-empty graphs. We say that  $(H_1, H_2, \dots, H_k)$  is  $(p_1, p_2, \dots, p_k)$ -common if, for any graphons  $W_1, W_2, \dots, W_k$  such that  $\sum_{i=1}^k W_i = 1$ ,

$$\sum_{i=1}^k \frac{t(H_i, W_i)}{e(H_i)p_i^{e(H_i)-1}} \geq \sum_{i=1}^k \frac{p_i}{e(H_i)}.$$

Following [25, 31], a graph  $H$  is said to be  $k$ -common if the  $k$ -tuple  $(H, \dots, H)$  is  $(1/k, \dots, 1/k)$ -common. As is the case with the ordinary notion of  $k$ -common graphs, it is more difficult to satisfy Definition 8.1 when  $k$  is large. The following lemma is an asymmetric analogue of [25, Theorem 13].

**Lemma 8.2.** Let  $k \geq 2$  and  $p_1, \dots, p_k, q_{k+1} \in (0, 1)$  such that  $\sum_{i=1}^k p_i = 1$ , define  $q_i = (1 - q_{k+1})p_i$  for  $1 \leq i \leq k$  and let  $H_1, \dots, H_k, H_{k+1}$  be graphs. If  $(H_1, \dots, H_k)$  is not  $(p_1, \dots, p_k)$ -common, then  $(H_1, \dots, H_{k+1})$  is not  $(q_1, \dots, q_{k+1})$ -common.

*Proof.* Using the fact that  $(H_1, \dots, H_k)$  is not  $(p_1, \dots, p_k)$ -common, we let  $W_1, \dots, W_k$  be graphons such that  $\sum_{i=1}^k W_i = 1$  and

$$\sum_{i=1}^k \frac{t(H_i, W_i)}{e(H_i)p_i^{e(H_i)-1}} < \sum_{i=1}^k \frac{p_i}{e(H_i)}.$$

Define  $W'_i := (1 - q_{k+1})W_i$  for  $1 \leq i \leq k$  and  $W'_{k+1} = q_{k+1}$ . Then  $\sum_{i=1}^{k+1} W'_i = 1$  and

$$\sum_{i=1}^{k+1} \frac{t(H_i, W'_i)}{e(H_i)q_i^{e(H_i)-1}} = \sum_{i=1}^k \frac{(1 - q_{k+1})^{e(H_i)} t(H_i, W_i)}{e(H_i)(1 - q_{k+1})^{e(H_i)-1} p_i^{e(H_i)-1}} + \frac{q_{k+1}^{e(H_{k+1})}}{e(H_{k+1})q_{k+1}^{e(H_{k+1})-1}} = \sum_{i=1}^{k+1} \frac{q_i}{e(H_i)}.$$

This completes the proof.  $\square$

A simple convexity argument proves that, if  $H_1, \dots, H_k$  are Sidorenko, then  $(H_1, \dots, H_k)$  is  $(p_1, \dots, p_k)$  common for any  $p_1, \dots, p_k \in (0, 1)$  such that  $\sum_{i=1}^k p_i = 1$ . The first non-bipartite examples of  $k$ -common graphs for  $k \geq 3$  were obtained recently in [31]. It is also known that a graph  $H$  is  $k$ -common for all  $k \geq 2$  if and only if it is Sidorenko [31, Theorem 2]. Apart from this, we do not know of any examples of  $(p_1, p_2, \dots, p_k)$ -common tuples of graphs; it would be interesting to explore this concept in more depth.

We close this section with a necessary condition on  $(p_1, p_2, p_3)$ -common tuples graphs in terms of their girth; note that, by Lemma 8.2, this lemma implies a necessary condition on  $(p_1, \dots, p_k)$ -common graphs for any  $k \geq 3$ . This result essentially generalizes [31, Theorem 3], which is a strengthening of the main result of [12].

**Theorem 8.3.** *Let  $H_1, H_2$  and  $H_3$  be graphs and let  $p_1, p_2, p_3 \in (0, 1)$  such that  $p_1 + p_2 + p_3 = 1$ . If  $\min\{g(H_1), g(H_2), g(H_3)\}$  is odd, then  $(H_1, H_2, H_3)$  is not  $(p_1, p_2, p_3)$ -common.*

*Proof.* Let  $k = \min\{g(H_1), g(H_2), g(H_3)\}$ . Without loss of generality, we can assume that

$$\frac{c_k(H_1)}{e(H_1)p_1^{k-1}} \geq \frac{c_k(H_2)}{e(H_2)p_2^{k-1}} \geq \frac{c_k(H_3)}{e(H_3)p_3^{k-1}}. \quad (8.4)$$

For  $i \in \{1, 2, 3\}$ , let  $b_i$  be the number of subgraphs of  $H_i$  with at least  $k + 1$  non-isolated vertices.

Now, let  $B$  be the kernel as in the proof of Theorem 3.6, let  $p = \frac{1}{2} \min\{p_1, p_2, p_3\}$ , let  $\delta$  be a small real number to be specified later and define  $U_1 = 2p \cdot B^\delta$  and  $U_2 = U_3 = -p \cdot B^\delta$ . Define  $W_1 = p_1 + U_1$  and  $W_2 = W_3 = p_2 + U_2$ . By Observation 2.3,

$$\sum_{i=1}^3 \frac{t(H_i, W_i)}{e(H_i)p_i^{e(H_i)-1}} = \sum_{i=1}^3 \sum_{E \subseteq E(H_i)} \frac{t(H_i[E], U_i)}{e(H_i)p_i^{|E|-1}}. \quad (8.5)$$

Since  $U_1, U_2$  and  $U_3$  are all 0-regular, we get that  $t(F, U_i) = 0$  for any acyclic graph  $F$  and  $i \in \{1, 2, 3\}$ . In particular,  $t(H_i[E], U_i) = 0$  whenever  $H_i[E]$  has at most  $k - 1$  non-isolated vertices. If  $H_i[E]$  has exactly  $k$  non-isolated vertices, then  $t(H_i[E], U_i)$  is non-zero if and only if  $H_i[E]$  is a cycle of length  $k$ . Also, as in the proof of Theorem 3.6, we have  $t(C_k, U_1) = -(2p)^k \delta^k$  and  $t(C_k, U_2) = t(C_k, U_3) = p^k \delta^k$ . If  $F$  has at least  $k + 1$  non-isolated vertices, then  $t(F, U_i) \leq \delta^{k+1}$  for all  $i \in \{1, 2, 3\}$  by Lemma 3.2. Thus, the expression on the right side of (8.5) is bounded above by

$$\begin{aligned} & \sum_{i=1}^3 \frac{c_k(H_i) \cdot t(C_k, U_i)}{e(H_i)p_i^{k-1}} + \sum_{i=1}^3 \frac{b_i \delta^{k+1}}{e(H_i)p_i^{|E(H_i)|-1}} \\ &= \left( \frac{-c_k(H_1)(2p)^k}{e(H_1)p_1^{k-1}} + \frac{c_k(H_2)p^k}{e(H_2)p_2^{k-1}} + \frac{c_k(H_3)p^k}{e(H_3)p_3^{k-1}} \right) \delta^k + \sum_{i=1}^3 \frac{b_i \delta^{k+1}}{e(H_i)p_i^{|E(H_i)|-1}}. \end{aligned}$$

Now, by (8.4) and the fact that  $2^k > 2$ , we see that, if  $\delta$  is chosen sufficiently small with respect to  $H_1, H_2, H_3, p_1, p_2$  and  $p_3$ , the above expression is negative.  $\square$

**Remark 8.6.** Analogously to Remark 3.18, we remark that the proof of Theorem 8.3 can be adapted to show that  $(H_1, H_2, H_3)$  is not  $(p_1, p_2, p_3)$ -common in a local sense by simply scaling  $U_1, U_2$  and  $U_3$  by a factor of  $\varepsilon$  for some arbitrarily small  $\varepsilon$  and letting  $\delta$  depend additionally on  $\varepsilon$ . After applying this modification, Theorem 8.3 now generalizes [31, Theorem 3].

## 9 Open Problems

We conclude by stating a few open problems. We start with the following two conjectures which link asymmetric commonness to the standard notion of common graphs.

**Conjecture 9.1.** *If  $(H_1, H_2)$  is  $(p, 1 - p)$ -common for some  $p \in (0, 1)$ , then either  $H_1$  is common or  $H_2$  is common.*

**Conjecture 9.2.** *There exists a pair  $(H_1, H_2)$  of graphs and  $p \in (0, 1)$  such that  $H_1$  is not common, but  $(H_1, H_2)$  is  $(p, 1 - p)$ -common.*

A promising candidate for the second conjecture seems to be the pair  $(H_1, H_2)$  where  $H_1$  is the paw graph and  $H_2$  is the diamond graph. Flag algebra calculations on 5-vertex graphs strongly suggest that this pair is  $(p, 1 - p)$ -common for  $p = \frac{2\sqrt{10}-5}{3}$ ; however, we have been unable to round the matrices to turn the computer output into a rigorous proof. Next, we propose two possible converses to Corollary 4.10 of different strengths (both of which could be true).

**Question 9.3.** Suppose that  $H$  is a graph such that  $(H, H)$  is  $(p, 1 - p)$ -common for all  $p \in (0, 1)$ . Does it follow that  $H$  is Sidorenko? What if  $(H, H)$  is  $(p, 1 - p)$ -common for *some*  $p \in (0, 1) \setminus \{1/2\}$ ?

Given a pair  $H_1$  and  $H_2$  of graphs, it would be very interesting to have a better understanding of the set of all  $p$  such that  $(H_1, H_2)$  is  $(p, 1 - p)$ -common. As we have seen, it can be  $(0, 1)$  if  $H_1$  and  $H_2$  are Sidorenko, it can be empty if  $H_1$  contains a  $K_4$ , it can be a singleton if  $H_1$  and  $H_2$  have the same odd girth, and it can be a non-trivial subset of  $(0, 1)$  with more than one element if  $(H_1, H_2) = (C_4, C_5)$ . It seems very likely that it is an interval in general.

**Conjecture 9.4.** *Let  $H_1$  and  $H_2$  be graphs. The set*

$$\{p \in (0, 1) : (H_1, H_2) \text{ is } (p, 1 - p)\text{-common}\}$$

*is always an interval.*

As a first step in the direction of the previous question, it would be interesting to determine the answer for the pair  $(C_4, C_5)$ .

**Problem 9.5.** Determine the set of all  $p \in (0, 1)$  such that  $(C_4, C_5)$  is  $(p, 1 - p)$ -common.

As we have seen, the property of being strongly common is useful in obtaining examples of  $(p_1, p_2)$ -common pairs  $(H_1, H_2)$  when  $(p_1, p_2) \neq (1/2, 1/2)$ , in addition to being natural in its own right. It would be interesting to classify graphs which satisfy this stronger property. We are currently unaware of any graph which is common but not strongly common.

**Problem 9.6.** Classify strongly common graphs.

We are not aware of any examples of non-bipartite strongly common graphs apart from  $K_3$  and  $C_5$ . Given the results of Section 5, it seems highly likely that every odd cycle is strongly common, but we have been unable to prove this in general.

**Conjecture 9.7.** *Every odd cycle is strongly common.*<sup>1</sup>

Our next few problems concern the strong barrier to commonality posed by  $K_4$  subgraphs. It seems to us that  $K_4$  may be the only subgraph which automatically prevents a graph from being common.

**Conjecture 9.8.** *For any  $K_4$ -free graph  $F$ , there exists a connected common graph  $H$  containing  $F$  as a subgraph. There is also a connected common graph  $H$  containing  $F$  as an induced subgraph.*

Some strong evidence in favour of Conjecture 9.8 comes from the recent paper of Král', Volec and Wei [32], where it is shown that every graph of girth at least 50 is an induced subgraph of a common graph.

An idea used in [31, 32] is that the presence of large complete bipartite subgraphs in a graph  $H$  tends to “drive up” the homomorphism density. Thus, a natural approach to Conjecture 9.8 is to simply consider a *blow-up* of the graph  $F$ ; that is, a graph obtained by replacing each vertex of  $F$  with an independent set and adding all edges between two such sets if their corresponding vertices are adjacent. We conjecture that this strategy works.

**Conjecture 9.9.** *If  $F$  is  $K_4$ -free, then there exists a blow-up of  $F$  which is common.*

It is known that, for every bipartite graph  $F$ , there exists a blow-up of  $F$  that is Sidorenko [8]; therefore, Conjecture 9.9 holds for bipartite graphs (in a strong sense). We remark that a recent related conjecture of Grzesik et al. [21, Conjecture 1.3] asserts that any graph obtained from a Sidorenko graph by adding an arbitrary number of apex vertices is common; Sidorenko [46] proved that this is true when just one apex vertex is added.

We pose one more conjecture regarding the exceptional nature of  $K_4$ .

**Conjecture 9.10.** *If  $H_1$  is  $K_4$ -free, then there exists a graph  $H_2$  and  $p \in (0, 1)$  such that  $(H_1, H_2)$  is  $(p, 1 - p)$ -common.*

A particularly interesting special case would be to prove this conjecture when  $H_1$  is bipartite. To date, there are no examples of bipartite graphs that are known to be common but not known to be Sidorenko. Could there be graphs  $H_1$  and  $H_2$  such that  $H_1$  is bipartite

---

<sup>1</sup>Conjecture 9.7 has now been solved by Kim and Lee [28].

and not known to be Sidorenko and the pair  $(H_1, H_2)$  is  $(p, 1 - p)$ -common for some  $p$ ? We remark that the smallest bipartite graph which is not currently known to satisfy Sidorenko's Conjecture is the graph obtained from  $K_{5,5}$  by deleting the edges of a 10-cycle.

By Theorem 3.6, if  $H_1$  contains a triangle and  $(H_1, H_2)$  is  $(p, 1 - p)$ -common, then  $H_2$  must also contain a triangle and  $\frac{p}{1-p}$  must be the square root of a rational number. Thus, in particular,  $p$  must be algebraic. However, if  $H_1$  is triangle-free, then it seems unlikely that there would be any such restriction on  $p$ .

**Conjecture 9.11.** *If  $H_1$  is  $K_3$ -free, then, for every  $p \in (0, 1)$ , there exists a graph  $H_2$  such that  $(H_1, H_2)$  is  $(p, 1 - p)$ -common.*

If Conjecture 9.11 is too general, then it seems that it would already be interesting to find a *single* example of a graph  $H_1$  which satisfies it.

**Acknowledgements.** The authors would like to thank Elena Moss for helping us make the pictures of flags in Section 7.

## References

- [1] G. Blekherman and A. Raymond. A path forward: Tropicalization in extremal combinatorics. E-print arXiv:2108.06377v2, 2022.
- [2] G. Blekherman, A. Raymond, M. Singh, and R. R. Thomas. Tropicalization of graph profiles. E-print arXiv:2004.05207v2, 2022.
- [3] S. A. Burr and V. Rosta. On the Ramsey multiplicities of graphs—problems and recent results. *J. Graph Theory*, 4(4):347–361, 1980.
- [4] D. Conlon. On the Ramsey multiplicity of complete graphs. *Combinatorica*, 32(2):171–186, 2012.
- [5] D. Conlon, J. Fox, and B. Sudakov. An approximate version of Sidorenko's conjecture. *Geom. Funct. Anal.*, 20(6):1354–1366, 2010.
- [6] D. Conlon, J. Fox, and B. Sudakov. Recent developments in graph Ramsey theory. In *Surveys in combinatorics 2015*, volume 424 of *London Math. Soc. Lecture Note Ser.*, pages 49–118. Cambridge Univ. Press, Cambridge, 2015.
- [7] D. Conlon, J. H. Kim, C. Lee, and J. Lee. Some advances on Sidorenko's conjecture. *J. Lond. Math. Soc. (2)*, 98(3):593–608, 2018.
- [8] D. Conlon and J. Lee. Finite reflection groups and graph norms. *Adv. Math.*, 315:130–165, 2017.
- [9] D. Conlon and J. Lee. Sidorenko's conjecture for blow-ups. *Discrete Anal.*, pages Paper No. 2, 13, 2021.

- [10] L. N. Coregliano and A. A. Razborov. Biregularity in Sidorenko’s Conjecture. E-print arXiv:2108.06599v2, 2021.
- [11] E. Csóka, T. Hubai, and L. Lovász. Locally common graphs. E-print arXiv:1912.02926v1, 2019.
- [12] J. Cummings and M. Young. Graphs containing triangles are not 3-common. *J. Comb.*, 2(1):1–14, 2011.
- [13] P. Erdős. On the number of complete subgraphs contained in certain graphs. *Magyar Tud. Akad. Mat. Kutató Int. Közl.*, 7:459–464, 1962.
- [14] P. Erdős and M. Simonovits. Supersaturated graphs and hypergraphs. *Combinatorica*, 3(2):181–192, 1983.
- [15] P. Erdős. Some remarks on the theory of graphs. *Bull. Amer. Math. Soc.*, 53:292–294, 1947.
- [16] C. Even-Zohar and N. Linial. A note on the inducibility of 4-vertex graphs. *Graphs Combin.*, 31(5):1367–1380, 2015.
- [17] J. Fox. There exist graphs with super-exponential Ramsey multiplicity constant. *J. Graph Theory*, 57(2):89–98, 2008.
- [18] J. Fox, H. T. Pham, and Y. Zhao. Common and Sidorenko linear equations. *Q. J. Math.*, 72(4):1223–1234, 2021.
- [19] J. Fox and Y. Widgerson. Ramsey multiplicity and the Turán coloring. E-print arXiv:2207.07775v1, 2022.
- [20] A. W. Goodman. On sets of acquaintances and strangers at any party. *Amer. Math. Monthly*, 66:778–783, 1959.
- [21] A. Grzesik, J. Lee, B. Lidický, and J. Volec. On tripartite common graphs. E-print arXiv:2012.02057v1, 2020.
- [22] R. Hancock, D. Král’, M. Krnc, and J. Volec. Towards characterizing locally common graphs. E-print arXiv:2011.02562v1, 2020.
- [23] H. Hatami. Graph norms and Sidorenko’s conjecture. *Israel J. Math.*, 175:125–150, 2010.
- [24] H. Hatami, J. Hladký, D. Král’, S. Norine, and A. Razborov. Non-three-colourable common graphs exist. *Combin. Probab. Comput.*, 21(5):734–742, 2012.
- [25] C. Jagger, P. Štoviček, and A. Thomason. Multiplicities of subgraphs. *Combinatorica*, 16(1):123–141, 1996.

- [26] N. Kamčev, A. Liebenau, and N. Morrison. On uncommon systems of equations. E-print arXiv:2106.08986v2, 2021.
- [27] J. H. Kim, C. Lee, and J. Lee. Two approaches to Sidorenko’s conjecture. *Trans. Amer. Math. Soc.*, 368(7):5057–5074, 2016.
- [28] J. S. Kim and J. Lee. Extended commonality of paths and cycles via Schur convexity. E-print arXiv:2210.00977v1, 2022.
- [29] S. Ko and J. Lee. Common graphs with arbitrary connectivity and chromatic number. E-print arXiv:2207.09427v1, 2022.
- [30] S. Kopparty and B. Rossman. The homomorphism domination exponent. *European J. Combin.*, 32(7):1097–1114, 2011.
- [31] D. Král’, J. A. Noel, S. Norin, J. Volec, and F. Wei. Non-bipartite  $k$ -common graphs. *Combinatorica*, 42(1):87–114, 2022.
- [32] D. Král’, J. Volec, and F. Wei. Common graphs with arbitrary chromatic number. E-print arXiv:2206.05800v1, 2022.
- [33] J. Lee. On some graph densities in locally dense graphs. *Random Structures Algorithms*, 58(2):322–344, 2021.
- [34] L. Lovász. Subgraph densities in signed graphons and the local Simonovits-Sidorenko conjecture. *Electron. J. Combin.*, 18(1):Paper 127, 21, 2011.
- [35] L. Lovász. *Large networks and graph limits*, volume 60 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2012.
- [36] L. Lovász and B. Szegedy. Limits of dense graph sequences. *J. Combin. Theory Ser. B*, 96(6):933–957, 2006.
- [37] D. S. Mitrinović. *Analytic inequalities*. Die Grundlehren der mathematischen Wissenschaften, Band 165. Springer-Verlag, New York-Berlin, 1970. In cooperation with P. M. Vasić.
- [38] J. A. Noel. Non-bipartite  $k$ -common graphs. Online seminar talk, [https://youtu.be/BY1\\_3g1ImS0](https://youtu.be/BY1_3g1ImS0), September 2020. Extremal and Probabilistic Combinatorics Webinar, Organizers: J. Hladký, D. Piguet, J. Volec and L. Yepremyan.
- [39] O. Parczyk, S. Pokutta, C. Spiegel, and T. Szabó. New ramsey multiplicity bounds and search heuristics. E-print arXiv:2206.04036v1, 2022.
- [40] S. P. Radziszowski. Small Ramsey numbers. *Electron. J. Combin.*, 1:Dynamic Survey 1, 30, 1994.



- [41] F. P. Ramsey. On a Problem of Formal Logic. *Proc. London Math. Soc. (2)*, 30(4):264–286, 1929.
- [42] A. A. Razborov. Flag algebras. *J. Symbolic Logic*, 72(4):1239–1282, 2007.
- [43] A. Saad and J. Wolf. Ramsey multiplicity of linear patterns in certain finite abelian groups. *Q. J. Math.*, 68(1):125–140, 2017.
- [44] A. Sah, M. Sawhney, D. Stoner, and Y. Zhao. A reverse Sidorenko inequality. *Invent. Math.*, 221(2):665–711, 2020.
- [45] A. Sidorenko. A correlation inequality for bipartite graphs. *Graphs Combin.*, 9(2):201–204, 1993.
- [46] A. Sidorenko. Randomness friendly graphs. *Random Structures Algorithms*, 8(3):229–241, 1996.
- [47] A. Sidorenko. Inequalities for doubly nonnegative functions. *Electron. J. Combin.*, 28(1):Paper No. 1.32, 16, 2021.
- [48] A. F. Sidorenko. Cycles in graphs and functional inequalities. *Mat. Zametki*, 46(5):72–79, 104, 1989.
- [49] B. Szegedy. An information theoretic approach to Sidorenko’s conjecture. E-print arXiv:1406.6738v3, 2015.
- [50] A. Thomason. A disproof of a conjecture of Erdős in Ramsey theory. *J. London Math. Soc. (2)*, 39(2):246–255, 1989.
- [51] A. Thomason. Graph products and monochromatic multiplicities. *Combinatorica*, 17(1):125–134, 1997.
- [52] L. Versteegen. Common and Sidorenko equations in Abelian groups. E-print arXiv:2109.04445v2, 2021.
- [53] L. Versteegen. Linear configurations containing 4-term arithmetic progressions are uncommon. E-print arXiv:2106.06846v3, 2021.

## A Classical Convexity Inequalities

In this paper, we frequently use Bernoulli’s Inequality to deal with the different coefficients in Definition 1.2.

**Theorem A.1** (Bernoulli’s Inequality, see [37, Section 2.4]). *Let  $x \geq -1$ . If  $r \geq 1$ , then*

$$(1 + x)^r \geq 1 + rx.$$

On the other hand, if  $0 \leq r \leq 1$ , then

$$(1+x)^r \leq 1+rx.$$

We also require Hölder's Inequality in the following form.

**Theorem A.2** (Generalized Hölder's Inequality, see [37, Section 2.8]). *Let  $A = (a_{i,j})$  be an  $n \times m$  matrix of positive reals and let  $\alpha_1, \alpha_2, \dots, \alpha_m$  be positive reals such that  $\sum_{i=1}^m \alpha_i = 1$ . Then*

$$\sum_{i=1}^n \left( \prod_{j=1}^m a_{i,j}^{\alpha_j} \right) \leq \prod_{j=1}^m \left( \sum_{i=1}^n a_{i,j} \right)^{\alpha_j}.$$

By combining Theorems A.1 and A.2, we get the following.

**Lemma A.3.** *Let  $b_1, b_2, p_1, p_2, \ell_1, \ell_2$  be positive reals and let  $y$  be such that  $-p_1 \leq y \leq p_2$ . If  $d$  is an integer such that  $d \geq \max\{\ell_1, \ell_2\} + 1$ , then*

$$\frac{b_1}{(1+p_1^{-1}y)^{\ell_1}} + \frac{b_2}{(1-p_2^{-1}y)^{\ell_2}} \geq \frac{\left( p_1 \ell_1^{-1} (b_1 p_1^{-1} \ell_1)^{1/d} + p_2 \ell_2^{-1} (b_2 p_2^{-1} \ell_2)^{1/d} \right)^d}{(p_1 \ell_1^{-1} + p_2 \ell_2^{-1})^{d-1}}.$$

*Proof.* We start by applying Theorem A.2 with  $n = 2$ ,  $m = d$  and  $\alpha_j = 1/d$  for  $1 \leq j \leq d$ . In this application, we define  $\alpha_{1,1} = \frac{b_1}{(1+p_1^{-1}y)^{\ell_1}}$ ,  $\alpha_{2,1} = \frac{b_1}{(1+p_1^{-1}y)^{\ell_1}}$  and, for  $2 \leq j \leq m$ , let  $\alpha_{1,j} = p_1 \ell_1^{-1} (1+p_1^{-1}y)^{\ell_1/(d-1)}$  and  $\alpha_{2,j} = p_2 \ell_2^{-1} (1-p_2^{-1}y)^{\ell_2/(d-1)}$ . We get

$$\begin{aligned} & \left( \frac{b_1}{(1+p_1^{-1}y)^{\ell_1}} + \frac{b_2}{(1-p_2^{-1}y)^{\ell_2}} \right)^{1/d} \left( p_1 \ell_1^{-1} (1+p_1^{-1}y)^{\ell_1/(d-1)} + p_2 \ell_2^{-1} (1-p_2^{-1}y)^{\ell_2/(d-1)} \right)^{(d-1)/d} \\ & \geq \frac{b_1^{1/d} p_1^{(d-1)/d} \ell_1^{-(d-1)/d} (1+p_1^{-1}y)^{\ell_1/d}}{(1+p_1^{-1}y)^{\ell_1/d}} + \frac{b_2^{1/d} p_2^{(d-1)/d} \ell_2^{-(d-1)/d} (1-p_2^{-1}y)^{\ell_2/d}}{(1-p_2^{-1}y)^{\ell_2/d}} \\ & = p_1 \ell_1^{-1} (b_1 p_1^{-1} \ell_1)^{1/d} + p_2 \ell_2^{-1} (b_2 p_2^{-1} \ell_2)^{1/d}. \end{aligned}$$

Rearranging this expression yields

$$\frac{b_1}{(1+p_1^{-1}y)^{\ell_1}} + \frac{b_2}{(1-p_2^{-1}y)^{\ell_2}} \geq \frac{\left( p_1 \ell_1^{-1} (b_1 p_1^{-1} \ell_1)^{1/d} + p_2 \ell_2^{-1} (b_2 p_2^{-1} \ell_2)^{1/d} \right)^d}{(p_1 \ell_1^{-1} (1+p_1^{-1}y)^{\ell_1/(d-1)} + p_2 \ell_2^{-1} (1-p_2^{-1}y)^{\ell_2/(d-1)})^{d-1}}.$$

Thus, our final task is to bound the denominator from above. By Theorem A.1 and the fact that  $d \geq \max\{\ell_1, \ell_2\} + 1$ ,

$$\begin{aligned} & p_1 \ell_1^{-1} (1+p_1^{-1}y)^{\ell_1/(d-1)} + p_2 \ell_2^{-1} (1-p_2^{-1}y)^{\ell_2/(d-1)} \\ & \leq p_1 \ell_1^{-1} \left( 1 + \frac{\ell_1 y}{p_1(d-1)} \right) + p_2 \ell_2^{-1} \left( 1 - \frac{\ell_2 y}{p_2(d-1)} \right) = p_1 \ell_1^{-1} + p_2 \ell_2^{-1}. \end{aligned}$$

This completes the proof.  $\square$