

ZOLOTAREV POLYNOMIALS OF DEGREE 5, 6 AND 7 WITH SIMPLE CRITICAL POINTS AND THEIR MODULI SPACES

YURY KOCHETKOV

ABSTRACT. A polynomial $p \in \mathbb{C}[z]$ with three finite values is called the Zolotarev polynomial. For a class of such polynomials with the given degree, given passport and simple critical points we define a *combinatorial moduli space*. A combinatorial moduli space have the same essential properties as analytical moduli space, but much easier to construct. We study these objects for Zolotarev polynomials of degree 5, 6 and 7.

1. INTRODUCTION

Let $p \in \mathbb{C}[z]$ be a Zolotarev polynomial, i.e. a polynomial with three finite pairwise different critical values α , β and γ . Points α, β, γ are vertices of a triangle Δ in the complex plane and the inverse image $p^{-1}(\Delta)$ is a system of curvilinear triangles (ovals). Any two ovals in this system are either disjoint or have one common vertex (this common vertex is a critical point of p). A simple critical point is a common vertex of exactly two different ovals. In what follows such systems will be called cacti [2], [3].

Remark 1.1. In what follows by Galois group we understand *the geometric Galois group* [1].


Example 1.1. Let

$$p = 6 \int (x^2 + 1)(x - 1)(x - a) dx, \text{ where } a = -\frac{2}{3} + \frac{\sqrt{35}i}{3}.$$

Then

$$p(i) \approx -6.1 - 0.4i, p(-i) \approx 7 + 5.3i, p(1) = p(a) \approx -3 + 5.7i.$$

In the figure below the cactus $p^{-1}(\Delta)$ is presented.



Definition 1.1. Let p be a Zolotarev polynomial with simple critical points and let α, β, γ be its critical values. If the inverse image of α contains k critical points, the inverse image of β contains l critical points and the inverse image of γ contains m critical points, then the array (k, l, m) will be called the *passport* of p .

Thus, the passport of the polynomial in the above example is $(1, 1, 2)$.


Let us note now that the space of Zolotarev polynomials with the given passport (k, l, m) depends on one complex parameter. Indeed, let $0, x_1, \dots, x_{k-1}$ be critical points with the same critical value, $1, y_1, \dots, y_{l-1}$ be

¹Department of Applied Mathematics, Higher School of Economics, Moscow, Russia


critical points with another critical value and $z_1, \dots, z_m$ be critical points with the third critical value. Thus we have $k + l + m - 2$ parameters. But we have $k + l + m - 3$ relations: $k - 1$ relations of the type $p(x_i) = p(0)$, $l - 1$ relations of the type $p(y_i) = p(1)$ and $m - 1$ relations of the type $p(z_i) = p(z_1)$. Thus, each polynomial from this space is defined by some value of complex parameter t and the domain of t is an algebraic curve C , maybe reducible. If C' is a connected component of C , then the change of t in it induces a change of the triangle Δ .

The generic situation is as follows. There is a system L of one-dimensional lines in the domain C' . When $t \in L$, then the triangle Δ degenerates into a line segment: some critical value (α , for example) is strictly inside the line segment $[\beta, \gamma]$ bounded by two other critical values. In this case the cactus degenerates into a tree with the number of edges equal to the double polynomial degree. Thus, we have a map in C' : L is the set of edges that split C' into disjoint areas — cells. If t_1 and t_2 belong to one cell, then the corresponding cactuses are combinatorially equivalent. The passing of t from one cell into another induces a transformation of the cactus.

Example 1.2. We will denote vertices of Δ by three symbols $*$, $\circ$ and $\bullet$ and inverse images of vertices will be denoted by the same symbols. Let us consider the cactus




The initial cactus can be reconstructed from this tree in the following way: to triples $*, \circ, \bullet$ we add arcs in such way that the cyclic order of vertices $*, \circ, \bullet$ in each thus obtained ovals is the same as in the initial cactus. Naturally, the transformed cactus is constructed from the tree in the same way, only the cyclic order in each constructed oval must be opposite to the initial one.




In the generic case the border of a cell in C' consists of three arcs, because Δ can be degenerated into the segment $[\circ, \bullet]$, or into the segment $[\bullet, *]$, or into the segment $[\circ, *]$. To each arc a tree, i.e. a degenerated cactus is corresponded. These degenerated cactuses constitute the *combinatorial border* of the initial cactus and the corresponded transformed cactuses constitute the *combinatorial neighborhood* of the initial cactus.

Example 1.3. Let us find the combinatorial border and neighborhood of the cactus




We have:



Further degeneration transforms a line segment with a vertex inside (i.e. a degenerated Δ) into a line segment. Thus, the degenerated triangle Δ $\circ \rightarrow *$ can be transformed into segment $\rightarrow *$, or into segment $\circ \rightarrow *$. In the first case in a degenerated cactus we contract subsegments $\circ \rightarrow *$. And in the second case we contract in it

subsegments $\star \rightarrow \cdot$. It means that we transform a 3-colored tree (a degenerated cactus) into 2-colored tree. Thus, from the degenerated cactus



Remark 1.2. A polynomial p with exactly two finite critical values is called the Shabat polynomial. If α and β are its critical values, then the inverse image $p^{-1}[\alpha, \beta]$ is a tree with number of edges equal to the degree of p . Zolotarev polynomial became Shabat polynomials exactly for those $t \in C'$ that are intersection points of two curves from L .

Given the degree and the passport of polynomials, we construct the set of cacti. For each cactus we construct its combinatorial border — three degenerated cactuses, i.e. 3-colored trees with number of edges equal to the double degree of polynomials. Two cactuses are neighbors, if some degenerated cactus belongs to both borders. Given a degenerated cactus we construct two 2-colored trees with the number of edges equal to the degree of polynomials. Thus, we have a graph: edges are enumerated by degenerated cactuses, vertices — by 2-colored trees, that can be obtained from degenerated cactuses by contraction half of edges. A vertex and an edge are incident to each other, if the corresponding 2-colored tree can be obtained by contraction from 3-colored tree.

Definition 1.2. Let Z be some class of Zolotarev polynomials of the given degree, with given passport and with simple critical points. And let G be the graph of its degenerated cactuses and 2-colored trees. *The combinatorial moduli space* of Z is a map in some surface S , generated by the embedding G into S . Moreover, faces of this map are enumerated by cactuses and relations "being neighbors" and "being element of a border" are preserved.


Our aim is to construct combinatorial moduli spaces for some classes of Zolotarev polynomials. Main results are as follows.

- A combinatorial moduli space of Zolotarev polynomials of degree 5 and passport $(1, 1, 2)$ is a projective map with 5 faces, 8 edges and 4 vertices (Theorem 2.1).
- A combinatorial moduli space of Zolotarev polynomials of degree 6, with passport $(2, 2, 1)$ and the Galois group S_6 is a spherical map with 12 faces, 18 edges and 8 vertices (Theorem 4.1).
- A combinatorial moduli space of Zolotarev polynomials of degree 7, with passport $(2, 2, 2)$ and the Galois group A_7 consists of two components. Each component is a projective map with 7 faces, 11 edges and 5 vertices (Theorem 6.1).


2. DEGREE 5

Here we study Zolotarev polynomials of degree 5 and with the passport $(1, 1, 2)$. The vertex of Δ that is the image of two critical points will be denoted by star $\star$. The vertex following $\star$ in the counterclockwise going around Δ will be denoted by white circle $\circ$ and the third vertex will be denoted by black circle $\bullet$. Our cactus contains 5 ovals and 4 points of their contact. Two of these points are inverse images of $\star$. In what follows inverse images of the star vertex will be denoted by $\star$. The same is true for inverse images of $\circ$ and $\bullet$.


We can construct 2 cacti from 5 ovals:




In the first case positions of contact points $\star$ can be




In the second case we also have two possibilities



In result we have 5 cacti:



Let us construct combinatorial borders of our cacti. We do not mark vertices of degenerated cacti, because it is easy to recover them.



Contractions of degenerated cacti give us four trees:

α : obtained by contraction of degenerated cacti 1, 3, 4 and 7.


β : obtained by contraction of degenerated cacti 2, 3 and 5.

γ : obtained by contraction of degenerated cacti 1, 2, 4, 5, 6 and 8.

δ : obtained by contraction of degenerated cacti 7 and 8.

Now we can formulate the statement.

Theorem 2.1. *The combinatorial moduli space of Zolotarev polynomials of degree 5 and with the passport $(1, 1, 2)$ is a projective map with 5 faces, 8 edges and 4 vertices.*



3. DEGREE 5, ANALYTICAL APPROACH


For our polynomials (degree 5, passport $(1, 1, 2)$) we can easily construct analytical moduli space. Indeed, let $p = 20 \int (x^2 - 1)(x - b)(x - c) dx$. The condition $p(1) = p(-1)$ is satisfied when $5bc = -1$. Then

$$p(1) = p(-1) = 5b - \frac{1}{b}, \quad p(b) = -b^5 + 3b^3 + 2b, \quad p(c) = \frac{1}{3125b^5} - \frac{3}{125b^3} - \frac{2}{5b}.$$

The triangle Δ degenerates, if

$$\begin{aligned}
 & xy(15625x^{14} + 78125x^{12}y^2 + 140625x^{10}y^4 + 78125x^8y^6 - 78125x^6y^8 - 140625x^4y^{10} - 78125x^2y^{12} - 15625y^{14} - \\
 & - 26250x^{12} - 142500x^{10}y^2 - 333750x^8y^4 - 435000x^6y^6 - 333750x^4y^8 - 142500x^2y^{10} - 26250y^{12} + \\
 & + 5775x^{10} + 16125x^8y^2 + 10350x^6y^4 - 10350x^4y^6 - 16125x^2y^8 - 5775y^{10} + \\
 & + 4660x^8 + 20480x^6y^2 + 31512x^4y^4 + 20480x^2y^6 + 4660y^8 + \\
 & + 231x^6 + 183x^4y^2 - 183x^2y^4 - 231y^6 - 42x^4 - 60x^2y^2 - 42y^4 + x^2 - y^2) = 0
 \end{aligned}$$

where $b = x + iy$. The curve, defined by the above equation, is symmetric with respect to x and y axes. In the figure below the plot of this curve is presented.




The curve intersects x axis in the origin and in points $\pm \frac{1}{5}$ and ± 1 and y axis in points $\pm i/\sqrt{3}$. If $b = \pm 1, \pm \frac{1}{5}$, then p is a Shabat polynomial that defines the tree $\text{---}\overset{\uparrow}{\bullet}\text{---}$. If $b = \pm i/\sqrt{3}$, then p is a Shabat polynomial that defines the tree $\text{---}\overset{\uparrow}{\bullet}\text{---}$. When $b \approx \pm 0.10$ and $b \approx \pm 1.89$ (these points are marked with crosses $\times$), then $p(b) = p(c)$ and p is a Shabat polynomial that defines the tree $\text{---}\bullet\text{---}\bullet\text{---}$. We see, that an analytical approach gives us an "excessive" moduli space, while combinatorial construction describes the structure of the cell adjacency in more "compact" way.

4. DEGREE 6


Here we will construct the combinatorial moduli space for Zolotarev polynomials of degree 6 and with passport $(2, 2, 1)$. The star $*$ will denote the vertex of the triangle Δ that is the image of only one critical point. The vertex following $*$ in the counterclockwise going around Δ will be denoted by white circle $\circ$ and the last vertex will be denoted by black circle $\bullet$.

In all there are 15 cacti, but three of them




have a nontrivial symmetry group: they admit the rotation on 180° .


We will consider "non symmetrical" cacti with the Galois group S_6 . There are 12 of them:



Also we have 18 degenerated cacti:



And at last we have 8 trees — contractions of degenerated cacti:




For each degenerated cactus we: a) enumerate cacti to which border it belongs; b) enumerate trees that can be obtained from it by contraction.

- | | | | | | |
|-----------------------------|------------------------------|-------------------------------|--------------------------|----------------------------|----------------------------|
| 1 : $A, J; \lambda, \nu.$ | 2 : $A, E; \alpha, \lambda;$ | 3 : $A, H; \alpha, \nu.$ | 4 : $B, K; \mu, \nu.$ | 5 : $B, I; \alpha, \nu.$ | 6 : $B, E; \alpha, \mu.$ |
| 7 : $C, G; \beta, \lambda.$ | 8 : $C, D; \beta, \delta.$ | 9 : $C, L; \delta, \lambda.$ | 10 : $D, F; \beta, \mu.$ | 11 : $D, L; \delta, \mu.$ | 12 : $E, L; \lambda, \mu.$ |
| 13 : $F, G; \beta, \gamma.$ | 14 : $F, K; \gamma, \mu.$ | 15 : $G, J; \gamma, \lambda.$ | 16 : $H, I; \nu, \rho.$ | 17 : $H, I; \alpha, \rho.$ | 18 : $J, K; \gamma, \nu.$ |

As we have 12 cacti, 18 degenerated cacti and 8 trees, then our map is a spherical map.

Theorem 4.1. *The combinatorial moduli space of Zolotarev polynomials of degree 6, with passport $(2, 2, 1)$ and the Galois group S_6 is a spherical map presented in figure below (here A is the external face):*



5. DEGREE 6, ANALYTICAL APPROACH


Let $p = \int (x^2 - 1)(x^2 + ax + b)(x - c) dx$. If $p(1) = p(-1)$ and values of p in roots of $x^2 + ax + b$ coincide, then p is a Zolotarev polynomial with the passport $(2, 2, 1)$. The set of solutions of the system

$$\begin{cases} p(1) = p(-1) \\ \text{the remainder } r, \text{ where } p = s \cdot (x^2 + ax + b) + r \text{ has degree } 0 \end{cases}$$

has a rational parametrization:

$$a = \frac{\sqrt{3}(z^2 + 1)}{2z}, \quad b = \frac{z^2 - 1}{2}, \quad c = \frac{\sqrt{3}(z^2 + 1)}{z(5z^2 - 3)}.$$


In the figure below curves of the system L in the first quadrant are presented. Axes x and y belong to L . If a segment at axis x (or y) separates two cells, then corresponding cactuses are mirror symmetric.




Vertices have the following coordinates: $\beta \approx 0.44i$; $\gamma \approx 1.73$; $\delta = i$; $\lambda \approx 0.39 + 0.38i$; $\mu \approx 1.1 + 0.65i$; $\nu \approx 0.47$ and ≈ 1.14 . At point $w = \sqrt{3/5} c$ becomes infinite.

6. DEGREE 7


Here we will study Zolotarev polynomials of degree 7 and with passport $(2, 2, 2)$. As above, vertices of Δ in the counterclockwise going around Δ are in the order $*$, $\circ$, $\bullet$. We will mark by $*$, $\circ$, $\bullet$ only contact points of cacti. We have 20 of them:






The Galois group of D, E, G, I, K, N is $PSL_2(7)$ of order 168. Other 14 cacti have group A_7 as the Galois group. We will construct the combinatorial moduli space exactly for these 14 cacti.

Now let us enumerate degenerated cacti:



and enumerate trees:




As in the case of degree 6, we for each degenerated cactus will: a) enumerate cacti to which border it belongs; b) enumerate trees that can be obtained from it by contraction.

- | | | | | | |
|----------------------------------|----------------------------------|--------------------------------|--------------------------------|---------------------------------|--------------------------------|
| 1 : A, L ; α, λ . | 2 : A, Q ; λ, μ . | 3 : A, O ; α, μ . | 4 : B, F ; β, μ . | 5 : B, S ; δ, μ . | 6 : B, C ; β, δ . |
| 7 : C, R ; δ, λ . | 8 : C, H ; β, λ . | 9 : F, T ; β, δ . | 10 : F, S ; μ, ρ . | 11 : H, R ; λ, ρ . | 12 : H, T ; β, ρ . |
| 13 : J, P ; γ, ν . | 14 : J, M ; α, γ . | 15 : J, O ; α, ν . | 16 : L, Q ; λ, ν . | 17 : L, M ; α, ν . | 18 : M, P ; γ, ν . |
| 19 : O, Q ; μ, ν . | 20 : P, P ; γ, γ . | 21 : R, S ; δ, ρ . | 22 : T, T ; ρ, ρ . | | |

Theorem 6.1. *Combinatorial moduli space of Zolotarev polynomials of degree 7 with passport $(2, 2, 2)$ and the Galois A_7 has two components. Each component is a projective map of 7 faces, 11 edges and 5 vertices.*

The first component: faces A, J, L, M, O, P, Q ; edges 1, 2, 3, 13, 14, 15, 16, 17, 18, 19, 20; vertices $\alpha, \gamma, \lambda, \mu, \nu$. The second component: faces B, C, F, H, R, S, T ; edges 4, 5, 6, 7, 8, 9, 10, 11, 12, 21, 22; vertices $\beta, \delta, \lambda, \mu, \rho$. In figure

below we present maps of both components.



Remark 6.1. Let $p = \int (x^2 - 1)(x^2 + ax + b)(x^2 + cx + d) dx$. The elimination of variables c and d from the system

$$p(1) = p(-1)$$

the remainder r , where $p = s \cdot (x^2 + ax + b) + r$ has degree 0

the remainder g , where $p = h \cdot (x^2 + cx + d) + g$ has degree 0

gives us the polynomial in a and b of degree 20. This polynomial factorizes over $\mathbb{Q}$ in two factors of degrees 6 and 14, respectively. The factor of degree 6 describes polynomials with the Galois group $PSL_2(7)$. The factor of degree 14 factorizes over $\mathbb{Q}(\sqrt{21})$ in two factors of degree 7. Each of them defines a curve of genus 1.

REFERENCES

- [1] V.B. Alekseev, *Abel's Theorem in Problems and Solutions*, Kluwer Academic Publishers, 2004.
- [2] Goulden I.P. and Jackson D.M., The combinatorial relationship between trees, cacti and certain connection coefficients for the symmetric group, *European J. Combin.*, 1992, V.13, p. 357-365.
- [3] S.K. Lando, A.K. Zvonkin, *Graphs on surfaces and their applications*, Springer, 2004.

Email address: yukochetkov@hse.ru, yuyukochetkov@gmail.com