

Sombor-index-like invariants of some graphs

Nima Ghanbari* Saeid Alikhani

December 9, 2022

Department of Informatics, University of Bergen, P.O. Box 7803, 5020 Bergen,
Norway

Department of Mathematical Sciences, Yazd University, 89195-741, Yazd, Iran
Nima.ghanbari@uib.no, alikhani@yazd.ac.ir

Abstract

The Sombor index (SO) is a vertex-degree-based graph invariant, defined as the sum over all pairs of adjacent vertices of $\sqrt{d_i^2 + d_j^2}$, where d_i is the degree of the i -th vertex. It has been conceived using geometric considerations. Recently, a series of new SO-like degree-based graph invariants (denoted by SO_1, SO_2, \dots, SO_6) is taken into consideration, when the geometric background of several classical topological indices (Zagreb, Albertson) has considered. In this paper, we compute and study these new indices for some graphs, cactus chains and polymers.

Keywords: sombor index, degree, vertex-degree-based graph invariant, polymer.

AMS Subj. Class.: 05C07; 05C09.

1 Introduction

Let $G = (V, E)$ be a finite, connected and simple graph. We denote the degree of a vertex v in G by d_v . A topological index of G is a real number related to G . It does not depend on the labeling or pictorial representation of a graph. In the mathematical and chemical literature, several dozens of vertex-degree-based (VDB) graph invariants such as the first and second Zagreb index, the Albertson index and the Sombor index have been introduced and extensively studied [10, 11, 19]. Their general formula is

$$TI(G) = \sum_{ij \in E} F(d_i, d_j),$$

where $F(x, y)$ is some function with the property $F(x, y) = F(y, x)$.

*Corresponding author

The Sombor index, is defined as

$$SO(G) = \sum_{uv \in E(G)} \sqrt{d_u^2 + d_v^2}.$$

This index soon attracted much attention, and its numerous mathematical [12, 13, 15, 16, 18, 20] and chemical [1, 2, 4, 5, 8, 14, 17, 18] applications have been established. Recently, Gutman [9] intended to point out the great variety of Sombor-index-like VDB graph invariants that can be constructed by means of geometric arguments and so generated a number of new Sombor-index-like VDB invariants, denoted below by SO_1, SO_2, \dots, SO_6 .

$$SO_1(G) = \frac{1}{2} \sum_{uv \in E(G)} |d_u^2 - d_v^2|.$$

$$SO_2(G) = \sum_{uv \in E(G)} \left| \frac{d_u^2 - d_v^2}{d_u^2 + d_v^2} \right|.$$

$$SO_3(G) = \sum_{uv \in E(G)} \sqrt{2} \frac{d_u^2 + d_v^2}{d_u + d_v} \pi.$$

$$SO_4(G) = \frac{1}{2} \sum_{uv \in E(G)} \left(\frac{d_u^2 + d_v^2}{d_u + d_v} \right)^2 \pi.$$

$$SO_5(G) = \sum_{uv \in E(G)} \frac{2|d_u^2 - d_v^2|}{\sqrt{2} + 2\sqrt{d_u^2 + d_v^2}} \pi.$$

$$SO_6(G) = \sum_{uv \in E(G)} \left[\frac{d_u^2 - d_v^2}{\sqrt{2} + 2\sqrt{d_u^2 + d_v^2}} \right]^2 \pi.$$

Polymer graphs can be decomposed into subgraphs that we call monomer units. Blocks of graphs are particular examples of monomer units, but a monomer unit may consist of several blocks.

Let G be a connected graph constructed from pairwise disjoint connected graphs G_1, \dots, G_k as follows. Select a vertex of G_1 , a vertex of G_2 , and identify these two vertices. Then continue in this manner inductively. Note that the graph G constructed in this way has a tree-like structure, the G_i 's being its building stones (see Figure 1). Usually we say that G is a polymer graph, obtained by point-attaching from G_1, \dots, G_k and that G_i 's are the monomer units of G . A particular case of this construction is the decomposition of a connected graph into blocks (see [3, 6, 7]).

In Section 2, We compute the Sombor-like degree-based indices for certain graphs such as path, cycle, complete bipartite, friendship, ladder, book, Dutch-windmill and cactus chains. In Section 3, we consider polymers and study some of $SO_i(G)$ based their monomer units G_i . In Section 4, we obtain some relations between indices $SO_i(G)$ ($i = 2, 3, 4, 5, 6$) and index $SO_1(G)$.




Figure 1: A polymer graph with monomer units G_1, \dots, G_k .

2 Sombor-like degree-based indices of certain graphs

In this section, we compute the Sombor-index-like invariants indices for certain graphs, such as paths, friendship graph, grid graphs and cactus chains. By the definitions of $SO_1(G)$, $SO_2(G)$, $SO_5(G)$ and $SO_6(G)$, these indices are 0 for every regular graphs.

2.1 Sombor-like degree-based indices of specific graphs

We begin with the following theorem:

Theorem 2.1 (i) For every $n \in \mathbb{N} - \{1, 2\}$, $SO_1(P_n) = 3$.

(ii) For every $m, n \in \mathbb{N}$, $SO_1(K_{1,n}) = \frac{n(n-1)(n+1)}{2}$ and $SO_1(K_{m,n}) = \frac{mn(m-n)(m+n)}{2}$, if $m \geq n$.

(iii) For the wheel graph $W_{n+1} = C_n \vee K_1$, $SO_1(W_{n+1}) = \frac{n(n-3)(n+3)}{2}$.

(iv) For the ladder graph $L_n = P_n \square K_2$, ($n \geq 3$), $SO_1(L_n) = 10$.

(v) For the friendship graph $F_n = K_1 \vee nK_2$, $SO_1(F_n) = 4n(n-1)(n+1)$.

(vi) For the book graph $B_n = K_{1,n} \square K_2$, and for $n \geq 3$, $SO_1(B_n) = n(n+3)(n-1)$.

(vii) For the Dutch windmill graph $D_n^{(m)}$, and for every $n \geq 3$ and $m \geq 2$,

$$SO_1(D_n^{(m)}) = 4m(m-1)(m+1).$$

Proof. The proof of all parts are easy and similar. For instance we state the proof of Part (i).

- (i) There are two edges with endpoints of degree 1 and 2. Also there are $n-3$ edges with endpoints of degree 2. Therefore we have the result. \square

Similarly, we can compute SO_2, \dots, SO_6 for these graphs. Table 1 and 2 show the exact values for these Sombor-like degree-based indices.

G	$SO_2(G)$	$SO_3(G)$	$SO_4(G)$
P_n	$\frac{6}{5}$	$\frac{10\sqrt{2}\pi}{3} + (n-3)\sqrt{2}(2\pi)$	$\frac{25\pi}{9} + (n-3)(2\pi)$
$K_{1,n}$	$\frac{n(n^2-1)}{n^2+1}$	$\frac{\sqrt{2}n\pi(n^2+1)}{n+1}$	$\frac{n\pi}{2} \left(\frac{n^2+1}{n+1}\right)^2$
$K_{m,n}$	$\frac{mn(m^2-n^2)}{m^2+n^2}$	$\frac{\sqrt{2}nm\pi(m^2+n^2)}{m+n}$	$\frac{nm\pi}{2} \left(\frac{n^2+m^2}{n+m}\right)^2$
W_n	$\frac{(n-1)(n^2-2n-8)}{n^2-2n+10}$	$\sqrt{2}\pi \left(\frac{(n-1)(n^2-2n+10)}{n+2} + 3n-3\right)$	$\frac{\pi}{2} \left((n-1) \left(\frac{n^2-2n+10}{n+2}\right)^2 + 9n-9\right)$
L_n	$\frac{20}{13}$	$\frac{\sqrt{2}\pi}{5}(45n-48)$	$\frac{\pi}{50}(675n-924)$
F_n	$\frac{2n^3-2n}{n^2+1}$	$\sqrt{2}\pi \left(\frac{4n^3+2n^2+6n}{n+1}\right)$	$2n\pi \left(\frac{2n^4+5n^2+2n+3}{n^2+2n+1}\right)$
B_n	$\frac{2n^3+4n^2-6n}{n^2+2n+5}$	$\sqrt{2}\pi \left(3n+1+2n \left(\frac{n^2+2n+5}{n+3}\right)\right)$	$\frac{\pi}{2} \left(4n+(n+1)^2+2n \left(\frac{n^2+2n+5}{n+3}\right)^2\right)$
$D_n^{(m)}$	$\frac{2m(m^2-1)}{m^2+1}$	$\sqrt{2}\pi \left(2m(n-2)+4m \left(\frac{m^2+1}{m+1}\right)\right)$	$\frac{\pi}{2} \left(4m(n-2)+8m \left(\frac{m^2+1}{m+1}\right)^2\right)$

Table 1. The exact values of $SO_i(G)$ ($2 \leq i \leq 4$), for $m \geq n \geq 3$.

G	$SO_5(G)$	$SO_6(G)$
P_n	$\frac{12\pi}{\sqrt{2+2\sqrt{5}}}$	$\frac{18\pi}{(\sqrt{2+2\sqrt{5}})^2}$
$K_{1,n}$	$\frac{2n\pi(n^2-1)}{\sqrt{2+2\sqrt{n^2+1}}}$	$n\pi \left(\frac{n^2-1}{\sqrt{2+2\sqrt{n^2+1}}}\right)^2$
$K_{m,n}$	$\frac{2nm\pi(m^2-n^2)}{\sqrt{2+2\sqrt{n^2+m^2}}}$	$nm\pi \left(\frac{m^2-n^2}{\sqrt{2+2\sqrt{n^2+m^2}}}\right)^2$
W_n	$2(n-1)\pi \left(\frac{n^2-2n-8}{\sqrt{2+2\sqrt{n^2-2n+10}}}\right)$	$\pi(n-1) \left(\frac{n^2-2n-8}{\sqrt{2+2\sqrt{n^2-2n+10}}}\right)^2$
L_n	$\frac{40\pi}{\sqrt{2+2\sqrt{13}}}$	$\frac{100\pi}{(\sqrt{2+2\sqrt{13}})^2}$
F_n	$4n\pi \left(\frac{4n^2-4}{\sqrt{2+2\sqrt{4n^2+4}}}\right)$	$2n\pi \left(\frac{4n^2-4}{\sqrt{2+2\sqrt{4n^2+4}}}\right)^2$
B_n	$4n\pi \left(\frac{n^2+2n-3}{\sqrt{2+2\sqrt{n^2+2n+5}}}\right)$	$2n\pi \left(\frac{n^2+2n-3}{\sqrt{2+2\sqrt{n^2+2n+5}}}\right)^2$
$D_n^{(m)}$	$16m\pi \left(\frac{m^2-1}{\sqrt{2+2\sqrt{4m^2+4}}}\right)$	$32m\pi \left(\frac{m^2-1}{\sqrt{2+2\sqrt{4m^2+4}}}\right)^2$

Table 2. The exact values of $SO_5(G)$ and $SO_6(G)$, for $m \geq n \geq 3$.

Here, we consider the grid graph $(P_n \square P_m)$, and compute the Sombor-like degree-based indices for it.

Theorem 2.2 *Let $P_m \square P_n$ be the grid graph (Figure 2). For every $n \geq 6$ and $m \geq 6$,*

$$SO_1(P_m \square P_n) = 14m + 14n - 16,$$

$$SO_2(P_m \square P_n) = \frac{40}{13} + \frac{14m + 14n - 56}{25},$$

$$SO_3(P_m \square P_n) = \sqrt{2}\pi \left(\frac{104}{5} + \frac{50n + 50m - 200}{7}\right),$$




Figure 2: Grid graph $P_m \square P_n$




Figure 3: Chain triangular cactus T_n

$$SO_4(P_m \square P_n) = \frac{\pi}{2} \left(\frac{1352}{25} + \frac{1250m + 1250n - 5000}{49} \right),$$

$$SO_5(P_m \square P_n) = 2\pi \left(\frac{40}{\sqrt{2} + 2\sqrt{13}} + \frac{14m + 14n - 56}{\sqrt{2} + 10} \right),$$

$$SO_6(P_m \square P_n) = \pi \left(\frac{200}{(\sqrt{2} + 2\sqrt{13})^2} + \frac{98m + 98n - 392}{(\sqrt{2} + 10)^2} \right).$$

Proof. There are eight edges with endpoints of degree 2 and 3 and there are $m + n - 4$ edges with endpoints of degree 3. Also there are $2m + 2n - 8$ edges with endpoints of degree 3 and 4 and there are $2nm - 5n - 5m - 12$ edges with endpoints of degree 4. Therefore we have the result. \square

2.2 Sombor-like degree-based indices of cactus chains

In this subsection, we consider a class of simple linear polymers called cactus chains. Cactus graphs were first known as Husimi tree, they appeared in the scientific literature some sixty years ago in papers by Husimi and Riddell concerned with cluster integrals in the theory of condensation in statistical mechanics [12, 20, 18]. We refer the reader to papers [8, 16] for some aspects of parameters of cactus graphs.

Theorem 2.3 (i) Let T_n be the chain triangular graph (See Figure 3) of order n . Then for every $n \geq 2$, $SO_1(T_n) = 12n$.




Figure 4: Para-chain square cactus Q_n




Figure 5: Para-chain square cactus O_n and Ortho-chain graph O_n^h

- (ii) Let Q_n be the para-chain square cactus graph (See Figure 4) of order n . Then for every $n \geq 2$, $SO_1(Q_n) = 24n - 24$.
- (iii) Let O_n be the para-chain square cactus (See Figure 5) graph of order n . Then for every $n \geq 2$, $SO_1(O_n) = 12n$.
- (iv) Let O_n^h be the Ortho-chain graph (See Figure 5) of order n . Then for every $n \geq 2$, $SO_1(O_n^h) = 12n$.
- (v) Let L_n be the para-chain hexagonal cactus graph (See Figure 6) of order n . Then for every $n \geq 2$, $SO_1(L_n) = 24n - 24$.
- (vi) Let M_n be the Meta-chain hexagonal cactus graph (See Figure 6) of order n . Then for every $n \geq 2$, $SO_1(M_n) = 24n - 24$.

Proof.

- (i) There are two edges with endpoints of degree 2. Also there are $2n$ edges with endpoints of degree 2 and 4 and there are $n - 2$ edges with endpoints of degree 4. Therefore we have the result.
- (ii) There are four edges with endpoints of degree 2. Also there are $4n - 4$ edges with endpoints of degree 2 and 4. Therefore the result follows.




Figure 6: Para-chain L_n and Meta-chain M_n

- (iii) There are $n + 2$ edges with endpoints of degree 2. Also there are $2n$ edges with endpoints of degree 2 and 4 and there are $n - 2$ edges with endpoints of degree 4. Therefore we have the result.
- (iv) There are $3n + 2$ edges with endpoints of degree 2. Also there are $2n$ edges with endpoints of degree 2 and 4 and there are $n - 2$ edges with endpoints of degree 4. Therefore the result follows.
- (v) There are $2n + 4$ edges with endpoints of degree 2. Also there are $4n - 4$ edges with endpoints of degree 2 and 4. Therefore we have the result.
- (vi) There are $2n + 4$ edges with endpoints of degree 2. Also there are $4n - 4$ edges with endpoints of degree 2 and 4. Therefore the result follows. \square

Similarly, we can compute SO_2, \dots, SO_6 for cactus chains. Table 3 shows the exact value for these Sombor-like degree-based indices.

G	$SO_2(G)$	$SO_3(G)$	$SO_4(G)$	$SO_5(G)$	$SO_6(G)$
T_n	$\frac{6n}{5}$	$3\sqrt{2}\pi \left(\frac{44n-36}{3}\right)$	$\frac{\pi}{2} \left(\frac{776n-1080}{9}\right)$	$\frac{48n\pi}{\sqrt{2}+2\sqrt{20}}$	$\frac{288n\pi}{(\sqrt{2}+2\sqrt{20})^2}$
Q_n	$\frac{12n-12}{5}$	$\sqrt{2}\pi \left(\frac{40n-16}{3}\right)$	$\frac{\pi}{2} \left(\frac{400n-256}{9}\right)$	$\frac{24\pi(4n-4)}{\sqrt{2}+2\sqrt{20}}$	$\frac{144\pi(4n-4)}{(\sqrt{2}+2\sqrt{20})^2}$
O_n	$\frac{6n}{5}$	$\sqrt{2}\pi \left(\frac{38n-12}{3}\right)$	$\frac{\pi}{2} \left(\frac{380n-216}{9}\right)$	$\frac{48n\pi}{\sqrt{2}+2\sqrt{20}}$	$\frac{288n\pi}{(\sqrt{2}+2\sqrt{20})^2}$
O_n^h	$\frac{6n}{5}$	$\sqrt{2}\pi \left(\frac{50n-12}{3}\right)$	$\frac{\pi}{2} \left(\frac{452n-216}{9}\right)$	$\frac{48n\pi}{\sqrt{2}+2\sqrt{20}}$	$\frac{288n\pi}{(\sqrt{2}+2\sqrt{20})^2}$
L_n	$\frac{12n-12}{5}$	$\sqrt{2}\pi \left(\frac{52n-26}{3}\right)$	$\frac{\pi}{2} \left(\frac{472n-256}{9}\right)$	$\frac{24\pi(4n-4)}{\sqrt{2}+2\sqrt{20}}$	$\frac{144\pi(4n-4)}{(\sqrt{2}+2\sqrt{20})^2}$
M_n	$\frac{12n-12}{5}$	$\sqrt{2}\pi \left(\frac{52n-26}{3}\right)$	$\frac{\pi}{2} \left(\frac{472n-256}{9}\right)$	$\frac{24\pi(4n-4)}{\sqrt{2}+2\sqrt{20}}$	$\frac{144\pi(4n-4)}{(\sqrt{2}+2\sqrt{20})^2}$

Table 3. The exact values of $SO_i(G)$ ($2 \leq i \leq 6$), for $n \geq 2$.

Corollary 2.4 *Meta-chain hexagonal cactus graphs and para-chain hexagonal cactus graphs of the same order, have the same Sombor-like degree-based indices.*

3 Study of $SO_1(G)$ for polymers and their monomer's unit

In this section, we study the index $SO_1(G) = \frac{1}{2} \sum_{uv \in E(G)} |d_u^2 - d_v^2|$, where G is a polymer graph. We begin with the following proposition which gives an upper bound for $SO_1(G - e)$:

Proposition 3.1 *Let $G = (V, E)$ be a non-regular graph and $e = uv \in E$. Also let d_w , Δ and δ , be the degree of vertex w , maximum degree and minimum degree of vertices in G , respectively. Then,*

$$SO_1(G) > SO_1(G - e) + \frac{1}{2}(\delta^2 - \Delta^2).$$




Figure 7: Link of n graphs G_1, G_2, \dots, G_n

Proof. First we remove edge e and find $SO_1(G - e)$. Obviously, by adding edge e to $G - e$ and $\frac{1}{2}|d_u^2 - d_v^2|$ to $SO_1(G - e)$, the value $SO_1(G)$ is greater than that. So we have

$$SO_1(G) > SO_1(G - e) + \frac{1}{2}|d_u^2 - d_v^2| \geq SO_1(G - e) + \frac{1}{2}(d_u^2 - d_v^2),$$

and therefore we have the result. \square

Here, we study the Sombor-like degree-based indices of some polymer graphs.

Theorem 3.2 *Let G be a polymer graph with composed of monomers $\{G_i\}_{i=1}^k$ with respect to the vertices $\{x_i, y_i\}_{i=1}^k$, and Δ_i and δ_i be the maximum degree and minimum degree of vertices in G_i , respectively. Let $G = L(G_1, \dots, G_n)$ be the link of graphs (see Figure 7). If G and $L(G_1, \dots, G_i)$ are non-regular for $i \geq 2$, then,*

$$SO_1(G) > \sum_{i=1}^k SO_1(G_i) + \frac{1}{2} \sum_{i=1}^{k-1} \delta_i^2 - \frac{1}{2} \sum_{i=2}^k \Delta_i^2.$$

Proof. First we remove edge y_1x_2 (Figure 7). Similar to the proof of Proposition 3.1, we have

$$SO_1(G) > SO_1(G - y_1x_2) + \frac{1}{2}(d_{y_1}^2 - d_{x_2}^2).$$

Let G' be the link graph related to graphs $\{G_i\}_{i=2}^k$ with respect to the vertices $\{x_i, y_i\}_{i=2}^k$. Then we have,

$$SO_1(G - y_1x_2) = SO_1(G_1) + SO_1(G'),$$

and therefore,

$$\begin{aligned} SO_1(G) &> SO_1(G_1) + SO_1(G') + \frac{1}{2}(d_{y_1}^2 - d_{x_2}^2) \\ &\geq SO_1(G_1) + SO_1(G') + \frac{1}{2}(\delta_1^2 - \Delta_2^2). \end{aligned}$$

By continuing this process, we have the result. \square

As an immediate result of theorem 3.2, we have:

Corollary 3.3 *Let G be a polymer graph with composed of monomers $\{G_i\}_{i=1}^k$ with respect to the vertices $\{x_i, y_i\}_{i=1}^k$. If G , $L(G_1, \dots, G_i)$ are non-regular for $i \geq 2$, and Δ and δ are the maximum degree and the minimum degree of vertices in G . Then,*

$$SO_1(G) > \frac{k-1}{2} (\delta^2 - \Delta^2) + \sum_{i=1}^k SO_1(G_i).$$

4 Relations between $SO_i(G)$ ($i = 2, 3, 4, 5, 6$) and $SO_1(G)$

In this section, we obtain bounds for $SO_i(G)$ ($i = 2, 3, 4, 5, 6$) based $SO_1(G)$. We begin with the following proposition which gives a lower and upper bounds for $SO_2(G)$ based $SO_1(G)$.

Proposition 4.1 *If G is a graph with minimum degree δ and maximum degree Δ , then*

$$\frac{SO_1(G)}{\Delta^2} \leq SO_2(G) \leq \frac{SO_1(G)}{\delta^2}.$$

Proof. First we find the lower bound. We have

$$\begin{aligned} SO_2(G) &= \sum_{uv \in E(G)} \left| \frac{d_u^2 - d_v^2}{d_u^2 + d_v^2} \right| \geq \sum_{uv \in E(G)} \frac{|d_u^2 - d_v^2|}{2\Delta^2} \\ &\geq \frac{1}{\Delta^2} \left(\frac{1}{2} \sum_{uv \in E(G)} |d_u^2 - d_v^2| \right) = \frac{SO_1(G)}{\Delta^2}. \end{aligned}$$

Similarly, we have the result for the upper bound. □

The following proposition gives an upper bound for $SO_2(G - e)$:

Theorem 4.2 *Let $G = (V, E)$ be a non-regular graph and $e = uv \in E$. Also let Δ and δ , be the maximum degree and minimum degree of vertices in G , respectively. Then,*

$$SO_2(G) > \frac{\delta^2}{\Delta^2} \left(SO_2(G - e) + \frac{1}{2} - \frac{\Delta^2}{2\delta^2} \right).$$

Proof. By using Propositions 3.1 and 4.1, we have

$$\begin{aligned} SO_2(G) &\geq \frac{SO_1(G)}{\Delta^2} \\ &> \frac{1}{\Delta^2} \left(SO_1(G - e) + \frac{1}{2}(\delta^2 - \Delta^2) \right) \\ &\geq \frac{1}{\Delta^2} \left(\delta^2 SO_2(G - e) + \frac{1}{2}(\delta^2 - \Delta^2) \right), \end{aligned}$$

and therefore we have the result. □

The following proposition gives a lower and upper bounds for $SO_3(G)$ based $SO_1(G)$, δ and Δ .

Proposition 4.3 *Let $G = (V, E)$ be a graph with $|E| = m$ and minimum degree δ and maximum degree Δ . Then*

$$\sqrt{2\pi} \left(\frac{SO_1(G) + m\delta^2}{\Delta} \right) \leq SO_3(G) \leq \sqrt{2\pi} \left(\frac{SO_1(G) + m\Delta^2}{\delta} \right).$$

Proof. First we obtain the upper bound. We have

$$\begin{aligned}
SO_3(G) &= \sum_{uv \in E(G)} \sqrt{2} \frac{d_u^2 + d_v^2}{d_u + d_v} \pi \\
&\leq \sqrt{2} \pi \left(\sum_{uv \in E(G)} \frac{|d_u^2 - d_v^2| + 2\Delta^2}{2\delta} \right) \\
&\leq \sqrt{2} \pi \left(\frac{1}{\delta} \left(\frac{1}{2} \sum_{uv \in E(G)} |d_u^2 - d_v^2| \right) + \frac{m\Delta^2}{\delta} \right) \\
&= \sqrt{2} \pi \left(\frac{SO_1(G) + m\Delta^2}{\delta} \right).
\end{aligned}$$

Similarly, we have the result for the lower bound. \square

By similar argument as Theorem 4.2, by using Propositions 3.1 and 4.3, we have

Theorem 4.4 *Let $G = (V, E)$ be a graph and $e = uv \in E$ and $|E| = m$. Also let Δ and δ , be the maximum degree and minimum degree of vertices in G , respectively. Then,*

$$SO_3(G) > \frac{\delta}{\Delta} (SO_3(G - e)) + \frac{(2m + 1)\pi}{\sqrt{2}\Delta} (\delta^2 - \Delta^2).$$

Proposition 4.5 *Let $G = (V, E)$ be a graph with $|E| = m$ and minimum degree δ and maximum degree Δ . Then*

$$\frac{\pi\delta^2}{2\Delta^2} (m\delta^2 + SO_1(G)) \leq SO_4(G) \leq \frac{\pi\Delta^2}{2\delta^2} (m\Delta^2 + SO_1(G)).$$

Proof. First we obtain the upper bound. We have

$$\begin{aligned}
SO_4(G) &= \frac{1}{2} \sum_{uv \in E(G)} \left(\frac{d_u^2 + d_v^2}{d_u + d_v} \right)^2 \pi \\
&= \frac{\pi}{2} \sum_{uv \in E(G)} \left(\frac{(d_u^2 + d_v^2)(d_u^2 + d_v^2)}{(d_u + d_v)^2} \right) \\
&\leq \frac{\pi}{2} \sum_{uv \in E(G)} \left(\frac{(d_u^2 + d_v^2)(2\Delta^2)}{(2\delta)^2} \right) \\
&\leq \frac{\pi\Delta^2}{2\delta^2} \left(\frac{1}{2} \sum_{uv \in E(G)} 2\Delta^2 + |d_u^2 - d_v^2| \right) \\
&= \frac{\pi\Delta^2}{2\delta^2} (m\Delta^2 + SO_1(G)).
\end{aligned}$$

Similarly, we have the result for the lower bound. \square

By similar argument as Theorem 4.2, by using Propositions 3.1 and 4.3, we have

Theorem 4.6 *Let $G = (V, E)$ be a graph and $e = uv \in E$ and $|E| = m$. Also let Δ and δ , be the maximum degree and minimum degree of vertices in G , respectively. Then,*

$$SO_4(G) > SO_4(G - e) + \frac{(2m + 1)\pi\delta^2(\delta^2 - \Delta^2)}{2\Delta^2}.$$

By similar argument as Proposition 4.1, we have:

Proposition 4.7 *Let G be a graph with minimum degree δ and maximum degree Δ . Then*

$$\frac{2\sqrt{2}\pi SO_1(G)}{2\Delta + 1} \leq SO_5(G) \leq \frac{2\sqrt{2}\pi SO_1(G)}{2\delta + 1}.$$

By similar argument as Theorem 4.2, by using Propositions 3.1 and 4.7, we have

Theorem 4.8 *Let $G = (V, E)$ be a non-regular graph and $e = uv \in E$. Also let Δ and δ , be the maximum degree and minimum degree of vertices in G , respectively. Then,*

$$SO_5(G) > SO_5(G - e) + \frac{\sqrt{2}\pi}{2\Delta + 1} (\delta^2 - \Delta^2).$$

Proposition 4.9 *Let $G = (V, E)$ be a graph with $|E| = m$ and minimum degree δ and maximum degree Δ . Then*

$$SO_6(G) \leq \frac{2\pi(\Delta^2 - \delta^2)SO_1(G)}{(\sqrt{2} + 2\delta\sqrt{2})^2}.$$

Proof. We have

$$\begin{aligned} SO_6(G) &= \pi \sum_{uv \in E(G)} \left(\frac{d_u^2 - d_v^2}{\sqrt{2} + 2\sqrt{d_u^2 + d_v^2}} \right)^2 \\ &\leq \pi \sum_{uv \in E(G)} \frac{|d_u^2 - d_v^2|(\Delta^2 - \delta^2)}{(\sqrt{2} + 2\delta\sqrt{2})^2} \\ &= \frac{2\pi(\Delta^2 - \delta^2)}{(\sqrt{2} + 2\delta\sqrt{2})^2} \left(\frac{1}{2} \sum_{uv \in E(G)} |d_u^2 - d_v^2| \right) \\ &= \frac{2\pi(\Delta^2 - \delta^2)SO_1(G)}{(\sqrt{2} + 2\delta\sqrt{2})^2}. \end{aligned}$$

□

5 Acknowledgements

The first author would like to thank the Research Council of Norway and Department of Informatics, University of Bergen for their support.

References

- [1] S. Alikhani, N. Ghanbari, Sombor index of polymers, *MATCH Commun. Math. Comput. Chem.*, 86 (2021) 715-728.
- [2] S. Amin, A. Ur Rehman Virk, M.A. Rehman, N.A. Shah, Analysis of dendrimer generation by Sombor indices, *J. Chem.*, (2021) Article No. 9930645.
- [3] D. Emeric, S. Klavžar, Computing Hosoya polynomials of graphs from primary subgraphs, *MATCH Commun. Math. Comput. Chem.* 70 (2013) 627–644.
- [4] X. Fang, L. You, H. Liu, The expected values of Sombor indices in random hexagonal chains, phenylene chains and Sombor indices of some chemical graphs, *Int. J. Quantum Chem.*, 121(17) (2021) Article No. e26740.
- [5] N. Ghanbari, On the Sombor characteristic polynomial and Sombor energy of a graph, *Comp. Appl. Math.*, 41 (2022) 242.
- [6] N. Ghanbari, On the Graovac-Ghorbani and atom-bond connectivity indices of graphs from primary subgraphs, *Iranian J. Math. Chem.*, 13(1) (2022) 45-72.
- [7] N. Ghanbari, S. Alikhani, Mostar index and edge Mostar index of polymers, *Comp. Appl. Math.*, 40, 260 (2021).
- [8] N. Ghanbari, S. Alikhani, Sombor index of certain graphs, *Iranian J. Math. Chem.*, 12(1) (2021) 27-37.
- [9] I. Gutman, Sombor indices-back to geometry, *Open J. Disc. Appl. Math.*, 5(2) (2022) 1-5.
- [10] I. Gutman, Geometric approach to degree-based topological indices: Sombor indices. *MATCH Commun. Math. Comput. Chem.*, 86 (2021) 1116.
- [11] I. Gutman, Degreebased topological indices, *Croat. Chem. Acta*, 86 (2013) 351361.
- [12] I. Gutman, TEMO theorem for Sombor index, *Open J. Disc. Appl. Math.*, 5(1) (2022) 2528.
- [13] B. Horoldagva, C. Xu, On Sombor index of graphs, *MATCH Commun. Math. Comput. Chem.*, 86 (2021) 793713.
- [14] H. Liu, H. Chen, Q. Xiao, X. Fang, Z. Tang, More on Sombor indices of chemical graphs and their applications to the boiling point of benzenoid hydrocarbons, *Int. J. Quantum Chem.*, 121(17) (2021) Article no. e26689.
- [15] W. Ning, Y. Song, K. Wang, More on Sombor index of graphs, *Mathematics*, 10(3) (2022) Article No. 301.
- [16] J. Rada, J.M. Rodríguez, J.M. Sigarreta, General properties on Sombor indices, *Discret. Appl. Math.*, 299 (2021) 87-97.

- [17] I. Redžepović, Chemical applicability of Sombor indices, *J. Serbian Chem. Soc.*, 86 (2021) 445-457.
- [18] Y. Shang, Sombor index and degree-related properties of simplicial networks, *Appl. Math. Comput.*, 419 (2022) Article No. 126881.
- [19] R. Todeschini, V. Consonni, *Molecular Descriptors for Chemoinformatics*. Weinheim: (2009) WileyVCH.
- [20] S.O. Ünal, Sombor index over the tensor and Cartesian product of monogenic semigroup graphs, *Symmetry*, 14(5) (2022) Article No. 1071.