

# Partial-wave expansion of $\Lambda NN$ three-baryon interactions in chiral effective field theory

M. Kohno,<sup>1</sup> H. Kamada,<sup>2</sup> and K. Miyagawa<sup>1</sup>

<sup>1</sup>Research Center for Nuclear Physics, Osaka University, Ibaraki 567-0047, Japan

<sup>2</sup>Department of Physics, Faculty of Engineering,  
Kyushu Institute of Technology, Kitakyushu 804-8550, Japan

An expression of partial wave expansion of three-baryon interactions in chiral effective field theory is presented. The derivation follows the method by Hebeler *et al.* [Phys. Rev. C **91**, 044001 (2015)], but the final expression is more general. That is, a systematic treatment of the higher-rank spin-momentum structure of the interaction becomes possible. Using the derived formula, a  $\Lambda$ -deuteron folding potential is evaluated. This information is valuable for inferring the possible contribution of the  $\Lambda NN$  three-baryon forces to the hypertriton as the basis of further studies by sophisticated Faddeev calculations. A microscopic understanding of  $\Lambda NN$  three-baryon forces together with two-body  $\Lambda N$  interactions is essential for the description of hypernuclei and neutron-star matter.

## I. INTRODUCTION

Any description of two-body baryon-baryon interactions in which various degrees of freedom are eliminated or frozen is effective. When the interactions are applied in many-body systems, the appearance of three-body interactions is inevitable as induced interactions. The important role of three-body forces (3BFs) has been expected in nuclear physics, in particular, to properly reproduce nuclear binding and saturation properties [1–3]. The recent development of the construction of baryon-baryon interactions in chiral effective field theory (ChEFT) [4, 5] provides a systematic way to introduce three-body (and higher-body) forces in a power-counting scheme and therefore quantifies the role of 3BFs as opposed to simple phenomenological adjustment.

The inclusions of 3BFs in a microscopic description of nuclei often need partial-wave expansion in two Jacobi coordinates. An efficient method was developed by Hebeler *et al.* [6] for the local 3NFs. Here the local means that the interaction is a function of the momentum transfer of each Jacobi momentum except for the cutoff regularization function that does not depend on angle variables. In their method, the original eight-dimensional angular integration, though five-dimensional because of the rotational invariance, was reduced essentially to two-dimensional.

In this article, following the derivation in Ref. [6], a different expression for the partial-wave expansion of 3BFs is presented, which is more systematic for treating higher-rank coupling of spin and momentum vectors.

Before discussing the partial-wave expansion, the basic structure of leading-order 3BFs in ChEFT is summarized in Sec. II. The expression of partial wave decomposition of 3BFs in momentum space concerning the Jacobi coordinates is presented in Sec. III. As an application of the expression derived, a possible role of the  $\Lambda NN$  3BFs in the hypertriton is studied by calculating a  $\Lambda$ -deuteron folding potential from  $\Lambda NN$  3BFs. Summary follows in Sec. IV.

## II. STRUCTURE OF $\Lambda NN$ 3BFS IN CHEFT

Two-pion exchange  $\Lambda NN$  3BFs are considered as a concrete example, which is relevant for studying hypertriton. The structure of the  $\Lambda NN$  force in the leading-order, namely next-to-next-to-leading order (NNLO), is particularly simple because the  $\pi\Lambda\Lambda$  vertex is not present. The contribution is only from the diagram shown in Fig. 1.

Following the expression by Petschauer *et al.* [7], the Born amplitude of this diagram is written as

$$V_{TPE}^{\Lambda NN} = \frac{g_A^2}{3f_0^4} (\boldsymbol{\tau}_2 \cdot \boldsymbol{\tau}_3) \frac{(\boldsymbol{\sigma}_3 \cdot \mathbf{q}_{3d})(\boldsymbol{\sigma}_2 \cdot \mathbf{q}_{2d})}{(q_{3d}^2 + m_\pi^2)(q_{2d}^2 + m_\pi^2)} \times \{-(3b_0 + b_D)m_\pi^2 + (2b_2 + 3b_4)\mathbf{q}_{3d} \cdot \mathbf{q}_{2d}\}, \quad (1)$$

where the coordinate 1 is assigned to the  $\Lambda$  hyperon and  $\mathbf{q}_{2d}$  ( $\mathbf{q}_{3d}$ ) is the difference between the final and initial momenta at the nucleon line 2 (line 3):  $\mathbf{q}_{2d} = \mathbf{k}'_2 - \mathbf{k}_2$  and  $\mathbf{q}_{3d} = \mathbf{k}'_3 - \mathbf{k}_3$ .  $g_A$  is the axial coupling constant,  $f_0$  is the pion decay constant,  $m_\pi$  is the pion mass, and  $\boldsymbol{\sigma}$  and  $\boldsymbol{\tau}$  stand for the spin and isospin operators. The coupling constants  $b_0$ ,  $b_D$ ,  $b_2$ , and  $b_4$  inherit those in the underlying Lagrangian. These coupling constants are to be determined in parametrizing  $\Lambda N$  interactions in the next-to-next-to-leading-order. However, such an attempt




FIG. 1: Diagram of two-pion exchange  $\Lambda NN$  3BF. The small filled circle denotes the  $NN\pi$  vertex with the coupling constant  $g_A/f_0^2$ , and the large filled circle the  $NN\pi\pi$  vertex specified by the coupling constants  $3b_0 + b_D$  and  $2b_2 + 3b_4$  in Eq. (1).




FIG. 2: Three types of the Jacobi coordinate. The length of the vectors  $\mathbf{p}_i$  and  $\mathbf{q}_i$  does not correspond to the distance in the figure.

is not possible at present. In this paper, we use the estimation by Petshauer *et al.* [7].

In the following, particle 1 is assigned to the  $\Lambda$  hyperon and the case of the Jacobi coordinate  $(\mathbf{p}_1, \mathbf{q}_1)$  that is depicted by the leftmost diagram of Fig. 2 is considered. In the center-of-mass frame,  $\mathbf{k}_1 = \mathbf{q}_1$ ,  $\mathbf{k}_2 = \mathbf{p}_1 - r_{NN}\mathbf{q}_1$ , an  $\mathbf{k}_3 = -\mathbf{p}_1 - r_{NN}\mathbf{q}_1$  with  $r_{NN} \equiv 1/2$ . Then,  $\mathbf{q}_{1d} = \mathbf{k}'_1 - \mathbf{k}_1 = \mathbf{q}'_1 - \mathbf{q}_1$ ,  $\mathbf{q}_{2d} = \mathbf{k}'_2 - \mathbf{k}_2 = \mathbf{p}_1 - \mathbf{p}_1 - r_{NN}(\mathbf{q}'_1 - \mathbf{q}_1)$ , and  $\mathbf{q}_{3d} = \mathbf{k}'_3 - \mathbf{k}_3 = -(\mathbf{p}_1 - \mathbf{p}_1) - r_{NN}(\mathbf{q}'_1 - \mathbf{q}_1)$ .  $V_{TPE}^{ANN}$  is a function of  $\mathbf{p} \equiv \mathbf{p}'_1 - \mathbf{p}_1$  and  $\mathbf{q} \equiv \mathbf{q}'_1 - \mathbf{q}_1$  and can be organized in the following tensor-product representation.

$$V_{TPE}^{ANN}(\mathbf{p}, \mathbf{q}) = 4\pi(\boldsymbol{\tau}_2 \cdot \boldsymbol{\tau}_3) \sum_{K=0,1,2} \sum_{\ell_a, \ell_b} V_{TPE}^{(K, \ell_a, \ell_b)}(p, q) \times [[Y_{\ell_a}(\hat{\mathbf{p}}) \times Y_{\ell_b}(\hat{\mathbf{q}})]^K \times [\boldsymbol{\sigma}_2 \times \boldsymbol{\sigma}_3]^K]_0^0, \quad (2)$$

where the standard notation is employed for the tensor product:

$$[Y_{\ell_a}(\hat{\mathbf{a}}) \times Y_{\ell_b}(\hat{\mathbf{b}})]_{m_c}^{\ell_c} = \sum_{m_a m_b} (\ell_a m_a \ell_b m_b | \ell_c m_c) \times Y_{\ell_a m_a}(\hat{\mathbf{a}}) Y_{\ell_b m_b}(\hat{\mathbf{b}}). \quad (3)$$

The explicit expressions of  $V_{TPE}^{(K, \ell_a, \ell_b)}(p, q)$  for the Jacobi coordinate  $(\mathbf{p}_1, \mathbf{q}_1)$  are given in Appendix A. It is straightforward to obtain a similar representation for the other two sets of the Jacobi coordinate,  $(\mathbf{p}_2, \mathbf{q}_2)$  and  $(\mathbf{p}_3, \mathbf{q}_3)$  for the 3BFs of Eq. (1). It is noted that the third sigma operator  $\boldsymbol{\sigma}_3$  may appear in a general 3BF. In that case, the rank of  $K = 3$  can appear.

As for the cutoff, the following regulator function is introduced for the initial and final Jacobi coordinates,  $(\mathbf{a}, \mathbf{b}) = (\mathbf{p}_1, \mathbf{q}_1)$  or  $(\mathbf{p}'_1, \mathbf{q}'_1)$ , with the scale of  $\Lambda = 550$  MeV in present calculations:

$$f_\Lambda(\mathbf{a}, \mathbf{b}) = \exp\left\{-\left(\mathbf{a}^2 + \frac{3}{4}\mathbf{b}^2\right)/\Lambda^4\right\}. \quad (4)$$

Because this function does not depend on the angles, it does not affect the calculation of the partial-wave expansion, which is discussed in the next section.

### III. PARTIAL-WAVE EXPANSION

The partial-wave expansion is an operation of angle integrations concerning four Jacobi-coordinates  $\mathbf{p}'_1, \mathbf{q}'_1,$

$\mathbf{p}_1$ , and  $\mathbf{q}_1$  for a 3BF  $V_{3BF}(\mathbf{p}'_1, \mathbf{q}'_1, \mathbf{p}_1, \mathbf{q}_1)$  with 4 spherical-harmonic functions multiplied:

$$\frac{1}{\{(2\pi)^{3/2}\}^4} \int \cdots \int d\hat{\mathbf{p}}'_1 d\hat{\mathbf{q}}'_1 d\hat{\mathbf{p}}_1 d\hat{\mathbf{q}}_1 Y_{\ell'_p m'_p}^*(\hat{\mathbf{p}}'_1) \times Y_{\ell'_q m'_q}^*(\hat{\mathbf{q}}'_1) Y_{\ell_p m_p}(\hat{\mathbf{p}}_1) Y_{\ell_q m_q}(\hat{\mathbf{q}}_1) V_{3BF}(\mathbf{p}'_1, \mathbf{q}'_1, \mathbf{p}_1, \mathbf{q}_1). \quad (5)$$

To make the expression compact, the following notation is used in the subsequent derivation, in which the angular momenta  $\ell_p$  and  $\ell_q$  are coupled to  $L$ , and  $\ell'_p$  and  $\ell'_q$  to  $L'$ , respectively. That is, supplementing the summation

$$\sum_{m'_p m'_q} \sum_{m_p m_q} (\ell'_p m'_p \ell'_q m'_q | L' M'_L) (\ell_p m_p \ell_q m_q | L M_L), \quad (6)$$

the angular integration on a 3BF  $V_{3BF}$  is expressed as

$$\hat{I}_{L'L} V_{3BF} \equiv \frac{1}{(2\pi)^6} \langle [Y_{\ell'_p}(\hat{\mathbf{p}}'_1) \times Y_{\ell'_q}(\hat{\mathbf{q}}'_1)]_{M'_L}^{L'} | V_{3BF} \times [Y_{\ell_p}(\hat{\mathbf{p}}_1) \times Y_{\ell_q}(\hat{\mathbf{q}}_1)]_{M_L}^L \rangle. \quad (7)$$

For the case of a local 3BF  $V_{3BF}(\mathbf{p}'_1, \mathbf{q}'_1, \mathbf{p}_1, \mathbf{q}_1) = V_{3BF}(\mathbf{p}'_1 - \mathbf{p}_1, \mathbf{q}'_1 - \mathbf{q}_1)$ , the subtle manipulation [6] of adding a radial part to the angle integration with keeping the absolute value by a delta function is helpful:

$$d\hat{\mathbf{p}}'_1 \rightarrow d\mathbf{p}'_1 \frac{\delta(p''_1 - p'_1)}{p_1'^2} \quad \text{and} \quad d\hat{\mathbf{q}}'_1 \rightarrow d\mathbf{q}'_1 \frac{\delta(q''_1 - q'_1)}{q_1'^2}. \quad (8)$$

Changing the variables  $\mathbf{p}''_1$  and  $\mathbf{q}''_1$  to  $\mathbf{p}$  and  $\mathbf{q}$  by the shifts of  $\mathbf{p}''_1 = \mathbf{p} + \mathbf{p}_1$  and  $\mathbf{q}''_1 = \mathbf{q} + \mathbf{q}_1$ , Eq. (7) is modified as

$$\hat{I}_{L'L} V_{3BF} = \frac{1}{(2\pi)^6} \int_0^\infty p^2 dp \int_0^\infty q^2 dq \times \langle [Y_{\ell'_p}(\widehat{\mathbf{p}}_1 + \mathbf{p}) \times Y_{\ell'_q}(\widehat{\mathbf{q}}_1 + \mathbf{q})]_{M'_L}^{L'} | \times \frac{\delta(|\mathbf{p} - \mathbf{p}_1| - p'_1)}{p_1'^2} \frac{\delta(|\mathbf{q} - \mathbf{q}_1| - q'_1)}{q_1'^2} V_{3BF} \times [Y_{\ell_p}(\hat{\mathbf{p}}_1) \times Y_{\ell_q}(\hat{\mathbf{q}}_1)]_{M_L}^L \rangle. \quad (9)$$

The delta function can be written as follows by using a Legendre polynomial of the first kind  $P_k$ :

$$\delta(|\mathbf{p} - \mathbf{p}_1| - p'_1) = \frac{p'_1}{pp_1} \delta\left(\cos \widehat{\mathbf{p}}_1 \mathbf{p} - \frac{p_1'^2 - p_1^2 - p^2}{2p_1 p}\right) = 2\pi \frac{p'_1}{pp_1} \sum_{k'} P_{k'}(c_p) (-1)^k \sqrt{\hat{k}'} [Y_{k'}(\hat{\mathbf{p}}_1) \times Y_{k'}(\hat{\mathbf{p}})]_0^0, \quad (10)$$

$$\delta(|\mathbf{q} - \mathbf{q}_1| - q'_1) = \frac{q'_1}{qq_1} \delta\left(\cos \widehat{\mathbf{q}}_1 \mathbf{q} - \frac{q_1'^2 - q_1^2 - q^2}{2q_1 q}\right) = 2\pi \frac{q'_1}{qq_1} \sum_k P_k(c_q) (-1)^k \sqrt{\hat{k}} [Y_k(\hat{\mathbf{q}}_1) \times Y_k(\hat{\mathbf{q}})]_0^0, \quad (11)$$

where  $\hat{k} \equiv 2k + 1$ ,  $c_p \equiv \frac{p_1'^2 - p_1^2 - p^2}{2p_1 p}$  and  $c_q \equiv \frac{q_1'^2 - q_1^2 - q^2}{2q_1 q}$ . These  $\delta$ -functions restrict the  $p$ - and  $q$ -integrations as

$$p_{min} \equiv |p'_1 - p_1| \leq p \leq p_{max} \equiv p'_1 + p_1, \quad (12)$$

$$q_{min} \equiv |q'_1 - q_1| \leq q \leq q_{max} \equiv q'_1 + q_1. \quad (13)$$

The spherical-harmonic functions  $Y_{\ell_p}(\widehat{\mathbf{p}_1 + \mathbf{p}})$  and  $Y_{\ell_q}(\widehat{\mathbf{q}_1 + \mathbf{q}})$  are also expanded as follows:

$$Y_{\ell_p m'_p}(\widehat{\mathbf{p}_1 + \mathbf{p}}) = \sum_{j_p + j'_p = \ell_p} \sqrt{\frac{4\pi(\ell'_p)!}{(\hat{j}_p)!(\hat{j}'_p)!} \frac{p_1^{j_p} p^{j'_p}}{p_1^{\ell'_p}}} \times [Y_{j_p}(\hat{\mathbf{p}}_1) \times Y_{j'_p}(\hat{\mathbf{p}})]_{m'_p}^{\ell'_p}, \quad (14)$$

$$Y_{\ell'_q m'_q}(\widehat{\mathbf{q}_1 + \mathbf{q}}) = \sum_{j_q + j'_q = \ell'_q} \sqrt{\frac{4\pi(\ell'_q)!}{(\hat{j}_q)!(\hat{j}'_q)!} \frac{q_1^{j_q} q^{j'_q}}{q_1^{\ell'_q}}} \times [Y_{j_q}(\hat{\mathbf{q}}_1) \times Y_{j'_q}(\hat{\mathbf{q}})]_{m'_q}^{\ell'_q}. \quad (15)$$

A straightforward evaluation of the recoupling of these decoupled spherical harmonics finally gives

$$\begin{aligned} \hat{I}_{L'L} V_{3BF} &= \frac{1}{(2\pi)^6} (-1)^{\ell'_p + \ell'_q - L'} \sum_{JM} (-1)^{M_L} (L' - M_L L M_L | JM) \frac{1}{p_1' q_1' p_1 q_1} \int_{p_{min}}^{p_{max}} pdp \int_{q_{min}}^{q_{max}} qdq \\ &\times (2\pi)^2 \sum_{k'k} \hat{k}' \hat{k} P_{k'}(c_p) P_k(c_q) \sum_{j_p + j'_p = \ell_p} \sqrt{\frac{(\ell'_p)!}{(\hat{j}_p)!(\hat{j}'_p)!} \frac{p_1^{j_p} p^{j'_p}}{p_1^{\ell'_p}}} \sum_{j_q + j'_q = \ell'_q} (-1)^{j_p + j_q} \sqrt{\frac{(\ell'_q)!}{(\hat{j}_q)!(\hat{j}'_q)!} \frac{q_1^{j_q} q^{j'_q}}{q_1^{\ell'_q}}} \sqrt{\hat{\ell}'_p \hat{\ell}'_q \hat{L} \hat{L}'} \\ &\times \frac{1}{4\pi} \sum_{L_p L_q} \sqrt{\hat{j}'_p \hat{j}_q \hat{j}_p \hat{j}'_q} (k' 0 j'_p 0 | L_p 0) (k 0 j_q 0 | L_q 0) (k' 0 j_p 0 | \ell_p 0) (k 0 j_q 0 | \ell_q 0) \\ &\times \left\{ \begin{matrix} \ell_q & \ell'_q & L_q \\ j'_q & k & j_q \end{matrix} \right\} \left\{ \begin{matrix} \ell_p & \ell'_p & L_p \\ j'_p & k' & j_p \end{matrix} \right\} \left\{ \begin{matrix} \ell'_p & \ell'_q & L' \\ \ell_p & \ell_q & L \\ L_p & L_q & J \end{matrix} \right\} \int d\hat{\mathbf{p}} d\hat{\mathbf{q}} [Y_{L_p}(\hat{\mathbf{p}}) \times Y_{L_q}(\hat{\mathbf{q}})]_M^J V_{3BF}(\mathbf{p}, \mathbf{q}) \end{aligned} \quad (16)$$

Observing  $P_\ell(\cos \widehat{\mathbf{p}\mathbf{q}}) = (-1)^\ell \frac{4\pi}{\sqrt{\ell}} [Y_\ell(\hat{\mathbf{p}}) \times Y_\ell(\hat{\mathbf{q}})]_0^0$ , the above expression with  $J = 0$  and  $L_p = L_q = \ell$  corresponds to Eq. (6) of Ref. [6]. The evaluation of  $\hat{I}_{L'L} V_{3BF}$  with  $J > 0$  is straightforward and transparent.

The angle integration in Eq. (16) for 3BFs in the form of Eq. (2) needs

$$\begin{aligned} &\int d\hat{\mathbf{p}} d\hat{\mathbf{q}} [Y_{L_p}(\hat{\mathbf{p}}) \times Y_{L_q}(\hat{\mathbf{q}})]_M^J [Y_{\ell_a}(\hat{\mathbf{p}}) \times Y_{\ell_b}(\hat{\mathbf{q}})]_\mu^K \\ &= (-1)^{k+\mu} \delta_{JK} \delta_{M, -\mu} \delta_{L_p \ell_a} \delta_{L_q \ell_b} (-1)^{\ell_a + \ell_b}. \end{aligned} \quad (17)$$

This means that the angular momentum  $J$  in Eq. (16) corresponds to the rank  $K$  of the tensor-product structure of the angular-momentum coupling in 3BFs.

For actual calculations of the matrix element of 3BFs in various situations, it is convenient to define a reduced matrix element in a form similar to the Wigner-Eckart theorem.

$$\begin{aligned} &\langle [Y_{\ell'_p}(\hat{\mathbf{p}}_1) \times Y_{\ell'_q}(\hat{\mathbf{q}}_1)]_{M_L'}^{L'} [Y_{\ell_a}(\hat{\mathbf{p}}) \times Y_{\ell_b}(\hat{\mathbf{q}})]_\mu^K [Y_{\ell_p}(\hat{\mathbf{p}}_1) \times Y_{\ell_q}(\hat{\mathbf{q}}_1)]_{M_L}^L \rangle \\ &= (L M_L K \mu | L' M_L') \langle [Y_{\ell'_p}(\hat{\mathbf{p}}_1) \times Y_{\ell'_q}(\hat{\mathbf{q}}_1)]^{L'} || [Y_{\ell_a}(\hat{\mathbf{p}}) \times Y_{\ell_b}(\hat{\mathbf{q}})]^K || [Y_{\ell_p}(\hat{\mathbf{p}}_1) \times Y_{\ell_q}(\hat{\mathbf{q}}_1)]^L \rangle_{pwe}. \end{aligned} \quad (18)$$

From Eq. (16), the reduced matrix element is found as

$$\begin{aligned} &\langle [Y_{\ell'_p}(\hat{\mathbf{p}}_1) \times Y_{\ell'_q}(\hat{\mathbf{q}}_1)]^{L'} || [Y_{\ell_a}(\hat{\mathbf{p}}) \times Y_{\ell_b}(\hat{\mathbf{q}})]^K || [Y_{\ell_p}(\hat{\mathbf{p}}_1) \times Y_{\ell_q}(\hat{\mathbf{q}}_1)]^L \rangle_{pwe} \\ &= \frac{1}{4\pi} \frac{1}{(2\pi)^4} \sqrt{\frac{\hat{K}}{\hat{L}'}} (-1)^{\ell_p + \ell_q} \frac{1}{p_1' q_1' p_1 q_1} \sqrt{\hat{\ell}'_p \hat{\ell}'_q \hat{\ell}_p \hat{\ell}_q \hat{L} \hat{L}' \hat{\ell}_a \hat{\ell}_b} \int_{p_{min}}^{p_{max}} pdp \int_{q_{min}}^{q_{max}} qdq \sum_{k'k} \hat{k}' \hat{k} P_{k'}(c_p) P_k(c_q) \\ &\times \sum_{j_p + j'_p = \ell_p} \sqrt{\frac{(\ell'_p)!}{(\hat{j}_p)!(\hat{j}'_p)!} \frac{p_1^{j_p} p^{j'_p}}{p_1^{\ell'_p}}} \sum_{j_q + j'_q = \ell'_q} (-1)^{j_p + j_q} \sqrt{\frac{(\ell'_q)!}{(\hat{j}_q)!(\hat{j}'_q)!} \frac{q_1^{j_q} q^{j'_q}}{q_1^{\ell'_q}}} (k' 0 \ell_a 0 | j'_p 0) (k 0 \ell_b 0 | j_q 0) \\ &\times (k' 0 \ell_p 0 | j_p 0) (k 0 \ell_q 0 | j_q 0) \left\{ \begin{matrix} \ell_q & \ell'_q & \ell_b \\ j'_q & k & j_q \end{matrix} \right\} \left\{ \begin{matrix} \ell_p & \ell'_p & \ell_a \\ j'_p & k' & j_p \end{matrix} \right\} \left\{ \begin{matrix} \ell'_p & \ell'_q & L' \\ \ell_p & \ell_q & L \\ \ell_a & \ell_b & K \end{matrix} \right\}. \end{aligned} \quad (19)$$

It should be reminded that this expression is supposed for the 3BF with  $\mathbf{p} = \mathbf{p}'_1 - \mathbf{p}_1$  and  $\mathbf{q}'_1 = \mathbf{q}_1$ . An introduction of the cutoff regularization function as a function of  $p'_q$ ,  $q'_1$ ,  $p_1$ , and  $q_1$  does not affect the angular integration.

#### IV. $\Lambda$ -DEUTERON FOLDING POTENTIAL

The construction of hyperon-nucleon interactions in the strangeness  $S = -1$  sector has a difficulty in lacking sufficient scattering data. The fact that there is no two-body  $\Lambda N$  bound state enhances the difficulty. The hypertriton is, therefore, an important hyper-nuclear system [8, 9] for adjusting the interaction, where the tuning of  $\Lambda N$  interactions in the spin-singlet and triplet states together with the strength of  $\Lambda N$ - $\Sigma N$  coupling can be carried out. However, the possible role of the  $\Lambda NN$  3BFs has been investigated little. Expecting the settlement of the current controversy over the shallow binding energy [10], it is important to describe the hypertriton including  $\Lambda NN$  3BFs. Before considering Faddeev calculations for the hypertriton including  $\Lambda NN$  3BFs, it is instructive to calculate  $\Lambda$ -deuteron folding potential due to the  $\Lambda NN$  interactions, applying the expression derived in the previous section, to obtain some idea about the contribution of the 3BFs.

The folding potential is evaluated by the following integration:

$$U_{\Lambda-d}^{J_t}(q'_1, q_1) = \iint d\mathbf{p}'_1 d\mathbf{p}_1 \langle [\Psi_d(\mathbf{p}'_1), (\ell'_\Lambda 1/2) j_\Lambda] J_t | V_{TPE}^{\Lambda NN} \times | [\Psi_d(\mathbf{p}_1), (\ell_\Lambda 1/2) j_\Lambda] J_t \rangle, \quad (20)$$

$$\Psi_d(\mathbf{p}_1) = \sum_{\ell_d=0,2} \frac{1}{p_1} \phi_{\ell_d}(p_1) [Y_{\ell_d}(\hat{\mathbf{p}}_1) \times \chi_{d,m}^{11}]. \quad (21)$$

The above expression is in an abbreviated notation. A more detailed calculational procedure is given in Appendix B.

Deuteron wave functions, the  $s$ - and  $d$ -components, are those of the  $N^3\text{LO}$  ChEFT interactions with the cutoff of 550 MeV [11]. This scale of the wave functions and the 3BFs may not be soft enough to permit a perturbative treatment. Nevertheless, without strong short-range singularities, the resulting folding potential helps intuitively infer the  $\Lambda$ -deuteron interaction and therefore the possible role of the  $\Lambda NN$  3BFs to the hypertriton. It is noted that it is not appropriate to employ deuteron wave functions of other  $NN$  interactions having strong short-range singularities together with the ChEFT 3BFs. For comparison, deuteron wave functions in momentum space are shown in Fig. 3 in which those of other  $NN$  interactions, AV18 [12], CD-Bonn [13], and Paris [14], are compared.

Calculated  $s$ -wave ( $\ell'_\Lambda = \ell_\Lambda = 0$  in Eq. (20))  $\Lambda$ -deuteron folding potential with  $J_t = 1/2$  from the leading-order  $\Lambda NN$  interactions is presented in Fig. 4. The upper panel shows the contribution from the  $s$ -wave component of the deuteron wave function, in which

$V_{TPE}^{(K=0, \ell_a, \ell_b)}$  participates. The lower panel presents the sum of the remaining contributions from the  $s$ - $d$ ,  $d$ - $s$ , and  $d$ - $d$  pairs of the deuteron wave function.  $V_{TPE}^{(K=2, \ell_a, \ell_b)}$  contributes in the  $s$ - $d$  and  $d$ - $s$  pairs. Both  $V_{TPE}^{(K=0, \ell_a, \ell_b)}$  and  $V_{TPE}^{(K=2, \ell_a, \ell_b)}$  contribute in the case of the  $d$ - $d$  pair. The potential with  $J_t = 3/2$  is identical to that with  $J_t = 1/2$ .

The parameters employed are taken from the estimation by Petshauer *et al.* [7]; that is,  $(3b_0 + b_D) = 0$  and  $(2b_2 + 3b_4) = 3.0 \times 10^{-3}$  MeV. In principle, parameters of the two-pion exchange 3BFs should be determined in fitting the two-body  $\Lambda N$  interactions. However, the present experimental situation of the strangeness  $S = -1$  sector does not allow such an investigation.

The experimental separation energy of the hypertriton is very shallow, though the actual value is still controversial. Therefore, the  $\Lambda NN$  3BF contribution may be argued to be hindered. However, these results shown in Fig. 4 indicate that they may not be negligible, though the magnitude depends on the coupling constants. Thus, it suggests that further study of the hypertriton in Faddeev formalism with incorporating the 3BFs is necessary. The repulsive bump structure seen beyond  $q \simeq 2.5$  fm implies that the scale of 550 MeV employed for the ChEFT description still has remnants of shorter-range singularities, which should be treated in a Faddeev framework.




FIG. 3: Deuteron  $s$ - and  $d$ -wave functions in momentum space described by various  $NN$  interactions: ChEFT [11], AV18 [12], CD-Bonn [13], and Paris [14].  $u$  and  $w$  correspond to  $\phi_0$  and  $\phi_2$  in Eq. (21), respectively.

## V. SUMMARY

We have presented an expression of partial-wave decomposition of 3BFs concerning the relevant Jacobi coordinates. The derivation essentially follows that of Hebeler *et al.* [6], but the final formula is different that can systematically treat the higher-rank spin and angular-momentum tensor-product structure of 3BFs. Although the consideration is inspired specifically for  $\Lambda NN$  3BFs and one set of the Jacobi coordinate, the formula is general, as far as 3BFs are a function of the momentum transfer in each Jacobi coordinate. Even if a regularization function is angle-dependent, the 3BFs can be expressed in the form of Eq. (2) and then the result of Eq. (19) is applicable.

As an application of the derived expression, the  $\Lambda$ -

deuteron folding potential from NNLO  $\Lambda NN$  3BFs is evaluated. At the present stage, the construction of baryon-baryon interactions in the strangeness  $S = -1$  sector in ChEFT is practiced up to the NLO level [15, 16]. Even at this low order, it is difficult to unambiguously determine coupling constants because of the lack of sufficient experimental data, and therefore a plausible assumption of the SU(3) symmetry has to be called for. In addition, there is no conclusive way to fix the parameters of the two-pion exchange  $\Lambda NN$  3BFs that are basically of the NNLO. This means that experimental and theoretical investigations are needed in the future. It is essentially important to quantitatively establish the contribution of  $\Lambda NN$  3BFs in hypernuclei, which is relevant also for the understanding of the appearance or absence of  $\Lambda$  hyperons in neutron star matter [17, 18].




FIG. 4:  $\Lambda$ -deuteron folding potential with  $\ell_\Lambda = 0$  from two-pion exchange  $\Lambda NN$  interactions. The upper panel shows the contribution of the deuteron  $s$ -state pair. The lower panel is the remaining contributions of the  $sd$ ,  $ds$ , and  $dd$  pairs.

In particular, the investigation of the hypertriton is basic. Before doing full Faddeev calculations for the hypertriton including  $\Lambda NN$  3BFs, it is worthwhile to estimate the possible role of the 3BFs for the hypertriton. The present folding potential calculation indicates the necessity of considering  $\Lambda NN$  3BFs in the theoretical study of the hypertriton because the quantitative estimation for it affects the parametrization of  $\Lambda N$  two-body interactions.

*Acknowledgements.* This work is supported by JSPS KAKENHI Grants No. JP19K03849 and No. JP22K03597.

### Appendix A: Tensor-product decomposition of $\Lambda NN$ three-body interactions

$V_{TPE}^{(k,\ell_a,\ell_b)}(p, q)$  in Eq. (2) for  $k = 0, 1$  and  $2$  are:

$$V_{TPE}^{k=0,\ell_a,\ell_b}(p, q) = (-1)^{\ell_a} \delta_{\ell_a\ell_b} \delta_{\ell_a, \text{even}} \sqrt{\frac{\hat{\ell}_a \{C_0 m_\pi^2 + C_1(-p^2 + r_{NN}^2 q^2)\}}{3}} \frac{(p^2 - r_{NN}^2 q^2)}{2pr_{NN}q(p^2 + r_{NN}^2 q^2 + m_\pi^2)} Q_{\ell}(z_{pq}), \quad (\text{A1})$$

$$V_{TPE}^{k=1,\ell_a,\ell_b}(p, q) = \delta_{\ell_a\ell_b} \delta_{\ell_a, \text{odd}} \frac{\{C_0 m_\pi^2 + C_1(-p^2 + r_{NN}^2 q^2)\}}{(p^2 + r_{NN}^2 q^2 + m_\pi^2)} \frac{1}{\hat{\ell}_a} \sqrt{\frac{\ell_a(\ell_a + 1)}{6\hat{\ell}_a}} \{Q_{\ell_a+1}(z_{pq}) - Q_{\ell_a-1}(z_{pq})\}, \quad (\text{A2})$$

$$V_{TPE}^{k=2,\ell_a,\ell_b}(p, q) = -\sqrt{\frac{2}{15}} \frac{\{C_0 m_\pi^2 + C_1(-p^2 + r_{NN}^2 q^2)\}}{2pr_{NN}q(p^2 + r_{NN}^2 q^2 + m_\pi^2)} \sqrt{\hat{\ell}_a \hat{\ell}_b} (\ell_a 0 \ell_b 0 | 20) \{p^2 Q_{\ell_b}(z_{pq}) - r_{NN}^2 q^2 Q_{\ell_a}(z_{pq})\}, \quad (\text{A3})$$

where  $C_0 = -\frac{g_A^2}{3f_0^2}(3b_0 + b_D)$ ,  $C_1 = \frac{g_A^2}{3f_0^2}(2b_2 + 3b_4)$ , and  $z_{pq} = \frac{p^2 + r_{NN}^2 q^2 + m_\pi^2}{2pr_{NN}q}$ .  $Q_\ell$  is a Legendre polynomial of the second kind.

### Appendix B: Evaluation of $\Lambda$ -deuteron folding potential from $\Lambda NN$ 3BFs

An explicit calculational procedure of the  $\Lambda$ -deuteron folding potential, Eq. (20), is provided. The angular-momentum recoupling in Eq. (20) yields

$$\begin{aligned} & \langle [[Y_{\ell'_d}(\hat{\mathbf{p}}'_1) \times \chi_d^1]^1 \times [Y_{\ell'_\Lambda}(\hat{\mathbf{q}}'_1) \times \chi_\Lambda^{1/2}]^{j_\Lambda}]_{M_t}^{J_t} | [[Y_{\ell_a}(\hat{\mathbf{p}}) \times Y_{\ell_b}(\hat{\mathbf{q}})]^K \times [\boldsymbol{\sigma}_2 \times \boldsymbol{\sigma}_3]^K]_0^0 | [[Y_{\ell_d}(\hat{\mathbf{p}}_1) \times \chi_d^1]^1 \times [Y_{\ell_\Lambda}(\hat{\mathbf{q}}_1) \times \chi_\Lambda^{1/2}]^{j_\Lambda}]_{M_t}^{J_t} \rangle \\ &= \sum_{L'L} \sum_{S'S} 3\hat{j}_\Lambda \sqrt{\hat{L}'\hat{S}'\hat{L}\hat{S}} \begin{Bmatrix} \ell'_d & \ell_{\Lambda'} & L' \\ 1 & 1/2 & S' \\ 1 & j_{\Lambda'} & J_t \end{Bmatrix} \begin{Bmatrix} \ell_d & \ell_\Lambda & L \\ 1 & 1/2 & S \\ 1 & j_{\Lambda'} & J_t \end{Bmatrix} \langle [[Y_{\ell'_d}(\hat{\mathbf{p}}'_1) \times Y_{\ell'_\Lambda}(\hat{\mathbf{q}}'_1)]^{L'} \times [\chi_d^1 \times \chi_\Lambda^{1/2}]^{S'}]_{M_t}^{J_t} | \\ & \quad \times [[Y_{\ell_a}(\hat{\mathbf{p}}) \times Y_{\ell_b}(\hat{\mathbf{q}})]^K \times [\boldsymbol{\sigma}_2 \times \boldsymbol{\sigma}_3]^K]_0^0 | [[Y_{\ell_d}(\hat{\mathbf{p}}_1) \times Y_{\ell_\Lambda}(\hat{\mathbf{q}}_1)]^L \times [\chi_d^1 \times \chi_\Lambda^{1/2}]^S]_{M_t}^{J_t} \rangle \\ &= \sum_{L'L} \sum_{S'S} 3\hat{j}_\Lambda \sqrt{\hat{L}'\hat{S}'\hat{L}\hat{S}} \begin{Bmatrix} \ell'_d & \ell_{\Lambda'} & L' \\ 1 & 1/2 & S' \\ 1 & j_{\Lambda'} & J_t \end{Bmatrix} \begin{Bmatrix} \ell_d & \ell_\Lambda & L \\ 1 & 1/2 & S \\ 1 & j_{\Lambda'} & J_t \end{Bmatrix} \sqrt{\hat{J}_t \hat{L}' \hat{S}'} \begin{Bmatrix} J_t & J_t & 0 \\ L' & L & K \\ S' & S & K \end{Bmatrix} \\ & \quad \times \langle [Y_{\ell'_d}(\hat{\mathbf{p}}'_1) \times Y_{\ell'_\Lambda}(\hat{\mathbf{q}}'_1)]^{L'} | [[Y_{\ell_a}(\hat{\mathbf{p}}) \times Y_{\ell_b}(\hat{\mathbf{q}})]^K | [[Y_{\ell_d}(\hat{\mathbf{p}}_1) \times Y_{\ell_\Lambda}(\hat{\mathbf{q}}_1)]^L]_{pwe} \\ & \quad \times 18\sqrt{\hat{S}\hat{K}} (-1)^{K+3/2+S} \begin{Bmatrix} S' & S & K \\ 1 & 1 & 1/2 \end{Bmatrix} \begin{Bmatrix} 1 & 1 & K \\ 1/2 & 1/2 & 1 \\ 1/2 & 1/2 & 1 \end{Bmatrix}, \quad (\text{B1}) \end{aligned}$$

where  $\mathbf{p} = \mathbf{p}'_1 - \mathbf{p}_1$  and  $\mathbf{q} = \mathbf{q}'_1 - \mathbf{q}_1$ . Then, Eq. (19) is applied in this expression.

- 
- [1] A. Akmal, V. R. Pandharipande, and D. G. Ravenhall, "Equation of state of nuclear matter and neutron star structure," *Phys. Rev. C* **58**, 1804 (1998).
  - [2] S. C. Pieper, V. R. Pandharipande, R. B. Wiringa, and J. Carlson, "Realistic models of pion-exchange three-nucleon interactions," *Phys. Rev. C* **64**, 014001 (2001).
  - [3] M. Baldo and G. F. Burgio, "Properties of the nuclear medium," *Rep. Prog. Phys.* **75**, 026301 (2012).
  - [4] E. Epelbaum, H.-W. Hammer, and U.-G. Meißner, "Modern theory of nuclear force," *Rev. Mod. Phys.* **81**, 1773 (2009).
  - [5] R. Machleidt and D. R. Entem, "Chiral effective field theory and nuclear forces," *Phys. Rep.* **503**, 1 (2011).
  - [6] K. Hebeler, H. Krebs, E. Epelbaum, J. Golak, and R. Skibinski, "Efficient calculation of chiral three-nucleon forces up to N3LO for ab initio studies," *Phys. Rev. C* **91**, 044001 (2015).
  - [7] S. Petschauer, N. Kaiser, J. Haidenbauer, U.-G. Meißner, and W. Weise, "Leading three-baryon forces from SU(3) chiral effective field theory," *Phys. Rev. C* **93**, 014001 (2016).
  - [8] K. Miyagawa, H. Kamada, W. Glöckle, V. Stoks, "Properties of the bound  $\Lambda(\Sigma)NN$  system and hyperon-nucleon interactions", *Phys. Rev. C* **51**, 2905 (1995).
  - [9] J. Haidenbauer, U.-G. Meißner, and A. Nogga, "Hyperon-nucleon interaction within chiral effective field

- theory revisited”, *Eur. Phys. J. A* **56**, 91 (2020).
- [10] J. Adam *et al.* (STAR Collaboration), ”Measurement of the mass difference and the binding energy of the hypertriton and antihypertriton”, *Nature Phys.* **16**, 409 (2020).
- [11] E. Epelbaum, W. Glöckle, and U.-G. Meißner, ”The two-nucleon system at next-to-next-to-next-to-leading order,” *Nucl. Phys. A* **747**, 362 (2005).
- [12] R. B. Wiringa, V. G. J. Stoks, and R. Schiavilla, ”Accurate nucleon-nucleon potential with charge-independence breaking,” *Phys. Rev. C* **51**, 38 (1995).
- [13] R. Machleidt, ”High-precision, charge-dependent Bonn nucleon-nucleon potential,” *Phys. Rev. C* **63**, 024001 (2001).
- [14] M. Lacombe, B. Loiseau, R. Vinh Mau, J. Côté, P. Pirés, and R. de Tourreil, ”Parametrization of the deuteron wave function of the Paris potential,” *Phys. Lett.* **101B**, 139 (1981).
- [15] J. Haidenbauer, S. Petschauer, N. Kaiser, U.-G. Meißner, A. Nogga, and W. Weise, ”Hyperon-nucleon interaction at next-to-leading order in chiral effective field theory,” *Nucl. Phys. A* **915**, 24 (2013).
- [16] J. Haidenbauer, U.-G. Meißner, and A. Nogga, ”Hyperon-nucleon interaction within chiral effective field theory revisited,” *Eur. Phys. J. A* **56**, 91 (2020).
- [17] M. Kohno, ”Single-particle potential of the hyperon in nuclear matter with chiral effective field theory NLO interactions including effects of YNN three-baryon interactions,” *Phys. Rev. C* **97**, 035206 (2018).
- [18] D. Gerstung, N. Kaiser, and W. Weise, ”Hyperon-nucleon three-body forces and strangeness in neutron stars,” *Eur. Phys. J. A* **56**, 175 (2020).