

# Exact results for generalized extremal problems forbidding an even cycle

Ervin Győri<sup>1</sup>, Zhen He<sup>1,3</sup>, Zequn Lv<sup>1,3</sup>, Nika Salia<sup>1</sup>, Casey Tompkins<sup>1</sup>, Kitti Varga<sup>1,4</sup>,  
and Xiutao Zhu<sup>1,2</sup>

<sup>1</sup>Alfréd Rényi Institute of Mathematics, Hungarian Academy of Sciences.

<sup>2</sup>Department of Mathematics, Nanjing University.

<sup>3</sup>Department of Mathematical Sciences, Tsinghua University.

<sup>4</sup>Department of Computer Science and Information Theory, Budapest University of Technology and Economics.

## Abstract

We determine the maximum number of copies of  $K_{s,s}$  in a  $C_{2s+2}$ -free  $n$ -vertex graph for all integers  $s \geq 2$  and sufficiently large  $n$ . Moreover, for  $s \in \{2, 3\}$  and any integer  $n$  we obtain the maximum number of cycles of length  $2s$  in an  $n$ -vertex  $C_{2s+2}$ -free bipartite graph.

## 1 Introduction

**Notation.** For a graph  $G$ , the vertex and edge set of  $G$  are denoted by  $V(G)$  and  $E(G)$ , respectively. Furthermore, we let  $v(G) = |V(G)|$  and  $e(G) = |E(G)|$ . The maximum degree of a vertex in  $G$  is denoted by  $\Delta(G)$ . For  $S \subseteq V(G)$ , we denote by  $G[S]$  the subgraph of  $G$  induced on the vertex set  $S$ . For a vertex  $v$ , we denote by  $G - v$  the graph  $G[V(G) \setminus \{v\}]$ . Moreover, for a subgraph  $H$  of  $G$  we write  $G - H$  for  $G[V(G) \setminus V(H)]$ . For  $v \in V(G)$ , we denote by  $N(v)$  the neighborhood of  $v$  and we write  $N_H(v)$  for  $N(v) \cap V(H)$ . For a vertex  $v$ , the set  $N(v) \cup \{v\}$  is denoted by  $N[v]$ . For a set  $S$ , we denote the set of  $k$ -element subsets of  $S$  by  $\binom{S}{k}$ . By a copy of  $H$  in a graph  $G$  we mean a subgraph of  $G$  isomorphic to  $H$ . For graphs  $G$  and  $H$ , we denote by  $G + H$  the join of  $G$  and  $H$ , that is the graph obtained by connecting each pair of vertices between a vertex disjoint copy of  $G$  and  $H$ .

The path, cycle and complete graph on  $k$  vertices are denoted by  $P_k$ ,  $C_k$  and  $K_k$ , respectively. Let  $S_k$  be a  $k$ -vertex complete bipartite graph with a color class of size 1. We sometimes refer to a  $k$ -vertex path by a sequence of vertices  $v_1 v_2 \dots v_k$ , where  $v_i v_{i+1}$  is an edge for all  $1 \leq i \leq k - 1$ , and we sometimes refer to a  $k$ -vertex cycle by a sequence of the form  $v_1 v_2 \dots v_k v_1$  where  $v_1 v_k$  and  $v_i v_{i+1}$  are edges for all  $1 \leq i \leq k - 1$ . We denote by  $K_{a,b}$  the complete bipartite graph with parts of size  $a$  and  $b$ , and we denote by  $K_{a,b}^+$  the join of a clique of size  $a$  and an independent set of size  $b$ . A star is a tree with at most one vertex of degree at least two; this vertex is referred to as the central vertex of the star. A forest in which every connected component is a star is called a star forest. A double star is a tree consisting of two adjacent vertices  $u$  and  $v$  such that every vertex in  $(N(u) \setminus \{v\}) \cup (N(v) \setminus \{u\})$  is a leaf; such vertices  $u$  and  $v$  are referred to as central vertices of the double star. For any two graphs  $H$  and  $G$ , we denote the number of copies of  $H$  in  $G$  by  $H(G)$ . For given graphs  $F$  and  $G$ , we say that  $G$  is  $F$ -free if it does not contain  $F$  as a subgraph (not necessarily induced). The maximum of  $H(G)$  across  $n$ -vertex  $F$ -free graphs  $G$  is denoted by  $\text{ex}(n, H, F)$ . The maximum of  $H(G)$  across bipartite  $n$ -vertex  $F$ -free graphs  $G$  is denoted by  $\text{ex}_{\text{bip}}(n, H, F)$ .

**Background.** Generalized extremal problems have a long history dating back to a result of Zykov [16] (and later independently Erdős [3]), who determined for all  $s$  and  $t$  the value of

$\text{ex}(n, K_s, K_t)$ , extending the classical theorem of Turán [15]. Following this initial result, there has been extensive work determining the value of  $\text{ex}(n, H, F)$  for various pairs of graphs  $H$  and  $F$ . An important early result in this direction is a result of Győri, Pach and Simonovits [12] which determined the value of  $\text{ex}(n, H, K_3)$  for all bipartite graphs  $H$  containing a matching on all but at most one of its vertices. The generalized extremal function  $\text{ex}(n, H, F)$  for arbitrary pairs of graphs was introduced by Alon and Shikhelman [1]. In addition to the function  $\text{ex}(n, H, F)$ , we will consider the analogous function under the assumption that the ground graph is bipartite, which we denote by  $\text{ex}_{\text{bip}}(n, H, F)$ .

In the present paper we will be interested in the setting when the forbidden graph is an even cycle. Determining  $\text{ex}(n, C_5, C_3)$  was a long standing open problem of Erdős [4]. A construction is obtained by taking a blow-up of  $C_5$  consisting of almost equal classes. This problem was finally settled by Grzesik [9] and independently by Hatami, Hladký, Král, Norine and Razborov [14]. This result has recently been extended Grzesik and Kielak [10] who determined  $\text{ex}(n, C_{2k+1}, C_{2k-1})$  asymptotically for all  $k > 2$ . The dual problem of determining  $\text{ex}(n, C_3, C_5)$  was introduced by Bollobás and Győri [2] who proved an upper bound, and this upper bound was subsequently improved in [1], [6] and [5]. Bounds on  $\text{ex}(n, C_3, C_{2k+1})$  were obtained by Győri and Li [11]. An exact result for  $\text{ex}(n, C_5, C_7)$  was obtained by Górski and Grzesik [13]. The order of magnitude of the  $\text{ex}(n, C_k, C_\ell)$  for general  $k$  and  $\ell$  was determined by Gishboliner and Shapira [8] and independently for even  $k$  and  $\ell$  by Gerbner, Győri, Methuku and Vizer [7].

In [7] an asymptotic result was obtained for  $\text{ex}(n, C_4, C_6)$  and  $\text{ex}_{\text{bip}}(n, C_6, C_8)$ . We determine the exact value of  $\text{ex}_{\text{bip}}(n, C_4, C_6)$ ,  $\text{ex}_{\text{bip}}(n, C_6, C_8)$  for all  $n$ , and the exact value of  $\text{ex}(n, C_4, C_6)$  for sufficiently large  $n$  in the following three theorems. Moreover, we determine the structure of the extremal graphs in each case.

**Theorem 1.** *For all positive integers  $n$ , we have*

$$\text{ex}_{\text{bip}}(n, C_4, C_6) = \binom{n-2}{2},$$

*and equality holds only for  $K_{2, n-2}$ .*

**Theorem 2.** *For all positive integers  $n$ , we have*

$$\text{ex}_{\text{bip}}(n, C_6, C_8) = 6 \binom{n-3}{3},$$

*and equality holds only for  $K_{3, n-3}$ .*

**Theorem 3.** *For  $n > 3 \binom{31}{4}$ , we have*

$$\text{ex}(n, C_4, C_6) = \binom{n-2}{2} + 2,$$

*and equality holds only for a graph obtained from  $K_{2, n-2}$  by adding a single edge in the both independent sets.*

Finally, we turn our attention to exact results about the generalized extremal numbers of complete bipartite graphs  $K_{s,s}$  in graphs without a copy of  $C_{2s+2}$ . We prove the following exact result.

**Theorem 4.** *For integers  $n$  and  $s$  such that  $s \geq 3$  and  $n \geq 3s + 1 + \binom{2s+1}{s} \frac{s+1}{2}$ , we have*

$$\text{ex}(n, K_{s,s}, C_{2s+2}) = \binom{n-s}{s}.$$

*Equality holds only for graphs containing  $K_{s, n-s}$  and contained in the graph  $K_s + H$ , where  $H$  is an  $(n-s)$ -vertex graph with exactly one edge.*

The proofs of Theorems 1 and 2 are given in Section 2, and the proofs of Theorems 3 and 4 are given in Section 3.

## 2 Proofs of theorems about bipartite graphs

*Proof of Theorem 1.* Consider the graph  $K_{2,n-2}$ . It contains no cycle of length six and  $\binom{n-2}{2}$  cycles of length four. Hence we have  $\text{ex}_{\text{bip}}(n, C_4, C_6) \geq \binom{n-2}{2}$ .

Let  $G$  be an  $n$ -vertex  $C_6$ -free bipartite graph. Let  $C$  be a cycle of length four in  $G$  such that  $C = v_1v_2v_3v_4v_1$ . Then since  $G$  is a bipartite  $C_6$ -free graph, either there is no cycle of length four containing the vertices  $v_1, v_3$  distinct from  $C$  or there is no cycle of length four containing the vertices  $v_2, v_4$  distinct from  $C$ . Hence each  $C_4$  in  $G$  contains a pair of vertices from the same color class of  $G$  which is unique for that cycle.

We define an auxiliary graph  $G'$  on the same set of vertices as  $G$  by taking the edge  $uv$  in  $G'$  if and only if there is exactly one  $C_4$  in  $G$  with opposite vertices  $u$  and  $v$ . Note that since  $G$  is bipartite, the vertices  $u$  and  $v$  belong to the same color class of  $G$ . The number of copies of  $C_4$  in  $G$  is at most the number of edges in  $G'$  since each copy of  $C_4$  in  $G$  contains a pair of opposite vertices which is unique for the cycle. If the color classes of  $G$  have sizes  $a$  and  $n - a$  such that  $2 \leq a \leq \frac{n}{2}$ , then the number of edges of  $G'$  is at most  $\binom{a}{2} + \binom{n-a}{2}$ . Note that if  $a > 2$ , then  $\binom{a}{2} + \binom{n-a}{2} < \binom{n-2}{2}$  when  $n > 6$ . Therefore if  $n > 6$  and  $G$  contains at least  $\binom{n-2}{2}$  copies of  $C_4$ , then it is a bipartite graph with color classes of sizes  $n - 2$  and  $2$ , and it follows that the number of copies of  $C_4$  in  $G$  is at most  $\binom{n-2}{2}$  with equality only when  $G$  is isomorphic to  $K_{2,n-2}$ . For  $n \leq 6$  it is straightforward to verify that the theorem holds.  $\square$

*Proof of Theorem 2.* The graph  $K_{3,n-3}$  contains no cycle of length eight and contains  $6\binom{n-3}{3}$  cycles of length six. Hence we have  $\text{ex}_{\text{bip}}(n, C_6, C_8) \geq 6\binom{n-3}{3}$ .

Let  $G$  be an  $n$ -vertex bipartite graph with color classes  $V_1$  and  $V_2$ , containing no cycle of length eight and containing the maximum number of cycles of length six. Note that we may assume that  $n \geq 8$  since the theorem is simple to verify for  $n < 8$ . In what follows we are going to find a function  $\phi$  from the set of copies of  $C_6$  in  $G$  to unordered monochromatic triples of the vertices of  $G$ . Then we are going to show that the inverse image of each triple has size at most six. This implies that

$$C_6(G) < 6\binom{|V_1|}{3} + 6\binom{|V_2|}{3}.$$

If  $|V_1|, |V_2| > 3$ , then we have  $C_6(G) \leq 6\binom{|V_1|}{3} + 6\binom{|V_2|}{3} < 6\binom{n-3}{3}$  since  $n \geq 8$ , a contradiction. We may therefore assume  $|V_1| = 3$ , and it follows that  $G$  is isomorphic to  $K_{3,n-3}$  since it maximizes the number of cycles of length six. Hence we are done if such a function  $\phi$  exists.

We are going to define the function  $\phi$  recursively. Let  $C$  be a subgraph of  $G$  be isomorphic to  $C_6$ , such that  $C = v_1v_2v_3v_4v_5v_6v_1$  with color classes  $B = \{v_1, v_3, v_5\}$  and  $R = \{v_2, v_4, v_6\}$ , and assume that  $\phi(C)$  is not yet defined for  $C$ . If all of the cycles of length six containing  $B$  as a color class contain  $R$  as the other color class, then we take  $\phi(C) := B$ ; if all of the cycles of length six containing  $R$  as a color class contain  $B$  as the other color class, then we take  $\phi(C) := R$ . If both of the above occur, then we take  $\phi(C) := B$ . Otherwise, without loss of generality, we may assume there is a vertex  $v'_2$  incident to  $v_1$  and  $v_3$ , and there is a vertex  $v'_5$  incident to  $v_4$  and  $v_6$ .

Consider a subgraph  $G'$  of  $G$  containing all cycles of length six with opposite vertices  $v_1$  and  $v_4$  such that each edge of  $G'$  is in a cycle of length six with opposite vertices  $v_1$  and  $v_4$ . Let the neighborhood of  $v_1$  in  $G'$  excluding  $v_4$  (if they are incident) be denoted by  $U$ , and let the neighborhood of  $v_4$  in  $G'$  excluding  $v_1$  be denoted by  $W$ . Note that  $|U|, |W| \geq 3$  since  $\{v_2, v'_2, v_6\} \subseteq U$  and  $\{v_3, v_5, v'_5\} \subseteq W$ . Since  $G'$  is  $C_8$ -free, it is straightforward to deduce that the structure of  $G'$  is one of the following:

- If  $v_3v_6$  is an edge, then  $v_1v_4$  is also an edge since  $G'$  is  $C_8$ -free and every edge is in a copy of  $C_6$ . Then the induced graph  $G'[U \cup W]$  is a double star with central vertices  $v_3$  and  $v_6$ . Note that every cycle of length six in  $G'$  contains a vertex from  $U \setminus \{v_6\}$  and a vertex from  $W \setminus \{v_3\}$ . Even more, for each pair consisting of a vertex from  $U \setminus \{v_6\}$  and a vertex from

$W \setminus \{v_3\}$ , there are exactly two cycles of length six in  $G'$ . Hence the number of cycles of length six in  $G'$  is  $2(|U| - 1)(|W| - 1)$ .

- Otherwise  $v_3v_6$  is not an edge of  $G'$ , and the graph  $G'[U \cup W]$  is a star forest. In particular, it contains at least two stars, one with central vertex  $v_3$  and the other with central vertex  $v_6$ . Note that every  $C_6$  in  $G'$  contains  $v_1$  and  $v_4$  and at least two leaves of  $G'[U \cup W]$ . Each pair of the leaves is contained in at most one cycle of length six in  $G'$ . Hence the number of cycles of length six in  $G'$  is at most  $\binom{|U|+|W|-2}{2}$ .

Therefore we have the following bound on the number of cycles of length six in  $G'$ :

$$C_6(G') \leq \max \left( \binom{|U|+|W|-2}{2}, 2(|U| - 1)(|W| - 1) \right).$$

Let us consider the set of triples

$$X_{v_1, v_4} := \{\{v_1, w_1, w_2\} : \{w_1, w_2\} \subseteq W\} \cup \{\{v_4, u_1, u_2\} : \{u_1, u_2\} \subseteq U\}.$$

Note that  $|X_{v_1, v_4}| = \binom{|U|}{2} + \binom{|W|}{2}$ , hence  $2|X_{v_1, v_4}| > C_6(G')$ .

**Observation 5.** *There is no vertex in  $V(G - G')$  incident to two vertices of  $G'$  since  $G$  is  $C_8$ -free. Hence the only vertices adjacent to more than one vertex from  $\{v_1\} \cup W$  are  $v_4$  and  $v_6$ .*

*There is no path of length three between any two vertices of  $G'$  containing a vertex outside of  $G'$  since  $G$  is  $C_8$ -free and there is no vertex in  $V(G - G')$  incident to two vertices of  $G'$ .*

*By the previous two claims,  $v_6$  and  $w \in W$  do not appear as opposite vertices in any copy of  $C_6$  which is not contained in  $G'$ . Hence if  $X_{v_1, v_4} \cap X_{u, v} \neq \emptyset$  for some opposite pair of vertices  $u, v$ , then every six-cycle with opposite vertices  $u$  and  $v$  are six cycles of  $G'$ . Hence  $G'$  doesn't contain any 6-cycle for which  $\phi$  is already defined.*

Let us denote the set of 6-cycles in  $G'$  by  $\mathcal{C}$ . We define  $\phi$  on  $\mathcal{C}$  in the following way:  $\phi(\mathcal{C}) \subseteq X_{v_1, v_4}$  and  $|\phi^{-1}(x) \cap \mathcal{C}| \leq 2$  for every  $x \in X_{v_1, v_4}$ .

We repeat the procedure iteratively for all cycles of length six in  $G$  for which  $\phi$  is not yet defined. By Observation 5, for each triple  $\{v_1, w_1, w_2\} \in X_{v_1, v_4}$  we have  $|\phi^{-1}(\{v_1, w_1, w_2\})| \leq 2 < 6$ .  $\square$

### 3 Proofs of theorems about general graphs

#### 3.1 Maximizing cycles of length four

First we prove an essential lemma which we will need in the proof of Theorem 3.

**Lemma 6.** *Let  $G$  be a  $K_{2, s+1}$ -free and  $C_6$ -free graph for some integer  $s \geq 2$ , and let  $v \in V(G)$ . Then the number of copies of  $C_4$  incident to  $v$  is at most*

$$\max \left( 3|N_G(v)|, \frac{(s-1)(s+2)}{2(s+1)} |N_G(v)| \right).$$

*If  $G$  is also  $K_5$ -free, then the number of copies of  $C_4$  incident to  $v$  is at most*

$$\max \left( 2|N_G(v)|, \frac{(s-1)(s+2)}{2(s+1)} |N_G(v)| \right).$$

*Proof.* Let  $G'$  be the subgraph of  $G$  spanned by the edges which are in a copy of  $C_4$  incident to  $v$ . By the linearity of the desired upper bound in Lemma 6, we may assume that  $G'$  is 2-connected. Let us denote the degree of  $v$  in  $G'$  by  $x$ .

First we consider the case when  $v(G') = x + 1$ . Observe that  $G' - v$  is a connected graph since  $G'$  is 2-connected, and moreover we have that  $G'[N_{G'}(v)] = G' - v$ . Note that the number

of paths of length two in  $G' - v$  is equal to the number of copies of  $C_4$  incident to  $v$  in  $G$ . The graph  $G' - v$  is  $P_5$ -free since  $G'$  is  $C_6$ -free. Even more, we have  $\Delta(G' - v) \leq s$  since  $G$  is  $K_{2,s+1}$ -free. If  $G' - v$  is a tree, then it is either a star or a double star, and therefore the number of copies of  $C_4$  incident to  $v$  in  $G$  is at most  $\frac{(s-1)x}{2} < \frac{(s-1)(s+2)}{2(s+1)} |N_G(v)|$ . If  $G' - v$  contains a cycle of length four, then the number of copies of  $C_4$  incident to  $v$  is at most  $12 \leq 3x$ . Note that if  $G$  is  $K_5$ -free (and  $G' - v$  contains a  $C_4$ ), then the number of copies of  $C_4$  incident to  $v$  is at most  $8 \leq 2x$ . If  $G' - v$  contains a triangle and no 4-cycle, then  $G' - v$  is a graph obtained from a triangle by adding  $x - 3$  pendant edges to one of its vertices. In this case we have at most  $\binom{x-1}{2} + 2$  cycles of length 4 incident to  $v$ . Note that  $x - 1 \leq s$ , and a simple calculation shows that the required upper bound in Lemma 6 holds.

From here we will assume that  $v(G') > x + 1$ . Hence  $V(G') \setminus N_{G'}[v]$  is nonempty. Note that  $G'[V(G') \setminus N_{G'}[v]]$  is an independent set, and every vertex from  $V(G') \setminus N_{G'}[v]$  has at least two neighbors in  $N_{G'}(v)$ . Even more, if  $|V(G') \setminus N_{G'}[v]| > 1$ , then for all vertices  $u, u' \in V(G') \setminus N_{G'}[v]$  we have  $N_{G'}(u) = N_{G'}(u')$  and  $|N_{G'}(u)| = 2$  since  $G'$  is a 2-connected  $C_6$ -free graph.

If  $G'[N_{G'}(v)]$  is connected, then  $G'[N_{G'}(v)]$  is either a star or a triangle. It follows that the number of copies of  $C_4$  incident to  $v$  in  $G'$  is at most

$$\max \left( 2x, \frac{(s-1)(s+2)}{2(s+1)} x \right).$$

Note that if  $G'[N_{G'}(v)]$  is a star of  $s + 1$  vertices and  $G'[V(G') \setminus N_{G'}[v]]$  is an independent set of size  $s - 1$  incident to the same two vertices in  $G'[N_{G'}(v)]$  inducing an edge, then there are exactly  $\frac{(s-1)(s+2)}{2(s+1)} x$  copies of  $C_4$  incident to  $v$ .

If  $N_{G'}(v)$  is not connected, then  $G'[N_{G'}(v)]$  consists of isolated vertices and a star. If  $G'[V(G') \setminus N_{G'}[v]]$  consists of just a vertex, denoted by  $u$ , then in  $G'$  all of the  $y$  isolated vertices are incident to  $u$ , therefore  $y < s$  and  $x - y - 1 \leq s$ . The star  $S$  in  $G'[N_{G'}(v)]$  is isomorphic to  $S_{x-y}$ . Since every edge of  $G'$  is in a copy of  $C_4$  incident to  $v$ , the star  $S$  is not an edge. The only vertex of  $S$  incident to  $u$  is the central vertex of the star. Then the number of copies of  $C_4$  incident to  $v$  is  $\binom{y+1}{2} + \binom{x-y-1}{2} \leq \frac{(s-1)x}{2} < \max \left( 2x, \frac{(s-1)(s+2)}{2(s+1)} x \right)$  since  $x - y - 2 \leq s - 1$  and  $y \leq s - 1$ . If  $G'[V(G') \setminus N_{G'}[v]]$  contains at least two vertices, then each of the vertices have at least two neighbors in  $N_{G'}(v)$ . Hence  $G'[N_{G'}(v)]$  contains two connected components since  $G'$  is 2-connected. Therefore  $G'[N_{G'}(v)]$  consists of an isolated vertex  $w$  and a star isomorphic to  $S_{x-1}$  with central vertex  $w'$ . Note that  $S_{x-1}$  is not an edge and every vertex of  $G'[V(G') \setminus N_{G'}[v]]$  is incident to  $w$  and  $w'$ . Then the number of copies of  $C_4$  is at most  $s - 1 + \binom{x-2}{2} \leq \frac{(s-1)x}{2} < \max \left( 2x, \frac{(s-1)(s+2)}{2(s+1)} x \right)$ .  $\square$

*Proof of Theorem 3.* Let  $G$  be an  $n$ -vertex graph with no cycle of length six. Let  $K$  be a subgraph of  $G$  isomorphic to  $K_{2,s}$ , where  $s$  is the maximum integer for which  $G$  contains a copy of  $K_{2,s}$ . Let the color classes of  $K$  of size 2 and  $s$  be  $A$  and  $B$ , respectively.

**Observation 7.** For  $s > 2$ , let  $v$  and  $u$  be distinct vertices of  $V(G - K)$ . Then the following properties hold.

1. The vertex  $v$  is not adjacent to both vertices in  $A$  by the maximality of  $s$ .
2. Since  $G$  is  $C_6$ -free,  $v$  is not adjacent to more than one vertex from  $B$ .
3. By Items 1 and 2, we have that  $v$  and  $u$  are adjacent to at most 2 vertices of  $K$ . If  $v$  and  $u$  both have two neighbors in  $V(K)$ , then  $N(v) \cap V(K) = N(u) \cap V(K)$  since  $G$  is  $C_6$ -free.
4. For  $s > 3$ , the graph  $G[B]$  contains at most one edge since  $G$  is  $C_6$ -free.

In the following proof we need to classify the copies of  $C_4$  depending how they meet with  $K$ . For simplicity, we introduce the following technical definitions. If there exists a vertex  $v \in V(G - K)$  with two neighbors  $x$  and  $y$  in  $V(K)$ , then we call  $xy$  a red edge. Note that there is at most one red edge in  $G[V(K)]$  by Item 3 of Observation 7. We refer to any edge in  $G[A]$  or  $G[B]$  as a blue edge. Note that there are at most two blue edges in  $G[V(K)]$  for  $s > 3$  by Item 4 of Observation 7. .

Let  $C$  be a subgraph of  $G$  isomorphic to  $C_4$ .

- We say  $C$  is of *Type 1* if  $|V(K) \cap V(C)| = 1$ .
- We say  $C$  is of *Type 2* if  $|V(K) \cap V(C)| = 2$ . We will further divide the class of Type 2 copies of  $C_4$  into two subclasses. Each copy of  $C_4$  using a blue edge of  $G[V(K)]$  will be referred to as a *blue Type 2  $C_4$* , and each copy of  $C_4$  with two opposite vertices belonging to  $V(K)$  (thus, inducing a red edge in  $K$  as its diagonal) will be referred to as a *red Type 2  $C_4$* . Note that there is no  $C_4$  of Type 2 which is neither blue nor red.
- We say  $C$  is of *Type 3* if  $|V(K) \cap V(C)| = 3$ . Note that such a copy of  $C_4$  is incident to a blue edge, and it induces a red edge of  $K$  as its diagonal.
- We say  $C$  is of *Type 4* if  $|V(K) \cap V(C)| = 4$ .

**Observation 8.** *It is important to note that if there is a blue Type 2  $C_4$ , then  $K$  contains exactly one blue edge since  $G$  is  $C_6$ -free. Thus there are at most  $s - 1$  blue Type 2  $C_4$ 's since  $G$  is  $C_6$ - and  $K_{2,s+1}$ -free. Even more, if there is a blue edge and a red edge, then the blue edge is in the color class  $A$ . All Type 3  $C_4$ 's share the same blue edge, hence they share the same three vertices of  $K$ . Therefore the number of Type 3  $C_4$ 's is at most  $s - 1$ .*

In the following we show that for all  $n \geq 31$  either we have  $C_4(G) \leq \binom{n-2}{2} + 2$ , and equality holds if and only if  $G$  is isomorphic to  $K_{2,n-2}^+$  or  $C_4(G) < \text{ex}(n-1, C_4, C_6) + n - 3$ . For a vertex  $v$  and for a copy  $K$  of  $K_{2,s}$  with  $v \in V(K)$ , the collection of copies of  $C_4$  incident to  $v$  in  $G$  can be partitioned into five sets: Type 4, Type 3, red Type 2, blue Type 2 and Type 1  $C_4$ 's. For convenience in the following we will bound the number of copies of  $C_4$  incident to a given vertex by the sum of five terms where each term represents an upper bound on the number of  $C_4$ 's of Type 4, Type 3, red Type 2, blue Type 2 and Type 1 in this given order.

If  $s = 2$ , then for each pair of vertices there is at most one  $C_4$  containing this pair as opposite vertices. Therefore we have  $C_4(G) \leq \frac{\binom{n}{2}}{2} < \binom{n-2}{2} + 2$  for  $n > 5$ .

If  $s = 3$  and  $G$  contains a  $K_5$ , then without loss of generality we may assume  $G[V(K)]$  is isomorphic to  $K_5$ . Then every  $C_4$  incident to a vertex of  $K$  is either of Type 4 or Type 1 since  $G$  is  $C_6$ -free. Note that every vertex of  $G - K$  is adjacent to at most one vertex of  $K$ , hence by the pigeonhole principle there is a vertex  $v$  of  $K$  with at most  $\frac{n-5}{5}$  neighbors in  $V(G - K)$ . Hence by applying Lemma 6 for the vertex  $v$  in  $G[V(G - K) \cup \{v\}]$ , we have that the number of copies of  $C_4$  incident to  $v$  in  $G$  is at most

$$12 + 0 + 0 + 0 + 3 \cdot \frac{n-5}{5} < n - 3,$$

and the last inequality holds for  $n > 30$ .

If  $s = 3$  and  $G$  is  $K_5$ -free, then there are at least three vertices of  $K$  not contained in any blue Type 2  $C_4$  by Observation 8. Hence by the pigeonhole principle, there is a vertex  $v$  of  $K$  not contained in a blue Type 2  $C_4$  with at most  $\frac{n-5}{3}$  neighbors in  $V(G - K)$ . Thus by applying Lemma 6 and Observation 8 for the vertex  $v$  in  $G[V(G - K) \cup \{v\}]$ , we have that the number of copies of  $C_4$  incident to  $v$  is at most

$$12 + 2 + 1 + 0 + 2 \cdot \frac{n-5}{5} < n - 3,$$

and the last inequality holds for  $n \geq 27$ .

If  $s \geq 4$  and there is a red edge, then there is no blue edge incident to any of the vertices of  $B$ . Let us fix a vertex  $v$  in  $B$  not incident to the red edge and with a neighborhood of the smallest size in  $V(G - K)$ . Then by the pigeonhole principle, we have that  $|N_G(v) \cap V(G - K)| \leq \frac{n-s-2-1}{s-1}$  since there are  $n - s - 2$  vertices in  $G - K$  and at least one is incident to the red edge. Since there are no red or blue edges incident to the vertex  $v$ , there are no Type 2 and Type 3  $C_4$ 's incident to  $v$ . Hence we have that the number of copies of  $C_4$  incident to  $v$  is at most

$$(s-1) + 0 + 0 + 0 + \max\left(3 \cdot \frac{n-s-3}{s-1}, \frac{(s-1)(s+2)}{2(s+1)} \cdot \frac{n-s-3}{s-1}\right) < n-3$$

for all  $n > 6$ .

Assume  $4 \leq s < \frac{n}{3}$  and there is no red edge. Let us fix a vertex  $v$  in  $B$  with a neighborhood of the smallest size in  $V(G - K)$ . Then by the pigeonhole principle, we have  $|N_G(v) \cap V(G - K)| \leq \frac{n-s-2}{s}$ . Thus by applying Lemma 6 and Observation 8 for the vertex  $v$  in  $G[V(G - K) \cup \{v\}]$ , we have that the number of copies of  $C_4$  incident to  $v$  is at most

$$(s-1) + 0 + 0 + (s-1) + \max\left(3 \cdot \frac{n-s-2}{s}, \frac{(s-1)(s+2)}{2(s+1)} \cdot \frac{n-s-2}{s}\right) < n-3$$

for all  $n > 18$ .

If  $\frac{n}{3} \leq s < n-2$  and there is no red edge in  $K$ , then there is a vertex  $v$  in  $B$  not incident to a blue  $C_4$  such that it has at most one neighbor in  $V(G - K)$ . Therefore the number of  $C_4$ 's incident to  $v$  is at most  $(s-1) + 0 + 0 + 0 + 0 < n-3$ .

If  $s = n-2$ , then  $G$  contains a copy of  $K_{2,n-2}$ . In this case we have  $C_4(G) \leq \binom{n-2}{2} + 2$ , and equality holds only for a graph obtained from a graph isomorphic to  $K_{2,n-2}$  by adding an edge in each independent set.

For all  $n > 3\binom{31}{4}$ , we have either  $C_4(G) \leq \binom{n-2}{2} + 2$  and equality holds only for a graph obtained from  $K_{2,n-2}$  by adding an edge in each independent set, or

$$C_4(G) \leq 3\binom{31}{4} + \sum_{i=31}^n (i-4) < \binom{n-2}{2}. \quad \square$$

### 3.2 Maximizing complete bipartite graphs

*Proof of Theorem 4.* Since  $K_{s,n-s}$  contains  $\binom{n-s}{s}$  copies of  $K_{s,s}$ , we have  $\text{ex}(n, K_{s,s}, C_{2s+2}) \geq \binom{n-s}{s}$ . In the following we prove a matching upper bound for all integers  $n$  and  $s$  such that  $s \geq 4$  and  $n \geq 3s + 2 + \binom{2s+1}{s} \frac{s+1}{2}$ . Let  $G$  be an  $n$ -vertex  $C_{2s+2}$ -free graph with the maximum number of copies of  $K_{s,s}$ .

Let  $K$  be a subgraph of  $G$  isomorphic to  $K_{s,s}$  with color classes  $A$  and  $B$ . If there is a subgraph  $K'$  of  $G$  isomorphic to  $K_{s,s}$  with color classes  $A'$  and  $B'$  different from  $K$  such that  $A = A'$ , then there exists a vertex  $b \in B' \setminus B$  incident to all vertices of  $A$ . Similarly if there is a subgraph  $K''$  of  $G$  isomorphic to  $K_{s,s}$  with color classes  $A''$  and  $B''$  different from  $K$  such that  $B = B''$ , then there exists a vertex  $a \in A'' \setminus A$  incident to all vertices of  $B$ . Since  $G$  is  $C_{2s+2}$ -free, we have  $a = b$  and  $K$  is contained in a subgraph of  $G$  isomorphic to  $K_{s,s} + K_1$ . Thus for each subgraph  $K$  of  $G$  isomorphic to  $K_{s,s}$ , either there is a color class  $A$  of  $K$  such that the only  $K_{s,s}$  containing  $A$  as a color class is  $K$  or there is a vertex in  $G$  incident to all vertices of  $K$ . Observe that the graph  $K_{s,s} + K_1$  contains cycles of every length  $l \in \{3, 4, \dots, 2s+1\}$ .

In the following we give a classification of all subgraphs of  $G$  isomorphic to  $K_{s,s}$ .

- For each subgraph  $M$  of  $G$  such that  $M \cong K_{s,s} + K_1$ , let us label all copies of  $K_{s,s}$  which are subgraphs of  $G[V(M)]$  by  $G[V(M)]$ . We call each such label a *Type 1* label.

- For each maximal subgraph  $M$  of  $G$  such that  $M \cong K_{s,t}$  for some  $t \geq s + 2$  let us label all copies of  $K_{s,s}$  which are subgraphs of  $G[V(M)]$  by  $G[V(M)]$ . We call each such label a *Type 2* label.
- Let  $M$  be a maximal subgraph of  $G$  such that  $M \cong K_{s,t}$  for some  $t = s$  or  $t = s + 1$  and  $K_{s,s} + K_1 \not\subseteq G[V(M)]$ . If  $t = s + 1$ , let us label all copies of  $K_{s,s}$  which are subgraphs of  $G[V(M)]$  by  $G[V(M)]$ . If  $t = s$ , let  $V'$  be the set of vertices from  $V(G - M)$  incident to a vertex from both color classes of  $M$  and incident to at least two vertices from some color class of  $M$ . Note that  $|V'| \leq 1$  since  $G$  is  $C_{2s+2}$ -free. We label each copy of  $K_{s,s}$  which is a subgraph of  $G[V(M) \cup V']$  by  $G[V(M) \cup V']$ , and we call each such label a *Type 3* label.

It is easy to see that each copy of  $K_{s,s}$  is assigned at least one label. Even more, we will now show that each copy of  $K_{s,s}$  in  $G$  receives exactly one label. If a copy  $K$  of  $K_{s,s}$  has a Type 1 label  $G[V(M)]$ , then it is immediate from the definition of the labels that it has no Type 3 label. Since  $G$  is  $C_{2s+2}$ -free, there is no vertex from  $V(G - M)$  incident to at least two vertices of  $K$ . Therefore  $K$  has no Type 1 label distinct from  $G[V(M)]$  and no Type 2 label either. By the definition of Type 2 and Type 3 labels, it is impossible to have both types of labels for a copy of  $K_{s,s}$ . If a copy of  $K_{s,s}$  received a Type 2 label, then it contains a color class  $A$  such that the only  $K_{s,s}$  containing  $A$  as a color class is this  $K_{s,s}$ , hence no  $K_{s,s}$  receives two different Type 2 labels. Finally, no copy of  $K_{s,s}$  receives two distinct Type 3 labels, say  $G[V(K) \cup \{v_1\}]$  and  $G[V(K) \cup \{v_2\}]$ , for otherwise  $G[V(K) \cup \{v_1, v_2\}]$  contains a cycle of length  $2s + 2$ , a contradiction.

**Claim 9.** *Let  $K$  and  $K'$  be two copies of  $K_{s,s}$  in  $G$  with color classes  $A, B$  and  $A', B'$ , respectively, such that  $K$  and  $K'$  have different labels. Then we have  $A \not\subseteq A' \cup B'$  and  $B \not\subseteq A' \cup B'$ .*

*Proof.* Suppose by way of contradiction that  $A \subseteq A' \cup B'$ . First note that since the labels are distinct, neither  $A = A'$  nor  $A = B'$ . Hence we may assume without loss of generality that  $|A \cap A'| \geq 2$  and  $|A \cap B'| \geq 1$ . Thus there are at most  $2s + 1$  vertices in  $V(K) \cup V(K')$ , otherwise there would be at least two vertices in  $B \setminus (A' \cup B')$  incident to two vertices in the class  $A'$  and to a vertex in  $B'$ , yielding a cycle of length  $2s + 2$ . Note that  $V(K) \neq V(K')$  since they have different labels. Hence we have  $|V(K) \cup V(K')| = 2s + 1$ . We denote the only vertex in  $V(K) \setminus V(K')$  by  $v'$ . This vertex has at least two neighbors in the color class  $A'$ , which we denote by  $a_1, a_2$ , and at least one neighbor in the color class  $B'$ , which we denote by  $u$ .

Since  $K'$  has a different label than  $K$  and  $v'$  has at least two neighbors in  $A$  and a neighbor in  $B$ , there is a vertex  $v''$  from  $V((G - K') - K)$  incident to all vertices of  $A'$  or all vertices of  $B'$ . Since  $G$  is  $C_{2s+2}$ -free and  $v'$  has at least two neighbors in  $A'$ , the vertex  $v''$  is incident to all vertices of  $A'$ . Even more,  $v''$  has exactly one neighbor in  $B'$ , which was previously denoted by  $u$ . Observe that  $B \cap B'$  is nonempty; let us denote an arbitrary vertex from  $B \cap B'$  by  $u'$ . Since  $A'$  and  $(B' \setminus \{u\}) \cup \{v''\}$  induce a complete bipartite graph, there exists a cycle of length  $2s$  using the edge  $u'a_1$ ; let us denote this cycle by  $C$ . Note that there is a path  $P = u'uv'a_1$  of length three. By replacing the edge  $u'a_1$  of  $C$  with  $P$ , we obtain a cycle of length  $2s + 2$ , a contradiction.  $\square$

**Claim 10.** *Let  $K$  and  $K'$  be two copies of  $K_{s,s}$  in  $G$  with color classes  $A, B$  and  $A', B'$ , respectively, such that  $K$  and  $K'$  have different labels. Then we have either*

$$|V(K) \cap V(K')| \leq 1,$$

*or  $|V(K) \cap V(K')| = 2$  and the vertices of  $V(K) \cap V(K')$  are in the same color class in one of the copies of  $K_{s,s}$  and in different color classes in the other copy of  $K_{s,s}$ .*

*Proof.* First we prove  $|A \cap A'| \leq 1$ . Suppose by way of contradiction that  $\{a, a'\} \subseteq A \cap A'$ , where  $a \neq a'$ . Then by Claim 9, there exist vertices  $a_1$  and  $b_1$  such that  $a_1 \in A' \setminus V(K)$  and  $b_1 \in B' \setminus V(K)$ . Since  $B'$  does not contain the vertices  $a$  and  $a'$ , there is a vertex  $b_2 \in B' \setminus (A \cup \{b_1\})$ .



Note that there exists a  $2s$ -cycle  $C$  in  $K$  such that  $a$  and  $a'$  are at distance two. Even more, if  $b_2 \in B$ , then let  $C$  be a  $2s$ -cycle in which the vertex  $b_2$  is incident to  $a$  and  $a'$ . Then by changing the path of length two in  $C$  between the vertices  $a$  and  $a'$  with the path  $ab_1a_1b_2a'$ , we obtain a cycle of length  $2s + 2$ , a contradiction.

In a similar way, we have that if  $|A \cap A'| = 1$ , then  $|B \cap B'| = 0$ . This implies the claim.  $\square$

Let us fix a maximal copy  $M$  of  $K_{s,t}$  in  $G$  with color classes  $D$  of size  $s$  and  $D'$  of size  $t$ . If  $s = 3$ , we may assume that  $t > 4$ , otherwise for every pair of vertices  $u, v$  there are at most 20 copies of  $K_{3,3}$  containing  $v$  and  $u$  in the same color class by Claim 10, since the number of  $K_{3,3}$ 's in a  $K_7$  containing a fixed pair of vertices in the same color class is 20. Then

$$K_{3,3}(G) \leq \frac{20}{6} \binom{n}{2} < \binom{n-3}{3}$$

since  $n$  is sufficiently large.

In what follows, we find an injective mapping  $\phi$  from the set of copies of  $K_{s,s}$  in  $G$  to  $\binom{V(G) \setminus D}{s}$ . Finding such a mapping implies the desired upper bound.

Let  $L_1$  be a copy of  $K_{s,s} + K_1$  in  $G$ , if such a copy exists. Note that  $G[V(L_1)]$  is a Type 1 label. If  $s > 3$  and  $V(L_1) \neq V(M)$ , then let  $A'$  be a subset of  $V(L_1) \setminus M$  of size  $s - 1$ . Such a subset exists by Claim 10. If  $s > 3$  and  $V(L_1) = V(M)$ , let  $A'$  be a subset of  $V(L_1) \setminus D$  of size  $s - 1$ . Let  $X$  be a set of all copies of  $K_{s,s}$  in  $G[V(L_1)]$ . Note that  $|X| \leq K_{s,s}(K_{2s+1}) = \frac{s+1}{2} \binom{2s+1}{s}$ . Let  $Y$  be a subset of  $V(G) \setminus (V(L_1) \cup D)$  of size  $|X|$ ; such a subset exists since  $n$  is large. We define  $\phi$  on the set of copies of  $K_{s,s}$  with label  $G[V(L_1)]$  in the following way

$$\phi(X) := \{A' \cup \{v\} : v \in Y\}.$$

If  $s = 3$ , then  $V(L_1) \neq V(M)$  since  $t > 4$ . Then by Claim 10, we have  $|V(L_1) \cap D| \leq 1$  and  $|V(L_1) \cap D'| \leq 1$ . Let  $X$  be a set of all copies of  $K_{s,s}$  in  $G[V(L_1)]$ . Note that  $|X| \leq 70$ . If  $|V(L_1) \cap D| = 0$ , then let  $Y$  be a set of size  $|X|$  containing distinct triples from  $\binom{V(L_1)}{3}$  and  $\binom{V(L_1)}{2}$  with a vertex from  $D' \setminus V(L_1)$ . Note that  $|D' \setminus V(L_1)| \geq 2$  by Claim 10. Such a set  $Y$  exists since  $|D' \setminus V(L_1)| \geq 2$  and  $|X| \leq 70 < \binom{7}{3} + 2\binom{7}{2}$ . If  $|V(L_1) \cap D| = 1$ , then let  $Y$  be a set of size  $|X|$  containing distinct triples from  $\binom{V(L_1) \setminus D}{3}$  and  $\binom{V(L_1) \setminus D}{2}$  with vertices from  $D' \setminus V(L_1)$ . Note that  $|D' \setminus V(L_1)| \geq 4$  by Claim 10 and  $70 < \binom{6}{3} + 4\binom{6}{2}$ . We define  $\phi$  on the set of copies of  $K_{s,s}$  with label  $G[V(L_1)]$  by

$$\phi(X) := \{A' \cup \{v\} : v \in Y\}.$$

Let  $L_2$  be a maximal subgraph of  $G$  isomorphic to  $K_{s,t'}$  for some  $t' \geq s + 2$  with color classes  $A$  of size  $s$  and  $B$  of size  $t'$ , if such a subgraph  $L_2$  exists. Note that  $G[V(L_2)]$  is a Type 2 label. Since  $G$  is  $C_{2s+2}$ -free, each subgraph  $K$  of  $G[V(L_2)]$  isomorphic to  $K_{s,s}$  has color classes  $A$  and a subset  $B'$  of  $B$  of size  $s$ . Then by Claim 10, we have  $|B \cap D| \leq 1$ . If  $|B \cap D| = 0$ , then let  $\phi(K) := B'$  and if  $|B \cap D| \leq 1$ , then let  $\phi(K) := (B \cup \{v_{L_2}\}) \setminus D$ , where  $v_{L_2}$  is any fixed vertex of  $D' \setminus V(L_2)$ .

Let  $L_3$  be a subgraph of  $G$  such that  $G[V(L_3)]$  is a Type 3 label if such a subgraph  $L_3$  exists. We define  $\phi$  for the copies of  $K_{s,s}$  in  $G[V(L_3)]$  as we did for all copies of  $K_{s,s}$  with Type 1 label. Note that if  $s = 3$  and  $|V(L_3) \setminus D| \leq 1$  the number of  $K_{3,3}$  with such a label is at most 10, and the number of triples  $\binom{V(L_3) \setminus D}{3}$  is at least 10. Therefore for the image of  $\phi$  we do not need to use any vertex outside of this label. If  $s = 3$  and  $|V(L_3) \setminus D| = 2$  (by Claim 10 it is at most 2), then  $L_3$  induces a  $K_{3,3}$  since  $G$  is  $C_8$ -free. Therefore for the image of  $\phi$  we do not need to use any vertex outside of this label.

Let  $K$  and  $K'$  be copies of  $K_{s,s}$  in  $G$ . Note that if  $K$  and  $K'$  have the same label, then we have

$$\phi(K) \neq \phi(K').$$

If  $K$  and  $K'$  have different labels and  $s > 3$ , then since each triple  $\phi(K)$  and  $\phi(K')$  contains at most one vertex outside of its label and we have  $\phi(K) \neq \phi(K')$  by Claim 10.

If  $K$  and  $K'$  have different labels and  $s = 3$  suppose by way of contradiction that  $\phi(K) = \phi(K') = \{a, d, d'\}$ . By Claim 10 we have that  $a$  is a vertex of both labels,  $d$  is a vertex of label of  $K$ ,  $d'$  is a vertex of label of  $K'$  and  $d, d' \in D'$ . There is a path  $dPd'$  of length four going through the vertex  $a$  in  $G[V(K) \cup V(K')]$ . There is a path of length 5 in  $G[M]$  from  $d$  to  $d'$  internally disjoint from  $dPd'$  by Claim 10. These two paths constitute a cycle of length 8, a contradiction. Hence  $\phi$  is an injective function and we obtained the extremal number  $ex(n, C_6, C_8)$ .

Note that if there is more than one label, then for each label there is room to choose an extra set from  $\binom{V(G) \setminus D}{s}$  which will not be an image of  $\phi$  at the end of the procedure. Therefore equality holds only for graphs  $G$  containing exactly one label. Hence  $K_{s, n-s} \subseteq G$ . It is easy to see that  $G \subseteq K_s + H$ , where  $H$  is an  $(n - 3)$ -vertex graph with exactly one edge since  $G$  is  $C_8$ -free.  $\square$

## 4 Acknowledgements

The research of Győri and Salia was supported by the National Research, Development and Innovation Office NKFIH, grants K132696 and SNN-135643. The research of Tompkins was supported by NKFIH grant K135800.

## References

- [1] Noga Alon and Clara Shikhelman. Many  $T$  copies in  $H$ -free graphs. *Journal of Combinatorial Theory, Series B*, 121:146–172, 2016.
- [2] Béla Bollobás and Ervin Győri. Pentagons vs. triangles. *Discrete Mathematics*, 308(19):4332–4336, 2008.
- [3] Paul Erdős. On the number of complete subgraphs contained in certain graphs. *Magyar Tud. Akad. Mat. Kutató Int. Közl.*, 7(3):459–464, 1962.
- [4] Paul Erdős. On some problems in graph theory, combinatorial analysis and combinatorial number theory. *Graph theory and combinatorics (Cambridge, 1983)*, pages 1–17, 1984.
- [5] Beka Ergemlidze and Abhishek Methuku. Triangles in  $C_5$ -free graphs and hypergraphs of girth six. *arXiv preprint arXiv:1811.11873*, 2018.
- [6] Beka Ergemlidze, Abhishek Methuku, Nika Salia, and Ervin Győri. A note on the maximum number of triangles in a  $C_5$ -free graph. *Journal of Graph Theory*, 90(3):227–230, 2019.
- [7] Dániel Gerbner, Ervin Győri, Abhishek Methuku, and Máté Vizer. Generalized Turán problems for even cycles. *Journal of Combinatorial Theory, Series B*, 145:169–213, 2020.
- [8] Lior Gishboliner and Asaf Shapira. A generalized Turán problem and its applications. *International Mathematics Research Notices*, 2020(11):3417–3452, 2020.
- [9] Andrzej Grzesik. On the maximum number of five-cycles in a triangle-free graph. *Journal of Combinatorial Theory, Series B*, 102(5):1061–1066, 2012.
- [10] Andrzej Grzesik and Bartłomiej Kielak. On the maximum number of odd cycles in graphs without smaller odd cycles. *Journal of Graph Theory*, 99(2):240–246, 2022.
- [11] Ervin Győri and Hao Li. The maximum number of triangles in  $C_{2k+1}$ -free graphs. *Combinatorics, Probability and Computing*, 21(1-2):187–191, 2012.

- [12] Ervin Győri, János Pach, and Miklós Simonovits. On the maximal number of certain subgraphs in  $K_r$ -free graphs. *Graphs and Combinatorics*, 7(1):31–37, 1991.
- [13] Mateusz Górski and Andrzej Grzesik. On the maximum number of 5-cycles in graphs without 7-cycles. *Manuscript*, 2022.
- [14] Hamed Hatami, Jan Hladký, Daniel Král, Sergey Norine, and Alexander Razborov. On the number of pentagons in triangle-free graphs. *Journal of Combinatorial Theory, Series A*, 120(3):722–732, 2013.
- [15] Paul Turán. On an external problem in graph theory. *Mat. Fiz. Lapok*, 48:436–452, 1941.
- [16] Alexander Aleksandrovich Zykov. On some properties of linear complexes. *Matematicheskii sbornik*, 66(2):163–188, 1949.

*E-mail addresses:*

E. Győri: gyori.ervin@renyi.hu  
Z. He: hz18@mails.tsinghua.edu.cn  
J. Lv: lvzq19@mails.tsinghua.edu.cn  
N. Salia: nikasalia@yahoo.com  
C. Tompkins: ctompkins496@gmail.com  
K. Varga: vkitti@renyi.hu  
X. Zhu: zhuxt@smail.nju.edu.cn