

UDK 512.552+ 512.643+519.173.1+519.173.5

MSC 13A70+05C12+05C25+05C40

O. G. Styrť

# Orthogonality graphs of matrices over commutative rings

The paper is devoted to studying the orthogonality graph of the matrix ring over a commutative ring. It is proved that the orthogonality graph of the ring of matrices with size greater than 1 over a commutative ring with zero-divisors is connected and has diameter 3 or 4; a criterion for each value is obtained. It is also shown that each of its vertices has distance at most 2 from some scalar matrix.

**Key words:** associative ring with identity, commutative ring, zero-divisor, matrix ring, zero-divisor graph, orthogonality graph.

## § 1. Introduction

Researching properties of associative rings in terms of graphs of some naturally occurring algebraic binary relations takes an important place in modern mathematics. Thus, a *zero-divisor graph* was first defined in 1986 by Beck [1] for a commutative ring. Its vertices were all zero-divisors, and edges connected exactly all pairs of distinct elements giving zero in product. But since 1999 one uses its more convenient interpretation introduced by Anderson and Livingston in [2] via excluding the zero element of the ring from its vertex set. It is also proved in [2] that the zero-divisor graph of a commutative ring is connected and has diameter at most three; in the former treatment of the graph these statements would be trivial. A number of further papers also studies various characteristics of the zero-divisor graph: center and radius [8], concepts of planarity [4] and uniqueness of determining the ring by the graph up to an isomorphism [3, 5]. For non-commutative rings, there are several types of graphs defined by zero-divisors:

№	Name	Edge orientation	Vertices	Edge from $x$ to $y$	See
1)	Directed zero-divisor graph	Yes	One- and two-sided zero-divisors	$xy = 0$	[6, 7]
2)	(Undirected) zero-divisor graph	No	Nonzero one- and two-sided zero-divisors	$\begin{cases} xy = 0 \\ yx = 0 \end{cases}$	[7]
3)	Orthogonality graph	No	Nonzero two-sided zero-divisors	$\begin{cases} xy = 0 \\ yx = 0 \end{cases}$	[9, 10]

The main results for orthogonality graphs of non-commutative rings found by now concern primarily matrix rings. Thus, in the case of the basic ring being a skew field, the following properties of the orthogonality graph of the  $(n \times n)$ -matrix ring are obtained: once  $n = 2$ , it is disconnected and all its connected components have diameters at most 2, and, once  $n \geq 3$ , it is connected and has diameter 4. These statements are proved in 2014 for a field [9] and later, in 2017 — for an arbitrary skew field [10]; they can also be easily generalized to integral domains (by reducing to the field of fractions).

In this paper, there will be the orthogonality graph of the matrix ring over a commutative ring with zero-divisors studied and the following main result proved.

**Theorem 1.1.** *Let  $R$  be a commutative ring with zero-divisor set  $Z_R \neq \{0\}$ . Then, for any  $n > 1$ , the orthogonality graph of the ring of  $(n \times n)$ -matrices over  $R$  is connected and has diameter 3 or 4, the value 3 being equivalent to the relation*

$$\forall a_0 \in Z_R \quad \exists a_1, a_2 \in R \setminus \{0\} \quad \forall i, j \in \{0, 1, 2\}, i \neq j: \quad a_i a_j = 0, \quad (1.1)$$

*and each of its vertices has distance at most 2 from some scalar matrix.*

**Theorem 1.2.** *Let  $r$  be the radius of the graph under conditions of Theorem 1.1. Then*

- 1)  $2 \leq r \leq 4$ ;
- 2) if (1.1) holds, then  $r \in \{2, 3\}$ ;

3)  $r = 2$  if and only if there exists an element  $c \in R \setminus \{0\}$  such that

$$\forall a \in Z_R \quad \text{Ann}(c) \cap \text{Ann}(a) \neq 0. \quad (1.2)$$

## § 2. Auxiliary agreements

In the paper, the following notations and agreements will be used.

1) Set-theoretical:

- While listing elements of a disordered set, figured brackets are used. As for elements of an ordered tuple, they are listed in round brackets and can be repeated.
- $D^n := \underbrace{D \times \dots \times D}_n$  is the  $n$ -ary Cartesian power of a set  $D$ .

2) General algebraic:

- All rings considered are supposed to be associative and with identity.
- $R$  is an arbitrary ring.
- For any subset  $D \subset R$ , define  $D^* := D \setminus \{0\}$ . In particular, by  $R^*$  denote the subset of all nonzero (not necessarily invertible as in standard interpretation) elements of  $R$ .
- An ideal in  $R$  is *proper* if it does not equal  $R$ .
- $M_{m \times n}(R)$  is the  $R$ -module of  $(m \times n)$ -matrices over  $R$ ;  $M_n(R)$  is the ring  $M_{n \times n}(R)$ . If in the brackets the ring is replaced with some of its subsets  $D$ , then the subset of all matrices with entries from  $D$  is meant.
- $0_n^m$  is the zero  $(m \times n)$ -matrix;  $0_n := 0_n^n$ ;  $E_n$  is the identity  $(n \times n)$ -matrix;  $J_r$  is the Jordan cell of size  $r$  with eigenvalue 0. If the matrix sizes are clear from the context, then the indices can be omitted.
- $E_{kl}$  is the matrix unit  $(a_{ij})$ ,  $a_{ij} := \delta_{ki}\delta_{lj}$ .
- For a square matrix  $A$  over a commutative ring:  $\tilde{A}$  is its cofactor matrix;  $\hat{A} := (\tilde{A})^T$ .
- If  $A = (a_{k_1, k_2}) \in M_{n_1 \times n_2}(R)$ ,  $P_i \in \{1, \dots, n_i\}^{m_i}$  ( $i = 1, 2$ ), then  $A_{P_2}^{P_1}$  is the matrix  $(b_{l_1, l_2}) \in M_{m_1 \times m_2}(R)$ ,  $b_{l_1, l_2} := a_{k_1(l_1), k_2(l_2)}$ , where  $k_i(l_i)$  is the  $l_i$ -th element of  $P_i$ . If numbers are repeated neither in  $P_1$ , nor in  $P_2$ , then  $A_{P_2}^{P_1}$  is the submatrix of  $A$  with row and column numbers from  $P_1$  and  $P_2$  respectively.

3) On zero-divisor types:

- An element  $a \in R$  is called
  - a *left* (resp. *right*) *zero-divisor* if there exists an element  $b \in R^*$  such that  $ab = 0$  (resp.  $ba = 0$ );
  - a *zero-divisor* if it is either left or right zero-divisor;
  - a *two-sided zero-divisor* if it is both left and right zero-divisor.

At that,

- in a commutative ring, the concepts of all zero-divisor types are equivalent;
  - zero is a two-sided zero-divisor; if there are no other zero-divisors, then  $R$  is called a *ring without zero-divisors*.
  - An *integral domain* is a commutative ring without zero-divisors.
- 4) From general graph theory:
- All graphs considered are assumed to be undirected.
  - $\Gamma = (V, E)$  is an arbitrary graph;  $V$  and  $E$  are its vertex and edge sets respectively. In doing so, one can (usually with more convenience) define  $E$  via a symmetric binary relation on  $V$ .
  - Two vertices are *adjacent* if they are connected with an edge.
  - A *subgraph* is a graph with vertex set  $V' \subset V$  and, unless otherwise stated, with the same binary relation restricted on  $V'$ .
  - A *path* is a sequence of vertices where any two neighbor ones are adjacent.
  - The *length* of a path is the number of its edges.
  - The *distance* between vertices  $v$  and  $w$  (not.  $d(v, w)$ ) is the minimum of lengths of paths between them; if they do not exist, then set  $d(v, w) := +\infty$ ; the sign is obvious in this context and therefore will be omitted. Clearly,  $(d(v, w) = 0) \Leftrightarrow (v = w)$ .
  - The *distance* from a vertex  $v$  to a subset  $W \subset V$  (not.  $d(v, W)$ ) is the number<sup>\*</sup>

$$\min\{d(v, w) : w \in W\}.$$
  - $d(v) := \sup\{d(v, w) : w \in W\}$  ( $v \in V$ ).
  - The *diameter* of  $\Gamma$  is the number<sup>\*</sup>

$$\text{diam}(\Gamma) := \sup\{d(v, w) : v, w \in V\} = \max\{d(v) : v \in W\}.$$
  - 
  - The *radius* of  $\Gamma$  is the number<sup>\*</sup>  $\text{rad}(\Gamma) := \min\{d(v) : v \in W\}$ . Clearly,
$$\text{rad}(\Gamma) \leq \text{diam}(\Gamma) \leq 2 \cdot \text{rad}(\Gamma). \quad (2.1)$$
  - A graph is *connected* if there exists a path between any two of its vertices.
- Remark.** It is easy to see that a graph with finite diameter is connected. The converse fails; an example is the set of positive integers with the neighborhood relation.
- 5) On special graphs in algebraic structures:
- $O(R)$  is the orthogonality graph of the ring  $R$  (for a commutative ring it is the same as the zero-divisor graph).
  - Vertices of  $O(R)$  are all nonzero two-sided zero-divisors of  $R$ ; the orthogonality relation  $(xy = yx = 0)$  is written as  $(x \perp y)$ ;  $O_R(x)$  is the set of all vertices orthogonal to  $x$ .

---

<sup>\*</sup>Possibly  $\infty$ .

### § 3. Proofs of the results

Consider an arbitrary commutative ring  $R$ . Denote by  $\text{Ann}(a)$  ( $a \in R$ ) the ideal  $\{x \in R: ax = 0\}$  and by  $Z_R$  the set  $\{a \in R: \text{Ann}(a) \neq 0\}$  of all zero-divisors. Further, let  $S$  be the ring  $M_n(R)$  ( $n > 1$ ). Via the natural ring embedding  $R \hookrightarrow S$ ,  $a \rightarrow aE$ , identify  $R$  with the subring  $RE \subset S$  (and, thus,  $O(R)$  — with a subgraph of the graph  $O(S)$ ). For  $A \in S$ , set  $I_A := \text{Ann}(\det A) \triangleleft R$ .

The graph  $O(R)$  is connected and has diameter at most 3 (see Theorem 2.3 in [2, §2]). Besides, if  $R$  is a skew body, then

- 1) once  $n = 2$ , the graph  $O(S)$  is disconnected and all its connected components have diameters  $\leq 2$ ;
- 2) once  $n \geq 3$ , the graph  $O(S)$  is connected and has diameter 4.

These results are obtained in [9, §4] for fields (Lemma 4.1 and Theorem 4.5 respectively), and in [10, §2] are generalized to arbitrary skew-fields (Lemma 2.2 and Theorem 2.1 respectively). They are also shifted to integral domains (by reducing to the field of fractions).

**Theorem 3.1.** *For any matrix  $A \in S$  and proper ideal  $I \triangleleft R$  containing  $\det A$ , there exists a matrix  $B \in S \setminus (M_n(I))$  such that  $AB, BA \in M_n(I)$ .*

□ For  $m \in \mathbb{N}$ , set  $Q_m := \{1, \dots, m\}$  and  $P_m := (1, \dots, m) \in \mathbb{N}^m$ .

Consider all triples  $(k, P', P'')$  ( $k \geq 0$ ,  $P', P'' \in (Q_n)^k$ ) satisfying the relation  $\det(A_{P''}^{P'}) \notin I$ . For each of them, numbers are repeated neither in  $P_1$ , nor in  $P_2$ , and, by condition,  $k < n$ . Besides, at least one of such triples exists: for  $k := 0$  and empty tuples  $P', P''$ , the corresponding  $(0 \times 0)$ -matrix has determinant  $1 \notin I$ . Hence, we can fix one of these triples with the largest possible  $k$ , and then  $0 \leq k < n$ ,  $m := k + 1 \in Q_n$ .

**Case 1).**  $P' = P'' = P_k$ .

By construction,  $\det(A_{P_k}^{P_k}) \notin I$ . Further, set  $C := A_{P_m}^{P_m} \in M_m(R)$ ,

$$B := \begin{pmatrix} \hat{C} & 0_{n-m}^m \\ 0_m^{n-m} & 0_{n-m} \end{pmatrix} \in S.$$

Then  $b_{m,m} = \det(A_{P_k}^{P_k}) \notin I$  implying  $B \notin M_n(I)$ . Show that  $AB, A^T B^T \in M_n(I)$ , i. e. that, for any  $p, q \in Q_n$ , the matrix entries  $(AB)_{p,q}$  and  $(A^T B^T)_{p,q}$  belong to  $I$ . Assume that  $p \in Q_n$  and  $q \in Q_m$  (otherwise  $(AB)_{p,q} = (A^T B^T)_{p,q} = 0$ ). Let  $P \in (Q_n)^m$  be the tuple obtained from  $P_m$  by changing the  $q$ -th element with  $p$ . Due to maximality of  $k$  and the inequality  $m > k$ , we have  $\det(A_P^P), \det(A_{P^m}^P) \in I$ ,

$$\begin{aligned} (AB)_{p,q} &= \sum_{i \in Q_n} (a_{p,i} b_{i,q}) = \sum_{i \in Q_m} (a_{p,i} (\hat{C})_{i,q}) = \sum_{i \in Q_m} (a_{p,i} (\tilde{C})_{q,i}) = \det(A_{P^m}^P) \in I; \\ (A^T B^T)_{p,q} &= \sum_{i \in Q_n} ((A^T)_{p,i} (B^T)_{i,q}) = \sum_{i \in Q_m} (a_{i,p} (\tilde{C})_{i,q}) = \det(A_P^{P^m}) \in I. \end{aligned}$$

Thereby, it is proved that  $AB, (BA)^T = A^T B^T \in M_n(I)$  implying  $BA \in M_n(I)$ .

**Case 2).**  $P', P'' \in (Q_n)^k$  are arbitrary tuples.

In each of the tuples  $P'$  and  $P''$  all numbers are distinct. Hence, via suitable permutations of rows and columns, one can obtain from  $A$  a matrix  $A_0$  satisfying Case 1) with the same  $k$ . By proved above, there exists a matrix  $B_0 \in S \setminus (M_n(I))$  such that  $A_0 B_0, B_0 A_0 \in M_n(I)$ . At that, there exist monomial (therefore, invertible) matrices  $C_1, C_2 \in S$  such that  $A = C_1 A_0 C_2^{-1}$ . Left (resp. right) multiplying a matrix by a monomial one permutes its rows (resp. columns), and, consequently,  $B := C_2 B_0 C_1^{-1} \in S \setminus (M_n(I))$ ,  $AB = C_1 (A_0 B_0) C_1^{-1} \in M_n(I)$ ,  $BA = C_2 (B_0 A_0) C_2^{-1} \in M_n(I)$ . ■

**Corollary 3.1.** *If  $A \in S$  and  $c \in I_A^*$ , then, in the subset  $(cS)^* \subset S$ , there exists an element orthogonal to  $A$ .*

□ By condition,  $I := \text{Ann}(c) \triangleleft R$  is a proper ideal containing  $\det A$ . According to Theorem 3.1, there exists a matrix  $B \in S \setminus (M_n(I))$  such that  $AB, BA \in M_n(I)$ . Thus,  $C := cB \neq 0$  and  $c(AB) = c(BA) = 0$ , i. e.  $C \in (cS)^*$  and  $AC = CA = 0$ . ■

**Lemma 3.1.** *For any  $A \in S$ , the following conditions are equivalent:*

- 1)  $\det A \in Z_R$ ;
- 2)  $I_A \neq 0$ ;
- 3) in  $S^*$ , there exists an element orthogonal to  $A$ ;
- 4)  $A$  is a two-sided zero-divisor;
- 5)  $A$  is a zero-divisor.

□ The implications 1)  $\Leftrightarrow$  2) and 3)  $\Rightarrow$  4)  $\Rightarrow$  5) obviously follow from definitions, and the implication 2)  $\Rightarrow$  3) — from Corollary 3.1.

Prove the implication 5)  $\Rightarrow$  1). Suppose that, without loss of generality,  $A$  is a left zero-divisor, i. e. that  $AB = 0$  for some  $B \in S^*$ . Then  $\hat{A}A = (\det A)E$  implying  $(\det A)B = \hat{A}AB = 0$ . It remains to use non-triviality of  $B$ . ■

**Corollary 3.2.** *All zero-divisors in  $S$  are two-sided.*

Let  $Z_S \subset S$  be the subset of all elements  $A \in S$  satisfying each of the equivalent conditions 1)–5) of Lemma 3.1, i. e. the set of all zero-divisors of the ring  $S$ . Then the vertex set of the graph  $O(S)$  is  $Z_S^*$ .

Further, we will assume that  $Z_R^* \neq \emptyset$ .

**Statement 3.1.** *If  $I \triangleleft R$  and  $I \neq 0$ , then  $Z_R \cap I \neq \{0\}$ .*

□ Suppose that  $Z_R \cap I = \{0\}$ . There exist elements  $b \in I^*$  and  $c \in Z_R^*$ ; then  $bc \in Z_R \cap I = \{0\}$ . So,  $bc = 0 \neq c$  that implies  $b \in Z_R \cap I^* = \emptyset$ , a contradiction. ■

**Lemma 3.2.** *If, for a subset  $D \subset S$ , the ideal  $I := \bigcap_{A \in D} I_A \triangleleft R$  is nonzero, then there exist elements  $b \in Z_R^*$  and  $C_A \in S^*$ ,  $A \in D$ , such that  $bE \perp C_A \perp A$  ( $A \in D$ ).*

□ According to Statement 3.1, the ideal  $I$  contains an element  $c \in Z_R^*$ . Then  $bc = 0$  where  $b \in Z_R^*$ . Further, for any  $A \in D$ , we have  $c \in I_A^*$  and, by Corollary 3.1, there exist an element  $C_A \in (cS)^*$  orthogonal to  $A$ ; at that,  $bC_A \in bcS = 0$ ,  $bE \perp C_A$ . ■

**Corollary 3.3.**

- 1) For any  $A \in Z_S^*$ , we have  $d(A, O(R)) \leq 2$ .
- 2) If  $A_1, A_2 \in Z_S^*$  and  $I_{A_1} \cap I_{A_2} \neq 0$ , then  $d(A_1, A_2) \leq 4$ .

□ It suffices to apply Lemma 3.2 to the subsets  $\{A\}, \{A_1, A_2\} \subset S$ . ■

**Lemma 3.3.** If  $A_i \in Z_S^*$ ,  $c_i \in I_{A_i}^*$  ( $i = 1, 2$ ) and  $c_1c_2 = 0$ , then  $d(A_1, A_2) \leq 3$ .

□ By Corollary 3.1, for each  $i = 1, 2$ , there exists an element  $C_i \in (c_iS)^*$  such that  $C_i \perp A_i$ . In this case,  $C_1C_2, C_2C_1 \in c_1c_2S = 0$ ,  $C_1 \perp C_2$ . ■

**Definition.** We will say that an ideal  $I \triangleleft R$  does not have zero-divisors if  $I^*I^* \not\cong 0$ , i. e. if the ring\*  $I$  does not have zero-divisors.

**Lemma 3.4.** If  $A_1, A_2 \in Z_S^*$  and  $d(A_1, A_2) > 3$ , then  $I_{A_i}$  ( $i = 1, 2$ ) is the same ideal without zero-divisors.

□ According to Lemma 3.3,  $I_{A_1}^* I_{A_2}^* \not\cong 0$ . It remains to prove that  $I_{A_1} = I_{A_2}$ .

Suppose that  $I_{A_1} \neq I_{A_2}$ . Without loss of generality, assume that there exists an element  $c \in I_{A_1} \setminus I_{A_2}$ . Setting  $a := \det A_2$ , we have  $I_{A_2} = \text{Ann}(a)$ ,  $b := ca \in I_{A_1}^*$  and  $bI_{A_2} = caI_{A_2} = 0$  implying  $bI_{A_2}^* \subset \{0\} \cap (I_{A_1}^* I_{A_2}^*) = \emptyset$ ,  $I_{A_2}^* = \emptyset$ ,  $I_{A_2} = 0$ , a contradiction. ■

**Theorem 3.2.** The graph  $O(S)$  is connected and has diameter at most 4.

□ Suppose that there exist elements  $A_1, A_2 \in Z_S^*$  satisfying the inequality  $d(A_1, A_2) > 4$ . By Lemma 3.4,  $0 \neq I_{A_1} = I_{A_2} = I_{A_1} \cap I_{A_2}$  that contradicts with Corollary 3.3. ■

**Theorem 3.3.** We have  $\text{diam}(O(S)) \geq 3$ , the strict inequality being equivalent to the existence of an ideal  $\text{Ann}(a) \triangleleft R$  ( $a \in Z_R$ ) without zero-divisors.

□ Similarly with examples from [9, 10] giving lower estimates of the diameter, for an arbitrary  $a \in Z_R$ , set  $I := \text{Ann}(a) \triangleleft R$  and  $A := J_n + aE_{n1} \in S$ . Note that

- $A, A^T \in Z_S^*$ ,  $O_S(A) = I^*E_{1n}$ ,  $O_S(A^T) = I^*E_{n1}$ ;
- $a_{12} = 1 \neq a_{21}$ ,  $(AA^T)_{11} = 1$  and  $O_S(A) \cap O_S(A^T) = \emptyset$ , that implies  $d(A, A^T) \geq 3$ ;
- if  $I^*I^* \not\cong 0$ , then  $(O_S(A))(O_S(A^T)) = (I^*I^*)E_{11} \not\cong 0$  and, hence,  $d(A, A^T) \geq 4$ .

Due to mentioned above,  $\text{diam}(O(S)) \geq 3$ , the strict inequality following from the existence of an ideal  $\text{Ann}(a) \triangleleft R$  ( $a \in Z_R$ ) without zero-divisors. Conversely, in the case of the strict inequality, by Lemma 3.4, for some elements  $A \in Z_S$  and  $a := \det A \in Z_R$ , the ideal  $I_A = \text{Ann}(a) \triangleleft R$  does not have zero-divisors. ■

Now the main Theorem 1.1 follows from Theorems 3.2 and 3.3, and Corollary 3.3. It implies (see (2.1)) the statements 1) and 2) of Theorem 1.2. Let us prove 3).

---

\*In general, without identity.

Suppose that  $\text{rad}(O(S)) = 2$ . There exist elements  $C \in Z_S^*$ ,  $c \in R^*$  and  $k, l \in Q_n$  such that  $d(C, A) \leq 2$  ( $A \in Z_S^*$ ) and  $c_{kl} = c$ . Further, there exists a permutation  $\sigma \in S_n$  such that  $m := \sigma(k) \neq l$ .

Let  $a \in Z_R$  be an arbitrary element.

Set  $I := \text{Ann}(a) \triangleleft R$  and  $A := \left( \sum_{i \neq k} E_{i, \sigma(i)} \right) + aE_{km} \in S$ . Note that

- $A \in Z_S^*$ ,  $O_S(A) = I^*E_{mk}$ ;
- $A \neq C$  (otherwise  $a_{kl} = c \neq 0$ ,  $m = l$ );
- $(m, k) \neq (k, l)$  (otherwise  $m = k = l$ ), that implies  $C \notin O_S(A)$ .

Thus,  $d(C, A) = 2$ , so, there exists an element  $B \in Z_S^*$  orthogonal to  $C$  and  $A$ . We have  $B = bE_{mk}$  where  $b \in I^*$ . Meanwhile,  $BC = 0$ ,  $0 = (BC)_{ml} = bc$ ,  $b \in \text{Ann}(c) \cap I^*$ .

Due to arbitrariness of  $a \in Z_R$ , the element  $c \in R^*$  satisfies (1.2).

Conversely, assume that (1.2) holds for some  $c \in R^*$ . Show that the element  $C := cE \in S^*$  satisfies, for each  $A \in Z_S^*$ , the inequality  $d(C, A) \leq 2$ .

Let  $A \in Z_S^*$  be an arbitrary element. Then  $\det A \in Z_R$ , and, by (1.2), there exists an element  $b \in I_A^*$  such that  $cb = 0$ . Further, according to Corollary 3.1, there exists an element  $B \in (bS)^*$  orthogonal to  $A$ ; in this case,  $cB \in cbS = 0$ ,  $C \in Z_S^*$ ,  $C \perp B \perp A$ ,  $d(C, A) \leq 2$ .

So, Theorem 1.2 is completely proved.

## Acknowledgements

The author is grateful to Prof. E. B. Vinberg for exciting interest to algebra.

The author dedicates the article to E. N. Troshina.

## References

- [1] Beck I. Coloring of commutative rings // J. Algebra. 1988. Vol. 116. Pp. 208–226.
- [2] Anderson D. F., Livingston P. S. The zero-divisor graph of a commutative ring // J. Algebra. 1999. Vol. 217. Pp. 434–447.
- [3] Anderson D. F., Frazier A., Lauve A., Livingston P. S. The zero-divisor graph of a commutative ring, II // Lect. Notes Pure Appl. Math. 2001. Vol. 220, Marcel Dekker, New York. Pp. 61–72.
- [4] Akbari S., Maimani H. R., Yassemi S. When zero-divisor graph is planar or a complete  $r$ -partite graph // J. Algebra. 2003. Vol. 270. Pp. 169–180.



- [5] Akbari S., Mohammadian A. On the zero-divisor graph of a commutative ring // J. Algebra. 2004. Vol. 274. Pp. 847–855.
- [6] Akbari S., Mohammadian A. Zero-divisor graphs of non-commutative rings // J. Algebra. 2006. Vol. 296. Pp. 462–479.
- [7] Akbari S., Mohammadian A. On zero-divisor graphs of finite rings // J. Algebra. 2007. Vol. 314. Pp. 168–184.
- [8] Redmond S. P. Central sets and radii of the zero-divisor graphs of commutative rings // Comm. in Algebra. 2006. Vol. 34 (is. 7). Pp. 2389–2401.
- [9] Bakhadly B. R., Guterman A. E., Markova O. V. Graphs defined by orthogonality // Computational methods and algorithms. Part XXVII, Zap. Nauchn. Sem. POMI, St. Petersburg. 2014. Vol. 428. Pp. 49–80; J. Math. Sci. 2015. Vol. 207 (is. 5). Pp. 698–717.
- [10] Guterman A. E., Markova O. V. Orthogonality graphs of matrices over skew fields // Computational methods and algorithms. Part XXX, Zap. Nauchn. Sem. POMI, St. Petersburg. 2017. Vol. 463. Pp. 81–93; J. Math. Sci. 2018. Vol. 232 (is. 6). Pp. 797–804.