

Search for Coincident Gravitational Wave and Long Gamma-Ray Bursts from 4-OGC and the Fermi/Swift Catalog

YI-FAN WANG (王一帆)^{1,2} ALEXANDER H. NITZ^{1,2} COLLIN D. CAPANO^{3,1,2} XIANGYU IVY WANG^{4,5} YU-HAN YANG^{4,5}
AND BIN-BIN ZHANG^{4,5}

¹*Max-Planck-Institut für Gravitationsphysik (Albert-Einstein-Institut), D-30167 Hannover, Germany*

²*Leibniz Universität Hannover, D-30167 Hannover, Germany*

³*Department of Mathematics, University of Massachusetts, Dartmouth, MA 02747, USA*

⁴*School of Astronomy and Space Science, Nanjing University, Nanjing 210093, China*

⁵*Key Laboratory of Modern Astronomy and Astrophysics (Nanjing University), Ministry of Education, China*

ABSTRACT

The recent discovery of a kilonova associated with an apparent long-duration gamma-ray burst has challenged the typical classification that long gamma-ray bursts originate from the core collapse of massive stars and short gamma-ray bursts are from compact binary coalescence. The kilonova indicates a neutron star merger origin and suggests the viability of gravitational-wave and long gamma-ray burst multimessenger astronomy. Gravitational waves play a crucial role by providing independent information for the source properties. This work revisits the archival 2015-2020 LIGO/Virgo gravitational-wave candidates from the 4-OGC catalog which are consistent with a binary neutron star or neutron star-black hole merger and the long-duration gamma-ray bursts from the Fermi and Swift catalogs. We search for spatial and temporal coincidence with up to 10 s time lag between gravitational-wave candidates and the onset of long-duration GRBs. The most significant candidate association has only a false alarm rate of once every two years; given the LIGO/Virgo observational period, this is consistent with a null result. We report an exclusion distance for each search candidate for a fiducial gravitational-wave signal and conservative viewing angle assumptions.

Keywords: gravitational wave — gamma-ray burst — binary neutron star — neutron star-black hole

1. INTRODUCTION

The detection of a gravitational wave (GW) from the binary neutron star merger GW170817 together with an electromagnetic counterpart began a new era of multimessenger astronomy (Abbott et al. 2017a,b,c). The initial report of the GW by Advanced LIGO (Aasi et al. 2015) and Virgo (Acernese et al. 2015) and the short gamma-ray burst (GRB) by the Fermi Gamma-ray Burst Monitor (Meegan et al. 2009) triggered an extensive campaign of follow-up observations across the electromagnetic (EM) spectrum. This multimessenger event provided a great wealth of knowledge for astrophysics and fundamental physics. For example, the speed of GW propagation was constrained to deviate by no more than a factor of $[-3 \times 10^{-15}, 7 \times 10^{-16}]$ from the speed of light (Abbott et al. 2017b, 2019a). Be-

sides, direct measurement of the Hubble constant was found to be $H_0 = 70.0_{-8.0}^{+12.0}$ km s⁻¹ Mpc⁻¹ within the 68% credible interval (Abbott et al. 2017d) (see also e.g., Guidorzi et al. 2017; Hotokezaka et al. 2019, for later improvements). It also confirmed the long hypothesis that short-duration GRBs originate from the merger of compact objects involving neutron stars (Goodman 1986; Paczynski 1986; Eichler et al. 1989; Narayan et al. 1992), and provided the first confirmation for a kilonova associated with a neutron star merger and r-process nucleosynthesis (e.g., Abbott et al. 2017e).

The correlation analysis of signals from different observations underlies GW/EM multimessenger astronomy. Since the beginning of the third observational run in April 2019, LIGO/Virgo have released real-time public trigger alerts (Abbott et al. 2021a), enabling swift follow-up observation of EM-band or neutrino signals. Later, archival GW/EM searches are performed to dig deeper for associated signals missed by the low-latency searches. For instance, Abbott et al. (2017f, 2019b,

2021b, 2022) and Nitz et al. (2019) have studied temporal and spatial coincidence between GW and GRB using archival LIGO/Virgo data and Fermi/Swift (Meegan et al. 2009; Gehrels et al. 2004) data. The LIGO Scientific Collaboration et al. (2022) and Wang & Nitz (2022) have searched for GWs coincident with observations from the fast radio burst catalog released by the CHIME collaboration (Amiri et al. 2018, 2021).

Conventionally, GRBs are separated into two classes based on their duration and spectral characteristics, i.e., short-duration hard-spectral and long-duration soft-spectral (Dezalay et al. 1992; Kouveliotou et al. 1993). As confirmed by GW170817, short GRBs are known to originate from neutron star mergers, while many long GRBs are generated by the core collapse of massive stars as determined by the co-observation of supernova (Galama et al. 1998; Hjorth et al. 2003; Stanek et al. 2003). Based on this knowledge, previous searches (Abbott et al. 2017f, 2019b, 2021b, 2022; Nitz et al. 2019) for GW/GRB multimessenger have only targeted short GRBs with a template-based match-filtering method to look for associated GW signals from compact binary coalescence. The GW candidates potentially associated with long GRBs were analyzed with a template-free generic search, with the aim to detect GWs from asymmetric core-collapse massive stars.

However, the separation of GRBs based on their observed light-curve duration is not precise (Gehrels et al. 2006; Gal-Yam et al. 2006; Zhang et al. 2021; Ahumada et al. 2021). In particular, Rastinejad et al. (2022) recently reported a kilonova located at 350 Mpc associated with GRB 211211A, an apparent long-duration GRB with $T_{90} = 51.37$ s given by Swift/BAT (2018), challenging the previous paradigm for GRB classification and progenitor. Additional interesting features were reported, including a quasi-periodic oscillated precursor that occurred ~ 1 s prior to the main emission (Xiao et al. 2022), and a high energy (> 100 MeV) gamma-ray afterglow lasting $\sim 2 \times 10^4$ s that occurred ~ 1000 s after the burst (Mei et al. 2022; Zhang et al. 2022). Several models have been proposed to explain the emission mechanism of GRB 211211A; for instance, fast-cooling synchrotron radiation after a binary neutron star merger (Gompertz et al. 2022), magnetic barrier effect involving a magnetar progenitor (Gao et al. 2022), a neutron star-white dwarf merger (Yang et al. 2022), or thermal emission from heated dust as an alternative scenario to kilonova (Waxman et al. 2022). The associated kilonova clearly indicates the progenitor of GRB 211211A is a binary merger with at least one neutron star, suggesting the prospect of detecting a GW signal from events of this kind (however Waxman et al. 2022, for an alter-

native explanation). GW observation provides unique information about the source’s mass and spin (Veitch et al. 2015), which may help disentangle different models (Gompertz et al. 2022; Gao et al. 2022; Yang et al. 2022; Waxman et al. 2022). Unfortunately, none of the GW detectors were in observation mode at the time of the GRB 211211A.

Nevertheless, we revisit the archival GW data from LIGO/Virgo and long GRB candidates from the Fermi (Meegan et al. 2009) and Swift (Gehrels et al. 2004) observation motivated by the kilonova/GRB 211211A association. We focus on GW candidates, both significant and subthreshold, that are consistent with mergers involving at least one neutron star and study their temporal and spatial coincidence with confident long GRB observations recorded by Fermi and Swift.

2. CANDIDATE SELECTION AND RANKING STATISTIC

This section describes the selection of GW and long GRB candidates, and our algorithm to rank the GW/long-GRB temporal and spatial coincidence.

Advanced LIGO and Virgo have completed three observational runs from 2015 to 2020 and released around ninety GW events in the Gravitational Wave Transient Catalog (Abbott et al. 2021c). Additional detections were reported by independent groups (Nitz et al. 2021; Olsen et al. 2022). This work utilizes the GW search results from the fourth Open Gravitational-wave Catalog (4-OGC, Nitz et al. 2021). 4-OGC has searched for the entire three observation runs of Advanced LIGO/Virgo using the PyCBC toolkit (Nitz et al. 2018) and detected ninety-four confident GW events from compact binary coalescences. The subthreshold triggers’ search results are also publicly released (Nitz & Kumar 2021).

To select potential GW signals comprising at least one neutron star, we choose those confident and subthreshold candidates in 4-OGC where the secondary mass m_2 (the mass of the lighter component of the binary) of the associated gravitational-wave template is within $[1, 3] M_\odot$. Given the observation of low-spin neutron star – black hole mergers (Abbott et al. 2021d), we also extend the mass range up to total mass $m_1 + m_2$ of $10 M_\odot$. Since we primarily target EM counterpart from binary neutron star mergers, we constrain the effective spin χ_{eff} in $[-0.2, 0.2]$. Effective spin is the primary spin parameter characterizing the GW signal (Ajith et al. 2011) and is defined as

$$\chi_{\text{eff}} = \frac{m_1 \chi_1 + m_2 \chi_2}{m_1 + m_2}, \quad (1)$$

where $\chi_{1/2}$ is the dimensionless component spin aligned with the orbital angular momentum direction. We

limit χ_{eff} because observations show that galactic binary neutron stars would have small spin ($\chi_{\text{eff}} \lesssim 0.05$) at merger (Zhu et al. 2018); the binary neutron star events GW170817 and GW190425 are consistent with zero spin (Abbott et al. 2019c, 2020). Also, note that $[1,2] M_{\odot}$ is considered the binary neutron star search region in 4-OGC, and χ_{eff} in the search template is constrained to $[-0.05, 0.05]$. While we expect high-spin black hole – neutron star mergers to experience greater disruption (Foucart et al. 2018; Capano et al. 2020), our candidate population may have significant observational biases if the spin-axis is misaligned with the orbital axis (Dhurkunde & Nitz 2022), which can be improved in future GW search by accounting for spin precession. We also expect that for neutron star – black hole mergers with a mass ratio greater than $\sim 10:1$ there is not likely to be sufficient mass ejecta to produce a GRB (Foucart et al. 2018; Capano et al. 2020).

We further require the candidates to trigger at least two GW detectors, i.e., no single detector triggers are considered (Nitz et al. 2020). The 4-OGC catalog assigns a search ranking statistic, λ_{gw} , to each GW trigger, which is the natural logarithm of the ratio of rate densities for signal and noise hypothesis (Nitz et al. 2017; Davies et al. 2020). We only consider those triggers with $\lambda_{\text{gw}} \geq 0$. These choices resulted in $\sim 5 \times 10^6$ triggers from the three observational runs in 2015-2022.

For each GW candidate, we use **PyCBC Inference** (Biwer et al. 2019) to estimate the Bayesian posterior of the sky localization. To save the computation resources, we fix the mass and spin parameters to be those reported by 4-OGC (Nitz & Kumar 2021) from the PyCBC search. We do not expect this procedure would significantly bias the sky position estimation because of the decoupling of intrinsic parameters (the source parameters independent of the observer orientation) and extrinsic parameters (the source parameters dependent on the observer orientation) (Singer & Price 2016). Therefore, the variables to be estimated include the luminosity distance, right ascension, declination, polarization angle, inclination angle between the source orbital angular momentum and the line of sight of observatories, and the phase and time of coalescence. The posterior is numerically estimated using the dynamical nested sampler *dynesty* (Speagle 2020) using the standard GW likelihood Finn (1992) assuming stationary and Gaussian noise.

For long GRBs, we select those from the Fermi/GBM and Swift/BAT, with $T_{90} - \delta T_{90} > 4\text{s}$, where T_{90} and δT_{90} are the time duration containing 90% of the burst fluence and its associated error, respectively. The sky maps of these long-GRBs are released by Fermi and

Swift, respectively (Meegan et al. 2009; Gehrels et al. 2004).

We describe the algorithm to quantify the temporal and spatial coincidence of GW and long GRB. The time lag between GW and long GRB is allowed to be within $[0, 10]$ s. GW/long GRB pairs which meet this condition are considered temporally associated. We further compute the sky overlap probability for the associated pairs using the posterior overlap integral following Eq. 11 in Ashton et al. (2018) as

$$\mathcal{B}_{\text{overlap}} = \int \frac{P(\vec{\theta}|d_{\text{GW}}, \mathcal{H})P(\vec{\theta}|d_{\text{GRB}}, \mathcal{H})}{P(\vec{\theta}|\mathcal{H})} d\vec{\theta}, \quad (2)$$

where $\vec{\theta}$ is the sky localization (right ascension and declination), $P(\vec{\theta}|d_{\text{GW}}, \mathcal{H})$ and $P(\vec{\theta}|d_{\text{GRB}}, \mathcal{H})$ are the sky location posterior estimated by GW and GRB data, respectively, $P(\vec{\theta}|\mathcal{H})$ is the prior that we choose to be isotropic. \mathcal{H} is the underlying hypothesis characterizing the GW and GRB data analysis.

Given the search ranking statistic λ_{gw} from PyCBC, we design the following ranking statistic for GW/long-GRB association

$$\lambda_{\text{gw+lgrb}} = \lambda_{\text{gw}} + \ln \mathcal{B}_{\text{overlap}}. \quad (3)$$

To measure the statistical significance for a specific associate GW/long-GRB pair, we time shift the GRB data with respect to GW data with a time stride larger than 10 s (we use 200 s) and calculate the ranking statistics of the new associated pairs. Any coincidence obtained after the time-shifting is non-astrophysical and thus considered as the background distribution from the null hypothesis that GW and GRB candidates are only chance associations. Performing the time-shifting multiple times, we effectively created a background observation of ~ 1000 years. The false alarm rate for a foreground association is quantified by the rate of background signals with an equal or more significant ranking statistic. Following the conventions in the literature (e.g., Abbott et al. 2016), we consider GW candidate events that have a false alarm rate less than 1 in 100 years to be a confident detection.

3. SEARCH RESULTS AND IMPLICATIONS

As described, we target the long GRB signals that occurred at most 10 s after a GW candidate and rank their spatial overlap and statistical significance. The complete search results are illustrated in Fig. 1 for the cumulative number of candidates and the inverse false alarm rate from search and expected from noise background, and Table 1 for more detailed source information for the associated pairs.




Figure 1. Cumulative number of GW/long-GRB association candidates versus the inverse false alarm rate. The dashed line and shadows show the expected results from background noise fluctuation and one-, two- and three-sigma uncertainty from a Poisson process. The most significant associated pair 200205.201716 GRB 200205845 has ranking statistic $\lambda_{\text{gw+grb}}=5.53$ which corresponds to a false alarm rate of 1 in 2.2 years.

There are thirty-two associated GW/long-GRB candidates from 2015-2020 when the GW and GRB detectors were both in observation mode. The most significant candidate is from the GW candidate 200205.201716 and the long-duration GRB 200205845, which occurs 6.7 s after the GW coalescence time. The sky overlap integral is $\ln B_{\text{overlap}} = 1.35$ and visualized in Fig. 2. However, the false alarm rate is once every 2.2 years; given the total observation time, this is consistent with the null hypothesis. Since none of the search candidates meet the criteria that the false alarm rate is lower than 1-in-100 years, we conclude no statistically significant GW/long-GRB associations are found in our search.




Figure 2. The posterior distribution of sky location for 200205.201716 from Bayesian inference and GRB 200205845 from the Fermi burst catalog. The inner and outer contours correspond to 50% and 90% credible intervals, respectively. The sky map overlap integral is $\ln B_{\text{overlap}} = 1.35$.

We further retrieved any follow-up observations for the top-5 long GRB candidates in Table 1 and noticed Swift/UVOT had detected an afterglow 109 s after the trigger time of GRB 170330A (Swift/UVOT team 2017), the second top in our search results. We are unaware of any additional follow-up observations associated with these long GRBs. In addition, a visual inspection of the light curves of the GRBs in Table 1 shows that GRB 170626A seems the only event that resembles the morphology of the light curve of GRB 211211A, i.e., with the main emission phase followed by some extended emission. Unfortunately, no firm conclusions can be drawn due to the lack of information, such as the GRB’s host galaxy (thus, the redshift). Nevertheless, future improvements along this line can incorporate the GRB light curve properties into the ranking statistic as our knowledge evolves for long GRBs as the result of compact binary mergers.

Given the non-detection, we define an effective exclusion distance d_{ex} for each GW/long-GRB pair as the distance at which a 1.4-1.4 M_{\odot} (source frame mass) binary neutron star merger would have an signal-to-noise ratio (SNR) of 8 at the GW trigger time while conservatively assuming the viewing angle is 30° . An SNR of 8 is a conservative detection criteria for coincident gravitational-wave candidates. We marginalize the assumed source’s location over the sky map given by Fermi/Swift and over the gravitational-wave polarization angle, assuming a uniform $[0, 2\pi]$ interval. The physical meaning of d_{ex} is the critical distance at which a source would have been conservatively detected, assuming it is indeed from a 1.4-1.4 M_{\odot} binary neutron star merger with sky map inferred from Fermi/Swift. For other component mass and inclination viewing angle assumptions, one can straightforwardly rescale d_{ex} . The exclusion distance for all the GW/long-GRB search candidates is presented in the last column of Table 1.

4. DISCUSSION AND CONCLUSION

Inspired by the discovery of a kilonova located at 350 Mpc associated with a long GRB event (Rastinejad et al. 2022), this work performs the first multimessenger search for GW from binary neutron star or neutron star-black hole mergers and potentially associated long GRB signals. We set a 10 s time lag window for long GRB compared to a GW candidate and rank their correlation by their spatial overlap and the confidence of GW as real signals. We present the complete search results in Fig. 1 and Table 1, and do not find statistically significant associated pairs and thus conclude no detection. With the null results, we report the exclusion distance as a threshold for the GW/GRB candidates, closer than that any

1.4-1.4 M_{\odot} binary neutron star mergers with an inclina-

tion angle 30° should have been detected through GW signals.

Table 1. The complete list of the search results. We list the name of the GW candidates, their chirp mass $\mathcal{M}_c = (m_1 m_2)^{3/5} / (m_1 + m_2)^{1/5}$, the name of the long GRB candidates, their trigger time in UTC (coordinated universal time), the GW search statistic λ_{gw} , the ranking statistic of GW/long-GRB association $\lambda_{\text{gw+grb}}$, the inverse false alarm rate (IFAR), the GW observatories which record the data (H, L, and V stand for LIGO Hanford, LIGO Livingston and Virgo, respectively) and the exclusion distance d_{ex} . The nomenclature of GW candidates is in the form of YYMMDD_HHMMSS which is the coalescence time in UTC.

	GW candidate	\mathcal{M}_c/M_{\odot}	long GRB	t_{GRB}	λ_{gw}	$\lambda_{\text{gw+grb}}$	IFAR/yr	GWobs	d_{ex}/Mpc
1	200205_201716	1.01	GRB 200205845	2020-02-05 20:17:23.32	4.2	5.5	2.20	HL	281
2	170330_222948	1.05	GRB 170330A	2017-03-30 22:29:51.34	6.9	6.7	1.99	HL	191
3	191213_060532	1.01	GRB 191213254	2019-12-13 06:05:33.02	4.1	4.1	0.91	HLV	286
4	151001_082025	1.12	GRB 151001348	2015-10-01 08:20:35.16	3.2	5.2	0.73	HL	156
5	190404_070114	1.03	GRB 190404293	2019-04-04 07:01:21.92	1.7	3.9	0.67	HLV	341
6	191110_140525	1.20	GRB 191110587	2019-11-10 14:05:34.99	0.3	3.2	0.54	HLV	231
7	191213_040623	1.53	GRB 191213A	2019-12-13 04:06:23.92	2.7	1.8	0.29	HLV	363
8	190701_094513	1.03	GRB 190701A	2019-07-01 09:45:20.83	3.4	1.9	0.27	HLV	295
9	170723_161524	0.97	GRB 170723677	2017-07-23 16:15:27.85	1.8	2.2	0.21	HL	176
10	170409_024157	0.88	GRB 170409112	2017-04-09 02:42:00.49	1.9	2.1	0.20	HL	194
11	190613_040717	1.19	GRB 190613A	2019-06-13 04:07:18.31	1.7	0.6	0.16	HLV	343
12	200216_090724	0.88	GRB 200216380	2020-02-16 09:07:25.03	0.6	-0.0	0.14	HLV	274
13	190827_111245	1.57	GRB 190827467	2019-08-27 11:12:48.54	1.3	-0.5	0.11	HL	76
14	190508_234118	1.24	GRB 190508987	2019-05-08 23:41:24.14	0.6	-0.5	0.11	HLV	389
15	190628_123052	1.55	GRB 190628521	2019-06-28 12:30:55.31	0.4	-0.8	0.10	HL	313
16	170402_065048	2.42	GRB 170402285	2017-04-02 06:50:54.39	1.1	-0.7	0.09	HL	184
17	190726_152445	0.87	GRB 190726642	2019-07-26 15:24:53.60	0.7	-1.3	0.09	HLV	183
18	151029_074936	1.40	GRB 151029A	2015-10-29 07:49:38.96	4.2	-0.5	0.09	HL	65
19	191119_061605	1.27	GRB 191119261	2019-11-19 06:16:07.17	5.1	-2.4	0.07	HL	192
20	170424_101224	1.03	GRB 170424425	2017-04-24 10:12:30.75	1.7	-1.7	0.07	HL	129
21	190623_110326	1.46	GRB 190623461	2019-06-23 11:03:27.09	0.7	-2.7	0.07	HLV	238
22	170626_093721	1.17	GRB 170626A	2017-06-26 09:37:23.12	0.4	-2.6	0.06	HL	137
23	200317_004025	0.91	GRB 200317028	2020-03-17 00:40:30.48	0.6	-4.4	0.06	HLV	131
24	200117_122400	1.05	GRB 200117517	2020-01-17 12:24:06.53	2.9	-4.9	0.06	HLV	213
25	190919_181958	1.34	GRB 190919764	2019-09-19 18:20:02.65	0.7	-6.7	0.05	HLV	394
26	190805_044554	1.84	GRB 190805199	2019-08-05 04:46:00.97	3.2	-10.8	0.04	HLV	204
27	190422_160459	2.95	GRB 190422670	2019-04-22 16:05:04.52	4.2	-11.1	0.04	HLV	370
28	170825_120003	1.12	GRB 170825500	2017-08-25 12:00:05.99	0.2	-6.7	0.04	HLV	186
29	170323_012316	2.04	GRB 170323058	2017-03-23 01:23:23.26	4.6	-9.5	0.04	HL	121
30	190824_144634	1.19	GRB 190824A	2019-08-24 14:46:39.57	0.8	-inf	0.04	HLV	292
31	151027_224040	1.90	GRB 151027B	2015-10-27 22:40:40.66	7.4	-inf	0.03	HL	182
32	170629_125329	2.17	GRB 170629A	2017-06-29 12:53:33.15	3.6	-inf	0.03	HL	195

The observation reported by [Rastinejad et al. \(2022\)](#) opens up a promising future for a new type of GW multimessenger astronomy. Richer structures of the burst, such as a precursor ([Xiao et al. 2022](#)) and an afterglow ([Mei et al. 2022](#); [Zhang et al. 2022](#)) are reported. GW observation can provide irreplaceable information for the source properties of compact objects,

including their mass, spin, tidal deformability, luminosity distance, and inclination angle, which is crucial to distinguishing different scenarios between binary neutron star, neutron star-black hole (e.g., [Gompertz et al. 2022](#); [Gao et al. 2022](#); [Yang et al. 2022](#)). Even with non-detection, GW can constrain the exclusion distance and provide evidence for alternative explanations for the en-

gine of GRB 211211A, such as neutron star-white dwarf merger Yang et al. (2022). The next observation run of LIGO/Virgo/KAGRA is scheduled to start in March 2023 with further improved horizon distance. Future (non-) detections will shed more light on the source property mystery of the GRB 211211A-like events.

We release all the data and scripts necessary to reproduce this work in <https://github.com/gwastro/gw-longgrb>

ACKNOWLEDGMENTS

YFW, AHN, and CDC acknowledge the Max Planck Gesellschaft and the Atlas cluster computing team at AEI Hannover for support. BBZ acknowledges the support by the National Key Research and Development Programs of China (2018YFA0404204), the National Natural Science Foundation of China (Grant Nos. 11833003, U2038105, 12121003), the science research grants from the China Manned Space Project with NO.CMS-CSST-2021-B11, and the Program for Innovative Talents, Entrepreneur in Jiangsu. This research has made use of data, software and/or web tools obtained from the Gravitational Wave Open Science Center (<https://www.gw-openscience.org>), a service of LIGO Laboratory, the LIGO Scientific Collaboration and the Virgo Collaboration. LIGO is funded by the U.S. National Science Foundation. Virgo is funded by the French Centre National de Recherche Scientifique (CNRS), the Italian Istituto Nazionale della Fisica Nucleare (INFN) and the Dutch Nikhef, with contributions by Polish and Hungarian institutes. We acknowledge the use of public data from the Fermi Science Support Center (FSSC) and the Swift data archive.

REFERENCES

- Aasi, J., et al. 2015, *Class. Quantum Grav.*, 32, 074001, doi: [10.1088/0264-9381/32/7/074001](https://doi.org/10.1088/0264-9381/32/7/074001)
- Abbott, B., et al. 2017a, *Phys. Rev. Lett.*, 119, 161101, doi: [10.1103/PhysRevLett.119.161101](https://doi.org/10.1103/PhysRevLett.119.161101)
- . 2017b, *Astrophys. J. Lett.*, 848, L13, doi: [10.3847/2041-8213/aa920c](https://doi.org/10.3847/2041-8213/aa920c)
- . 2020, *Astrophys. J. Lett.*, 892, L3, doi: [10.3847/2041-8213/ab75f5](https://doi.org/10.3847/2041-8213/ab75f5)
- Abbott, B. P., et al. 2016, *Phys. Rev. X*, 6, 041015, doi: [10.1103/PhysRevX.6.041015](https://doi.org/10.1103/PhysRevX.6.041015), [10.1103/PhysRevX.8.039903](https://doi.org/10.1103/PhysRevX.8.039903)
- . 2017c, *Astrophys. J.*, 848, L12, doi: [10.3847/2041-8213/aa91c9](https://doi.org/10.3847/2041-8213/aa91c9)
- . 2017d, *Nature*, 551, 85, doi: [10.1038/nature24471](https://doi.org/10.1038/nature24471)
- . 2017e, *Astrophys. J. Lett.*, 850, L39, doi: [10.3847/2041-8213/aa9478](https://doi.org/10.3847/2041-8213/aa9478)
- . 2017f, *Astrophys. J.*, 841, 89, doi: [10.3847/1538-4357/aa6c47](https://doi.org/10.3847/1538-4357/aa6c47)
- . 2019a, *Phys. Rev. Lett.*, 123, 011102, doi: [10.1103/PhysRevLett.123.011102](https://doi.org/10.1103/PhysRevLett.123.011102)
- . 2019b, *Astrophys. J.*, 886, 75, doi: [10.3847/1538-4357/ab4b48](https://doi.org/10.3847/1538-4357/ab4b48)
- . 2019c, *Phys. Rev. X*, 9, 011001, doi: [10.1103/PhysRevX.9.011001](https://doi.org/10.1103/PhysRevX.9.011001)
- Abbott, R., et al. 2021a, *Phys. Rev. X*, 11, 021053, doi: [10.1103/PhysRevX.11.021053](https://doi.org/10.1103/PhysRevX.11.021053)
- . 2021b, *Astrophys. J.*, 915, 86, doi: [10.3847/1538-4357/abee15](https://doi.org/10.3847/1538-4357/abee15)

- . 2021c. <https://arxiv.org/abs/2111.03606>
- . 2021d, *Astrophys. J. Lett.*, 915, L5, doi: [10.3847/2041-8213/ac082e](https://doi.org/10.3847/2041-8213/ac082e)
- . 2022, *Astrophys. J.*, 928, 186, doi: [10.3847/1538-4357/ac532b](https://doi.org/10.3847/1538-4357/ac532b)
- Acernese, F., et al. 2015, *Class. Quantum Grav.*, 32, 024001, doi: [10.1088/0264-9381/32/2/024001](https://doi.org/10.1088/0264-9381/32/2/024001)
- Ahumada, T., et al. 2021, *Nature Astron.*, 5, 917, doi: [10.1038/s41550-021-01428-7](https://doi.org/10.1038/s41550-021-01428-7)
- Ajith, P., Hannam, M., Husa, S., et al. 2011, *Phys. Rev. Lett.*, 106, 241101, doi: [10.1103/PhysRevLett.106.241101](https://doi.org/10.1103/PhysRevLett.106.241101)
- Amiri, M., et al. 2018, doi: [10.3847/1538-4357/aad188](https://doi.org/10.3847/1538-4357/aad188)
- . 2021, *Astrophys. J. Supp.*, 257, 59, doi: [10.3847/1538-4365/ac33ab](https://doi.org/10.3847/1538-4365/ac33ab)
- Ashton, G., Burns, E., Canton, T. D., et al. 2018, *Astrophys. J.*, 860, 6, doi: [10.3847/1538-4357/aabfd2](https://doi.org/10.3847/1538-4357/aabfd2)
- Biwer, C. M., Capano, C. D., De, S., et al. 2019, *Publ. Astron. Soc. Pac.*, 131, 024503, doi: [10.1088/1538-3873/aaef0b](https://doi.org/10.1088/1538-3873/aaef0b)
- Capano, C. D., Tews, I., Brown, S. M., et al. 2020, *Nature Astron.*, 4, 625, doi: [10.1038/s41550-020-1014-6](https://doi.org/10.1038/s41550-020-1014-6)
- Davies, G. S., Dent, T., Tápai, M., et al. 2020, *Phys. Rev. D*, 102, 022004, doi: [10.1103/PhysRevD.102.022004](https://doi.org/10.1103/PhysRevD.102.022004)
- Dezalay, J. P., Barat, C., Talon, R., et al. 1992, in *American Institute of Physics Conference Series*, Vol. 265, American Institute of Physics Conference Series, ed. W. S. Paciesas & G. J. Fishman, 304
- Dhurkunde, R., & Nitz, A. H. 2022. <https://arxiv.org/abs/2207.14645>
- Eichler, D., Livio, M., Piran, T., & Schramm, D. N. 1989, *Nature*, 340, 126, doi: [10.1038/340126a0](https://doi.org/10.1038/340126a0)
- Finn, L. S. 1992, *Phys. Rev.*, D46, 5236, doi: [10.1103/PhysRevD.46.5236](https://doi.org/10.1103/PhysRevD.46.5236)
- Foucart, F., Hinderer, T., & Nissanke, S. 2018, *Phys. Rev. D*, 98, 081501, doi: [10.1103/PhysRevD.98.081501](https://doi.org/10.1103/PhysRevD.98.081501)
- Gal-Yam, A., et al. 2006, *Nature*, 444, 1053, doi: [10.1038/nature05373](https://doi.org/10.1038/nature05373)
- Galama, T. J., Vreeswijk, P. M., van Paradijs, J., et al. 1998, *Nature*, 395, 670, doi: [10.1038/27150](https://doi.org/10.1038/27150)
- Gao, H., Lei, W.-H., & Zhu, Z.-P. 2022. <https://arxiv.org/abs/2205.05031>
- Gehrels, N., et al. 2004, *Astrophys. J.*, 611, 1005, doi: [10.1086/422091](https://doi.org/10.1086/422091)
- . 2006, *Nature*, 444, 1044, doi: [10.1038/nature05376](https://doi.org/10.1038/nature05376)
- Gompertz, B. P., et al. 2022. <https://arxiv.org/abs/2205.05008>
- Goodman, J. 1986, *ApJL*, 308, L47, doi: [10.1086/184741](https://doi.org/10.1086/184741)
- Guidorzi, C., et al. 2017, *Astrophys. J. Lett.*, 851, L36, doi: [10.3847/2041-8213/aaa009](https://doi.org/10.3847/2041-8213/aaa009)
- Hjorth, J., Sollerman, J., Møller, P., et al. 2003, *Nature*, 423, 847, doi: [10.1038/nature01750](https://doi.org/10.1038/nature01750)
- Hotokezaka, K., Nakar, E., Gottlieb, O., et al. 2019, *Nature Astron.*, 3, 940, doi: [10.1038/s41550-019-0820-1](https://doi.org/10.1038/s41550-019-0820-1)
- Kouveliotou, C., Meegan, C. A., Fishman, G. J., et al. 1993, *ApJL*, 413, L101, doi: [10.1086/186969](https://doi.org/10.1086/186969)
- Meegan, C., Lichti, G., Bhat, P. N., et al. 2009, *ApJ*, 702, 791, doi: [10.1088/0004-637X/702/1/791](https://doi.org/10.1088/0004-637X/702/1/791)
- Mei, A., et al. 2022. <https://arxiv.org/abs/2205.08566>
- Narayan, R., Paczynski, B., & Piran, T. 1992, *ApJL*, 395, L83, doi: [10.1086/186493](https://doi.org/10.1086/186493)
- Nitz, A. H., Dent, T., Dal Canton, T., Fairhurst, S., & Brown, D. A. 2017, *Astrophys. J.*, 849, 118, doi: [10.3847/1538-4357/aa8f50](https://doi.org/10.3847/1538-4357/aa8f50)
- Nitz, A. H., Dent, T., Davies, G. S., & Harry, I. 2020, *Astrophys. J.*, 897, 169, doi: [10.3847/1538-4357/ab96c7](https://doi.org/10.3847/1538-4357/ab96c7)
- Nitz, A. H., Kumar, S., Wang, Y.-F., et al. 2021. <https://arxiv.org/abs/2112.06878>
- Nitz, A. H., & Kumar, S. e. a. 2021, 4-OGC Open Gravitational-wave Catalog, www.github.com/gwastro/4-ogc, GitHub
- Nitz, A. H., Nielsen, A. B., & Capano, C. D. 2019, *Astrophys. J. Lett.*, 876, L4, doi: [10.3847/2041-8213/ab18a1](https://doi.org/10.3847/2041-8213/ab18a1)
- Nitz, A. H., Harry, I. W., Willis, J. L., et al. 2018, *PyCBC Software*, <https://github.com/gwastro/pycbc>, GitHub
- Olsen, S., Venumadhav, T., Mushkin, J., et al. 2022. <https://arxiv.org/abs/2201.02252>
- Paczynski, B. 1986, *ApJL*, 308, L43, doi: [10.1086/184740](https://doi.org/10.1086/184740)
- Rastinejad, J. C., et al. 2022. <https://arxiv.org/abs/2204.10864>
- Singer, L. P., & Price, L. R. 2016, *Phys. Rev. D*, 93, 024013, doi: [10.1103/PhysRevD.93.024013](https://doi.org/10.1103/PhysRevD.93.024013)
- Speagle, J. S. 2020, *Monthly Notices of the Royal Astronomical Society*, 493, 3132, doi: [10.1093/mnras/staa278](https://doi.org/10.1093/mnras/staa278)
- Stanek, K. Z., Matheson, T., Garnavich, P. M., et al. 2003, *ApJL*, 591, L17, doi: [10.1086/376976](https://doi.org/10.1086/376976)
- Swift/BAT. 2018, GRB 211211A, https://swift.gsfc.nasa.gov/archive/grb_table/211211A/
- Swift/UVOT team. 2017, GRB 170330A: Swift/UVOT Detection of a Fading Afterglow, <https://gcn.gsfc.nasa.gov/gcn/gcn3/20959.gcn3>
- The LIGO Scientific Collaboration, the Virgo Collaboration, the KAGRA Collaboration, et al. 2022, arXiv e-prints, arXiv:2203.12038. <https://arxiv.org/abs/2203.12038>
- Veitch, J., Raymond, V., Farr, B., et al. 2015, *Phys. Rev. D*, 91, 042003, doi: [10.1103/PhysRevD.91.042003](https://doi.org/10.1103/PhysRevD.91.042003)

- Wang, Y.-F., & Nitz, A. H. 2022.
<https://arxiv.org/abs/2203.17222>
- Waxman, E., Ofek, E. O., & Kushnir, D. 2022.
<https://arxiv.org/abs/2206.10710>
- Xiao, S., et al. 2022. <https://arxiv.org/abs/2205.02186>
- Yang, J., Zhang, B.-B., Ai, S. K., et al. 2022.
<https://arxiv.org/abs/2204.12771>
- Zhang, B. B., et al. 2021, *Nature Astron.*, 5, 911,
doi: [10.1038/s41550-021-01395-z](https://doi.org/10.1038/s41550-021-01395-z)
- Zhang, H.-M., Huang, Y.-Y., Zheng, J.-H., Liu, R.-Y., &
Wang, X.-Y. 2022, doi: [10.3847/2041-8213/ac7b23](https://doi.org/10.3847/2041-8213/ac7b23)
- Zhu, X., Thrane, E., Osłowski, S., Levin, Y., & Lasky, P. D.
2018, *Phys. Rev. D*, 98, 043002,
doi: [10.1103/PhysRevD.98.043002](https://doi.org/10.1103/PhysRevD.98.043002)