

Inflation Alternative via the Gravitational Field of a Singularity

Michael Zlotnikov^a

^aDepartment of Physics, Center for Theoretical Physics, Columbia University, 538 West 120th Street, New York, NY 10027, USA.

E-mail: mz2737@columbia.edu

Abstract. We explore the scenario that the observable universe emerged from the ergosphere of a negative mass ring singularity, and all content of the universe travels at the same group velocity close to the speed of light on a geodesic trajectory along the axis of rotation of the singularity. In appropriate coordinate parametrization and evaluated on the trajectory, we find that the metric tensor in the vicinity of the trajectory exhibits a conformal scale factor $\mathbf{a}(\eta)$ with contraction and subsequent expansion properties that solve the horizon problem. We then introduce a static flow of gravitating radiation along the trajectory (perturbatively with respect to the mass scale of the singularity) to model a homogeneous radiation dominated universe. Solving the Einstein field equations with a physically motivated ansatz of metric perturbation then reveals that the effective conformal scale factor indeed grows asymptotically with the same power law as expected in a conventional radiation dominated universe.

Contents

1	Introduction	1
2	Preliminaries	2
3	Empty Universe	3
4	Radiation Dominated Universe	7
5	Conclusion and Outlook	12

1 Introduction

The horizon problem [1] is concerned with the fact that different regions of the Cosmic Microwave Background (CMB) [2, 3], emitted at the surface of last scattering, possess surprisingly uniform temperature. Assuming a generic cosmic evolution, these regions could not have been in causal contact with each other, and yet their almost equal temperatures suggest that they must have thermalized at some point in the past. A possible solution to this problem is the inflationary universe paradigm [4, 5], in which the universe undergoes a period of accelerated expansion at early times, taking space-time regions that previously interacted with each other out of causal contact, which then re-enter the cosmological horizon upon later cosmic evolution. The process of inflation is typically assumed to be driven by a space-filling inflaton field (see for example [5–18]). At the end of inflation all other fields are thinned out and the universe is re-populated with standard model constituents through the decay process of the inflaton field called reheating [19, 20].

In this work we explore an alternative way to resolve the horizon problem by assuming that the observable universe has been emitted from the ergosphere of a negative mass Kerr-like solution of the Einstein field equations,¹ and all content of the universe travels at approximately the same group velocity close to the speed of light along a radial geodesic parallel to the Kerr spin axis. The naked singularity in this setting is hidden from view because the universe is opaque at early times. The relative motion with respect to the singularity manifests itself locally on the geodesic in terms of an effective cosmological scale factor with specific contraction and expansion properties solving the horizon problem.

Subsequently, introducing a static flow of gravitating radiation, with energy density parametrically smaller than the mass of the singularity, we show that a leading order perturbative correction to the metric tensor leads to the same asymptotic late conformal time scaling of the effective scale factor as expected in a Friedmann–Lemaître–Robertson–Walker (FLRW) radiation dominated universe. The transition to a matter dominated universe could then be made in a conventional formalism, without invoking a singularity in the far past.

This paper is organized as follows:

In section 2 we recall the basic tools of General Relativity to be used in the following.

Section 3 introduces an appropriate cylindrical change of coordinates, parameterizing the vicinity of a radial geodesic trajectory parallel to the axis of rotation in a negative mass

¹Negative mass regions are commonplace in exact solutions of General relativity, and can be argued to be physically meaningful. See e.g. [21] for a motivation.

Kerr space-time. Evaluating the metric on the geodesic trajectory and expressing the solution in conformal time reveals that at leading order in the vicinity of the trajectory the metric possesses an effective conformal scale factor \mathbf{a} . The scale factor \mathbf{a} starts at unity, contracts to a parametrically small number, depending on the Kerr spin parameter and mass $a/M \rightarrow 0$, and subsequently asymptotically expands back to unity in the case of an empty universe. The initial contraction and subsequent expansion of the scale factor explains how seemingly separated regions entering the cosmological horizon during the expansion phase were in causal contact previously.

In section 4 we perturbatively add a static flow of gravitating radiation traveling along the radial geodesic. With an appropriate perturbation ansatz for the metric tensor, we solve the Einstein equations to leading order in the vicinity of the radial trajectory and small Kerr spin parameter $a/M \rightarrow 0$. The resulting correction to the effective conformal scale factor turns out to be given by an integral over ratios of hypergeometric functions. At late conformal time $\eta \gg M$, the effective conformal scale factor then turns out to scale as $\sim \eta$, which is the expected scaling for a radiation dominated universe.

We conclude in section 5, and offer a few suggestions for further investigation that goes beyond the scope of this paper.

2 Preliminaries

In this section we recall the basic tools of General Relativity (see e.g. [22]), to be applied in subsequent sections. We use the Einstein index summation convention $a^\mu b_\mu \equiv \sum_{\mu=0}^3 a^\mu b_\mu$ throughout.

Given a set of coordinates x^μ describing a space-time, along with the corresponding metric tensor $g_{\mu\nu}$ (lowering indices $x_\mu = g_{\mu\nu}x^\nu$) and inverse metric tensor $g^{\mu\nu}$ (raising indices $x^\mu = g^{\mu\nu}x_\nu$), the Christoffel symbols are determined as follows

$$\Gamma^\sigma{}_{\mu\nu} = \frac{1}{2}g^{\sigma\beta} \left(\frac{\partial g_{\beta\mu}}{\partial x^\nu} + \frac{\partial g_{\beta\nu}}{\partial x^\mu} - \frac{\partial g_{\mu\nu}}{\partial x^\beta} \right). \quad (2.1)$$

The Ricci curvature tensor $R_{\mu\nu}$ and Ricci scalar R are then constructed as

$$R_{\mu\nu} = \frac{\partial \Gamma^\beta{}_{\mu\nu}}{\partial x^\beta} - \frac{\partial \Gamma^\beta{}_{\beta\nu}}{\partial x^\mu} + \Gamma^\beta{}_{\beta\gamma} \Gamma^\gamma{}_{\mu\nu} - \Gamma^\beta{}_{\mu\gamma} \Gamma^\gamma{}_{\beta\nu} \quad \text{and} \quad R = g^{\mu\nu} R_{\mu\nu}. \quad (2.2)$$

These quantities enter the Einstein field equations, along with energy momentum tensor $T^{\mu\nu}$

$$G^{\mu\nu} \equiv R^{\mu\nu} - \frac{R}{2}g^{\mu\nu} + \Lambda g^{\mu\nu} = \kappa T^{\mu\nu}, \quad (2.3)$$

where $G^{\mu\nu}$ is the Einstein tensor, and Λ is a potentially non-zero cosmological constant. In the following we will be working in geometrized units with the Einstein gravitational constant set to unity $\kappa = 1$.

For physical matter/radiation space-time content, the covariant divergence of the energy momentum tensor must vanish

$$0 = \nabla_\nu T^{\mu\nu} = \frac{\partial T^{\mu\nu}}{\partial x^\nu} + \Gamma^\mu{}_{\sigma\nu} T^{\sigma\nu} + \Gamma^\nu{}_{\sigma\nu} T^{\mu\sigma}. \quad (2.4)$$

Free falling particles in a given space-time follow geodesic trajectories that are solutions to the geodesic equation

$$\frac{d^2 x^\mu}{d\tau^2} + \Gamma^\mu_{\beta\beta} \frac{dx^\beta}{d\tau} \frac{dx^\beta}{d\tau} = 0, \quad (2.5)$$

where τ is the proper time (or affine parameter) parameterizing the trajectory. Sometimes, it is also convenient to employ the proper time equation

$$-d\tau^2 = g_{\mu\nu} dx^\mu dx^\nu \quad (2.6)$$

to solve for a geodesic trajectory.

3 Empty Universe

We start with the *negative mass* region of the Kerr metric [23] in Boyer-Lindquist coordinates, employing the parameter $M > 0$:²

$$-d\tau^2 = -\frac{\Delta_r}{\Sigma_r} (dt - a d\phi \sin^2(\theta))^2 + \Sigma_r \left(\frac{dr^2}{\Delta_r} + d\theta^2 \right) + \frac{\sin^2(\theta)}{\Sigma_r} (adt - d\phi (a^2 + r^2))^2, \quad (3.1)$$

where

$$\Delta_r = a^2 + 2Mr + r^2 \quad \text{and} \quad \Sigma_r = a^2 \cos^2(\theta) + r^2, \quad (3.2)$$

and a is the spin parameter. Since there is no relative minus sign in the Δ_r factors, this space-time does not exhibit any black hole horizons. Therefore, particles can escape from the ergosphere to infinity, which will be an important feature in the following.

We work towards understanding how the metric behaves locally, when evaluated on an outgoing radial geodesic trajectory and expressed in conformal coordinates at leading order in the vicinity of the trajectory. This shall facilitate a comparison with a conventional cosmological metric of an unperturbed universe.

As a first step, we introduce a cylindrical type of coordinates, along with an expansion around small cylindrical radius, to describe the vicinity of the straight line trajectory. To that end, we perform the following change of coordinates $(r, \theta) \rightarrow (\mathbf{r}, z)$ for $z \geq 0$, and an expansion around small $\mathbf{r} \ll M$:³

$$\begin{aligned} r &= z + \mathbf{r}^2 \frac{(M+z)}{2(a^2 + z(2M+z))} + \mathbf{r}^4 \frac{((2M+3z)a^4 + 2z^2(4M+z)a^2 - z^4(2M+z))}{8(a^2 + z^2)^2 (a^2 + z(2M+z))^2} + \mathcal{O}(\mathbf{r}^5), \\ \theta &= \mathbf{r} \frac{1}{\sqrt{a^2 + z(2M+z)}} + \mathbf{r}^3 \frac{a^4 + (3M-z)za^2 - z^3(3M+2z)}{6(a^2 + z^2)^2 (a^2 + z(2M+z))^{3/2}} + \mathcal{O}(\mathbf{r}^5), \end{aligned} \quad (3.3)$$

²Compared to the regular Kerr black hole metric, this differs by the sign in front of the mass parameter M , and is equally an exact solution of the vacuum Einstein field equations.

³Here, small cylindrical radius \mathbf{r} corresponds to small θ , while the original spherical radius r approaches z in that limit.

resulting in the following differential relations

$$dr = dz + r \frac{M+z}{a^2+z(2M+z)} dr - r^2 \frac{-a^2+2M^2+z^2+2Mz}{2(a^2+z(2M+z))^2} dz + r^3 \frac{(2M+3z)a^4+2z^2(4M+z)a^2-z^4(2M+z)}{2(a^2+z^2)^2(a^2+z(2M+z))^2} dr + \mathcal{O}(r^4), \quad (3.4)$$

$$d\theta = \frac{dr}{\sqrt{a^2+z^2+2Mz}} - \frac{dz(M+z)r}{(a^2+z(2M+z))^{3/2}} + \frac{dr(a^4+(3M-z)za^2-z^3(3M+2z))r^2}{2(a^2+z^2)^2(a^2+z(2M+z))^{3/2}} - r^3 \frac{3a^6+(M^2+12zM+4z^2)a^4+z^2(12M^2+6zM-z^2)a^2-z^4(5M^2+6zM+2z^2)}{2(a^2+z^2)^3(a^2+z(2M+z))^{5/2}} z dz + \mathcal{O}(r^4). \quad (3.5)$$

Introducing these coordinates into the metric (3.1), we obtain⁴

$$-d\tau^2 = -\frac{a^2+z(2M+z)}{a^2+z^2} dt^2 + \frac{a^2+z^2}{a^2+z(2M+z)} (dr^2 + r^2 d\phi^2 + dz^2) + r^2 M \left(\frac{(z^2(M+z) - a^2(M+3z)) dt^2 + 4az(a^2+z^2) dt d\phi}{(a^2+z^2)^2(a^2+z(2M+z))} + \frac{z^3 - a^2(2M+3z)}{(a^2+z(2M+z))^3} dz^2 + \frac{(a^4(M+6z) + 2a^2z^2(5M+2z) - z^4(3M+2z))}{(a^2+z^2)(a^2+z(2M+z))^3} dr^2 \right) + dz dr \mathcal{O}(r^3) + r^4 \frac{2aM(a-z)(a+z)(a^2(M+3z) + 3z^2(M+z))}{(a^2+z^2)^3(a^2+z(2M+z))^2} dt d\phi + \mathcal{O}(r^4). \quad (3.6)$$

The first line in (3.6) contains the leading order metric and features homogeneous scaling of generic cylindrical spatial coordinates by a z -dependent factor. The subsequent lines in (3.6) contain sub-leading metric terms sufficient to ensure that the overall metric remains a vacuum solution of the Einstein field equations (2.3) (with $\Lambda = 0$) in the vicinity ($r/M \rightarrow 0$) of the trajectory of interest to order $\mathcal{O}(r^2)$:

$$R_{\mu\nu} - \frac{R}{2} g_{\mu\nu} = \mathcal{O}(r^2). \quad (3.7)$$

Note in particular, that a $dz dr \mathcal{O}(r^3)$ term as well as several $\mathcal{O}(r^4)$ terms are neglected in (3.6), since they contribute to (3.7) at order $\mathcal{O}(r^2)$ or higher and are therefore irrelevant for our purposes.

What is the utility of formulating the metric as in (3.6)?

By the first postulate of General Relativity, the relativity principle, a point-like observer at any point in any space-time perceives the metric in their immediate vicinity as the flat Minkowski metric, such that local physics is governed by Special Relativity. In other words, at any point in any physical space-time one can find normal coordinates, such that the leading term is given by the flat Minkowski metric, and first derivatives of the metric with respect to all coordinates vanish. Therefore in our case of interest, a point-like observer traveling from

⁴The change of variables (3.3) is set up specifically such that there is no $dz dr$ term at order $\mathcal{O}(r^1)$, the scale factor in front of all spatial coordinates is homogeneous at $\mathcal{O}(r^0)$, and there is no non-trivial $d\phi^2$ term at order $\mathcal{O}(r^4)$.

the ergosphere to infinity along a radial trajectory finds themselves in such normal coordinates at any given instant. If we desire to keep track of how the normal coordinates at a given point relate to the respective normal coordinates at a different point visited previously, a fruitful course of action is to relax the condition of normality for the single coordinate aligned with the path of trajectory, and by doing so treat coordinates at all visited points on an equal footing. The form of the metric (3.6) achieves precisely that, while maintaining homogeneous scaling of spatial coordinates at leading order in small $\mathbf{r} \ll M$. Normality is relaxed for the z -direction, which parametrizes different points on the trajectory, while normality is maintained in t, \mathbf{r}, ϕ coordinates (perturbatively in $\mathbf{r} \ll M$, and up to constant rescaling at fixed z), with their first derivatives vanishing locally. This is analogous to the situation in the FLRW metric of cosmology, where the scale factor $\mathbf{a}(t)$ breaks the normality of the metric in the t -direction, while the rest of the coordinates are normal (up to constant rescaling at fixed t), and spatial coordinates experience homogeneous scaling.

So far, we have chosen a particular parametrization (3.6) for the space-time metric in the vicinity of a radial trajectory. To compare this metric with a typical cosmological metric of an unperturbed universe, we evaluate the metric along the radial trajectory parametrized in conformal time η .

Due to the asymptotic flatness of the Kerr space-time, all sub-leading terms in (3.6) are guaranteed to vanish for increasing z values. We therefore concentrate on the leading order metric in (3.6) to model the observable universe, and transform the time variable to conformal time $t = f(\eta)$, $dt = f'(\eta)d\eta$:

$$-d\tau^2 = -\frac{a^2 + z(2M + z)}{a^2 + z^2} f'(\eta)^2 d\eta^2 + \frac{a^2 + z^2}{a^2 + z(2M + z)} (d\mathbf{r}^2 + \mathbf{r}^2 d\phi^2 + dz^2) + \mathcal{O}(\mathbf{r}^2), \quad (3.8)$$

with the appropriate function $f(\eta)$ to be determined. In these variables, the zero'th component of the geodesic equation (2.5) reads

$$\frac{d^2 x^0}{d\tau^2} + \Gamma^0_{\beta\beta} \frac{dx^\beta}{d\tau} \frac{dx^\beta}{d\tau} = \frac{2M(a-z)(a+z)\eta'(\tau)z'(\tau)}{(a^2+z^2)(a^2+z(2M+z))} + \frac{f''(\eta)\eta'(\tau)^2}{f'(\eta)} + \eta''(\tau) = 0. \quad (3.9)$$

After dividing by $\eta'(\tau)$, this directly integrates to

$$\eta'(\tau) = \frac{c(a^2 + z(\tau)^2)}{(a^2 + z(\tau)(2M + z(\tau))) f'(\eta(\tau))}, \quad (3.10)$$

with an integration constant c . To find $z'(\tau)$ in the case of a massive particle, we use (3.8) with $\mathbf{r}'(\tau) = 0$, $\phi'(\tau) = 0$, plug in (3.10), and solve selecting the positive square root solution

$$-1 = -\frac{a^2 + z(2M + z)}{a^2 + z^2} f'(\eta)^2 \eta'^2 + \frac{a^2 + z^2}{a^2 + z(2M + z)} z'^2 \quad \Rightarrow \quad z'(\tau) = \sqrt{c^2 - 1 - \frac{2Mz(\tau)}{a^2 + z(\tau)^2}}.$$

Note that in order for $z'(\tau)$ to be real valued on the whole range of $0 \leq z(\tau) \leq \infty$, the integration constant c must satisfy $c \geq \sqrt{1 + M/a}$. Therefore, we can set c to e.g. the minimum admissible value

$$c = \sqrt{1 + \frac{M}{a}} \quad \Rightarrow \quad z'(\tau) = \sqrt{\frac{M}{a} - \frac{2Mz(\tau)}{a^2 + z(\tau)^2}}. \quad (3.11)$$

Finally, to obtain $z'(\eta)$ we use the chain rule $z'(\tau) = z'(\eta)\eta'(\tau)$, plug in (3.10, 3.11) and solve

$$z'_m(\eta) = \sqrt{\frac{M(a - z_m)^2}{(a + M)(a^2 + z_m^2)} \frac{a^2 + z_m(2M + z_m)}{a^2 + z_m^2}} f'(\eta), \quad (3.12)$$

where we used the subscript m to emphasize that this is the solution for a massive particle.

To find $z'(\eta)$ in case of a massless particle, we use (3.8) with $d\mathbf{r} = 0$, $d\phi = 0$, $d\tau = 0$, choose the positive square root solution and obtain

$$z'_\gamma(\eta) = \frac{a^2 + z_\gamma(2M + z_\gamma)}{a^2 + z_\gamma^2} f'(\eta), \quad (3.13)$$

where the subscript γ emphasizes that this is the solution for a massless particle.

Recall that we introduced the change of time coordinate $t = f(\eta)$ in (3.8) in order for η to be conformal time. We can enforce this when we evaluate the metric on the trajectory parametrized by η , such that $z = z(\eta)$. In conformal time, we expect e.g. the equality $-g_{00} = g_{33}$ between metric components. Therefore, choosing the positive square root solution, we define $f(\eta)$ such that

$$-g_{00} = \frac{a^2 + z(2M + z)}{a^2 + z^2} f'(\eta)^2 = \frac{a^2 + z^2}{a^2 + z(2M + z)} = g_{33} \quad \Rightarrow \quad f'(\eta) = \frac{a^2 + z(\eta)^2}{a^2 + z(\eta)(2M + z(\eta))}. \quad (3.14)$$

We will see shortly that to model an effective inflation alternative we should take the limit $a \ll M$. In that limit the massive and massless relations (3.12) and (3.13) become equivalent, meaning that massive particles must travel close to the speed of light to successfully emerge from the ergosphere. Therefore, in the following we focus on the massless case without loss of generality.

Plugging (3.14) into (3.13), we observe

$$z'(\eta) = 1, \quad \text{the simplest solution being } z(\eta) = \eta. \quad (3.15)$$

Thus, transformed to conformal time and evaluated on the trajectory of interest, the leading order metric reads

$$-d\tau_{\text{local}}^2 = \frac{a^2 + \eta^2}{a^2 + \eta(2M + \eta)} (-d\eta^2 + d\mathbf{r}^2 + \mathbf{r}^2 d\phi^2 + dz^2) + \mathcal{O}(\mathbf{r}^2), \quad (3.16)$$

where all finite factors have been evaluated to their appropriate η -dependence, while differentials $d\eta$, $d\mathbf{r}$, $d\phi$, dz refer to local space-time properties along the trajectory at a given moment in conformal time η .

Metric (3.16) is analogous to a conventional FLRW metric in conformal time, with an effective conformal scale factor

$$\mathbf{a}(\eta) = \sqrt{\frac{a^2 + \eta^2}{a^2 + \eta(2M + \eta)}}. \quad (3.17)$$

The scale factor equals $\mathbf{a}(0) = 1$ at the initial ergosphere surface, and falls to a minimum of $\mathbf{a}(a) = \sqrt{a/(a + M)}$ when conformal time reaches the value of the Kerr spin parameter $\eta = a$. Subsequently, the scale factor starts growing and asymptotes back to $\mathbf{a}(\infty) = 1$. There




Figure 1. Acceleration $\frac{d^2}{dt^2}\mathbf{a}$ of the scale factor (3.17) as a function of the scale factor \mathbf{a} itself, evaluated at $M = 1$, $a = 10^{-1}$. The curve starts at $\mathbf{a}(a) = \sqrt{a/(a+M)}$, where it assumes its maximum value $\frac{1}{2}Ma^{-3}\sqrt{a/(a+M)}$. After a brief period of positive acceleration, the expansion starts decelerating, until it asymptotically comes to a halt when the scale factor reaches unity $\mathbf{a}(\infty) = 1$.

is a brief period of accelerated expansion at the beginning of the growth phase, see Figure 1 for a qualitative plot. However, the fact that the scale factor starts at unity and decreases in magnitude initially is sufficient to ensure prior causal contact between regions reentering the horizon during the later expansion phase. The smaller the spin parameter a , the smaller the scale factor becomes at its lowest point. Therefore, we concentrate on the small spin case $0 < a \ll M$ in the following, which ensures that any amount of relative expansion starting from the lowest point $\mathbf{a}(a) = \sqrt{a/(a+M)}$ can be accommodated. This demonstrates how non-trivial time evolution of the cosmological scale factor solving the horizon problem can be achieved dynamically e.g. by traversing the gravitational field created by a super-galactic ring singularity in the far past, instead of the conventional approach of producing inflation by introducing space-filling energy-momentum at the particular time when the scale factor is evolving.

So far, no gravitating energy momentum content is traveling along the geodesic trajectory, so that the above metric evolves into the flat Minkowski metric for $\eta \rightarrow \infty$. In the next section we add gravitating radiation emerging from the ergosphere ($z = 0$ surface), to model a radiation dominated unperturbed universe in this setting.

4 Radiation Dominated Universe

In this section we introduce a gravitating radiation energy density emerging from the ergosphere and flowing along a radial outgoing geodesic parallel to the axis of rotation. We

assume the energy momentum flow to be constant at the time scale of interest, and are therefore looking for a static perturbative correction to the metric tensor.

The ergosphere collapses to a point if the Kerr spin parameter a is set to zero exactly, in which case the flow of radiation should vanish. Therefore we assume that the emerging radiation density is proportional to a :

$$\rho_{\gamma,0} \propto \frac{a}{M}. \quad (4.1)$$

Furthermore, as observed in the previous section, the smaller the Kerr spin parameter a is, the more the effective scale factor (3.17) contracts before the expansion phase. We can accommodate any relative amount of expansion from the lowest point if we consider the small spin parameter regime $0 < a \ll M$.

Introducing a non-zero energy momentum flow of radiation causes the geometry of space-time to shift. In order to account for these changes, we take the following ansatz for the perturbation of metric (3.6):

$$\begin{aligned} -d\tau^2 = & -\frac{a^2 + z(2M + z)}{a^2 + z^2} dt^2 \left(1 - \frac{a}{M}(1 + \beta) h_1(z) + \frac{a\mathbf{r}^2}{M^3} h_3(z) \right) + 2dtdz \frac{a\mathbf{r}^2}{M^3} h_2(z) \quad (4.2) \\ & + \frac{a^2 + z^2}{a^2 + z(2M + z)} (d\mathbf{r}^2 + \mathbf{r}^2 d\phi^2 + dz^2) \left(1 + \frac{a}{M} h_1(z) + \frac{a\mathbf{r}^2}{M^3} h_4(z) \right) \\ & + \mathbf{r}^2 M \left(\frac{(z^2(M + z) - a^2(M + 3z)) dt^2 + 4az(a^2 + z^2) dt d\phi}{(a^2 + z^2)^2 (a^2 + z(2M + z))} + \frac{z^3 - a^2(2M + 3z)}{(a^2 + z(2M + z))^3} dz^2 \right. \\ & \quad \left. + \frac{(a^4(M + 6z) + 2a^2 z^2(5M + 2z) - z^4(3M + 2z))}{(a^2 + z^2)(a^2 + z(2M + z))^3} d\mathbf{r}^2 \right) + dzd\mathbf{r} \mathcal{O}(\mathbf{r}^3) \\ & + \mathbf{r}^4 \frac{2aM(a - z)(a + z)(a^2(M + 3z) + 3z^2(M + z))}{(a^2 + z^2)^3 (a^2 + z(2M + z))^2} dt d\phi + \mathcal{O}(\mathbf{r}^4), \end{aligned}$$

where β is a constant and $h_1(z), h_2(z), h_3(z), h_4(z)$ are perturbation functions to be determined. We focus on the case where purely spatial deformations are homogeneous. As discussed above, each perturbation is accompanied by a factor of $\frac{a}{M}$, making sure that for $a = 0$ the perturbations vanish along with the emerging radiation. We are seeking $h_i(z)$ with $i = 1, 2, 3, 4$ solutions of the Einstein field equations at leading order in $a/M \rightarrow 0$ as well as $\mathbf{r}/M \rightarrow 0$. We assume that at leading order $\mathcal{O}(a^1, \mathbf{r}^0)$ the metric deformation can be parametrized by a single function $h_1(z)$ similar to a scale factor; however, for more generality we allow a relative constant factor $-(1 + \beta)$ in the dt^2 metric component. Note that we have introduced a $dt dz$ perturbation into the metric at the sub-leading order $\mathcal{O}(\mathbf{r}^2)$ to account for non-zero energy momentum flow in the z -direction without breaking the diagonal form of the leading order metric.

Radiation *fluid* typically possesses non-zero pressure. However, in our case we are instead interested in radiation propagating at the speed of light in the z -direction. This is equivalent to energy momentum of a superposition of plane waves and takes the familiar form

$$T^{\mu\nu} = \rho(z) U^\mu U^\nu, \quad \text{with four-velocity } U^\mu = \left(\frac{1}{\sqrt{-g_{00}}}, 0, 0, \frac{1}{\sqrt{g_{33}}} \right), \quad (4.3)$$

where we allow the energy density $\rho(z)$ to vary statically with z along the geodesic trajectory. Here, the four-velocity U^μ is lightlike $g_{\mu\nu} U^\mu U^\nu = 0$ for $\mathbf{r}/M \rightarrow 0$ by construction.

At leading order in $\mathbf{r}/M \rightarrow 0$, the covariant derivative vanishing condition (2.4) leads to a single non-trivial equation:

$$\nabla_\nu T^{0\nu} \propto \nabla_\nu T^{3\nu} \propto \partial_z [\rho(z) (ah_1(z) + M) (M - a(1 + \beta)h_1(z))] = 0. \quad (4.4)$$

This directly integrates to⁵

$$\rho = \frac{\rho_{\gamma,0}}{(ah_1 + M) (M - a(1 + \beta)h_1)} = \frac{a}{M (ah_1 + M) (M - a(1 + \beta)h_1)}, \quad (4.5)$$

where the initial energy density (4.1) appears as an integration constant. Making use of (4.5), the other two components of (2.4) vanish or are sub-leading in $a \ll M$

$$\nabla_\nu T^{1\nu} = 0, \quad \nabla_\nu T^{2\nu} = \mathcal{O}(a^2, \mathbf{r}^0). \quad (4.6)$$

We utilize (4.2), (4.3) and (4.5) in the Einstein field equations (2.3). The single non-trivial off-diagonal equation expanded to leading order $\mathcal{O}(a^1, \mathbf{r}^0)$ fixes h_2 in terms of other quantities

$$R^{03} - \frac{R}{2}g^{03} = T^{03} \quad \Rightarrow \quad h_2 = \frac{z}{4M + 2z}. \quad (4.7)$$

All other off-diagonal elements either vanish exactly, or are of sub-leading order $\mathcal{O}(a^2, \mathbf{r}^0)$ and are therefore irrelevant for our purposes.

The diagonal elements $R^{\mu\mu} - \frac{R}{2}g^{\mu\mu} = T^{\mu\mu}$ (no summation over μ) of the Einstein equations (2.3) at order $\mathcal{O}(a^1, \mathbf{r}^0)$ (and an order $\mathcal{O}(a^1, \mathbf{r}^{-2})$ term for $\mu = 3$) then lead to the following three non-trivial linearly independent constraints

$$0 = M^3(3M + 4z)h_1 - z(2M + z) (M^3h_1' + z^2 + (2M + z) (M^2zh_1'' + 4zh_4)), \quad (4.8)$$

$$0 = z(2M + z) ((2M + z) [2z(h_3 + h_4) - \beta M^2zh_1''] + 2(1 + \beta)M^3h_1' - 2M^3h_1((1 + \beta)(M + z) + z)), \quad (4.9)$$

$$0 = z(2M + z) (z(2(h_3 + h_4)(2M + z) - z) - (1 + \beta)M^3h_1' - M^3h_1((2\beta + 5)M + 2(2 + \beta)z)). \quad (4.10)$$

Note that no derivatives of h_3 and h_4 functions make an appearance, which allows us to simply solve two out of the three equations algebraically:

$$h_3 = \frac{z(2M + z) (z((2\beta + 1)M^2(2M + z)h_1'' + z) - (4\beta + 3)M^3h_1') + M^3h_1((4\beta + 1)M + 4(1 + \beta)z)}{4z^2(2M + z)^2},$$

$$h_4 = \frac{M^3h_1(3M + 4z) - z(2M + z) (M^3h_1' + z(M^2(2M + z)h_1'' + z))}{4z^2(2M + z)^2}. \quad (4.11)$$

The third equation then reduces to a differential equation in h_1 only:

$$0 = z(2M + z) (z(z - \beta M^2(2M + z)h_1'') + 3(1 + \beta)M^3h_1') + 3M^4h_1. \quad (4.12)$$

⁵Note that ρ has units of $[\rho] = L^{-2}$ as appropriate in the geometrized units convention.

Despite its simple appearance, this equation is rather complicated and, according to *Mathematica*, its general solution can be given in terms of integrals over hypergeometric functions

$$\begin{aligned}
h_1 = & z^{\frac{5\beta-R_1+3}{4\beta}} (2M+z)^{\frac{R_2-\beta-3}{4\beta}} \left[z^{\frac{R_1}{2\beta}} {}_2F_1 \left(\begin{matrix} \frac{R_1+R_2}{4\beta}, \frac{R_1+R_2}{4\beta}+1 \\ \frac{R_1}{2\beta}+1 \end{matrix}; -\frac{z}{2M} \right) \times \right. \\
& \times \left(c_1 + \int_a^z dx \frac{8(7\beta(\beta+2)+3)x^{\frac{3\beta-R_1-3}{4\beta}} (2M+x)^{\frac{3-3\beta-R_2}{4\beta}}}{MH(x)} {}_2F_1 \left(\begin{matrix} \frac{R_2-R_1}{4\beta}, \frac{R_2-R_1}{4\beta}+1 \\ 1-\frac{R_1}{2\beta} \end{matrix}; -\frac{x}{2M} \right) \right) \\
& + {}_2F_1 \left(\begin{matrix} \frac{R_2-R_1}{4\beta}, \frac{R_2-R_1}{4\beta}+1 \\ 1-\frac{R_1}{2\beta} \end{matrix}; -\frac{z}{2M} \right) \times \\
& \left. \times \left(c_2 - \int_a^z dx \frac{8(7\beta(\beta+2)+3)x^{\frac{3\beta+R_1-3}{4\beta}} (2M+x)^{\frac{3-3\beta-R_2}{4\beta}}}{MH(x)} {}_2F_1 \left(\begin{matrix} \frac{R_1+R_2}{4\beta}, \frac{R_1+R_2}{4\beta}+1 \\ \frac{R_1}{2\beta}+1 \end{matrix}; -\frac{x}{2M} \right) \right) \right], \tag{4.13}
\end{aligned}$$

where integration constants c_1, c_2 can be used to alter the lower boundaries of integration on each of the two integrals respectively.⁶ Abbreviations R_1, R_2 denote

$$R_1 = \sqrt{\beta(25\beta+42)+9}, \quad R_2 = \sqrt{\beta(\beta+18)+9}, \tag{4.14}$$

and the function $H(x)$ is given by

$$\begin{aligned}
H(x) = & x \left(\beta(8\beta-3R_1+7R_2+8) + 2(7R_2-5R_1) + 3(R_2-R_1) \right) \times \\
& \times {}_2F_1 \left(\begin{matrix} \frac{R_2-R_1}{4\beta}+1, \frac{R_2-R_1}{4\beta}+2 \\ 2-\frac{R_1}{2\beta} \end{matrix}; -\frac{x}{2M} \right) {}_2F_1 \left(\begin{matrix} \frac{R_1+R_2}{4\beta}, \frac{R_1+R_2}{4\beta}+1 \\ \frac{R_1}{2\beta}+1 \end{matrix}; -\frac{x}{2M} \right) \\
& + {}_2F_1 \left(\begin{matrix} \frac{R_2-R_1}{4\beta}, \frac{4\beta-R_1+R_2}{4\beta} \\ 1-\frac{R_1}{2\beta} \end{matrix}; -\frac{x}{2M} \right) \left[4(7\beta(\beta+2)+3)R_1M {}_2F_1 \left(\begin{matrix} \frac{R_1+R_2}{4\beta}, \frac{R_1+R_2}{4\beta}+1 \\ \frac{R_1}{2\beta}+1 \end{matrix}; -\frac{x}{2M} \right) \right. \\
& \left. - x \left(\beta(8\beta+3R_1+7R_2+8) + 2(5R_1+7R_2) + 3(R_1+R_2) \right) {}_2F_1 \left(\begin{matrix} \frac{R_1+R_2}{4\beta}+1, \frac{R_1+R_2}{4\beta}+2 \\ \frac{R_1}{2\beta}+2 \end{matrix}; -\frac{x}{2M} \right) \right]. \tag{4.15}
\end{aligned}$$

In order for the perturbations to make physical sense, the sub-leading terms h_2, h_3, h_4 may not grow large for $z \rightarrow \infty$, while the leading term h_1 is expected to grow with z . Making use of standard hypergeometric expansions as well as the initial differential equation (4.12), it is straightforward to determine that for $\beta \neq 0$:⁷

$$h_1|_{z \rightarrow \infty} = \frac{1}{2M^2\beta} z^2 + \mathcal{O}(z^2), \tag{4.16}$$

while equations (4.7) and (4.11) then reveal

$$\begin{aligned}
h_2|_{z \rightarrow \infty} &= \frac{1}{2} + \mathcal{O}(z^0), \\
h_3|_{z \rightarrow \infty} &= \frac{1}{4} \left(3 + \frac{1}{\beta} \right) + \mathcal{O}(z^0), \\
h_4|_{z \rightarrow \infty} &= -\frac{1+\beta}{4\beta} + \mathcal{O}(z^0), \tag{4.17}
\end{aligned}$$

⁶By default we have chosen lower boundaries of integration to be equal to the Kerr spin parameter a for $c_1 = c_2 = 0$, such that the metric deformation is smallest at the minimum of the effective scale factor (3.17).

⁷The degenerate case $\beta = 0$ causes the h_1 solution to scale with a power of z not consistent with the expectation of a radiation dominated universe. Therefore, we discard that case as unphysical.

where the small $\mathcal{O}(x^n)$ notation denotes terms that grow strictly slower than x^n . For the leading order perturbation to be positive at large z (4.16), we must enforce $\beta > 0$. The sub-leading perturbations (4.17) indeed do not grow with z , as required.

Next, we evaluate the metric on the radial geodesic trajectory, transformed to locally conformal time $t = f(\eta)$, $dt = f'(\eta)d\eta$. As in the previous section, we focus on the leading order metric only to model the observable universe

$$-d\tau^2 = -\frac{a^2 + z(2M + z)}{a^2 + z^2} f'(\eta)^2 \left(1 - \frac{a}{M}(1 + \beta) h_1(z)\right) d\eta^2 + \frac{a^2 + z^2}{a^2 + z(2M + z)} \left(1 + \frac{a}{M} h_1(z)\right) (d\mathbf{r}^2 + \mathbf{r}^2 d\phi^2 + dz^2) + \mathcal{O}(\mathbf{r}^2). \quad (4.18)$$

To find $z'(\eta)$ in case of a massless particle, we use (4.18) with $d\mathbf{r} = 0$, $d\phi = 0$, $d\tau = 0$, choose the positive square root solution and obtain⁸

$$z'(\eta) = \frac{a^2 + z(2M + z)}{a^2 + z^2} f'(\eta) \sqrt{\frac{1 - \frac{a}{M}(1 + \beta) h_1(z)}{1 + \frac{a}{M} h_1(z)}}. \quad (4.19)$$

Solving $-g_{00} = g_{33}$ for $f'(\eta)$ when we evaluate the metric on the trajectory $z = z(\eta)$ as in (3.14), we find

$$f'(\eta) = \frac{a^2 + z^2}{a^2 + z(2M + z)} \sqrt{\frac{1 + \frac{a}{M} h_1(z)}{1 - \frac{a}{M}(1 + \beta) h_1(z)}}. \quad (4.20)$$

Plugging (4.20) into (4.19), the relation simplifies analogously to (3.15):

$$z'(\eta) = 1, \quad \text{the simplest solution being } z(\eta) = \eta. \quad (4.21)$$

Therefore, transformed to conformal time and evaluated on the trajectory of interest, the perturbed leading order metric reads

$$-d\tau^2 = \frac{a^2 + \eta^2}{a^2 + \eta(2M + \eta)} \left(1 + \frac{a}{M} h_1(\eta)\right) (-d\eta^2 + d\mathbf{r}^2 + \mathbf{r}^2 d\phi^2 + dz^2) + \mathcal{O}(\mathbf{r}^2). \quad (4.22)$$

This means, that due to the introduction of the gravitating flow of radiation along the radial geodesic, the effective scale factor (3.17) is deformed to

$$\mathbf{a}(\eta) = \sqrt{\frac{a^2 + \eta^2}{a^2 + \eta(2M + \eta)} \left(1 + \frac{a}{M} h_1(\eta)\right)}. \quad (4.23)$$

From (4.16) we see that for late η this implies

$$\mathbf{a}(\eta)|_{\eta \gg M} \approx \left(\frac{a}{2\beta M^3}\right)^{\frac{1}{2}} \eta. \quad (4.24)$$

For a radiation dominated universe we expect $\mathbf{a}(t) \propto t^{1/2}$ and the transformation to conformal time therefore requires $t'(\eta)^2 \propto t(\eta)$, which is solved by $t(\eta) \propto \eta^2$. The result (4.24) is thus precisely consistent with our expectation for a radiation dominated universe. The free parameter β can be used to fit the proportionality factor to observation. See Figure 2 for a visualization of the effective scale factor.

⁸The relative minus sign in the square root looks like it may pose a problem. However, note that we introduced h_1 as a perturbation that is small thanks to the $a/M \rightarrow 0$ factor. In particular, we assume that M is large enough and a small enough such that $h_1(z) \ll M/a$ for all z of interest.




Figure 2. Comparison of effective scale factors (3.17) and (4.23), for demonstration evaluated at $c_1 = c_2 = 0$, $M = 1$, $a = 10^{-1}$, $\beta = 1$. Note that, due to only a single order of magnitude difference in M and a for visualization purposes, the radiation contribution appears dramatically exaggerated.

5 Conclusion and Outlook

In this work we have explored the scenario that our observable universe has been emitted from the ergosphere in a negative mass Kerr space-time, and is traveling along a radial geodesic parallel to the axis of rotation of the ring singularity. In appropriate cylindrical coordinates \mathbf{r}, z , assuming large enough mass parameter M such that the observable universe is well described by the leading terms in \mathbf{r}/M at large z , we have shown that the metric on the trajectory exhibits a particular conformal scale factor. The inherent contraction and expansion properties of the scale factor solve the horizon problem. Subsequently, we introduced a gravitating stream of radiation flowing statically along the trajectory, to model a radiation dominated universe. With an appropriate ansatz for metric perturbation, we then demonstrated that the resulting asymptotic growth of the scale factor indeed matches the expected result in a radiation dominated universe. As the features stemming from the black hole in the past die off asymptotically at large conformal time, the transition to a matter dominated universe could then be made utilizing the conventional FLRW metric.

One objection to working with the negative mass Kerr space-time is the existence of unbounded modes in linear time-dependent metric perturbations [24, 25], which make the space-time unstable. However, just the presence of an instability alone does not preclude a meta-stable state that may have a life-time of the age of the universe. Additionally, the fact that we are specifically working with only a patch of the space-time in the vicinity of the trajectory along the axis of rotation means that, especially in the case when radiation is present, a different global completion of the metric may exist that does not exhibit the same

instability.

The sub-leading terms in metric (3.6) vanish (or asymptote to a constant in metric (4.2), negligible with respect to the growing leading terms) for large z , which is achieved for late conformal time η on the trajectory. However, at early conformal time $\eta \approx 0$ the sub-leading terms can be relevant away from the absolute center of the trajectory. Therefore, it would be interesting to investigate if the sub-leading terms can imprint some signature onto the streaming radiation at early times, which could be measured at late times. Technically, the initial change of variables (3.3) is not completely unique. Choosing a slightly different coordinate transformation can change the early conformal time behavior of sub-leading terms and therefore potentially imprint different signatures onto the radiation. Similarly, a re-parametrization of the z -coordinate to any monotonously growing function starting at zero and linearly asymptoting to infinity may possibly change the quantitative details but not the qualitative picture.

Furthermore, the ansatz of metric perturbation (4.2) is physically motivated with the target form of the resulting leading order metric in mind, but not unique. It would be interesting to explore modifications of the ansatz, to potentially determine an exhaustive equivalence class of qualitatively similar perturbative solutions.

In the conventional inflationary paradigm, quantum fluctuations of the inflaton field and the metric tensor blow up to macroscopic size and seed the inhomogeneities that lead to structure formation such as galaxies. It would be interesting to investigate if introducing quantum fluctuations in the flowing radiation as well as the metric tensor in the scenario discussed in this paper can similarly lead to an appropriate amount of inhomogeneity to seed structure formation.

Acknowledgments

This work was supported by DOE grant DE-SC0011941.

References

- [1] W. Rindler, “Visual horizons in world-models,” *Mon. Not. Roy. Astron. Soc.* **116**, 662-677 (1956) doi:10.1023/A:1015347106729 [2?]
- [2] R. H. Dicke, “The Measurement of Thermal Radiation at Microwave Frequencies,” *Review of Scientific Instruments* **17**, 268 (1946) doi:10.1063/1.1770483
- [3] R. H. Dicke, P. J. E. Peebles, P. G. Roll and D. T. Wilkinson, “Cosmic Black-Body Radiation,” *Astrophys. J.* **142**, 414-419 (1965) doi:10.1086/148306
- [4] A. A. Starobinsky, “A New Type of Isotropic Cosmological Models Without Singularity,” *Phys. Lett. B* **91**, 99-102 (1980) doi:10.1016/0370-2693(80)90670-X
- [5] A. H. Guth, “The Inflationary Universe: A Possible Solution to the Horizon and Flatness Problems,” *Phys. Rev. D* **23**, 347-356 (1981) doi:10.1103/PhysRevD.23.347
- [6] D. Kazanas, “Dynamics of the Universe and Spontaneous Symmetry Breaking,” *Astrophys. J. Lett.* **241**, L59-L63 (1980) doi:10.1086/183361
- [7] K. Sato, “First Order Phase Transition of a Vacuum and Expansion of the Universe,” *Mon. Not. Roy. Astron. Soc.* **195**, 467-479 (1981) NORDITA-80-29.
- [8] A. D. Linde, “A New Inflationary Universe Scenario: A Possible Solution of the Horizon, Flatness, Homogeneity, Isotropy and Primordial Monopole Problems,” *Phys. Lett. B* **108**, 389-393 (1982) doi:10.1016/0370-2693(82)91219-9

- [9] A. Albrecht and P. J. Steinhardt, “Cosmology for Grand Unified Theories with Radiatively Induced Symmetry Breaking,” *Phys. Rev. Lett.* **48**, 1220-1223 (1982)
doi:10.1103/PhysRevLett.48.1220
- [10] A. D. Linde, “Chaotic Inflation,” *Phys. Lett. B* **129**, 177-181 (1983)
doi:10.1016/0370-2693(83)90837-7
- [11] V. F. Mukhanov and G. V. Chibisov, “Quantum Fluctuations and a Nonsingular Universe,” *JETP Lett.* **33**, 532-535 (1981)
- [12] L. Boubekeur and D. H. Lyth, “Hilltop inflation,” *JCAP* **07**, 010 (2005)
doi:10.1088/1475-7516/2005/07/010 [arXiv:hep-ph/0502047 [hep-ph]].
- [13] G. R. Dvali, Q. Shafi and S. Solganik, “D-brane inflation,” [arXiv:hep-th/0105203 [hep-th]].
- [14] F. C. Adams, J. R. Bond, K. Freese, J. A. Frieman and A. V. Olinto, “Natural inflation: Particle physics models, power law spectra for large scale structure, and constraints from COBE,” *Phys. Rev. D* **47**, 426-455 (1993) doi:10.1103/PhysRevD.47.426
[arXiv:hep-ph/9207245 [hep-ph]].
- [15] E. Silverstein and A. Westphal, “Monodromy in the CMB: Gravity Waves and String Inflation,” *Phys. Rev. D* **78**, 106003 (2008) doi:10.1103/PhysRevD.78.106003 [arXiv:0803.3085 [hep-th]].
- [16] F. L. Bezrukov and M. Shaposhnikov, “The Standard Model Higgs boson as the inflaton,” *Phys. Lett. B* **659**, 703-706 (2008) doi:10.1016/j.physletb.2007.11.072 [arXiv:0710.3755 [hep-th]].
- [17] H. Murayama, H. Suzuki, T. Yanagida and J. Yokoyama, “Chaotic inflation and baryogenesis by right-handed sneutrinos,” *Phys. Rev. Lett.* **70**, 1912-1915 (1993)
doi:10.1103/PhysRevLett.70.1912
- [18] J. Ellis, D. V. Nanopoulos and K. A. Olive, “No-Scale Supergravity Realization of the Starobinsky Model of Inflation,” *Phys. Rev. Lett.* **111**, 111301 (2013) [erratum: *Phys. Rev. Lett.* **111**, no.12, 129902 (2013)] doi:10.1103/PhysRevLett.111.111301 [arXiv:1305.1247 [hep-th]].
- [19] L. Kofman, A. D. Linde and A. A. Starobinsky, “Towards the theory of reheating after inflation,” *Phys. Rev. D* **56**, 3258-3295 (1997) doi:10.1103/PhysRevD.56.3258
[arXiv:hep-ph/9704452 [hep-ph]].
- [20] B. A. Bassett, S. Tsujikawa and D. Wands, “Inflation dynamics and reheating,” *Rev. Mod. Phys.* **78**, 537-589 (2006) doi:10.1103/RevModPhys.78.537 [arXiv:astro-ph/0507632 [astro-ph]].
- [21] G. Chardin, “Motivations for antigravity in general relativity,” *Hyperfine Interact.* **109**, no.1-4, 83-94 (1997) doi:10.1023/A:1012688930074
- [22] S. M. Carroll, “space-time and Geometry,” Cambridge University Press, 2019, ISBN 978-0-8053-8732-2, 978-1-108-48839-6, 978-1-108-77555-7
- [23] R. P. Kerr, “Gravitational field of a spinning mass as an example of algebraically special metrics,” *Phys. Rev. Lett.* **11**, 237-238 (1963) doi:10.1103/PhysRevLett.11.237
- [24] G. Dotti, “Linear Stability of Black Holes and Naked Singularities,” *Universe* **8**, no.1, 38 (2022)
doi:10.3390/universe8010038 [arXiv:2201.03640 [gr-qc]].
- [25] V. Cardoso and M. Cavaglia, “Stability of naked singularities and algebraically special modes,” *Phys. Rev. D* **74**, 024027 (2006) doi:10.1103/PhysRevD.74.024027 [arXiv:gr-qc/0604101 [gr-qc]].