# On Divisibility Property of Type 2 (p,q)-Analogue of r-Whitney Numbers of the Second Kind

Roberto B. Corcino Cristina B. Corcino

Research Institute for Computational Mathematics and Physics Cebu Normal University Cebu City, Philippines 6000 rcorcino@yahoo.com

August 10, 2022

#### Abstract

In this paper, the divisibility property of the type 2 (p, q)-analogue of the *r*-Whitney numbers of the second kind is established. More precisely, a congruence relation modulo pq for this (p, q)-analogue is derived.

**Keywords**: *r*-Whitney numbers, *r*-Dowling numbers, *A*-tableaux, convolution identities, binomial transform, Hankel transform.

### 1 Introduction

The *r*-Whitney numbers of the second kind were introduced by Mezo [22] as coefficients of the following generating function:

$$(mx+r)^n = \sum_{k=0}^n m^k W_{m,r}(n,k) x^{\underline{k}},$$

where  $x^{\underline{k}} = x(x-1)\dots(x-k+1)$ . These numbers satisfy the following properties:

1. the exponential generating function

$$\sum_{n=0}^{\infty} W_{m,r}(n,k) \frac{z^n}{n!} = \frac{e^{rz}}{k!} \left(\frac{e^{mz}-1}{m}\right)^k,$$

2. the explicit formula

$$W_{m,r}(n,k) = \frac{1}{m^k k!} \sum_{i=0}^k \binom{k}{i} (-1)^{k-i} (mi+r)^n,$$

3. the triangular recurrence relation

$$W_{m,r}(n,k) = W_{m,r}(n-1,k-1) + (km+r)W_{m,r}(n-1,k).$$

These properties are exactly the same properties that the  $(r, \beta)$ -Stirling numbers in [9] have possessed. This implies that the *r*-Whitney numbers of the second kind and the  $(r, \beta)$ -Stirling numbers are equivalent. More properties of these numbers can be found in [9, 22, 7].

One of the early studies on q-analogue of Stirling numbers of the second kind was introduced by Carlitz in [2] in connection with a problem in abelian groups. This is known as q-Stirling numbers of the second kind and is defined in terms of the following recurrence relation

$$S_q[n,k] = S_q[n-1,k-1] + [k]_q S_q[n-1,k], \quad [k]_q = \frac{1-q^k}{1-q}$$

such that, when  $q \to 1$ , this gives the triangular recurrence relation for the classical Stirling numbers of the second kind S(n,k)

$$S(n,k) = S(n-1,k-1) + kS(n-1,k).$$

Another version of definition of this q-analogue was adapted in [18] as follows

$$S_q[n,k] = q^{k-1}S_q[n-1,k-1] + [k]_qS_q[n-1,k].$$
(1)

Through this definition, the Hankel transform of q-exponential polynomials and numbers was successfully established, which may be considered as the Hankel transform of a certain q-analogue of Bell polynomials and numbers.

There are many ways to define q-analogue of Stirling-type and Bell-type numbers (see [14, 9, 12, 6, 8, 24]). However, in the desire to establish the Hankel transform of q-analogue of generalized Bell numbers, Corcino et al. [13] were motivated to define a q-analogue of r-Whitney numbers of the second kind parallel to that in (1) as follows:

$$W_{m,r}[n,k]_q = q^{m(k-1)-r} W_{m,r}[n-1,k-1]_q + [mk-r]_q W_{m,r}[n-1,k]_q.$$
(2)

Two more forms of this q-analogue, denoted by  $W_{m,r}^*[n,k]_q$  and  $\widetilde{W}_{m,r}[n,k]_q$ , were respectively defined by

$$W_{m,r}^*[n,k]_q := q^{-kr+m\binom{k}{2}} W_{m,r}[n,k]_q,$$
  
$$\widetilde{W}_{m,r}[n,k]_q := q^{-kr} W_{m,r}^*[n,k]_q = q^{-m\binom{k}{2}} W_{m,r}[n,k]$$

The corresponding q-analogues of generalized Bell numbers, also known as q-analogues of r-Dowling numbers, were also defined in three forms as (see [11, 13, 5])

$$D_{m,r}[n]_q := \sum_{k=0}^n W_{m,r}[n,k]_q,$$
$$D_{m,r}^*[n]_q := \sum_{k=0}^n W_{m,r}^*[n,k]_q,$$

and

$$\widetilde{D}_{m,r}[n]_q := \sum_{k=0}^n \widetilde{W}_{m,r}[n,k]_q.$$

where  $D_{m,r}[n]_q, D^*_{m,r}[n]_q$  and  $\widetilde{D}_{m,r}[n]_q$  denote the first, second and third form of the qanalogues of r-Dowling numbers, respectively. The Hankel transforms of  $D_{m,r}[n]_q, D^*_{m,r}[n]_q$ and  $\widetilde{D}_{m,r}[n]_q$  were successfully established in [5, 11, 13].

To extend these research studies, a certain (p,q)-analogue of r-Whitney numbers of the second kind, denoted by  $W_{m,r}[n,k]_{p,q}$ , was defined in [15] as coefficients of the following generating function:

$$[mt+r]_{p,q}^{n} = \sum_{k=0}^{n} W_{m,r}[n,k]_{p,q}[mt|m]_{p,q}^{k}$$
(3)

where

$$[t|m]_{p,q}^{\underline{n}} = \prod_{j=0}^{n-1} [t - jm]_{p,q}.$$
(4)

The orthogonality and inverse relations, an explicit formula, and a kind of exponential generating function of  $W_{m,r}[n,k]_{p,q}$  were already obtained. Unfortunately, its Hankel transform was not successfully established using the method applied in [5, 11, 13]. This motivated Corcino et al. [16] to define the type 2 (p,q)-analogue of r-Whitney numbers of the second kind, denoted by  $W_{m,r}[n,k;t]_{p,q}$ , as follows:

$$W_{m,r}[n+1,k;t]_{p,q} = q^{m(k-1)+r}W_{m,r}[n,k-1;t]_{p,q} + [mk+r]_{p,q}p^{mt-km}W_{m,r}[n,k;t]_{p,q}.$$
 (5)

The second form was then defined as follows:

$$W_{m,r}^*[n,k;t]_{p,q} := q^{-kr - m\binom{k}{2}} W_{m,r}[n,k;t]_{p,q}.$$
(6)

Several properties of these (p, q)-analogues were established in [16] including their Hankel transforms, which are given by

$$\det (W_{m,r}[s+i+j,s+j;t]_{p,q})_{0 \le i,j \le n} = \prod_{k=0}^{n} q^{m\binom{s+k}{2} + (s+k)r} p^{nmt} [m(s+k)+r]_{p,q}^{k}$$
$$\det (W_{m,r}^{*}[s+i+j,s+j;t]_{p,q})_{0 \le i,j \le n} = \prod_{k=0}^{n} p^{nmt} [m(s+k)+r]_{p,q}^{k}.$$

On the other hand, the first, second and third forms of type 2 (p, q)-analogue of the *r*-Dowling numbers, denoted by  $D_{m,r}[n]_{p,q}$ ,  $D^*_{m,r}[n]_{p,q}$  and  $\widetilde{D}_{m,r}[n]_{p,q}$  were defined respectively in [16] as follows:

$$D_{m,r}[n]_{p,q} := \sum_{k=0}^{n} W_{m,r}[n,k;t]_{p,q},$$
$$D_{m,r}^{*}[n]_{p,q} := \sum_{k=0}^{n} W_{m,r}^{*}[n,k;t]_{p,q},$$
$$\widetilde{D}_{m,r}[n]_{p,q} := \sum_{k=0}^{n} \widetilde{W}_{m,r}[n,k;t]_{p,q},$$

where  $\widetilde{W}_{m,r}[n,k;t]_{p,q} = q^{kr}W^*_{m,r}[n,k;t]_{p,q}$  denotes the third form of the (p,q)-analogue of the *r*-Whitney numbers of the second kind. Among these three forms, only the second form was provided a Hankel transform, which is given by

$$H(D_{m,r}^*[n]_{p,q}) = \left(\frac{q}{p}\right)^{\frac{n(n^2+3n+8)}{6}+r-1\binom{n}{2}} ([m]_{\frac{q}{p}})^{\binom{n}{2}} \prod_{k=0}^{n-1} [k]_{\left(\frac{q}{p}\right)^m}!$$

The main objective of this study is to establish additional property of the type 2 (p, q)analogues of the *r*-Whitney numbers of the second kind. More precisely, the divisibility
property of these type 2 (p, q)-analogues will be discussed thoroughly.

### 2 Preliminary Results

This section provides a brief discussion on some relations that are necessary in deriving the divisibility property of the type 2 (p,q)-analogue of the *r*-Whitney numbers of the second kind  $W_{m,r}^*[n,k;t]_{p,q}$ .

Multiplying both sides of the recurrence relation in (5) by  $q^{-kr-m\binom{k}{2}}$  yields

$$q^{-kr-m\binom{k}{2}}W_{m,r}[n+1,k;t]_{p,q} = q^{-kr-m\binom{k}{2}}q^{m(k-1)+r}W_{m,r}[n,k-1;t]_{p,q} + q^{-kr-m\binom{k}{2}}[mk+r]_{p,q}p^{mt-km}W_{m,r}[n,k;t]_{p,q} q^{-kr-m\binom{k}{2}}W_{m,r}[n+1,k;t]_{p,q} = q^{-(k-1)r-m\binom{k-1}{2}}W_{m,r}[n,k-1;t]_{p,q} + [mk+r]_{p,q}p^{mt-km}q^{-kr-m\binom{k}{2}}W_{m,r}[n,k;t]_{p,q}$$

Applying (6) consequently gives

$$W_{m,r}^*[n+1,k;t]_{p,q} = W_{m,r}^*[n,k-1;t]_{p,q} + [mk+r]_{p,q}p^{mt-km}W_{m,r}^*[n,k;t]_{p,q}.$$
(7)

This relation can be used to generate the following first few values of  $W_{m,r}^*[n,k;t]_{p,q}$ :

n/k	0	1	2	3
0	1			
1	$[r]_{p,q}p^{mt}$	1		
2	$[r]_{p,q}^2 p^{2mt}$	$[r]_{p,q}p^{mt} + [m+r]_{p,q}p^{m(t-1)}$	1	
3	$[r]_{p,q}^{3}p^{3mt}$	$[P, p, q]^{r}$ $[P, q]^{r}$ $[P, q]^{r}$	$[r]_{p,q}p^{mt} + 2[m+r]_{p,q}p^{m(t-1)}$	1
		$+[m+r]_{p,q}^2 p^{2m(t-1)}$		

Table 1. The First Values of  $W_{m,r}^*[n,k;t]_{p,q}$ 

By repeated application of (7), we can easily derive the following vertical recurrence relation.

**Theorem 2.1.** For nonnegative integers n and k, and real number r, the (p,q)-analogue of r-Whitney numbers of the second kind satisfies the following vertical recurrence relation

$$W_{m,r}^*[n+1,k+1;t]_{p,q} = \sum_{j=k}^n [m(k+1)+r]_{p,q}^{n-j} p^{(n-j)[mt-(k+1)m]} W_{m,r}^*[j,k;t]_{p,q}.$$
 (8)

One can easily verify relation (8) using the values of  $W_{m,r}^*[n,k;t]_{p,q}$  in Table 1.

Now, let us derive the rational generating function for  $W_{m,r}^*[n,k;t]_{p,q}$ . Suppose that

$$\Psi_k^*(x) = \sum_{n=k}^{\infty} W_{m,r}^*[n,k;t]_{p,q} x^{n-k}.$$

When k = 0, (7) reduces to

$$W_{m,r}^*[n+1,0;t]_{p,q} = [r]_{p,q} p^{mt} W_{m,r}^*[n,0;t]_{p,q}$$

By repeated application of (7), this inductively gives

$$W_{m,r}^{*}[n+1,0;t]_{p,q} = [r]_{p,q}p^{mt}W_{m,r}^{*}[n,0;t]_{p,q} = \left([r]_{p,q}p^{mt}\right)^{2}W_{m,r}^{*}[n-1,0;t]_{p,q}$$
  
$$\vdots$$
  
$$= \left([r]_{p,q}p^{mt}\right)^{n+1}W_{m,r}^{*}[0,0;t]_{p,q} = \left([r]_{p,q}p^{mt}\right)^{n+1}.$$

Hence,

$$\Psi_0^*(x) = \sum_{n=0}^{\infty} W_{m,r}^*[n,0;t]_{p,q} x^n = \frac{1}{(1-xp^{mt}[r]_{p,q})}$$

When k > 0 and applying the triangular recurrence relation in (5), we have

$$\Psi_k^*(x) = \sum_{n=k}^{\infty} W_{m,r}^*[n,k;t]_{p,q} x^{n-k}$$
  
=  $\sum_{n-1=k-1}^{\infty} W_{m,r}^*[n-1,k-1;t]_{p,q} x^{(n-1)(k-1)}$   
+  $xp^{mt-km}[mk+r]_{p,q} \sum_{n-1=k}^{\infty} W_{m,r}^*[n-1,k;t]_{p,q} x^{n-1-k}$   
=  $\Psi_{k-1}^*(x) + xp^{m(t-k)}[mk+r]_{p,q} \Psi_k^*(x)$ 

Solving for  $\Psi_k^*(t)$  yields

$$\Psi_k^*(x) = \frac{1}{1 - x p^{m(t-k)} [mk+r]_{p,q}} \Psi_{k-1}^*(x).$$

Applying backward substitution gives the following rational generating function for  $W_{m,r}[n,k;t]_{p,q}$ .

**Theorem 2.2.** For nonnegative integers n and k, and real number r, the (p,q)-analogue  $W_{m,r}[n,k;t]_{p,q}$  satisfies the following rational generating function

$$\Psi_k^*(x) = \sum_{n=k}^{\infty} W_{m,r}^*[n,k;t]_{p,q} x^{n-k} = \frac{1}{\prod_{j=0}^k (1 - xp^{m(t-j)}[mj+r]_{p,q})}.$$
(9)

**Remark 2.3.** This rational generating function plays an important role in proving the main result of the paper.

# 3 Divisibility Property

In this section, the congruence relation modulo pq for the type 2 (p,q)-analogue of the *r*-Whitney numbers of the second kind  $W_{m,r}^*[n,k;t]_{p,q}$  will be established using the rational generating function in (9).

Using the values of  $W_{m,r}^*[n,k;t]_{p,q}$  in Table 1, we observe that, with

$$[t]_{p,q} = p^{t-1} + p^{t-2}q + p^{t-3}q^2 + \ldots + pq^{t-2} + q^{t-1},$$

the polynomial expressions of  $W_{m,r}^*[n,k]_q$  from row 0 to row 3, if they are divided by pq, the remainders form the following triangle of expressions in p:

$$\begin{array}{c}
1\\
p^{mt+r-1} & 1\\
p^{2(mt+r-1)} & 2p^{mt+r-1} & 1\\
p^{3(mt+r-1)} & 3p^{2(mt+r-1)} & 3p^{mt+r-1} & 1.
\end{array}$$

This can further be written as

$$\begin{pmatrix} \begin{pmatrix} 0\\0 \end{pmatrix} \\ \begin{pmatrix} 1\\0 \end{pmatrix} p^{mt+r-1} & \begin{pmatrix} 1\\1 \end{pmatrix} \\ \begin{pmatrix} 2\\0 \end{pmatrix} p^{2(mt+r-1)} & \begin{pmatrix} 2\\1 \end{pmatrix} p^{mt+r-1} & \begin{pmatrix} 2\\2 \end{pmatrix} \\ \begin{pmatrix} 3\\0 \end{pmatrix} p^{3(mt+r-1)} & \begin{pmatrix} 3\\1 \end{pmatrix} p^{2(mt+r-1)} & \begin{pmatrix} 3\\2 \end{pmatrix} p^{mt+r-1} & \begin{pmatrix} 3\\3 \end{pmatrix},$$

To generalize this observation, the next theorem contains the divisibility property of  $W_{m,r}^*[n,k;t]_{p,q}$ .

**Theorem 3.1.** For nonnegative integers n and k, the type 2 (p,q)-analogue of the r-Whitney numbers of the second kind  $W_{m,r}[n,k;t]_{p,q}$  satisfies the following congruence relation

$$W_{m,r}^*[n,k;t]_{p,q} \equiv \binom{n}{k} p^{(n-k)(mt+r-1)} \mod pq.$$

$$\tag{10}$$

*Proof.* The polynomial  $[t]_{p,q}$  can be written as

$$[t]_{p,q} = p^{t-1} + q^{t-1} + pqy,$$

where y is a polynomial in p and q. Then, we have

$$\frac{1}{\prod_{j=0}^{k} (1 - xp^{m(t-j)}[mj+r]_{p,q})} = \sum_{n=0}^{\infty} \left( xp^{m(t-j)}[mj+r]_{p,q} \right)^{n}$$
$$= \sum_{n=0}^{\infty} p^{nm(t-j)} (p^{mj+r-1} + q^{mj+r-1} + pqy)^{n} x^{n}$$
$$= \sum_{n=0}^{\infty} p^{n(mt+r-j)} x^{n} + pq \sum_{n=0}^{\infty} \hat{z}_{n} x^{n},$$

where  $\hat{z}_n$  is a polynomial in p and q. It follows that

$$\frac{1}{\prod_{j=0}^{k} (1 - xp^{m(t-j)}[mj+r]_{p,q})} = \sum_{n=0}^{\infty} p^{n(mt+r-j)} x^n \mod pq$$
$$= \frac{1}{1 - p^{mt+r-1}x} \mod pq.$$

Thus, using (9), we have

$$\begin{split} \sum_{n=k}^{\infty} W_{m,r}^*[n,k;t]_{p,q} x^{n-k} &\equiv \frac{1}{(1-p^{mt+r-1}x)^{k+1}} \mod pq \\ &\equiv \sum_{n=0}^{\infty} \binom{n+(k+1)-1}{n} p^{n(mt+r-1)} x^n \mod pq \\ &\equiv \sum_{n=k}^{\infty} \binom{n}{k} p^{(n-k)(mt+r-1)} x^{n-k} \mod pq. \end{split}$$

Comparing the coefficients of  $x^{n-k}$  completes the proof of the theorem.

**Remark 3.2.** Using (6) and Theorem 3.1, the first form of the of the type 2(p,q)-analogues of the r-Whitney numbers of the second kind satisfies the following congruence relation modulo pq:

$$W_{m,r}[n,k;t]_{p,q} \equiv \binom{n}{k} p^{(n-k)(mt+r-1)} q^{kr+m\binom{k}{2}} \mod pq$$
$$\equiv \begin{cases} q^{nr+m\binom{n}{2}} \mod pq, & \text{for } n=k\\ 0 \mod pq, & \text{otherwise} \end{cases}$$

## References

- Aigner, M., A Characterization of the Bell Numbers. Discrete Math. 205 (1999), 207– 210.
- [2] Carlitz, L., q-Bernoulli numbers and polynomials. Duke Math. J. 15 (1948) 987-1000.
- [3] Cheon, G.C. and Jung, J.H., r-Whitney Numbers of Dowling Lattices. Discrete Math. 312 (2012), 2337–2348.
- [4] Comtet, L., Advanced Combinatorics, Reidel, Dordrecht, The Netherlands, 1974.
- [5] Corcino R.B., Hankel Transform of the first form (q, r)-Dowling Numbers. Eur. J. Pure Appl. Math. 14(3) (2021), 746–759.
- [6] Corcino, R.B. and Corcino, C.B., The Hankel Transform of Generalized Bell Numbers and Its q-Analogue, Utilitas Mathematica, 89 (2012), 297-309.
- [7] Corcino, R.B., Corcino, C.B. and Aldema, R., Asymptotic Normality of the  $(r, \beta)$ -Stirling Numbers, Ars Combin., **81** (2006), 81-96.
- [8] Corcino, R.B., Corcino, C.B., Ontolan, J.M., Perez, C.M. and Cantallopez, E.R., The Hankel Transform of q-Noncentral Bell Numbers. International Journal of Mathematics and Mathematical Sciences. 2015 (2015)
- [9] Corcino, R.B., The  $(r,\beta)$ -Stirling numbers. Mindanao Forum. 14(2) (1999), 91–100.
- [10] Corcino, R.B. and Corcino, C.B., The Hankel Transform of Generalized Bell Numbers and Its q-Analogue, Utilitas Mathematica, 89 (2012), 297-309.
- [11] Corcino R.B., Ontolan, J.M. and Rama, G., Hankel Transform of the Second Form (q, r)-Dowling Numbers, Eur. J. Pure Appl. Math. 12(4) (2019), 1676-1688.
- [12] Corcino R.B., Ontolan, J.M., Cañete, J. and Latayada, M. R., A q-analogue of r-Whitney numbers of the second kind and its Hankel transform, J. Math. Computer Sci. 21 (2020), 258–272.
- [13] Corcino R.B., Latayada, M.R. and Vega, M.P. Hankel Transform of (q, r)-Dowling Numbers. Eur. J. Pure Appl. Math. 12(2) (2019), 279–293.
- [9] Corcino, R. B. and Barrientos, C., Some Theorems on the q-Analogue of the Generalized Stirling Numbers, Bulletin of the Malaysian Mathematical Sciences Society, 34(3) (2011), 487-501.
- [14] Corcino, R.B., Hsu, L.C., Tan, E.L., A q-Analogue of Generalized Stirling Numbers, Fibonacci Quart., 44 (2006), 154-165.
- [15] Corcino R.B., Vega, M.P. and Dibagulun, A.M., A (p,q)-analogue of Qi-type formula for r-Dowling numbers, J. Math. Computer Sci. 24 (2021), 273–286.

- [16] Corcino R.B., Vega, M.P. and Dibagulun, A.M., A (p,q)-analogue of Qi-type formula for r-Dowling numbers, Axioms. 10 (2021), 343.
- [17] de Médicis, A. and Leroux, P., Generalized Stirling Numbers, Convolution Formulae and (p, q)-analogues, Can. J. Math 47(3) (1995), 474–499.
- [18] Ehrenborg, R., Determinants involving q-Stirling numbers, Adv. in Appl. Math., 31 (2003), 630–642.
- [19] Layman, J.W., The Hankel Transform and Some of its Properties. Journal of Integer Sequences, 4 (2001), Article 01.1.5.
- [20] Mangontarum, M.M., Macodi-Ringia, A.P. and Abdulcarim, N.S. The Translated Dowling Polynomials and Numbers. *International Scholarly Research Notices*. 2014 (2014), Article ID 678408, 8 pages
- [21] Mangontarum, M.M., Cauntongan, O.I. and Macodi-Ringia, A.P., The Noncentral Version of the Whitney Numbers: A Comprehensive Study. *International Journal of Mathematics and Mathematical Sciences.* **2016** (2016), Article ID 6206207, 16 pages
- [22] Mező, I., A New Formula for the Bernoulli Polynomials. *Result Math.* **58**(3) (2010), 329–335.
- [23] Mező, I., The r-Bell numbers, J. Integer Seq., 14 (2011)
- [24] Montero, C.B. and Corcino, R.B., A q-analogue of Rucinski-Voigt numbers. ISRN Discrete Mathematics, 2012 (2012) 592818, 18 pages.
- [25] Sloane N.J., Least positive sequence with Hankel transform {1, 1, 1, 1, 1, 1, ...}. The On-Line Encyclopedia of Integer Sequences, http://oeis.org., 2000.
- [26] Spivey, M.Z. and Steil, L.L., The k-binomial transform and Hankel transform. J. Integer Seq,9 (2006) Article 06.1.1
- [27] Riordan, J., Combinatorial Identities, Wiley, New York, NY, USA, 1968.