

Precise Calculations of Nucleosynthesis Parameters and Electron Self-mass

Samina Masood and Jaskeerat Singh
Department of Physical and Applied Sciences
University of Houston Clear Lake

Abstract

We study the effect of temperature on beta decay rate during primordial nucleosynthesis. Using thermal contributions to the renormalized mass of electron, we re-compute thermal effects to the nucleosynthesis parameters in the early universe in relation to the thermal self-mass of electron. In this study we show how the presence of fermions in a medium cause the variation in nucleosynthesis parameters with temperature. Before and after nucleosynthesis, temperature contribution from the electron self-mass goes away. The temperature dependence of beta decay rate, helium abundance and energy density of the universe are calculated as a function of temperature during nucleosynthesis. The values of these nucleosynthesis parameters before and after the nucleosynthesis are also calculated.

PACS Nos: 11.10 Wx;, 11.10 Gh; 12.20 -m; 98.80-k; 14.60 Cd

Key words: Electron self-mass, Beta Decay, Primordial nucleosynthesis, Early Universe

1. Introduction

Beta decay is a process of spontaneous decay of neutron into proton and its cross-section is simply related to the mass of electron. The statistical background contribution to electron self-mass [1-4] is calculated up to the two-loop level in literature [5] and has a measurable effect on particle processes including beta decay, which in turn contributes to helium synthesis [6-7] in the primordial universe. The radiative corrections to electroweak processes [8] are not significant in a hot media in the absence of required distribution of hot electrons. The statistical contribution to the self-mass of electron is calculated using the renormalization scheme [9] of QED (quantum electrodynamics) in real-time formalism [10-12]. In this scheme of calculations, the order-by-order cancellation of singularities can be shown by simply adding all of the same order diagrams. Real-time formalism is the only formalism to incorporate thermal effects and still show the cancellation of singularities clearly. We can prove Kinoshita-Lee-Nauenberg (KLN) theorem [11-12] in QED only in real-time formalism. Renormalization cannot be proved at each order of perturbation in the imaginary time formalism. Another benefit of this formalism is that we can separate out thermal corrections from T independent corrections easily. At high temperatures, including higher order corrections, electroweak theory has to be incorporated for beta decay. These contributions are significant in the early universe when temperature is above the mass of the electron (which correspond to roughly 10^9 K).

Nucleosynthesis in the early universe took place around 1 MeV temperature and densities were low enough to be ignored. Thermal corrections were needed to be incorporated for precise calculation of beta decay rate, and the related processes [13--14] for the correct estimate of helium synthesis in the early universe. Self-mass of electron clearly has the major thermal correction as neutrino do not directly couple

with radiation. Therefore, detailed study of self-mass of electron during nucleosynthesis is required for better estimate of beta decay rate and consequently helium synthesis, energy density and even the expansion rate of the universe during that period. For a better understanding of self-mass effects we need to revisit the electron self-mass during nucleosynthesis in the early universe.

Next section is devoted to a quick review of the calculation of electron self-mass at finite temperature. A detailed study of the electron self-mass during nucleosynthesis shows that the fermion background cannot be there even around the temperatures close to electron mass. The behavior of electron self-mass navigates the behavior of other parameters in the early universe; such as beta decay rate, helium abundance, energy density of the universe and the expansion rate of the universe during nucleosynthesis. We study thermal dependence of these parameters, during nucleosynthesis, in Section 3. The last Section 4 is devoted to the discussion of results in this paper.

2. Self-mass of the electron at Finite Temperature

Self-mass of the electron in the early universe is computed using the renormalization scheme of QED in real-time formalism. We calculated the dependence of self-mass on temperature in terms of Masood's "abc" functions which were first introduced in Ref. [2]. Calculation of the most dominant contributions of the medium (at high energies) to the electron mass for high temperatures and ignorable densities, up to the first order in alpha, has been computed as [2]

$$\frac{\delta m}{m} = \frac{\alpha \pi T^2}{3m^2} [\{1 - c(m\beta)\} + b(m\beta) + a(m\beta)], \quad (1)$$

which is found to be applicable to all temperatures higher than the electron mass, whereas Masood's "abc" functions are calculated in detail at high temperatures [2] and have been used in several published papers earlier. To calculate thermal corrections, covariant hot loop integrals are evaluated exactly. These abc functions can be written as

$$a(m\beta) = \ln(1 + e^{-m\beta}), \quad (2a)$$

$$b(m\beta) = \sum_{n=1}^{\infty} (-1)^n Ei(-nm\beta), \quad (2b)$$

$$c(m\beta) = \sum_{n=1}^{\infty} (-1)^n \frac{e^{-nm\beta}}{n^2}. \quad (2c)$$

These functions can give a negative contribution for $T \leq m$ and may give a negative contribution to self-mass which is an unphysical situation and is discussed later in detail.

Using the renormalization scheme of QED, the renormalized mass of electron in a medium is expressed as the physical mass, which is given as:

$$m_{phys} = m \left(1 + \frac{\delta m}{m}\right) \quad (3)$$

The evaluation of how self-mass of the electron in different thermal regions helps to understand the behavior of electrons in various statistical environments, which helps to compute the effect of electron mass change under statistical conditions. The above equations reveal the self-mass contribution in the extremely hot condition of the early universe above and below the nucleosynthesis temperatures, with and

without the fermion contributions [2], respectively. We can then write the self-mass, after nucleosynthesis ($T \leq m$), due to the significant radiation background, that is derived from the general T Eq. (1)

$$\frac{\delta m}{m} = \frac{\alpha \pi T^2}{3m^2} \quad (4a)$$

. Whereas, for temperatures sufficiently larger than electron mass ($T \gg m$ before nucleosynthesis), we incorporate the hot electron background along with radiation. The electron self-mass corrections are expressed as:

$$\frac{\delta m}{m} = \frac{\alpha \pi T^2}{2m^2} \quad (4b)$$

However, “abc” functions describe the functional dependence of self-mass and the relevant impact on the beta decay rates during nucleosynthesis. In this way, electron mass, beta decay rate, and helium abundance change with respect to their vacuum values, and all become functions of temperature.

The general contribution of self-mass of electron in units of the rest-mass of electron is given in Eq. (1). We use this general equation to compute thermal contributions for the entire range of temperatures and plot it as a solid line in Fig. (1). A comparison with the low temperature effect (with no fermion background) is calculated from Eq. (4a) and the high temperature contribution (including fermion background) to electron mass, using Eq. (4b). Fig. (1) shows a direct comparison of high temperature and low temperature behaviors of the electron self-mass with the generally computed self-mass values from “abc” functions. As mentioned earlier, the negative contribution of “abc” functions gives negative values to self-mass for $T \leq m$, thus revealing that free hot fermions will not radiatively interact in the background at lower T . This happens because enough concentration of free hot electrons is not found at low temperatures. Exponential dependence of the Fermion distribution function on temperature contributes negatively to overcome the thermal correction of photon background ($\frac{\alpha \pi T^2}{3m^2}$), giving an overall negative result until $T > 0.6$ MeV.




Fig.1: Low T line correspond to Eq.(4a) High T line to Eq. (4b) and general T to Eq. (1)

This behavior is changed for T slightly greater than m . The significance of “abc” functions is to describe the thermal behavior during nucleosynthesis, mainly. A plot of these functions in Fig. (1) shows that the fermion contribution from the medium seems to coincide with high T behavior after $T=1.2$ MeV. This is around 2.5 times the electron mass. At this point, Fermion contribution is almost comparable to photon

background contributions and contribution of $c(m\beta)$ dominates over the contributions of $b(m\beta)$ and $a(m\beta)$ [2].

A consolidated graph of self-mass as a function of temperature is shown in Fig. (2), where we calculate the low temperature contribution from Eq. (4a) and high temperature contribution from Eq. (4b). A little disconnected region indicated by an arrow shows the point where low temperature and high temperature behavior start to differ. This is the region where nucleosynthesis actually took place. For a heating system, going from lower temperature, neutrino emission starts due to beta decay processes. This is the region that is studied in detail in this paper using “abc” functions.




Fig. (2): Thermal correction to self-mass of lepton as a function of temperature using Eqs. (4a and 4b) in the relevant ranges. m corresponds to lepton mass and T will vary for every value of m .

Self-mass below the temperature of electron mass is almost a straight line and is only a very small fraction of the particle’s rest mass, whereas the self-mass above the electron mass increases exponentially for some time until the temperature is too high for nucleosynthesis. Beyond $T = 2$ MeV (the decoupling temperature), the nucleosynthesis stops and self-mass increases quadratically on the ratio between temperature and electron mass ($\sim T^2/m^2$). So, the electron mass increases significantly at high temperatures before nucleosynthesis. During nucleosynthesis, at $T \sim 1$ MeV, the rise in electron temperature is not ignorable as it is related to the burning of Primordial Hydrogen into Helium. However, this graph depicts that the thermal corrections to electron mass are easily ignorable after nucleosynthesis stops. Table 1 shows a comparison of calculated values of self-mass for various temperatures at T/m greater than 1, indicating that the dominant high temperature contribution is a quadratic function of T as given in Eq. (4b). These corrections are extremely small for T/m up to 1 and it increases almost exponentially during nucleosynthesis.

The computed contributions are using Masood’s functions, or in other words, are including the fermion contribution from “abc” functions. Whereas the low temperature contribution is mainly coming from the interaction of radiation with electrons only. The degenerate pressure of fermions may not allow the creation of electrons at low temperature beyond certain limit. That explains why the self-mass reduces the physical mass at temperatures below the electron mass. Therefore, at low temperature, the presence of

electrons reduces the self-mass instead of increasing. It means that at low temperatures, the free electrons are not allowed to stay inside a system.

We have compared the contributions of Masood's "abc" functions during nucleosynthesis, as compared to the high temperature expansion of Eq. (4b) for $T \gg m$, as prior to the start of nucleosynthesis in the early universe. This is shown in Table 1. It shows that the significance of self-mass corrections depends a lot on "abc" functions during nucleosynthesis and their contribution is required for the precise calculations of decay rate. All these values are plotted in Fig. (3) which shows that the self-mass contribution via "abc" functions is maximized around the temperatures near the electron mass and just approximately reduced to quadratic dependence of T/m around 1.5 MeV.

T	Computed with "abc" functions	Calculated High T value	Percentage Difference
0.6	0.001683382	0.015814	89.35511572
0.7	0.008703485	0.021524	59.56381249
0.8	0.018555127	0.028113	33.99805428
0.9	0.0284381	0.035581	20.07504005
1	0.038655771	0.043927	11.99997496
1.1	0.049383453	0.053152	7.090132074
1.2	0.06072877	0.063255	3.993723816
1.3	0.072761123	0.074237	1.988061209
1.4	0.085527131	0.086097065	0.66196774
1.5	0.099059208	0.098835917	0.225920572

Table 1: Computed values of electron self-mass near primordial nucleosynthesis temperatures using "abc" functions and the approximate values for $T < m$ and $T > m$ in units of m

This temperature may exactly indicate the temperature (~ 1.2 MeV) where nucleosynthesis started to contribute, significantly, to helium synthesis in the early universe. This temperature is reached right after the universe is cooled to neutrino decoupling temperature ($T \sim 2$ MeV).




Fig.3: Percentage deviation from quadratic dependence of self-mass on temperature

The overall behavior of the self-mass of electrons, as a function of temperature for $0.6\text{MeV} < T < 1.6\text{ MeV}$, is plotted in Fig. (4) to closely study nucleosynthesis. Fermion background starts to add thermal mass around 0.6 MeV . The increase in mass decreases the decay rate according to the regular calculations of decay rates.




Fig. (4): Plot of the self-mass of electron during nucleosynthesis. Increase of self-mass with temperature corresponds to decrease in physical mass of electron with cooling of the universe.

In the next section, we study the impact of temperature on nucleosynthesis parameters including beta decay and helium yield in the early universe. We can also see the impact of mass-change on the energy density of the universe and the Hubble parameter. The temperature dependence is induced to all nucleosynthesis parameters through the thermal mass of the electron. Beta type decay processes are possible for higher leptonic generations, but they are suppressed by the large masses of the muon and tauon, even in a vacuum. Thermal contributions come from the medium and the existence of enough concentration of fermions depends on their mass. The existence of enough free muons and heavier particles cannot be ruled out in the very early universe where temperatures were possible to express in muon mass and tauon mass, however nucleosynthesis does not occur there. Similarly, the mass of the

proton is taken constant as its direct interaction with radiation and charge is significantly suppressed by heavy baryonic mass near nucleosynthesis.

3. Temperature Dependence of Nucleosynthesis Parameters in the Early Universe

It is well-understood that the increase in electron mass reduces the decay rate for heavier mass and increased temperature decreases the possibility of decay rate and hence decreases the rate of nucleosynthesis. The beta decay rate in the early universe is proportional to the thermally corrected self-mass of the electron. A direct calculation of nucleosynthesis parameters as functions of self-mass are already calculated in Ref. [9] and have been used later in literature.

Thermal contributions to electron mass are found to be related to beta decay rate $\frac{\delta\lambda}{\lambda}$, and hence the helium abundance $\frac{\delta Y}{Y}$ in the early universe. We use these relations from Ref. [6,7,9] and the references provided there. There are several processes that can contribute to net beta decay such as neutrino capture or inverse beta decay.

$$\frac{\delta\lambda}{\lambda} = -0.2 \frac{\delta m}{m} \left(\frac{m}{T}\right)^2 \quad (6)$$

$$\frac{\delta Y}{Y} = -0.2 \frac{\delta\lambda}{\lambda} = 0.04 \frac{\delta m}{m} \left(\frac{m}{T}\right)^2 \quad (7)$$

Eqs. (6) and (7) show that the beta decay rate and the helium abundance parameter are proportional to the self-mass and vary relatively slowly with temperature, respectively. These functions are plotted as a function of temperature in Fig (5).




Fig. (5): Plot of the thermal contribution to beta decay rate during nucleosynthesis. The beta decay rate increases with lowering of temperature and then reduces because nucleosynthesis slows down around 0.5 MeV

Fig. (5) shows a plot of beta decay as a function of temperature. However, thermal effects to helium synthesis are not necessarily ignorable. It is known that helium yield is around 25% in the early universe. Therefore, $0.003/0.25$ gives a contribution between 1-2 percent after the completion of nucleosynthesis. A plot of thermal corrections to helium yield Y is given in Fig. (6)




Fig. (6): Plot of the thermal contribution to helium abundance during nucleosynthesis. The beta decay rate increases with the lowering of temperature and then reduces because nucleosynthesis slows down around 0.5 MeV.

On the other hand, the modification in the energy density of the universe with respect to temperature becomes:

$$\frac{\delta \rho}{\rho} = -\frac{\delta m}{m} \left(\frac{m}{4T} \right)^2 \quad (8)$$

The plot of the variation in energy density of the universe during nucleosynthesis is shown in Fig. (7). This shows the increase in energy density with the decrease in temperature. However, energy density is independent of temperature before or after nucleosynthesis as well. Similar behavior is expected by the Hubble parameter as well and is given by Fig. (7), which gives a plot of equation (9).




Fig. (7): Plot of the thermal contribution to the energy density of the universe during nucleosynthesis. Thermal contribution to the energy density increases with lowering of temperature and then then become independent of time after nucleosynthesis stops.

The Hubble constant H changes with respect to temperature as compared to the calculated value of H as:

$$\frac{\delta H}{H} = -0.5 \frac{\delta m}{m} \left(\frac{m}{4T} \right)^2 \quad (9)$$

These plots show that the electron background introduces thermal contribution to nucleosynthesis due to the temperature dependence of electron mass on temperature.




Fig. (8): Plot of the thermal contribution to the expansion rate of the universe during nucleosynthesis. The contribution to Hubble parameter is reduced after nucleosynthesis stops.

In this paper we concentrate on the study of the effect of thermal mass on beta decay and its impact on nucleosynthesis parameters in the early universe.

4. Results and Discussions

A detailed study of the electron self-mass during nucleosynthesis shows that the fermion background cannot be there even around the temperatures close to electron mass. At $T \leq m$, electrons interact with the hot photons in the background only. The effect of fermion background gives an additive contribution at higher temperatures, that is slightly above the electron rest mass (0.511 MeV), i.e; $T > 0.6$ MeV

Thermal contributions during nucleosynthesis in the early universe are mainly expressed by “abc” functions. High temperature and low temperature values of the beta decay rate and the helium abundance parameters, before and after nucleosynthesis, become independent of temperature. Equations (6-9) reduce to the following equations after cancellation of temperature dependence before nucleosynthesis as:

$$\frac{\delta \lambda}{\lambda} = -0.2 \frac{\alpha \pi}{2} \quad (10a)$$

$$\frac{\delta Y}{Y} = 0.04 \frac{\alpha \pi}{2} \quad (10b)$$

$$\frac{\delta \rho}{\rho} = -\frac{\alpha \pi}{48} \quad (10c)$$

$$\frac{\delta H}{H} = -\frac{\alpha \pi}{16} \quad (10d)$$

After the nucleosynthesis is stopped in the early universe, we obtain the following set of equations:

$$\frac{\delta\lambda}{\lambda} = -0.2 \frac{\alpha\pi}{3} \quad (11a)$$

$$\frac{\delta Y}{Y} = 0.04 \frac{\alpha\pi}{3} \quad (11b)$$

$$\frac{\delta\rho}{\rho} = -\frac{\alpha\pi}{32} \quad (11c)$$

$$\frac{\delta H}{H} = -\frac{\alpha\pi}{12} \quad (11d)$$

Actual changes in the values of nucleosynthesis parameters and the contribution to the rate of expansion due to the hot fermion background in the early universe are tabulated in Table (2). Temperature independence of Eqs. (10-11) explain the flatness of all graphs in Figs (5-8) below and above nucleosynthesis temperatures as well. The constant values of these parameters are calculated in Table (2) as:

Quantity	Symbol	Before Nucleosynthesis	After Nucleosynthesis
Electron Self-mass	$\frac{\delta m}{m}$	$\frac{\alpha\pi T^2}{2m^2}$	$\frac{\alpha\pi T^2}{3m^2}$
Beta Decay Rate	$\frac{\delta\lambda}{\lambda}$	-0.002294056	-0.001529371
Helium Abundance	$\frac{\delta Y}{Y}$	0.000458811	0.000305874
Energy Density	$\frac{\delta\rho}{\rho}$	-0.0007166893	-0.000477928
Hubble Parameter	$\frac{\delta H}{H}$	-0.000358446	-0.000238964

Table 2: Values of nucleosynthesis parameters, before and after nucleosynthesis.

It shows that without the estimation of correct thermal effects at a particular temperature, just considering hot fermion background gives a correction factor that is not the same at $T < m$ and $T > m$. can then be calculated that all the nucleosynthesis parameters change before and after nucleosynthesis due to the modification of the self-mass of the electron. However, thermal contribution is very small.

Our results show that the fermionic concentration changes significantly during nucleosynthesis due to beta decay and that is understandable that why that occurred around the mass of electron due to their pivotal role in nucleosynthesis.

References

1. Landsman, Nicolaas P., and Ch G. Van Weert. "Real-and imaginary-time field theory at finite temperature and density." *Physics Reports* 145, no. 3-4 (1987): 141-249.
2. Ahmed, K., and Samina Saleem. "Renormalization and radiative corrections at finite temperature reexamined." *Physical Review D* 35, no. 6 (1987): 1861.
3. Ahmed, K., and Samina Saleem. "Finite-temperature and-density renormalization effects in QED." *Physical Review D* 35, no. 12 (1987): 4020.
4. Haseeb, Mahnaz Q., and Samina S. Masood. "Two loop low temperature corrections to electron self energy." *Chinese Physics C* 35, no. 7 (2011): 608.
5. Masood, Samina S. "Renormalization of QED near decoupling temperature." *Physics Research International* 2014 (2014).
6. Masood, Samina S. "Nucleosynthesis in hot and dense media." *arXiv preprint arXiv:1405.1239* (2014).
7. Saleem, Samina. "Finite-temperature and-density effects on electron self-mass and primordial nucleosynthesis." *Physical Review D* 36, no. 8 (1987): 2602.
8. Masood, Samina S., and Mahnaz Qader. "Finite-temperature and-density corrections to electroweak processes." *Physical Review D* 46, no. 11 (1992): 5110.
9. Schwartz, Matthew D. *Quantum field theory and the standard model*. Cambridge University Press, 2014.
10. Dolan, Louise, and Roman Jackiw. "Symmetry behavior at finite temperature." *Physical Review D* 9, no. 12 (1974): 3320.
11. Kinoshita, Toichiro. "Mass singularities of Feynman amplitudes." *Journal of Mathematical Physics* 3, no. 4 (1962): 650-677.
12. Lee, Tsung-Dao, and Michael Nauenberg. "Degenerate systems and mass singularities." *Physical Review* 133, no. 6B (1964): B1549.
13. Kolb, Edward W., and Michael S. Turner. *The early universe*. CRC press, 2018.
14. Weinberg, Steven. "Gravitation and cosmology: principles and applications of the general theory of relativity." (1972).