

Testing for homogeneous treatment effects in linear and nonparametric instrumental variable models

Jad BEYHUM ^{*}

jad.beyhum@gmail.com

Elia LAPENTA [¶]

elia.lapenta@gmail.com

Jean-Pierre FLORENS [§]

jean-pierre.florens@tse-fr.eu

Ingrid VAN KEILEGOM ^{*}

ingrid.vankeilegom@kuleuven.be

April 17, 2023

Abstract

The hypothesis of homogeneous treatment effects is central to the instrumental variables literature. This assumption signifies that treatment effects are constant across all subjects. It allows to interpret instrumental variable estimates as average treatment effects over the whole population of the study. When this assumption does not hold, the bias of instrumental variable estimators can be larger than that of naive estimators ignoring endogeneity. This paper develops two tests for the assumption of homogeneous treatment effects when the treatment is endogenous and an instrumental variable is available. The tests leverage a covariable that is (jointly with the error terms) independent of a coordinate of the instrument. This covariate does not need to be exogenous. The first test assumes that the potential outcomes are linear in the regressors and is computationally simple. The second test is nonparametric and relies on Tikhonov regularization. The treatment can be either discrete or continuous. We show that the tests have asymptotically correct level and asymptotic power equal to one against a range of alternatives. Simulations demonstrate that the proposed tests attain excellent finite sample performances. The methodology is also applied to the evaluation of returns to schooling and the effect of price on demand in a fish market.

KEYWORDS: Nonparametric Instrumental variables, Hypothesis testing, Homogeneous treatment effects.

^{*}ORSTAT, KU Leuven. Financial support from the European Research Council (2016-2021, Horizon 2020 / ERC grant agreement No. 694409) is gratefully acknowledged.

[§]Toulouse School of Economics, Université Toulouse Capitole. Jean-Pierre Florens acknowledges funding from the French National Research Agency (ANR) under the Investments for the Future program (Investissements d'Avenir, grant ANR-17-EURE-0010).

[¶]CREST, ENSAE Paris, Institut Polytechnique de Paris

1 Introduction

We are interested in the effect of a (possibly continuous) random treatment Z (with support $\mathcal{Z} \subset \mathbb{R}^p$) on a scalar outcome Y . Let $Y(z)$ be the scalar potential outcome of the outcome Y under treatment status $z \in \mathcal{Z}$. The potential outcomes are not observed, but $Y = Y(Z)$ is part of the data. We write

$$Y(z) = \varphi(z) + U(z),$$

where $U(z)$ is an error term such that $\mathbb{E}[U(z)] = 0$. The goal is to identify the structural function φ , which allows to obtain the average treatment effect of changing treatment level from $z \in \mathcal{Z}$ to $z' \in \mathcal{Z}$, i.e. $\mathbb{E}[Y(z') - Y(z)] = \varphi(z') - \varphi(z)$.

We say that Z is endogenous when the structural function φ is not characterized by the distribution of Y conditional on Z . Even in this setting, it is possible to identify φ thanks to an instrumental variable W with support $\mathcal{W} \subset \mathbb{R}^q$. Loosely speaking, an instrumental variable satisfies two conditions. First, it is (in some sense specified in the paper) independent of $\{U(z)\}_z$. Second, it is sufficiently related to Z .

Our goal is to test the following hypothesis

$$H_0 : U(z) = U \text{ for all } z \in \mathcal{Z}.$$

We call this assumption “homogeneous treatment effects” since it stipulates that the treatment effect $Y(z') - Y(z)$ is a deterministic variable. Under H_0 , we have $Y = \varphi(Z) + U$, which corresponds to the standard nonparametric instrumental variable (NPIV) model, see [Newey and Powell \(2003a\)](#); [Horowitz and Lee \(2007\)](#); [Darolles et al. \(2011\)](#); [Chen and Pouzo \(2012\)](#) and the book [Li and Racine \(2007\)](#), among others.

There are two reasons why one would be interested in testing H_0 . First, under the instrumental variable conditions, the aforementioned works in the instrumental variable literature show that the structural function φ is identified if H_0 holds. When H_0 is not satisfied, instrumental variable estimators can be more biased than naive estimators ignoring confounding since the error term $U(Z)$ of the model can be dependent of W (through Z). In this case, the techniques developed in the nonparametric instrumental variable literature outlined above should not be used. Second, if H_0 is true, the treatment effects are constant (i.e. $Y(z') - Y(z)$ is a deterministic variable). This allows to estimate all counterfactuals for each observation

in the dataset (individualized predictions) using an estimator of φ . It also means that the treatment does not distort the ranks of potential outcomes, which has implications in terms of inequalities.

The contributions of the paper are as follows. We propose two tests for H_0 . The tests rely on a scalar covariate X such that $(\{U(z)\}_z, X)$ is independent of one component W_k of the instrument W . Importantly, the covariable X is allowed to be dependent of $\{U(z)\}_z$ (even conditional on W). The first test relies on the assumption that the potential outcomes are linear in the regressors. In a first step, it estimates U as the residual of a two stage least-squares regression. Then, it tests that the estimator of U is uncorrelated with $X(W_k - \mathbb{E}[W_k])$, which is an implication of homogeneous treatment effects under our assumptions. The second test is nonparametric. In this case, the error term U is estimated by the residual of a Tikhonov regression. The second step of the test is then the same as that of the linear test. In both cases, the treatment can be discrete or continuous. We study the asymptotic distribution of the test statistic and the power of the test. We discuss how to choose X : to maximize the likelihood that the conditions on X are satisfied, we argue in the paper that X should be chosen so as to be independent of W . Finally, the empirical performance of the tests is assessed through simulations and illustrated thanks to applications on returns to schooling and price elasticity of demand in a fish market.

Related literature Our paper is related to a large literature on testing for (various implications of) homogeneous treatment effects: see [Koenker and Xiao \(2002\)](#); [Crump et al. \(2008\)](#); [Chernozhukov and Fernández-Val \(2005\)](#); [Ding et al. \(2016\)](#); [Hsu \(2017\)](#); [Goldman and Kaplan \(2018\)](#); [Chung and Olivares \(2021\)](#); [Sant’Anna \(2021\)](#) and [Dai and Stern \(2022\)](#) among others. Apart from [Sant’Anna \(2021\)](#), none of these papers considers the case where there is endogeneity. [Sant’Anna \(2021\)](#) only studies the setting where both the treatment and the instrument are binary and a monotonicity assumption such as in the literature on local average treatment effects holds (see [Angrist et al. \(1996\)](#)). Instead, our paper allows for confounding and continuous treatments.

Another related literature is that on the hypothesis of rank invariance (see [Chernozhukov and Hansen \(2005\)](#)). This assumption is the counterpart of our homogeneous treatment effects condition in nonseparable instrumental variable models (in contrast with the present

paper which studies separable models). [Chernozhukov and Hansen \(2005\)](#) notes that identification of φ can be obtained under a weaker (but less interpretable) form of rank invariance called rank similarity. We could define an assumption analogous to rank similarity in our case of separable models. Our tests would also work for this assumption. We chose to focus solely on homogeneous treatment effects to simplify the exposition. Moreover, [Dong and Shen \(2018\)](#), [Frandsen and Lefgren \(2018\)](#) and [Kim and Park \(2022\)](#) develop tests of the rank invariance assumption in nonseparable instrumental variable models in the case where the treatment and the instrument are binary. The case with continuous treatment is much more challenging because regularization is needed for nonparametric estimation. Note also that [Dong and Shen \(2018\)](#) and [Kim and Park \(2022\)](#) rely on a monotonicity assumption from local average treatment effects literature (see [Angrist et al. \(1996\)](#)), while our approach does not require it. The test developed in [Frandsen and Lefgren \(2018\)](#) bears similarities with our tests since they are based on equivalent single restrictions. However, our models and the first steps of our tests are different. We provide a more in depth analysis of the underlying assumptions of the method and study it in a different context with models for average treatment effects and allow for continuous treatments.

Outline Section 2 presents the linear test along with theory, simulations, and an application to returns to schooling. Then, in Section 3, we introduce the nonparametric test, discuss its asymptotic properties, evaluate its finite sample performance through numerical experiments, and illustrate the test with an application to demand estimation.

2 Linear test

2.1 Model

In this section, we develop theory for a test assuming that the mapping φ is linear and the instrument W is uncorrelated with $\{U(z)\}_z$, i.e. there exists $\beta \in \mathbb{R}^p$ such that

$$Y(z) = z^\top \beta + U(z), \quad \mathbb{E}[WU(z)] = 0,$$

for all $z \in \mathcal{Z}$. We also impose the usual relevance assumption for instrumental variables in the linear model, i.e. $\mathbb{E}[WZ^\top]$ has rank equal to p . The aim is to recover β , since the

average treatment effect of changing treatment level from $z \in \mathcal{Z}$ to $z' \in \mathcal{Z}$ can be expressed as $\mathbb{E}[Y(z') - Y(z)] = (z' - z)^\top \beta$.

Let the symbol \perp stand for statistical independence. To test for H_0 , we leverage a scalar covariate X satisfying the following condition:

Assumption 2.1 *There exists $k \in \{1, \dots, q\}$ such that, for all $z \in \mathcal{Z}$, $(\{U(z)\}_z, X) \perp W_k$.*

This assumption allows X and $\{U(z)\}_z$ to be dependent (even conditional on W). Hence X does not need to be exogenous or satisfy an exclusion restriction. In particular, X is not a second instrument.

We argue that Assumption 2.1 is very likely to hold for some X if W_k is a good instrument. Indeed, if W_k is independent of the unobserved heterogeneity of the model, there is every reason to believe that it is also independent of some observable variables. Note that Assumption 2.1 is equivalent to the standard instrumental variable condition $\{U(z)\}_z \perp W_k$ and the additional condition $X \perp W_k | \{U(z)\}_z$. Hence, the condition we add to the standard instrumental variable literature is really $X \perp W_k | \{U(z)\}_z$.

The tests of H_0 that we propose depend crucially on Assumption 2.1. Contrarily to H_0 , hypothesis 2.1 can be heuristically justified. The econometrician should try to select a covariable X which is likely to be (jointly with $\{U(z)\}_z$) independent of W_k . In applications, we recommend to pick a variable X independent of W_k since this should make Assumption 2.1 more likely to hold as $X \perp W_k$ is a necessary condition for Assumption 2.1. The independence of X and W_k can be assessed by a statistical test of independence such as χ^2 or Kolmogorov-Smirnov independence tests. Our empirical applications illustrate possible choices of X .

Note that Assumption 2.1 could hold for several components of W and a multidimensional X . The extension of the test would be straightforward in this case. We focus on the scalar case to simplify the exposition and because the validity of this assumption for a single component of W is less restrictive.

Let us now define the population analog of the two-stage least squares estimator (henceforth, TSLS) in this context:

$$\beta^{TSLS} = [\Gamma^\top \mathbb{E}[WW^\top] \Gamma]^{-1} \Gamma^\top \mathbb{E}[WY],$$

where $\Gamma = \mathbb{E}[WW^\top]^{-1}\mathbb{E}[WZ^\top]$. In this case, under H_0 , β can be estimated by TSLS, i.e. $\beta^{TSLS} = \beta$. However, when H_0 does not hold, then the bias of the TSLS estimator can be larger than that of the ordinary least squares (henceforth, OLS) estimator as illustrated by the following example.

Example 1. We study the case of a randomized experiment with two-sided noncompliance and monotonicity (see Angrist et al. (1996)). Let $W = (1, W_2)^\top$, where W_2 is a Bernoulli random variable with $\mathbb{P}(W_2 = 1) = 1/2$ and $Z = (1, Z_2)^\top$, with $Z_2 = W_2 1\{\epsilon \geq 1/2\} + (1 - W_2) 1\{1/2 \leq \epsilon < 3/4\}$, where $\epsilon \perp\!\!\!\perp W$ follows a uniform distribution on the interval $[0, 1]$. Note that $\mathbb{P}(Z_2 = 1|W_2 = 0) = 1/4$ and $\mathbb{P}(Z_2 = 1|W_2 = 1) = 1/2$. For some $\alpha > 0$, we also impose $\beta_1 = 0$ and $\beta_2 = \alpha/4$ (i.e. $Y(1, z_2) = (\alpha/4)z_2 + U(Z)$ and $U(1, z_2) = z_2(1\{\epsilon \geq 3/4\} - (1/4))\alpha$ for $z_2 = 0, 1$). This definition ensures that $E[U(z)] = 0$, for $z = 0, 1$. This model satisfies all the usual instrumental variable assumptions except H_0 . The average treatment effect over the whole population is

$$\Delta = \mathbb{E}[Y(1, 1) - Y(1, 0)] = \frac{\alpha}{4}.$$

We show in the supplementary material that the population analog of the OLS estimator of β_2 is given by

$$\beta_2^{OLS} = \mathbb{E}[Y|Z_2 = 1] - \mathbb{E}[Y|Z_2 = 0] = \frac{\alpha}{3}.$$

Instead, in the context of the present example, it is known (Angrist et al. (1996)) that β_2^{TSLS} is equal to the average treatment effects on the population of compliers. The compliers are the subjects who change treatment status with the instrument, or, equivalently, with $\epsilon \geq 3/4$. As a result, it holds that $\beta_2^{TSLS} = \alpha$. Hence, the bias of the TSLS estimator is larger than that of the OLS estimator for Δ and can even go to infinity as $\alpha \rightarrow \infty$.

2.2 Testable implication

Let $U^{TSLS} = Y - Z^\top \beta^{TSLS}$. Under the null hypothesis H_0 , $U(z)$ will not depend on the treatment level z and $U^{TSLS} = U$, which yields $(U^{TSLS}, X) \perp\!\!\!\perp W_k$ (by Assumption 2.1). This last independence condition implies that

$$\mathbb{E}[U^{TSLS} X (W_k - \mathbb{E}[W_k])] = 0. \quad (2.1)$$

This is the testable implication of H_0 that we use to construct the test. In particular, our test estimates the moment (2.1) and checks if the latter is close to 0. Remark that the role of the covariable X appears clearly in (2.1). Indeed, when W contains an intercept, the equality

$$\mathbb{E}[U^{TSLS}(W - \mathbb{E}[W])] = 0 \quad (2.2)$$

always holds by definition of U^{TSLS} , regardless of the validity of H_0 . Hence, a test based on (2.2) would have no power. The role of X is therefore to give power to our test.

A few remarks are in order. First, there are multiple reasons why we choose to focus on testing the moment condition (2.1) instead of any of the other implications of H_0 . This is because, under our assumptions, (2.1) is equivalent to saying that the coefficients of the linear projection of $W_k - \mathbb{E}[W_k]$ onto a constant, U^{TSLS} , X , and the product XU^{TSLS} are all null. Indeed, the latter coefficients are “proportional” to $\mathbb{E}[(W_k - \mathbb{E}[W_k])(1, U^{TSLS}, X, U^{TSLS}X)]$, with $\mathbb{E}[(W_k - \mathbb{E}[W_k])(1, U^{TSLS})] = 0$ by construction and $\mathbb{E}[(W_k - \mathbb{E}[W_k])X] = 0$ by Assumption 2.1, so that the coefficients are null if and only if (2.1) holds (under Assumption 2.1). We therefore focus on (2.1) because of this nice interpretation.

Second, we acknowledge that instead of focusing on (2.1) we could directly test the independence condition $(U^{TSLS}, X) \perp\!\!\!\perp W_k$. However, we prefer to focus on (2.1) for three main reasons. First, as we show in the next section, our test based on (2.1) is very simple to implement, and hence it will be attractive to practitioners. Differently, building a test for the independence condition $(U^{TSLS}, X) \perp\!\!\!\perp W_k$ would require performing a multi-step estimation, where in a first step we estimate the error U^{TSLS} , in a second step we should estimate the cumulative distribution functions of W_k and (U^{TSLS}, X) , and in a third step we should compare these cumulative distribution functions on the basis of a certain distance. A test of this type would be less attractive to implement in practice. Second, as we show in Section 2.4, a test based on the moment condition (2.1) will have power against a wide range of alternatives. We also remark that focusing on the moment condition in (2.1) is not necessarily a limitation: while we do not have consistency against all fixed alternatives, we focus the power towards the specific direction of the moment (2.1). Instead, a test comparing the nonparametric cumulative distribution functions of (U^{TSLS}, X) and W would spread the power along all directions, and may have lower power in the direction of our moment (2.1) and, hence, in the direction of the linear projection of $W_k - \mathbb{E}[W_k]$ onto a constant, U^{TSLS} ,

X and the product XU^{TSLS} , as discussed above.

Third, remark that when X is binary (support equal to $\{0, 1\}$), (2.1) becomes $\mathbb{E}[U^{TSLS}(W_k - \mathbb{E}[W_k])|X = 1] = 0$. Hence, in this case our test checks if U^{TSLS} and W_k are uncorrelated (that is W_k is exogenous) in the subpopulations for which $X = 0$ and $X = 1$.

2.3 Sample test

Let us formally outline the test. Consider an i.i.d. sample $\{Y_i, Z_i, W_i, X_i\}_{i=1}^n$ generated from the model of Section 2.1. Let $\hat{\beta}^{TSLS}$ be the TSLS estimator in this sample, given by

$$\hat{\beta}^{TSLS} = \left[\hat{\Gamma}^\top \left(\frac{1}{n} \sum_{i=1}^n W_i W_i^\top \right) \hat{\Gamma} \right]^{-1} \hat{\Gamma}^\top \left(\frac{1}{n} \sum_{i=1}^n W_i Y_i \right),$$

where $\hat{\Gamma} = \left(\frac{1}{n} \sum_{i=1}^n W_i W_i^\top \right)^{-1} \frac{1}{n} \sum_{i=1}^n W_i Z_i^\top$. An estimator for U_i is therefore

$$\hat{U}_i = Y_i - Z_i^\top \hat{\beta}^{TSLS}.$$

Let $\bar{W}_k = n^{-1} \sum_{i=1}^n W_{ki}$. The test statistic is then

$$T_n = \frac{1}{\sqrt{n}} \sum_{i=1}^n \hat{U}_i X_i (W_{ki} - \bar{W}_k).$$

We use bootstrap to obtain the p-value of the test, since the variance of the statistic has a tedious expression and variance estimators based on analytical formulas tend to perform poorly in the presence of heteroscedasticity (see e.g. [MacKinnon and White \(1985\)](#)). The procedure to compute the p-value is as follows.

1. Draw n observations with replacement from the sample $\{Y_i, Z_i, W_i, X_i\}_{i=1}^n$
2. Compute the bootstrapped statistic $T_{n,b}^*$ on the bootstrapped sample
3. Repeat steps 1-2 B times (with B large) so as to get the collection of bootstrapped statistics $\{T_{n,b}^* : b = 1 \dots, B\}$
4. Compute the *symmetric* p-value as $\frac{1}{B} \sum_{b=1}^B 1 \{ |T_{n,b}^* - T_n| > |T_n| \}$.

If the p-value so obtained is smaller than α (the nominal size of the test), the null hypothesis is rejected at the α nominal level. Notice that in the above procedure we are computing a *symmetric* p-value. An alternative procedure is to compute an *equal-tailed* p-value, but in our simulations the symmetric one has a satisfying performance.

2.4 Power analysis

Informally, when H_0 does not hold, U^{TSLS} will, in general, not be equal to some $U(z)$ for a fixed $z \in \mathcal{Z}$. Therefore, in this case, there is no reason that $(U^{TSLS}, X) \perp\!\!\!\perp W_k$, and hence, that (2.1) holds. This gives power to our test. Going further, we argue in this section that it is overwhelmingly likely that (2.1) is wrong when H_0 does not hold. The test will have asymptotic power equal to 1 under alternative hypotheses for which

$$\mathbb{E}[U^{TSLS}X(W_k - \mathbb{E}[W_k])] \neq 0. \quad (2.3)$$

Since $U^{TSLS} = Y - Z^\top \beta^{TSLS} = U(Z) + Z^\top(\beta - \beta^{TSLS})$, Equation (2.3) is equivalent to

$$\mathbb{E}[U(Z)X(W_k - \mathbb{E}[W_k])] + \mathbb{E}[X(W_k - \mathbb{E}[W_k])Z^\top](\beta - \beta^{TSLS}) \neq 0. \quad (2.4)$$

We see that there are two reasons why the test would have power:

- (1) the variable W_k is correlated with $U(Z)X$;
- (2) it holds that $\beta_\ell \neq \beta_\ell^{TSLS}$ and $\mathbb{E}[X(W_k - \mathbb{E}[W_k])Z^\top]_\ell \neq 0$ for at least one ℓ in $\{1, \dots, p\}$.

We argue that (1) and (2) are likely to hold in applications. Indeed, statement (1) is probable since W and Z are correlated and $U(Z)$ depends on Z . The fact that $\mathbb{E}[X(W_k - \mathbb{E}[W_k])Z^\top] \neq 0$ is likely under the assumption that $\mathbb{E}[ZW^\top]$ has rank p . When H_0 does not hold, β and β^{TSLS} should be different, they would be equal only under very specific values of $U(Z)$.

Notice that it would be possible for (1) and (2) to hold while the test does not have power. This happens when the two terms in (2.4) compensate each other. This case, however, requires very specific data generating processes.

Overall, although there are cases where the test does not have power, the above discussion suggests that the test has power against a wide range of alternative hypotheses. The following example illustrates this claim.

Example 2. Let $(W, E, X)^\top$ be a 3×1 Gaussian vector with mean zero and variance equal to the identity matrix. Let also $Z = W + WE + \rho WX$, $U(Z) = Z(E + \rho X)$, where $\rho \in \mathbb{R}$,

and $\beta = 0$ so that $Y = U(Z) = Z(E + \rho X)$. In this case, we have

$$\begin{aligned}\mathbb{E}[YW] &= \mathbb{E}[WZ(E + \rho X)] \\ &= \mathbb{E}[W^2E + W^2E^2 + \rho W^2XE + \rho(W^2X + W^2EX + \rho W^2X^2)] = 1 + \rho^2.\end{aligned}$$

Moreover, $\mathbb{E}[ZW] = \mathbb{E}[W^2 + W^2E + \rho W^2X] = 1$. In the present context, it is well-known that $\beta^{TSLS} = \mathbb{E}[YW]/\mathbb{E}[ZW]$, which yields $\beta^{TSLS} = 1 + \rho^2$. We have $U^{TSLS} = Y - Z\beta^{TSLS} = Z\{(E + \rho X) - (1 + \rho^2)\}$. Hence, we get

$$\begin{aligned}\mathbb{E}[U^{TSLS}WX] &= \mathbb{E}[(W + WE + \rho WX)\{(E + \rho X) - (1 + \rho^2)\}WX] \\ &= \rho\mathbb{E}[(WX)^2] - \rho(1 + \rho^2)\mathbb{E}[(WX)^2] = -\rho^3.\end{aligned}$$

As a result, the test has no power only when X and U are independent, i.e. $\rho = 0$, which is a degenerate case.

In this previous parametric example, we see that the alternatives under which the test has no power have measure equal to 0 for the uniform measure. The example also shows the role of X in providing power to the test, since the latter does not have power only when X and $(Z, \{U(z)\}_z)$ are independent.

2.5 Asymptotic theory

In this section, we state the asymptotic properties of the test statistic. We make the following assumption, which ensures the convergence of the TSLS estimator.

Assumption 2.2 $\mathbb{E}[U(Z)^2 + X^2 + \|W\|_2^2] < \infty$, $\mathbb{E}[ZW^\top]$ exists and has rank p and $\mathbb{E}[WW^\top]$ exists and has full rank.

We have the following theorem.

Theorem 2.1 *Let Assumptions 2.1 and 2.2 hold. Then, T_n converges in distribution to a zero mean Gaussian distribution under H_0 , while $|T_n|/\sqrt{n} \xrightarrow{\mathbb{P}} C \neq 0$ when (2.3) holds.*

The influence function representation of the test statistic is given in Lemma 1.1 in the supplementary material. The asymptotic variance of the test statistic can be derived from this result. Note that the fact that $|T_n|/\sqrt{n} \rightarrow C \neq 0$ when (2.3) holds implies that the power of the test goes to 1 as n goes to ∞ under alternative hypotheses satisfying (2.3).

2.6 Simulations

We study the following data generating process. Let (W_2, E, X, Z_3) follow a standard 4-dimensional Gaussian distribution. We define $Z_2 = W_2 + E + X$, $Z = (1, Z_2, Z_3)^\top$, $W = (1, W_2, Z_3)^\top$. The variable Z_2 suffers from confounding and is instrumented by W_2 , while the variable Z_3 is exogenous. We let $U(z) = (1 + \rho z_2)(E + X)$, for all $z \in \{1\} \times \mathbb{R}^2$, where $\rho \in \mathbb{R}$. The null hypothesis H_0 holds when $\rho = 0$. The outcome is $Y = U(Z)$ so that the causal regression function of interest is $\varphi = 0$. We set $k = 2$, i.e. we use the second component of W to compute the test statistic $T_n = n^{-1/2} \sum_{i=1}^n \hat{U}_i X_i (W_{2i} - \bar{W}_2)$.

We generate data with sample sizes $n \in \{100, 200, 500, 1000\}$. We investigate the empirical size of the test when $\rho = 0$ (Table 1). The results are averages over 10000 replications using $B = 1000$ bootstrap resamples. We also study the empirical power of the test when $\rho = -1, -0.9, \dots, 1$, with a thousand replications and bootstrap resamples (See Figure 1). The empirical size of the test is almost nominal even for low sample sizes. The power of the test increases as the deviation from the null ($|\rho|$), or the sample size, become larger.

n	100	250	500	1000
Empirical size at 5%	0.0397	0.0512	0.0506	0.0506
Empirical size at 10%	0.1038	0.1050	0.1038	0.1014

Table 1: Empirical size of the test of theoretical size 5% and 10% for various sample sizes.

2.7 Application on returns to schooling

In this section, we test for homogeneous treatment effects in a famous application on returns to schooling from Card (1995). The database is available in the R package *ivreg* and is an extract from the 1976 National Longitudinal Survey (NLS) of young men. Here, $Y = \log(wage)$, $Z = (\mathbf{1}, education, experience, experience^2, smsa, south, ethnicity)^\top$, where $\mathbf{1}$ denotes a constant, *education* and work *experience* are measured in years, and *smsa*, *south*, *ethnicity* are controls whose definition can be found in the *ivreg* package. *Education* and *experience* are endogenous since they depend on individual's ability which is unobserved. The instrument is $W = (\mathbf{1}, nearcollege, age, age^2, smsa, south, ethnicity)^\top$, where *nearcollege* is an indicator whose value is equal to 1 when the individual grew up near an accredited four-year college. The validity of the instrument is documented in Card (1995).




Figure 1: Empirical power of the test of theoretical size 5% as a function of ρ .

This corresponds to one of the specifications in the original paper of [Card \(1995\)](#). The variable X is an indicator for being married. It is unlikely that being married is (jointly with the unobserved heterogeneity of the model) dependent from growing up near a four-year college. In fact, a χ^2 test does not reject the hypothesis of independence of $W_2 = \text{nearcollege}$ and $X = \text{married}$ (p-value equal to 0.55). Hence, we can reasonably assume that $(\{U(z)\}_z, X) \perp\!\!\!\perp W_2$ for all $z \in \mathcal{Z}$, which corresponds to Assumption 2.1. Using 10,000 bootstrap resamplings, the p-value of the test is equal to 0.0965, so that the null hypothesis of homogeneous treatment effects can be (borderline) rejected at the 10% level. This result cautions against interpreting the estimates of [Card \(1995\)](#) as causal effects.

3 Nonparametric test

In this section we extend the linear framework to consider a nonparametric treatment function. We assume that Z and W are unidimensional to simplify the exposition (it is rare to apply nonparametric procedures with multidimensional variables). We continue supposing

that Assumption 2.1 holds and we keep the notation previously introduced. We write

$$Y(z) = \varphi(z) + U(z), \mathbb{E}[U(z)|W] = 0, \text{ for all } z \in \mathcal{Z},$$

where the conditional mean independence $\mathbb{E}[U(z)|W] = 0$ is a direct consequence of Assumption 2.1. The functional form of φ is unknown. Let $L^2(Z)$ be the set of functions that are square integrable with respect to the distribution of Z . We maintain the following assumption which is also standard in the NPIV literature.

Assumption 3.3 (a) For all $\phi \in L^2(Z)$, $\mathbb{E}[\phi(Z)|W] = 0 \Rightarrow \phi \equiv 0$. (b) There exists at least one $\tilde{\varphi} \in L^2(Z)$ satisfying $\mathbb{E}[Y|W] = \mathbb{E}[\tilde{\varphi}(Z)|W]$.

The first part of the assumption is the completeness condition standard in the NPIV literature. It means that W is sufficiently associated with Z . Under the null hypothesis H_0 , such a condition is necessary and sufficient for identifying φ (see, among others, Carrasco et al. (2007), Darolles et al. (2011), Newey and Powell (2003b), D'Haultfoeuille (2011)). The second part states that the NPIV model in the above equation ($\mathbb{E}[Y|W] = \mathbb{E}[\tilde{\varphi}(Z)|W]$) is well specified. We will discuss this assumption in Section 3.2.

Recall that the hypothesis that we want to test is

$$H_0 : U(z) = U \text{ for all } z \in \mathcal{Z}.$$

Under the null hypothesis of homogeneous treatment effects $\mathbb{E}[U(Z)|W] = \mathbb{E}[U(z)|W] = \mathbb{E}[U(z)] = 0$, so the treatment function φ must satisfy the following (functional) equation in $\phi \in L^2(Z)$

$$\mathbb{E}[Y|W] = \mathbb{E}[\phi(Z)|W]. \tag{3.1}$$

Under the completeness condition (Assumption 3.3 (a)) the solution to the above equation is *unique* and can be consistently estimated (we introduce an estimator below). Since under the null hypothesis of homogeneous treatment effects the treatment function φ will satisfy the above equation, φ will be identified as its unique solution. Thus, under the null hypothesis of homogeneous treatment effects we can consistently estimate the nonparametric treatment function φ . Differently, when H_0 does not hold, the solution to Equation (3.1) will be different, in general, from the treatment function of interest φ . This solution, denoted with

$\tilde{\varphi}$, is introduced in Assumption 3.3 (b). So, if H_0 is not true, although we will be able to estimate consistently the solution to Equation (3.1), we will not in general obtain a consistent estimator of φ . It is therefore crucial to test for the homogeneous treatment effects hypothesis.

Let $\tilde{U}(Z) := Y(Z) - \tilde{\varphi}(Z)$ and $\tilde{\varphi}$ denote the unique solution to Equation (3.1) (which exists by Assumption 3.3 (ii)). Under the null of homogeneous treatment effects $\tilde{U}(Z) = U$ and $(X, \tilde{U}(Z)) \perp\!\!\!\perp W$. Thus, by arguing similarly as in the previous section, we test

$$\mathbb{E}[\tilde{U}(Z)X(W - \mathbb{E}[W])] = 0 \quad (3.2)$$

Remark that, in the NPIV model, instead of conducting a test based on the moment condition in (3.2), we could build a test to directly check that $(\tilde{U}(Z), X) \perp\!\!\!\perp W$. However, we choose to focus on the single moment condition in (3.2) for several reasons. First, our test based on a single moment restriction has power under a wide range of alternatives. Indeed, a power analysis similar to that in Section 2.4 could be carried out in the present nonparametric case. Some examples of alternative hypotheses under which the test has power are given in Section S2.2 of the supplementary material. Second, as argued in Section 2.4, our test avoids to spread power across multiple directions. Third, focusing on a single restriction allows to propose a test which is simple to implement. In contrast, building a test for the condition $(\tilde{U}(Z), X) \perp\!\!\!\perp W$ would be theoretically intricate. In fact, it would require to estimate the error $\tilde{U}(Z)$ in a first step and then to estimate the cumulative distribution functions of W and $(\tilde{U}(Z), W)$ in a second step. It is likely that in this case we would need an $n^{-1/4}$ convergence rate for the first-step NPIV estimator of $\tilde{\varphi}$, as it is required in multi-step estimations with a nonlinear second step, see Chen et al. (2003). This would require a slow convergence rate of the regularization parameter to allow the variance of the first-step NPIV estimator to converge fast enough to zero, see Assumption 3.6 ahead. At the same time, we would need the regularization bias of $\tilde{\varphi}$ to disappear fast enough, and this would require that the regularization parameter converges quickly to zero, see Assumption 3.6 ahead. Therefore, there would be some tensions between these two conditions and the construction of the test would become intricate. In particular, the rates of convergences obtained with the Tikhonov regularization scheme would not allow to obtain a consistent nonparametric test. A more complicated first-step estimator would be required.

Let $\hat{\varphi}$ be an estimator for φ (below we define our estimator), and let us define the estimated

residuals $\widehat{U}_i = Y_i - \widehat{\varphi}(Z_i)$, for $i = 1, \dots, n$. Our statistic is the empirical counterpart of the above moment

$$S_n = \frac{1}{\sqrt{n}} \sum_{i=1}^n \widehat{U}_i (W_i - \overline{W}) X_i,$$

where $\overline{W} = n^{-1} \sum_{i=1}^n W_i$ is the empirical mean of W . In the following section, we explain how to compute the above statistic and implement the test based on it. In Section 3.2, we present the assumptions and the results for the validity of the test. Next, in Section 3.3, we provide evidence about the finite sample performance of the test based on S_n . Finally, we illustrate the procedure by an application to demand estimation in Section 3.4.

3.1 Construction of the statistic and implementation of the test

In this section, we outline the computation of the statistic S_n and implement the test. To this end, we need (a) to compute $\widehat{\varphi}(Z_i)$ for $i = 1, \dots, n$, and (b) to obtain the p-value necessary for testing.

Computation of $\widehat{\varphi}$. Let π and τ be positive functions on \mathbb{R} . We denote with $L_\pi^2(\mathbb{R})$ the space of square integrable functions with respect to π , and let $L_\tau^2(\mathbb{R})$ be similarly defined. The functions π and τ are introduced for the aim of generality and for technical reasons. In simulations they are set to Gaussian density functions. Let us briefly explain their roles. First, from a technical point of view it would be ideal to work with the spaces $L^2(Z)$ and $L^2(W)$, i.e. the spaces of square integrable functions with respect to the distribution of Z and W . However, since we do not know such distributions, it is convenient to replace $L^2(Z)$ and $L^2(W)$ with $L_\tau^2(\mathbb{R})$ and $L_\pi^2(\mathbb{R})$ and work with the latter spaces. Second, by working with $L_\tau^2(\mathbb{R})$ and $L_\pi^2(\mathbb{R})$ we can obtain that the estimator of the dual of A (defined below) is actually the dual of the estimator of A . This is an important property for establishing the asymptotic normality of our statistic S_n under H_0 .

Let f_W stand for the density of W with respect to the Lebesgue measure, and assume that $\tau(w) > 0$ whenever $f_W(w) > 0$. Then, Equation (3.1) is equivalent to

$$\mathbb{E}[Y|W] \frac{f_W(W)}{\tau(W)} = \mathbb{E}[\widehat{\varphi}(Z)|W] \frac{f_W(W)}{\tau(W)}. \quad (3.3)$$

To build an estimator for $\tilde{\varphi}$ we start from the above integral equation. Let us denote by f_{ZW} the joint density of (Z, W) with respect to the Lebesgue measure and introduce $L^2_{\pi \otimes \tau}(\mathbb{R}^2)$, which is the set of functions from \mathbb{R}^2 to \mathbb{R} square integrable with respect to the product measure $\pi \otimes \tau$ (the measure generated by the product density $\pi \cdot \tau$). By assuming that $f_{ZW}/(\pi \tau) \in L^2_{\pi \otimes \tau}(\mathbb{R}^2)$, we define $A : L^2_{\pi}(\mathbb{R}) \mapsto L^2_{\tau}(\mathbb{R})$ as the operator

$$(A\phi)(\cdot) = \int \phi(z) f_{ZW}(z, \cdot) dz \frac{1}{\tau(\cdot)}.$$

Its Hilbert adjoint $A^* : L^2_{\tau}(\mathbb{R}) \mapsto L^2_{\pi}(\mathbb{R})$ is given by

$$(A^*\psi)(\cdot) = \int \psi(w) f_{ZW}(\cdot, w) dw \frac{1}{\pi(\cdot)}.$$

We also define the “left hand side” of the integral equation (3.3) as

$$r(\cdot) = \mathbb{E}[Y|W = \cdot] \frac{f_W(\cdot)}{\tau(\cdot)}.$$

We estimate A and A^* by replacing f_{ZW} with its kernel estimator

$$\hat{f}_{ZW}(z, w) = \frac{1}{n h_Z h_W} \sum_{i=1}^n K_Z \left(\frac{Z_i - z}{h_Z} \right) K_W \left(\frac{W_i - w}{h_W} \right),$$

where K_Z and K_W are two kernel functions while (h_Z, h_W) denote two bandwidths converging to zero as the sample size increases. The mapping r is instead estimated by

$$\hat{r}(\cdot) := \frac{1}{n h_W} \sum_{i=1}^n Y_i K_W \left(\frac{W_i - \cdot}{h_W} \right) \frac{1}{\tau(\cdot)}.$$

To estimate $\tilde{\varphi}$, i.e. the solution to Equation (3.3), we use a Tikhonov regularization scheme

$$\hat{\varphi} = \left(\lambda I + \hat{A}^* \hat{A} \right)^{-1} \hat{A}^* \hat{r},$$

where I stands for the identity operator, while $\lambda > 0$ denotes the Tikhonov regularization parameter. See, e.g., Carrasco et al. (2007) and Darolles et al. (2011).

To compute $\hat{\varphi}$ we need to select two bandwidths (h_Z, h_W) and a regularization parameter λ . We will explain below how to do it. For the moment, let us outline the computation of $\hat{\varphi}$ in the previous display for given (h_Z, h_W, λ) . Although $\hat{\varphi}$ seems to have an abstract expression, its computation is straightforward as it reduces to matrix products. To see this, we first

approximate \hat{A} and \hat{A}^* by using a common bias computation similarly as in [Centorrino et al. \(2017\)](#) :

$$\begin{aligned}
(\hat{A}\phi)(w) &= \frac{1}{nh_Z h_W} \sum_{i=1}^n K\left(\frac{W_i - w}{h_W}\right) \frac{1}{\tau(w)} \underbrace{\int K\left(\frac{Z_i - z}{h_Z}\right) \phi(z) dz}_{\approx h_Z \phi(Z_i)} \\
&\approx \frac{1}{nh_W} \sum_{i=1}^n K\left(\frac{W_i - w}{h_W}\right) \frac{1}{\tau(w)} \phi(Z_i) \\
(\hat{A}^*\psi)(z) &= \frac{1}{nh_Z h_W} \sum_{j=1}^n K\left(\frac{Z_j - z}{h_Z}\right) \frac{1}{\pi(z)} \underbrace{\int K\left(\frac{W_j - w}{h_W}\right) \psi(w) dw}_{\approx h_W \psi(W_j)} \\
&\approx \frac{1}{nh_Z} \sum_{j=1}^n K\left(\frac{Z_j - z}{h_Z}\right) \frac{1}{\pi(z)} \psi(W_j).
\end{aligned}$$

The above approximations hold for $h_Z, h_W \rightarrow 0$. Let M_Z be the $n \times n$ matrix having on the i th row and j th column the element $K((Z_i - Z_j)/h_Z)/[\pi(Z_i)nh_Z]$ and let M_W be the $n \times n$ matrix having on the i th row and j th column the element $K((W_i - W_j)/h_W)/[\tau(W_i)nh_W]$. Let $\vec{\hat{\varphi}} := (\hat{\varphi}(Z_1), \dots, \hat{\varphi}(Z_n))^T$. For a generic function ψ of W let $\vec{\hat{A}^*\psi} := ((\hat{A}^*\psi)(W_1), \dots, (\hat{A}^*\psi)(W_n))^T$. Let $\vec{\hat{r}}$ and $\vec{\hat{A}\phi}$ (for a generic function ϕ) be similarly defined. By the expression of $\hat{\varphi}$ and the above approximations we get

$$\vec{\hat{r}} = \lambda \vec{\hat{\varphi}} + \hat{A}^*(\hat{A}\hat{\varphi}) \approx \lambda \vec{\hat{\varphi}} + M_Z \vec{\hat{A}\hat{\varphi}} \approx \lambda \vec{\hat{\varphi}} + M_Z M_W \vec{\hat{\varphi}},$$

so that

$$\vec{\hat{\varphi}} \approx (\lambda I + M_Z M_W)^{-1} M_Z \vec{\hat{r}} \quad (3.4)$$

up to an approximation error. Thus, the computation of $\hat{\varphi}$ at the data points reduces to a simple matrix computation and the residuals can be easily computed as $\hat{U}_i = Y_i - \hat{\varphi}(Z_i)$ with $i = 1, \dots, n$.

Let us turn now to the selection of the smoothing parameters (h_Z, h_W) and the regularization parameter λ . The bandwidths h_Z and h_W are selected by the Silverman Rule of Thumb, so that $h_Z = \hat{\sigma}_Z n^{-1/5}$ and $h_W = \hat{\sigma}_W n^{-1/5}$, where $\hat{\sigma}_Z$ and $\hat{\sigma}_W$ denote the sample standard deviations of Z and W . Finally, to select the Tikhonov regularization parameter, we use the Cross-Validation method developed in [Centorrino et al. \(2017\)](#). Thus,

$$\hat{\lambda} := \arg \min_{\lambda} \sum_{i=1}^n \left[(\hat{A}\hat{\varphi}^\lambda)_{-i}(W_i) - \hat{r}(W_i) \right]^2, \quad (3.5)$$

where $(\widehat{A}\widehat{\varphi}^\lambda)_{-i}(W_i) := \left[\widehat{A}_{-i}(\lambda I + \widehat{A}^* \widehat{A})^{-1} \widehat{A}_{-i}^* \widehat{r} \right] (W_i)$. Here \widehat{A}_{-i} and \widehat{A}_{-i}^* denote the “leave-one-out” versions of \widehat{A} and \widehat{A}^* that use the entire sample except the i th observation. By the approximations of \widehat{A} and \widehat{A}^* previously outlined, the vector $\{(\widehat{A}\widehat{\varphi}^\lambda)_{-i}(W_i) : i = 1, \dots, n\}$ is computed as

$$\begin{aligned} \left[(\widehat{A}\widehat{\varphi}^\lambda)_{-i}(W_i) \right]_{i=1, \dots, n} &= \left[\widehat{A}_{-i}(\lambda I + \widehat{A}^* \widehat{A})^{-1} \widehat{A}_{-i}^* \widehat{r} \right]_{i=1, \dots, n} \\ &\approx (M_W - \text{diag}(M_W)) (\lambda I + M_Z M_W)^{-1} (M_Z - \text{diag}(M_Z)) \vec{\widehat{r}}, \end{aligned}$$

where $\text{diag}(M_W)$ denotes the diagonal matrix having the same main diagonal as M_W , and $\text{diag}(M_Z)$ is similarly defined. The objective function in (3.5) has a U-shaped form as a function of λ and can be minimized in a simple way by numerical methods, see Centorrino et al. (2017).

For the sake of clarity, we sum up the steps needed for computing S_n as follows:

1. Select $\widehat{\lambda}$ according to the Cross-validation method in Equation (3.5)
2. Use $\widehat{\lambda}$ to compute $\{\widehat{\varphi}(Z_i) : i = 1, \dots, n\}$ as in (3.4) and compute the residuals as $\{\widehat{U}_i = Y_i - \widehat{\varphi}(Z_i) : i = 1 \dots, n\}$
3. Compute the statistic as $S_n = (1/\sqrt{n}) \sum_{i=1}^n \widehat{U}_i(W_i - \overline{W})X_i$.

Implementation of the test. To implement the test, we are just left with the computation of the p-value. As we show in the next section, under the null hypothesis the statistic S_n is asymptotically normal, but the asymptotic covariance has an intricate expression. So, to obtain the p-value necessary for testing we rely on the pairwise bootstrap. The validity of the pairwise bootstrap is confirmed by our simulations and is not surprising given that we show asymptotic normality. In fact, under different conditions, Chen et al. (2003) show the validity of the bootstrap for a *real-valued* estimator based on a first step *infinite-dimensional* estimator, as is $\widehat{\varphi}$ in this paper. The procedure goes as follows:

1. Draw n observations with replacement from the sample $\{Y_i, Z_i, W_i, X_i\}_{i=1}^n$
2. Compute the bootstrapped statistic $S_{n,b}^*$ on the bootstrapped sample using the same bandwidths (h_Z, h_W) and regularization parameter λ as in the original sample

3. Repeat steps 1-2 B times (with B large) to obtain the collection of bootstrapped statistics $\{S_{n,b}^* : b = 1 \dots, B\}$
4. Compute the *symmetric* p-value as $\frac{1}{B} \sum_{b=1}^B 1 \{|S_{n,b}^* - S_n| > |S_n|\}$.

3.2 Asymptotic behavior

In this section, we outline the assumptions under which we obtain the asymptotic properties of the test statistics. The following definition introduces some regularity features that the joint density f_{ZW} and the NPIV function $\tilde{\varphi}$ must satisfy.

Definition 3.1 *For a given function γ and for $\alpha \geq 0, s > 0$, the space $\mathcal{B}_\gamma^{s,\alpha}(\mathbb{R}^\ell)$ is the class of functions $g : \mathbb{R}^\ell \mapsto \mathbb{R}$ satisfying: g is everywhere $(m-1)$ -times partially differentiable for $m-1 < s \leq m$; for some $R > 0$ and for all x , the inequality*

$$\sup_{y: \|y-x\| < R} \frac{|g(y) - g(x) - Q(y-x)|}{\|y-x\|^s} \leq \psi(x)$$

holds true, where $Q = 0$ when $m = 1$ and Q is an $(m-1)$ th degree homogeneous polynomial in $(y-x)$ with coefficients the partial derivatives of g at x of orders 1 through $m-1$ when $m > 1$; ψ is uniformly bounded by a constant when $\alpha = 0$ and the functions g and ψ have finite α th moments with respect to $1/\gamma$ when $\alpha > 0$, i.e. $\int |g(x)|^\alpha / \gamma(x) dx < \infty$ and $\int |\psi(x)|^\alpha / \gamma(x) dx < \infty$.

Let f_Z denote the density of Z with respect to the Lebesgue measure and let us define

$$g(z) := \mathbb{E}[(W - \mathbb{E}[W])X | Z = z] f_Z(z).$$

Assumption 3.4 $\tilde{\varphi} \in \mathcal{B}_1^{\rho,0}(\mathbb{R}) \cap L_\pi^2(\mathbb{R})$ and $f_{ZW}/(\pi\tau) \in \mathcal{B}_1^{\rho,0}(\mathbb{R}^2) \cap L_{\pi \otimes \tau}^2(\mathbb{R}^2)$ for a ρ specified below, $\mathbb{E}[U^2 | W = \cdot] f_W / \tau$ is bounded, $\mathbb{E}[X | Z = \cdot] f_Z / \pi \in L_\pi^2(\mathbb{R})$, $\mathbb{E}[X^2(W - \mathbb{E}[W])^2 | Z = \cdot] f_Z / \pi$ is bounded, and $\mathbb{E}[X^2(W - \mathbb{E}[W])^2 | Z = \cdot]$ is bounded.

Assumption 3.4 is a common regularity condition allowing for several degrees of integrability and differentiability (Florens et al., 2012). Under the above assumption $\hat{A} : L_\pi^2(\mathbb{R}) \mapsto L_\tau^2(\mathbb{R})$, $\hat{A}^* : L_\tau^2(\mathbb{R}) \mapsto L_\pi^2(\mathbb{R})$, and $\hat{r} \in L_\tau^2(\mathbb{R})$. Notice that \hat{A}^* is actually the Hilbert adjoint of \hat{A} . This aspect is used multiple times in the proofs.

Assumption 3.5 K_Z and K_W are symmetric kernels of order ρ with bounded support.

The kernel orders in Assumption 3.5 are assumed to be equal only for notational simplicity. To simplify our theoretical exposition, we further assume that $h_W = h_Z$ and denote each bandwidth by h . However, our proofs also hold when such smoothing parameters are set to different values (and the kernels have different orders).

The order ρ of the kernels K_Z and K_W , the bandwidth h and the regularization parameter λ satisfy the following assumption.

Assumption 3.6 $nh^2\lambda^{3/2} \rightarrow \infty$, $n\lambda^2 \rightarrow 0$, $h^\rho\lambda^{-3/4} \rightarrow 0$.

Such conditions allow us to control the regularization bias and the variance of $\hat{\varphi}$ appearing in the expansion of S_n . In particular, the condition on $nh^2\lambda^{3/2}$ allows controlling the "variance term". The conditions on $n\lambda^2$ and $h^\rho\lambda^{-3/4}$ allow us to show that the regularization bias that appears in the expansion of S_n is negligible. This bias is due to the ill-posed nature of the inverse problem.

Let us denote with $\langle \cdot, \cdot \rangle$ the inner product of either $L_\pi^2(\mathbb{R})$ or $L_\tau^2(\mathbb{R})$, and let $\|\cdot\|$ be the norm induced by such an inner product. The specific inner product or norm we refer to will be clear at each time from the context. Let $(\lambda_j, \varphi_j, \psi_j)_j$ be the Singular Value decomposition of the operator A , where $(\lambda_j)_j$ is the sequence of singular values in \mathbb{R} , $(\varphi_j)_j$ is an orthonormal sequence in $L_\pi^2(\mathbb{R})$, and $(\psi_j)_j$ is an orthonormal sequence in $L_\tau^2(\mathbb{R})$ (see Kress (1999)). The following assumptions introduces the usual source conditions.

Assumption 3.7 Let $\langle \cdot, \cdot \rangle$ denote the inner product on $L_\pi^2(\mathbb{R})$. (a) For some $\eta \geq 2$, $\sum_j \frac{|\langle g, \varphi_j \rangle|^2}{\lambda_j^{2\eta}} < \infty$; (b) For some $\theta \geq 2$, $\sum_j \frac{|\langle \tilde{\varphi}, \varphi_j \rangle|^2}{\lambda_j^{2\theta}} < \infty$.

Source conditions are standard in the NPIV literature (see Carrasco et al. (2007) or Darolles et al. (2011)). In general, the source condition is imposed only on $\tilde{\varphi}$ when the interest is on estimating $\tilde{\varphi}$. Here, we need a source condition on both $\tilde{\varphi}$ and g to establish the asymptotic properties of the statistic based on the residuals from the nonparametric regression. To this end, we also need some regularity conditions on the estimator $\hat{\varphi}$. So, let $N_{[\cdot]}(\epsilon, \Phi, \|\cdot\|_{\mathbb{P}})$ be the bracketing number of size ϵ of a class of functions Φ , where $\|\cdot\|_{\mathbb{P}}$ denotes the L_2 norm with respect to the probability law of Z , see Van der Vaart (1998).

Assumption 3.8 *There exists a class of functions Φ such that $\int_0^1 \sqrt{\log N_{[\cdot]}(\epsilon, \Phi, \|\cdot\|_{\mathbb{P}})} d\epsilon < \infty$ and $\mathbb{P}(\hat{\varphi} \in \Phi) \rightarrow 1$.*

The above condition is a high-level assumptions allowing us to handle an empirical process in the expansion of our statistic. It essentially introduces a Donsker feature for the function $\hat{\varphi}$. Such an assumption has been used among others by [Rothe \(2009\)](#), [Escanciano et al. \(2016\)](#), and [Mammen et al. \(2016\)](#). Sufficient conditions for it can be found in [Van der Vaart \(1998\)](#) or [Van der Vaart and Wellner \(1996\)](#). We have also derived alternative proofs based on sample splitting or cross-fitting that avoid such a high-level condition. We have run simulations with cross-fitting and sample-splitting, but we did not notice any improvement in terms of finite sample performance. Thus, we choose to keep the above assumption and avoid sample splitting (or cross-fitting) for a simple implementation of the test. Alternatively such a high-level condition could be avoided by using a Sobolev penalized method for estimating φ . This however would produce much longer proofs and complicate the implementation of the test.

Let \mathbb{P}_n be the empirical mean operator. We have the following theorem.

Theorem 3.2 *Let Assumptions [2.1](#) and [3.3](#) to [3.8](#) hold. Under H_0 ,*

$$\begin{aligned} S_n = & \sqrt{n} \mathbb{P}_n X (W - \mathbb{E}[W]) U - \mathbb{E}[UX] \sqrt{n} \mathbb{P}_n (W - \mathbb{E}[W]) \\ & - \sqrt{n} \mathbb{P}_n U (A(A^* A)^{-1} g)(W) + o_P(1). \end{aligned}$$

If instead $\mathbb{E}[\tilde{U} X (W - \mathbb{E}[W])] \neq 0$, then $|S_n|/\sqrt{n} \rightarrow C \neq 0$.

The proof of the Theorem is in Section S2.1 of the supplementary material. Let us briefly comment on the asymptotic expansion (influence-function representation) of the statistic. The first term on the right hand side is the version of our statistic that we could use had we observed the error U and $\mathbb{E}[W]$. The second term arises because the expectation $\mathbb{E}[W]$ is unobserved and is replaced by its estimated counterpart. Similarly, the third term arises because the error U is unobserved and to estimate it we replace the true function φ with its estimator $\hat{\varphi}$. More in detail, such a term originates from a (nonparametric) bias involving the difference $\hat{\varphi} - \varphi$ that appears in the expansion of S_n . This inflates the asymptotic variance of the statistic, as compared to the case where U is known, and hence represents the price to pay for not observing the error term. To handle this bias term, we rely on decompositions

from Darolles et al. (2011) and Babii and Florens (2017).

The expansion in Theorem 3.2 allows us to obtain the asymptotic normality of our statistic under the null hypothesis. To this end, notice that the first term has zero expectation under the null of homogeneous treatment effects. The second term has clearly a null expectation. Finally, since $\mathbb{E}[U|W] = 0$, the last term also has zero expectation. Therefore, a standard Central Limit Theorem implies that $S_n \xrightarrow{d} \mathcal{N}(0, \Psi)$, where Ψ is a covariance matrix defined by the influence function representation in Theorem 3.2. Since such a covariance matrix has an intricate expression, it would be inconvenient to estimate it in practice. Thus, we suggest to bootstrap the statistic S_n by the pairwise scheme to obtain the p-values of the test. As for the linear test, the fact that $|S_n|/\sqrt{n} \rightarrow C \neq 0$ when $\mathbb{E}[\tilde{U}X(W - \mathbb{E}[W])] \neq 0$ holds implies that the power of the test goes to 1 as n goes to ∞ under alternative hypotheses satisfying $\mathbb{E}[\tilde{U}X(W - \mathbb{E}[W])] \neq 0$.

Notice that we are keeping Assumption 3.3 both under the null of homogeneous treatment effects and under the alternative. We have chosen to do this to simplify the exposition. However, it is possible that if $\mathbb{E}[\tilde{U}X(W - \mathbb{E}[W])] \neq 0$ (and treatment effects are not homogeneous), the NPIV model might be misspecified and/or completeness might not hold. In such a case we would need to modify the proof about the power of the test. Let us briefly discuss these modifications. First, when the model is misspecified and completeness does not hold, it is possible to show that the Tikhonov regularized estimator $\hat{\varphi}$ converges to $\tilde{\varphi}^\perp$ that is the element of $\mathcal{N}(A)^\perp$ (the orthogonal complement of the null space of A) that solves $\min_{\phi \in \mathcal{N}(A)^\perp} \|r - A\phi\|$. In this case, we could define \tilde{U} as $Y - \tilde{\varphi}^\perp(Z)$ and modify the statement of Theorem 3.2 accordingly.

3.3 Simulations

We set $\varphi(z) := (1 + \exp(-z))^{-2}$; $X \sim \mathcal{N}(-0.5, 1)$; $W \sim \mathcal{N}(0, 1)$; $V \sim \mathcal{N}(0, 1)$; $Z = 0.4W + 0.2V$; $U(z) = (V + X)(1 + \gamma z)$ for all $z \in \mathbb{R}$, where X, W, V are mutually independent. The outcome Y is equal to $\varphi(Z) + U(Z)$. When $\gamma = 0$ we are under the null of homogeneous treatment effects, while for $\gamma \neq 0$ we are under the alternative hypothesis. So, γ represents the magnitude of the departure from the null.

To check the robustness of our test with respect to the choices of the smoothing and reg-

ularization parameters, we let them vary around benchmark values. So, we set $(h_Z, h_W) = C_h(h_Z^*, h_W^*)$ and $\lambda = C_\lambda \lambda^*$, where $(h_Z^*, h_W^*) = (\text{sd}(Z)n^{-1/5}, \text{sd}(W)n^{-1/5})$, where $\text{sd}(Z)$ (resp. $\text{sd}(W)$) is the empirical standard deviation of Z (resp. W), while λ^* is the regularization parameter selected according to the Cross-Validation method, as seen in Section 3.1.. Note that (h_Z^*, h_W^*) are the bandwidths set according to a version of Silverman’s Rule of Thumb (Silverman (1986)). C_h and C_λ are fixed constants. We run simulations for $C_h, C_\lambda \in \{0.5, 1, 2\}$ and we consider the modest sample sizes of $n = 100, 250, 500$. As a kernel we use the standard Gaussian density. We set π to the Gaussian density with mean equal to the sample mean of Z and variance equal to twice the sample variance of Z . Similarly, the measure τ is set to the Gaussian density with mean equal to the sample mean of W and variance equal to twice the sample variance of W . Florens et al. (2012) used a similar strategy to select π and τ in their simulation setting. Intuitively, since π and τ have to be two densities and at the same time appear at the denominators in \hat{A} and \hat{A}^* , we want them (a) to integrate to 1 and (b) to converge towards zero sufficiently slowly on the tails. To speed up computations, we use the warp-speed method by Giacomini et al. (2013): for each Monte Carlo iteration we draw a single bootstrap sample, and we use the whole set of bootstrap statistics to compute the bootstrap p-values associated with each original statistic. We perform a very large number of Monte Carlos iterations equal to 10,000.

The results under the null hypothesis of homogeneous treatment effects ($\gamma = 0$) are reported in Table 2. The tests are implemented at 5 and 10 percent nominal levels. The nominal sizes of each of the tests are in bold. The result show that the test is reasonably stable with respect to the choices of λ , h_Z , and h_W and controls well the size under the null hypothesis. Also, as expected, the error in the rejection probability (i.e. the difference between the empirical rejection proportions and the nominal size of the test) tends to become smaller as the sample size increases.

To check the power properties, we run simulations under the alternative hypothesis for different values of γ . The results are shown in Figure 2. We only report results for the benchmark values of h_Z , h_W , and λ and for the nominal size of 5 percent. For the other choices of h_Z , h_W , and λ and for the 10 percent nominal level the results are qualitatively similar. The test shows good power under the alternative hypothesis. As the departure from the null increases, i.e. γ gets further from 0, the test rejects the null hypothesis with an

increasing frequency. Also, such a rejection frequency increases with the sample size at every value of γ .

To sum up, these simulations show that (a) for the moderate sample sizes of $n = 100, 250, 500$ the tests display a good performance both in terms of size control and in terms of power, and (b) the results are stable with respect to the choices of the smoothing and regularization parameters.

		λ^*		$0.5 \lambda^*$		$2 \lambda^*$	
		0.05	0.10	0.05	0.10	0.05	0.10
$n=100$	h^*	0.0642	0.1250	0.0635	0.1246	0.0656	0.1227
	$0.5 h^*$	0.0803	0.1465	0.0800	0.1461	0.0761	0.1500
	$2 h^*$	0.0618	0.1240	0.0612	0.1201	0.0619	0.1277
$n=250$	h^*	0.0583	0.1136	0.0553	0.1101	0.0592	0.1139
	$0.5 h^*$	0.0640	0.1246	0.0661	0.1228	0.0634	0.1247
	$2 h^*$	0.0557	0.1155	0.0553	0.1111	0.0577	0.1163
$n=500$	h^*	0.0509	0.1070	0.0511	0.1046	0.0546	0.1091
	$0.5 h^*$	0.0568	0.1057	0.0556	0.1074	0.0568	0.1072
	$2 h^*$	0.0457	0.0991	0.0449	0.0964	0.0496	0.1011

Table 2: Empirical rejections of the tests under the null hypothesis $\gamma = 0$.

3.4 Application to real data

In this section we apply the test of homogeneous treatment effects to a real dataset. Note that the application of returns to schooling is not well-suited to the nonparametric test because, in this application, the endogenous variable (education) has a support much larger than the instrument (college proximity) which is binary.

Instead, we focus on estimating a fish demand equation. This describes the relationship between the demanded quantity of fish and its price. It is common practice in econometrics to assume that prices are endogenous and estimate demand equations by instrumental variable regressions.

The data we use come from [Graddy \(2006\)](#) and contain daily observations at the New York Fulton fish market about the log of the price, the log of the sold quantity, an indicator for each day of the week, and the wind speed registered in the previous day. The dataset can




Figure 2: Power curve for 5% level tests.

be downloaded at “<https://users.ox.ac.uk/~nuff0078/EconometricModeling/>”. The total number of observations is 111.

We study a demand equation. The outcome Y is equal to the logarithm of the quantity sold and Z is the logarithm of the market price. Market price is likely to be endogenous since it also depends on expected demand, which itself also affects the quantity. Following the literature (Graddy (2006)), we choose a variable related to weather as an instrument, so we set W equal to the wind speed recorded on the day corresponding to the observation. Such a variable is viewed as sufficiently correlated with the price (Z), since the weather affects the ability to fish and is at the same time exogenous with respect to the errors $\{U(z)\}_z$ because it is probably unrelated to demand shocks related to human factors.

The choice of the covariable X is crucial. We set X equal to the indicator for week days (Monday to Thursday). Such a variable should not be correlated with the wind speed since the weather does not depend on the day of the week. In fact, a Kolmogorov-Smirnov test does not reject the hypothesis of independence between X and W (p-value of 0.17). Hence, we can reasonably assume that $(\{U(z)\}_z, X) \perp\!\!\!\perp W_k$, which corresponds to Assumption 2.1.

We perform the nonparametric test using the cross-validated penalty λ^* and bandwidth $(h_Z^*, h_W^*) = (\text{sd}(Z)n^{-1/5}, \text{sd}(W)n^{-1/5})$, where $\text{sd}(Z)$ (resp. $\text{sd}(W)$) is the empirical standard deviation of Z (resp. W) and n is the sample size. This is a version of Silverman’s Rule of Thumb (Silverman (1986)). We use 1,000 bootstrap resamplings. The p-value of the test is 0.0362. Hence, we reject the null hypothesis of homogeneous treatment effects at the 5 percent nominal level. So, we conclude that the causal interpretation of the separable NPIV regression should be doubted for this specific example. This is a reason to use a different approach such as instrumental variable quantile regression (Chernozhukov and Hansen (2008)) which allows heterogeneity of treatment effects by quantiles.

Supporting information

All the proofs along with some examples regarding the analysis of the power of the nonparametric test can be found in the online supplement.

References

- Angrist, J. D., G. W. Imbens, and D. B. Rubin (1996). Identification of causal effects using instrumental variables. *Journal of the American statistical Association* 91, 444–455.
- Babii, A. and J.-P. Florens (2017). Are unobservables separable? *arXiv preprint arXiv:1705.01654*.
- Card, D. (1995). Using geographic variation in college proximity to estimate the return to schooling. In: Christofides, L.N., Grant, K.E., Swidinsky, R. (Eds.), *Aspects of Labour Market Behaviour: Essays in Honour of John Vanderkamp*, 201–222.
- Carrasco, M., J.-P. Florens, and E. Renault (2007, January). Chapter 77 Linear Inverse Problems in Structural Econometrics Estimation Based on Spectral Decomposition and Regularization. In J. J. Heckman and E. E. Leamer (Eds.), *Handbook of Econometrics*, Volume 6, pp. 5633–5751. Elsevier.

- Centorrino, S., F. Feve, and J.-P. Florens (2017, January). Additive Nonparametric Instrumental Regressions: A Guide to Implementation. *Journal of Econometric Methods* 6, 1–25.
- Chen, X., O. Linton, and I. Van Keilegom (2003). Estimation of semiparametric models when the criterion function is not smooth. *Econometrica* 71, 1591–1608.
- Chen, X. and D. Pouzo (2012). Estimation of nonparametric conditional moment models with possibly nonsmooth generalized residuals. *Econometrica* 80, 277–321.
- Chernozhukov, V. and I. Fernández-Val (2005). Subsampling inference on quantile regression processes. *Sankhyā: The Indian Journal of Statistics* 67, 253–276.
- Chernozhukov, V. and C. Hansen (2005). An IV model of quantile treatment effects. *Econometrica* 73, 245–261.
- Chernozhukov, V. and C. Hansen (2008). Instrumental variable quantile regression: A robust inference approach. *Journal of Econometrics* 142(1), 379–398.
- Chung, E. and M. Olivares (2021). Permutation test for heterogeneous treatment effects with a nuisance parameter. *Journal of Econometrics* 225, 148–174.
- Crump, R. K., V. J. Hotz, G. W. Imbens, and O. A. Mitnik (2008). Nonparametric tests for treatment effect heterogeneity. *The Review of Economics and Statistics* 90, 389–405.
- Dai, M. and H. S. Stern (2022). A U-statistic-based test of treatment effect heterogeneity. *Journal of Nonparametric Statistics* 34, 141–163.
- Darolles, S., Y. Fan, J.-P. Florens, and E. Renault (2011). Nonparametric instrumental regression. *Econometrica* 79, 1541–1565.
- D’Haultfoeuille, X. (2011). On the Completeness Condition in Nonparametric Instrumental Problems. *Econometric Theory* 27, 460–471.
- Ding, P., A. Feller, and L. Miratrix (2016). Randomization inference for treatment effect variation. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 78, 655–671.

- Dong, Y. and S. Shen (2018). Testing for rank invariance or similarity in program evaluation. *Review of Economics and Statistics* 100, 78–85.
- Escanciano, J. C., D. Jacho-Chávez, and A. Lewbel (2016). Identification and estimation of semiparametric two-step models. *Quantitative Economics* 7, 561–589.
- Florens, J.-P., J. Johannes, and S. V. Bellegem (2012). Instrumental regression in partially linear models. *Econometrics Journal* 15(2), 304–324.
- Frandsen, B. R. and L. J. Lefgren (2018). Testing rank similarity. *Review of Economics and Statistics* 100, 86–91.
- Giacomini, R., D. N. Politis, and H. White (2013, June). A Warp-Speed Method for Conducting Monte Carlo Experiments Involving Bootstrap Estimators. *Econometric Theory* 29, 567–589.
- Goldman, M. and D. M. Kaplan (2018). Comparing distributions by multiple testing across quantiles or CDF values. *Journal of Econometrics* 206, 143–166.
- Graddy, K. (2006). Markets: the fulton fish market. *Journal of Economic Perspectives* 20, 207–220.
- Horowitz, J. L. and S. Lee (2007). Nonparametric instrumental variables estimation of a quantile regression model. *Econometrica* 75, 1191–1208.
- Hsu, Y.-C. (2017). Consistent tests for conditional treatment effects. *The Econometrics Journal* 20, 1–22.
- Kim, J. H. and B. G. Park (2022). Testing rank similarity in the local average treatment effects model. *Econometric Reviews*, 1–22.
- Koenker, R. and Z. Xiao (2002). Inference on the quantile regression process. *Econometrica* 70, 1583–1612.
- Kress, R. (1999, December). *Linear Integral Equations*. Springer Science & Business Media.
- Li, Q. and J. S. Racine (2007). *Nonparametric econometrics: theory and practice*. Princeton University Press.

- MacKinnon, J. G. and H. White (1985). Some heteroskedasticity-consistent covariance matrix estimators with improved finite sample properties. *Journal of Econometrics* 29, 305–325.
- Mammen, E., C. Rothe, and M. Schienle (2016, October). Semiparametric Estimation with Generated Covariates. *Econometric Theory* 32, 1140–1177.
- Newey, W. K. and J. L. Powell (2003a). Instrumental variable estimation of nonparametric models. *Econometrica* 71, 1565–1578.
- Newey, W. K. and J. L. Powell (2003b). Instrumental Variable Estimation of Nonparametric Models. *Econometrica* 71, 1565–1578.
- Rothe, C. (2009, November). Semiparametric estimation of binary response models with endogenous regressors. *Journal of Econometrics* 153, 51–64.
- Sant’Anna, P. H. (2021). Nonparametric tests for treatment effect heterogeneity with duration outcomes. *Journal of Business & Economic Statistics* 39(3), 816–832.
- Silverman, B. W. (1986). *Density Estimation for Statistics and Data Analysis*, Volume 26 of *Monographs on Statistics and Applied Probability*. London: Chapman & Hall/CRC.
- Van der Vaart, A. W. (1998). *Asymptotic Statistics*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge: Cambridge University Press.
- Van der Vaart, A. W. and J. Wellner (1996). *Weak Convergence and Empirical Processes: With Applications to Statistics*. Springer Series in Statistics. New York: Springer-Verlag.

Online supplement to “Testing for homogeneous treatment effects in linear and nonparametric instrumental variable models”

Jad BEYHUM ^{*}
jad.beyhum@kuleuven.be

Jean-Pierre FLORENS [§]
jean-pierre.florens@tse-fr.eu

Elia LAPENTA [¶]
elia.lapenta@gmail.com

Ingrid VAN KEILEGOM ^{*}
ingrid.vankeilegom@kuleuven.be

April 17, 2023

Section 1 is concerned with the linear test. It contains the proofs of a result regarding Example 1 (1.1) and of Theorem 2.1 (1.2). Section 2 treats the nonparametric test. It includes the proof of the results of Section 3 in the main text (2.1) and a theoretical analysis of the power of the test through examples (2.2).

1 On the linear test

1.1 On example 1

Let us show that $\beta_2^{OLS} = \alpha/3$ in this example. We write

$$Y = \mathbb{E}[Y|Z_2 = 0] + Z_2(\mathbb{E}[Y|Z_2 = 1] - \mathbb{E}[Y|Z_2 = 0]) + V. \quad (1.1)$$

^{*}ORSTAT, KU Leuven. Financial support from the European Research Council (2016-2021, Horizon 2020 / ERC grant agreement No. 694409) is gratefully acknowledged.

[§]Toulouse School of Economics, Université Toulouse Capitole. Jean-Pierre Florens acknowledges funding from the French National Research Agency (ANR) under the Investments for the Future program (Investissements d’Avenir, grant ANR-17-EURE-0010).

[¶]CREST, ENSAE Paris, Institut Polytechnique de Paris

It holds that

$$\begin{aligned}\mathbb{E}[V] &= \mathbb{E}[Y - \mathbb{E}[Y|Z_2 = 0] - Z_2(\mathbb{E}[Y|Z_2 = 1] - \mathbb{E}[Y|Z_2 = 0])] \\ &= \mathbb{E}[Y - \mathbb{E}[Y|Z_2 = 0]|Z_2 = 0]\mathbb{P}(Z_2 = 0) + \mathbb{E}[Y - \mathbb{E}[Y|Z_2 = 1]|Z_2 = 1]\mathbb{P}(Z_2 = 1) = 0\end{aligned}$$

and

$$\begin{aligned}\mathbb{E}[Z_2 V] &= \mathbb{E}[Z_2\{Y - \mathbb{E}[Y|Z_2 = 0] - Z_2(\mathbb{E}[Y|Z_2 = 1] - \mathbb{E}[Y|Z_2 = 0])\}] \\ &= \mathbb{E}[Y - \mathbb{E}[Y|Z_2 = 1]|Z_2 = 1]\mathbb{P}(Z_2 = 1) = 0.\end{aligned}$$

Hence, (1.1) is a linear projection of Y on Z . Since this linear projection is unique (because $E[Z_2^2] > 0$), we get

$$\begin{aligned}\beta_1^{OLS} &= \mathbb{E}[Y|Z_2 = 0] \\ \beta_2^{OLS} &= \mathbb{E}[Y|Z_2 = 1] - \mathbb{E}[Y|Z_2 = 0].\end{aligned}$$

Now, let us compute β_2^{OLS} . We have $\mathbb{E}[Y|Z_2 = 0] = 0$ because $U(Z) = Z_2(1\{\epsilon \geq \frac{3}{4}\} - \frac{1}{4})\alpha$ and

$$\begin{aligned}\mathbb{E}[Y|Z_2 = 1] &= \alpha \mathbb{P}\left(\epsilon \geq \frac{3}{4} \middle| Z_2 = 1\right) \\ &= \alpha \frac{\mathbb{P}(\{\epsilon \geq \frac{3}{4}\} \cap \{Z_2 = 1\})}{\mathbb{P}(Z_2 = 1)} \\ &= \alpha \frac{\mathbb{P}(\{\epsilon \geq \frac{3}{4}\} \cap \{W_2 = 1\})}{3/8} = \alpha \frac{1/8}{3/8} = \frac{\alpha}{3}.\end{aligned}$$

1.2 Proof of Theorem 2.1

Theorem 1 is a direct corollary of the following lemma, which gives the influence function representation of the test statistic.

Lemma 1.1 *Let Assumptions 2.1 and 2.2 in the main text hold. Then, we have*

$$T_n = \Psi \frac{1}{\sqrt{n}} \left(\sum_{i=1}^n V_i \right) + o_P \left(\frac{1}{\sqrt{n}} \right),$$

where

$$V_i = \begin{pmatrix} U_i^{TSLS} X_i \widetilde{W}_{ki} \\ \widetilde{W}_{ki} \\ U_i^{TSLS} W_i \end{pmatrix}; \quad \Psi = \left(I_q, -\mathbb{E}[U^{TSLS} X] I_q, -\mathbb{E}[X \widetilde{W}_k Z^\top] \Sigma^{TSLS} \right);$$

$$\Sigma^{TSLS} = (\mathbb{E}[ZW^\top] \mathbb{E}[WW^\top]^{-1} \mathbb{E}[ZW^\top]^\top)^{-1} \mathbb{E}[ZW^\top] \mathbb{E}[WW^\top]^{-1}; \quad \widetilde{W} = W - E[W].$$

Proof: Since $\widehat{U}_i = Y_i - Z_i^\top \widehat{\beta}^{TSLS} = U_i^{TSLS} - Z_i^\top (\widehat{\beta}^{TSLS} - \beta^{TSLS})$, it holds that

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n \widehat{U}_i X_i (W_{ki} - \overline{W}_k) \\ &= \frac{1}{n} \sum_{i=1}^n U_i^{TSLS} X_i (W_{ki} - \overline{W}_k) - \left(\frac{1}{n} \sum_{i=1}^n X_i (W_{ki} - \overline{W}_k) Z_i^\top \right) (\widehat{\beta}^{TSLS} - \beta^{TSLS}). \end{aligned} \quad (1.2)$$

We deal with the first term on the right-hand side of (1.2). We have

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n U_i^{TSLS} X_i (W_{ki} - \overline{W}_k) \\ &= \frac{1}{n} \sum_{i=1}^n U_i^{TSLS} X_i \widetilde{W}_{ki} + \frac{1}{n} \sum_{i=1}^n U_i^{TSLS} X_i (E[W_k] - \overline{W}_k) \\ &= \frac{1}{n} \sum_{i=1}^n U_i^{TSLS} X_i \widetilde{W}_{ki} - \left[\frac{1}{n} \sum_{i=1}^n U_i^{TSLS} X_i \right] (\overline{W}_k - E[W_k]) \\ &= \frac{1}{n} \sum_{i=1}^n U_i^{TSLS} X_i \widetilde{W}_{ki} - \mathbb{E}[U^{TSLS} X] \left[\frac{1}{n} \sum_{j=1}^n \widetilde{W}_{kj} \right] + o_P \left(\frac{1}{\sqrt{n}} \right). \end{aligned}$$

Then, we handle the second term on the right-hand side of (1.2). It holds that

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n X_i (W_{ki} - \overline{W}_k) Z_i^\top \\ &= \frac{1}{n} \sum_{i=1}^n X_i \widetilde{W}_{ki} Z_i^\top + \frac{1}{n} \sum_{i=1}^n X_i (E[W_k] - \overline{W}_k) Z_i^\top \\ &= \frac{1}{n} \sum_{i=1}^n X_i \widetilde{W}_{ki} Z_i^\top - (E[W_k] - \overline{W}_k) \left[\frac{1}{n} \sum_{i=1}^n X_i Z_i^\top \right]. \end{aligned}$$

Since $\widehat{\beta}^{TSLS} - \beta = O_P(1/\sqrt{n})$, and by the law of large numbers

$$(E[W_k] - \overline{W}_k) \left[\frac{1}{n} \sum_{i=1}^n X_i Z_i^\top \right] = o_P(1),$$

we get

$$\begin{aligned} \left(\frac{1}{n} \sum_{i=1}^n X_i(W_{ki} - \bar{W}_k) Z_i^\top \right) (\hat{\beta}^{TSLs} - \beta) &= \frac{1}{n} \sum_{i=1}^n X_i \widetilde{W}_{ki} Z_i^\top (\hat{\beta}^{TSLs} - \beta) + o_P \left(\frac{1}{\sqrt{n}} \right) \\ &= \mathbb{E}[X(W_k - \mathbb{E}[W_k]) Z^\top] (\hat{\beta}^{TSLs} - \beta) + o_P \left(\frac{1}{\sqrt{n}} \right). \end{aligned}$$

Moreover, by standard computations, we have

$$\hat{\beta}^{TSLs} = \Sigma^{TSLs} \left(\frac{1}{n} \sum_{i=1}^n U_i^{TSLs} W_i \right) + \beta + o_P \left(\frac{1}{\sqrt{n}} \right),$$

which leads to the result. □

2 On the nonparametric test

2.1 Proof of the results of Section 3

Recall that \mathbb{P}_n represents the empirical mean operator. Let us denote with \mathbb{P} the population mean operator, so $\mathbb{P}f(\xi) = \mathbb{E}f(\xi)$ for any random variable ξ . We finally denote with $\|\cdot\|_\pi$ the norm of the $L_\pi^2(\mathbb{R})$ space, i.e. $\|f\|_\pi^2 = \int |f(z)|^2 \pi(dz)$ for any $f \in L_\pi^2(\mathbb{R})$.

2.1.1 Proof of Theorem 3.1

Let us decompose the statistic as follows:

$$\begin{aligned} \sqrt{n} \mathbb{P}_n \hat{U} X(W - \bar{W}) &= \sqrt{n} \mathbb{P}_n U X(W - \mathbb{E}[W]) \\ &\quad + \sqrt{n} (\mathbb{P}_n - \mathbb{P})(\varphi - \hat{\varphi}) X(W - \mathbb{E}[W]) \\ &\quad + \sqrt{n} \mathbb{P}(\varphi - \hat{\varphi}) X(W - \mathbb{E}[W]) \\ &\quad - \sqrt{n} \mathbb{P}_n \hat{U} X(\bar{W} - \mathbb{E}[W]). \end{aligned} \tag{2.3}$$

We start by showing that the last term on the RHS equals $\mathbb{E}[UX] \cdot \sqrt{n} \mathbb{P}_n(W - \mathbb{E}[W]) + o_P(1)$. Notice that $\sqrt{n} \mathbb{P}_n \hat{U} X(\bar{W} - \mathbb{E}[W]) = \left[\mathbb{P}_n \hat{U} X \right] \cdot \sqrt{n} (\mathbb{P}_n - \mathbb{P}) W$. By Assumption 3.6 in the main text, with probability approaching one

$$\left| (\mathbb{P}_n - \mathbb{P}) \hat{U} X \right| \leq \sup_{\phi \in \Upsilon} |(\mathbb{P}_n - \mathbb{P}) \phi|,$$

where $\Upsilon := \{(u, z, x) \mapsto (y - \phi(z))x : \phi \in \Phi\}$. It follows from the definition of bracketing number that $N_{[\cdot]}(\epsilon, \Upsilon, \|\cdot\|_{\mathbb{P}}) \leq N_{[\cdot]}(\mathbb{E}[X^2] \cdot \epsilon, \Phi, \|\cdot\|_{\mathbb{P}})$. Assumption 3.6 ensures that the RHS of this inequality is finite for any ϵ . So, the class Υ is Glivenko-Cantelli and by Theorem 19.4 in [Van der Vaart \(1998\)](#) we get that $(\mathbb{P}_n - \mathbb{P})\phi = o_P(1)$ uniformly in $\phi \in \Upsilon$. This and the previous display lead to

$$\mathbb{P}_n \widehat{U} X = \mathbb{P} U X + \mathbb{P}(\varphi - \widehat{\varphi}) X + o_P(1).$$

To show the negligibility of the second term on the RHS, by the Law of Iterated Expectations and the Cauchy-Schwartz inequality we get

$$\begin{aligned} |\mathbb{P}(\widehat{\varphi} - \varphi)X|^2 &= \left| \int (\widehat{\varphi}(z) - \varphi(z)) \mathbb{E}[X|Z=z] \frac{f_Z(z)}{\pi(z)} \pi(dz) \right|^2 \\ &\leq \|\widehat{\varphi} - \varphi\|_{\pi}^2 \cdot \|\mathbb{E}[X|Z=\cdot] f_Z/\pi\|_{\pi}^2 \end{aligned}$$

where $\|\cdot\|_{\pi}$ denotes the norm on $L_{\pi}^2(\mathbb{R})$. The second factor on the RHS is finite by Assumption 3.2 in the main text. The first factor is instead $o_P(1)$ by Lemma 2.6. By putting together these results, we find that

$$\sqrt{n} \mathbb{P}_n \widehat{U} X (\overline{W} - \mathbb{E}[W]) = \mathbb{E}[UX] \cdot \sqrt{n} \mathbb{P}_n (W - \mathbb{E}[W]) + o_P(1). \quad (2.4)$$

We now show the negligibility of the second term on the RHS of Equation (2.3). Combining the Law of Iterated Expectations and Assumption 3.2 leads to

$$\begin{aligned} \|(\widehat{\varphi} - \varphi)X(W - \mathbb{E}[W])\|_P^2 &= \int |\widehat{\varphi}(z) - \varphi(z)|^2 \frac{\mathbb{E}[X^2(W - \mathbb{E}[W])^2|Z=z] f_Z(z)}{\pi(z)} \pi(dz) \\ &\leq C \|\widehat{\varphi} - \varphi\|_{\pi}^2, \end{aligned}$$

with $\|\widehat{\varphi} - \varphi\|_{\pi}^2 = o_P(1)$, as previously found. So, by Assumption 3.6 and the boundedness of $\mathbb{E}[X^2(W - \mathbb{E}[W])^2|Z=\cdot]$ (see Assumption 3.2), the conditions of Lemma 2.4 are satisfied and we obtain

$$\sqrt{n}(\mathbb{P}_n - \mathbb{P})(\widehat{\varphi} - \varphi) X (W - \mathbb{E}[W]) = o_P(1). \quad (2.5)$$

Gathering together Equations (2.3), (2.5), (2.4), and the result of Lemma 2.2 ahead delivers the desired result under the null hypothesis of error invariance.

To show the behavior of the statistic under the alternative, we can proceed as at the beginning of this proof to obtain

$$\mathbb{P}_n \widehat{U} X (W_k - \mathbb{E}[W_k]) = \mathbb{E}[\widetilde{U} X (W - \mathbb{E}[W])] + o_P(1)$$

where $\tilde{U} = Y - \tilde{\varphi}(Z)$ and the first leading term is different from 0 under H_1 .

2.1.2 Auxiliary lemma

In this section we provide an auxiliary lemma we used in the proofs of Theorem 3.1. Let us denote with $\langle \cdot, \cdot \rangle$ the inner product on either $L_\pi^2(\mathbb{R})$ or $L_\tau^2(\mathbb{R})$, and let $\| \cdot \|$ denote the norm induced by such an inner product. Whether $\langle \cdot, \cdot \rangle$ and $\| \cdot \|$ refer to either $L_\pi^2(\mathbb{R})$ or $L_\tau^2(\mathbb{R})$ will be clear by their arguments. Also, given the operators A and A^* we define

$$\|A\|_{op}^2 := \sup_{\phi \in L_\pi^2(\mathbb{R}): \|\phi\|=1} \|A\phi\|^2 \quad \text{and} \quad \|A^*\|_{op}^2 := \sup_{\psi \in L_\tau^2(\mathbb{R}): \|\psi\|=1} \|A^*\psi\|^2.$$

Since A and A^* are linear bounded operators, both $\|A\|_{op}^2$ and $\|A^*\|_{op}^2$ are finite. Also, since A^* is the Hilbert adjoint of A , it holds that $\|A\|_{op}^2$ and $\|A^*\|_{op}^2$ are equal.

Lemma 2.2 *Under Assumptions 3.1 to 3.6 in the main text, we have*

$$\sqrt{n} \mathbb{P}(\hat{\varphi} - \tilde{\varphi})g = \sqrt{n} \mathbb{P}_n \tilde{U} [A(A^*A)^{-1}g](W) + o_P(1).$$

Proof: Let us start with the following decomposition

$$\hat{\varphi} - \tilde{\varphi} = \Xi_1 + \Xi_2 + \Xi_3 + \Xi_4 + (\tilde{\varphi}_\lambda - \tilde{\varphi}) \tag{2.6}$$

where

$$\begin{aligned} \Xi_1 &:= (\lambda I + A^*A)^{-1} A^*(\hat{r} - \hat{A}\tilde{\varphi}) \\ \Xi_2 &:= (\lambda I + A^*A)^{-1} (\hat{A}^* - A^*)(\hat{r} - \hat{A}\tilde{\varphi}) \\ \Xi_3 &:= \left[(\lambda I + \hat{A}^*\hat{A})^{-1} - (\lambda I + A^*A)^{-1} \right] \hat{A}^*(\hat{r} - \hat{A}\tilde{\varphi}) \\ \Xi_4 &:= (\lambda I + \hat{A}^*\hat{A})^{-1} \hat{A}^*\hat{A}(\tilde{\varphi} - \tilde{\varphi}_\lambda) \\ \tilde{\varphi}_\lambda &:= (\lambda I + A^*A)^{-1} A^*r. \end{aligned}$$

The previous decomposition holds because

$$\begin{aligned} \Xi_1 + \Xi_2 + \Xi_3 &= (\lambda I + A^*A)^{-1} A^*(\hat{r} - \hat{A}\tilde{\varphi}) \\ &\quad + (\lambda I + A^*A)^{-1} \hat{A}^*(\hat{r} - \hat{A}\tilde{\varphi}) - (\lambda I + A^*A)^{-1} A^*(\hat{r} - \hat{A}\tilde{\varphi}) \\ &\quad + (\lambda I + \hat{A}^*\hat{A})^{-1} \hat{A}^*(\hat{r} - \hat{A}\tilde{\varphi}) - (\lambda I + A^*A)^{-1} \hat{A}^*(\hat{r} - \hat{A}\tilde{\varphi}) \\ &= \hat{\varphi} - (\lambda I + \hat{A}^*\hat{A})^{-1} \hat{A}^*\hat{A}\tilde{\varphi}. \end{aligned}$$

Now,

$$\begin{aligned}
|\langle \Xi_2, g \rangle| &= \left| \left\langle (\hat{A}^* - A^*)(\hat{r} - \hat{A}\tilde{\varphi}), (\lambda I + A^*A)^{-1}g \right\rangle \right| \\
&\leq \left\| \hat{A} - A \right\|_{op} \cdot \left\| \hat{r} - \hat{A}\tilde{\varphi} \right\| \cdot \|(\lambda I + A^*A)^{-1}g\| \\
&\leq C \left\| \hat{A} - A \right\|_{op} \cdot \left\| \hat{r} - \hat{A}\tilde{\varphi} \right\| \lambda^{\frac{\eta \wedge 2}{2} - 1},
\end{aligned}$$

where the second line follows from the Cauchy-Schwartz inequality and the continuity of $\hat{A} - A$, while the third line follows from Assumption 3.5(a) and Lemma 2.5(d). Assumption 3.5(a) ensures that $\eta \geq 2$. As already noticed before, $\|\hat{A} - A\|_{op}/\sqrt{\lambda} = o_P(1)$ and $\|\hat{r} - \hat{A}\tilde{\varphi}\|/\sqrt{\lambda} = o_P(1)$. So, by the above display and Assumption 3.4 we find

$$\sqrt{n} \langle \Xi_2, g \rangle = O_P(\sqrt{n\lambda^2}) = o_P(1). \quad (2.7)$$

We now handle the term $\sqrt{n} \langle \tilde{\varphi}_\lambda - \tilde{\varphi}, g \rangle$. By Assumption 3.5(b) and Lemma 2.5(f) $\|\tilde{\varphi}_\lambda - \tilde{\varphi}\| = O(\lambda^{\frac{\theta \wedge 2}{2}})$ with $\theta \geq 2$. Combining this rate with the Cauchy-Schwartz inequality gives

$$\sqrt{n} |\langle \tilde{\varphi}_\lambda - \tilde{\varphi}, g \rangle| \leq \sqrt{n} \cdot \|\tilde{\varphi}_\lambda - \tilde{\varphi}\| \cdot \|g\| = O(\sqrt{n\lambda^2}) = o(1). \quad (2.8)$$

To show the negligibility of $\sqrt{n} \langle \Xi_3, g \rangle$, notice that

$$\begin{aligned}
\Xi_3 &= (\lambda I + A^*A)^{-1} \left[(\lambda I + A^*A) - (\lambda I + \hat{A}^*\hat{A}) \right] (\lambda I + \hat{A}^*\hat{A})^{-1} \hat{A}^*(\hat{r} - \hat{A}\tilde{\varphi}) \\
&= (\lambda I + A^*A)^{-1} \left[A^*(A - \hat{A}) + (A^* - \hat{A}^*)\hat{A} \right] (\lambda I + \hat{A}^*\hat{A})^{-1} \hat{A}^*(\hat{r} - \hat{A}\tilde{\varphi}) \\
&= (\lambda I + A^*A)^{-1} A^*(A - \hat{A})(\lambda I + \hat{A}^*\hat{A})^{-1} \hat{A}^*(\hat{r} - \hat{A}\tilde{\varphi}) \\
&\quad + (\lambda I + A^*A)^{-1} (A^* - \hat{A}^*)\hat{A}(\lambda I + \hat{A}^*\hat{A})^{-1} \hat{A}^*(\hat{r} - \hat{A}\tilde{\varphi}).
\end{aligned}$$

In view of the above decomposition and the Cauchy-Schwartz inequality

$$\begin{aligned}
\sqrt{n} |\langle \Xi_3, g \rangle| &\leq \sqrt{n} \left| \left\langle \hat{r} - \hat{A}\tilde{\varphi}, \hat{A}(\lambda I + \hat{A}^*\hat{A})^{-1} (A^* - \hat{A}^*)A(\lambda I + A^*A)^{-1}g \right\rangle \right| \\
&\quad + \sqrt{n} \left| \left\langle \hat{r} - \hat{A}\tilde{\varphi}, \hat{A}(\lambda I + \hat{A}^*\hat{A})^{-1} \hat{A}^*(A - \hat{A})(\lambda I + A^*A)^{-1}g \right\rangle \right| \\
&\leq \sqrt{n} \left\| \hat{r} - \hat{A}\tilde{\varphi} \right\| \cdot \left\| \hat{A}(\lambda I + \hat{A}^*\hat{A})^{-1} (A^* - \hat{A}^*)A(\lambda I + A^*A)^{-1}g \right\| \\
&\quad + \sqrt{n} \left\| \hat{r} - \hat{A}\tilde{\varphi} \right\| \cdot \left\| \hat{A}(\lambda I + \hat{A}^*\hat{A})^{-1} \hat{A}^*(A - \hat{A})(\lambda I + A^*A)^{-1}g \right\| \\
&\leq \sqrt{n} \cdot \left\| \hat{r} - \hat{A}\tilde{\varphi} \right\| \cdot \left\| \hat{A}(\lambda I + \hat{A}^*\hat{A})^{-1} \right\|_{op} \cdot \left\| \hat{A} - A \right\|_{op} \cdot \|A(\lambda I + A^*A)^{-1}g\| \\
&\quad + \sqrt{n} \cdot \left\| \hat{r} - \hat{A}\tilde{\varphi} \right\| \cdot \left\| \hat{A}(\lambda I + \hat{A}^*\hat{A})^{-1} \hat{A}^* \right\|_{op} \cdot \left\| \hat{A} - A \right\|_{op} \cdot \|(\lambda I + A^*A)^{-1}g\|.
\end{aligned}$$

Lemma 2.5(a) and (c) imply that $\|\widehat{A}(\lambda I + \widehat{A}^* \widehat{A})^{-1} \widehat{A}^*\|_{op} \leq 1$ and $\|\widehat{A}(\lambda I + \widehat{A}^* \widehat{A})^{-1}\|_{op} \lesssim 1/\sqrt{\lambda}$. Combining Assumption 3.5(a) and Lemma 2.5(d) and (e) gives $\|(\lambda I + A^* A)^{-1} g\| \lesssim \lambda^{\frac{\eta \wedge 2}{2} - 1}$ and $\|A(\lambda I + A^* A)^{-1} g\| \lesssim \lambda^{\frac{\eta+1}{2} \wedge 1 - 1}$ with $\eta \geq 2$. Finally, Assumption 3.4 and Lemma 2.3 ensure that $\|\widehat{A} - A\|_{op} \lambda^{-3/4} = o_P(1)$ and $\|\widehat{r} - \widehat{A} \widetilde{\varphi}\| \lambda^{-3/4} = o_P(1)$. Hence, by these rates and the previous display we obtain

$$\sqrt{n} |\langle \Xi_3, g \rangle| \leq \sqrt{n} o_P(\lambda) = o_P(1). \quad (2.9)$$

To show that $\sqrt{n} \langle \Xi_4, g \rangle$ is negligible, we first define the operator

$$f \mapsto (A^* A)^{\eta/2} f := \sum_j \lambda_j^\eta \langle f, \varphi_j \rangle \varphi_j, \quad (A^* A)^{\eta/2} : L^2(\mathbb{R}) \mapsto L^2(\mathbb{R}), \quad (2.10)$$

and let

$$\widetilde{f} := \sum_j \lambda_j^{-\eta} \langle g, \varphi_j \rangle \varphi_j. \quad (2.11)$$

Notice that by Assumption 3.5, $\|\widetilde{f}\| < \infty$ for some $\eta \geq 2$. Also, $(A^* A)^{\eta/2} \widetilde{f} = g$. By using these for $\eta = 2$ we obtain

$$\begin{aligned} \sqrt{n} |\langle \Xi_4, g \rangle| &= \sqrt{n} \left| \left\langle \Xi_4, (A^* A)^{\eta/2} \widetilde{f} \right\rangle \right| \\ &= \sqrt{n} \left| \left\langle \Xi_4, (A^* A) \widetilde{f} \right\rangle \right| \\ &\leq \sqrt{n} \left| \left\langle \Xi_4, (A^* - \widehat{A}^*) A \widetilde{f} \right\rangle \right| + \sqrt{n} \left| \left\langle \Xi_4, \widehat{A}^* A \widetilde{f} \right\rangle \right| \\ &= \sqrt{n} \left| \left\langle \Xi_4, (A^* - \widehat{A}^*) A \widetilde{f} \right\rangle \right| + \sqrt{n} \left| \left\langle \widehat{A} \Xi_4, A \widetilde{f} \right\rangle \right| \\ &\leq \sqrt{n} \cdot \|\Xi_4\| \cdot \|\widehat{A} - A\|_{op} \cdot \|A \widetilde{f}\| + \sqrt{n} \cdot \|\widehat{A} \Xi_4\| \cdot \|A \widetilde{f}\|, \end{aligned} \quad (2.12)$$

where the second equality follows from $(A^* A)^{\eta/2} = A^* A$ for $\eta = 2$. Now, to bound the RHS

we decompose Ξ_4 as

$$\begin{aligned}
\Xi_4 &= (\lambda I + \hat{A}^* \hat{A})^{-1} \hat{A}^* \hat{A} \tilde{\varphi} - (\lambda I + A^* A)^{-1} A^* A \tilde{\varphi} \\
&= \left[(\lambda I + \hat{A}^* \hat{A})^{-1} (\lambda I + \hat{A}^* \hat{A} - \lambda I) - (\lambda I + A^* A)^{-1} (\lambda I + A^* A - \lambda I) \right] \tilde{\varphi} \\
&= \lambda \left[(\lambda I + A^* A)^{-1} - (\lambda I + \hat{A}^* \hat{A})^{-1} \right] \tilde{\varphi} \\
&= \lambda (\lambda I + \hat{A}^* \hat{A})^{-1} \left[(\lambda I + \hat{A}^* \hat{A}) - (\lambda I + A^* A) \right] (\lambda I + A^* A)^{-1} \tilde{\varphi} \\
&= \lambda (\lambda I + \hat{A}^* \hat{A})^{-1} \left[\hat{A}^* (\hat{A} - A) + (\hat{A}^* - A^*) A \right] (\lambda I + A^* A)^{-1} \tilde{\varphi} \\
&= \lambda (\lambda I + \hat{A}^* \hat{A})^{-1} \hat{A}^* (\hat{A} - A) (\lambda I + A^* A)^{-1} \tilde{\varphi} \\
&\quad + \lambda (\lambda I + \hat{A}^* \hat{A})^{-1} (\hat{A}^* - A^*) A (\lambda I + A^* A)^{-1} \tilde{\varphi} .
\end{aligned} \tag{2.13}$$

By the above decomposition, we find that

$$\begin{aligned}
\|\Xi_4\| &\leq \|\lambda (\lambda I + \hat{A}^* \hat{A})^{-1}\|_{op} \cdot \|\hat{A}^*\|_{op} \cdot \|\hat{A} - A\|_{op} \cdot \|(\lambda I + A^* A)^{-1} \tilde{\varphi}\| \\
&\quad + \|(\lambda (\lambda I + \hat{A}^* \hat{A})^{-1})\|_{op} \cdot \|\hat{A} - A\|_{op} \cdot \|A (\lambda I + A^* A)^{-1} \tilde{\varphi}\| \\
&\leq C \|\hat{A}^*\|_{op} \cdot \|\hat{A} - A\|_{op} \lambda^{\frac{\theta \wedge 2}{2} - 1} + \|\hat{A} - A\|_{op} \lambda^{\frac{\theta+1}{2} \wedge 1 - 1} ,
\end{aligned} \tag{2.14}$$

where the last inequality follows from $\|(\lambda (\lambda I + \hat{A}^* \hat{A})^{-1})\|_{op} \leq 2$ (see Lemma 2.5(b)), $\|(\lambda I + A^* A)^{-1} \tilde{\varphi}\| \leq C \lambda^{\frac{\theta \wedge 2}{2} - 1}$ (see Lemma 2.5(d) and Assumption 3.5(b)), and $\|A (\lambda I + A^* A)^{-1} \tilde{\varphi}\| \leq C \lambda^{\frac{\theta+1}{2} \wedge 1 - 1}$ (see Lemma 2.5(e) and Assumption 3.5(b)).

Similarly, the decomposition in (2.13) leads to

$$\begin{aligned}
\|\hat{A} \Xi_4\| &\leq \|\hat{A} (\lambda I + \hat{A}^* \hat{A})^{-1} \hat{A}^*\|_{op} \cdot \|\hat{A} - A\|_{op} \cdot \lambda \cdot \|(\lambda I + A^* A)^{-1} \tilde{\varphi}\| \\
&\quad + \lambda \cdot \|\hat{A} (\lambda I + \hat{A}^* \hat{A})^{-1}\|_{op} \cdot \|\hat{A}^* - A^*\|_{op} \cdot \|A (\lambda I + A^* A)^{-1} \tilde{\varphi}\| \\
&\leq \|\hat{A} - A\|_{op} \cdot \lambda^{\frac{\theta \wedge 2}{2}} + \|\hat{A} - A\|_{op} \cdot \lambda^{\frac{\theta+1}{2} \wedge 1 - 1/2} ,
\end{aligned} \tag{2.15}$$

where the last equality follows from Lemma 2.5(a)(c)(d)(e) and Assumption 3.5(b). So, recalling that $\theta \geq 2$, that $\|\hat{A} - A\|_{op} / \sqrt{\lambda} = o_P(1)$, and putting together (2.15), (2.14), and (2.12) gives

$$\sqrt{n} |\langle \Xi_4, g \rangle| = O_P \left(\sqrt{n} \|\hat{A} - A\|_{op}^2 + \sqrt{n} \lambda \right) = o_P(\sqrt{n} \lambda) = o_P(1) , \tag{2.16}$$

where in the last equality we have used $n \lambda^2 = o(1)$ (see Assumption 3.4).

By gathering (2.16), (2.9), (2.8), (2.7), and the decomposition in Equation (2.6), we obtain

$$\sqrt{n} \langle \hat{\varphi} - \tilde{\varphi}, g \rangle = \sqrt{n} \langle \Xi_1, g \rangle + o_P(1) . \quad (2.17)$$

So, to show the desired result it suffices to obtain an Influence Function Representation for the leading term of the above display. Now, by the definition of Ξ_1 we get

$$\begin{aligned} \sqrt{n} \langle \Xi_1, g \rangle &= \left\langle \sqrt{n} A^* (\hat{r} - \hat{A} \tilde{\varphi}), (A^* A)^{-1} g \right\rangle \\ &\quad + \left\langle \sqrt{n} A^* (\hat{r} - \hat{A} \tilde{\varphi}), [(\lambda I + A^* A)^{-1} - (A^* A)^{-1}] g \right\rangle . \end{aligned} \quad (2.18)$$

By using a change of variable, Assumption 3.2 about $\tilde{\varphi}$, and the ρ th order of the kernel K_Z , we obtain

$$\begin{aligned} \hat{r}(w) - (\hat{A} \tilde{\varphi})(w) &= \frac{1}{nh} \sum_{i=1}^n Y_i K_W \left(\frac{w - W_i}{h} \right) \frac{1}{\tau(w)} \\ &\quad - \frac{1}{nh} \sum_{i=1}^n K_W \left(\frac{w - W_i}{h} \right) \frac{h^{-1}}{\tau(w)} \int \tilde{\varphi}(z) K_Z \left(\frac{z - Z_i}{h} \right) dz \\ &= \frac{1}{nh} \sum_{i=1}^n U_i K_W \left(\frac{w - W_i}{h} \right) \frac{1}{\tau(w)} \\ &\quad - \frac{1}{nh} \sum_{i=1}^n K_W \left(\frac{w - W_i}{h} \right) \frac{1}{\tau(w)} \int [\tilde{\varphi}(Z_i + vh) - \tilde{\varphi}(Z_i)] K_Z(v) dv \\ &= \frac{1}{nh^1} \sum_{i=1}^n U_i K_W \left(\frac{w - W_i}{h} \right) \frac{1}{\tau(w)} \\ &\quad - \frac{h^\rho}{nh^1} \sum_{i=1}^n K_W \left(\frac{w - W_i}{h} \right) \frac{1}{\tau(w)} S_n(Z_i) \text{ with } |S_n(z)| \leq C , \end{aligned}$$

By the previous display and the definition of A^* we obtain (see the comments below)

$$\begin{aligned}
\sqrt{n} \left[A^*(\hat{r} - \hat{A}\tilde{\varphi}) \right] (z) &= \frac{1}{\sqrt{n}} \sum_{i=1}^n \tilde{U}_i h^{-1} \int \frac{f_{ZW}(z, w)}{\pi(z) \tau(w)} K_W \left(\frac{w - W_i}{h} \right) dw \\
&\quad + \frac{h^\rho}{\sqrt{n}} \sum_{i=1}^n S_n(Z_i) h^{-1} \int \frac{f_{ZW}(z, w)}{\pi(z) \tau(w)} K_W \left(\frac{w - W_i}{h} \right) dw \\
&= \frac{1}{\sqrt{n}} \sum_{i=1}^n \tilde{U}_i \int \frac{f_{ZW}(z, W_i + vh)}{\pi(z) \tau(W_i + vh)} K_W(v) dv \\
&\quad + \frac{h^\rho}{\sqrt{n}} \sum_{i=1}^n S_n(Z_i) \int \frac{f_{ZW}(z, W_i + vh)}{\pi(z) \tau(W_i + vh)} K_W(v) dv \\
&= \frac{1}{\sqrt{n}} \sum_{i=1}^n \tilde{U}_i \frac{f_{ZW}(z, W_i)}{\pi(z) \tau(W_i)} + \frac{h^\rho}{\sqrt{n}} \sum_{i=1}^n U_i S_n^{(2)}(W_i, z) \\
&\quad - \frac{h^\rho}{\sqrt{n}} \sum_{i=1}^n S_n(Z_i) \frac{f_{ZW}(z, W_i)}{\pi(z) \tau(W_i)} - \frac{h^{2\rho}}{\sqrt{n}} \sum_{i=1}^n S_n(Z_i) S_n^{(3)}(W_i, z) \quad (2.19) \\
&\quad \text{with } |S_n^{(2)}(w, z)| \leq C \text{ and } |S_n^{(3)}(w, z)| \leq C,
\end{aligned}$$

where in the second equality we have used a change of variable, while in the third equality we have combined Assumption 3.2 with the ρ th order of the kernel K_W . By the iid assumption and since $\mathbb{E}[\tilde{U}|W] = 0$,

$$\begin{aligned}
\mathbb{E} \|\text{1st term RHS of (2.19)}\|^2 &= \mathbb{E} \frac{1}{n} \sum_{i,j} \tilde{U}_i \tilde{U}_j \int \frac{f_{ZW}(z, W_i)}{\pi(z) \tau(W_i)} \frac{f_{ZW}(z, W_j)}{\pi(z) \tau(W_j)} \pi(dz) \\
&= \int \left[\frac{f_{ZW}(z, w)}{\pi(z) \tau(w)} \right]^2 \frac{\sigma_W(w)^2 f_W(w)}{\tau(w)} \pi(dz) \otimes \tau(dw),
\end{aligned}$$

where $\sigma_W^2 := \mathbb{E}[\tilde{U}^2|W = \cdot]$. Assumption 3.2 ensures that $\sigma_W^2 f_W/\tau$ is bounded and $f_{ZW}/(\pi \tau) \in L_{\pi \otimes \tau}^2$, so the RHS of the above display is finite. Similarly,

$$\begin{aligned}
\mathbb{E} \|\text{2nd term RHS of (2.19)}\|^2 &= h^{2\rho} \int \frac{\sigma_W^2(w) f_W(w)}{\tau(w)} |S_n^{(2)}(w, z)|^2 \pi(dz) \otimes \tau(dw) \\
&= O(h^{2\rho}) = o(1),
\end{aligned}$$

where the last equality has used the fact that $|S_n^{(2)}(w, z)| \leq C$. Similar arguments show that

$$\begin{aligned}
\mathbb{E} \|\text{3rd term RHS of (2.19)}\|^2 &= \frac{h^{2\rho}}{n} \sum_{i,j} \mathbb{E}[S_n(Z_i)S_n(Z_j)] \int \frac{f_{ZW}(z, W_i)}{\pi(z)\tau(W_i)} \frac{f_{ZW}(z, W_j)}{\pi(z)\tau(W_j)} \pi(dz) \\
&\leq C h^{2\rho} \int \left[\frac{f_{ZW}(z, w)}{\pi(z)\tau(w)} \right]^2 \frac{f_W(w)}{\tau(w)} \pi(dz) \otimes \tau(dw) \\
&\quad + C h^{2\rho} \frac{n(n-1)}{n} \int \left\{ \int \frac{f_{ZW}(z, w)}{\pi(z)\tau(w)} \frac{f_W(w)}{\tau(w)} \tau(dw) \right\}^2 \pi(dz) \\
&\leq C h^{2\rho} n \int \left[\frac{f_{ZW}(z, w)}{\pi(z)\tau(w)} \right]^2 \pi(dz) \otimes \tau(dw) \\
&= o(1),
\end{aligned}$$

where for the first inequality we have used the boundedness of S_n , for the second inequality we have used the boundedness of f_W/τ (see Assumption 3.2) and the Cauchy-Schwartz inequality, while for the last equality we have used Assumption 3.4.

By proceeding along the same lines we find that

$$\mathbb{E} \|\text{4th term RHS of (2.19)}\|^2 \leq C h^{4\rho} n = o(1).$$

We are now able to obtain the Influence function representation for $\sqrt{n} \langle \Xi_1, g \rangle$ in Equation (2.18). By the previous five displays and since $\|(A^*A)^{-1}g\| < \infty$, the first term on the RHS of (2.18) equals

$$\begin{aligned}
\left\langle \sqrt{n}A^*(\hat{r} - \hat{A}\tilde{\varphi}), (A^*A)^{-1}g \right\rangle &= \left\langle \frac{1}{\sqrt{n}} \sum_{i=1}^n \tilde{U}_i \frac{f_{ZW}(\cdot, W_i)}{\pi(\cdot)\tau(W_i)}, (A^*A)^{-1}g \right\rangle + o_P(1) \\
&= \frac{1}{\sqrt{n}} \sum_{i=1}^n \tilde{U}_i \int \frac{f_{ZW}(z, W_i)}{\tau(W_i)} [(A^*A)^{-1}g](z) dz + o_P(1) \\
&= \frac{1}{\sqrt{n}} \sum_{i=1}^n \tilde{U}_i [A(A^*A)^{-1}g](W_i) + o_P(1).
\end{aligned}$$

To conclude the proof, notice first that Equation (2.19) and the four displays after it ensure that $\|\sqrt{n}A^*(\hat{r} - \hat{A}\tilde{\varphi})\| = O_P(1)$. So, the norm of the second term on the RHS of (2.18) is upperbounded as follows

$$\left| \left\langle \sqrt{n}A^*(\hat{r} - \hat{A}\tilde{\varphi}), [(\lambda I + A^*A)^{-1} - (A^*A)^{-1}]g \right\rangle \right| \leq O_P(\|[(\lambda I + A^*A)^{-1} - (A^*A)^{-1}]g\|).$$

By recalling that we have set $\eta = 2$, we have $(A^*A)^{\eta/2} = A^*A$. Thus, by definition of \tilde{f} we get $g = (A^*A)^{\eta/2}\tilde{f} = A^*A\tilde{f}$ and

$$[(\lambda I + A^*A)^{-1} - (A^*A)^{-1}]g = (\lambda I + A^*A)^{-1}A^*(A\tilde{f}) - \tilde{f}.$$

Since $(\lambda I + A^*A)^{-1}A^*$ is a regularization scheme, the RHS of the above display converges to 0 as $\lambda \rightarrow 0$ (see [Kress \(1999\)](#)). This concludes the proof. \square

2.1.3 Auxiliary Results

The following lemma is borrowed from [Florens et al. \(2012\)](#) (see their Lemma A.1(b)).

Lemma 2.3 *Under Assumptions 3.2 to 3.4 in the main text, we have*

$$\|\hat{A} - A\|_{op} = O_P\left(\frac{1}{\sqrt{nh}} + h^\rho\right) \text{ and } \|\hat{r} - r\| = O_P\left(\frac{1}{\sqrt{nh}} + h^\rho\right)$$

The following lemma is well established in the literature on empirical process theory. A proof can be found in [Andrews \(1994\)](#).

Lemma 2.4 *Let T be a random variable such that $\mathbb{E}[T|Z]$ is bounded and let \mathcal{F} be a class of functions of Z such that $\int_0^1 \sqrt{N_{[]}(\epsilon, \mathcal{F}, \|\cdot\|_{\mathbb{P}})} d\epsilon < \infty$. If $\|(\hat{f} - f_0)T\|_{2,P} = o_P(1)$ and $\mathbb{P}(\hat{f} \in \mathcal{F}) \rightarrow 1$, then*

$$\sqrt{n} (\mathbb{P}_n - \mathbb{P}) \left(\hat{f} - f_0 \right) T = o_P(1).$$

The lemma that follows gathers several useful results about inequalities of norms involving compact operators. Its proof can be found in [Florens et al. \(2011\)](#) and [Carrasco et al. \(2007\)](#).

Lemma 2.5 *Let \mathcal{X} and \mathcal{Y} be two Hilbert spaces and $K : \mathcal{X} \mapsto \mathcal{Y}$ be a linear compact operator with Singular Value Decomposition given by $(\tilde{\lambda}_j, \tilde{\varphi}_j, \tilde{\psi}_j)_j$.*

(a)

$$\|K(\lambda I + K^*K)^{-1}K^*\|_{op} \leq 1$$

(b)

$$\|\lambda(\lambda I + K^*K)^{-1}\|_{op} \leq 2$$

(c)

$$\|(\lambda I + K^*K)^{-1}K^*\|_{op} \leq \frac{1}{2\sqrt{\lambda}}$$

(d) If $\|\phi\|_\gamma^2 := \sum_j \tilde{\mu}_j^{-2\gamma} |\langle \phi, \tilde{\varphi}_j \rangle|^2 < \infty$ then

$$\|\lambda(\lambda I + K^*K)^{-1}\phi\| \leq C\|\phi\|_\gamma \lambda^{\frac{\gamma \wedge 2}{2}}$$

(e) Under the same conditions of (d) it holds that

$$\|\lambda K(\lambda I + K^*K)^{-1}\phi\| \leq C\|\phi\|_\gamma \lambda^{\frac{\gamma+1}{2} \wedge 1}$$

(f) If $\|\phi\|_\gamma^2 < \infty$ then

$$\|(\lambda I + K^*K)^{-1}K^*K\phi - \phi\| = O(\lambda^{\frac{\gamma \wedge 2}{2}})$$

Lemma 2.6 Under Assumptions 3.1 to 3.6 in the main text, we have

$$\|\hat{\varphi} - \tilde{\varphi}\|_\pi = O_P \left(\frac{\|\hat{A} - A\|}{\sqrt{\lambda}} + \frac{\|\hat{r} - r\|}{\sqrt{\lambda}} + \lambda^{\frac{\theta \wedge 2}{2}} \right)$$

where $\|\cdot\|$ denotes the $L^2(\mathbb{R})$ norm

Proof: The proof readily follows from the decomposition in (2.6) and Lemma 2.5. □

2.2 Power analysis in the nonparametric framework

We consider the scalar case $p = q = 1$. It is clear that our test will have asymptotic power equal to one against alternatives for which $\mathbb{E}[\tilde{U}(W - \mathbb{E}[W])X] \neq 0$. Remark that $\mathbb{E}[\tilde{U}(W - \mathbb{E}[W])X]$ can take in principle any value. So, we argue that the alternatives against which we do not have power are degenerate. Let us give some examples.

Example 3. Let E be uniformly distributed on the interval $[-1, 1]$. Let also X and W be Bernoulli random variables with parameters equal to $1/2$, respectively. The variables E, X, W are mutually independent. The treatment is generated as

$$Z = WX I(E \geq 0).$$

The residual $U(0)$ and $U(1)$ follow

$$U(0) = \rho X E;$$

$$U(1) = 0,$$

where $\rho > 0$. We assume that $\varphi(0) = \varphi(1) = 0$, so that $Y = U(Z)$. Notice that $Y = 0$ when $Z = 0$ and $Y = \rho X E$ when $Z = 1$. The mapping $\tilde{\varphi}$ solves

$$\mathbb{E}[Y|W] = \mathbb{E}[\tilde{\varphi}(Z)|W]. \quad (2.20)$$

Using the fact that, on the event $\{W = 0\}$, we have $Z = 0$, we get

$$\tilde{\varphi}(0) = \mathbb{E}[Y|W = 0] = 0.$$

Next, we have

$$\mathbb{E}[Y|W = 1] = \frac{1}{4} \mathbb{E}[Y|E \geq 0, X = 1, W = 1] = \frac{\rho}{8}. \quad (2.21)$$

Since $\tilde{\varphi}(0) = 0$ and $\mathbb{P}(Z = 1|W = 1) = \frac{1}{4}$, it also holds that $\mathbb{E}[\tilde{\varphi}(Z)|W = 1] = \frac{1}{4}\tilde{\varphi}(1)$. By (2.21), this leads to $\tilde{\varphi}(1) = -\frac{\rho}{2}$. As a result, we obtain $\tilde{U} = Y - \tilde{\varphi}(Z) = Z(\rho X E - \frac{\rho}{2})$. Hence, we have

$$\begin{aligned} \mathbb{E}[\tilde{U}WX] &= \frac{1}{4} \mathbb{E}[\tilde{U}|W = 1, X = 1] \\ &= \frac{1}{4} \mathbb{E} \left[Z \left(\rho X E - \frac{\rho}{2} \right) \middle| W = 1, X = 1 \right] \\ &= \frac{\rho}{8} (\mathbb{E}[E|E \geq 0, W = 1, X = 1] - 1) = -\frac{\rho}{16}. \end{aligned}$$

Therefore, in this example, our test does not have power only when $\rho = 0$, which is a degenerate case.

Example 4. Let $(W, E, X)^\top$ be a 3×1 Gaussian vector with mean zero and variance equal to the identity matrix. Let also $Z = W + E + X$, $U(Z) = Z(E + \rho X)$, where $\rho \in \mathbb{R}$, and $\varphi \equiv 0$ so that $Y = U(Z) = Z(E + \rho X)$. In this case, we have

$$\mathbb{E}[Y|W] = \mathbb{E}[Z(E + \rho X)|W] = \mathbb{E}[WE + E^2 + XE + \rho(WX + EX + X^2)|W] = 1 + \rho.$$

Hence, $\tilde{\varphi} \equiv 1 + \rho$ solves the equation. It is the unique solution since Z is strongly complete conditional on W (see [Newey and Powell \(2003\)](#) for a discussion of conditional completeness in the Gaussian case). As a result, we have $\tilde{U} = Z(E + \rho X) - 1 - \rho$. Hence, we get

$$\begin{aligned} \mathbb{E}[\tilde{U}WX] &= \mathbb{E}[(W + E + X)(E + \rho X)WX] - (1 + \rho)\mathbb{E}[WX] \\ &= \rho\mathbb{E}[(WX)^2] = \rho. \end{aligned}$$

As a result, the test does not have power only when X is not correlated with U , i.e. $\rho = 0$, which is a degenerate case.

References

- Andrews, D. W. K. (1994, January). Chapter 37 Empirical process methods in econometrics. In *Handbook of Econometrics*, Volume 4, pp. 2247–2294. Elsevier.
- Carrasco, M., J.-P. Florens, and E. Renault (2007, January). Chapter 77 Linear Inverse Problems in Structural Econometrics Estimation Based on Spectral Decomposition and Regularization. In J. J. Heckman and E. E. Leamer (Eds.), *Handbook of Econometrics*, Volume 6, pp. 5633–5751. Elsevier.
- Florens, J.-P., J. Johannes, and S. V. Bellegem (2012). Instrumental regression in partially linear models. *Econometrics Journal* 15(2), 304–324.
- Florens, J.-P., J. Johannes, and S. Van Bellegem (2011). Identification and estimation by penalization in a nonparametric instrumental regression. *Econometric Theory* 27, 472–496.
- Kress, R. (1999, December). *Linear Integral Equations*. Springer Science & Business Media.
- Newey, W. K. and J. L. Powell (2003). Instrumental variable estimation of nonparametric models. *Econometrica* 71, 1565–1578.

Van der Vaart, A. W. (1998). *Asymptotic Statistics*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge: Cambridge University Press.