

Hadronic molecules composed of a doubly charmed tetraquark state and a charmed meson

Ya-Wen Pan,¹ Tian-Wei Wu,^{2,3} Ming-Zhu Liu,^{4,1,*} and Li-Sheng Geng^{5,1,6,7,†}

¹*School of Physics, Beihang University, Beijing 100191, China*

²*School of Fundamental Physics and Mathematical Sciences,*

Hangzhou Institute for Advanced Study,

University of Chinese Academy of Sciences, Hangzhou 310024, China

³*University of Chinese Academy of Sciences, Beijing 100049, China*

⁴*School of Space and Environment, Beihang University, Beijing 100191, China*

⁵*Peng Huanwu Collaborative Center for Research and Education,*

Beihang University, Beijing 100191, China

⁶*Beijing Key Laboratory of Advanced Nuclear Materials and Physics,*

Beihang University, Beijing 100191, China

⁷*School of Physics and Microelectronics,*

Zhengzhou University, Zhengzhou, Henan 450001, China

(■Dated: August 11, 2022)

Abstract

The three pentaquark states, $P_c(4312)$, $P_c(4440)$ and $P_c(4457)$, discovered by the LHCb Collaboration in 2019, can be arranged into a complete heavy quark spin symmetry multiplet of hadronic molecules of $\bar{D}^{(*)}\Sigma_c^{(*)}$. In the heavy quark mass limit, the $\Sigma_c^{(*)}$ baryons can be related to the doubly charmed tetraquark states of isospin 1, i.e., $T_{\bar{c}\bar{c}}^{(*)}(T_{\bar{c}\bar{c}}^0, T_{\bar{c}\bar{c}}^1, T_{\bar{c}\bar{c}}^2)$, via heavy antiquark diquark symmetry, which dictates that the $\bar{D}^{(*)}\Sigma_c^{(*)}$ interactions are the same as the $\bar{D}^{(*)}T_{\bar{c}\bar{c}}^{(*)}$ interactions up to HADS breakings. In this work, we employ the contact-range effective field theory to systematically study the $\bar{D}^{(*)}T_{\bar{c}\bar{c}}^{(*)}$ systems, and we show the existence of a complete heavy quark spin symmetry multiplet of hadronic molecules composed of a doubly charmed tetraquark state and a charmed meson. These are a new kind of hadronic molecules and, if discovered, can lead to a better understanding of the many exotic hadrons discovered so far. In addition, we find that the $\bar{\Omega}_{ccc}p$ system can bind by the Coulomb force, which is analogous to a hydrogenlike atom.

* zhengmz11@buaa.edu.cn

† lisheng.geng@buaa.edu.cn

I. INTRODUCTION

A hadronic molecule is a bound or resonant state composed of conventional hadrons that is held together by the residual strong interaction, which is analogous to the deuteron that is bound by the nuclear force. The studies on hadronic molecules containing charm quarks began with the proposal that $\psi(4040)$ is a $\bar{D}^* D^*$ hadronic molecule [1, 2]. Although nowadays $\psi(4040)$ is widely viewed as a conventional charmonium state, the idea motivated a lot of theoretical studies on heavy hadronic molecules [3–6]. Such studies were revived after the discovery of $X(3872)$ by the Belle Collaboration in 2003 [7]. Treated as a $D\bar{D}^*$ bound state, the lower mass (than the conventional quark model prediction) and the isospin violation of the strong decays of $X(3872)$ can be easily understood, which implies that $X(3872)$ contains a large molecular component [8–10]. Since then a series of states beyond the naive quark model picture have been discovered and many of them can be regarded as molecular candidates. See Refs. [11–19] for some recent reviews.

Symmetry is a particularly important and useful concept in particle physics. The SU(3)-flavor symmetry has already achieved great successes, e.g., in classifying the ground-state mesons and baryons. Symmetry plays an important role in studies of hadronic molecules as well, such as SU(3)-flavor and heavy quark spin symmetry (HQSS). Assuming that $D_{s0}(2317)$ discovered by the BarBar Collaboration [20] is a DK molecule, it is natural to expect a D^*K molecule in terms of HQSS, corresponding to the exotic state $D_{s1}(2460)$ [21–24]. The $Z_c(3900)$ and $Z_{cs}(3985)$ states, discovered by the BESIII Collaboration [25, 26], are often interpreted as $\bar{D}D^*$ and $\bar{D}D_s^*/\bar{D}_sD^*$ molecules, and are related to each other by SU(3)-flavor symmetry [27–33]. Another example is the three hidden charm pentaquark states, $P_c(4312)$, $P_c(4440)$ and $P_c(4457)$, discovered by the LHCb Collaboration [34]. They can be arranged into a complete HQSS multiplet of $\bar{D}^{(*)}\Sigma_c^{(*)}$ hadronic molecules [35–41]. The existence of such multiplets of hadronic molecules not only enriches the hadron spectrum, but also can help verify the molecular nature of some exotic states [42, 43].

The heavy antiquark diquark symmetry(HADS) dictates that a pair of heavy quarks behaves like a heavy antiquark from the perspective of the color degree of freedom [44]. Therefore, the charmed antimesons are the same as the doubly charmed baryons. Via HADS, one can derive the following relation between the mass splitting of the doubly charmed baryon doublet and that of the charmed meson doublet, i.e., $m_{\Xi_{cc}^*} - m_{\Xi_{cc}} = \frac{3}{4}(m_{\bar{D}^*} - m_{\bar{D}})$ [45], which has been verified by a series of lattice QCD simulations [46–49]. In addition, the coupling of a doubly charmed baryon

to a pion can be related to that of a charmed meson to a pion via HADS, which is found to be consistent with the quark model prediction [50, 51]. Applying HADS to the $\Sigma_c^{(*)}$ baryons, it is natural to expect the existence of a set of compact doubly charmed tetraquark states [52], which have received renewed interests since the recent discovery of the doubly charmed tetraquark T_{cc} by the LHCb Collaboration [53, 54]. In Ref. [55], HADS has been utilized to predict the existence of a series of compact doubly charmed tetraquark states as well.

The HADS has also been utilized to study hadronic molecules. In Refs. [43, 56], the authors related the $D^{(*)}\Xi_{cc}^{(*)}$ and $\Xi_{cc}^{(*)}\Xi_{cc}^{(*)}$ systems with the $\bar{D}^{(*)}D^{(*)}$ system via HADS, and investigated the likely existence of hadronic molecules of $D^{(*)}\Xi_{cc}^{(*)}$ and $\Xi_{cc}^{(*)}\Xi_{cc}^{(*)}$. Given that there exists a complete multiplet of HQSS hadronic molecules of $\bar{D}^{(*)}\Sigma_c^{(*)}$, it is natural to expect the existence of $\Xi_{cc}^{(*)}\Sigma_c^{(*)}$ hadronic molecules by replacing $\bar{D}^{(*)}$ with $\Xi_{cc}^{(*)}$ via HADS [57, 58]. In this work, following Ref. [58], we relate the $\bar{D}^{(*)}T_{cc}^{(*)}$ system to the $\bar{D}^{(*)}\Sigma_c^{(*)}$ system via HADS as shown in Fig. 1, and then employ the contact-range effective field theory (EFT) to investigate the likely existence of $\bar{D}^{(*)}T_{cc}^{(*)}$ hadronic molecules. Such molecules composed of a charmed meson and a compact doubly charmed tetraquark state are a new kind of triply charmed hexaquark states in addition to the dibaryon states $\Xi_{cc}^{(*)}\Sigma_c^{(*)}$ [58] and three-body molecules $D^{(*)}D^{(*)}D^{(*)}$ [59, 60].




FIG. 1. From the $\bar{D}^{(*)}\Sigma_c^{(*)}$ system to the $\bar{D}^{(*)}T_{cc}^{(*)}$ system via HADS.

This paper is organized as follows. In Sec. II, we explain how to derive the contact potential for the $\bar{D}^{(*)}T_{cc}^{(*)}$ system constrained by HQSS, solve the Lippmann-Schwinger equation, and search for poles which correspond to $\bar{D}^{(*)}T_{cc}^{(*)}$ molecules. In Sec. III, we present the pole positions obtained with both single-channel and coupled-channel potentials. Finally, this paper is ended with a short summary in Sec. IV.

II. THEORETICAL FORMALISM

In this section, we explain how to derive the contact-range potential for the $\bar{D}^{(*)}T_{\bar{c}\bar{c}}^{(*)}$ system constrained by HQSS. The $\Sigma_c^{(*)}$ baryons are made of a charm quark and a pair of light quarks in the conventional quark model. HADS dictates that the charm quark of a $\Sigma_c^{(*)}$ baryon can be replaced with a pair of anti-charm quarks, and then it transforms to a doubly charmed tetraquark state as shown in Fig. 1, where the pair of anti-charm quarks is in the color $\bar{3}$ state of spin 1. Combining with the spin and color configurations of a pair of light quarks, the doubly charmed tetraquark states can have spin 0, 1, or 2, i.e.,

$$\begin{aligned} |T_{\bar{c}\bar{c}}(0^+)\rangle &= 1_H^3 \otimes 1_L^{\bar{3}}, \\ |T_{\bar{c}\bar{c}}(1^+)\rangle &= 1_H^3 \otimes 1_L^{\bar{3}}, \\ |T_{\bar{c}\bar{c}}(2^+)\rangle &= 1_H^3 \otimes 1_L^{\bar{3}}, \end{aligned} \tag{1}$$

where 1_H^3 and $1_L^{\bar{3}}$ represent an anti-charm quark pair and a light quark pair, respectively. In the following, we denote $T_{\bar{c}\bar{c}}(0^+)$, $T_{\bar{c}\bar{c}}(1^+)$, and $T_{\bar{c}\bar{c}}(2^+)$ as $T_{\bar{c}\bar{c}}^0$, $T_{\bar{c}\bar{c}}^1$, and $T_{\bar{c}\bar{c}}^2$, respectively. Since these states have not been discovered experimentally, we turn to the theoretical works [55, 61–63] for their masses, which are given in Table I. In the following numerical study, we use their averages given in the last column of Table I.

TABLE I. Masses (in units of MeV) of $T_{\bar{c}\bar{c}}^1$, $T_{\bar{c}\bar{c}}^2$, and $T_{\bar{c}\bar{c}}^3$ in several models and their averages.

Tetraquark	[55]	[61]	[62]	[63]	A.V
$T_{\bar{c}\bar{c}}^0$	3999.8	4132	4032	3969.2	4033.3
$T_{\bar{c}\bar{c}}^1$	4124.0	4151	4117	4053.2	4111.3
$T_{\bar{c}\bar{c}}^2$	4194.9	4185	4179	4123.8	4170.7

The HQSS dictates that the interactions between light quarks are independent of the spin of the heavy quark(s). In the following, we apply HQSS to derive the $\bar{D}^{(*)}T_{\bar{c}\bar{c}}^{(*)}$ contact-range potentials. First, we express the spin wave functions of the $\bar{D}^{(*)}T_{\bar{c}\bar{c}}^{(*)}$ pairs in terms of the spins of the heavy quarks s_{1H} and s_{2H} and those of the light quark(s) (often referred to as brown muck [64, 65]) s_{1L}

and s_{2L} , where 1 and 2 denote $\bar{D}^{(*)}$ and $T_{\bar{c}\bar{c}}^{(*)}$, respectively, via the following spin coupling formula,

$$|s_{1l}, s_{1h}, j_1; s_{2l}, s_{2h}, j_2; J\rangle = \sqrt{(2j_1 + 1)(2j_2 + 1)(2s_L + 1)(2s_H + 1)} \begin{pmatrix} s_{1l} & s_{2l} & s_L \\ s_{1h} & s_{2h} & s_H \\ j_1 & j_2 & J \end{pmatrix} |s_{1l}, s_{2l}, s_L; s_{1h}, s_{2h}, s_H; J\rangle, \quad (2)$$

where j_1 and j_2 represent the spin of $\bar{D}^{(*)}$ and $T_{\bar{c}\bar{c}}^{(*)}$, s_L and s_H represent the total spin of the light quarks and heavy quarks, and J represent the total angular momentum of the $\bar{D}^{(*)}T_{\bar{c}\bar{c}}^{(*)}$ system. Ten $\bar{D}^{(*)}T_{\bar{c}\bar{c}}^{(*)}$ states can be decomposed into the light quark basis S_L and heavy quark basis S_H , which are explicitly given in Appendix A.

In the heavy quark mass limit, the $\bar{D}^{(*)}T_{\bar{c}\bar{c}}^{(*)}$ potentials can be parameterized by two coupling constants describing the interactions between light quarks of spin 1/2 and 3/2, respectively, i.e., $F_{1/2} = \langle 1/2_L | V | 1/2_L \rangle$ and $F_{3/2} = \langle 3/2_L | V | 3/2_L \rangle$, which are explicitly given in Appendix A. The $\bar{D}^{(*)}T_{\bar{c}\bar{c}}^{(*)}$ potentials can be rewritten as a combination of C_a and C_b , i.e. $F_{1/2} = C_a - 2C_b$ and $F_{3/2} = C_a + C_b$ [66]. For the $T_{\bar{c}\bar{c}}^0 \bar{D} - T_{\bar{c}\bar{c}}^1 \bar{D}^*$ coupled channels, the contact-range potentials V in matrix form read

$$V_{T_{\bar{c}\bar{c}}^0 \bar{D} - T_{\bar{c}\bar{c}}^1 \bar{D}^*}^{J=0} = \begin{pmatrix} C_a & -\sqrt{2}C_b \\ -\sqrt{2}C_b & C_a - C_b \end{pmatrix}. \quad (3)$$

For the $T_{\bar{c}\bar{c}}^1 \bar{D} - T_{\bar{c}\bar{c}}^0 \bar{D}^* - T_{\bar{c}\bar{c}}^1 \bar{D}^* - T_{\bar{c}\bar{c}}^2 \bar{D}^*$ coupled channels, the contact-range potentials V in matrix form read

$$V_{T_{\bar{c}\bar{c}}^1 \bar{D} - T_{\bar{c}\bar{c}}^0 \bar{D}^* - T_{\bar{c}\bar{c}}^1 \bar{D}^* - T_{\bar{c}\bar{c}}^2 \bar{D}^*}^{J=1} = \begin{pmatrix} C_a & \sqrt{\frac{2}{3}}C_b & -\frac{1}{\sqrt{2}}C_b & -\sqrt{\frac{5}{6}}C_b \\ \sqrt{\frac{2}{3}}C_b & C_a & \frac{2}{\sqrt{3}}C_b & 0 \\ -\frac{1}{\sqrt{2}}C_b & \frac{2}{\sqrt{3}}C_b & C_a - \frac{1}{2}C_b & \frac{1}{2}\sqrt{\frac{5}{3}}C_b \\ -\sqrt{\frac{5}{6}}C_b & 0 & \frac{1}{2}\sqrt{\frac{5}{3}}C_b & C_a - \frac{3}{2}C_b \end{pmatrix}. \quad (4)$$

For the $T_{\bar{c}\bar{c}}^2 \bar{D} - T_{\bar{c}\bar{c}}^1 \bar{D}^* - T_{\bar{c}\bar{c}}^2 \bar{D}^*$ coupled channels, the contact-range potentials V in matrix form read

$$V_{T_{\bar{c}\bar{c}}^2 \bar{D} - T_{\bar{c}\bar{c}}^1 \bar{D}^* - T_{\bar{c}\bar{c}}^2 \bar{D}^*}^{J=2} = \begin{pmatrix} C_a & \frac{1}{\sqrt{2}}C_b & -\frac{3}{\sqrt{6}}C_b \\ \frac{1}{\sqrt{2}}C_b & C_a + \frac{1}{2}C_b & \frac{\sqrt{3}}{2}C_b \\ -\frac{3}{\sqrt{6}}C_b & \frac{\sqrt{3}}{2}C_b & C_a - \frac{1}{2}C_b \end{pmatrix}. \quad (5)$$

For the $T_{cc}^2 \bar{D}^*$ single channel, the contact-range potential is $V_{T_{cc}^2 \bar{D}^*}^{J=3} = C_a + C_b$.

With the above potentials, we can solve the Lippmann-Schwinger equation

$$T = (1 - VG)^{-1}V, \quad (6)$$

where V is the coupled-channel potential determined by the contact-range EFT approach described above, and G is the two-body propagator. In evaluating the loop function G , we introduce a regulator of Gaussian form e^{-2q^2/Λ^2} in the integral as

$$G(s) = \int \frac{d^3q}{(2\pi)^3} \frac{e^{-2q^2/\Lambda^2}}{\sqrt{s} - m_1 - m_2 - q^2/(2\mu_{12}) + i\varepsilon}, \quad (7)$$

where \sqrt{s} is the total energy in the center-of-mass (c.m.) frame of m_1 and m_2 , $\mu_{12} = \frac{m_1 m_2}{m_1 + m_2}$ is the reduced mass, and Λ is the momentum cutoff. Following our previous works [66, 67], we take $\Lambda = 0.75$ GeV and $\Lambda = 1.5$ GeV in the present work, which reflects the internal structure of the involved hadrons and provides an estimate of the uncertainties of the numerical results. The dynamically generated states correspond to poles in the unphysical sheet. In this sheet, the loop function of Eq. (7) becomes

$$G^{II}(s, m_1, m_2) = G(s, m_1, m_2) + i \frac{4m_1 m_2}{8\pi} \frac{p}{\sqrt{s}} e^{-2p^2/\Lambda^2}, \quad (8)$$

where the c.m. momentum p is

$$p = \frac{\sqrt{s - (m_1 + m_2)^2} \sqrt{s - (m_1 - m_2)^2}}{2\sqrt{s}}. \quad (9)$$

III. RESULTS AND DISCUSSIONS

Following our previous works [58, 66], the parameters C_a and C_b of the contact-range potentials are determined by reproducing the masses of $P_c(4440)$ and $P_c(4457)$ in two scenarios. In Scenario A, $P_c(4440)$ and $P_c(4457)$ have $J^P = 1/2^-$ and $J^P = 3/2^-$, while in Scenario B the assignment interchanges. In Table II, we present the values of C_a and C_b obtained in Scenario A and B with $\Lambda = 1.5$ GeV and $\Lambda = 0.75$ GeV. One should note that the uncertainty of HADS is expected to be of the order of $\Lambda_{QCD}/(m_Q v)$ [44], where v is the velocity of the heavy quark pair. In general, the QCD scale is taken to be $\Lambda_{QCD} \sim 200 - 300$ MeV and $m_Q v$ is estimated to be 800 MeV for charm quarks. The breaking of HADS is then estimated to be 25 – 40% [58]. The mass splitting between the $J = 1/2$ and $J = 3/2$ doubly charmed baryons obtained in lattice QCD simulations

TABLE II. Couplings C_a and C_b (in units of GeV^{-2}) in Scenario A and B obtained with $\Lambda = 0.75 \text{ GeV}$ and $\Lambda = 1.50 \text{ GeV}$.

Scenario	Λ (GeV)	C_a	C_b	Λ (GeV)	C_a	C_b
A	1.50	-12.18	1.07	0.75	-30.80	5.10
B	1.50	-12.89	-1.07	0.75	-34.20	-5.10

TABLE III. Binding energies (in units of MeV) of the $\bar{D}^{(*)}T_{cc}^{(*)}$ molecules in Scenario A and Scenario B obtained with single-channel potentials. The numbers inside and outside the brackets correspond to $\Lambda = 0.75 \text{ GeV}$ and $\Lambda = 1.5 \text{ GeV}$. The superscripts and subscripts are obtained by allowing for a 25% breaking of HADS.

Molecule	J^P	Threshold	B.E.(Scenario A)	B.E.(Scenario B)
$\bar{D}T_{cc}^0$	0^-	5900.3	$24.1^{+39.4}_{-22.5}(16.5^{+16.1}_{-12.1})$	$32.1^{+44.9}_{-28.3}(23.3^{+19.3}_{-15.4})$
$\bar{D}T_{cc}^1$	1^-	5978.3	$24.7^{+39.7}_{-23.0}(16.8^{+16.2}_{-12.2})$	$32.8^{+45.3}_{-28.8}(23.6^{+19.4}_{-15.5})$
$\bar{D}T_{cc}^2$	2^-	6037.7	$25.2^{+40.0}_{-23.3}(17.0^{+16.2}_{-12.3})$	$33.4^{+45.5}_{-29.1}(23.8^{+19.4}_{-15.6})$
$\bar{D}^*T_{cc}^0$	1^-	6042.3	$29.5^{+42.1}_{-26.2}(18.6^{+16.7}_{-13.0})$	$38.2^{+47.6}_{-32.0}(25.6^{+19.9}_{-16.3})$
$\bar{D}^*T_{cc}^1$	0^-	6120.3	$43.7^{+50.7}_{-35.3}(29.7^{+21.6}_{-18.0})$	$26.1^{+39.6}_{-23.8}(15.6^{+15.2}_{-11.4})$
$\bar{D}^*T_{cc}^1$	1^-	6120.3	$36.8^{+46.6}_{-31.0}(24.1^{+19.2}_{-15.6})$	$32.4^{+43.8}_{-28.1}(20.6^{+17.6}_{-13.9})$
$\bar{D}^*T_{cc}^1$	2^-	6120.3	$24.2^{+38.2}_{-22.3}(14.0^{+14.4}_{-10.6})$	$46.1^{+52.1}_{-36.7}(31.6^{+22.4}_{-18.8})$
$\bar{D}^*T_{cc}^2$	1^-	6179.7	$51.7^{+55.0}_{-39.8}(35.8^{+24.0}_{-20.5})$	$20.9^{+35.7}_{-19.8}(11.2^{+12.8}_{-9.0})$
$\bar{D}^*T_{cc}^2$	2^-	6179.7	$37.4^{+46.8}_{-31.3}(24.3^{+19.3}_{-15.6})$	$32.9^{+44.0}_{-28.5}(20.8^{+17.7}_{-14.0})$
$\bar{D}^*T_{cc}^2$	3^-	6179.7	$19.0^{+34.3}_{-18.3}(9.8^{+12.0}_{-8.1})$	$54.3^{+56.4}_{-41.3}(37.8^{+24.8}_{-21.3})$

are 20–25% smaller than the HADS prediction [46–49]. Therefore, we settle on a 25% uncertainty for the $\bar{D}^{(*)}T_{cc}^{(*)}$ potentials.

First, we search for poles of $\bar{D}^{(*)}T_{cc}^{(*)}$ with single-channel potentials. In Table III, we present the poles found in Scenario A and Scenario B, where the values inside and outside the brackets correspond to the cutoff $\Lambda = 0.75 \text{ GeV}$ and $\Lambda = 1.5 \text{ GeV}$, respectively. One can see that our results are weakly dependent on the cutoff. Even taking into account the 25% uncertainty of HADS, all

of these states are below the $\bar{D}^{(*)}T_{cc}^{(*)}$ mass thresholds. Obviously, we obtain a complete HQSS multiplet of hadronic molecules composed of a $\bar{D}^{(*)}$ meson and a $T_{cc}^{(*)}$ tetraquark state. In other words, the existence of $\bar{D}^{(*)}\Sigma_c^{(*)}$ hadronic molecules implies the existence of $\bar{D}^{(*)}T_{cc}^{(*)}$ molecules, if HQSS is approximately satisfied. We note in passing that the existence of $\bar{D}^{(*)}\Sigma_c^{(*)}$ molecules also implies the existence of a complete HQSS multiplet of $\Xi_{cc}^{(*)}\Sigma_c^{(*)}$ hadronic molecules in terms of HADS [58], which agrees with the lattice QCD simulations [68].

TABLE IV. Pole positions relative to the corresponding two-body thresholds (in units of MeV), $M - i\Gamma/2$, for the $\bar{D}^{(*)}T_{cc}^{(*)}$ molecules in Scenario A and Scenario B obtained with coupled-channel potentials. The numbers inside and outside brackets correspond to $\Lambda = 0.75$ GeV and $\Lambda = 1.5$ GeV. The superscripts and subscripts represent corrections induced by a 25% breaking of HADS.

Molecule	J^P	Threshold	$M_{threshold} - M(\text{Scenario A})$	$M_{threshold} - M(\text{Scenario B})$
$\bar{D}T_{cc}^0$	0^-	5900.3	$26.3_{-24.4}^{+42.6}(17.9_{-13.0}^{+17.4})$	$34.2_{-30.0}^{+47.4}(24.6_{-16.2}^{+20.4})$
$\bar{D}T_{cc}^1$	1^-	5978.3	$29.0_{-26.7}^{+46.4}(19.4_{-14.0}^{+18.9})$	$36.4_{-31.6}^{+49.6}(25.9_{-17.0}^{+21.3})$
$\bar{D}T_{cc}^2$	2^-	6037.7	$29.2_{-26.7}^{+46.0}(19.4_{-13.9}^{+18.7})$	$36.8_{-31.8}^{+49.6}(26.0_{-17.0}^{+21.3})$
$\bar{D}^*T_{cc}^0$	1^-	6042.3	$31.7_{-28.0}^{+43.4} + 1.2_{-0.8}^{-1.2}i(19.7_{-13.7}^{+17.8} + 1.0_{-0.5}^{+0.3}i)$	$40.9_{-34.0}^{+51.6} + 0.5_{-0.2}^{-0.5}i(27.2_{-17.3}^{+21.4} + 0.5_{-0.1}^{-0.1}i)$
$\bar{D}^*T_{cc}^1$	0^-	6120.3	$42.0_{-34.0}^{+48.5} + 2.0_{-1.3}^{+1.0}i(28.0_{-17.1}^{+20.4} + 0.9_{-0.5}^{+0.6}i)$	$24.8_{-22.7}^{+37.9} + 1.5_{-1.1}^{+0.9}i(14.3_{-10.6}^{+14.1} + 0.6_{-0.4}^{+0.4}i)$
$\bar{D}^*T_{cc}^1$	1^-	6120.3	$36.5_{-30.8}^{+46.4} + 1.8_{-0.9}^{-1.8}i(23.3_{-15.1}^{+19.0} + 1.4_{-0.6}^{+0.3}i)$	$30.9_{-26.8}^{+41.2} + 2.4_{-1.5}^{-2.0}i(19.0_{-12.9}^{+16.2} + 1.7_{-0.9}^{+0.7}i)$
$\bar{D}^*T_{cc}^1$	2^-	6120.3	$25.7_{-23.6}^{+40.4} + 1.1_{-0.9}^{+0.2}i(14.7_{-11.1}^{+15.1} + 0.8_{-0.5}^{+0.5}i)$	$48.1_{-38.1}^{+53.9} + 0.3_{-0.1}^{-0.3}i(32.8_{-19.5}^{+23.4} + 0.3_{-0.1}^{-0.0}i)$
$\bar{D}^*T_{cc}^2$	1^-	6179.7	$49.8_{-38.2}^{+49.5} + 1.6_{-0.8}^{+0.3}i(34.1_{-19.5}^{+22.5} + 1.2_{-0.5}^{+0.1}i)$	$19.8_{-18.8}^{+34.4} + 0.9_{-0.7}^{+0.1}i(10.2_{-8.3}^{+12.0} + 0.5_{-0.3}^{+0.3}i)$
$\bar{D}^*T_{cc}^2$	2^-	6179.7	$25.3_{-28.9}^{+41.6} + 2.4_{-1.4}^{-1.0}i(21.7_{-14.0}^{+17.1} + 1.7_{-0.8}^{+0.5}i)$	$30.1_{-26.2}^{+39.6} + 2.4_{-1.5}^{-0.2}i(18.3_{-12.4}^{+15.6} + 1.7_{-0.8}^{+0.6}i)$
$\bar{D}^*T_{cc}^2$	3^-	6179.7	$19.0_{-18.3}^{+34.3}(9.8_{-8.1}^{+12.0})$	$54.3_{-41.3}^{+56.4}(37.8_{-21.3}^{+24.8})$

Next we take into account the inelastic potentials, and search for poles with coupled-channel potentials. We present the results in Table IV, where some poles are located in the complex plane instead of in the real axis due to the contribution of the inelastic channels. We find that the resulting widths are rather small, similar to the $\bar{D}^{(*)}\Sigma_c^{(*)}$ molecules [36]. We stress that the number of $\bar{D}^{(*)}T_{cc}^{(*)}$ molecules remain unchanged after including the inelastic potentials, and the masses of the $\bar{D}^{(*)}T_{cc}^{(*)}$ molecules change little compared with the results shown in Table III. That is to

say, even with coupled-channel potentials, we still obtain a complete HQSS multiplet of $\bar{D}^{(*)}T_{\bar{c}\bar{c}}^{(*)}$ hadronic molecules. One should note that as we reduce the strength of the $\bar{D}^{(*)}T_{\bar{c}\bar{c}}^{(*)}$ potential, all the poles are still located in the same Riemann sheet. However, if we increase the strength of the potentials, some poles move to different Riemann sheets, leading to two poles located in the same Riemann sheet.




FIG. 2. Three possible configurations of triply charmed hexaquark states with the same quark contents: three charmed mesons $\bar{D}^{(*)}\bar{D}^{(*)}\bar{D}^{(*)}$ (a), a charmed meson and a doubly charmed tetraquark $\bar{D}^{(*)}T_{\bar{c}\bar{c}}^{(*)}$ (b), and a triply charmed anti-baryon and a proton $\bar{\Omega}_{ccc}p$ (c).

The $\bar{D}^{(*)}T_{\bar{c}\bar{c}}^{(*)}$ molecules are a new kind of hadronic molecules, which are composed of a conventional meson and a compact tetraquark state. The parity and quark content of the $\bar{D}^{(*)}T_{\bar{c}\bar{c}}^{(*)}$ molecules are the same as those of the three-body hadronic molecules of $\bar{D}^{(*)}\bar{D}^{(*)}\bar{D}^{(*)}$ as well as the two-body $\bar{\Omega}_{ccc}p$ system as shown in Fig. 2. The existence of $D^{(*)}D^{(*)}D^{(*)}$ depend on the molecular nature of T_{cc} recently discovered by the LHCb Collaboration [59, 60]. Similarly, the existence of the $\bar{D}^{(*)}T_{\bar{c}\bar{c}}^{(*)}$ molecules depend on the molecular nature of the three pentaquark states, $P_c(4312)$, $P_c(4440)$, and $P_c(4457)$, discovered by the LHCb Collaboration in 2019. It is important to note that the configurations of (a) and (b) can mix with each other while they both can decay into configuration (c) because of their respective masses. For the $\bar{\Omega}_{ccc}p$ system, it is quite difficult to bind by the strong interaction due to the Okubo-Zweig-Iizuka (OZI) rule, while it can bind with the Coulomb force as shown below.

The electromagnetic interaction of the $\bar{\Omega}_{ccc}p$ system reads

$$V_{\bar{\Omega}_{ccc}p}^C = -2\frac{\alpha}{r}, \quad (10)$$

where $\alpha = 1/137$ is the fine structure constant. Since the Ω_{ccc} baryon has not been observed so far, we take the mass $m_{\Omega_{ccc}} = 4795.6$ MeV predicted by lattice QCD [69]. Solving the Schrödinger equation via the Gaussian expansion approach, we find that there exists a shallow $\bar{\Omega}_{ccc}p$ bound state with a binding energy of 83.6 keV for the orbit angular momentum $l = 0$. Recall for a Hydrogenlike atom, the energy level is given by

$$E_n = -\frac{\mu e^4 Z^2}{2\hbar n^2}, \quad (11)$$

where μ is the reduced mass of the hydrogen, e is the electron charge, and Z is the charge number. Applying the above formula to the $\bar{\Omega}_{ccc}p$ system, one obtains the binding energy 83.6 keV for the ground state, in agreement with our numerical result. In Fig. 3, we present the energy spectrum of the $\bar{\Omega}_{ccc}p$ system, which is typical of a two-body system bound by the electromagnetic force. We note that the $\bar{\Omega}_{ccc}p$ molecule is similar to the $D^\pm D^{*\mp}$ hadronic atom of Ref. [70] but differs in that the strong interaction between $\bar{\Omega}_{ccc}$ and p is more reduced because of the OZI rule.




FIG. 3. Energy spectrum of the $\bar{\Omega}_{ccc}p$ molecule for the orbit angular momentum $l = 0$.

According to HADS, the existence of $\bar{D}^{(*)}\Sigma_c^{(*)}$ hadronic molecules indicates the existence of $\Xi_{cc}^{(*)}\Sigma_c^{(*)}$ hadronic molecules [58], which are also triply charmed hexaquark states. The quark contents of the $\Xi_{cc}^{(*)}\Sigma_c^{(*)}$ molecules are the same as those of the $\bar{\Omega}_{ccc}p$ system, while the latter can

not bind due to the repulsive Coulomb force and the OZI rule. As a result, the triply charmed hexaquark states in the molecular picture can be classified into two groups: $\Xi_{cc}^{(*)}\Sigma_c^{(*)}$ molecules with baryon number $B = 2$ and $(\bar{D}^{(*)}\bar{D}^{(*)}\bar{D}^{(*)}, \bar{D}^{(*)}T_{cc}^{(*)}, \bar{\Omega}_{ccc}p)$ molecules with baryon number zero. In addition, we find that the $\bar{D}^{(*)}\bar{D}^{(*)}\bar{D}^{(*)}$ and $\bar{D}^{(*)}T_{cc}^{(*)}$ molecules are significantly different from the $\bar{\Omega}_{ccc}p$ molecule due to their different binding mechanisms. As for the $\bar{D}^{(*)}\bar{D}^{(*)}\bar{D}^{(*)}$ and $\bar{D}^{(*)}T_{cc}^{(*)}$ molecules, the number of $\bar{D}^{(*)}\bar{D}^{(*)}\bar{D}^{(*)}$ molecules are less than that of $\bar{D}^{(*)}T_{cc}^{(*)}$ molecules due to the constraint of Bose-Einstein statistics. In addition, the $\bar{D}^{(*)}T_{cc}^{(*)}$ molecules can decay into three charmed mesons [52], while the $\bar{D}^{(*)}\bar{D}^{(*)}\bar{D}^{(*)}$ molecules are more likely to decay into three charmed mesons together with a π or a γ [59], which can be used to discriminate the $\bar{D}^{(*)}T_{cc}^{(*)}$ and $\bar{D}^{(*)}\bar{D}^{(*)}\bar{D}^{(*)}$ molecules.

IV. SUMMARY AND CONCLUSION

The pentaquark states, $P_c(4312)$, $P_c(4440)$, and $P_c(4457)$, discovered by the LHCb Collaboration, have been suggested to be $\bar{D}^{(*)}\Sigma_c$ hadronic molecules. Together with the other four states dictated by HQSS, a complete HQSS multiplet of hadronic molecules may exist. HADS allows one to relate the $\bar{D}^{(*)}T_{cc}^{(*)}$ system to the $\bar{D}^{(*)}\Sigma_c^{(*)}$ system. The $T_{cc}^{(*)}$ are isovector doubly charmed compact tetraquark states, which have been extensively studied after the discovery of T_{cc}^+ by the LHCb Collaboration. Since the masses of the isovector tetraquark states $T_{cc}^{(*)}$ are unknown, we took the average masses estimated by several theoretical models, which are the main uncertainty of our present work.

In terms of HADS, the $\bar{D}^{(*)}T_{cc}^{(*)}$ potentials are the same as the $\bar{D}^{(*)}\Sigma_c^{(*)}$ potentials. We determined the unknown parameters of the $\bar{D}^{(*)}T_{cc}^{(*)}$ potentials by reproducing the masses of $P_c(4440)$ and $P_c(4457)$ with two spin assignments. We then employed the Lippmann-Schwinger equation to search for poles near the mass thresholds of $\bar{D}^{(*)}T_{cc}^{(*)}$ with single-channel and coupled-channel potentials, respectively. We find that, even taking into account a 25% breaking of HADS, there exists a complete HQSS multiplet of $\bar{D}^{(*)}T_{cc}^{(*)}$ hadronic molecules composed of a charmed meson and a doubly charmed compact tetraquark state. Such states are a new kind of hadronic molecules, which, if discovered, will further enlarge the family of hadronic molecules. At last, we predicted the existence of a different kind of triply charmed hadronic molecule of $\bar{\Omega}_{ccc}p$ held together solely by the Coulomb force. It is interesting to note that the same mechanism could lead to the existence of other hadronic atoms as well, such as $\Omega_{ccc}\Omega_{sss}$, $\Omega_{bbb}p$, and $\Omega_{bbb}\Omega_{ccc}$.

V. ACKNOWLEDGMENTS

This work is supported in part by the National Natural Science Foundation of China under Grants No.11975041, No.11735003, and No.11961141004. Ming-Zhu Liu acknowledges support from the National Natural Science Foundation of China under Grant No.1210050997 and China Postdoctoral Science Foundation under Grants No. 2022M710317, and No. 2022T150036. Tian-Wei Wu acknowledges support from the National Natural Science Foundation of China under Grant No.12147152.

Appendix A: Contact-range potentials for the $T_{cc}^{(*)}\bar{D}^{(*)}$ system

The $T_{cc}^{(*)}\bar{D}^{(*)}$ states with spin $J = 0, 1, 2, 3$ are expressed in terms of the total spin of the light quark and heavy quark pairs as follows:

$$\begin{aligned}
|T_{cc}^0\bar{D}(0^-)\rangle &= \frac{1}{\sqrt{3}}1/2_H \otimes 1/2_L + \sqrt{\frac{2}{3}}3/2_H \otimes 3/2_L, \\
|T_{cc}^1\bar{D}(1^-)\rangle &= \frac{\sqrt{2}}{3}1/2_H \otimes 1/2_L - \frac{1}{3}1/2_H \otimes 3/2_L + \frac{1}{3}3/2_H \otimes 1/2_L + \frac{\sqrt{5}}{3}3/2_H \otimes 3/2_L, \\
|T_{cc}^2\bar{D}(2^-)\rangle &= \frac{1}{\sqrt{3}}3/2_H \otimes 1/2_L - \frac{1}{\sqrt{3}}1/2_H \otimes 3/2_L + \frac{1}{\sqrt{3}}3/2_H \otimes 3/2_L, \\
|T_{cc}^0\bar{D}^*(1^-)\rangle &= -\frac{1}{3\sqrt{3}}1/2_H \otimes 1/2_L + \frac{2\sqrt{\frac{2}{3}}}{3}1/2_H \otimes 3/2_L - \frac{2\sqrt{\frac{2}{3}}}{3}3/2_H \otimes 1/2_L + \frac{\sqrt{\frac{10}{3}}}{3}3/2_H \otimes 3/2_L, \\
|T_{cc}^1\bar{D}^*(0^-)\rangle &= \sqrt{\frac{2}{3}}1/2_H \otimes 1/2_L - \frac{1}{\sqrt{3}}3/2_H \otimes 3/2_L, \\
|T_{cc}^1\bar{D}^*(1^-)\rangle &= \frac{1}{\sqrt{2}}1/2_H \otimes 3/2_L + \frac{1}{\sqrt{2}}3/2_H \otimes 1/2_L, \\
|T_{cc}^1\bar{D}^*(2^-)\rangle &= -\frac{1}{\sqrt{6}}3/2_H \otimes 1/2_L + \frac{1}{\sqrt{6}}1/2_H \otimes 3/2_L + \sqrt{\frac{2}{3}}3/2_H \otimes 3/2_L, \\
|T_{cc}^2\bar{D}^*(1^-)\rangle &= \frac{2\sqrt{\frac{5}{3}}}{3}1/2_H \otimes 1/2_L + \frac{\sqrt{\frac{5}{6}}}{3}1/2_H \otimes 3/2_L - \frac{\sqrt{\frac{5}{6}}}{3}3/2_H \otimes 1/2_L - \frac{\sqrt{\frac{2}{3}}}{3}3/2_H \otimes 3/2_L, \\
|T_{cc}^2\bar{D}^*(2^-)\rangle &= \frac{1}{\sqrt{2}}1/2_H \otimes 3/2_L + \frac{1}{\sqrt{2}}3/2_H \otimes 1/2_L, \\
|T_{cc}^2\bar{D}^*(3^-)\rangle &= 3/2_H \otimes 3/2_L.
\end{aligned} \tag{A1}$$

In the heavy quark limit, the interactions are only dependent on the degree of freedom of light quarks. As a result, we can write the matrix elements of the potentials as $F_{1/2} = \langle 1/2_L | V | 1/2_L \rangle$ and $F_{3/2} = \langle 3/2_L | V | 3/2_L \rangle$, then the potentials for the $T_{cc}^{(*)}\bar{D}^{(*)}$ system can be expressed in terms

of two parameters, e.g., $F_{1/2}$ and $F_{3/2}$. For the $T_{cc}^0\bar{D} - T_{cc}^1\bar{D}^*$ coupled channels, the contact-range potentials V in matrix form read

$$V_{T_{cc}^0\bar{D}-T_{cc}^1\bar{D}^*}^{J=0} = \begin{pmatrix} \frac{1}{3}F_{1/2} + \frac{2}{3}F_{3/2} & \frac{\sqrt{2}}{3}(F_{1/2} - F_{3/2}) \\ \frac{\sqrt{2}}{3}(F_{1/2} - F_{3/2}) & \frac{2}{3}F_{1/2} + \frac{1}{3}F_{3/2} \end{pmatrix} \quad (\text{A2})$$

For the $T_{cc}^1\bar{D} - T_{cc}^0\bar{D}^* - T_{cc}^1\bar{D}^* - T_{cc}^2\bar{D}^*$ coupled channels, the contact-range potentials V in matrix form read $V_{T_{cc}^1\bar{D}-T_{cc}^0\bar{D}^*-T_{cc}^1\bar{D}^*-T_{cc}^2\bar{D}^*}^{J=1} =$

$$\begin{pmatrix} \frac{1}{3}F_{1/2} + \frac{2}{3}F_{3/2} & -\frac{1}{3}\sqrt{\frac{2}{3}}(F_{1/2} - F_{3/2}) & \frac{1}{3\sqrt{2}}(F_{1/2} - F_{3/2}) & \frac{1}{3}\sqrt{\frac{5}{6}}(F_{1/2} - F_{3/2}) \\ -\frac{1}{3}\sqrt{\frac{2}{3}}(F_{1/2} - F_{3/2}) & \frac{1}{3}F_{1/2} + \frac{2}{3}F_{3/2} & -\frac{2}{3\sqrt{3}}(F_{1/2} - F_{3/2}) & 0 \\ \frac{1}{3\sqrt{2}}(F_{1/2} - F_{3/2}) & -\frac{2}{3\sqrt{3}}(F_{1/2} - F_{3/2}) & \frac{1}{2}F_{1/2} + \frac{1}{2}F_{3/2} & -\frac{1}{6}\sqrt{\frac{5}{3}}(F_{1/2} - F_{3/2}) \\ \frac{1}{3}\sqrt{\frac{5}{6}}(F_{1/2} - F_{3/2}) & 0 & -\frac{1}{6}\sqrt{\frac{5}{3}}(F_{1/2} - F_{3/2}) & \frac{5}{6}F_{1/2} + \frac{1}{6}F_{3/2} \end{pmatrix}. \quad (\text{A3})$$

For the $T_{cc}^2\bar{D} - T_{cc}^1\bar{D}^* - T_{cc}^2\bar{D}^*$ coupled channels, the contact-range potentials V in matrix form read

$$V_{T_{cc}^2\bar{D}-T_{cc}^1\bar{D}^*-T_{cc}^2\bar{D}^*}^{J=2} = \begin{pmatrix} \frac{1}{3}F_{1/2} + \frac{2}{3}F_{3/2} & -\frac{1}{3\sqrt{2}}(F_{1/2} - F_{3/2}) & \frac{1}{\sqrt{6}}(F_{1/2} - F_{3/2}) \\ -\frac{1}{3\sqrt{2}}(F_{1/2} - F_{3/2}) & \frac{1}{6}F_{1/2} + \frac{5}{6}F_{3/2} & -\frac{1}{2\sqrt{3}}(F_{1/2} - F_{3/2}) \\ \frac{1}{\sqrt{6}}(F_{1/2} - F_{3/2}) & -\frac{1}{2\sqrt{3}}(F_{1/2} - F_{3/2}) & \frac{1}{2}F_{1/2} + \frac{1}{2}F_{3/2} \end{pmatrix}. \quad (\text{A4})$$

For the $T_{cc}^3\bar{D}^*$ single channel, the contact-range potential is $V_{T_{cc}^3\bar{D}^*}^{J=3} = F_{3/2}$.

-
- [1] A. De Rujula, H. Georgi, and S. L. Glashow, Phys. Rev. Lett. **38**, 317 (1977).
 - [2] M. B. Voloshin and L. B. Okun, JETP Lett. **23**, 333 (1976).
 - [3] N. A. Tornqvist, Phys. Rev. Lett. **67**, 556 (1991).
 - [4] N. A. Tornqvist, Z. Phys. **C61**, 525 (1994), arXiv:hep-ph/9310247 [hep-ph].
 - [5] A. V. Manohar and M. B. Wise, Nucl. Phys. B **399**, 17 (1993), arXiv:hep-ph/9212236.
 - [6] T. E. O. Ericson and G. Karl, Phys. Lett. B **309**, 426 (1993).
 - [7] S. K. Choi et al. (Belle), Phys. Rev. Lett. **91**, 262001 (2003), arXiv:hep-ex/0309032.
 - [8] D. Gamermann and E. Oset, Phys. Rev. **D80**, 014003 (2009), arXiv:0905.0402 [hep-ph].
 - [9] N. Li and S.-L. Zhu, Phys. Rev. D **86**, 074022 (2012), arXiv:1207.3954 [hep-ph].

- [10] Q. Wu, D.-Y. Chen, and T. Matsuki, *Eur. Phys. J. C* **81**, 193 (2021), arXiv:2102.08637 [hep-ph].
- [11] H.-X. Chen, W. Chen, X. Liu, and S.-L. Zhu, *Phys. Rept.* **639**, 1 (2016), arXiv:1601.02092 [hep-ph].
- [12] A. Hosaka, T. Hyodo, K. Sudoh, Y. Yamaguchi, and S. Yasui, *Prog. Part. Nucl. Phys.* **96**, 88 (2017), arXiv:1606.08685 [hep-ph].
- [13] R. F. Lebed, R. E. Mitchell, and E. S. Swanson, *Prog. Part. Nucl. Phys.* **93**, 143 (2017), arXiv:1610.04528 [hep-ph].
- [14] F.-K. Guo, C. Hanhart, U.-G. Meißner, Q. Wang, Q. Zhao, and B.-S. Zou, *Rev. Mod. Phys.* **90**, 015004 (2018), arXiv:1705.00141 [hep-ph].
- [15] S. L. Olsen, T. Skwarnicki, and D. Zieminska, *Rev. Mod. Phys.* **90**, 015003 (2018), arXiv:1708.04012 [hep-ph].
- [16] A. Ali, J. S. Lange, and S. Stone, *Prog. Part. Nucl. Phys.* **97**, 123 (2017), arXiv:1706.00610 [hep-ph].
- [17] N. Brambilla, S. Eidelman, C. Hanhart, A. Nefediev, C.-P. Shen, C. E. Thomas, A. Vairo, and C.-Z. Yuan, *Phys. Rept.* **873**, 1 (2020), arXiv:1907.07583 [hep-ex].
- [18] Y.-R. Liu, H.-X. Chen, W. Chen, X. Liu, and S.-L. Zhu, *Prog. Part. Nucl. Phys.* **107**, 237 (2019), arXiv:1903.11976 [hep-ph].
- [19] H.-X. Chen, W. Chen, X. Liu, Y.-R. Liu, and S.-L. Zhu, (2022), arXiv:2204.02649 [hep-ph].
- [20] B. Aubert *et al.* (BaBar), *Phys. Rev. Lett.* **90**, 242001 (2003), arXiv:hep-ex/0304021.
- [21] M. Altenbuchinger, L. S. Geng, and W. Weise, *Phys. Rev.* **D89**, 014026 (2014), arXiv:1309.4743 [hep-ph].
- [22] F.-K. Guo, P.-N. Shen, and H.-C. Chiang, *Phys. Lett.* **B647**, 133 (2007), arXiv:hep-ph/0610008 [hep-ph].
- [23] G. S. Bali, S. Collins, A. Cox, and A. Schäfer, *Phys. Rev. D* **96**, 074501 (2017), arXiv:1706.01247 [hep-lat].
- [24] M.-W. Hu, X.-Y. Lao, P. Ling, and Q. Wang, *Chin. Phys. C* **45**, 021003 (2021), arXiv:2008.06894 [hep-ph].
- [25] M. Ablikim *et al.* (BESIII), *Phys. Rev. Lett.* **110**, 252001 (2013), arXiv:1303.5949 [hep-ex].
- [26] M. Ablikim *et al.* (BESIII), *Phys. Rev. Lett.* **126**, 102001 (2021), arXiv:2011.07855 [hep-ex].
- [27] L. Meng, B. Wang, and S.-L. Zhu, *Phys. Rev. D* **102**, 111502 (2020), arXiv:2011.08656 [hep-ph].
- [28] Z. Yang, X. Cao, F.-K. Guo, J. Nieves, and M. P. Valderrama, *Phys. Rev. D* **103**, 074029 (2021), arXiv:2011.08725 [hep-ph].
- [29] B. Wang, L. Meng, and S.-L. Zhu, *Phys. Rev. D* **103**, L021501 (2021), arXiv:2011.10922 [hep-ph].

- [30] M.-J. Yan, F.-Z. Peng, M. Sánchez Sánchez, and M. Pavon Valderrama, Phys. Rev. D **104**, 114025 (2021), arXiv:2102.13058 [hep-ph].
- [31] M.-L. Du, M. Albaladejo, F.-K. Guo, and J. Nieves, Phys. Rev. D **105**, 074018 (2022), arXiv:2201.08253 [hep-ph].
- [32] V. Baru, E. Epelbaum, A. A. Filin, C. Hanhart, and A. V. Nefediev, Phys. Rev. D **105**, 034014 (2022), arXiv:2110.00398 [hep-ph].
- [33] Q.-Y. Zhai, M.-Z. Liu, J.-X. Lu, and L.-S. Geng, (2022), arXiv:2205.03878 [hep-ph].
- [34] R. Aaij et al. (LHCb), Phys. Rev. Lett. **122**, 222001 (2019), arXiv:1904.03947 [hep-ex].
- [35] C. W. Xiao, J. Nieves, and E. Oset, Phys. Rev. **D100**, 014021 (2019), arXiv:1904.01296 [hep-ph].
- [36] S. Sakai, H.-J. Jing, and F.-K. Guo, Phys. Rev. **D100**, 074007 (2019), arXiv:1907.03414 [hep-ph].
- [37] Y. Yamaguchi, H. García-Tecocoatzi, A. Giachino, A. Hosaka, E. Santopinto, S. Takeuchi, and M. Takizawa, Phys. Rev. D **101**, 091502 (2020), arXiv:1907.04684 [hep-ph].
- [38] M.-Z. Liu, T.-W. Wu, M. Sánchez Sánchez, M. P. Valderrama, L.-S. Geng, and J.-J. Xie, Phys. Rev. D **103**, 054004 (2021), arXiv:1907.06093 [hep-ph].
- [39] M. Pavon Valderrama, Phys. Rev. **D100**, 094028 (2019), arXiv:1907.05294 [hep-ph].
- [40] L. Meng, B. Wang, G.-J. Wang, and S.-L. Zhu, Phys. Rev. **D100**, 014031 (2019), arXiv:1905.04113 [hep-ph].
- [41] M.-L. Du, V. Baru, F.-K. Guo, C. Hanhart, U.-G. Meißner, J. A. Oller, and Q. Wang, Phys. Rev. Lett. **124**, 072001 (2020), arXiv:1910.11846 [hep-ph].
- [42] M.-Z. Liu, Y.-W. Pan, and L.-S. Geng, Phys. Rev. D **103**, 034003 (2021), arXiv:2011.07935 [hep-ph].
- [43] M.-Z. Liu and L.-S. Geng, Eur. Phys. J. C **81**, 179 (2021), arXiv:2012.05096 [hep-ph].
- [44] M. J. Savage and M. B. Wise, Phys. Lett. **B248**, 177 (1990).
- [45] J. Hu and T. Mehen, Phys. Rev. **D73**, 054003 (2006), arXiv:hep-ph/0511321 [hep-ph].
- [46] M. Padmanath, R. G. Edwards, N. Mathur, and M. Peardon, Phys. Rev. D **91**, 094502 (2015), arXiv:1502.01845 [hep-lat].
- [47] Y.-C. Chen and T.-W. Chiu (TWQCD), Phys. Lett. B **767**, 193 (2017), arXiv:1701.02581 [hep-lat].
- [48] C. Alexandrou and C. Kallidonis, Phys. Rev. D **96**, 034511 (2017), arXiv:1704.02647 [hep-lat].
- [49] N. Mathur and M. Padmanath, Phys. Rev. D **99**, 031501 (2019), arXiv:1807.00174 [hep-lat].
- [50] M.-Z. Liu, T.-W. Wu, J.-J. Xie, M. Pavon Valderrama, and L.-S. Geng, Phys. Rev. **D98**, 014014 (2018), arXiv:1805.08384 [hep-ph].
- [51] M.-Z. Liu, Y. Xiao, and L.-S. Geng, Phys. Rev. **D98**, 014040 (2018), arXiv:1807.00912 [hep-ph].

- [52] E. J. Eichten and C. Quigg, Phys. Rev. Lett. **119**, 202002 (2017), arXiv:1707.09575 [hep-ph].
- [53] R. Aaij *et al.* (LHCb), Nature Phys. **18**, 751 (2022), arXiv:2109.01038 [hep-ex].
- [54] R. Aaij *et al.* (LHCb), Nature Commun. **13**, 3351 (2022), arXiv:2109.01056 [hep-ex].
- [55] J.-B. Cheng, S.-Y. Li, Y.-R. Liu, Z.-G. Si, and T. Yao, Chin. Phys. C **45**, 043102 (2021), arXiv:2008.00737 [hep-ph].
- [56] F.-K. Guo, C. Hidalgo-Duque, J. Nieves, and M. P. Valderrama, Phys. Rev. **D88**, 054014 (2013), arXiv:1305.4052 [hep-ph].
- [57] M.-Z. Liu, F.-Z. Peng, M. Sánchez Sánchez, and M. P. Valderrama, Phys. Rev. **D98**, 114030 (2018), arXiv:1811.03992 [hep-ph].
- [58] Y.-W. Pan, M.-Z. Liu, F.-Z. Peng, M. Sánchez Sánchez, L.-S. Geng, and M. Pavon Valderrama, Phys. Rev. D **102**, 011504 (2020), arXiv:1907.11220 [hep-ph].
- [59] T.-W. Wu, Y.-W. Pan, M.-Z. Liu, S.-Q. Luo, L.-S. Geng, and X. Liu, Phys. Rev. D **105**, L031505 (2022), arXiv:2108.00923 [hep-ph].
- [60] S.-Q. Luo, T.-W. Wu, M.-Z. Liu, L.-S. Geng, and X. Liu, Phys. Rev. D **105**, 074033 (2022), arXiv:2111.15079 [hep-ph].
- [61] Y. Kim, M. Oka, and K. Suzuki, Phys. Rev. D **105**, 074021 (2022), arXiv:2202.06520 [hep-ph].
- [62] W.-X. Zhang, H. Xu, and D. Jia, Phys. Rev. D **104**, 114011 (2021), arXiv:2109.07040 [hep-ph].
- [63] X.-Z. Weng, W.-Z. Deng, and S.-L. Zhu, Chin. Phys. C **46**, 013102 (2022), arXiv:2108.07242 [hep-ph].
- [64] N. Isgur and M. B. Wise, Adv. Ser. Direct. High Energy Phys. **10**, 234 (1992).
- [65] J. M. Flynn and N. Isgur, J. Phys. G **18**, 1627 (1992), arXiv:hep-ph/9207223.
- [66] M.-Z. Liu, Y.-W. Pan, F.-Z. Peng, M. Sánchez Sánchez, L.-S. Geng, A. Hosaka, and M. Pavon Valderrama, Phys. Rev. Lett. **122**, 242001 (2019), arXiv:1903.11560 [hep-ph].
- [67] J.-M. Xie, X.-Z. Ling, M.-Z. Liu, and L.-S. Geng, (2022), arXiv:2204.12356 [hep-ph].
- [68] P. Junnarkar and N. Mathur, Phys. Rev. Lett. **123**, 162003 (2019), arXiv:1906.06054 [hep-lat].
- [69] Y. Lyu, H. Tong, T. Sugiura, S. Aoki, T. Doi, T. Hatsuda, J. Meng, and T. Miyamoto, Phys. Rev. Lett. **127**, 072003 (2021), arXiv:2102.00181 [hep-lat].
- [70] Z.-H. Zhang and F.-K. Guo, Phys. Rev. Lett. **127**, 012002 (2021), arXiv:2012.08281 [hep-ph].