Chapter 18

Quantum Glasses

Leticia F. Cugliandolo^{1,2} and Markus Müller³

¹Sorbonne Université, Laboratoire de Physique Théorique et Hautes Energies, CNRS UMR 7589, 4 Place Jussieu, 75252 Paris 05, France ²Institut Universitaire de France, 1 rue Descartes, 75231 Paris 05, France ³Laboratory for Theoretical and Computational Physics, Paul Scherrer Institute, Villigen CH-5232, Switzerland

18.1. Introduction

The interplay of disorder, interactions and quantum mechanics leads to a host of interesting phenomena. A particularly intriguing aspect is the breakdown of ergodicity in a way that differs significantly from that of standard spontaneous symmetry breaking. This can arise as a consequence of quenched disorder, as in spin glasses, or structural disorder, think of amorphous density order. Both quenched and self-generated disorder entail a clustering in phase space associated with glass attributes like memory effects and extremely slow relaxations [1–3]. The replica method allows one to access the phase space structure, and replica symmetry breaking (RSB) signals the existence of a large number of metastable states separated by substantial free energy barriers. Even though barriers may be finite in low dimensions, preventing a genuine thermodynamic glass transition, the replica method has proven useful for these cases as well.

As a result of frustration and the competition between many nearly degenerate ordering patterns, the glassy order is often rather soft and gives rise to an unusually high density of low energy excitations. In or close to a quantum glass phase those may have a strong influence on electronic transport and have often been invoked as candidates to explain the non-Fermi liquid behavior of the resistivity of complex materials, such as high temperature superconductors (cuprates), heavy fermion systems, but also systems close to a Mott-Anderson metal-insulator transition.

Apart from this "glassy" route, very slow dynamics and non-ergodicity in quantum systems can be due to other mechanisms too, especially in low dimensions where the effects of disorder are very strong. On the one hand, under a real space renormalization group treatment, disordered quantum spin chains typically flow to fixed points characterized by infinite randomness [4], with extended regions of parameter space where Griffiths–McCoy singularities are very important [5, 6]. Thereby the low frequency response is dominated by nearly isolated degrees of freedom only weakly coupled to others further away.

On the other hand, non-interacting quantum particles in one dimension subject to a disordered potential Anderson localize and stop diffusing on the scale of the elastic mean free path [7, 8]. Its interacting counterpart, many-body localization (MBL) [9–12], relies on the discreteness of non-interacting excitations and the rareness of resonant interactions, which then fail to induce diffusion of energy. Thus, MBL can only occur in isolation, with no coupling to the continuum of a bath which would reinstate transport and ergodicity. In contrast, the above discussed glass phases are stable against a bath, as they rely on the rugged structure of the free-energy landscape. A further difference between quantum glasses and MBL systems is that the latter are highly susceptible to local inclusions where disorder is weak. Those act as sources of dynamical chaos, and it is believed that only in low-dimensional, discrete lattices such nucleation centers of local baths do not destroy MBL [13-15] (although the existence of MBL even in 1d has been questioned recently [16–18]). In contrast, glasses, like standard symmetry broken phases, are stabilized in high dimensions (and RSB is certain to occur above a d_c [19]). In summary, while both quantum glassiness and MBL entail non-ergodicity and impede full thermalization, they do so for fundamentally different reasons.

Other obstructions to full ergodicity exist, e.g., integrable systems with an extensive number of conserved quantities [20], systems hosting many-body scars (a set of eigenstates that violate the eigenstate thermalization hypothesis) [21] or cases with fragmented Hilbert spaces [22]. However, none of these is robust against generic perturbations of the Hamiltonian, as even modest ones reinstate equilibrating dynamics rather quickly. We do not consider them as quantum glassy and we will not discuss them here. A further way to kill ergodicity is the sufficiently strong coupling to an Ohmic bath of a single quantum degree of freedom, which can get dressed with many bath modes, so as to acquire an infinite mass, and its dynamics freeze out [23, 24]. We shall not discuss this effect of non-glassy "local ergodicity" breaking either, except for noting that the coupling of quantum systems to a bath usually results in the reduction of fluctuations, and thus extends the regime of stability of the glass [25, 26].

In this review we focus on quantum glasses that arise in high dimensions, where disorder is so relevant as to produce amorphous order and phase space clusterization, but does not flow to infinite randomness. This structure withstands both thermal and quantum fluctuations, which soften the energy landscape and eventually melt the glass. For certain types of glasses the quantum melting differs drastically from the thermal route.

Since this chapter is part of a book devoted to replica symmetry breaking, we focus the discussion on equilibrium aspects and we structure it as follows. In Sec. 18.2 we overview some general aspects of quantum glassy dynamics. Section 18.3 reminds the gist of the replica treatment of mean field glasses. In Sec. 18.4 we classify different universality classes of (mean field) glasses according to the type of interactions and environments, and discuss their phenomenology, in particular the nature of their low energy collective modes which we obtain both from Landau theory and an effective potential approach in Sec. 18.5. In Sec. 18.6 we discuss whether an isolated quantum glass can explore its clusterized phase space by tunneling or stays localized. The interplay between the glass and localization routes to non-equilibrium is a multi-faceted topic

briefly reviewed in Sec. 18.7. Section 18.8 presents some experimental glassy materials. Finally, in Sec. 18.9 we survey fermionic models closely connected to the paramagnetic phase of Heisenberg spin glasses, a.k.a. the Sachdev-Ye-Kitaev (SYK) model, that may fertilize the study of spin glasses via interesting dynamical analogies.

18.2. Dynamics in quantum glassy phases

A particularly interesting aspect of quantum glasses is their dynamics, both in and out of equilibrium. We broadly survey them here and discuss selected questions in more detail in later sections.

18.2.1. Short-time dynamics

Distinct quantum aspects of glasses appear on microscopic time scales. The high-frequency dynamics depends crucially on the symmetries of the order parameter and the nature of the quantum fluctuations. In the low-frequency limit, instead, the excitations feel the softness of the glassy landscape, which, in the mean-field limit, exhibits surprisingly universal spectral features whose frequency scaling merely depends on the glass being metallic or insulating. We will review the salient results in Secs. 18.4, 18.5 and 18.9.

18.2.2. Moderately long-time dynamics

The intermediate to long-time dynamics of quantum glassy systems are essentially identical to those of their classical counterparts. Indeed, one expects that on time scales $t \gg \hbar/T$ all degrees of freedom decohere and any quantum effects are washed out. After a quench from the disordered phase, the glass hovers over saddles in the free energy landscape and slowly relaxes its energy without ever getting deeply trapped into rare deep minima, in a manner dubbed weak ergodicity breaking [27–31]. The response slows down increasingly with time (the glass "ages"). In solvable mean field models, the fluctuation-dissipation relation evaluated on those long time scales takes the classical form, albeit with an effective temperature $T_{\rm eff}$ that exceeds the bath temperature (even at T=0). The latter is formally related to Parisi's order parameter Q(x) as in the classical limit [1]. Moreover, the equations governing the slow evolution of the response and correlation functions are invariant under time reparametrizations, $t \to f(t)$ [32]. Such an invariance also occurs in the large M limit of the SU(M) Heisenberg and the SYK models. Recent progress in this direction will be reviewed in Sec. 18.9. Finally, as in the classical limit, one can tune the initial state to be within one of the valleys in the free-energy landscape and show that the subsequent dynamics will remain confined to it (at least for times exponentially large in system size).

A short summary of the real-time dynamics of quantum disordered systems on these time scales can be found in Ref. [33]. We will not extend this discussion further here, our aim being to focus mainly on equilibrium properties.

18.2.3. Extremely long-time dynamics

On extremely long time scales, crossing over or tunneling under barriers between metastable valleys should be possible. In mean-field models barriers scale with system size and thermal activation over them is exponentially slow. Since the latter is the root cause for the failure of simulated annealing in finding the ground state of NP-hard problems, there was hope that quantum annealing could achieve a significant speed-up [34]. The idea was to initialize the system under strong quantum fluctuations (e.g. a transverse field) which induce a simple ground state. Then, by invoking the adiabatic theorem, one should reach the classical ground state upon turning off the fluctuations sufficiently slowly. The caveat is that exponentially slow switch-off rates are required to follow the ground state while it delicately hybridizes distant low-energy valleys in phase space. Still, one may ask which of the exponentially long times (barrier crossing or adiabatic ground state preparation) is more efficient to explore complex landscapes [35], an idea that was explored experimentally in quantum Ising spin glasses [36], cf. Sec. 18.8.1. However, an even bigger obstacle to quantum annealing [37, 38] is posed by the fact that NP-hard problems have quantum first order transitions [39, 40] (see Sec. 18.4.1.2).

A related dynamical question arises for an isolated quantum spin glass – the setting considered in MBL: after preparation in a deep valley, can it tunnel to others or does it remain localized? We discuss it in Sec. 18.6.

18.3. Replica treatment of quantum mean-field glasses

Most theoretical insights on quantum glasses have been obtained from mean-field SU(M) models, where all $i=1,\ldots,N$ spins interact with all others via random couplings with variance proportional to J^2 and conveniently scaled with N and M. While this looks remote from any real physical glass, long range couplings can be emulated via light-matter coupling in multi-mode cavities, see Sec. 18.8.3. Their equilibrium properties were derived with a replicated imaginary-time path integral formulation of the partition function [41]. A saddle-point evaluation allows one to define the glass order parameter*

$$Q_{ab}(\tau) = \frac{1}{NM^2} \sum_{i} [\langle T \, \boldsymbol{\sigma}_i^a(\tau) \cdot \boldsymbol{\sigma}_i^b(0) \rangle]$$
 (18.1)

where $a,b=1,\ldots,n$ are the replica indices, τ the imaginary time and T the time-ordering operator. Depending on the complexity of the model, the free-energy density can either be just a functional of the saddle-point $Q_{ab}(\tau)$, or the partition function can be transformed into one for a single spin with $Q_{ab}(\tau)$ self-consistently given by the expectation value of its imaginary-time correlations (see, e.g., [42]). In all cases, $Q_{a\neq b}$ are τ -independent, vanish in the disordered phase, and adopt an RSB structure in the glass phase. The diagonal elements $Q_{aa}(\tau) \equiv q_d(\tau)$ are independent of the replica index but depend on time, being $\beta\hbar$ periodic, in the whole phase diagram. In certain cases (typically models in which also the single-spin partition function can be calculated

^{*}The normalization factor M^2 captures the leading dependence for $M \to \infty$. In general one should naturally divide by the number of spin components $(M^2 - 1)$.

exactly), Q_{ab} satisfies a generic differential equation of the form

$$G_0^{-1}(\tau)_{ac}Q_{cb}(\tau) = \delta_{ab}\delta(\tau) + \int_0^{\beta\hbar} d\tau' \; \Sigma_{ac}(\tau - \tau')Q_{cb}(\tau') \;,$$
 (18.2)

with a sum over repeated indices. G_0^{-1} is a differential operator, typically diagonal in replica indices, and its order (first or second time derivative, further time dependence) depends on the kind of degrees of freedom and whether the system is coupled to a bath (microscopic dynamics). Σ is a self-energy which typically depends on time only via~Q. Note though that this equation is not causal, and is simpler to solve in frequency space. We will specify G_0^{-1} and Σ in several concrete cases below. The physical properties follow from Q_{ab} . Of special importance is the out-of-phase local susceptibility

$$\chi_{\text{loc}}^{"}(\omega) = \text{Im}\,\tilde{q}_d(\omega + i0^+)\,\,,\tag{18.3}$$

with \tilde{q}_d the Fourier transform of q_d . The right-hand-side is the analytic continuation to real ω of $\tilde{q}_d(\omega_n)$, with ω_n the Matsubara frequencies. χ''_{loc} contains essential information on the spectrum of collective excitations since

$$\chi_{\text{loc}}''(\omega) = \frac{\pi}{N} \sum_{im} |\langle \psi_m | \sigma_i^1 | \psi_0 \rangle|^2 \left[\delta(\omega - E_m + E_0) - \delta(\omega + E_m - E_0) \right],$$

(at T=0) where (ψ_m, E_m) are the many-body eigenstates and energies, with 0 referring to the ground state. A continuous glass transition is identified by the condition $J\chi_{loc} = 1$, with χ_{loc} the zero frequency limit of the local susceptibility, $\chi_{loc} = \int_0^{\beta\hbar} d\tau \, q_d(\tau)$.

18.4. Classification of mean-field quantum glasses

The local degrees of freedom, their interactions and the environment they couple to define different mean-field classes, distinguished by their phase diagrams and properties. We list three distinguishing aspects of spin models and discuss them in turn below.

- The number of interacting spins decides on the organization of phase space as elucidated, for instance, by the replica analysis and its symmetry breaking scheme.
 This distinction drastically affects the nature and the order of the quantum glass transition.
- The symmetry group in spin space is important. It matters whether the various spin-spin interaction terms mutually commute (as in Ising and rotor models) or not (as in Heisenberg systems). In the latter case quantum fluctuations are substantially stronger, entailing a weaker glassy order and softer excitations, both at low and high T.
- The low-frequency dynamics depend on whether the quantum glass is embedded in an insulating, gapped host, or couples to a gapless bath with Ohmic spectrum, as in metallic glasses. For a given environment – insulating or metallic – the dynamically relevant states of all mean-field models, in spite of quantitative differences, display a striking universality in their low-frequency response within the glass phase (Sec. 18.5).

Note that these spin glass models do not cover all possible glassy quantum systems. Some other cases are pinned elastic systems (including vortex lattices in superconductors, Wigner crystals, charge and spin-density waves or disordered liquid crystals) in which frustrating disorder originates, e.g., from substrate impurities [43]. The interested reader may consult Ref. [44]. Nonetheless, once coarse-grained at the collective pinning scale, these systems resemble short-range coupled spin systems in random fields.

18.4.1. Pair vs multi-spin interactions

18.4.1.1. Pair interactions (p = 2) - continuous glass transition

Usually one considers Ising-like interactions when discussing the distinction between pair and multi-spin interactions. Ising-type glasses arise in numerous contexts, as they describe generic interacting two level systems (TLS). The simplest representative is the transverse field Ising model (TFIM)

$$\hat{\mathcal{H}} = -\sum_{ij} J_{ij} \,\hat{\sigma}_i^z \hat{\sigma}_j^z + \sum_i \mathbf{H}_i \cdot \hat{\boldsymbol{\sigma}}_i \tag{18.4}$$

where $\hat{\sigma}_i = (\hat{\sigma}_i^x, \hat{\sigma}_i^y, \hat{\sigma}_i^z)$, with the usual Pauli matrices acting in the Hilbert space of the *i*'th TLS. In finite dimensions the first sum extends over nearest neighbours of a lattice. The interaction strengths J_{ij} are drawn from a probability distribution, $P(J_{ij})$, typically chosen to be Gaussian with zero mean $[J_{ij}] = 0$ and variance $[J_{ij}^2] = J^2/(2c)$. J is O(1) and c is the connectivity of the lattice. Mean values over P are indicated with square brackets. The \mathbf{H}_i are local fields. Most often one restricts to a homogeneous transverse field, $\mathbf{H}_i = \Gamma \mathbf{e}_x$. Since the last term does not commute with the interactions, it induces non-trivial quantum dynamics. In the absence of longitudinal fields H_i^z , the Hamiltonian possesses an Ising symmetry which is spontaneously broken at sufficiently low T and weak transverse fields $\Gamma < \Gamma_c$. For $H_i^z \neq 0$ (as in many experimental systems of interest) the only potential phase transition is an RSB one, which however occurs only in high enough dimensions.

Allowing each spin to interact with all others, $c \to N-1$, places the spins on a complete graph. The scaling of the variance, $[J_{ij}^2] \approx J^2/(2N)$, ensures a non-trivial thermodynamic limit, $N \to \infty$. The Hamiltonian becomes the quantum extension [41] of the Sherrington-Kirkpatrick (SK) spin-glass (TFSK), which is tractable with the replica trick. Again, only the replica diagonal, $q_d(\tau) \equiv Q_{aa}(\tau) = [\langle T\sigma_a^z(\tau)\sigma_a^z(0)\rangle]$ depends on the imaginary time τ , while the RSB structure of the off-diagonal elements remains basically the same as in the classical limit. Glassy freezing is signalled by a finite Edwards-Anderson parameter $q_{\rm EA} \equiv \lim_{\tau \to \infty} q_d(\tau)$ (at T=0). The resulting single-site problem, with the coupling to the rest encapsulated self-consistently in Q_{ab} , is of similar difficulty as the impurity problems of Dynamical Mean Field Theory [45].

From the stability criterion $1 = J\chi_{\rm loc}$, a $(T/J, \Gamma/J)$ phase diagram with a second order phase transition between a paramagnetic and a spin-glass phase is predicted [41]. Quantum fluctuations depress the transition temperature but do not destroy the transition. For $\Gamma \to 0$ the classical SK $T_c = J$ is recovered and for $T \to 0$ a quantum critical point at $\Gamma_c = 1.52J$ is found [46]. Order sets in when the entropy loss and/or the loss in transverse field energy are compensated by the gain in interaction energy.

Close to the phase transition both scale like $O(m^2)$, where $m \ll 1$ is a typical ordered moment of a single spin. The glass transition, whether classical or quantum, can thus be viewed as a condensation into the first magnetization mode to become unstable. The glass transition differs crucially from standard ordering phenomena, however, because a large number of modes turn soft almost simultaneously. This entails many local minima with different ordering patterns. Moreover, throughout the entire glass phase there are permanently some gapless, critical modes at the verge of condensing.

Interestingly enough, a "static approximation" in which $q_d(\tau)$ is replaced by a τ -independent variational parameter captures the essence of the phase transition. A more detailed dynamical analysis for the Ising case [47] and related rotor models with $M \gg 1$ components [48] shows that the spectral gap closes as $\Delta \sim [(\Gamma - \Gamma_c)/\ln(\Gamma - \Gamma_c)]^{1/2}$ at Γ_c , assuming a gapless spectral function $\chi''_{loc}(\omega) \sim \omega$, associated with algebraic decay in imaginary time, $q_d(\tau) \sim 1/\tau^2$.

Determining Q_{ab} and $q_d(\tau)$ deeper in the glass phase requires the solution of a quantum impurity problem in a frozen field whose distribution is controlled by full RSB in the off-diagonal Q_{ab} [49]. The latter ensures marginal stability, which in turn implies a gapless spectrum *everywhere* in the glass phase. Moreover, the limit of small Γ admits a scaling form for the mean field solution. From this it follows that the low-frequency spectral function becomes independent of Γ ,

$$\chi_{\text{loc}}''(\omega; \Gamma \ll J) = 0.59 \,\omega/J^2,\tag{18.5}$$

which we will interpret physically in Sec. 18.5.

18.4.1.2. p > 2 - discontinuous onset of glassy order

Models in which p > 2 classical spins interact simultaneously are particularly interesting since at the mean field level their dynamics are described by equations identical to those arising in the Mode-Coupling Theory of structural glasses. It is therefore believed that they capture the physics of the structural glass transition and the glassy phase [1, 2]. Moreover, such multi-spin Hamiltonians are ubiquitous in combinatorial optimization problems when translated into questions about ground states of statistical mechanics systems. Defined on diluted graphs, the p-spin models are, e.g., closely connected to the K-satisfiability problem [50].

A typical multi-spin Hamiltonian reads [25, 26, 40, 51–53]

$$\hat{\mathcal{H}} = -\sum_{i_1 \neq \dots \neq i_p} J_{i_1 \dots i_p} \hat{\sigma}_{i_1}^z \dots \hat{\sigma}_{i_p}^z + \sum_i \Gamma_i \hat{\sigma}_i^x , \qquad (18.6)$$

with the parameter p taking any integer value $p \geq 3$. The model is defined on a hypergraph, the sum running over all p-uplets. The couplings are random independent variables with variance $p!J^2/(2N^{p-1})$.

The important difference w.r.t. pairwise interacting models lies in the scaling of the loss of entropy or transverse field energy as compared to the interaction energy gain with a putative small emerging magnetization pattern $m_i = \langle \sigma_i^z \rangle$. The former still scale as $O(m^2)$, but the energy gain is only $O(m^p)$ and cannot compensate for this loss if m is small. The only possibility for a phase transition is that the order

parameter, $q_{\rm EA}=1/N\sum m_i^2$ jumps discontinuously to a finite value at a critical point. Technically, this is reflected by a one-step RSB. Part of the liquid entropy is transferred to configurational entropy without a thermodynamic phase transition (the free energy is smooth). The Gibbs weight distributes over an exponential number of metastable states. When quantum fluctuations are switched on, and temperature is sufficiently low, the transition necessarily has to change nature since the configurational entropy cannot contribute to the free energy. The phase transition thus becomes truly first order [29] and the ground states on the two sides of the transition are essentially unrelated. For this reason quantum annealing is not expected to be useful for (p>2)-spin systems and other systems with a 1-step RSB.

In quantum and classical glasses of this type alike, the proliferation of metastable states affects the dynamics, but does not necessarily change the static observables. Over a finite range of free-energy densities one finds exponentially many local minima, the fewer the lower the free-energy. As temperature decreases the Gibbs weight concentrates on lower energy states. A thermodynamic freeze-out transition occurs once the lowest O(1) states dominate. Consequently, the dynamical glassy phase extends beyond the thermodynamic one [27, 28, 54].

The p-spin potential was also used to model a quantum particle in a random potential, in which the interactions are between p-uplets of coordinate positions on an N-dimensional sphere. Conventional kinetic energy is given to the particle. In these cases an equation of the kind of Eq. (18.2) applies with $G_{0\ ab}^{-1} = (d_{\tau}^2 + \mu)\delta_{ab}$ with μ a Lagrange multiplier enforcing the spherical constraint and $\Sigma_{ab} = J^2 Q_{ab}^{\bullet(p-1)}$ where \bullet denotes a normal power. Self-energies of this very same kind appear in the SYK model, Sec. 18.9.

18.4.2. Commuting vs non-commuting interactions

18.4.2.1. Commuting interactions

The disordered quantum rotor model is defined by [48]

$$\hat{\mathcal{H}} = \frac{g}{2M} \sum_{i} \hat{\mathbb{L}}_{i}^{2} + \frac{M}{\sqrt{N}} \sum_{i < j} J_{ij} \,\hat{\mathbf{n}}_{i} \cdot \hat{\mathbf{n}}_{j} \qquad \hat{\mathbf{n}}_{i}^{2} = 1 \quad \forall i .$$
 (18.7)

The M components \hat{n}_i^{μ} of the ith unit-length rotor $\hat{\mathbf{n}}_i$ commute with each other, unlike the components of quantum spins. As a consequence, all interaction terms mutually commute. $\hat{L}_i^{\mu\nu}$ (with $\mu < \nu, \, \mu, \nu = 1, \ldots, M$) are the M(M-1)/2 components of the angular-momentum generator $\hat{\mathbb{L}}_i$ in rotor space, and $[L_i^{\mu\nu}, n_j^{\sigma}] = i\delta_{ij}(\delta_{\mu\sigma}n_j^{\nu} - \delta_{\nu\sigma}n_j^{\mu})$. The J_{ij} are O(1) randomly distributed uncorrelated exchange constants. As $g \to 0$ the model reduces to the classical, infinite-range, M-component spin glass, the limit $M \to 1$ being very similar to the Ising model.

There is no Berry phase in the real action of the path integral representation of the partition function, and the O(M) symmetric saddle-point [48] in the $M \to \infty$ limit yields a self-consistency equation, Eq. (18.2), with $G_{0\ ab}^{-1} = g^{-1}(d^2/d\tau^2 + \mu)\delta_{ab}$, where μ enforces the constraint $\hat{\mathbf{n}}_i^2 = 1$, and one has $Q_{ab} = q_d\delta_{ab}$ and $\Sigma_{ab} = Q_{ab}$. In the glass phase the same equation is obeyed by $q_d(\tau) - q_{\mathrm{EA}}$, while $Q_{a\neq b} = q_{\mathrm{EA}}$ becomes non-

zero, without however breaking the replica symmetry. The integro-differential equation is easy to solve in frequency space and yields spectral functions that we will review in Sec. 18.5. The quantum critical behavior was found to be in the TFSK universality class, which was rationalized by showing that 1/M corrections do not modify the critical exponents and the low frequency spectrum.

18.4.2.2. Non-commuting interactions - SU(M) spins

The most relevant model in this class is the quantum Heisenberg spin glass,

$$\hat{\mathcal{H}} = -\sum_{i \neq j} J_{ij} \,\hat{\boldsymbol{\sigma}}_i \cdot \hat{\boldsymbol{\sigma}}_j \ . \tag{18.8}$$

A static approximation used to evaluate the instability condition for S=1/2 [41] yields a surprisingly good estimate for $T_g \approx \sqrt{3}/12 \approx 0.14 J$, obtained with Quantum Monte Carlo calculations [55]. The substantial reduction with respect to the classical $T_g = J/4$ is due to quantum fluctuations.

With the static approximation, though, one cannot access the spin dynamics. A crucial step forward was taken in [56] where SU(2) was promoted to SU(M) and a Schwinger boson representation of the spins, with the number of bosons constrained to be $n_b = SM$, was adopted. A first analysis in the (M,S) parameter space was performed in the large M limit. Soon after, a complete solution with $M \to \infty$ and fixed S was derived, and global aspects, some of them also valid for SU(2), discussed [57, 58]. The solution is formulated in terms of the Green functions of the bosons, $G_B^{ab}(\tau) \equiv -M^{-1} \sum_{\mu} [\langle \mathrm{T} \hat{b}^a_{\mu}(\tau) \hat{b}^{b\dagger}_{\mu}(0) \rangle]$ and their self-consistency equations are again of the form of Eq. (18.2) with

$$G_{0\ ab}^{-1} = (d_{\tau} + \mu^{a})\delta_{ab} , \qquad \Sigma_{B}^{ab}(\tau) = J^{2}G_{B}^{ab}(\tau)G_{B}^{ab}(\tau)G_{B}^{ab}(-\tau) .$$
 (18.9)

 μ^a is a chemical potential which fixes $G_B^{aa}(\tau=0)=-S$. The disorder-averaged spin-spin correlator is obtained from G_B by

$$q_d(\tau) \equiv \frac{1}{M^2} [\langle \hat{\boldsymbol{\sigma}}^a(\tau) \cdot \hat{\boldsymbol{\sigma}}^a(0) \rangle] = G_B^{aa}(\tau) G_B^{aa}(-\tau) . \tag{18.10}$$

For large S one finds an essentially classical glass transition at $T_g \sim S^2 J$, where the bosonic spinons condense. The 4-spinon interaction term resembles the p=4-spin problem, and for similar reasons one finds a 1-step RSB transition. This leaves open the choice of the size of replica blocks, which selects the energy of the targeted metastable states. Usually one uses the condition of marginal stability (vanishing of the "replicon" mode), as those are reached dynamically [31] and exhibit a gapless spectrum. Extremization of the free-energy density instead describes dynamically inaccessible states with a gap in $\chi''_{loc}(\omega)$.

At large S, quantum fluctuations introduce a frequency scale $\sim SJ$, which sets the crossover temperature where collective quantum dynamics emerges. Consistent with the Landau analysis of Sec. 18.5 marginal states are found to have $\chi''_{loc}(\omega) \sim \omega$ and a low-T specific heat that scales as T^3 [59, 60] (contrary to the linear T dependence originally claimed in [57, 58]).

The case of small $S \ll 1$ exhibits much stronger quantum fluctuations and is thus more interesting. Here, the spin-fluid, paramagnetic regime survives down to much lower temperatures $T \ll J$ and is radically different from the classical paramagnet of the high T and large S limits. Below $T \sim J$, the system enters a gapless, quantum critical regime in which the Green function assumes long-time tails, $G_B^{aa}(\tau) \sim 1/(J\tau)^{1/2}$, while the melon-bubble self-energy is unusually large, $\Sigma(\omega) \sim (\omega J)^{1/2} \gg \omega$, reflecting the strong scattering among spinons. From the spin-spin correlation $q_d(\tau) \sim 1/(J\tau)$ one extracts a susceptibility very similar to the one of marginal Fermi liquids: $J\chi''_{\rm loc}(\omega) \simeq \tanh(\hbar\omega/2T)$ implying $\chi'_{\rm loc} \simeq \ln(J/T)$. One further finds linear in T specific heat and a residual low-temperature entropy. This holds in the paramagnetic regime $T_g < T < J$, for an extended range of S all the way up to $S \simeq 1$. These properties are very similar to those of black holes to which this model and SYK is related.

From Eq. (18.10) one sees that the onset of glassy order modifies the Green functions beyond the time scale $\tau^* = (\omega^*)^{-1} = (q_{\rm EA}J)^{-1}$ where $G_B(\tau)G_B(-\tau) \sim q_{\rm EA}$. At that scale the spinons get confined, and $Q(\tau)$ crosses over to $Q(\tau) - q_{\rm EA} \sim \tau^*/(J\tau^2)$. This corresponds to a linear spectral function $J\chi''_{\rm loc}(\omega) \sim \omega/(q_{\rm EA}J)$, as is found in all insulating mean-field glasses, cf. Sec. 18.5. Note, however, that weak glass order implies a potentially large prefactor $\sim 1/q_{\rm EA}$, reflecting very slow collective modes.

The bosonic representation is a priori best suited for large S. Fermionic SU(M) representations are instead believed to capture better the physically most relevant case S=1/2. Such an approach yields essentially identical results for the paramagnetic phase [61], though no glass transition is found at $M=\infty$ in this case [56], since quantum fluctuations are much stronger. Yet, a 1/M expansion [62] yields an instability to a glass phase where a fermionic bilinear condenses at $T_g \sim J \exp(-\sqrt{\pi M})$ (now with continuous RSB – as the effective action is quadratic in the emerging order parameter, like in p=2 models). One expects a similar suppression of the order parameter, $q_{\rm EA} \sim T_g/J$ (which ensures that at $T \sim T_g$ the low-T linear spectral function matches that of the paramagnetic regime). Exact diagonalization for all-to-all coupled SU(2) spins [63, 64] indeed yields $q_{\rm EA} \approx 0.02$. However, currently accessible system sizes of ~ 20 spins do not allow to disentangle the regime $\omega < \omega^*$ from the broadened peak $\sim q_{\rm EA} \delta(\omega)$ reflecting static spin glass order. Nonetheless, they do exhibit spectral features as predicted by the fractionalized large M approach for $\omega > \omega^*$.

18.4.2.3. Heisenberg glasses and Many-Body Localization

We close this section by remarking that Heisenberg spin chains with coupling J under strong, but random i.i.d. local fields $\mathbf{H}_i = H_i^x \mathbf{e}_x$ drawn from a distribution of width $W \gg J$, has become one of the standard models for MBL [10]. Indeed, the typically large mismatch between the local gaps H_i^x cannot be bridged resonantly by the weak couplings J, and thus coherent, bath-free dynamics is expected to remain stuck close to an arbitrary initial state. It is worth pointing out again that MBL is favored by completely opposite ingredients than a spin glass phase: it requires the non-commuting interactions to be weak, and the dimensionality should be low. In contrast, in the parameter range where a glass with non-trivial energy landscape exists, one expects rapid and full thermalization within any local minimum of the energy landscape, independently

Book Title

of the coupling to a bath.

18.4.3. Insulating vs metallic environment

Remarkably, all insulating mean-field quantum glasses in their marginally stable glassy states exhibit the same linear scaling $\chi''_{loc}(\omega) \sim \omega$, albeit with quantitatively different prefactors, and a low-T specific heat $c_V \sim T^3$. This contrasts with a spin glass coupled to a gapless bath, e.g. via Kondo coupling to itinerant conduction band electrons, as in heavy fermion materials [65]. The dissipation due to the low-energy bath degrees of freedom slows down the collective modes and thereby substantially enhances the lowfrequency spin spectral function and the spin contribution to the specific heat, as we will review via a Landau approach in the next section.

18.5. Landau theory & the low frequency spectrum

A Landau expansion was reviewed in [66] and we only sketch it here. The aim is to construct an effective action for the glass order parameter Q_{ab} by integrating out all other degrees of freedom. This approach is expected to succeed for phase transitions in which the spins undergo standard ordering, as e.g. in the TFIM and rotor models. However, in Heisenberg-type models, where at least for $M \gg 1$ the spins fractionalize into deconfined spinons in the paramagnetic phase, it will likely fail to capture the entire dynamical crossover functions correctly. Methods as the ones for deconfined quantum critical points [67] may instead be required. Still, even in these cases the Landau approach may capture well the glassy phase in its low-frequency, spinon-confined regime.

Considering symmetry restrictions and constraints on the replica structure, and assuming the absence of time-reversal symmetry-breaking fields, the resulting meanfield action close to the quantum glass transition takes the form [68]

$$n\beta \mathcal{F}[\delta Q] = \sum_{a} \int d\tau \, \mathcal{L}[\delta Q_{aa}] + u \sum_{a} \int d\tau \, [\delta Q_{aa}(\tau, \tau)]^2$$
$$-c_3 \sum_{abc} \int d\tau_1 d\tau_2 d\tau_3 \delta Q_{ab}(\tau_1, \tau_2) \delta Q_{bc}(\tau_2, \tau_3) \delta Q_{ca}(\tau_3, \tau_1) + O(\delta Q^4) ,$$

where $\delta Q(\tau, \tau') = Q_{ab}(\tau, \tau') - c\delta_{ab}\delta(\tau - \tau')$, with the constant c fixed to remove the uninteresting and non-universal short-time dynamics. The coefficients are bare correlations of the degrees of freedom coupling to Q, but only in the linear term their timedependence is relevant. The leading low-frequency dependencies, distinguish whether the degrees of freedom are gapped (insulating) or gapless like in a Fermi sea (metallic), and are

$$\mathcal{L}_{\text{ins}}[\delta Q_{aa}] = (r + \partial_{\tau_1} \partial_{\tau_2}) \delta Q_{aa}(\tau_1, \tau_2) ,$$

$$\mathcal{L}_{\text{met}}[\delta Q_{aa}] = [r' - (\tau_1 - \tau_2)^{-2}] \delta Q_{aa}(\tau_1, \tau_2) ,$$
(18.11)

where $\tau_1 = \tau_2 = \tau$. In the metallic case we dropped the subleading time derivatives, obtaining a leading $|\omega_n|$ dependence in Matsubara space, while the coefficient in insulating glasses displays a faster ω_n^2 dependence.

18.5.1. Insulating glasses

Minimizing \mathcal{L}_{ins} with respect to δQ close to the quantum transition at T=0 yields [68] $\delta \tilde{Q}_{aa}(\omega_n) = -\sqrt{\omega_n^2 + \Delta^2}$ (dropping numerical coefficients), which implies the spectral function

$$\chi''_{\rm loc}(\omega) = \operatorname{sgn}(\omega)\sqrt{\omega^2 - \Delta}\Theta(|\omega| - \sqrt{\Delta})$$
 (18.12)

The parameter-dependent energy Δ plays the role of a spectral gap, which closes, $\Delta \to 0$, upon approaching the transition from the paramagnetic side. However, remarkably, within the glassy phase the gap remains pinned to zero, $\Delta = 0$, which entails a linear low frequency spectral function:

$$\chi_{\rm loc}''(\omega) = \omega . {(18.13)}$$

This dynamical crossover agrees with the exact solution of large M quantum rotors and the critical behavior of the TFIM. Allowing for RSB, one finds marginally stable solutions with a vanishing replicon and gapless spectrum.

Note that, as anticipated above, the high-frequency spectrum predicted by the Landau approach differs qualitatively from the spin susceptibility $\chi''_{loc}(\omega) \sim sgn(\omega)$ of the large M limit of Heisenberg models, which hinges on spinon deconfinement.

18.5.2. Metallic glasses

Minimizing \mathcal{L}_{met} close to the transition one finds [65, 69], $\delta Q_{aa}(\omega_n) = -\sqrt{|\omega_n| + \Upsilon}$, whereby the crossover energy Υ vanishes again at the quantum transition, $\Upsilon \to 0$, and stays pinned to $\Upsilon = 0$ in the glassy phase. Here the spectral function is everywhere gapless and approaches

$$\chi_{\text{loc}}''(\omega) = \frac{1}{\sqrt{2}} \frac{\omega}{\sqrt{\Upsilon + \sqrt{\omega^2 + \Upsilon^2}}} \xrightarrow{\Upsilon \to 0} \text{sgn}(\omega) \frac{|\omega|^{1/2}}{\sqrt{2}}$$
(18.14)

in the glass phase, which, at low frequencies, is substantially stronger than the one of insulating glasses, Eq. (18.13). Again, the Landau approach does not capture the high-frequency features associated with spinon deconfinement [64].

18.5.3. Physical interpretation of glassy spectral functions

The low-frequency spectral functions in the glass phase can be interpreted as arising from of set of collective random spin-density modes that behave as either underdamped or overdamped harmonic oscillators.

In the glass phase a local minimum of the free-energy landscape (expressed as a functional of local magnetizations) has a Hessian with positive eigenvalues λ_k which may be interpreted as generalized spring constants. The marginal stability of the relevant glass states assures that their distribution is gapless, and random matrix theory suggests that at small λ , it behaves as $\rho(\lambda) \equiv N^{-1} \sum_{k} \delta(\lambda - \lambda_k) = c \sqrt{\lambda}$.

In the insulating case one expects the Hessian normal modes to behave as independent harmonic oscillators with an effective mass M, set by the typical microscopic time scale of the spin dynamics. The associated oscillator frequencies then scale as $\omega_k = \sqrt{\lambda_k/M}$, with a spectral density $\rho(\omega) \sim cM^{3/2}\omega^2$. This correctly predicts the specific heat to scale as $c_V(T) \sim T^3$ at low T, rationalizing the result of [59] (with the notable exception of quantum particles at jamming where $\rho(\lambda) \sim \lambda^{-1/2}$ [70]). The typical mean-square displacement of oscillator mode k scales as $\langle x_k^2 \rangle \sim \hbar/M\omega_k \sim \hbar/\sqrt{M\lambda_k}$, and from the Lehmann representation, Eq. (18.3), one expects the spectral function to scale as

$$\chi''_{\rm loc}(\omega) \sim \rho(\omega) \langle x^2 \rangle_{\omega} \sim c\sqrt{M}\omega$$
, (18.15)

which rationalizes the linear scaling in Eq. (18.13).

The random oscillator picture is further supported by the scaling of the ratio $\chi''_{loc}(\omega)/\omega$. Exact calculations for rotors [68] or a particle in a random environment [29] indeed obtained a ratio $\sim \sqrt{M}$. In the TFSK, an effective potential construction yields $c \sim \sqrt{\Gamma}/J^2$ for $\Gamma \ll J$, while one expects $M \sim 1/\Gamma$. This correctly predicts the Γ -independent ratio in Eq. (18.5).

For metallic environments, the oscillator dynamics is instead expected to be overdamped with a friction coefficient η proportional to the prefactor of $|\omega_n|\hat{Q}_{aa}$. In this case the characteristic frequency of mode k is given by the relaxation rate, $\omega_k \sim \lambda_k/\eta$, implying a mode density $\rho(\omega) \sim c\eta^{3/2}\omega^{1/2}$ with larger weight at low frequencies. Accordingly, one expects corrections $\delta c_V(T) \sim T^{3/2}$ to the specific heat, as indeed predicted in [65]. The typical oscillator displacement follows from the scaling $\hbar\omega_k \sim \lambda_k x_k^2$, that is, $\langle x_k^2 \rangle \sim \hbar/\eta$, independent of the mode. The spectral function should then scale

$$\chi''_{\rm loc}(\omega) \sim \langle x^2 \rangle_{\omega} \, \rho(\omega) \sim c\sqrt{\eta \omega} \,,$$
 (18.16)

which again correctly reproduces the result of the Landau theory given in Eq. (18.14), as well as the scaling with η .

Unfortunately, little is known about the collective modes in short-range glasses. Those may play an important role in electronic processes such as hopping transport in insulators, as potential glue for strong-coupling superconductors, or as inelastic scatterers affecting the electrical resistance in metallic glasses.

18.6. Tunneling and eigenstate localization

It is interesting to ask whether and how quantum tunneling allows one to explore the clusterized landscape of quantum models with a 1-step RSB pattern. Having initialized a quantum Ising glass in a deep energy valley, tunnelling might not take place, even in the thermodynamic limit, since the remaining discreteness of the eigenstates (within one valley) may prevent resonant intervalley coupling.

This problem was first analyzed for p > 2 models [71]. Computing the tunneling amplitude between valleys of equal internal energy E and equating it to the intra-valley level spacing determines a minimal transverse field $\Gamma_c(E)$, below which wavefunctions and thus quantum dynamics, remain valley-localized. This situation is globally non-ergodic, yet delocalized within one valley - an unusual situation, believed to be impossible in finite dimensional models of MBL. The delocalization threshold $\Gamma_c(E)$ was found to lie consistently below the threshold $\Gamma_d(E)$ where glassy dynamics sets in. Indeed, Γ_c only

senses the energy barriers in phase space, while dynamics can be exponentially trapped by entropic (*free* energy) barriers, in spite of percolation in the energy landscape.

At finite p, once quantum tunneling resonantly connects different valleys of energy E, the eigenstates hybridize all classical configurations having energy close to E, since the wavefunctions behave ergodically within a valley. This is not so in the strict limit $p \to \infty$ [72–74] where the random energy model results, and each valley contains only one single configuration. In this limit, only states in a very narrow energy shell hybridize, its width being the exponentially small tunneling between those states.

It was claimed that tunneling under barriers can be exploited for a quantum algorithm to efficiently find other valleys of equal internal energy [35].

18.7. Interplay of glassy order with localization phenomena

The interplay of glassy freezing with other quantum phenomena, especially with the localization of fermions or bosons, is a rich subject. Below, we discuss how (fermionic) electron glasses become metals, and how glasses of bosons may become superfluid.

18.7.1. Quantum electron glasses

The classical Coulomb glass is a disordered insulator with unscreened Coulomb interactions between localized electrons, realized in *e.g.* doped semiconductors [75–77]. A standard model is

$$\mathcal{H}_{Cb} = \sum_{i} \epsilon_{i} n_{i} + \sum_{i < j} \frac{n_{i} n_{j}}{r_{ij}}$$
(18.17)

with random energy ϵ_i on site *i* drawn from a box distribution [-W/2, W/2]. The local occupation number is $n_i = 0, 1$ and r_{ij} is the distance between sites *i* and *j*. The number of electrons is a fixed fraction of the sites, which are arranged on a lattice or at random.

The unscreened, long range 1/r interactions between localized electrons enforce the Efros-Shklovskii (ES) Coulomb gap in the single-particle density of states, $\rho(E)$. It has long been conjectured that the stability constraint $\rho(E) \leq \text{const.} \times E^{d-1}/e^{2d}$ (which assures that particle-hole recombinations do not lower the energy) is only marginally satisfied. A mean-field treatment with marginal full RSB predicts this critical property [78–80]. This phenomenon is directly analogous to the appearance of a linear density of local fields in the classical SK model, which is known to saturate similar stability bounds [81].

Upon increasing the density of dopants the hopping t_{ij} between the sites i becomes important and a term $\sum_{i < j} (t_{ij}c_j^{\dagger}c_i + h.c.)$ must be added to \mathcal{H}_{Cb} in Eq. (18.17). Single-particle eigenfunctions ψ_{α} now spread over several sites. When written in the basis of ψ_{α} , the interactions take the generic form $\sum_{\alpha,\beta,\gamma,\delta} U_{\alpha\beta\gamma\delta}c_{\alpha}^{\dagger}c_{\beta}^{\dagger}c_{\gamma}c_{\delta}$, which couples all wavefunctions ψ_{α} that overlap in real space. Close to the metal-insulator transition this term resembles the fermionic models reviewed in Sec. 18.9.

Eventually localization breaks down and the wavefunctions become extended [82]. At the same time screening sets in, the effective Coulomb interactions become short range and the density of states at the Fermi level becomes finite. Close to criticality

the growing Coulomb gap suppresses the occurrence of strong resonances for low-energy electrons, so that their wavefunctions concentrate less on rare paths. This results in a larger fractal dimension as compared to the ones of non-interacting critical wavefunctions [83, 84].

Already in the proximity of the insulating state one finds multiple solutions to Hartree-Fock equations. This may signal the emergence of metastability [85, 86], which then extends all the way into the insulating electron glass. Slow relaxation and aging in gated 2d electron gases in silicon [87, 88] at metallic densities was taken as an indication for electron glassy behavior. However, the existence of a genuine glass transition in d=3 remains debated even in the classical limit, as the random potential breaks a potential Ising symmetry so that only the replica symmetry is left to be broken. Numerical simulations have not fully clarified the situation [89, 90].

In Sec. 18.8.3.2 we will discuss the related case of fermionic atoms with photon-mediated long-range interactions. Interestingly, the latter allow for a genuine glass transition (with RSB) already in the metallic phase.

18.7.2. Superfluidity and glassy density order

Experiments that had suggested superflow in solid, but possibly amorphous Helium [91] had raised the interesting question whether and how (glassy) density order of bosons could coexist with the a priori competing superfluid order. For models with mean field-like, frustrated density interactions (such as e.g. realized in multimode cavities) and locally hopping bosons, it was indeed found that the two orders can coexist [92], while they try to avoid each other locally: if the local superfluid order parameter is high, the glassy density order is weak, and vice versa [93]. This phase is the bosonic analogue of the metallic glass of fermionic atoms discussed in Sec. 18.8.3.2. For a specific 3d model whose quantum Hamiltonian maps onto the Fokker-Planck equation of Brownian hard spheres, the existence of such a superglass phase could be inferred explicitly, drawing on the knowledge on the classical glass phase of hard spheres [94].

18.8. Experimental realizations of quantum glasses

Let us now review a few selected examples of quantum glasses in correlated materials and possible realizations in artificial structures involving light-matter coupling.

18.8.1. Quantum Ising spin glasses

For long, $\text{LiY}_{1-x}\text{Ho}_x\text{F}_4$ was considered to be "the" quantum spin-glass realization [95]. The rare earth magnetic ion Holmium (Ho) has an Ising doublet with a large magnetic moment of $5.4\mu_B$ as its crystal field ground state. As the content of Ho is decreased below $x \approx 44\%$, the material turns from a dipolar ferromagnet to a dipolar-interacting Ising spin glass. Quantum fluctuations are introduced by a transverse field H_t , which at second order in perturbation theory hybridizes the two states of the Ho ground-doublet, resulting in an effective transverse field for the Ising doublet $\Gamma \sim H_t^2/\Delta$, where Δ is the gap in the crystal field spectrum.

Initially, the variation of the susceptibility with Γ for x = 0.167 was interpreted in terms of a first order transition at low temperatures [96]. With decreasing Γ the linear susceptibility reflects a rather sudden slowdown of dynamics and the onset of low frequency 1/f noise. However, the non-linear susceptibility does not diverge, ruling out a continuous glass transition. A first order transition, instead, is at odds with the rule of thumb that two-body interactions usually entail continuous transitions. It was later argued [97] that no equilibrium glass transition can occur at large H_t , since the latter induces finite transverse moments whose internal longitudinal fields break the Ising symmetry explicitly. In the absence of a de Almeida-Thouless line, one could thus at best expect a regime of correlated spin clusters, whose size diverges as $\Gamma \to 0$. This might explain the flat maximum in the non-linear susceptibility as a function of Γ , and its decrease with T. However, to fully understand the dynamic response of $LiY_{1-x}Ho_xF_4$, it will be crucial to include the hyper-fine coupling to the Ho nuclear spin, especially in the quantum glassy low T regime [98].

Interestingly, the susceptibility at a given point (Γ, T) in the glassy regime depends strongly on the annealing protocol [36]. Quantum annealing results in significantly faster typical relaxation rates and larger $\chi'(\omega)$ than thermal annealing. Moreover, it results in a nearly Γ -independent dissipation $\chi''(\omega)$ at low ω , which was interpreted to signal a critical state. The authors suggested that faster response in quantum annealed samples implies more complete relaxation than under thermal annealing. However, the opposite conclusion seems equally possible: quantum annealing remains stuck closer to the marginal surface of the rugged energy landscape, leaving more fluctuating regions with small barriers and faster relaxation times. Instead the thermally annealed state, if the system managed to cross barriers, looks more "aged", being entrenched in deeper and more stable valleys, and thus displaying slower response.

18.8.2. Quantum Heisenberg spin glasses

Heisenberg quantum spin glasses attract a lot of interest because of the intriguing SYKlike spectral features they might display in their paramagnetic phase, as well as above a moderate frequency scale – provided the mean-field and large-M predictions survive for SU(2) spins in 3d. A prominent example is the cuprate $La_{2-x}Sr_xCuO_4$ [99] which was recently shown to host spin glass order at low temperatures, all the way up to optimal doping, albeit with strong short-range antiferromagnetic correlations. If critical spectral features as predicted for fully frustrated mean-field Heisenberg glasses also survive in such real materials, they might possibly be at the root of the linear temperature dependence of the resistivity often observed in this type of materials [100].

In heavy fermion compounds, such as e.g. $Y_{1-x}U_xPd_3$, the dilute, randomly positioned local moments of U couple to conduction electrons that mediate RKKY interactions. Such materials are promising candidates to find genuine metallic quantum glass phases and quantum glass transitions to a paramagnet, where Kondo screening of the moments dominates. A quantum spin glass phase and interesting non-Fermi liquid behavior at higher temperatures was indeed reported for 0.2 < x < 0.4 of the above compound [101]. For a more thorough review of quantum glass candidates and their magnetic properties we refer to [102].

18.8.3. Mean-field quantum glasses in optical cavities

Book Title

Random and long-range interactions among cold atoms can be generated via light-matter coupling in multi-mode cavities with sufficiently random mode functions [103, 104]. We distinguish realizations with non-intinerant and itinerant atoms, respectively.

18.8.3.1. Non-itinerant fermionic atoms

Laser-trapped, immobile atoms act as local TLS (Ising spins). They are coupled by an effective long range Ising interaction J_{ij} obtained from integrating out the photons of the enclosing optical cavity. In cavities with a large number of modes, the long-ranged J_{ij} are nearly random, owing to the complex structure of the mode amplitudes. Such systems effectively realize a TFSK model, whereby the interaction may assume a non-vanishing mean $[J_{ij}] = J_0$ [105], and can be tuned via the driving parameters of the optical cavity. Upon dialling up the average coupling J_0 a ferromagnetic component of the spin freezing may arise as in classical analogues. Here it essentially realizes Dicke's superradiant phase. Under off-equilibrium conditions, as introduced by the driving of the laser cavity and the leakage of photons, the (steady state) phase diagram remains robust, but the critical behavior is modified [106].

18.8.3.2. Itinerant fermionic atoms

If the atoms are able to move, an interesting interplay between amorphous charge order and atomic delocalization occurs. In contrast to the case of Coulomb interactions (Sec. 18.7.1), the interactions remain long-ranged across the glass and the localization transitions. This allows for a glassy density order to develop upon decreasing the kinetic energy of the fermions while they are still in a metallic phase. Glassy density order in turn generates increased local disorder which eventually localizes the fermions.

The instability of the ergodic metal to glassy order and the single particle localization threshold are well separated in a lattice model with one fermionic level per site and close to half-filling. Thus, a metallic glass intervenes between the insulator and the ergodic metal [107]. In contrast, for more dilute filling the transition between ergodic metal and glassy insulator becomes first order, and in an extended parameter regime the two phases can coexist. How one phase nucleates out of the less favorable one remains an interesting open question.

18.9. Fermion models and the Sachdev-Ye-Kitaev model

Kitaev [108] proposed to use the Majorana fermion system

$$\hat{\mathcal{H}}_{SYK} = \frac{1}{i^{q/2}} \sum_{i_i < i_2 < \dots < i_q}^{N} J_{i_i i_2 \dots i_q} \hat{\psi}_{i_1} \hat{\psi}_{i_2} \dots \hat{\psi}_{i_q} , \qquad (18.18)$$

with $\hat{\psi}_i = \hat{\psi}_i^{\dagger}$, $\{\hat{\psi}_i, \hat{\psi}_j\} = 2\delta_{ij}$, quenched random couplings with zero mean and $[J_{i_1...i_q}^2] = (q-1)!J^2/N^{q-1}$, and four fermion (q=4) interactions, as a toy model for near-extremal black holes. Being very similar to the fermionic representation of the

 $SU(M \to \infty)$ Heisenberg model, it has no glassy phase. The main reason to propose the connection with gravity is that it becomes approximately conformal in the infrared (dropping time-derivatives, low-frequency limit), with a time-reparametrization symmetry which is broken to SL(2,R), as expected in blackhole theories which develop a nearly AdS_2 background [100]. A "Schwarzian action" describes the cost of reparametrizations [109]. Moreover, there is non-zero entropy at $T \to 0$ (taken after $N \to \infty$) and the specific heat is linear in T – thermodynamic properties that are expected in the black hole context as well. Finally, this system is a "maximally chaotic/perfect scrambler" meaning that the bound $\lambda_T \leq 2\pi T/\hbar$ on the Lyapunov exponent (a consequence of the fluctuation-dissipation theorem [110]), defined from the exponential growth of out-of-time-order correlations [111], is saturated.

It was soon realized that Kitaev's proposal is very close to the Sachdev-Ye model, introduced and studied with the aim of describing non-Fermi liquid behavior in condensed matter systems hosting quantum spins (see Sec. 18.4.2.2). The above quenched disordered fermion system (with no glassy phase), subsequently called SYK, thus became a popular model in high-energy theory. Potential experimental realizations were proposed in disordered graphene flakes under strong magnetic fields [112, 113].

In order to make a connection with the formalism we described in the rest of the chapter, the relevant correlator in the SYK model is $Q(\tau) = -[\langle T\hat{\psi}_i(\tau)\hat{\psi}_j(0)\rangle]$ (the fermionic analogue of G_B of Sec. 18.4.2.2), which satisfies an equation like Eq. (18.2) in its replica-diagonal form (no need of off-diagonal terms here) with

$$G_0^{-1} = d_{\tau} , \qquad \qquad \Sigma(\tau) = J^2 Q^3(\tau) .$$
 (18.19)

Let us now comment on some interesting properties of the SYK model. Time-reparametrization symmetry also emerges in mean-field glass models, where sigma models for the reparametrizations were phenomenologically proposed (but not derived) [32]. Coupling two (or more) SYKs, popular in high-energy studies to mimic wormholes, is similar to coupling real replicas in disordered systems, a procedure that has long been used in the glass literature to access properties of the free-energy landscape [114]. Moreover, the relation of SYK with tensor models without quenched disorder [115], parallels the ideas of self-generated disorder put forward in the 90s to link p-spin models to the Mode-Coupling Theory for structural glasses [1]. In the SYK context, the trick is to use tensor models such that in a perturbative expansion only melon diagrams for the two-time function, and ladder diagrams for the four-time function, survive. With quenched disorder this structure occurs thanks to the average over randomness.

These results motivated numerous studies of other quantum spin glasses [116, 117], in particular those discussed in Sec. 18.4.1. Let us just mention here the analysis of chaos in the paramagnetic and marginal glassy phases of the spherical p-spin models [117]. Quantum fluctuations were found to make the paramagnetic phase less and the spin glass phase more chaotic. In the classical limit $\hbar \to 0$, a crossover from strong to weak chaos, as marked by a maximum in λ_T , arises well above T_d , concomitant with the onset of two-step slow relaxation.

Book Title

18.10. Conclusion and outlook

This short review on quantum glassy systems focuses on the discussion of their equilibrium properties. We have tried to tie connections between the different systems and models presented. On the analytic side, we mostly discussed results obtained in different mean-field limits, where the replica method and replica symmetry breaking schemes can be safely applied. The phenomenology of these disordered and glassy mean-field models is very rich and has a wide scope. Indeed, many applications beyond physics have been exploited, notably to optimization problems, but more recently fruitful connections with unexpected fields, like gravity, have been uncovered, too. As usual, finite-dimensional physically relevant problems are not amenable to an exact treatment and much of what is known about them is either phenomenological or numerical.

Putting this summary together we came across a number of open issues which would be interesting to study. We just mention three of them. Revisiting the quantum glass transitions with deconfining spinons, especially in insulating Heisenberg systems, should be within analytic reach. We have only superficially discussed the interplay between glassiness and localization. There are certainly many interesting open questions there: how and when does glassiness set in? How does a localized quantum glass melt into a metal? Finally, it would be welcome to establish deeper connections between SYK insights (such as the treatment of the reparametrization invariance) and glass physics.

References

- [1] J.-P. Bouchaud, L. F. Cugliandolo, J. Kurchan, and M. Mezard, *Spin-glasses and random fields*, vol. 12, chapter Out of equilibrium dynamics in spin-glasses and other glassy systems, p. 161. World Scientific, (1997).
- [2] L. Berthier and G. Biroli, Theoretical perspective on the glass transition and amorphous materials, Rev. Mod. Phys. 83, 587-645, (2011).
- [3] H. Westfahl, J. Schmalian, and P. G. Wolynes, Dynamical mean-field theory of quantum stripe glasses, *Phys. Rev. B.* **68**, 134203 (Oct, 2003).
- [4] D. S. Fisher, Random transverse field Ising spin chains, Phys. Rev. Lett. 69, 534-537, (1992).
- [5] H. Rieger and A. P. Young. Quantum spin glasses. In Complex Behaviour of Glassy Systems, vol. 492, Lecture Notes in Physics, p. 256. Springer Verlag, (1997).
- [6] F. Iglói and C. Monthus, Strong disorder RG approach of random systems, *Phys. Rep.* 412, 277–431, (2005).
- [7] P. W. Anderson, Absence of diffusion in certain random lattices, Phys. Rev. 109, 1492– 1505, (1958).
- [8] N. F. Mott and W. Twose, The theory of impurity conduction, Advances in Physics. 10, 107–163, (1961).
- [9] D. Basko, B. Altshuler, and I. Aleiner, Metal-insulator transition in a weakly interacting many-electron system with localized single-particle states, *Annals of Physics.* 321, 1126– 1205. (2006).
- [10] R. Nandkishore and D. A. Huse, Many body localization and thermalization in quantum statistical mechanics, Ann. Rev. Cond. Matt. Phys. 6, 15–38, (2015).
- [11] F. Alet and N. Laflorencie, Many-body localization: an introduction and selected topics, C. R. Physique. 19, 498–525, (2018).
- [12] D. A. Abanin, E. Altman, I. Bloch, and M. Serbyn, Colloquium: Many-body localization, thermalization, and entanglement, Rev. Mod. Phys. 91, 021001, (2019).

- [13] J. Imbrie, On many-body localization for quantum spin chains, J. Stat. Phys. 163, 998–1048, (2016).
- [14] W. de Roeck and F. Huveneers, Stability and instability towards delocalization in MBL systems, *Phys. Rev. B.* **95**, 155129, (2017).
- [15] W. de Roeck, F. Huveneers, M. Müller, and M. Schiulaz, Absence of many-body mobility edges, Phys. Rev. B. 93, 014203, (2016).
- [16] J. Šuntajs, J. Bonča, T. Prosen, and L. Vidmar, Quantum chaos challenges many-body localization, Phys. Rev. E. 102, 062144, (2020).
- [17] D. Sels, Markovian baths and quantum avalanches. (2021). URL https://doi.org/10. 48550/arXiv.2108.10796.
- [18] M. Kiefer-Emmanouilidis, R. Unanyan, M. Fleischhauer, and J. Sirker, Slow delocalization of particles in many-body localized phases, Phys. Rev. B. 103, 024203 (2021).
- [19] C. D. Dominicis, I. Kondor, and T. Temesvari, *Spin Glasses and Random Fields*, vol. 12, chapter Beyond the Sherrington-Kirkpatrick Model, p. 119. World Scientific, (1997).
- [20] P. Calabrese, F. H. L. Essler, and G. Mussardo, Introduction to 'quantum integrability in out of equilibrium systems', J. Stat. Mech. 2016, 064001, (2016).
- [21] C. Turner, A. Michailidis, and D. A. Abanin et al., Weak ergodicity breaking from quantum many-body scars, *Nature Phys.* 14, 745–749, (2018).
- [22] S. Moudgalya, B. A. Bernevig, and N. Regnault, Quantum many-body scars and Hilbert space fragmentation: a review of exact results, Rep. Prog. Phys. 85, 086501 (2022).
- [23] A. J. Leggett, S. Chakravarty, A. T. Dorsey, M. P. A. Fisher, A. Garg, and W. Zwerger, Dynamics of the dissipative two-state system, Rev. Mod. Phys. 59, 1–85, (1987).
- [24] M. Vojta, Impurity quantum phase transitions, Phil. Mag. 86, 1807, (2006).
- [25] L. F. Cugliandolo, D. R. Grempel, G. S. Lozano, H. Lozza, and C. A. da Silva Santos, Dissipative effects on quantum glassy systems, *Phys. Rev. B.* 66, 014444, (2002).
- [26] L. F. Cugliandolo, D. R. Grempel, G. S. Lozano, H. Lozza, and C. A. da Silva Santos, Effects of dissipation on disordered quantum spin models, *Phys. Rev. B.* 70, 024422, (2004).
- [27] L. F. Cugliandolo and G. S. Lozano, Quantum aging in mean-field models, Phys. Rev. Lett. 80, 4979, (1998).
- [28] L. F. Cugliandolo and G. S. Lozano, Real-time nonequilibrium dynamics of quantum glassy systems, Phys. Rev. B. 59, 915, (1999).
- [29] G. Biroli and L. F. Cugliandolo, Quantum Thouless-Anderson-Palmer equations for glassy systems, Phys. Rev. B. 64, 014206, (2001).
- [30] M. P. Kennett and C. Chamon, Time reparametrization group and the long time behavior in quantum glassy systems, *Phys. Rev. Lett.* 86, 1622, (2001).
- [31] G. Biroli and O. Parcollet, Out-of-equilibrium dynamics of a quantum Heisenberg spin glass, Phys. Rev. B. 65, 094414, (2002).
- [32] L. F. Cugliandolo and C. Chamon, Fluctuations in glassy systems, J. Stat. Mech. 2007, P07022, (2007).
- [33] L. F. Cugliandolo, Dissipative quantum disordered models, Int. J. Mod. Phys. B. 20, 2795, (2006).
- [34] T. Kadowaki and H. Nishimori, Quantum annealing in the transverse Ising model, Phys. Rev. E. 58, 5355, (1998).
- [35] C. L. Baldwin and C. R. Laumann, Quantum algorithm for energy matching in hard optimization problems, *Phys. Rev. B.* **97**, 224201, (2018).
- [36] J. Brooke, D. Bitko, T. F. Rosenbaum, and G. Aeppli, Quantum annealing of a disordered magnet, *Science.* **284**, 779, (1999).
- [37] T. Jörg, F. Krzakala, J. Kurchan, and A. J. Maggs, Simple glass models and their quantum annealing, Phys. Rev. Lett. 101, 147204, (2008).
- [38] V. Bapst, L. Foini, F. Krzakala, G. Semerjian, and F. Zamponi, The quantum adiabatic algorithm applied to random optimization problems: The quantum spin glass perspec-

- tive, Phys. Rep. **52**, 127, (2013).
- [39] V. Dobrosavljevic and D. Thirumalai, 1/p expansion for a p-spin interaction spin-glass model in a transverse field, Journal of Physics A: Mathematical and General. 23(15), L767-L774 (1990).
- [40] L. F. Cugliandolo, D. R. Grempel, and C. A. da Silva Santos, From second to first order transitions in a disordered quantum magnet, Phys. Rev. Lett. 85, 2589, (2000).
- [41] A. J. Bray and M. A. Moore, Replica theory of quantum spin-glasses, J. Phys. C. 13, L655, (1980).
- F. Krzakala, A. Rosso, G. Semerjian, and F. Zamponi, On the path integral representation for quantum spin models and its application to the quantum cavity method and to Monte Carlo simulations, *Phys. Rev. B.* **78**, 134428, (2008).
- L. F. Cugliandolo, T. Giamarchi, and P. L. Doussal, Dynamic compressibility and aging in wigner crystals and quantum glasses, Phys. Rev. Lett. 96, 217203, (1996)
- [44] T. Giamarchi and P. L. Doussal, Spin-glasses and random fields, vol. 12, chapter Statics and Dynamics of Disordered Elastic Systems, p. 321. World Scientific, (1997).
- [45] A. Georges, G. Kotliar, W. Krauth, and M. Rozenberg, Dynamical mean-field theory of strongly correlated fermion systems and the limit of infinite dimensions, Rev. Mod. Phys. 68, 13, (1996).
- [46] D. R. Grempel and M. J. Rozenberg, Dynamics of the infinite-range Ising spin-glass model in a transverse field, Phys. Rev. Lett. 81, 2550, (1998).
- [47] J. Miller and D. Huse, Zero-temperature critical behavior of the infinite-range quantum Ising spin glass, Phys. Rev. Lett. 70, 3147–3150, (1993).
- J. Ye, S. Sachdev, and N. Read, Solvable spin glass of quantum rotors, *Phys. Rev. Lett.* **70**, 4011, (1993).
- A. Andreanov and M. Müller, Long-range quantum Ising spin glasses at T=0: Gapless collective excitations and universality, Phys. Rev. Lett. 109, 177201, (2012).
- R. Monasson, R. Zecchina, S. Kirkpatrick, B. Selman, and L. Troyansky, Determining computational complexity from characteristic 'phase transitions', Nature. 400, 133–137, (1999)
- Y. Y. Goldschmidt, Solvable model of the quantum spin-glass in a transverse field, *Phys.* Rev. B. 41, 4858, (1990).
- T. M. Nieuwenhuizen and F. Ritort, Quantum phase transition in spin glasses with multi-spin interactions, *Physica A.* **250**, 89, (1998)
- L. F. Cugliandolo, D. R. Grempel, and C. A. da Silva Santos, Imaginary-time replica formalism study of a quantum spherical p-spin-glass model, Phys. Rev. B. 64, 014403, (2001).
- [54] L. F. Cugliandolo and J. Kurchan, Analytical solution of the off-equilibrium dynamics of a long-range spin-glass model, Phys. Rev. Lett. 71, 173, (1993).
- [55] D. R. Grempel and M. J. Rozenberg, Fluctuations in a quantum random Heisenberg paramagnet, Phys. Rev. Lett. 80, 389–392, (1998).
- S. Sachdev and J. Ye, Gapless spin-fluid ground state in a random quantum Heisenberg magnet, Phys. Rev. Lett. 70, 3339, (1993).
- A. Georges, O. Parcollet, and S. Sachdev, Mean field theory of a quantum Heisenberg spin glass, Phys. Rev. Lett. 85, 840, (2000).
- A. Georges, O. Parcollet, and S. Sachdev, Quantum fluctuations of a nearly critical Heisenberg spin glass, *Phys. Rev. B.* **63**, 134406, (2001).
- G. Schehr, T. Giamarchi, and P. L. Doussal, Specific heat of the quantum Bragg glass, EPL. 66, 538, (2004).
- G. Schehr, Low-temperature specific heat of some quantum mean-field glassy phases, Phys. Rev. B. 71, 184204, (2005).
- [61] O. Parcollet and A. Georges, Non-Fermi-liquid regime of a doped Mott insulator, Phys. Rev. B. 59, 5341, (1999).

- [62] M. Christos, F. M. Haehl, and S. Sachdev, Spin liquid to spin glass crossover in the random quantum Heisenberg magnet, Phys. Rev. B. 105, 085120, (2022).
- [63] L. Arrachea and M. J. Rozenberg, Dynamical response of quantum spin-glass models at T=0, Phys. Rev. Lett. 86, 5172–5175, (2001).
- [64] H. Shackleton, A. Wietek, A. Georges, and S. Sachdev, Quantum phase transition at nonzero doping in a random t-J model, *Phys. Rev. Lett.* **126**, 136602, (2021).
- [65] A. M. Sengupta and A. Georges, Non-Fermi-liquid behavior near a T=0 spin-glass transition, *Phys. Rev. B.* **52**, 10295, (1995).
- [66] N. Read and S. Sachdev, Landau theory of quantum spin glasses of rotors and ising spins, Phys. Rev. B. 52, 384, (1996).
- [67] T. Senthil, A. Vishwanath, L. Balents, S. Sachdev, and M. P. A. Fisher, Deconfined quantum critical points, *Science*. 303, 1490–1494, (2004).
- [68] N. Read, S. Sachdev, and J. Ye, Landau theory of quantum spin glasses of rotors and Ising spins, Phys. Rev. B. 52, 384–410, (1995).
- [69] S. Sachdev, N. Read, and R. Oppermann, Quantum field theory of metallic spin glasses, Phys. Rev. B. 52, 10286–10294, (1995).
- [70] S. Franz, T. Maimbourg, G. Parisi, and A. Scardicchio, Impact of jamming criticality on low-temperature anomalies in structural glasses, *Proc. Nat. Acad. Sc.* 116, 201820360 (2019).
- [71] C. L. Baldwin, C. R. Laumann, A. Pal, and A. Scardicchio, Clustering of nonergodic eigenstates in quantum spin glasses, *Phys. Rev. Lett.* 118, 127201, (2017).
- [72] C. L. Baldwin, C. R. Laumann, A. Pal, and A. Scardicchio, The many-body localized phase of the quantum random energy model, *Phys. Rev. B.* 93, 024202, (2016).
- [73] L. Faoro, M. Feigel'man, and L. Ioffe, Non-ergodic extended phase of the quantum random energy model, Annals of Physics. 409, 167916 (2019).
- [74] G. Biroli, D. Facoetti, M. Schiró, M. Tarzia, and P. Vivo, Out-of-equilibrium phase diagram of the quantum random energy model, *Phys. Rev. B.* 103, 014204, (2021).
- [75] A. L. Efros and B. I. Shklovskii, Coulomb gap and low temperature conductivity of disordered systems, J. Phys. C. 8, L49, (1975).
- [76] A. Vaknin, Z. Ovadyahu, and M. Pollak, Aging effects in an Anderson insulator, Phys. Rev. Lett. 84, 3402–3405 (Apr, 2000).
- [77] A. Vaknin, Z. Ovadyahu, and M. Pollak, Nonequilibrium field effect and memory in the electron glass, Phys. Rev. B. 65, 134208 (Mar, 2002).
- [78] M. Müller and L. B. Ioffe, Glass transition and the Coulomb gap in electron glasses, Phys. Rev. Lett. 93, 256403, (2004).
- [79] S. Pankov and V. Dobrosavljević, Nonlinear screening theory of the Coulomb glass, Phys. Rev. Lett. 94, 046402 (2005).
- [80] M. Müller and S. Pankov, Mean-field theory for the three-dimensional Coulomb glass, Phys. Rev. B. 75, 144201, (2007).
- [81] M. Müller and M. Wyart, Marginal stability in structural, spin and electron glasses, Ann. Rev. Cond. Matt. Phys. pp. 177–200, (2015).
- [82] F. Epperlein, M. Schreiber, and T. Vojta, Quantum Coulomb glass within a Hartree-Fock approximation, *Phys. Rev. B.* **56**, 5890–5896 (1997).
- [83] M. Amini, V. E. Kravtsov, and M. Müller, Multifractality and quantum-to-classical crossover in the Coulomb anomaly at the Mott–Anderson metal–insulator transition, New J. Phys. 16, 015022 (2014).
- [84] I.S. Burmistrov, I.V. Gornyi, and A.D. Mirlin, Multifractality at Anderson transitions with Coulomb interaction, *Phys. Rev. Lett.* 111, 066601, (2013).
- [85] A. A. Pastor and V. Dobrosavljević, Melting of the electron glass, Phys. Rev. Lett. 83, 4642–4645 (1999).
- [86] V. Dobrosavljević, D. Tanasković, and A. A. Pastor, Glassy behavior of electrons near metal-insulator transitions, Phys. Rev. Lett. 90, 016402 (2003).

- [87] S. Bogdanovich and D. Popović, Onset of glassy dynamics in a two-dimensional electron system in silicon, Phys. Rev. Lett. 88, 236401 (2002).
- [88] J. Jaroszyński and D. Popović, Nonequilibrium relaxations and aging effects in a twodimensional Coulomb glass, Phys. Rev. Lett. 99, 046405 (2007).
- [89] M. Goethe and M. Palassini, Phase diagram, correlation gap, and critical properties of the Coulomb glass, Phys. Rev. Lett. 103, 045702, (2009).
- [90] A. Barzegar, J. C. Andersen, M. Schechter, and H. G. Katzgraber, Numerical observation of a glassy phase in the three-dimensional Coulomb glass, *Phys. Rev. B.* 100, 104418, (2018).
- [91] E. Kim and M. H. W. Chan, Observation of superflow in solid helium, Science. 305 (5692), 1941–1944, (2004).
- [92] G. Carleo, M. Tarzia, and F. Zamponi, Bose-Einstein condensation in quantum glasses, Phys. Rev. Lett. 103, 215302, (2009).
- [93] X. Yu and M. Müller, Mean field theory of superglasses, Phys. Rev. B. 85, 104205 (2012).
- [94] G. Biroli, C. Chamon, and F. Zamponi, Theory of the superglass phase, Phys. Rev. B. 78, 224306, (2008).
- [95] W. Wu, B. Ellman, T. F. Rosenbaum, G. Aeppli, and D. H. Reich, From classical to quantum glass, Phys. Rev. Lett. 67, 2076, (1991).
- [96] W. Wu, D. Bitko, T. F. Rosenbaum, and G. Aeppli, Quenching of the nonlinear susceptibility at a T = 0 spin glass transition, Phys. Rev. Lett. 71, 1919, (1993).
- [97] M. Schechter and N. Laflorencie, Quantum spin glass and the dipolar interaction, Phys. Rev. Lett. 97, 137204, (2006).
- [98] M. Schechter and P. C. E. Stamp, Significance of the hyperfine interactions in the phase diagram of LiHo_xY_{1-x}F₄, Phys. Rev. Lett. 95, 267208, (2005).
- [99] M. Frachet et al, Hidden magnetism at the pseudogap critical point of a cuprate superconductor, Nat. Phys. 16, 1064–1068, (2020).
- [100] D. Chowdhury, A. Georges, O. Parcollet, and S. Sachdev, Sachdev-Ye-Kitaev models and beyond: A window into non-Fermi liquids, Rev. Mod. Phys. (2022).
- [101] D. A. Gajewski, R. Chau, and M. B. Maple, Metallurgical and low-temperature electronic properties of $M_{1-x}U_xPd_3$ (M = Sc, Y, La) alloys, *Phys. Rev. B.* **62**, 5496, (2000).
- [102] J. A. Mydosh, Spin glasses: Redux: An updated experimental/materials survey, Reports on Progress in Physics. 78(5), 052501-, (2015).
- [103] S. Gopalakrishnan, B. L. Lev, and P. M. Goldbart, Emergent crystallinity and frustration with Bose-Einstein condensates in multimode cavities, *Nature Phys.* 5, 845, (2009).
- [104] S. Gopalakrishnan, B. L. Lev, and P. M. Goldbart, Frustration and glassiness in spin models with cavity-mediated interactions, *Phys. Rev. Lett.* 107, 277201 (2011).
- [105] P. Strack and S. Sachdev, Dicke quantum spin glass of atoms and photons, Phys. Rev. Lett. 107, 277202 (2011).
- [106] M. Buchhold, P. Strack, S. Sachdev, and S. Diehl, Dicke-model quantum spin and photon glass in optical cavities: Nonequilibrium theory and experimental signatures, *Phys. Rev.* A. 87, 063622, (2013).
- [107] M. Müller, P. Strack, and S. Sachdev, Quantum charge glasses of itinerant fermions with cavity-mediated long-range interactions, *Phys. Rev. A.* 86, 023604 (2012).
- [108] A. Kitaev. A simple model of quantum holography in KITP strings seminar and entanglement program, (2015).
- [109] J. Maldacena and D. Stanford, Remarks on the Sachdev-Ye-Kitaev model, Phys. Rev. D 94, 106002, (2016).
- [110] S. Pappalardi, L. Foini, and J. Kurchan, Quantum bounds and fluctuation-dissipation relations, SciPost Phys. 12, 130, (2022).
- [111] J. Maldacena, S. H. Shenker, and D. Stanford, A bound on chaos, Journal of High Energy Physics. 2016, 106, (2016).
- [112] A. Chen, R. Ilan, F. de Juan, D. I. Pikulin, and M. Franz, Quantum holography in a

September 8, 2022 2:18

Book Title

- graphene flake with an irregular boundary, Phys. Rev. Lett. 121, 036403, (2018).
- [113] M. Brzezinska, Y. Guan, O. V. Yazyev, S. Sachdev, and A. Kruchkov. Engineering SYK interactions in disordered graphene flakes under realistic experimental conditions, (2022). URL https://arxiv.org/abs/2208.01032.
- [114] S. Franz and G. Parisi, Phase diagram of coupled glassy systems: A mean-field study, Phys. Rev. Lett. 79, 248, (1997).
- [115] I. R. Klebanov, F. Popov, and G. Tarnopolsky. TASI lectures on large N tensor models, TASI 2017, (2018).
- [116] T. Anous and F. M. Haehl, The quantum p-spin glass model: a user manual for holographers, J. Stat. Mech. 2021, 113101, (2021).
- [117] S. Bera, K. Y. V. Lokesh, and S. Banerjee, Quantum to classical crossover in many-body chaos in a glass, *Phys. Rev. Lett.* **128**, 115302, (2022).