

K_0 GROUPS OF NONCOMMUTATIVE $\mathbb{R}^{2n}$

REN GUAN

ABSTRACT. In this paper we show that the K_0 groups of noncommutative $\mathbb{R}^{2n}$ are $\mathbb{Z}$ for $\forall n \in \mathbb{N}^*$ and make an approach to the calculation of the smooth case, which will bring many new sequence problems relating to binomial numbers.

CONTENTS

1. Introduction	1
2. The nonsmooth case	4
3. The smooth case	7
3.1. $P_k(A^\infty(\mathbb{R}_\Theta^2))$ where $k \geq 0$.	9
3.2. $P_0(A^\infty(\mathbb{R}_\Theta^{2n}))$ where $n \geq 2$.	12
3.3. $P_k(A^\infty(\mathbb{R}_\Theta^4))$ where $k \geq 1$.	13
4. A representation of self-adjoint elements of $A^\infty(\mathbb{R}_\Theta^2)$	17
References	26

1. INTRODUCTION

For $n \in \mathbb{N}$, the noncommutative $\mathbb{R}^{2n}$, or noncommutative flat space-time, denoted by $A(\mathbb{R}_\Theta^{2n})$, which naturally arised from noncommutative field theory [4, 8, 10], are another important class of noncommutative differentiable manifolds besides noncommutative tori [4, 18]. When regarded as a subalgebra of the bounded operators $B(H)$ on a separable Hilbert space H , $A(\mathbb{R}_\Theta^{2n})$ can be described as a $*$ -algebra generated by $2n$ self-adjoint operators $x_1, x_2, \dots, x_{2n}$ satisfying

$$(1.1) \quad [x_p, x_q] = \begin{cases} -i\theta_{pq}, & 2|q, p = q - 1 \text{ or } 2|p, q = p - 1, \\ -i\theta_{pq} = 0, & \text{other cases,} \end{cases}$$

where $-\theta_{qp} = \theta_{pq} > 0$ for $2|q$ and $p = q - 1$, $1 \leq p, q \leq 2n$, $i = \sqrt{-1}$, $\Theta := \{\theta_{pq}\}_{1 \leq p, q \leq 2n}$ is a real skew-symmetric matrix. The relation (1.1) ensures that any product of $x_1, x_2, \dots, x_{2n}$ can be rearranged to a finite sum of the form

$$(1.2) \quad \sum a_{p_1, p_2, \dots, p_{2n}} x_1^{p_1} x_2^{p_2} \dots x_{2n}^{p_{2n}}$$

with $(p_1, p_2, \dots, p_{2n}) \in \mathbb{N}^{2n}$, $a_{p_1, p_2, \dots, p_{2n}} \in \mathbb{C}$ and we set $x_i^0 = I$, the identity operator of $B(H)$. For example,

$$(1.3) \quad \begin{aligned} x_4 x_2 x_3 x_1 &= x_2 x_1 x_4 x_3 = (x_1 x_2 + i\theta_{12})(x_3 x_4 + i\theta_{34}) \\ &= x_1 x_2 x_3 x_4 + i\theta_{34} x_1 x_2 + i\theta_{12} x_3 x_4 - \theta_{12} \theta_{34}. \end{aligned}$$

2020 *Mathematics Subject Classification.* 05A10, 19A49, 58B34.

Key words and phrases. self-adjoint elements, noncommutative $\mathbb{R}^{2n}$, projectors, K_0 groups.

Mimick the definition of the smooth noncommutative tori [6], the smooth version $A^\infty(\mathbb{R}_\Theta^{2n})$ of noncommutative $\mathbb{R}^{2n}$ is the C^* -algebra of formal series

$$(1.4) \quad \sum a_{p_1, p_2, \dots, p_{2n}} x_1^{p_1} x_2^{p_2} \dots x_{2n}^{p_{2n}}$$

where the coefficient function $\mathbb{N}^{2n} \ni (p_1, p_2, \dots, p_{2n}) \mapsto a_{p_1, p_2, \dots, p_{2n}} \in \mathbb{C}$ belongs to the Schwartz space $\mathcal{S}(\mathbb{N}^{2n})$, i.e., there is a constant C_r for every $r \geq 1$ such that

$$(1.5) \quad \sup_{(p_1, p_2, \dots, p_{2n}) \in \mathbb{N}^{2n}} \left(1 + \sum_{k=1}^{2n} p_k^2 \right)^r a_{p_1, p_2, \dots, p_{2n}} < C_r.$$

For a $*$ -algebra A , a *projector* (or *projection*) p of A is a matrix with entries in A and satisfies $p^2 = p = p^*$. We denote by $P(A)$ the set of projectors of A . For any two projectors $p, q \in P(A)$, define

$$(1.6) \quad p + q := \begin{pmatrix} p & 0 \\ 0 & q \end{pmatrix},$$

and call p, q *equivalent*, $p \sim q$, if there is a unitary $u \in M_n(A)$ for some suitable $n \in \mathbb{N}$ such that

$$(1.7) \quad \begin{pmatrix} p & 0 \\ 0 & 0 \end{pmatrix} = u \begin{pmatrix} q & 0 \\ 0 & 0 \end{pmatrix} u^*.$$

Then $P(A)/\sim$ forms a semigroup under the operation (1.6). The K_0 group $K_0(A)$ of A is defined as the Grothendieck group of the (direct sum) semigroup of isomorphism classes of finitely generated projective right modules over A [9, Section 3.2] [17], or equivalently, the Grothendieck group of $P(A)/\sim$. Computing the K_0 groups of an algebra is a natural question, in general it's not easy, even for commutative one.

K_0 groups are important for noncommutative geometry, Exel shows that Morita equivalent algebras have isomorphic K_0 groups [7], and the famous Baum-Connes conjecture [2] is related to the K -theory groups $K_j(C_r^*(G))$ where $j = 0, 1$ of the reduced C^* -algebra $C_r^*(G)$ of a locally compact, Hausdorff and second countable group G . See [13, 14, 20] for significant development in Baum-Connes conjecture, which also provide examples of calculating K_0 groups, and [11] for the calculation of relative algebraic K -theory of a truncated polynomial algebra over a perfect field k of positive characteristic p , [12] for division algebras over local fields, etc.

In [19], Rieffel and Schwarz give a equivalent condition for Morita equivalence of two noncommutative tori, and Elliott and Hanfeng Li discuss the smooth case [6]. Notably, Pimsner and Voiculescu construct a hexagonal exact sequence linking the K -groups of an algebra A and of $A \rtimes_\alpha \mathbb{Z}$, the crossed products of A by $\mathbb{Z}$ [16]. By applying this result and considering the noncommutative tori $\mathbb{T}_\Theta^n$ as an iterated crossed product $\mathbb{C} \rtimes_{\alpha_1} \mathbb{Z} \rtimes_{\alpha_2} \mathbb{Z} \rtimes \dots \rtimes_{\alpha_n} \mathbb{Z}$ (cf. [9, Proposition 12.8]), Elliott shows that $K_0(\mathbb{T}_\Theta^n) = K_1(\mathbb{T}_\Theta^n) = \mathbb{Z}^{2n-1}$, which are independent of the choices of Θ .

In this paper, We first compute $K_0(A(\mathbb{R}_\Theta^{2n}))$, and then make an approach to the computation of the smooth cases $K_0(A^\infty(\mathbb{R}_\Theta^{2n}))$. By the definition of the K_0 group, we need to characterize the projectors of $A(\mathbb{R}_\Theta^{2n})$ and $A^\infty(\mathbb{R}_\Theta^{2n})$ under the unitary equivalence respectively. Our first main result is

Theorem 1.1. $K_0(A(\mathbb{R}_\Theta^{2n})) = \mathbb{Z}$ for any $n \in \mathbb{N}^*$ and Θ .

But for the smooth case $A^\infty(\mathbb{R}_\Theta^{2n})$, even the characterization of the projective elements of $A^\infty(\mathbb{R}_\Theta^{2n})$ is a highly nontrivial problem. In this paper we consider the classification of a special class of projectors and the following conclusion is of great importance.

Conjecture 1.2. *Let $\{b_n\}_{n \geq 0}$ be a complex number sequence such that for some integer $l \geq 1$, $b_n b_{n+l} = 0$ for $\forall n \in \mathbb{N}$ and define*

$$a_n := \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} b_k.$$

If $\lim_{n \rightarrow \infty} a_n = 0$, then $b_n = 0$ for all n .

Currently I can't prove it or construct a counterexample, so I state it as a conjecture. We have the following theorem, which is proved in section 3.

Theorem 1.3. *If Conjecture 1.2 is true and*

$$\mathcal{P} := \sum_{p,q=0}^{\infty} a_{p,q} x^p y^q \in P(A^\infty(\mathbb{R}_\Theta^2))$$

such that $a_{p,q} = 0$ for all pairs (p, q) satisfying $|p - q| > k$ for some $k \in \mathbb{N}$, then $\mathcal{P} = 0$ or 1.

Remark 1.4. *Conjecture 1.2 does not hold if we weaken the constraint $b_n b_{n+l} = 0$. The following counterexample is given by Noam D. Elkies: Let*

$$(1.8) \quad b_n = \frac{\operatorname{sgn}(\sin \frac{n\pi}{3})}{2^n},$$

then $b_n = 0$ when $3|n$, and correspondingly

$$(1.9) \quad a_n \ll \left(\frac{3}{4}\right)^{\frac{n}{2}}.$$

In fact, by applying Noam D. Elkies' method, for any $q \geq 3$, we can construct a sequence $\{b_n\}_{n \geq 0}$ such that $\lim_{n \rightarrow \infty} a_n = 0$ and $b_n = 0$ wherever $q|n$.

By applying the Fabry Gap Theorem [1], Alexandre Eremenko shows that $b_n = O(\epsilon^n)$ for $\forall \epsilon > 0$. And if $|a_n|$ tends to zero with geometric speed, that is $|a_n| = O(\delta^n)$ for some $\delta \in (0, 1)$, we can conclude that $b_n = 0$ for all n . See <https://mathoverflow.net/questions/425948/a-number-sequence-problem-involving-binomial-coefficients> for details.

We also consider the higher-dimensional analogues $P_k(A^\infty(\mathbb{R}_\Theta^{2n}))$ in section 3, see Definition 3.5. We first consider $P_0(A^\infty(\mathbb{R}_\Theta^4))$, then make some primary discussion to $P_k(A^\infty(\mathbb{R}_\Theta^4))$ and left the $n \geq 3$ cases for future. Naturally we conjecture that:

Conjecture 1.5. $P_k(A^\infty(\mathbb{R}_\Theta^{2n})) = \{0, 1\}$ for $\forall k \geq 0, \forall n \geq 1$ and Θ .

The ultimate goal is the characterization of $P(A^\infty(\mathbb{R}_\Theta^{2n}))$ where $n \in \mathbb{N}^*$, which will tell us what $K_0(A^\infty(\mathbb{R}_\Theta^{2n}))$ looks like, is also left for the future. Mimiking Theorem 1.1, we propose the following conjecture:

Conjecture 1.6. $K_0(A^\infty(\mathbb{R}_\Theta^{2n})) = \mathbb{Z}$ for $\forall n \in \mathbb{N}^*$ and Θ .

In section 4, we give a characterization of self-adjoint elements of $A^\infty(\mathbb{R}_\theta^2)$. If $\sum_{p,q=0}^\infty a_{p,q}x^py^q \in A^\infty(\mathbb{R}_\theta^2)$ is self-adjoint, then the imaginary parts of the coefficients $a_{p,q}$ can be expressed by their real parts, see Theorem 4.1 for details. Projectors are always self-adjoint, so we state here for future use. See [15] for the similar work for the pseudo-unitary group $\mathbf{U}(p, p)$.

Through the whole paper we set all terms containing $n!$ with $n < 0$ to 0. For example,

$$\frac{n!}{(n-m+p)!} \binom{m}{p} a^{m-p} x^p y^{n-m+p} := 0 \text{ if } n-m+p < 0.$$

And unless stated otherwise, $i = \sqrt{-1}$, $j, k, l, m, n, p, q, r, s, t \in \mathbb{N}$, $a_{p_1, p_2, \dots, p_{2n}} \in \mathbb{C}$, and $a_{p_1, p_2, \dots, p_{2n}} = 0$ if one of $p_1, p_2, \dots, p_{2n}$ is less than zero. $\Theta := \{\theta_{pq}\}_{1 \leq p, q \leq 2n}$ is a real skew-symmetric matrix where $-\theta_{qp} = \theta_{pq} > 0$ for $2|q$, $p = q-1$ and $\theta_{pq} = 0$ for other cases, $1 \leq p, q \leq 2n$. Although some symbols ($\mathfrak{a}, \mathfrak{b}, \mathcal{P}$, etc.) are repeatedly used, the readers won't be confused.

Acknowledgement. This research is partially supported by NSFC grants 12201255.

2. THE NONSMOOTH CASE

Let's start from $A(\mathbb{R}_\theta^2)$, which is generated by two self-adjoint operators x and y satisfying

$$(2.1) \quad [x, y] = -i\theta$$

for a real number $\theta \geq 0$. For convenience, in the following we denote $a := i\theta$, so (2.1) is equivalent to

$$(2.2) \quad yx = xy + a.$$

The $*$ -operation on an element of $A(\mathbb{R}_\theta^2)$ is taking its adjoint as an operator in $B(H)$. Every element $T \in A(\mathbb{R}_\theta^2)$ can be written as

$$(2.3) \quad T = \sum_{k=0}^n \sum_{p+q=k} a_{p,q} x^p y^q$$

for some integer $n \in \mathbb{N}$ and $a_{p,q} \in \mathbb{C}$ for all p and q . For example,

$$\begin{aligned} y^2 x &= y(xy + a) = (xy + a)y + ay = 2ay + xy^2, \\ y^2 x^2 &= (2ay + xy^2)x = 2a(xy + a) + x(2ay + xy^2) = 2a^2 + 4axy + x^2 y^2. \end{aligned}$$

The smallest n in (2.3) is called the *degree* of T , denoted by $\deg(T)$, which means for such n , at least one of the complex numbers $\{a_{0,n}, a_{1,n-1}, \dots, a_{n,0}\}$ is nonzero. And we call

$$(2.4) \quad \sigma(T) := \sum_{p+q=n} a_{p,q} x^p y^q$$

the *principle symbol* of T in (2.3). For example $\sigma(y^2 x) = xy^2$ and $\sigma(y^2 x^2) = x^2 y^2$. The adjoint of T is

$$(2.5) \quad T^* = \sum_{k=0}^n \sum_{p+q=k} \bar{a}_{p,q} y^q x^p,$$

to transform T^* into the standard form (2.3), we should represent $y^q x^p$ as a linear combination of $x^m y^n$'s where $m, n \in \mathbb{N}$, like the above example $y^2 x$ and $y^2 x^2$. First, we have

Lemma 2.1. For $n \in \mathbb{N}$,

$$(2.6) \quad y^n x = n a y^{n-1} + x y^n.$$

Proof. The case $n = 1$ follows directly from the definition. If (2.6) holds for some $n \in \mathbb{N}$, then

$$y^{n+1}x = y^n(xy + a) = ay^n + (nay^{n-1} + xy^n)y = (n+1)ay^n + xy^{n+1}$$

and the conclusion follows by induction on n . $\square$

For slightly more complex situations, when $n \geq 3$, Lemma 2.1 immediately implies

$$\begin{aligned} y^n x^2 &= (nay^{n-1} + xy^n)x \\ &= na((n-1)ay^{n-2} + xy^{n-1}) + x(nay^{n-1} + xy^n) \\ &= n(n-1)a^2y^{n-2} + 2naxy^{n-1} + x^2y^n, \\ y^n x^3 &= (n(n-1)a^2y^{n-2} + 2naxy^{n-1} + x^2y^n)x \\ &= n(n-1)a^2((n-2)ay^{n-3} + xy^{n-2}) + 2nax((n-1)ay^{n-2} + xy^{n-1}) \\ &\quad + x^2(nay^{n-1} + xy^n) \\ &= n(n-1)(n-2)a^3y^{n-3} + 3n(n-1)a^2xy^{n-2} + 3nax^2y^{n-1} + x^3y^n, \end{aligned}$$

if $n < 3$, say $n = 1$, then

$$yx^3 = (xy + a)x^2 = ax^2 + x(xy + a)x = 2ax^2 + x^2(xy + a) = 3ax^2 + x^3y,$$

the second formula for $y^n x^3$ still holds. In fact, we have

Theorem 2.2. For all $m, n \in \mathbb{N}$,

$$(2.7) \quad y^n x^m = \sum_{p=0}^m \frac{n!}{(n-m+p)!} \binom{m}{p} a^{m-p} x^p y^{n-m+p}.$$

And hence $\sigma(y^n x^m) = x^m y^n$.

Proof. Lemma 2.1 confirm the $m = 1$ case. If (2.7) holds for some m , then

$$\begin{aligned} y^n x^{m+1} &= \sum_{p=0}^m \frac{n!}{(n-m+p)!} \binom{m}{p} a^{m-p} x^p y^{n-m+p} x \\ &= \sum_{p=0}^m \frac{n!}{(n-m+p)!} \binom{m}{p} a^{m-p} x^p ((n-m+p)ay^{n-m-1+p} + xy^{n-m+p}) \\ &= \sum_{p=0}^m \frac{n!}{(n-m+p-1)!} \binom{m}{p} a^{m-p+1} x^p y^{n-m+p-1} + \sum_{p=0}^m \frac{n!}{(n-m+p)!} \binom{m}{p} a^{m-p} x^{p+1} y^{n-m+p} \\ &= \sum_{p=1}^m \frac{n!}{(n-m+p-1)!} \left(\binom{m}{p} + \binom{m}{p-1} \right) a^{m-p+1} x^p y^{n-m+p-1} + \frac{n!}{(n-m-1)!} a^{m+1} y^{n-m-1} \\ &\quad + x^{m+1} y^n \\ &= \sum_{p=1}^m \frac{n!}{(n-m+p-1)!} \binom{m+1}{p} a^{m-p+1} x^p y^{n-m+p-1} + \frac{n!}{(n-m-1)!} a^{m+1} y^{n-m-1} + x^{m+1} y^n \\ &= \sum_{p=0}^{m+1} \frac{n!}{(n-m-1+p)!} \binom{m+1}{p} a^{m+1-p} x^p y^{n-m-1+p}, \end{aligned}$$

i.e., (2.7) also holds for $m+1$, so according to mathematical induction (2.7) holds for all $m, n \in \mathbb{N}$. $\square$

For any projectors $p \in P(A(\mathbb{R}_\theta^2))$, we have $p^2 = p = p^*$, then by applying Theorem 2.2, compare the principle symbol of entries of p^2 and p , we can show that in fact $p \in P(\mathbb{C})$, i.e., we have

Theorem 2.3. *If $p \in P(A(\mathbb{R}_\theta^2))$, then $p \in P(\mathbb{C})$.*

Proof. Let

$$p = \begin{pmatrix} p_{1,1} & \cdots & p_{1,n} \\ \vdots & \ddots & \vdots \\ p_{n,1} & \cdots & p_{n,n} \end{pmatrix} \in M_n(A(\mathbb{R}_\theta^2))$$

be a n -dimensional projector of $A(\mathbb{R}_\theta^2)$. Then $p^2 = p = p^*$, and

$$\begin{aligned} \begin{pmatrix} p_{1,1} & \cdots & p_{1,n} \\ \vdots & \ddots & \vdots \\ p_{n,1} & \cdots & p_{n,n} \end{pmatrix} &= p = p^2 = pp^* \\ &= \begin{pmatrix} p_{1,1} & \cdots & p_{1,n} \\ \vdots & \ddots & \vdots \\ p_{n,1} & \cdots & p_{n,n} \end{pmatrix} \cdot \begin{pmatrix} p_{1,1}^* & \cdots & p_{n,1}^* \\ \vdots & \ddots & \vdots \\ p_{1,n}^* & \cdots & p_{n,n}^* \end{pmatrix} \\ &= \begin{pmatrix} \sum_{j=1}^n p_{1,j} p_{1,j}^* & \cdots & \sum_{j=1}^n p_{1,j} p_{n,j}^* \\ \vdots & \ddots & \vdots \\ \sum_{j=1}^n p_{n,j} p_{1,j}^* & \cdots & \sum_{j=1}^n p_{n,j} p_{n,j}^* \end{pmatrix}, \end{aligned}$$

so for $1 \leq k \leq n$,

$$(2.8) \quad p_{k,k} = \sum_{j=1}^n p_{k,j} p_{k,j}^*.$$

Let $\deg(p_{k,l}) = N_{k,l}$ and

$$(2.9) \quad p_{k,l} = \sum_{j=0}^{N_{k,l}} \sum_{\alpha+\beta=j} a_{\alpha,\beta}^{k,l} x^\alpha y^\beta$$

where all $a_{\alpha,\beta}^{k,l} \in \mathbb{C}$. Without loss of generality we assume $N_{k,1} \geq N_{k,l}$ for $1 \leq l \leq n$. Then by applying Theorem 2.2,

$$\begin{aligned} \sigma \left(\sum_{k=1}^n p_{k,l} p_{k,l}^* \right) &= \sigma \left(\sum_{k=1}^n \left(\sum_{j=0}^{N_{k,l}} \sum_{\alpha+\beta=j} a_{\alpha,\beta}^{k,l} x^\alpha y^\beta \right) \left(\sum_{j=0}^{N_{k,l}} \sum_{\gamma+\delta=j} \bar{a}_{\alpha,\beta}^{k,l} y^\beta x^\alpha \right) \right) \\ &= \sigma \left(\sum_{k=1}^n \left(\sum_{\alpha+\beta=N_{k,l}} a_{\alpha,\beta}^{k,l} x^\alpha y^\beta \right) \left(\sum_{\gamma+\delta=N_{k,l}} \bar{a}_{\alpha,\beta}^{k,l} y^\beta x^\alpha \right) \right) \\ &= \sum_{k=1}^n \sum_{\alpha+\beta=N_{k,l}} \sum_{\gamma+\delta=N_{k,l}} a_{\alpha,\beta}^{k,l} \bar{a}_{\gamma,\delta}^{k,l} x^{\alpha+\gamma} y^{\beta+\delta} \\ &= \sum_{k=1}^n \sum_{j=0}^{2N_{k,l}} \sum_{\alpha=0}^j a_{\alpha, N_{k,l}-\alpha}^{k,l} \bar{a}_{j-\alpha, N_{k,l}-j+\alpha}^{k,l} x^j y^{2N_{k,l}-j}. \end{aligned}$$

If $N_{k,1} \geq 1$ and $\sigma\left(\sum_{k=1}^n p_{k,l} p_{k,l}^*\right) \neq 0$, then

$$(2.10) \quad \sigma\left(\sum_{k=1}^n p_{k,l} p_{k,l}^*\right) = 2N_{k,1} > N_{k,1} = \sigma(p_{k,k}),$$

contradicts to (2.8), so we must have

$$(2.11) \quad \sum_{k=1}^n \sum_{j=0}^{2N_{k,l}} \sum_{\alpha=0}^j a_{\alpha, N_{k,l}-\alpha}^{k,l} \bar{a}_{j-\alpha, N_{k,l}-j+\alpha}^{k,l} x^j y^{2N_{k,l}-j} = \sigma\left(\sum_{k=1}^n p_{k,l} p_{k,l}^*\right) = 0.$$

For $1 \leq k \leq n$, the coefficient of $y^{2N_{k,l}}$ is

$$(2.12) \quad 0 = \sum_{N_{k,j}=N_{k,l}} a_{0, N_{k,j}}^{k,j} \bar{a}_{0, N_{k,j}}^{k,j} = \sum_{N_{k,j}=N_{k,l}} |a_{0, N_{k,j}}^{k,j}|^2,$$

so $a_{0, N_{k,l}}^{k,l} = 0$ for all $1 \leq k \leq n$; Then the coefficient of $x^2 y^{2N_{k,l}-2}$ is

$$(2.13) \quad 0 = \sum_{N_{k,j}=N_{k,l}} \left(a_{0, N_{k,j}}^{k,j} \bar{a}_{2, N_{k,j}-2}^{k,j} + a_{1, N_{k,j}-1}^{k,j} \bar{a}_{1, N_{k,j}-1}^{k,j} + a_{2, N_{k,j}-2}^{k,j} \bar{a}_{0, N_{k,j}}^{k,j} \right) = \sum_{N_{k,j}=N_{k,l}} |a_{1, N_{k,j}}^{k,j}|^2,$$

so $a_{1, N_{k,l}-1}^{k,l} = 0$ for all $1 \leq k \leq n$; Next consider the coefficient of $x^4 y^{2N_{k,l}-4}$, finally we have $a_{\alpha, N_{k,l}-\alpha}^{k,l} = 0$ for all $1 \leq k \leq n$ and $0 \leq \alpha \leq N_{k,l}$, but this contradicts to the assumption $N_{k,1} \geq 1$. So we must have $N_{k,1} = 0$ and hence $p_{k,l} \in \mathbb{C}$ for all $1 \leq k, l \leq n$, which implies $p \in P(\mathbb{C})$. $\square$

Then by the definition of K_0 groups, we have

Corollary 2.4. $K_0(A(\mathbb{R}_\theta^2)) = K_0(\mathbb{C}) = \mathbb{Z}$.

For $A(\mathbb{R}_\theta^{2n})$, $n \geq 2$, note that $x_3, x_4, \dots, x_{2n}$ commute with x_1 and x_2 , so for any $p \in P(A(\mathbb{R}_\theta^{2n}))$, we temporarily regard $x_3, x_4, \dots, x_{2n}$ as constants, then by applying the above method, we can also prove that $p \in P(\mathbb{C})$. Hence we also have

Corollary 2.5. If $p \in P(A(\mathbb{R}_\theta^{2n}))$, then $p \in P(\mathbb{C})$ and hence $K_0(A(\mathbb{R}_\theta^{2n})) = K_0(\mathbb{C}) = \mathbb{Z}$ for all $n \in \mathbb{N}$.

We have finished the proof of Theorem 1.1.

3. THE SMOOTH CASE

For $A^\infty(\mathbb{R}_\theta^2)$, the characterization of projectors is much more difficult, even the projective elements of $A^\infty(\mathbb{R}_\theta^2)$. Again, we start from the simplest case, $A^\infty(\mathbb{R}_\theta^2)$. Define

$$\gamma := \frac{\sqrt{2\theta}(1+i)}{2},$$

then $\gamma^2 = i\theta = a$, $\gamma = i\bar{\gamma}$. Any element $T \in A^\infty(\mathbb{R}_\theta^2)$ can be represented as

$$(3.1) \quad T = \sum_{p,q=0}^{\infty} a_{p,q} \gamma^{-p-q} x^p y^q \in A^\infty(\mathbb{R}_\theta^2)$$

where $\{a_{p,q} \gamma^{-p-q}\} \in \mathcal{S}(\mathbb{Z}^2)$ (We will see the advantage of the representation (3.1) later). We can no longer use the method in the proof of Theorem 2.3 to study $P(A^\infty(\mathbb{R}_\theta^2))$ because we cannot define principle symbol for the infinite sum (3.1). First we have the following theorem, which provides the conditions that $\{a_{p,q}\}$ must satisfied when (3.1) is a projector.

Theorem 3.1. *If*

$$(3.2) \quad T = \sum_{p,q=0}^{\infty} a_{p,q} \gamma^{-p-q} x^p y^q \in A^{\infty}(\mathbb{R}_{\theta}^2)$$

satisfies $T = TT^*$, *we have*

$$(3.3) \quad \begin{aligned} a_{m,n} &= i^{m+n} \sum_{h=0}^{\infty} (-1)^h \binom{m+h}{m} \frac{(n+h)!}{n!} \bar{a}_{m+h,n+h} \\ &= \sum_{r=0}^{\infty} \sum_{h=0}^r \sum_{s=0}^n \binom{r}{h} \frac{(n+h-s)!}{(n-s)!} a_{m+h-r,n+h-s} a_{r,s}. \end{aligned}$$

Proof. If $T = TT^*$, then $T^* = (TT^*)^* = TT^* = T$, so $T \in P(A^{\infty}(\mathbb{R}_{\theta}^2))$. And Theorem 2.2 implies

$$\begin{aligned} T = T^* &= \sum_{p,q=0}^{\infty} \bar{a}_{p,q} \bar{\gamma}^{-p-q} y^q x^p \\ &= \sum_{p,q=0}^{\infty} \bar{a}_{p,q} i^{p+q} \gamma^{-p-q} \sum_{h=0}^p \frac{q!}{(q-p+h)!} \binom{p}{h} a^{p-h} x^h y^{q-p+h}, \\ T = T^2 &= \sum_{p,q=0}^{\infty} a_{p,q} \gamma^{-p-q} x^p y^q \sum_{r,s=0}^{\infty} a_{r,s} \gamma^{-r-s} x^r y^s \\ &= \sum_{p,q=0}^{\infty} \sum_{r,s=0}^{\infty} a_{p,q} a_{r,s} \gamma^{-p-q-r-s} x^p y^q x^r y^s \\ &= \sum_{p,q=0}^{\infty} \sum_{r,s=0}^{\infty} a_{p,q} a_{r,s} \gamma^{-p-q-r-s} x^p \sum_{j=0}^r \frac{q!}{(q-r+j)!} \binom{r}{j} a^{r-j} x^j y^{q-r+j+s} \\ &= \sum_{p,q=0}^{\infty} \sum_{r,s=0}^{\infty} \sum_{j=0}^r \gamma^{r-p-q-s-2j} \frac{q!}{(q-r+j)!} \binom{r}{j} a_{p,q} a_{r,s} x^{p+j} y^{q+s-r+j}. \end{aligned}$$

For $m, n \in \mathbb{N}$, compare the coefficients of $x^m y^n$ on both sides, in the first part of TT^* the indices j, p, q, r, s should satisfy the following constrains:

$$(3.4) \quad \begin{cases} p+j = m \\ q+s-r+j = n, \end{cases}$$

then

$$(3.5) \quad \begin{cases} p = m-j \\ q = n+r-j-s, \end{cases}$$

so we have

$$\begin{aligned} a_{m,n} \gamma^{-m-n} &= \sum_{h=0}^{\infty} i^{m+n+2h} \binom{m+h}{m} \frac{(n+h)!}{n!} \bar{a}_{m+h,n+h} \gamma^{-m-n-2h} a^h \\ &= \sum_{h=0}^{\infty} i^{m+n+2h} \binom{m+h}{m} \frac{(n+h)!}{n!} \bar{a}_{m+h,n+h} \gamma^{-m-n}, \\ a_{m,n} \gamma^{-m-n} &= \sum_{r,s=0}^{\infty} \sum_{j=0}^{r-1} \frac{(n+r-j-s)!}{(n-s)!} \binom{r}{j} a_{m-j,n+r-j-s} a_{r,s} \gamma^{-m-n} \\ &= \sum_{r=0}^{\infty} \sum_{j=0}^r \sum_{s=0}^n \binom{r}{j} \frac{(n+r-j-s)!}{(n-s)!} a_{m-j,n+r-j-s} a_{r,s} \gamma^{-m-n} \end{aligned}$$

$$= \sum_{r=0}^{\infty} \sum_{h=0}^r \sum_{s=0}^n \binom{r}{h} \frac{(n+h-s)!}{(n-s)!} a_{m+h-r, n+h-s} a_{r,s} \gamma^{-m-n},$$

and hence

$$\begin{aligned} a_{m,n} &= i^{m+n} \sum_{h=0}^{\infty} (-1)^h \binom{m+h}{m} \frac{(n+h)!}{n!} \bar{a}_{m+h, n+h} \\ &= \sum_{r=0}^{\infty} \sum_{h=0}^r \sum_{s=0}^n \binom{r}{h} \frac{(n+h-s)!}{(n-s)!} a_{m+h-r, n+h-s} a_{r,s}. \end{aligned}$$

□

3.1. $P_k(A^\infty(\mathbb{R}_\theta^2))$ **where** $k \geq 0$. It's too general to find all series $\{a_{m,n}\}$ satisfying (3.3). Let's first consider a simple case, $a_{m,n} = 0$ where $m \neq n$. We will need the following well-known binomial formula:

$$(3.6) \quad \sum_{k=0}^m \binom{m}{k} \binom{n}{l-k} = \binom{m+n}{l}.$$

It can be proved by comparing the coefficient of z^l in the polynomial $(1+z)^m(1+z)^n = (1+z)^{m+n}$. We have

Proposition 3.2. *If*

$$(3.7) \quad S := \sum_{p=0}^{\infty} a_{p,p} \gamma^{-2p} x^p y^p \in P(A^\infty(\mathbb{R}_\theta^2)),$$

then $S = 0$ or 1 .

Proof. Define $b_p := a_{p,p}$, then from (3.3) we have

$$(3.8) \quad b_m = \sum_{r=0}^m \sum_{h=0}^r \binom{r}{h} \frac{(m+h-r)!}{(m-r)!} b_{m+h-r} b_r.$$

Let $\mathfrak{b}_m := m! b_m$, then

$$(3.9) \quad \mathfrak{b}_m = \sum_{r=0}^m \sum_{h=0}^r \frac{m!}{h!(r-h)!(m-r)!} \mathfrak{b}_{m+h-r} \mathfrak{b}_r$$

and for $m \in \mathbb{N}$,

$$\begin{aligned} & \sum_{p=0}^m \binom{m}{p} \mathfrak{b}_p \\ &= \sum_{p=0}^m \binom{m}{p} \sum_{r=0}^p \sum_{h=0}^r \frac{p!}{h!(r-h)!(p-r)!} \mathfrak{b}_{p+h-r} \mathfrak{b}_r \\ &= \sum_{p=0}^m \sum_{r=0}^p \sum_{h=0}^r \frac{m!}{(m-p)!(p-r)!(r-h)!h!} \mathfrak{b}_{p+h-r} \mathfrak{b}_r \\ &= \sum_{p=0}^m \sum_{q=0}^m \sum_{h=0}^m \frac{m!}{(m+h-p-q)!(p-h)!(q-h)!h!} \mathfrak{b}_p \mathfrak{b}_q \\ &= \sum_{p=0}^m \sum_{q=0}^m \binom{m}{p} \sum_{h=0}^m \binom{m-p}{q-h} \binom{p}{h} \mathfrak{b}_p \mathfrak{b}_q \\ &= \sum_{p=0}^m \sum_{q=0}^m \binom{m}{p} \binom{m}{q} \mathfrak{b}_p \mathfrak{b}_q \end{aligned}$$

$$= \left(\sum_{p=0}^m \binom{m}{p} \mathfrak{b}_p \right)^2,$$

hence

$$(3.10) \quad \sum_{p=0}^m \binom{m}{p} \mathfrak{b}_p = 0 \text{ or } 1.$$

When $m \geq 1$,

$$(3.11) \quad \mathfrak{b}_m = - \sum_{p=0}^{m-1} \binom{m}{p} \mathfrak{b}_p \text{ or } 1 - \sum_{p=0}^{m-1} \binom{m}{p} \mathfrak{b}_p,$$

for $m = 0$, (3.10) implies $\mathfrak{b}_0 = 0$ or 1 , then combine with (3.11) we can see that $\mathfrak{b}_m \in \mathbb{N}$ for every $m \in \mathbb{N}$. Also, (3.3) implies

$$(3.12) \quad b_m = (-1)^m \sum_{h=0}^{\infty} (-1)^h \binom{m+h}{m} \frac{(m+h)!}{m!} \bar{b}_{m+h},$$

so

$$(3.13) \quad \mathfrak{b}_m = (-1)^m \sum_{h=0}^{\infty} (-1)^h \binom{m+h}{m} \bar{\mathfrak{b}}_{m+h}.$$

Note that

$$(3.14) \quad \lim_{h \rightarrow \infty} \binom{m+h}{m} = 1 \text{ or } +\infty,$$

so the sum (3.13) should be finite, otherwise it won't convergent. So in fact $S \in P(A(\mathbb{R}_\theta^2))$, and then Theorem 2.3 implies $S = 0$ or 1 . $\square$

On the basis of Proposition 3.2 we can consider more general cases.

Definition 3.3. For $k \in \mathbb{N}$, we define

$$P_k(A^\infty(\mathbb{R}_\theta^2)) := \left\{ \sum_{p,q=0}^{\infty} a_{p,q} \gamma^{-p-q} x^p y^q \in P(A^\infty(\mathbb{R}_\theta^2)) : a_{p,q} = 0 \text{ for } |p-q| > k \right\}.$$

Obviously we have $P_k(A^\infty(\mathbb{R}_\theta^2)) \subseteq P_l(A^\infty(\mathbb{R}_\theta^2))$ for $k \leq l$. Proposition 3.2 shows that $P_0(A^\infty(\mathbb{R}_\theta^2)) = \{0, 1\}$. With the help of Conjecture 1.2, we can prove that

Theorem 3.4. $P_k(A^\infty(\mathbb{R}_\theta^2)) = \{0, 1\}$ for all $k \in \mathbb{N}$.

Proof. The case of $k = 0$ can be derived from Proposition 3.2. If $P_{k-1}(A^\infty(\mathbb{R}_\theta^2)) = \{0, 1\}$ for a positive k , let

$$\mathcal{P} := \sum_{p,q=0}^{\infty} a_{p,q} \gamma^{-p-q} x^p y^q \in P_k(A^\infty(\mathbb{R}_\theta^2)),$$

then Theorem 3.1 and the definition of $P_k(A^\infty(\mathbb{R}_\theta^2))$ imply that

$$\begin{aligned} 0 = a_{m,m+2k} &= \sum_{r=0}^{\infty} \sum_{h=0}^r \sum_{s=0}^{m+2k} \binom{r}{h} \frac{(m+2k+h-s)!}{(m+2k-s)!} a_{m+h-r, m+2k+h-s} a_{r,s} \\ &= \sum_{r=0}^{m+k} \sum_{h=0}^r \binom{r}{h} \frac{(m-r+k+h)!}{(m-r+k)!} a_{m+h-r, m+h-r+k} a_{r,r+k}, \end{aligned}$$

hence for $n \in \mathbb{N}$,

$$\begin{aligned}
0 &= \sum_{p=0}^n \frac{n!}{(n-p)!} a_{p,p+2k} \\
&= \sum_{p=0}^n \frac{n!}{(n-p)!} \sum_{r=0}^{p+k} \sum_{h=0}^r \binom{r}{h} \frac{(p-r+k+h)!}{(p-r+k)!} a_{p+h-r,p+h-r+k} a_{r,r+k} \\
&= \sum_{q=0}^n a_{q,q+k} \sum_{p=q}^n \sum_{r=0}^{p+k} \frac{n!}{(n-p)!} \binom{r}{q+r-p} \frac{(q+k)!}{(p+k-r)!} a_{r,r+k} \\
&= \sum_{q=0}^n a_{q,q+k} \sum_{r=0}^{n+k} \left(\sum_{p=q}^{q+r} \frac{n!}{(n-p)!} \binom{r}{q+r-p} \frac{(q+k)!}{(p+k-r)!} \right) a_{r,r+k} \\
&= \sum_{q=0}^n \frac{n!}{(n-q)!} a_{q,q+k} \sum_{r=0}^{n+k} \left(\sum_{p=0}^r \frac{(n-q)!}{(n-p-q)!} \binom{r}{r-p} \frac{(q+k)!}{(p+q+k-r)!} \right) a_{r,r+k} \\
&= \sum_{q=0}^n \frac{n!}{(n-q)!} a_{q,q+k} \sum_{r=0}^{n+k} r! \sum_{p=0}^r \binom{n-q}{p} \binom{q+k}{r-p} a_{r,r+k} \\
&= \sum_{q=0}^n \frac{n!}{(n-q)!} a_{q,q+k} \sum_{r=0}^{n+k} \frac{(n+k)!}{(n+k-r)!} a_{r,r+k}. \quad \text{Here we apply (3.6).}
\end{aligned}$$

Let $\mathbf{a}_q := q! a_{q,q+k}$ and

$$(3.15) \quad b_n := \sum_{q=0}^n \frac{n!}{(n-q)!} a_{q,q+k} = \sum_{q=0}^n \binom{n}{q} \mathbf{a}_q,$$

then $b_n b_{n+k} = 0$, and the binomial transform formula tells us that

$$(3.16) \quad \mathbf{a}_n = \sum_{q=0}^n (-1)^{n-q} \binom{n}{q} b_q.$$

Also for $n \in \mathbb{N}$, Theorem 3.1 implies

$$(3.17) \quad a_{n,n+k} = i^{2n+k} \sum_{h=0}^{\infty} (-1)^h \binom{n+h}{n} \frac{(n+k+h)!}{(n+k)!} \bar{a}_{n+h,n+k+h},$$

hence

$$(3.18) \quad \mathbf{a}_0 = a_{0,k} = i^k \sum_{h=0}^{\infty} (-1)^h \frac{(h+k)!}{k!} \bar{a}_{h,h+k} = i^k \sum_{h=0}^{\infty} (-1)^h \binom{h+k}{k} \bar{\mathbf{a}}_h,$$

which means

$$(3.19) \quad \lim_{h \rightarrow \infty} \binom{h+k}{k} \bar{\mathbf{a}}_h = 0,$$

and so we must have $\lim_{h \rightarrow \infty} \mathbf{a}_h = 0$. Then if Conjecture 1.2 is true, we have $b_n \equiv 0$ and (3.16) implies $\mathbf{a}_n \equiv 0$, $a_{n,n+k} \equiv 0$. Similarly we can also prove that $a_{n+k,n} \equiv 0$. Hence in fact $\mathcal{P} \in P_{k-1}(A^\infty(\mathbb{R}_\theta^2))$, so $P_k(A^\infty(\mathbb{R}_\theta^2)) = \{0, 1\}$ and by induction on k , $P_k(A^\infty(\mathbb{R}_\theta^2)) = \{0, 1\}$ for any $k \in \mathbb{N}$. $\square$

What we have done is just a beginning, we left the study of more general cases, or all solutions of (3.3) for future. If that was finished, we may be able to characterize $P(A^\infty(\mathbb{R}_\theta^2))$, which will tell us what $K_0(A^\infty(\mathbb{R}_\theta^2))$ is.

3.2. $P_0(A^\infty(\mathbb{R}_\Theta^{2n}))$ **where** $n \geq 2$. We extend Definition 3.3 to higher-dimensional case.

Definition 3.5. For $k, n \in \mathbb{N}$, we define

$$P_k(A^\infty(\mathbb{R}_\Theta^{2n})) := \left\{ \sum_{p_1, \dots, p_{2n}=0}^{\infty} a_{p_1, \dots, p_{2n}} \gamma_1^{-p_1-p_2} \dots \gamma_n^{-p_{2n-1}-p_{2n}} x_1^{p_1} \dots x_{2n}^{p_{2n}} \in P(A^\infty(\mathbb{R}_\Theta^{2n})) : a_{p_1, \dots, p_{2n}} = 0 \right. \\ \left. \text{where } \max_{1 \leq r, s \leq 2n} |p_r - p_s| > k. \right\}$$

where

$$\gamma_m := \frac{\sqrt{2\theta_{2m-1, 2m}}(1+i)}{2}, \quad m = 1, 2, \dots, n.$$

Now, let's consider a slightly more complicated example, $P_0(A^\infty(\mathbb{R}_\Theta^4))$. If

$$\mathcal{P} := \sum_{p=0}^{\infty} b_p \gamma_1^{-2p} \gamma_2^{-2p} x_1^p x_2^p x_3^p x_4^p := \sum_{p=0}^{\infty} a_{p,p,p,p} \gamma_1^{-2p} \gamma_2^{-2p} x_1^p x_2^p x_3^p x_4^p \in P_0(A^\infty(\mathbb{R}_\Theta^4)),$$

from (3.8) we have

$$\begin{aligned} b_m \gamma_2^{-2m} x_3^m x_4^m &= \sum_{r=0}^m \sum_{h=0}^r \binom{r}{h} \frac{(m+h-r)!}{(m-r)!} b_{m+h-r} \gamma_2^{-2(m+h-r)} b_r \gamma_2^{-2r} x_3^{m+h-r} x_4^{m+h-r} x_3^r x_4^r \\ &= \sum_{r=0}^m \sum_{h=0}^r \binom{r}{h} \frac{(m+h-r)!}{(m-r)!} b_{m+h-r} b_r \gamma_2^{-2(m+h)} \sum_{j=0}^r \binom{r}{j} \frac{(m+h-r)!}{(m+h-2r+j)!} \\ &\quad \times \gamma_2^{2r-2j} x_3^{m+h-r+j} x_4^{m+h-r+j}, \end{aligned}$$

then

$$(3.20) \quad b_m = \sum_{r=0}^m \sum_{h=0}^r \left(\binom{r}{h} \frac{(m+h-r)!}{(m-r)!} \right)^2 b_{m+h-r} b_r.$$

This time we can no longer conclude that $\{m!b_m\}_{m \geq 0}$ are all integers, for instance,

$$(3.21) \quad \{b_0, b_1, b_2, b_3, \dots\} = \{0, 1, -\frac{1}{8}(7 + \sqrt{33}), \frac{1}{72}(46 + 9\sqrt{33} - \sqrt{2089 + 360\sqrt{33}}), \dots\}$$

is a set of solution of (3.20). $\mathcal{P} = \mathcal{P}^*$ implies

$$\begin{aligned} b_m \gamma_2^{-2m} x_3^m x_4^m &= (-1)^m \sum_{h=0}^{\infty} (-1)^h \binom{m+h}{m} \frac{(m+h)!}{m!} \overline{b_{m+h} \gamma_2^{-2(m+h)}} x_4^{m+h} x_3^{m+h} \\ &= \sum_{h=0}^{\infty} \binom{m+h}{m} \frac{(m+h)!}{m!} \bar{b}_{m+h} \gamma_2^{-2(m+h)} \sum_{j=0}^{m+h} \binom{m+h}{j} \frac{(m+h)!}{j!} \gamma_2^{2(m+h-j)} x_3^j x_4^j, \end{aligned}$$

so

$$(3.22) \quad b_m = \sum_{h=0}^{\infty} \left(\binom{m+h}{m} \frac{(m+h)!}{m!} \right)^2 \bar{b}_{m+h}.$$

Like before, if we define $\mathbf{b}_{m,2} := (m!)^2 b_m$, then

$$(3.23) \quad \mathbf{b}_{m,2} = \sum_{r=0}^m \sum_{h=0}^r \left(\frac{m!}{(m-r)!(r-h)!h!} \right)^2 \mathbf{b}_{m+h-r,2} \mathbf{b}_{r,2} = \sum_{h=0}^{\infty} \binom{m+h}{m}^2 \bar{\mathbf{b}}_{m+h,2}, \quad \forall m \in \mathbb{N}.$$

Similarly, every element of $P_0(A^\infty(\mathbb{R}_\Theta^{2n}))$ corresponds to a sequence $\{\mathbf{b}_{m,n}\}_{m \geq 0}$ satisfying

$$(3.24) \quad \mathbf{b}_{m,n} = \sum_{r=0}^m \sum_{h=0}^r \left(\frac{m!}{(m-r)!(r-h)!h!} \right)^n \mathbf{b}_{m+h-r,n} \mathbf{b}_{r,n} = \sum_{h=0}^{\infty} (-1)^{(m+h)n} \binom{m+h}{m}^n \bar{\mathbf{b}}_{m+h,n}, \quad \forall m \in \mathbb{N}.$$

It's quite possible that $P_0(A^\infty(\mathbb{R}_\Theta^{2n})) = \{0, 1\}$ for $\forall n \in \mathbb{N}$, so we propose the following conjecture.

Conjecture 3.6. *Let $n \in \mathbb{N}^*$ and if a sequence $\{\mathfrak{b}_{m,n}\}_{m \geq 0}$ satisfies (3.24), then*

$$\mathfrak{b}_{m,n} = \begin{cases} 0 \text{ or } 1, & m = 0, \\ 0, & m \geq 1, \end{cases}$$

which means $P_0(A^\infty(\mathbb{R}_\Theta^{2n})) = \{0, 1\}$ for $\forall n \in \mathbb{N}^*$.

3.3. $P_k(A^\infty(\mathbb{R}_\Theta^4))$ **where** $k \geq 1$. For the $k \geq 1$ case, as a warm-up, we first consider $P_1(A^\infty(\mathbb{R}_\Theta^4))$.

If

$$\mathcal{P} := \sum_{p_1, p_2, p_3, p_4=0}^{\infty} a_{p_1, p_2, p_3, p_4} \gamma_1^{-p_1-p_2} \gamma_2^{-p_3-p_4} x_1^{p_1} x_2^{p_2} x_3^{p_3} x_4^{p_4} \in P_1(A^\infty(\mathbb{R}_\Theta^4)),$$

rewrite $\mathcal{P}$ to

$$(3.25) \quad \sum_{p_1, p_2=0}^{\infty} \left(\sum_{p_3, p_4=0}^{\infty} a_{p_1, p_2, p_3, p_4} \gamma_2^{-p_3-p_4} x_3^{p_3} x_4^{p_4} \right) \gamma_1^{-p_1-p_2} x_1^{p_1} x_2^{p_2}$$

and set

$$(3.26) \quad \alpha_{p_1, p_2} := \sum_{p_3, p_4=0}^{\infty} a_{p_1, p_2, p_3, p_4} \gamma_2^{-p_3-p_4} x_3^{p_3} x_4^{p_4}, \quad \beta_n := \sum_{m=0}^n \frac{n!}{(n-p)!} \alpha_{p, p+1},$$

then from the proof of Theorem 3.4 we have $\beta_n \beta_{n+1} = 0$. Note that

$$(3.27) \quad \alpha_{m, m+1} = \sum_{p_3, p_4=0}^{\infty} a_{m, m+1, p_3, p_4} \gamma_2^{-p_3-p_4} x_3^{p_3} x_4^{p_4} = \sum_{r=0}^1 \sum_{s=0}^1 a_{m, m+1, m+r, m+s} \gamma_2^{-2m-r-s} x_3^{m+r} x_4^{m+s},$$

so

$$\begin{aligned} 0 &= \beta_n \beta_{n+1} \\ &= \sum_{p=0}^n \frac{n!}{(n-p)!} \sum_{r=0}^1 \sum_{s=0}^1 a_{p, p+1, p+r, p+s} \gamma_2^{-2p-r-s} x_3^{p+r} x_4^{p+s} \sum_{q=0}^{n+1} \frac{(n+1)!}{(n+1-q)!} \sum_{r=0}^1 \sum_{s=0}^1 a_{q, q+1, q+r, q+s} \\ &\quad \times \gamma_2^{-2q-r-s} x_3^{q+r} x_4^{q+s} \\ &= \sum_{p=0}^n \sum_{q=0}^{n+1} \sum_{r=0}^1 \sum_{s=0}^1 \sum_{u=0}^1 \sum_{v=0}^1 \frac{n!(n+1)!}{(n-p)!(n+1-q)!} a_{p, p+1, p+r, p+s} a_{q, q+1, q+u, q+v} \gamma_2^{-2p-2q-r-s-u-v} \\ &\quad \times x_3^{p+r} x_4^{p+s} x_3^{q+u} x_4^{q+v} \\ &= \sum_{p=0}^n \sum_{q=0}^{n+1} \sum_{r=0}^1 \sum_{s=0}^1 \sum_{u=0}^1 \sum_{v=0}^1 \frac{n!(n+1)!}{(n-p)!(n+1-q)!} a_{p, p+1, p+r, p+s} a_{q, q+1, q+u, q+v} \gamma_2^{-2p-2q-r-s-u-v} \\ &\quad \times \sum_{h=0}^{q+u} \binom{q+u}{h} \frac{(p+s)!}{(p+s-q-u+h)!} \gamma_2^{2q+2u-2h} x_3^{p+r+h} x_4^{p+s-u+v+h}, \end{aligned}$$

consider the coefficient of $x_3^m x_4^{m+2}$ where $m \in \mathbb{N}$, we have

$$(3.28) \quad \sum_{p=0}^m \sum_{q=m-p}^{m+1} \frac{n!(n+1)!}{(n-p)!(n+1-q)!} \binom{q}{m-p} \frac{(p+1)!}{(m+1-q)!} a_{p, p+1, p, p+1} a_{q, q+1, q, q+1} = 0.$$

Let $p = q = m/2$, (3.36) implies $a_{p, p+1, p, p+1}^2 = 0$, so $a_{p, p+1, p, p+1} = 0$, $\forall p \in \mathbb{N}$. Similarity, by considering the coefficients of $x_3^{m+2} x_4^m$ we have $a_{p, p+1, p+1, p} = 0$ for $\forall p \in \mathbb{N}$, and then

$$0 = \beta_n \beta_{n+1}$$

$$= \sum_{p=0}^n \sum_{q=0}^{n+1} \sum_{r=0}^1 \sum_{u=0}^1 \frac{n!(n+1)!}{(n-p)!(n+1-q)!} a_{p,p+1,p+r,p+r} a_{q,q+1,q+u,q+u} \sum_{h=0}^{q+u} \binom{q+u}{h} \frac{(p+r)!}{(p+r-q-u+h)!} \\ \times \gamma_2^{-2p-2r-2h} x_3^{p+r+h} x_4^{p+r+h}.$$

Then consider the coefficients of $x_3^m x_4^m$, we have

$$(3.29) \quad \sum_{p=0}^m \sum_{q=0}^{m+k} \frac{n!(n+1)!}{(n-p)!(n+1-q)!} \sum_{r=0}^1 \sum_{u=0}^1 \binom{q+u}{m-p-r} \frac{(p+r)!}{(m-q-u)!} \\ \times a_{p,p+1,p+r,p+r} a_{q,q+1,q+u,q+u} = 0, \quad \forall n \in \mathbb{N},$$

Let $p = q = m/2 - 1$, then $a_{p,p+1,p+1,p+1}^2 = 0$, so $a_{p,p+1,p+1,p+1} = 0$, $\forall p \in \mathbb{N}$. And next (3.29) implies $a_{p,p+1,p,p} = 0$ for $\forall p \in \mathbb{N}$. Similarly we can prove $a_{p+1,p,p,p} = a_{p+1,p,p+1,p} = a_{p+1,p,p,p+1} = a_{p+1,p,p+1,p+1} = 0$ for $\forall p \in \mathbb{N}$. Then from (3.8) we have

$$(3.30) \quad \alpha_{m,m} = \sum_{r=0}^m \sum_{h=0}^r \binom{r}{h} \frac{(m+h-r)!}{(m-r)!} \alpha_{m+h-r,m+h-r} \alpha_{r,r} = \sum_{h=0}^{\infty} \binom{m+h}{m} \frac{(m+h)!}{m!} \alpha_{m+h,m+h}^*.$$

Consider the coefficient of $x_3^{2m+2} x_4^{2m+2}$, we have $0 = m! a_{m,m,m+1,m+1}^2$, so $a_{m,m,m+1,m+1} = 0$, then consider the coefficients of $x_3^{2m} x_4^{2m+2}$ and $x_3^{2m+2} x_4^{2m}$ we have $a_{m,m,m,m+1} = a_{m,m,m+1,m} = 0$, set $m = 0$ we have $a_{0,0,0,0} = a_{0,0,0,0}^2$, so $a_{0,0,0,0} = 0$ or 1 .

For $m \geq 1$, consider the coefficient of $x_3^{2m} x_4^{2m}$, we can get $a_{m,m,m,m} = 0$, then consider the coefficient of $x_3^{2m-2} x_4^{2m}$ and $x_3^{2m} x_4^{2m-2}$, we get that $a_{m,m,m-1,m} = a_{m,m,m,m-1} = 0$. Now (3.30) implies

$$a_{m,m,m-1,m-1} \gamma_2^{-2m+2} x_3^{m-1} x_4^{m-1} \\ = \sum_{r=1}^{m-1} \sum_{h=0}^r \binom{r}{h} \frac{(m+h-r)!}{(m-r)!} a_{m+h-r,m+h-r,m+h-r-1,m+h-r-1} a_{r,r,r-1,r-1} \gamma_2^{-2m-2h+4} \\ \times x_3^{m+h-r-1} x_4^{m+h-r-1} x_3^{r-1} x_4^{r-1} + 2a_{0,0,0,0} a_{m,m,m-1,m-1} \gamma_2^{-2m+2} x_3^{m-1} x_4^{m-1} + \sum_{h=1}^m \binom{m}{h} h! \\ \times a_{h,h,h-1,h-1} a_{m,m,m-1,m-1} \gamma_2^{-2m-2h+4} x_3^{h-1} x_4^{h-1} x_3^{m-1} x_4^{m-1} \\ = \sum_{r=1}^{m-1} \sum_{h=0}^r \binom{r}{h} \frac{(m+h-r)!}{(m-r)!} a_{m+h-r,m+h-r,m+h-r-1,m+h-r-1} a_{r,r,r-1,r-1} \gamma_2^{-2m-2h+4} \sum_{j=0}^{r-1} \binom{r-1}{j} \\ \times \frac{(m+h-r-1)!}{(m+h-2r+j)!} \gamma_2^{2r-2j-2} x_3^{m+h-r+j-1} x_4^{m+h-r+j-1} + 2a_{0,0,0,0} a_{m,m,m-1,m-1} \gamma_2^{-2m+2} x_3^{m-1} x_4^{m-1} \\ + \sum_{h=1}^m \binom{m}{h} h! a_{h,h,h-1,h-1} a_{m,m,m-1,m-1} \sum_{j=0}^{m-1} \binom{m-1}{j} \frac{(h-1)!}{(h-m+j)!} \gamma_2^{-2h-2j+2} x_3^{h+j-1} x_4^{h+j-1},$$

denote $\mathbf{a}_{m,2} := a_{m,m,m-1,m-1}$ for $m \geq 1$ and $\mathbf{a}_{0,2} := a_{0,0,0,0}$, then

$$(3.31) \quad \mathbf{a}_{m,2} = \sum_{r=1}^{m-1} \sum_{h=1}^r \binom{r}{h} \frac{(m+h-r)!}{(m-r)!} \binom{r-1}{h-1} \frac{(m+h-r-1)!}{(m-r)!} \mathbf{a}_{m+h-r,2} \mathbf{a}_{r,2} + 2\mathbf{a}_{0,2} \mathbf{a}_{m,2} \\ + \sum_{h=1}^m \frac{m!(m-1)!}{(m-h)!^2} \mathbf{a}_{h,2} \mathbf{a}_{m,2}, \quad \forall m \geq 1.$$

For $m \geq 1$, (3.30) also implies

$$\begin{aligned} \sum_{r,s=0}^{\infty} a_{m,m,m-1,m-1} \gamma_2^{-2m-2} x_3^{m-1} x_4^{m-1} &= i^{m+m} \sum_{h=0}^{\infty} (-1)^h \binom{m+h}{m} \frac{(m+h)!}{m!} \sum_{r,s=0}^{\infty} \bar{a}_{m+h,m+h,r,s} i^{r+s} \\ &\quad \times \gamma_2^{-r-s} x_4^s x_3^r \\ &= i^{m+m} \sum_{h=0}^{\infty} (-1)^h \binom{m+h}{m} \frac{(m+h)!}{m!} \sum_{r,s=0}^{\infty} \bar{a}_{m+h,m+h,r,s} i^{r+s} \\ &\quad \times \gamma_2^{-r-s} \sum_{j=0}^r \binom{r}{j} \frac{s!}{(s-r+j)!} \gamma_2^{2r-2j} x_3^j x_4^{s-r+j}, \end{aligned}$$

hence

$$(3.32) \quad \mathbf{a}_{m,2} = \begin{cases} -\sum_{h=0}^{\infty} \binom{m+h}{m} \frac{(m+h)!}{m!} \binom{m+h-1}{m-1} \frac{(m+h-1)!}{(m-1)!} \bar{\mathbf{a}}_{m+h,2}, & m \geq 1, \\ \sum_{h=0}^{\infty} h!^2 \bar{\mathbf{a}}_{h,2}, & m = 0. \end{cases}$$

A sequence $\{\mathbf{a}_{m,2}\}_{m \geq 0}$ satisfying (3.31) is

$$0, 0, 0, \frac{1}{12}, -\frac{1}{288} \left(23 + \sqrt{241} \right), \frac{341 + 20\sqrt{241} - \sqrt{60281 + 4040\sqrt{241}}}{5760}, \dots$$

but currently I don't find a sequence satisfying (3.31) and (3.32) simultaneously except zero sequence. So we propose the following conjecture.

Conjecture 3.7. *There is only zero sequence which satisfies (3.31) and (3.32) simultaneously.*

If Conjecture 3.7 is true, then $\mathcal{P} \in P_0(A^\infty(\mathbb{R}_\Theta^4))$, which means $P_1(A^\infty(\mathbb{R}_\Theta^4)) = P_0(A^\infty(\mathbb{R}_\Theta^4))$. For $P_k(A^\infty(\mathbb{R}_\Theta^4))$ where $k \geq 2$, we can do similarly. If

$$\mathcal{P} := \sum_{p_1, p_2, p_3, p_4=0}^{\infty} a_{p_1, p_2, p_3, p_4} \gamma_1^{-p_1-p_2} \gamma_2^{-p_3-p_4} x_1^{p_1} x_2^{p_2} x_3^{p_3} x_4^{p_4} \in P_k(A^\infty(\mathbb{R}_\Theta^4)),$$

again we rewrite $\mathcal{P}$ to

$$(3.33) \quad \sum_{p_1, p_2=0}^{\infty} \left(\sum_{p_3, p_4=0}^{\infty} a_{p_1, p_2, p_3, p_4} \gamma_2^{-p_3-p_4} x_3^{p_3} x_4^{p_4} \right) \gamma_1^{-p_1-p_2} x_1^{p_1} x_2^{p_2}$$

and set

$$(3.34) \quad \alpha_{p_1, p_2} := \sum_{p_3, p_4=0}^{\infty} a_{p_1, p_2, p_3, p_4} \gamma_2^{-p_3-p_4} x_3^{p_3} x_4^{p_4}, \quad \beta_n := \sum_{m=0}^n \frac{n!}{(n-p)!} \alpha_{p, p+k},$$

then from the proof of Theorem 3.4 we have $\beta_n \beta_{n+k} = 0$. Note that

$$(3.35) \quad \alpha_{m, m+k} = \sum_{p_3, p_4=0}^{\infty} a_{m, m+k, p_3, p_4} \gamma_2^{-p_3-p_4} x_3^{p_3} x_4^{p_4} = \sum_{r=0}^k \sum_{s=0}^k a_{m, m+k, m+r, m+s} \gamma_2^{-2m-r-s} x_3^{m+r} x_4^{m+s},$$

so

$$\begin{aligned} 0 &= \beta_n \beta_{n+k} \\ &= \sum_{p=0}^n \frac{n!}{(n-p)!} \sum_{r=0}^k \sum_{s=0}^k a_{p, p+k, p+r, p+s} \gamma_2^{-2p-r-s} x_3^{p+r} x_4^{p+s} \sum_{q=0}^{n+k} \frac{(n+k)!}{(n+k-q)!} \sum_{r=0}^k \sum_{s=0}^k a_{q, q+k, q+r, q+s} \\ &\quad \times \gamma_2^{-2q-r-s} x_3^{q+r} x_4^{q+s} \end{aligned}$$

$$\begin{aligned}
&= \sum_{p=0}^n \sum_{q=0}^{n+k} \sum_{r=0}^k \sum_{s=0}^k \sum_{u=0}^k \sum_{v=0}^k \frac{n!(n+k)!}{(n-p)!(n+k-q)!} a_{p,p+k,p+r,p+s} a_{q,q+k,q+u,q+v} \gamma_2^{-2p-2q-r-s-u-v} \\
&\quad \times x_3^{p+r} x_4^{p+s} x_3^{q+u} x_4^{q+v} \\
&= \sum_{p=0}^n \sum_{q=0}^{n+k} \sum_{r=0}^k \sum_{s=0}^k \sum_{u=0}^k \sum_{v=0}^k \frac{n!(n+k)!}{(n-p)!(n+k-q)!} a_{p,p+k,p+r,p+s} a_{q,q+k,q+u,q+v} \gamma_2^{-2p-2q-r-s-u-v} \\
&\quad \times \sum_{h=0}^{q+u} \binom{q+u}{h} \frac{(p+s)!}{(p+s-q-u+h)!} \gamma_2^{2q+2u-2h} x_3^{p+r+h} x_4^{p+s-u+v+h},
\end{aligned}$$

consider the coefficient of $x_3^m x_4^{m+2k}$ where $m \in \mathbb{N}$, we have

$$(3.36) \quad \sum_{p=0}^m \sum_{q=m-p}^{m+k} \frac{n!(n+k)!}{(n-p)!(n+k-q)!} \binom{q}{m-p} \frac{(p+k)!}{(m+k-q)!} a_{p,p+k,p,p+k} a_{q,q+k,q,q+k} = 0, \forall n \in \mathbb{N}.$$

Let $p = q = m/2$, (3.36) implies $a_{p,p+k,p,p+k}^2 = 0$, so $a_{p,p+k,p,p+k} = 0$, $\forall p \in \mathbb{N}$. Then consider the coefficients of $x_3^m x_4^{m+2k-2}$, we have

$$\begin{aligned}
(3.37) \quad &\sum_{p=0}^m \sum_{q=0}^{m+k} \frac{n!(n+k)!}{(n-p)!(n+k-q)!} \sum_{r=0}^1 \sum_{u=0}^1 \binom{q+u}{m-p-r} \frac{(p+r+k-1)!}{(m+k-q-u-1)!} \\
&\times a_{p,p+k,p+r,p+r+k-1} a_{q,q+k,q+u,q+u+k-1} = 0, \forall n \in \mathbb{N},
\end{aligned}$$

so

$$\begin{aligned}
(3.38) \quad &\sum_{r=0}^1 \sum_{u=0}^1 \binom{q+u}{m-p-r} \frac{(p+r+k-1)!}{(m+k-q-u-1)!} a_{p,p+k,p+r,p+r+k-1} a_{q,q+k,q+u,q+u+k-1} = 0, \forall p, q \in \mathbb{N}.
\end{aligned}$$

First we let $p = q$, $m = p + q + 2$, then $a_{p,p+k,p+1,p+k}^2 = 0$, so $a_{p,p+k,p+1,p+k} = 0$, $\forall p \in \mathbb{N}$, then (3.38) implies $a_{p,p+k,p,p+k-1} = 0$, $\forall p \in \mathbb{N}$. Next consider the coefficients of $x_3^m x_4^{m+2k-4}$, we have

$$\begin{aligned}
(3.39) \quad &\sum_{p=0}^m \sum_{q=0}^{m+k} \frac{n!(n+k)!}{(n-p)!(n+k-q)!} \sum_{r=0}^2 \sum_{u=0}^2 \binom{q+u}{m-p-r} \frac{(p+r+k-2)!}{(m+k-q-u-2)!} \\
&\times a_{p,p+k,p+r,p+r+k-2} a_{q,q+k,q+u,q+u+k-2} = 0, \forall n \in \mathbb{N},
\end{aligned}$$

so

$$\begin{aligned}
(3.40) \quad &\sum_{r=0}^2 \sum_{u=0}^2 \binom{q+u}{m-p-r} \frac{(p+r+k-2)!}{(m+k-q-u-2)!} a_{p,p+k,p+r,p+r+k-2} a_{q,q+k,q+u,q+u+k-2} = 0, \forall p, q \in \mathbb{N}.
\end{aligned}$$

Let $p = q$, $m = p + q + 4$, then $a_{p,p+k,p+2,p+k}^2 = 0$, so $a_{p,p+k,p+2,p+k} = 0$, $\forall p \in \mathbb{N}$, then (3.40) implies

$$\begin{aligned}
(3.41) \quad &\sum_{r=0}^1 \sum_{u=0}^1 \binom{q+u}{m-p-r} \frac{(p+r+k-2)!}{(m+k-q-u-2)!} a_{p,p+k,p+r,p+r+k-2} a_{q,q+k,q+u,q+u+k-2} = 0, \forall p, q \in \mathbb{N}.
\end{aligned}$$

Let $p = q$, $m = p + q + 2$, then $a_{p,p+k,p+1,p+k-1}^2 = 0$, so $a_{p,p+k,p+1,p+k-1} = 0$, $\forall p \in \mathbb{N}$, then (3.41) implies $a_{p,p+k,p,p+k-2} = 0$, $\forall p \in \mathbb{N}$.

Continue the above procedure, we can prove that $a_{p,p+k,u,v} = 0$ where $p \leq u \leq v \leq p + k$. Conversely, by considering the coefficients of $x_3^{m+2k-2j} x_4^m$, $j = 0, 1, \dots, k$, we have $a_{p,p+k,u,v} = 0$ where $p \leq v \leq u \leq p + k$, which means $a_{p,p+k,u,v} = 0$ where $p \leq u, v \leq p + k$. Similarly $a_{p+k,p,u,v} = 0$ where $p \leq u, v \leq p + k$. In fact we have partially proved the following conjecture:

Conjecture 3.8. *If $k \geq 1$ and*

$$\mathcal{P} := \sum_{p_1, p_2, p_3, p_4=0}^{\infty} a_{p_1, p_2, p_3, p_4} \gamma_1^{-p_1-p_2} \gamma_2^{-p_3-p_4} x_1^{p_1} x_2^{p_2} x_3^{p_3} x_4^{p_4} \in P_k(A^\infty(\mathbb{R}_\Theta^4)),$$

then $a_{p_1, p_2, p_3, p_4} = 0$ where $\max_{1 \leq r, s \leq 4} |p_r - p_s| = k$, which implies $P_k(A^\infty(\mathbb{R}_\Theta^4)) = P_{k-1}(A^\infty(\mathbb{R}_\Theta^4)) = \dots = P_0(A^\infty(\mathbb{R}_\Theta^4))$.

It's not hard to see that Conjecture 3.7 is included in Conjecture 3.8, and the latter involves a couple of number sequences satisfying conditions like (3.31) and (3.32). We leave the study of Conjecture 3.8 and $P_k(A^\infty(\mathbb{R}_\Theta^{2n}))$ where $k \geq 1$ and $n \geq 3$ for future and naturally, we propose Conjecture 1.5. Finally, the study of $P(A^\infty(\mathbb{R}_\Theta^{2n}))$ where $n \geq 2$, which are equivalent to the calculation of $K_0(A^\infty(\mathbb{R}_\Theta^{2n}))$ and seem much more difficult than $P(A^\infty(\mathbb{R}_\Theta^2))$, are left as Conjecture 1.6 for future study.

4. A REPRESENTATION OF SELF-ADJOINT ELEMENTS OF $A^\infty(\mathbb{R}_\theta^2)$

In this section we give a characterization of self-adjoint elements of $A^\infty(\mathbb{R}_\theta^2)$. The main result of this section is the following theorem.

Theorem 4.1. *Let $\{B_m\}_{m \geq 0}$ be the Bernoulli number series and for $n \in \mathbb{N}$,*

$$(4.1) \quad a_n := \frac{(-1)^n (2^{2n+2} - 1) B_{2n+2}}{n+1}.$$

Then if

$$(4.2) \quad T = \sum_{p, q=0}^{\infty} a_{p, q} x^p y^q \in A^\infty(\mathbb{R}_\theta^2)$$

is self-adjoint, we have

$$(4.3) \quad \text{Im} a_{p, q} = \sum_{k=0}^{\infty} a_k \binom{p+2k+1}{2k+1} \frac{(q+2k+1)!}{q!} \theta^{2k+1} \text{Re} a_{p+2k+1, q+2k+1}, \quad \forall p, q \in \mathbb{N}.$$

For instance, if we set $\text{Re} a_{p, q} = e^{-p^2 - q^2}$, we will get a self-adjoint element of $A^\infty(\mathbb{R}_\theta^2)$.

Maybe it's surprising that Bernoulli numbers appear here. First let's consider the nonsmooth case first. Every element $T \in A(\mathbb{R}_\theta^2)$ can be written as

$$(4.4) \quad T = \sum_{k=0}^n \sum_{p+q=k} a_{p, q} x^p y^q$$

for some $n \in \mathbb{N}$. If $T = T^*$, then (2.7) implies

$$(4.5) \quad \sum_{k=0}^n \sum_{p+q=k} a_{p, q} x^p y^q = \sum_{k=0}^n \sum_{p+q=k} \bar{a}_{p, q} y^q x^p = \sum_{k=0}^n \sum_{p+q=k} \bar{a}_{p, q} \sum_{h=0}^p \binom{p}{h} \frac{q!}{(q-p+h)!} a^{p-h} x^h y^{q-p+h},$$

compare the coefficients, we have

$$(4.6) \quad a_{p, q} = \sum_{h=0}^n \binom{p+h}{h} \frac{(q+h)!}{q!} a^h \bar{a}_{p+h, q+h}.$$

Here we set $a_{p,q} = 0$ if $p + q > n$. For $p + q > n - 2$, we have $a_{p,q} = \bar{a}_{p,q}$, which means $a_{p,q} \in \mathbb{R}$. And for $k \in \mathbb{N}$, (4.6) implies

$$(4.7) \quad a_{p-k,q-k} = \bar{a}_{p-k,q-k} + \sum_{h=1}^k \binom{p-k+h}{h} \frac{(q-k+h)!}{(q-k)!} a^h \bar{a}_{p-k+h,q-k+h},$$

$a = i\theta$ is a pure imaginary number, when $k = 2l$ is even, (4.7) implies

$$\begin{aligned} 0 &= \sum_{h=1}^l \binom{p-2l+2h}{2h} \frac{(q-2l+2h)!}{(q-2l)!} a^{2h} \text{Re} a_{p-2l+2h,q-2l+2h} \\ &\quad - i \sum_{h=1}^l \binom{p-2l+2h-1}{2h-1} \frac{(q-2l+2h-1)!}{(q-2l)!} a^{2h-1} \text{Im} a_{p-2l+2h-1,q-2l+2h-1}, \\ 2i \text{Im} a_{p-2l,q-2l} &= -i \sum_{h=1}^l \binom{p-2l+2h}{2h} \frac{(q-2l+2h)!}{(q-2l)!} a^{2h} \text{Im} a_{p-2l+2h,q-2l+2h} \\ &\quad + \sum_{h=1}^l \binom{p-2l+2h-1}{2h-1} \frac{(q-2l+2h-1)!}{(q-2l)!} a^{2h-1} \text{Re} a_{p-2l+2h-1,q-2l+2h-1}. \end{aligned}$$

When $k = 2l - 1$ is odd, (4.7) implies

$$\begin{aligned} 0 &= \sum_{h=1}^{l-1} \binom{p-2l+1+2h}{2h} \frac{(q-2l+1+2h)!}{(q-2l+1)!} a^{2h} \text{Re} a_{p-2l+1+2h,q-2l+1+2h} \\ &\quad - i \sum_{h=1}^l \binom{p-2l+2h}{2h-1} \frac{(q-2l+2h)!}{(q-2l+1)!} a^{2h-1} \text{Im} a_{p-2l+2h,q-2l+2h}, \\ 2i \text{Im} a_{p-2l+1,q-2l+1} &= -i \sum_{h=1}^{l-1} \binom{p-2l+1+2h}{2h} \frac{(q-2l+1+2h)!}{(q-2l+1)!} a^{2h} \text{Im} a_{p-2l+1+2h,q-2l+1+2h} \\ &\quad + \sum_{h=1}^l \binom{p-2l+2h}{2h-1} \frac{(q-2l+2h)!}{(q-2l+1)!} a^{2h-1} \text{Re} a_{p-2l+2h,q-2l+2h}. \end{aligned}$$

We compute $\text{Im} a_{p-k,q-k}$ for small k , from which we can find the general formula. First,

$$(4.8) \quad a_{p-1,q-1} = \bar{a}_{p-1,q-1} + \binom{p}{1} \frac{q!}{(q-1)!} a \bar{a}_{p,q} = \bar{a}_{p-1,q-1} + p q a a_{p,q},$$

which means

$$(4.9) \quad \text{Im} a_{p-1,q-1} = \frac{1}{2} p q \theta a_{p,q}.$$

Also,

$$\begin{aligned} a_{p-2,q-2} &= \bar{a}_{p-2,q-2} + \binom{p-1}{1} \frac{(q-1)!}{(q-2)!} a \bar{a}_{p-1,q-1} + \binom{p}{2} \frac{q!}{(q-2)!} a^2 \bar{a}_{p,q} \\ &= \bar{a}_{p-2,q-2} + (p-1)(q-1) a \bar{a}_{p-1,q-1} + \frac{1}{2} p(p-1)q(q-1) a^2 a_{p,q}, \end{aligned}$$

take real and imaginary part of the above equality, we have

$$\begin{aligned} (p-1)(q-1)\theta \text{Im} a_{p-1,q-1} - \frac{1}{2} p(p-1)q(q-1)\theta^2 a_{p,q} &= 0, \\ 2\text{Im} a_{p-2,q-2} - (p-1)(q-1)\theta \text{Re} a_{p-1,q-1} &= 0, \end{aligned}$$

which means

$$(4.10) \quad \text{Im} a_{p-2,q-2} = \frac{1}{2} (p-1)(q-1)\theta \text{Re} a_{p-1,q-1}.$$

Continue the above procedure,

(4.11)

$$a_{p-3,q-3} = \bar{a}_{p-3,q-3} + \binom{p-2}{1} \frac{(q-2)!}{(q-3)!} a \bar{a}_{p-2,q-2} + \binom{p-1}{2} \frac{(q-1)!}{(q-3)!} a^2 \bar{a}_{p-1,q-1} + \binom{p}{3} \frac{q!}{(q-3)!} a^3 \bar{a}_{p,q},$$

again, take real and imaginary part of the above equality, we have

$$(p-2)(q-2)\theta \text{Im} a_{p-2,q-2} - \binom{p-1}{2} \frac{(q-1)!}{(q-3)!} \theta^2 \text{Re} a_{p-1,q-1} = 0,$$

$$2\text{Im} a_{p-3,q-3} - (p-2)(q-2)\theta \text{Re} a_{p-2,q-2} - \binom{p-1}{2} \frac{(q-1)!}{(q-3)!} \theta^2 \text{Im} a_{p-1,q-1} + \binom{p}{3} \frac{q!}{(q-3)!} \theta^3 a_{p,q} = 0,$$

so

$$(4.12) \quad \text{Im} a_{p-3,q-3} = \frac{1}{2} \binom{p-2}{1} \frac{(q-2)!}{(q-3)!} \theta \text{Re} a_{p-2,q-2} + \frac{1}{4} \binom{p}{3} \frac{q!}{(q-3)!} \theta^3 a_{p,q}.$$

For $a_{p-4,q-4}$,

$$a_{p-4,q-4} = \bar{a}_{p-4,q-4} + \binom{p-3}{1} \frac{(q-3)!}{(q-4)!} a \bar{a}_{p-3,q-3} + \binom{p-2}{2} \frac{(q-2)!}{(q-4)!} a^2 \bar{a}_{p-2,q-2} \\ + \binom{p-1}{3} \frac{(q-1)!}{(q-4)!} a^3 \bar{a}_{p-1,q-1} + \binom{p}{4} \frac{q!}{(q-4)!} a^4 \bar{a}_{p,q},$$

then

$$\binom{p-3}{1} \frac{(q-3)!}{(q-4)!} \theta \text{Im} a_{p-3,q-3} - \binom{p-2}{2} \frac{(q-2)!}{(q-4)!} \theta^2 \text{Re} a_{p-2,q-2} \\ - \binom{p-1}{3} \frac{(q-1)!}{(q-4)!} \theta^3 \text{Im} a_{p-1,q-1} + \binom{p}{4} \frac{q!}{(q-4)!} \theta^4 \text{Re} a_{p,q} = 0,$$

$$2\text{Im} a_{p-4,q-4} - \binom{p-3}{1} \frac{(q-3)!}{(q-4)!} \theta \text{Re} a_{p-3,q-3} - \binom{p-2}{2} \frac{(q-2)!}{(q-4)!} \theta^2 \text{Im} a_{p-2,q-2} \\ + \binom{p-1}{3} \frac{(q-1)!}{(q-4)!} \theta^3 \text{Re} a_{p-1,q-1} = 0,$$

so

$$\text{Im} a_{p-4,q-4} = \frac{1}{2} \binom{p-3}{1} \frac{(q-3)!}{(q-4)!} \theta \text{Re} a_{p-3,q-3} + \frac{1}{2} \binom{p-2}{2} \frac{(q-2)!}{(q-4)!} \theta^2 \\ \times \frac{1}{2} (p-1)(q-1) \theta \text{Re} a_{p-1,q-1} - \frac{1}{2} \binom{p-1}{3} \frac{(q-1)!}{(q-4)!} \theta^3 \text{Re} a_{p-1,q-1} \\ = \frac{1}{2} \binom{p-3}{1} \frac{(q-3)!}{(q-4)!} \theta \text{Re} a_{p-3,q-3} + \frac{1}{4} \binom{p-1}{3} \frac{(q-1)!}{(q-4)!} \theta^3 \text{Re} a_{p-1,q-1}.$$

For $a_{p-5,q-5}$,

$$a_{p-5,q-5} = \bar{a}_{p-5,q-5} + \binom{p-4}{1} \frac{(q-4)!}{(q-5)!} a \bar{a}_{p-4,q-4} + \binom{p-3}{2} \frac{(q-3)!}{(q-5)!} a^2 \bar{a}_{p-3,q-3} \\ + \binom{p-2}{3} \frac{(q-2)!}{(q-5)!} a^3 \bar{a}_{p-2,q-2} + \binom{p-1}{4} \frac{(q-1)!}{(q-5)!} a^4 \bar{a}_{p-1,q-1} + \binom{p}{5} \frac{q!}{(q-5)!} a^5 \bar{a}_{p,q},$$

then

$$2\text{Im} a_{p-5,q-5} = \binom{p-4}{1} \frac{(q-4)!}{(q-5)!} \theta \text{Re} a_{p-4,q-4} + \binom{p-3}{2} \frac{(q-3)!}{(q-5)!} \theta^2 \text{Im} a_{p-3,q-3} \\ - \binom{p-2}{3} \frac{(q-2)!}{(q-5)!} \theta^3 \text{Re} a_{p-2,q-2} - \binom{p-1}{4} \frac{(q-1)!}{(q-5)!} \theta^4 \text{Im} a_{p-1,q-1} \\ + \binom{p}{5} \frac{q!}{(q-5)!} \theta^5 a_{p,q}$$

$$\begin{aligned}
&= \binom{p-4}{1} \frac{(q-4)!}{(q-5)!} \theta \text{Rea}_{p-4,q-4} + \binom{p-3}{2} \frac{(q-3)!}{(q-5)!} \theta^2 \left(\frac{1}{2} \binom{p-2}{1} \frac{(q-2)!}{(q-3)!} \right. \\
&\quad \times \theta \text{Rea}_{p-2,q-2} + \frac{1}{4} \binom{p}{3} \frac{q!}{(q-3)!} \theta^3 a_{p,q} \Big) - \binom{p-2}{3} \frac{(q-2)!}{(q-5)!} \theta^3 \text{Rea}_{p-2,q-2} \\
&\quad - \binom{p-1}{4} \frac{(q-1)!}{(q-5)!} \theta^4 \times \frac{1}{2} p q \theta a_{p,q} + \binom{p}{5} \frac{q!}{(q-5)!} \theta^5 a_{p,q} \\
&= \binom{p-4}{1} \frac{(q-4)!}{(q-5)!} \theta \text{Rea}_{p-4,q-4} + \frac{1}{2} \binom{p-2}{3} \frac{(q-2)!}{(q-5)!} \theta^3 \text{Rea}_{p-2,q-2} \\
&\quad + \binom{p}{5} \frac{q!}{(q-5)!} \theta^5 a_{p,q},
\end{aligned}$$

so

$$\text{Ima}_{p-5,q-5} = \frac{1}{2} \binom{p-4}{1} \frac{(q-4)!}{(q-5)!} \theta \text{Rea}_{p-4,q-4} + \frac{1}{4} \binom{p-2}{3} \frac{(q-2)!}{(q-5)!} \theta^3 \text{Rea}_{p-2,q-2} + \frac{1}{2} \binom{p}{5} \frac{q!}{(q-5)!} \theta^5 a_{p,q}.$$

For $a_{p-6,q-6}$,

$$\begin{aligned}
a_{p-6,q-6} &= \bar{a}_{p-6,q-6} + \binom{p-5}{1} \frac{(q-5)!}{(q-6)!} a \bar{a}_{p-5,q-5} + \binom{p-4}{2} \frac{(q-4)!}{(q-6)!} a^2 \bar{a}_{p-4,q-4} \\
&\quad + \binom{p-3}{3} \frac{(q-3)!}{(q-6)!} a^3 \bar{a}_{p-3,q-3} + \binom{p-2}{4} \frac{(q-2)!}{(q-6)!} a^4 \bar{a}_{p-2,q-2} \\
&\quad + \binom{p-1}{5} \frac{(q-1)!}{(q-6)!} a^5 \bar{a}_{p-1,q-1} + \binom{p}{6} \frac{q!}{(q-6)!} a^6 \bar{a}_{p,q},
\end{aligned}$$

then,

$$\begin{aligned}
2\text{Ima}_{p-6,q-6} &= \binom{p-5}{1} \frac{(q-5)!}{(q-6)!} \theta \text{Rea}_{p-5,q-5} + \binom{p-4}{2} \frac{(q-4)!}{(q-6)!} \theta^2 \text{Ima}_{p-4,q-4} \\
&\quad - \binom{p-3}{3} \frac{(q-3)!}{(q-6)!} \theta^3 \text{Rea}_{p-3,q-3} - \binom{p-2}{4} \frac{(q-2)!}{(q-6)!} \theta^4 \text{Ima}_{p-2,q-2} \\
&\quad + \binom{p-1}{5} \frac{(q-1)!}{(q-6)!} \theta^5 \text{Rea}_{p-1,q-1} + \binom{p}{6} \frac{q!}{(q-6)!} \theta^6 \text{Ima}_{p,q} \\
&= \binom{p-5}{1} \frac{(q-5)!}{(q-6)!} \theta \text{Rea}_{p-5,q-5} + \binom{p-4}{2} \frac{(q-4)!}{(q-6)!} \theta^2 \left(\frac{1}{2} \binom{p-3}{1} \frac{(q-3)!}{(q-4)!} \theta \right. \\
&\quad \times \text{Rea}_{p-3,q-3} + \frac{1}{4} \binom{p-1}{3} \frac{(q-1)!}{(q-4)!} \theta^3 \text{Rea}_{p-1,q-1} \Big) - \binom{p-3}{3} \frac{(q-3)!}{(q-6)!} \theta^3 \text{Rea}_{p-3,q-3} \\
&\quad - \binom{p-2}{4} \frac{(q-2)!}{(q-6)!} \theta^4 \times \frac{1}{2} (p-1)(q-1) \theta \text{Rea}_{p-1,q-1} + \binom{p-1}{5} \frac{(q-1)!}{(q-6)!} \theta^5 \text{Rea}_{p-1,q-1} \\
&= \binom{p-5}{1} \frac{(q-5)!}{(q-6)!} \theta \text{Rea}_{p-5,q-5} + \frac{1}{2} \binom{p-3}{3} \frac{(q-3)!}{(q-6)!} \theta^3 \text{Rea}_{p-3,q-3} \\
&\quad + \binom{p-1}{5} \frac{(q-1)!}{(q-6)!} \theta^5 \text{Rea}_{p-1,q-1},
\end{aligned}$$

so

$$\begin{aligned}
\text{Ima}_{p-6,q-6} &= \frac{1}{2} \binom{p-5}{1} \frac{(q-5)!}{(q-6)!} \theta \text{Rea}_{p-5,q-5} + \frac{1}{4} \binom{p-3}{3} \frac{(q-3)!}{(q-6)!} \theta^3 \text{Rea}_{p-3,q-3} \\
&\quad + \frac{1}{2} \binom{p-1}{5} \frac{(q-1)!}{(q-6)!} \theta^5 \text{Rea}_{p-1,q-1}.
\end{aligned}$$

For $a_{p-7,q-7}$,

$$a_{p-7,q-7} = \bar{a}_{p-7,q-7} + \binom{p-6}{1} \frac{(q-6)!}{(q-7)!} a \bar{a}_{p-6,q-6} + \binom{p-5}{2} \frac{(q-5)!}{(q-7)!} a^2 \bar{a}_{p-5,q-5}$$

$$\begin{aligned}
 & + \binom{p-4}{3} \frac{(q-4)!}{(q-7)!} a^3 \bar{a}_{p-4,q-4} + \binom{p-3}{4} \frac{(q-3)!}{(q-7)!} a^4 \bar{a}_{p-3,q-3} \\
 & + \binom{p-2}{5} \frac{(q-2)!}{(q-7)!} a^5 \bar{a}_{p-2,q-2} + \binom{p-1}{6} \frac{(q-1)!}{(q-7)!} a^6 \bar{a}_{p-1,q-1} + \binom{p}{7} \frac{q!}{(q-7)!} a^7 \bar{a}_{p,q},
 \end{aligned}$$

then

$$\begin{aligned}
 2\text{Im}a_{p-7,q-7} &= \binom{p-6}{1} \frac{(q-6)!}{(q-7)!} \theta \text{Re}a_{p-6,q-6} + \binom{p-5}{2} \frac{(q-5)!}{(q-7)!} \theta^2 \text{Im}a_{p-5,q-5} \\
 & - \binom{p-4}{3} \frac{(q-4)!}{(q-7)!} \theta^3 \text{Re}a_{p-4,q-4} - \binom{p-3}{4} \frac{(q-3)!}{(q-7)!} \theta^4 \text{Im}a_{p-3,q-3} \\
 & + \binom{p-2}{5} \frac{(q-2)!}{(q-7)!} \theta^5 \text{Re}a_{p-2,q-2} + \binom{p-1}{6} \frac{(q-1)!}{(q-7)!} \theta^6 \text{Im}a_{p-1,q-1} \\
 & - \binom{p}{7} \frac{q!}{(q-7)!} \theta^7 a_{p,q} \\
 &= \binom{p-6}{1} \frac{(q-6)!}{(q-7)!} \theta \text{Re}a_{p-6,q-6} + \binom{p-5}{2} \frac{(q-5)!}{(q-7)!} \theta^2 \left(\frac{1}{2} \binom{p-4}{1} \frac{(q-4)!}{(q-5)!} \right. \\
 & \quad \times \theta \text{Re}a_{p-4,q-4} + \frac{1}{4} \binom{p-2}{3} \frac{(q-2)!}{(q-5)!} \theta^3 \text{Re}a_{p-2,q-2} + \frac{1}{2} \binom{p}{5} \frac{q!}{(q-5)!} \theta^5 a_{p,q} \Big) \\
 & - \binom{p-4}{3} \frac{(q-4)!}{(q-7)!} \theta^3 \text{Re}a_{p-4,q-4} - \binom{p-3}{4} \frac{(q-3)!}{(q-7)!} \theta^4 \left(\frac{1}{2} \binom{p-2}{1} \frac{(q-2)!}{(q-3)!} \right. \\
 & \quad \times \theta \text{Re}a_{p-2,q-2} + \frac{1}{4} \binom{p}{3} \frac{q!}{(q-3)!} \theta^3 a_{p,q} \Big) + \binom{p-2}{5} \frac{(q-2)!}{(q-7)!} \theta^5 \text{Re}a_{p-2,q-2} \\
 & + \binom{p-1}{6} \frac{(q-1)!}{(q-7)!} \theta^6 \times \frac{1}{2} p q \theta a_{p,q} - \binom{p}{7} \frac{q!}{(q-7)!} \theta^7 a_{p,q} \\
 &= \binom{p-6}{1} \frac{(q-6)!}{(q-7)!} \theta \text{Re}a_{p-6,q-6} + \frac{1}{2} \binom{p-4}{3} \frac{(q-4)!}{(q-7)!} \theta^3 \text{Re}a_{p-4,q-4} \\
 & + \binom{p-2}{5} \frac{(q-2)!}{(q-7)!} \theta^5 \text{Re}a_{p-2,q-2} + \frac{17}{4} \binom{p}{7} \frac{q!}{(q-7)!} \theta^7 a_{p,q},
 \end{aligned}$$

so

$$\begin{aligned}
 \text{Im}a_{p-7,q-7} &= \frac{1}{2} \binom{p-6}{1} \frac{(q-6)!}{(q-7)!} \theta \text{Re}a_{p-6,q-6} + \frac{1}{4} \binom{p-4}{3} \frac{(q-4)!}{(q-7)!} \theta^3 \text{Re}a_{p-4,q-4} \\
 & + \frac{1}{2} \binom{p-2}{5} \frac{(q-2)!}{(q-7)!} \theta^5 \text{Re}a_{p-2,q-2} + \frac{17}{8} \binom{p}{7} \frac{q!}{(q-7)!} \theta^7 a_{p,q}.
 \end{aligned}$$

...

From the above calculation we conjecturally set

$$(4.13) \quad \text{Im}a_{p-2l-1,q-2l-1} = \sum_{k=0}^l a_k \binom{p-2l+2k}{2k+1} \frac{(q-2l+2k)!}{(q-2l-1)!} \theta^{2k+1} \text{Re}a_{p-2l+2k,q-2l+2k},$$

for a sequence $\{a_n\}_{n \geq 0}$, and we can see that $a_0 = \frac{1}{2}$, $a_1 = \frac{1}{4}$, $a_2 = \frac{1}{2}$, $a_3 = \frac{17}{8}$. Note that

$$\begin{aligned}
 2\text{Im}a_{p-2l-1,q-2l-1} &= \sum_{k=1}^l (-1)^{k-1} \binom{p-2l+2k-1}{2k} \frac{(q-2l+2k-1)!}{(q-2l-1)!} \theta^{2k} \text{Im}a_{p-2l-1+2k,q-2l-1+2k} \\
 & + \sum_{k=0}^l (-1)^k \binom{p-2l+2k}{2k+1} \frac{(q-2l+2k)!}{(q-2l-1)!} \theta^{2k+1} \text{Re}a_{p-2l+2k,q-2l+2k},
 \end{aligned}$$

apply (4.13) to the above equality and compare the coefficients of $\text{Re}a_{p-2l+2k, q-2l+2k}$ where $0 \leq k \leq l$, we have for $m \leq l$,

$$(4.14) \quad 2a_m \binom{p-2l+2m}{2m+1} = \sum_{k=1}^m (-1)^{k-1} \binom{p-2l+2k-1}{2k} \binom{p-2l+2m}{2m-2k+1} a_{m-k} + (-1)^m \binom{p-2l+2m}{2m+1},$$

after simplification, we have

$$(4.15) \quad a_m = \sum_{k=0}^m (-1)^{k+1} \binom{2m+1}{2k} a_{m-k} + (-1)^m.$$

Similarly, we have

$$(4.16) \quad \text{Im}a_{p-2l, q-2l} = \sum_{k=0}^{l-1} a_k \binom{p-2l+1+2k}{2k+1} \frac{(q-2l+1+2k)!}{(q-2l)!} \theta^{2k+1} \text{Re}a_{p-2l+1+2k, q-2l+1+2k},$$

and in summary, for $l \in \mathbb{N}$, conjecturally we have

$$(4.17) \quad \text{Im}a_{p-l, q-l} = \sum_{k=0}^{\lfloor \frac{l-1}{2} \rfloor} a_k \binom{p-l+1+2k}{2k+1} \frac{(q-l+1+2k)!}{(q-l)!} \theta^{2k+1} \text{Re}a_{p-l+1+2k, q-l+1+2k}.$$

The first few elements of $\{a_n\}_{n \geq 0}$ satisfying (4.15) is

$$\frac{1}{2}, \frac{1}{4}, \frac{1}{2}, \frac{17}{8}, \frac{31}{2}, \frac{691}{4}, \frac{5461}{2}, \frac{929569}{16}, \frac{3202291}{2}, \frac{221930581}{4}, \frac{4722116521}{2}, \dots$$

and with the help of the On-Line Encyclopedia of Integer Sequences (OEIS), I find that the general term formula of $\{a_n\}_{n \geq 0}$ is

$$(4.18) \quad a_n = \frac{(-1)^n (2^{2n+2} - 1) B_{2n+2}}{n+1}$$

where $\{B_n\}_{n \geq 0}$ is the Bernoulli number series. We prove that this is indeed the case, even for the smooth case, i.e., (4.3) and (4.17) are both hold.

Proof of Theorem 4.1. For

$$(4.19) \quad T = \sum_{p, q=0}^{\infty} a_{p, q} x^p y^q \in A^{\infty}(\mathbb{R}_{\theta}^2),$$

if $T = T^*$, then

$$(4.20) \quad \sum_{p, q=0}^{\infty} a_{p, q} x^p y^q = \sum_{p, q=0}^{\infty} \bar{a}_{p, q} y^q x^p = \sum_{p, q=0}^{\infty} \bar{a}_{p, q} \sum_{h=0}^p \binom{p}{h} \frac{q!}{(q-p+h)!} a^{p-h} x^h y^{q-p+h},$$

compare the coefficients, we have

$$(4.21) \quad a_{p, q} = \sum_{h=0}^{\infty} \binom{p+h}{h} \frac{(q+h)!}{q!} \bar{a}_{p+h, q+h} a^h.$$

Take real and imaginary part of the above equality and note that $a = i\theta$, we have

$$\begin{aligned} \text{Re}a_{p, q} &= \sum_{h=0}^{\infty} \binom{p+2h}{2h} \frac{(q+2h)!}{q!} a^{2h} \text{Re}a_{p+2h, q+2h} - \sum_{h=0}^{\infty} \binom{p+2h+1}{2h+1} \frac{(q+2h+1)!}{q!} \\ &\quad \times a^{2h+1} i \text{Im}a_{p+2h+1, q+2h+1} \\ &= \sum_{h=0}^{\infty} (-1)^h \binom{p+2h}{2h} \frac{(q+2h)!}{q!} \theta^{2h} \text{Re}a_{p+2h, q+2h} + \sum_{h=0}^{\infty} (-1)^h \binom{p+2h+1}{2h+1} \frac{(q+2h+1)!}{q!} \\ &\quad \times \theta^{2h+1} \text{Im}a_{p+2h+1, q+2h+1}, \end{aligned}$$

$$\begin{aligned}
 \text{Ima}_{p,q} &= - \sum_{h=0}^{\infty} \binom{p+2h}{2h} \frac{(q+2h)!}{q!} a^{2h} \text{Ima}_{p+2h,q+2h} + \frac{1}{i} \sum_{h=0}^{\infty} \binom{p+2h+1}{2h+1} \frac{(q+2h+1)!}{q!} \\
 &\quad \times a^{2h+1} \text{Rea}_{p+2h+1,q+2h+1} \\
 &= - \sum_{h=0}^{\infty} (-1)^h \binom{p+2h}{2h} \frac{(q+2h)!}{q!} \theta^{2h} \text{Ima}_{p+2h,q+2h} + \sum_{h=0}^{\infty} (-1)^h \binom{p+2h+1}{2h+1} \frac{(q+2h+1)!}{q!} \\
 &\quad \times \theta^{2h+1} \text{Rea}_{p+2h+1,q+2h+1},
 \end{aligned}$$

so for $\forall k \in \mathbb{N}$,

$$\begin{aligned}
 &\sum_{h=0}^{\infty} (-1)^h \binom{p+2k+2h+1}{2h+1} \frac{(q+2k+2h+1)!}{(q+2k)!} \theta^{2h+1} \text{Ima}_{p+2k+2h+1,q+2k+2h+1} \\
 &= \sum_{h=0}^{\infty} (-1)^h \binom{p+2k+2h+2}{2h+2} \frac{(q+2k+2h+2)!}{(q+2k)!} \theta^{2h+2} \text{Rea}_{p+2k+2h+2,q+2k+2h+2}.
 \end{aligned}$$

Let $\{b_n\}_{n \geq 1}$ be a number series satisfying

$$(4.22) \quad \sum_{k=1}^n (-1)^{k-1} b_k \binom{p+2n+1}{2n+1-2k} = \binom{p+2n+1}{2n+1}.$$

Such a series exists. In fact, $b_1 = \frac{1}{6}(p+2)(p+1)$ and for $n \geq 2$,

$$(4.23) \quad b_n = \frac{(-1)^n}{p+2n+1} \left(\binom{p+2n+1}{2n+1} - \sum_{k=1}^{n-1} (-1)^{k-1} b_k \binom{p+2n+1}{2n+1-2k} \right).$$

Then

$$\begin{aligned}
 &(p+1)(q+1)\theta \text{Ima}_{p+1,q+1} \\
 &= \sum_{h=0}^{\infty} (-1)^h \binom{p+2h+1}{2h+1} \frac{(q+2h+1)!}{q!} \theta^{2h+1} \text{Ima}_{p+2h+1,q+2h+1} + \sum_{k=1}^{\infty} b_k \frac{(q+2k)!}{q!} \theta^{2k} \sum_{h=0}^{\infty} (-1)^h \\
 &\quad \times \binom{p+2k+2h+1}{2h+1} \frac{(q+2k+2h+1)!}{(q+2k)!} \theta^{2h+1} \text{Ima}_{p+2k+2h+1,q+2k+2h+1} \\
 &= \sum_{h=0}^{\infty} (-1)^h \binom{p+2h+2}{2h+2} \frac{(q+2h+2)!}{q!} \theta^{2h+2} \text{Rea}_{p+2h+2,q+2h+2} + \sum_{k=1}^{\infty} b_k \sum_{h=0}^{\infty} (-1)^h \binom{p+2k+2h+2}{2h+2} \\
 &\quad \times \frac{(q+2k+2h+2)!}{q!} \theta^{2k+2h+2} \text{Rea}_{p+2k+2h+2,q+2k+2h+2} \\
 &= \sum_{h=0}^{\infty} (-1)^h \binom{p+2h+2}{2h+2} \frac{(q+2h+2)!}{q!} \theta^{2h+2} \text{Rea}_{p+2h+2,q+2h+2} + \sum_{k=1}^{\infty} b_k \sum_{h=0}^{\infty} (-1)^h \binom{p+2k+2h+2}{2k+2h+2} \\
 &\quad \times \frac{(2k+2h+2)!p!}{(2h+2)!(p+2k)!} \frac{(q+2k+2h+2)!}{q!} \theta^{2k+2h+2} \text{Rea}_{p+2k+2h+2,q+2k+2h+2} \\
 &= \binom{p+2}{2} \frac{(q+2)!}{q!} \theta^2 \text{Rea}_{p+2,q+2} + \sum_{h=1}^{\infty} (-1)^h \left(1 + \sum_{k=1}^h (-1)^k b_k \frac{(2h+2)!p!}{(2h-2k+2)!(p+2k)!} \right) \binom{p+2h+2}{2h+2} \\
 &\quad \times \frac{(q+2h+2)!}{q!} \theta^{2h+2} \text{Rea}_{p+2h+2,q+2h+2} \\
 &= \sum_{h=0}^{\infty} (-1)^h \left(1 + \sum_{k=1}^h (-1)^k b_k \frac{(2h+2)!p!}{(2h-2k+2)!(p+2k)!} \right) \binom{p+2h+2}{2h+2} \frac{(q+2h+2)!}{q!} \theta^{2h+2} \text{Rea}_{p+2h+2,q+2h+2},
 \end{aligned}$$

hence for $p, q \in \mathbb{N}$,

$$\begin{aligned}
 & \text{Im} a_{p+1, q+1} \\
 (4.24) \quad &= \sum_{h=0}^{\infty} (-1)^h \left(\frac{1}{2h+2} + \sum_{k=1}^h (-1)^k b_k \frac{(2h+1)!p!}{(2h-2k+2)!(p+2k)!} \right) \binom{p+2h+2}{2h+1} \frac{(q+2h+2)!}{(q+1)!} \\
 & \quad \times \theta^{2h+1} \text{Re} a_{p+2h+2, q+2h+2}.
 \end{aligned}$$

Let

$$(4.25) \quad a_h := (-1)^h \left(\frac{1}{2h+2} + \sum_{j=1}^h (-1)^j b_j \frac{(2h+1)!p!}{(2h-2j+2)!(p+2j)!} \right),$$

then for $m \in \mathbb{N}$,

$$\begin{aligned}
 & a_m - \sum_{k=0}^m (-1)^{k+1} \binom{2m+1}{2k} a_{m-k} \\
 &= (-1)^m \left(\frac{1}{2m+2} + \sum_{k=1}^m (-1)^k b_k \frac{(2m+1)!p!}{(2m-2k+2)!(p+2k)!} \right) - \sum_{k=0}^m (-1)^{m+1} \binom{2m+1}{2k} \left(\frac{1}{2m-2k+2} \right. \\
 & \quad \left. + \sum_{j=1}^{m-k} (-1)^j b_j \frac{(2m-2k+1)!p!}{(2m-2k-2j+2)!(p+2j)!} \right) \\
 &= (-1)^m \left(\frac{1}{2m+2} + \sum_{k=0}^m \binom{2m+1}{2k} \frac{1}{2m-2k+2} \right) + (-1)^m \left(\sum_{k=1}^m (-1)^k b_k \frac{(2m+1)!p!}{(2m-2k+2)!(p+2k)!} \right. \\
 & \quad \left. + \sum_{k=0}^m \sum_{j=1}^{m-k} (-1)^j b_j \binom{2m+1}{2k} \frac{(2m-2k+1)!p!}{(2m-2k-2j+2)!(p+2j)!} \right) \\
 &= \frac{(-1)^m}{2m+2} \left(1 + \sum_{k=0}^m \binom{2m+2}{2k} \right) + (-1)^m (2m+1)!p! \left(\sum_{k=1}^m \frac{(-1)^k b_k}{(2m-2k+2)!(p+2k)!} \right. \\
 & \quad \left. + \sum_{k=0}^m \frac{1}{(2k)!} \sum_{j=1}^{m-k} \frac{(-1)^j b_j}{(2m-2k-2j+2)!(p+2j)!} \right) \\
 &= \frac{(-1)^m 2^{2m+1}}{2m+2} + (-1)^m (2m+1)!p! \sum_{k=1}^m \frac{(-1)^k b_k}{(p+2k)!} \left(\frac{1}{(2m-2k+2)!} + \sum_{j=0}^{m-k} \frac{1}{(2j)!(2m-2k-2j+2)!} \right) \\
 &= \frac{(-1)^m 2^{2m+1}}{2m+2} + (-1)^m (2m+1)!p! \sum_{k=1}^m \frac{(-1)^k 2^{2m-2k+1} b_k}{(2m-2k+2)!(p+2k)!}.
 \end{aligned}$$

The definition (4.22) of $\{b_n\}$ implies

$$\begin{aligned}
 2(4^n - n - 1) \binom{p+2n+2}{p} &= \sum_{m=1}^n \binom{p+2n+2}{2n+1-2m} \binom{p+2m+1}{2m+1} \\
 &= \sum_{m=1}^n \binom{p+2n+2}{2n+1-2m} \sum_{k=1}^m (-1)^{k-1} b_k \binom{p+2m+1}{2m+1-2k} \\
 &= \sum_{k=1}^n (-1)^{k-1} b_k \sum_{j=1}^{n+1-k} \binom{p+2n+2}{2n-2k-2j+3} \binom{p+2k+2j-1}{2j-1} \\
 &= \sum_{k=1}^n \frac{(-1)^{k-1} 2^{2n-2k+1} (p+2n+2)! b_k}{(2n-2k+2)!(p+2k)!},
 \end{aligned}$$

hence

$$\begin{aligned} a_m - \sum_{k=0}^m (-1)^{k+1} \binom{2m+1}{2k} a_{m-k} \\ = \frac{(-1)^m 2^{2m+1}}{2m+2} + (-1)^m (2m+1)! p! \sum_{k=1}^m \frac{(-1)^k 2^{2m-2k+1} b_k}{(2m-2k+2)!(p+2k)!} \\ = \frac{(-1)^m 2^{2m+1}}{2m+2} - (-1)^m (2m+1)! p! \frac{2(4^m - m - 1)}{(p+2m+2)!} \binom{p+2m+2}{p} = (-1)^m. \end{aligned}$$

Finally, from (4.24) we can see that for $h \geq 1$,

$$\text{Im}a_{2h,2h} = \sum_{k=0}^{\infty} a_k \binom{2h+2k+1}{2k+1} \frac{(2h+2k+1)!}{(2h)!} \theta^{2k+1} \text{Re}a_{2h+2k+1,2h+2k+1},$$

so

$$\begin{aligned} \text{Im}a_{0,0} &= - \sum_{h=0}^{\infty} (-1)^h \binom{2h}{2h} \frac{(2h)!}{0!} \theta^{2h} \text{Im}a_{2h,2h} + \sum_{h=0}^{\infty} (-1)^h \binom{2h+1}{2h+1} \frac{(2h+1)!}{0!} \theta^{2h+1} \text{Re}a_{2h+1,2h+1} \\ &= - \text{Im}a_{0,0} - \sum_{h=1}^{\infty} (-1)^h (2h)! \theta^{2h} \sum_{k=0}^{\infty} a_k \binom{2h+2k+1}{2k+1} \frac{(2h+2k+1)!}{(2h)!} \theta^{2k+1} \text{Re}a_{2h+2k+1,2h+2k+1} \\ &\quad + \sum_{h=0}^{\infty} (-1)^h (2h+1)! \theta^{2h+1} \text{Re}a_{2h+1,2h+1}, \\ &= - \text{Im}a_{0,0} + \sum_{h=0}^{\infty} \left(a_h + \sum_{k=0}^h (-1)^{k+1} \binom{2h+1}{2k} a_{h-k} + (-1)^h \right) (2h+1)! \theta^{2h+1} \text{Re}a_{2h+1,2h+1} \\ &= - \text{Im}a_{0,0} + 2 \sum_{h=0}^{\infty} a_h (2h+1)! \theta^{2h+1} \text{Re}a_{2h+1,2h+1}, \end{aligned}$$

hence

$$(4.26) \quad \text{Im}a_{0,0} = \sum_{k=0}^{\infty} a_k (2k+1)! \theta^{2k+1} \text{Re}a_{2k+1,2k+1}.$$

Similarly we can prove that

$$(4.27) \quad \text{Im}a_{0,1} = \sum_{k=0}^{\infty} a_k (2k+2)! \theta^{2k+1} \text{Re}a_{2k+1,2k+2}$$

and

$$(4.28) \quad \text{Im}a_{1,0} = \sum_{k=0}^{\infty} a_k (2k+2)! \theta^{2k+1} \text{Re}a_{2k+2,2k+1},$$

combine with (4.24), we can see that for $\forall p, q \in \mathbb{N}$, we have

$$(4.29) \quad \text{Im}a_{p,q} = \sum_{k=0}^{\infty} a_k \binom{p+2k+1}{2k+1} \frac{(q+2k+1)!}{q!} \theta^{2k+1} \text{Re}a_{p+2k+1,q+2k+1}.$$

Finally we need to determine the values of $\{a_n\}_{n \geq 0}$. Recall that the Bernoulli numbers $\{B_n\}_{n \in \mathbb{N}}$ are defined via

$$(4.30) \quad \frac{z}{e^z - 1} + \frac{z}{2} - 1 = \sum_{k=1}^{\infty} \frac{B_{2k}}{(2k)!} z^{2k},$$

then

$$(4.31) \quad \frac{2z}{e^{2z} - 1} + z - 1 = \sum_{k=1}^{\infty} \frac{2^{2k} B_{2k}}{(2k)!} z^{2k},$$

so

$$(4.32) \quad \sum_{k=1}^{\infty} \frac{(2^{2k}-1)B_{2k}}{(2k)!} z^{2k} = \frac{2z}{e^{2z}-1} + z - 1 - \left(\frac{z}{e^z-1} + \frac{z}{2} - 1 \right) = \frac{z(e^z-1)}{2(e^z+1)},$$

hence

$$(4.33) \quad \left(2 + \sum_{k=1}^{\infty} \frac{z^k}{k!} \right) \sum_{k=1}^{\infty} \frac{(2^{2k}-1)B_{2k}}{(2k)!} z^{2k} = \frac{1}{2} \sum_{k=1}^{\infty} \frac{z^{k+1}}{k!}.$$

For $n \in \mathbb{N}$, compare the coefficients of z^{2n+2} on both sides, we have

$$(4.34) \quad 2 \frac{(2^{2n+2}-1)B_{2n+2}}{(2n+2)!} + \sum_{k=1}^n \frac{(2^{2k}-1)B_{2k}}{(2n+2-2k)!(2k)!} = \frac{1}{2(2n+1)!},$$

which means

$$(4.35) \quad (2^{2n+2}-1)B_{2n+2} + \sum_{k=1}^{n+1} \binom{2n+2}{2k} (2^{2k}-1)B_{2k} = n+1.$$

For $n \in \mathbb{N}$ let

$$(4.36) \quad c_n := \frac{(-1)^n (2^{2n+2}-1)B_{2n+2}}{n+1},$$

then (4.35) implies

$$(4.37) \quad (-1)^n (n+1)c_n + \sum_{k=0}^n \binom{2n+2}{2k} (-1)^k (k+1)c_k = n+1,$$

so

$$(4.38) \quad c_n - \sum_{k=0}^n (-1)^{k+1} \binom{2n+1}{2k} c_{n-k} = (-1)^n.$$

Note that the series $\{a_n\}_{n \geq 0}$ also satisfies (4.38), so we must have

$$(4.39) \quad a_n = c_n = \frac{(-1)^n (2^{2n+2}-1)B_{2n+2}}{n+1}, \quad \forall n \in \mathbb{N}$$

and we finish the proof. $\square$

REFERENCES

- [1] Bieberbach. L. Analytische Fortsetzung. Springer-Verlag (1955).
- [2] Baum, P., Connes, A., Higson, N. Classifying Space for Proper Actions and K-Theory of Group C^* -algebras. 1994.
- [3] Bernstein, D. S. Scalar, Vector, and Matrix Mathematics: Theory, Facts, and Formulas - Revised and Expanded Edition. (2018).
- [4] Douglas, M. R., Nekrasov, N. A. Noncommutative Field Theory. International Journal of Theoretical Physics, 42(11) (2001), 2609-2704.
- [5] Elliott, G. A. On the K-theory of the C^* -algebra generated by a projective representation of a torsion-free discrete abelian group. in Operator Algebras and Group Representations 1, G. Arsene, ed., Pitman, London, (1984), 157-184.
- [6] Elliott, G. A., Li, H. F. Morita equivalence of smooth noncommutative tori. Acta Math 199 (2007), 1-27.
- [7] Exel, R. A Fredholm operator approach to Morita equivalence, K-Theory 7 (1993), 285-308.
- [8] Gracia-Bondía, J. M., Várilly, J. C. Algebras Of Distributions Suitable For Phase Space Quantum Mechanics. 1, J. Math. Phys. 29 (1988) 869.
- [9] Gracia-Bondía, J. M. and Várilly, J. C., Figueroa, H. Elements of noncommutative geometry. Birkhaeuser, Boston, USA. 2001.

- [10] Grosse, H., Wulkenhaar, R. Renormalisation of ϕ^4 -theory on noncommutative $\mathbb{R}^4$ in the matrix base. *Letters in Mathematical Physics* 71.1(2003), 13-26.
- [11] Hesselholt, L., Madsen, I. Cyclic polytopes and the K-theory of truncated polynomial algebras. *Inventiones mathematicae*. 130 (1997), 73–97.
- [12] Hesselholt, L., Larsen, M., Lindenstrauss, A.E. On the K-theory of division algebras over local fields. *Inventiones mathematicae*, 219 (2019), 281-329.
- [13] Higson, N., Lafforgue, V., Skandalis, G. Counterexamples to the Baum—Connes conjecture. *Geometric and Functional Analysis GAFA* 12.2(2002), 330-354.
- [14] Mineyev, I., Yu, G. The Baum-Connes conjecture for hyperbolic groups. *Inventiones mathematicae*. 149 (2002), 97–122.
- [15] Munshi, S., Yang, R. Self-adjoint elements in the pseudo-unitary group $\mathbf{U}(p, p)$. *Linear Algebra and its Applications*. 560, (2019), 100-113.
- [16] Pimsner, M. V., Voiculescu, D. Exact sequences for K-groups and Extgroups of certain cross-product C^* -algebras, *J Oper. Theory*. 4 (1980), 93-118.
- [17] Rosenberg, J. Algebraic K-theory and its applications. *Graduate Texts in Mathematics*. 1994.
- [18] Rieffel, M. A. Non-commutative tori — a case study of noncommutative differentiable manifolds, *Contemporary Math*. 105 (1990), 191–211.
- [19] Rieffel, M. A., Schwarz, A. Morita equivalence of multidimensional noncommutative tori. *International Journal of Mathematics* 10.02(1999), 289-299.
- [20] Voigt, C. The Baum-Connes conjecture for free orthogonal quantum groups. *Advances in Mathematics* 227. 5(2011), 1873-1913.

SCHOOL OF MATHEMATICS AND STATISTICS, JIANGSU NORMAL UNIVERSITY, XUZHOU 221100, CHINA
Email address: guanren@jsnu.edu.cn