

Boundedness of Calderón–Zygmund operators on ball Campanato-type function spaces

Yiqun Chen, Hongchao Jia and Dachun Yang*

Abstract Let X be a ball quasi-Banach function space on \mathbb{R}^n satisfying some mild assumptions. In this article, the authors first find a reasonable version \tilde{T} of the Calderón–Zygmund operator T on the ball Campanato-type function space $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$ with $q \in [1, \infty)$, $s \in \mathbb{Z}_+^n$, and $d \in (0, \infty)$. Then the authors prove that \tilde{T} is bounded on $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$ if and only if, for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \leq s$, $T^*(x^\gamma) = 0$, which is hence sharp. Moreover, \tilde{T} is proved to be the adjoint operator of T , which further strengthens the rationality of the definition of \tilde{T} . All these results have a wide range of applications. In particular, even when they are applied, respectively, to weighted Lebesgue spaces, variable Lebesgue spaces, Orlicz spaces, Orlicz-slice spaces, Morrey spaces, mixed-norm Lebesgue spaces, local generalized Herz spaces, and mixed-norm Herz spaces, all the obtained results are new. The proofs of these results strongly depend on the properties of the kernel of T under consideration and also on the dual theorem on $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$.

1 Introduction

Recall that, for any $q \in (0, \infty]$, the *Lebesgue space* $L^q(\mathbb{R}^n)$ is defined to be the set of all the measurable functions f on \mathbb{R}^n such that

$$\|f\|_{L^q(\mathbb{R}^n)} := \begin{cases} \left[\int_{\mathbb{R}^n} |f(x)|^q dx \right]^{\frac{1}{q}} & \text{if } q \in (0, \infty), \\ \operatorname{ess\,sup}_{x \in \mathbb{R}^n} |f(x)| & \text{if } q = \infty \end{cases}$$

is finite. It is well known that Lebesgue spaces play a fundamental role in the modern analysis of mathematics. Recall that the Calderón–Zygmund operator T is bounded on $L^p(\mathbb{R}^n)$ with $p \in (1, \infty)$, which no longer holds true for $L^\infty(\mathbb{R}^n)$. Indeed, T maps $L^\infty(\mathbb{R}^n)$ to the space $\operatorname{BMO}(\mathbb{R}^n)$ of functions with bounded mean oscillation, which was introduced by John and Nirenberg [38]. This implies that the space $\operatorname{BMO}(\mathbb{R}^n)$ is a reasonable replacement of $L^\infty(\mathbb{R}^n)$ when studying the boundedness of Calderón–Zygmund operators. Moreover, in recent years, the boundedness of Calderón–Zygmund operators on BMO -type spaces has attracted more and more attention. For instance,

2020 *Mathematics Subject Classification*. Primary 42B20; Secondary 42B25, 42B30, 42B35, 46E35, 47A30.

Key words and phrases. Calderón–Zygmund operator, Campanato-type space, ball quasi-Banach function space, Hardy-type space.

This project is partially supported by the National Natural Science Foundation of China (Grant Nos. 11971058 and 12071197) and the National Key Research and Development Program of China (Grant No. 2020YFA0712900).

*Corresponding author, E-mail: dcyang@bnu.edu.cn/August 12, 2022/Final version.

Nakai [45, 46] obtained the boundedness of Calderón–Zygmund operators on both Campanato-type spaces with variable growth conditions over spaces of homogeneous type, in the sense of Coifman and Weiss, and their predual spaces. We refer the reader also to [1, 24, 26, 47, 63, 64] for more studies on this.

Recall that Sawano et al. [50] introduced the ball quasi-Banach function space X and the related Hardy space $H_X(\mathbb{R}^n)$ which is a natural generalization of the classical Hardy space $H^p(\mathbb{R}^n)$. Ball quasi-Banach function spaces contain many well-known function spaces, for instance, the weighted Lebesgue space, the Morrey space, the mixed-norm Lebesgue space, and the Orlicz-slice space, but they may not be quasi-Banach function spaces (see [50, 72, 69] for the details). This implies that the studies on X (see, for instance, [8, 16, 27, 35, 50, 60, 61, 62, 72]) have more wide applications. We also refer the reader to [55, 56, 66, 65] for studies on (weak) Hardy spaces associated with X on spaces of homogeneous type in the sense of Coifman and Weiss.

On the other hand, Fefferman and Stein [21] obtained the well-known dual theorem that $(H^1(\mathbb{R}^n))^* = \text{BMO}(\mathbb{R}^n)$. Later, Taibleson and Weiss [57] generalized the above dual theorem to the classical Hardy space $H^p(\mathbb{R}^n)$ for any given $p \in (0, 1]$, namely, for any $q \in [1, \infty]$ and any integer $s \in [0, \infty) \cap [n(\frac{1}{p} - 1), \infty)$, the dual space of $H^p(\mathbb{R}^n)$ is $C_{\frac{1}{p}-1, q, s}(\mathbb{R}^n)$, the Campanato space introduced in [7], which coincides with $\text{BMO}(\mathbb{R}^n)$ when $p = 1$. Very recently, motivated by $(H^p(\mathbb{R}^n))^* = C_{\frac{1}{p}-1, q, s}(\mathbb{R}^n)$, Zhang et al. [69] studied the dual theorem of $H_X(\mathbb{R}^n)$. To be precise, Zhang et al. [69] proved that the dual space of $H_X(\mathbb{R}^n)$ is $\mathcal{L}_{X, q, s, d}(\mathbb{R}^n)$, the ball Campanato-type function space associated with X and, moreover, established the Carleson measure characterization of $\mathcal{L}_{X, q, s, d}(\mathbb{R}^n)$.

In this article, we study the boundedness of Calderón–Zygmund operators on $\mathcal{L}_{X, q, s, d}(\mathbb{R}^n)$. To be precise, let X be a ball quasi-Banach function space on \mathbb{R}^n satisfying some mild assumptions. We first find a reasonable version \widetilde{T} of the Calderón–Zygmund operator T on the ball Campanato-type function space $\mathcal{L}_{X, q, s, d}(\mathbb{R}^n)$ (see Definition 3.5 below), which was originally introduced in [42] in the case $s := 0$ and further studied by Nakai [45, 46] on spaces of homogeneous type. Then we prove that \widetilde{T} is bounded on $\mathcal{L}_{X, q, s, d}(\mathbb{R}^n)$ if and only if, for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \leq s$, $T^*(x^\gamma) = 0$, which is hence sharp. Moreover, \widetilde{T} is proved to be the adjoint operator of T , which further strengthens the rationality of the definition of \widetilde{T} . All these results have a wide range of applications. In particular, even when they are applied, respectively, to weighted Lebesgue spaces, variable Lebesgue spaces, Orlicz spaces, Orlicz-slice spaces, Morrey spaces, mixed-norm Lebesgue spaces, local generalized Herz spaces, and mixed-norm Herz spaces, all the obtained results are new. Due to the generality and the practicability, more applications of these main results of this article are predictable. The proofs of these results strongly depend on the properties of the kernel of T under consideration and also on the dual theorem that $(H_X(\mathbb{R}^n))^* = \mathcal{L}_{X, q, s, d}(\mathbb{R}^n)$.

To be precise, the remainder of this article is organized as follows.

In Section 2, we recall some basic concepts on the quasi-Banach function space X (see Definition 2.1 below) and its related ball Campanato-type function space (see Definition 2.6 below).

Section 3 is divided into two parts. In Subsection 3.1, we find a reasonable version \widetilde{T} of the Calderón–Zygmund operator T on $\mathcal{L}_{X, q, s, d}(\mathbb{R}^n)$. Then we establish the boundedness of \widetilde{T} on $\mathcal{L}_{X, q, s, d}(\mathbb{R}^n)$ (see Theorem 3.13 below). In Subsection 3.2, we prove that \widetilde{T} is the adjoint operator of T (see Theorem 3.24 below).

In Section 4, we apply all the above main results to eight concrete examples of ball quasi-

Banach function spaces, namely, the weighted Lebesgue space $L_w^p(\mathbb{R}^n)$, the variable Lebesgue space $L^{p(\cdot)}(\mathbb{R}^n)$, the Orlicz space $L^\Phi(\mathbb{R}^n)$, the Orlicz-slice space $(E_\Phi^r)_t(\mathbb{R}^n)$, the Morrey space $M_r^p(\mathbb{R}^n)$, the mixed-norm Lebesgue space $L^{\vec{p}}(\mathbb{R}^n)$, the local generalized Herz space $\mathcal{K}_{\omega,0}^{p,r}(\mathbb{R}^n)$, and the mixed-norm Herz space $\dot{E}_{\vec{q}}^{\vec{\alpha},\vec{\beta}}(\mathbb{R}^n)$, respectively. Therefore, all the boundedness of \tilde{T} on $\mathcal{L}_{L_w^p(\mathbb{R}^n),q,s,d}(\mathbb{R}^n)$, $\mathcal{L}_{L^{p(\cdot)}(\mathbb{R}^n),q,s,d}(\mathbb{R}^n)$, $\mathcal{L}_{L^\Phi(\mathbb{R}^n),q,s,d}(\mathbb{R}^n)$, and $\mathcal{L}_{(E_\Phi^r)_t(\mathbb{R}^n),q,s,d}(\mathbb{R}^n)$, $\mathcal{L}_{M_r^p(\mathbb{R}^n),q,s,d}(\mathbb{R}^n)$, $\mathcal{L}_{L^{\vec{p}}(\mathbb{R}^n),q,s,d}(\mathbb{R}^n)$, $\mathcal{L}_{\mathcal{K}_{\omega,0}^{p,r}(\mathbb{R}^n),q,s,d}(\mathbb{R}^n)$, and $\mathcal{L}_{\dot{E}_{\vec{q}}^{\vec{\alpha},\vec{\beta}}(\mathbb{R}^n),q,s,d}(\mathbb{R}^n)$ are obtained (see, respectively, Theorems 4.4, 4.8, 4.12, 4.16, 4.20, 4.24, 4.30, and 4.34 below).

Finally, we make some conventions on notation. Let $\mathbb{N} := \{1, 2, \dots\}$, $\mathbb{Z}_+ := \mathbb{N} \cup \{0\}$, and $\mathbb{Z}_+^n := (\mathbb{Z}_+)^n$. We always denote by C a *positive constant* which is independent of the main parameters, but it may vary from line to line. The symbol $f \lesssim g$ means that $f \leq Cg$. If $f \lesssim g$ and $g \lesssim f$, we then write $f \sim g$. If $f \leq Cg$ and $g = h$ or $g \leq h$, we then write $f \lesssim g \sim h$ or $f \lesssim g \lesssim h$, rather than $f \lesssim g = h$ or $f \lesssim g \leq h$. For any $s \in \mathbb{Z}_+$, we use $\mathcal{P}_s(\mathbb{R}^n)$ to denote the set of all the polynomials on \mathbb{R}^n with total degree not greater than s . We use $\mathbf{0}$ to denote the *origin* of \mathbb{R}^n . For any measurable subset E of \mathbb{R}^n , we denote by $\mathbf{1}_E$ its characteristic function. Moreover, for any $x \in \mathbb{R}^n$ and $r \in (0, \infty)$, let $B(x, r) := \{y \in \mathbb{R}^n : |y - x| < r\}$. Furthermore, for any $\lambda \in (0, \infty)$ and any ball $B(x, r) \subset \mathbb{R}^n$ with $x \in \mathbb{R}^n$ and $r \in (0, \infty)$, let $\lambda B(x, r) := B(x, \lambda r)$. Finally, for any $q \in [1, \infty]$, we denote by q' its *conjugate exponent*, namely, $\frac{1}{q} + \frac{1}{q'} = 1$. Also, when we prove a lemma, proposition, theorem, or corollary, we always use the same symbols in the wanted proved lemma, proposition, theorem, or corollary.

2 Ball quasi-Banach function spaces

In this section, we recall the definitions of ball quasi-Banach function spaces and their related ball Campanato-type function spaces. In what follows, let $\mathcal{M}(\mathbb{R}^n)$ denote the set of all measurable functions on \mathbb{R}^n and

$$\mathbb{B}(\mathbb{R}^n) := \{B(x, r) : x \in \mathbb{R}^n \text{ and } r \in (0, \infty)\}. \quad (2.1)$$

The following definition of ball quasi-Banach function spaces on \mathbb{R}^n is just [50, Definition 2.2].

Definition 2.1. Let $X \subset \mathcal{M}(\mathbb{R}^n)$ be a quasi-normed linear space equipped with a quasi-norm $\|\cdot\|_X$ which makes sense for all measurable functions on \mathbb{R}^n . Then X is called a *ball quasi-Banach function space* on \mathbb{R}^n if it satisfies:

- (i) if $f \in \mathcal{M}(\mathbb{R}^n)$, then $\|f\|_X = 0$ implies that $f = 0$ almost everywhere;
- (ii) if $f, g \in \mathcal{M}(\mathbb{R}^n)$, then $|g| \leq |f|$ almost everywhere implies that $\|g\|_X \leq \|f\|_X$;
- (iii) if $\{f_m\}_{m \in \mathbb{N}} \subset \mathcal{M}(\mathbb{R}^n)$ and $f \in \mathcal{M}(\mathbb{R}^n)$, then $0 \leq f_m \uparrow f$ almost everywhere as $m \rightarrow \infty$ implies that $\|f_m\|_X \uparrow \|f\|_X$ as $m \rightarrow \infty$;
- (iv) $B \in \mathbb{B}(\mathbb{R}^n)$ implies that $\mathbf{1}_B \in X$, where $\mathbb{B}(\mathbb{R}^n)$ is the same as in (2.1).

Moreover, a ball quasi-Banach function space X is called a *ball Banach function space* if it satisfies:

(v) for any $f, g \in X$,

$$\|f + g\|_X \leq \|f\|_X + \|g\|_X;$$

(vi) for any ball $B \in \mathbb{B}(\mathbb{R}^n)$, there exists a positive constant $C_{(B)}$, depending on B , such that, for any $f \in X$,

$$\int_B |f(x)| dx \leq C_{(B)} \|f\|_X.$$

Remark 2.2. (i) Let X be a ball quasi-Banach function space on \mathbb{R}^n . By [65, Remark 2.6(i)], we conclude that, for any $f \in \mathcal{M}(\mathbb{R}^n)$, $\|f\|_X = 0$ if and only if $f = 0$ almost everywhere.

(ii) As was mentioned in [65, Remark 2.6(ii)], we obtain an equivalent formulation of Definition 2.1 via replacing any ball B by any bounded measurable set E therein.

(iii) We should point out that, in Definition 2.1, if we replace any ball B by any measurable set E with finite measure, we obtain the definition of (quasi-)Banach function spaces which were originally introduced in [6, Definitions 1.1 and 1.3]. Thus, a (quasi-)Banach function space is also a ball (quasi-)Banach function space and the converse is not necessary to be true.

(iv) By [17, Theorem 2], we conclude that both (ii) and (iii) of Definition 2.1 imply that any ball quasi-Banach function space is complete.

We also recall the concepts of both the convexity and the concavity of ball quasi-Banach function spaces, which are a part of [50, Definition 2.6].

Definition 2.3. Let X be a ball quasi-Banach function space and $p \in (0, \infty)$.

(i) The p -convexification X^p of X is defined by setting

$$X^p := \{f \in \mathcal{M}(\mathbb{R}^n) : |f|^p \in X\}$$

equipped with the *quasi-norm* $\|f\|_{X^p} := \| |f|^p \|_X^{1/p}$ for any $f \in X^p$.

(ii) The space X is said to be *concave* if there exists a positive constant C such that, for any $\{f_k\}_{k \in \mathbb{N}} \subset \mathcal{M}(\mathbb{R}^n)$,

$$\sum_{k=1}^{\infty} \|f_k\|_X \leq C \left\| \sum_{k=1}^{\infty} |f_k| \right\|_X.$$

In particular, when $C = 1$, X is said to be *strictly concave*.

Next, we present the definition of ball Campanato-type function spaces associated with X introduced in [69, Definition 3.2]. In what follows, for any given $q \in (0, \infty]$, we use $L_{\text{loc}}^q(\mathbb{R}^n)$ to denote the set of all the measurable functions f on \mathbb{R}^n such that $f \mathbf{1}_E \in L^q(\mathbb{R}^n)$ for any bounded measurable set $E \subset \mathbb{R}^n$. For any $f \in L_{\text{loc}}^1(\mathbb{R}^n)$ and any finite measurable subset $E \subset \mathbb{R}^n$, let

$$f_E := \int_E f(x) dx := \frac{1}{|E|} \int_E f(x) dx.$$

Moreover, for any $s \in \mathbb{Z}_+$, $\mathcal{P}_s(\mathbb{R}^n)$ denotes the set of all the polynomials on \mathbb{R}^n with total degree not greater than s ; for any ball $B \in \mathbb{B}(\mathbb{R}^n)$ and any locally integrable function g on \mathbb{R}^n , $P_B^{(s)}(g)$

denotes the *minimizing polynomial* of g with total degree not greater than s , namely, $P_B^{(s)}(g)$ is the unique polynomial $f \in \mathcal{P}_s(\mathbb{R}^n)$ such that, for any $P \in \mathcal{P}_s(\mathbb{R}^n)$,

$$\int_B [g(x) - f(x)]P(x) dx = 0.$$

Recall that Yan et al. [67, Definition 1.11] introduced the following Campanato space $\mathcal{L}_{X,q,s}(\mathbb{R}^n)$ associated with the ball quasi-Banach function space X .

Definition 2.4. Let X be a ball quasi-Banach function space, $q \in [1, \infty)$, and $s \in \mathbb{Z}_+$. The *Campanato space* $\mathcal{L}_{X,q,s}(\mathbb{R}^n)$, associated with X , is defined to be the set of all the $f \in L_{\text{loc}}^q(\mathbb{R}^n)$ such that

$$\|f\|_{\mathcal{L}_{X,q,s}(\mathbb{R}^n)} := \sup_{B \subset \mathbb{R}^n} \frac{|B|}{\|\mathbf{1}_B\|_X} \left\{ \int_B |f(x) - P_B^{(s)}(f)(x)|^q dx \right\}^{1/q} < \infty,$$

where the supremum is taken over all balls $B \in \mathbb{B}(\mathbb{R}^n)$.

In what follows, by abuse of notation, we identify $f \in \mathcal{L}_{X,q,s}(\mathbb{R}^n)$ with $f + \mathcal{P}_s(\mathbb{R}^n)$.

Remark 2.5. Let $q \in [1, \infty)$, $s \in \mathbb{Z}_+$, and $\alpha \in [0, \infty)$. Recall that Campanato [7] introduced the following well-known *Campanato space* $\mathcal{C}_{\alpha,q,s}(\mathbb{R}^n)$ which is defined to be the set of all the $f \in L_{\text{loc}}^q(\mathbb{R}^n)$ such that

$$\|f\|_{\mathcal{C}_{\alpha,q,s}(\mathbb{R}^n)} := \sup |B|^{-\alpha} \left[\int_B |f(x) - P_B^{(s)}(f)(x)|^q dx \right]^{\frac{1}{q}} < \infty,$$

where the supremum is taken over all balls $B \subset \mathbb{R}^n$. It is well known that $\mathcal{C}_{\alpha,q,s}(\mathbb{R}^n)$ when $\alpha = 0$ coincides with the space $\text{BMO}(\mathbb{R}^n)$. Moreover, when $X := L^{\frac{1}{\alpha+1}}(\mathbb{R}^n)$, we have $\mathcal{L}_{X,q,s}(\mathbb{R}^n) = \mathcal{C}_{\alpha,q,s}(\mathbb{R}^n)$.

Very recently, Zhang et al. [69, Definition 3.2] introduced the following ball Campanato-type function space $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$ associated with the ball quasi-Banach function space X .

Definition 2.6. Let X be a ball quasi-Banach function space, $q \in [1, \infty)$, $d \in (0, \infty)$, and $s \in \mathbb{Z}_+$. Then the *ball Campanato-type function space* $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$, associated with X , is defined to be the set of all the $f \in L_{\text{loc}}^q(\mathbb{R}^n)$ such that

$$\begin{aligned} \|f\|_{\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)} := & \sup \left\| \left\{ \sum_{i=1}^m \left(\frac{\lambda_i}{\|\mathbf{1}_{B_i}\|_X} \right)^d \mathbf{1}_{B_i} \right\}^{\frac{1}{d}} \right\|_X^{-1} \\ & \times \sum_{j=1}^m \frac{\lambda_j |B_j|}{\|\mathbf{1}_{B_j}\|_X} \left[\int_{B_j} |f(x) - P_{B_j}^{(s)}(f)(x)|^q dx \right]^{\frac{1}{q}} \end{aligned}$$

is finite, where the supremum is taken over all $m \in \mathbb{N}$, $\{B_j\}_{j=1}^m \subset \mathbb{B}(\mathbb{R}^n)$, and $\{\lambda_j\}_{j=1}^m \subset [0, \infty)$ with $\sum_{j=1}^m \lambda_j \neq 0$.

In what follows, by abuse of notation, we identify $f \in \mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$ with $f + \mathcal{P}_s(\mathbb{R}^n)$.

Remark 2.7. Let $q \in [1, \infty)$, $s \in \mathbb{Z}_+$, $d \in (0, 1)$, and X be a concave ball quasi-Banach function space. In this case, it was proved in [69, Proposition 3.7] that $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n) = \mathcal{L}_{X,q,s}(\mathbb{R}^n)$ with equivalent quasi-norms.

3 Calderón–Zygmund operators on $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$

This section is devoted to studying the boundedness of Calderón–Zygmund operators on the space $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$. For this purpose, we divide this section into two subsections. In Subsection 3.1, we present a reasonable version \widetilde{T} of the Calderón–Zygmund operator T on $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$ and then obtain the boundedness of \widetilde{T} on $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$. In Subsection 3.2, we prove that \widetilde{T} is the adjoint operator of T , which further strengthens the rationality of the definition of \widetilde{T} .

3.1 Boundedness of Calderón–Zygmund operators on $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$

In this subsection, we establish the boundedness of Calderón–Zygmund operators on the space $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$. We begin with the concept of the s -order standard kernel (see, for instance, [54, Chapter III]). In what follows, for any $\gamma = (\gamma_1, \dots, \gamma_n) \in \mathbb{Z}_+^n$, any γ -order differentiable function $F(\cdot, \cdot)$ on $\mathbb{R}^n \times \mathbb{R}^n \setminus \{(w, w) : w \in \mathbb{R}^n\}$, and any $(x, y) \in \mathbb{R}^n \times \mathbb{R}^n \setminus \{(w, w) : w \in \mathbb{R}^n\}$, let

$$\partial_{(1)}^\gamma F(x, y) := \frac{\partial^{|\gamma|}}{\partial x_1^{\gamma_1} \dots \partial x_n^{\gamma_n}} F(x, y)$$

and

$$\partial_{(2)}^\gamma F(x, y) := \frac{\partial^{|\gamma|}}{\partial y_1^{\gamma_1} \dots \partial y_n^{\gamma_n}} F(x, y).$$

Definition 3.1. Let $s \in \mathbb{Z}_+$ and $\delta \in (0, 1]$. A measurable function K on $\mathbb{R}^n \times \mathbb{R}^n \setminus \{(w, w) : w \in \mathbb{R}^n\}$ is called an s -order standard kernel with regularity δ if there exists a positive constant C such that, for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \leq s$, the following hold true:

(i) for any $x, y \in \mathbb{R}^n$ with $x \neq y$,

$$\left| \partial_{(2)}^\gamma K(x, y) \right| \leq \frac{C}{|x - y|^{n+|\gamma|}}; \quad (3.1)$$

(ii) (3.1) still holds true for the first variable of K ;

(iii) for any $x, y, z \in \mathbb{R}^n$ with $x \neq y$ and $|x - y| \geq 2|y - z|$,

$$\left| \partial_{(2)}^\gamma K(x, y) - \partial_{(2)}^\gamma K(x, z) \right| \leq C \frac{|y - z|^\delta}{|x - y|^{n+|\gamma|+\delta}}; \quad (3.2)$$

(iv) (3.2) still holds true for the first variable of K .

In what follows, we use $\eta \rightarrow 0^+$ to denote $\eta \in (0, \infty)$ and $\eta \rightarrow 0$.

Definition 3.2. Let $s \in \mathbb{Z}_+$ and K be the same as in Definition 3.1. A linear operator T is called an s -order Calderón–Zygmund singular integral operator with kernel K if T is bounded on $L^2(\mathbb{R}^n)$ and, for any given $f \in L^2(\mathbb{R}^n)$ and for almost every $x \in \mathbb{R}^n$,

$$T(f)(x) = \lim_{\eta \rightarrow 0^+} T_\eta(f)(x),$$

where, for any $\eta \in (0, \infty)$,

$$T_\eta(f)(x) := \int_{\mathbb{R}^n \setminus B(x, \eta)} K(x, y) f(y) dy.$$

Remark 3.3. Let T be the same as in Definition 3.2. From a claim in [20, p. 102], we deduce that, for any $q \in (1, \infty)$, T is bounded on $L^q(\mathbb{R}^n)$ and, moreover, for any $f \in L^q(\mathbb{R}^n)$,

$$T(f) = \lim_{\eta \rightarrow 0^+} T_\eta(f)$$

both almost everywhere on \mathbb{R}^n and in $L^q(\mathbb{R}^n)$.

Now, we recall the following well-known concept of vanishing moments on T (see, for instance, [43, p. 23]).

Definition 3.4. Let $s \in \mathbb{Z}_+$. An s -order Calderón–Zygmund singular integral operator T is said to have the *vanishing moments up to order s* if, for any function $a \in L^2(\mathbb{R}^n)$ having compact support and satisfying that, for any $\gamma := (\gamma_1, \dots, \gamma_n) \in \mathbb{Z}_+^n$ with $|\gamma| \leq s$, $\int_{\mathbb{R}^n} a(x)x^\gamma dx = 0$, it holds true that

$$T^*(x^\gamma) := \int_{\mathbb{R}^n} T(a)(x)x^\gamma dx = 0,$$

here and thereafter, $|\gamma| := \gamma_1 + \dots + \gamma_n$ and, for any $x := (x_1, \dots, x_n) \in \mathbb{R}^n$, $x^\gamma := x_1^{\gamma_1} \dots x_n^{\gamma_n}$.

To show the boundedness of Calderón–Zygmund operators on both $\mathcal{L}_{X,q,s}(\mathbb{R}^n)$ and $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$, we now recall the following definition of s -order modified Calderón–Zygmund operators (see, for instance, [37, Definition 2.11]). In what follows, for any given $s \in \mathbb{Z}_+$ and $\delta \in (0, 1]$, we always need to consider a function $f \in L^1_{\text{loc}}(\mathbb{R}^n)$ satisfying that, for any ball $B(x, r) \in \mathbb{B}(\mathbb{R}^n)$ with $x \in \mathbb{R}^n$ and $r \in (0, \infty)$,

$$\int_{\mathbb{R}^n \setminus B(x,r)} \frac{|f(y) - P_{B(x,r)}^{(s)}(f)(y)|}{|x-y|^{n+s+\delta}} dy < \infty. \quad (3.3)$$

Definition 3.5. Let $s \in \mathbb{Z}_+$, $\delta \in (0, 1]$, and K be an s -order standard kernel. For any $x, y \in \mathbb{R}^n$ with $x \neq y$, let

$$\widetilde{K}(x, y) := K(y, x), \quad (3.4)$$

which is the formal adjoint kernel of K , and $B_0 := B(x_0, r_0) \in \mathbb{B}(\mathbb{R}^n)$ with center $x_0 \in \mathbb{R}^n$ and radius $r_0 \in (0, \infty)$. The s -order modified Calderón–Zygmund operator \widetilde{T}_{B_0} with kernel \widetilde{K} is defined by setting, for any $f \in L^1_{\text{loc}}(\mathbb{R}^n)$ satisfying (3.3) and for almost every $x \in \mathbb{R}^n$,

$$\widetilde{T}_{B_0}(f)(x) := \lim_{\eta \rightarrow 0^+} \widetilde{T}_{B_0, \eta}(f)(x), \quad (3.5)$$

where, for any $\eta \in (0, \infty)$,

$$\begin{aligned} \widetilde{T}_{B_0, \eta}(f)(x) := & \int_{\mathbb{R}^n \setminus B(x, \eta)} \left[\widetilde{K}(x, y) \right. \\ & \left. - \sum_{\{\gamma \in \mathbb{Z}_+^n : |\gamma| \leq s\}} \frac{\partial_{(1)}^\gamma \widetilde{K}(x_0, y)}{\gamma!} (x - x_0)^\gamma \mathbf{1}_{\mathbb{R}^n \setminus B_0}(y) \right] f(y) dy. \end{aligned}$$

Next, we show that $\widetilde{T}_{B_0}(f)$ in (3.5) is well defined; since its proof is quite similar to that of [37, Proposition 2.22], we only sketch some important steps.

Proposition 3.6. *Let all the symbols be the same as in Definition 3.5. Then, for any $f \in L^1_{\text{loc}}(\mathbb{R}^n)$ satisfying (3.3), $\widetilde{T}_{B_0}(f)$ in (3.5) is well defined almost everywhere on \mathbb{R}^n .*

Proof. Let $f \in L^1_{\text{loc}}(\mathbb{R}^n)$ satisfy (3.3), and $\widetilde{B} := B(z, r) \in \mathbb{B}(\mathbb{R}^n)$ with $z \in \mathbb{R}^n$ and $r \in (0, \infty)$. To show the present proposition, we write, for any given $\eta \in (0, r)$ [and hence $B(x, \eta) \subset 2\widetilde{B}$] and for any $x \in \widetilde{B}$,

$$\widetilde{T}_{B_0, \eta}(f)(x) = E_{\widetilde{B}, \eta}(x) + E_{\widetilde{B}}(x) + E_{\widetilde{B}, B_0, \eta}^{(1)}(x) + E_{\widetilde{B}, B_0, \eta}^{(2)}(x),$$

where

$$\begin{aligned} E_{\widetilde{B}, \eta}(x) &:= \int_{2\widetilde{B} \setminus B(x, \eta)} \widetilde{K}(x, y) \left[f(y) - P_{2\widetilde{B}}^{(s)}(f)(y) \right] dy, \\ E_{\widetilde{B}}(x) &:= \int_{\mathbb{R}^n \setminus 2\widetilde{B}} \left[\widetilde{K}(x, y) - \sum_{\{\gamma \in \mathbb{Z}_+^n: |\gamma| \leq s\}} \frac{\partial_{(1)}^\gamma \widetilde{K}(z, y)}{\gamma!} (x - z)^\gamma \right] \\ &\quad \times \left[f(y) - P_{2\widetilde{B}}^{(s)}(f)(y) \right] dy, \end{aligned} \tag{3.6}$$

$$E_{\widetilde{B}, B_0, \eta}^{(1)}(x) := \int_{\mathbb{R}^n \setminus B(x, \eta)} L_{\widetilde{B}, B_0}(x, y) \left[f(y) - P_{2\widetilde{B}}^{(s)}(f)(y) \right] dy$$

with

$$\begin{aligned} L_{\widetilde{B}, B_0}(x, y) &:= \sum_{\{\gamma \in \mathbb{Z}_+^n: |\gamma| \leq s\}} \frac{\partial_{(1)}^\gamma \widetilde{K}(z, y)}{\gamma!} (x - z)^\gamma \mathbf{1}_{\mathbb{R}^n \setminus 2\widetilde{B}}(y) \\ &\quad - \sum_{\{\gamma \in \mathbb{Z}_+^n: |\gamma| \leq s\}} \frac{\partial_{(1)}^\gamma \widetilde{K}(x_0, y)}{\gamma!} (x - x_0)^\gamma \mathbf{1}_{\mathbb{R}^n \setminus B_0}(y), \end{aligned}$$

and

$$\begin{aligned} E_{\widetilde{B}, B_0, \eta}^{(2)}(x) &:= \int_{\mathbb{R}^n \setminus B(x, \eta)} \left[\widetilde{K}(x, y) \right. \\ &\quad \left. - \sum_{\{\gamma \in \mathbb{Z}_+^n: |\gamma| \leq s\}} \frac{\partial_{(1)}^\gamma \widetilde{K}(x_0, y)}{\gamma!} (x - x_0)^\gamma \mathbf{1}_{\mathbb{R}^n \setminus B_0}(y) \right] P_{2\widetilde{B}}^{(s)}(f)(y) dy. \end{aligned}$$

Thus, to prove that $\widetilde{T}_{B_0}(f)$ is well defined almost everywhere on \widetilde{B} , it suffices to show that $E_{\widetilde{B}, \eta}$, $E_{\widetilde{B}}$, $E_{\widetilde{B}, B_0, \eta}^{(1)}$, and $E_{\widetilde{B}, B_0, \eta}^{(2)}$ are convergent almost everywhere on B as $\eta \rightarrow 0^+$, respectively.

We first consider $E_{\widetilde{B},\eta}^-$. Indeed, it is easy to show that $[f - P_{2\widetilde{B}}^{(s)}(f)]\mathbf{1}_{2\widetilde{B}} \in L^1(\mathbb{R}^n)$ and hence, for any $\eta \in (0, \infty)$, $E_{\widetilde{B},\eta}^-$ is well defined almost everywhere on \widetilde{B} . Moreover, let $T_{(\widetilde{K})}$ be the s -order Calderón–Zygmund singular integral operator with kernel \widetilde{K} . Then, by Remark 3.3, we know that

$$\lim_{\eta \rightarrow 0^+} E_{\widetilde{B},\eta}^- = T_{(\widetilde{K})} \left(\left[f - P_{2\widetilde{B}}^{(s)}(f) \right] \mathbf{1}_{2\widetilde{B}} \right) \quad (3.7)$$

almost everywhere on \widetilde{B} .

Now, we estimate $E_{\widetilde{B}}^-$. Observe that, since $E_{\widetilde{B}}$ is independent of η , we only need to prove that $E_{\widetilde{B}}^-$ is bounded on \widetilde{B} . Indeed, applying both an argument similar to that used in the estimation of [37, (2.33)] and (3.3), we conclude that, for any $x \in \widetilde{B}$,

$$|E_{\widetilde{B}}^-(x)| \lesssim r^{s+\delta} \int_{\mathbb{R}^n \setminus 2\widetilde{B}} \frac{|f(y) - P_{2\widetilde{B}}^{(s)}(f)(y)|}{|y - z|^{n+s+\delta}} dy < \infty. \quad (3.8)$$

Next, we consider both $E_{\widetilde{B},B_0,\eta}^{(1)}$ and $E_{\widetilde{B},B_0,\eta}^{(2)}$. Indeed, from [37, (2.39)], we deduce that

$$P_1(\widetilde{B}, B_0; f)(\cdot) := \lim_{\eta \rightarrow 0^+} E_{\widetilde{B},B_0,\eta}^{(1)}(\cdot) \in \mathcal{P}_s(B). \quad (3.9)$$

Moreover, similarly to the proof of [37, (2.40)], we conclude that there exists a polynomial $P_2(\widetilde{B}, B_0; f) \in \mathcal{P}_s(\mathbb{R}^n)$ such that, for almost every $x \in \widetilde{B}$,

$$\lim_{\eta \rightarrow 0^+} E_{\widetilde{B},B_0,\eta}^{(2)}(x) = P_2(\widetilde{B}, B_0; f)(x). \quad (3.10)$$

From (3.7), (3.8), (3.9), and (3.10), we deduce that, for any given ball $\widetilde{B} \in \mathbb{B}(\mathbb{R}^n)$ and for almost every $x \in \widetilde{B}$,

$$\begin{aligned} \widetilde{T}_{B_0}(f)(x) &= \lim_{\eta \rightarrow 0^+} \widetilde{T}_{B_0,\eta}(f)(x) \\ &= T_{(\widetilde{K})} \left(\left[f - P_{2\widetilde{B}}^{(d)}(f) \right] \mathbf{1}_{2\widetilde{B}} \right)(x) + E_{\widetilde{B}}^-(x) \\ &\quad + P_1(\widetilde{B}, B_0; f)(x) + P_2(\widetilde{B}, B_0; f)(x). \end{aligned} \quad (3.11)$$

Therefore, $\widetilde{T}_{B_0}(f)$ is well defined almost everywhere on \mathbb{R}^n , which then completes the proof of Proposition 3.6. \square

Remark 3.7. Let all the symbols be the same as in Proposition 3.6. By [37, Remark 2.23], we conclude that, for any f satisfying (3.3) and for any ball $B_1 \in \mathbb{B}(\mathbb{R}^n)$,

$$\widetilde{T}_{B_0}(f) - \widetilde{T}_{B_1}(f) \in \mathcal{P}_s(\mathbb{R}^n)$$

after changing values on a set of measure zero. Based on this, in what follows, we write \widetilde{T} instead of \widetilde{T}_{B_0} if there exists no confusion.

Now, we show that \widetilde{T} is well defined on both $\mathcal{L}_{X,q,s}(\mathbb{R}^n)$ and $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$. To this end, we need the following mild assumption about the boundedness of the Hardy–Littlewood maximal operator on ball quasi-Banach function spaces (see, for instance, [69, Assumption 2.6]). Recall that the *Hardy–Littlewood maximal operator* \mathcal{M} is defined by setting, for any $f \in L^1_{\text{loc}}(\mathbb{R}^n)$ and $x \in \mathbb{R}^n$,

$$\mathcal{M}(f)(x) := \sup_{B \ni x} \frac{1}{|B|} \int_B |f(y)| dy,$$

where the supremum is taken over all the balls $B \in \mathbb{B}(\mathbb{R}^n)$ containing x .

Assumption 3.8. *Let X be a ball quasi-Banach function space. Assume that there exists a $p_- \in (0, \infty)$ such that, for any given $p \in (0, p_-)$ and $u \in (1, \infty)$, there exists a positive constant C such that, for any $\{f_j\}_{j=1}^\infty \subset \mathcal{M}(\mathbb{R}^n)$,*

$$\left\| \left\{ \sum_{j \in \mathbb{N}} [\mathcal{M}(f_j)]^u \right\}^{\frac{1}{u}} \right\|_{X^{1/p}} \leq C \left\| \left(\sum_{j \in \mathbb{N}} |f_j|^u \right)^{\frac{1}{u}} \right\|_{X^{1/p}}.$$

Remark 3.9. Let both X and p_- satisfy Assumption 3.8, and let $d \in (0, \infty)$. From this and the fact that, for any $r \in (0, \min\{d, p_-\})$, any ball $B \in \mathbb{B}(\mathbb{R}^n)$, and any $\beta \in [1, \infty)$, $\mathbf{1}_{\beta B} \leq (\beta + 1)^{\frac{dn}{r}} [\mathcal{M}(\mathbf{1}_B)]^{\frac{d}{r}}$, we easily deduce that, for any $r \in (0, \min\{d, p_-\})$, any $\beta \in [1, \infty)$, any sequence $\{B_j\}_{j \in \mathbb{N}} \subset \mathbb{B}(\mathbb{R}^n)$, and any $\{\lambda_j\}_{j \in \mathbb{N}} \subset [0, \infty)$,

$$\left\| \left(\sum_{j \in \mathbb{N}} \lambda_j^d \mathbf{1}_{\beta B_j} \right)^{\frac{1}{d}} \right\|_X \leq (2\beta)^{\frac{n}{r}} \left\| \left(\sum_{j \in \mathbb{N}} \lambda_j^d \mathbf{1}_{B_j} \right)^{\frac{1}{d}} \right\|_X.$$

Particularly, for any $r \in (0, p_-)$ and any ball $B \in \mathbb{B}(\mathbb{R}^n)$, we have

$$\|\mathbf{1}_{\beta B}\|_X \leq (2\beta)^{\frac{n}{r}} \|\mathbf{1}_B\|_X. \quad (3.12)$$

We also need the following lemma which is a direct corollary of [10, Lemma 3.12]; we omit the details here.

Lemma 3.10. *Let both X and p_- satisfy Assumption 3.8. Let $\delta \in (0, 1]$, $s \in (\frac{n}{p_-} - n - \delta, \infty) \cap \mathbb{Z}_+$, $q \in [1, \infty)$, and $d \in (0, \infty)$. Then, for any $f \in \mathcal{L}_{X,q,s}(\mathbb{R}^n)$ [resp., $f \in \mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$], f satisfies (3.3).*

As a direct application of both Lemma 3.10 and Proposition 3.6, we immediately obtain the following conclusion; we omit the details here.

Theorem 3.11. *Let both X and p_- satisfy Assumption 3.8. Let $\delta \in (0, 1]$, $q \in (1, \infty)$, $s \in (\frac{n}{p_-} - n - \delta, \infty) \cap \mathbb{Z}_+$, and $d \in (0, \infty)$. Let \widetilde{T} be the same as in Remark 3.7. Then \widetilde{T} is well defined on both $\mathcal{L}_{X,q,s}(\mathbb{R}^n)$ and $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$.*

Next, we establish the boundedness of \widetilde{T} , respectively, on $\mathcal{L}_{X,q,s}(\mathbb{R}^n)$ and $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$. To be precise, we show the following two theorems.

Theorem 3.12. *Let both X and p_- satisfy Assumption 3.8. Let $\delta \in (0, 1]$, $q \in (1, \infty)$, $s \in (\frac{n}{p_-} - n - \delta, \infty) \cap \mathbb{Z}_+$, K be an s -order standard kernel with regularity δ , T the same as in Definition 3.1 with kernel K , \widetilde{K} the same as in (3.4), and \widetilde{T} the same as in Remark 3.7 with kernel \widetilde{K} . Then \widetilde{T} is bounded on $\mathcal{L}_{X,q,s}(\mathbb{R}^n)$, namely, there exists a positive constant C such that, for any $f \in \mathcal{L}_{X,q,s}(\mathbb{R}^n)$,*

$$\|\widetilde{T}(f)\|_{\mathcal{L}_{X,q,s}(\mathbb{R}^n)} \leq C\|f\|_{\mathcal{L}_{X,q,d}(\mathbb{R}^n)}$$

if and only if T has the vanishing moments up to order s .

Theorem 3.13. *Let both X and p_- satisfy Assumption 3.8. Let $\delta \in (0, 1]$, $q \in (1, \infty)$, $d \in (0, \infty)$, $s \in (\frac{n}{\min\{p_-, d\}} - n - \delta, \infty) \cap \mathbb{Z}_+$, K be an s -order standard kernel with regularity δ , T the same as in Definition 3.1 with kernel K , \widetilde{K} the same as in (3.4), and \widetilde{T} the same as in Remark 3.7 with kernel \widetilde{K} . Then \widetilde{T} is bounded on $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$, namely, there exists a positive constant C such that, for any $f \in \mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$,*

$$\|\widetilde{T}(f)\|_{\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)} \leq C\|f\|_{\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)}$$

if and only if T has the vanishing moments up to order s .

We only prove Theorem 3.13; since the proof of Theorem 3.12 is similar, we omit the details here. To this end, we first state two technique lemmas. The following conclusion is just [37, Proposition 2.16].

Lemma 3.14. *Let $s \in \mathbb{Z}_+$, K be an s -order standard kernel, and T an s -order Calderón–Zygmund singular integral operator with kernel K . Let \widetilde{K} be the same as in (3.4), $B_0 := B(x_0, r_0) \subset \mathbb{R}^n$ a given ball with $x_0 \in \mathbb{R}^n$ and $r_0 \in (0, \infty)$, and, for any $v \in \mathbb{Z}_+^n$ with $|v| \leq s$, $\widetilde{T}_{B_0}(y^v)$ the same as in (3.5) with f replaced by y^v . Then T has the vanishing moments up to order s if and only if, for any $v \in \mathbb{Z}_+^n$ with $|v| \leq s$, $\widetilde{T}_{B_0}(y^v)(\cdot) \in \mathcal{P}_s(\mathbb{R}^n)$ after changing values on a set of measure zero.*

Similarly to the proof of [37, Lemma 2.21], we have the following lemma; we omit the details here.

Lemma 3.15. *Let $q \in [1, \infty)$, $s \in \mathbb{Z}_+$, and $\lambda \in (s, \infty)$. Then there exists a positive constant C such that, for any $f \in L_{\text{loc}}^1(\mathbb{R}^n)$ and any ball $B(x, r)$ with $x \in \mathbb{R}^n$ and $r \in (0, \infty)$,*

$$\begin{aligned} & \int_{\mathbb{R}^n \setminus B(x,r)} \frac{|f(y) - P_{B(x,r)}^{(s)}(f)(y)|}{|x-y|^{n+\lambda}} dy \\ & \leq C \sum_{k \in \mathbb{N}} (2^k r)^{-\lambda} \left[\int_{2^k B(x,r)} |f(y) - P_{2^k B(x,r)}^{(s)}(f)(y)|^q dy \right]^{\frac{1}{q}}. \end{aligned}$$

Now, we show Theorem 3.13.

Proof of Theorem 3.13. From Theorem 3.11, we deduce that \widetilde{T} is well defined on $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$. We now show the necessity. Indeed, if \widetilde{T} is bounded on $\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$, then, for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \leq s$,

$$\|\widetilde{T}(x^\gamma)\|_{\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)} \lesssim \|x^\gamma\|_{\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)} = 0$$

and hence $\widetilde{T}(x^\gamma) \in \mathcal{P}_s(\mathbb{R}^n)$. Using this and Lemma 3.14, we find that, for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \leq s$, $T^*(x^\gamma) = 0$, which completes the proof of the necessity.

Next, we show the sufficiency. To this end, let $T_{(\widetilde{K})}$ be the s -order Calderón–Zygmund singular integral operator with kernel \widetilde{K} , $m \in \mathbb{N}$, $\{B_j\}_{j=1}^m := \{B(z_j, r_j)\}_{j=1}^m \subset \mathbb{B}(\mathbb{R}^n)$ with both $\{z_j\}_{j=1}^m \subset \mathbb{R}^n$ and $\{r_j\}_{j=1}^m \subset (0, \infty)$, and $\{\lambda_j\}_{j=1}^m \subset [0, \infty)$ with $\sum_{j=1}^m \lambda_j \neq 0$. It is easy to prove that, for any $g \in L_{\text{loc}}^1(\mathbb{R}^n)$ and any ball $B \in \mathbb{B}(\mathbb{R}^n)$,

$$\left[\int_B |g(x) - P_B^{(s)}(g)(x)|^q dx \right]^{\frac{1}{q}} \sim \inf_{P \in \mathcal{P}_s(B)} \left[\int_B |g(x) - P(x)|^q dx \right]^{\frac{1}{q}};$$

see, for instance, [36, (2.12)]. From this, (3.11), and $P_B^{(s)}(P) = P$ for any $P \in \mathcal{P}_s(\mathbb{R}^n)$ and any ball $B \in \mathbb{B}(\mathbb{R}^n)$, we infer that, for any given $f \in \mathcal{L}_{X,q,s,d}(\mathbb{R}^n)$,

$$\begin{aligned} & \sum_{j=1}^m \frac{\lambda_j |B_j|}{\|\mathbf{1}_{B_j}\|_X} \left[\int_{B_j} |f(x) - P_{B_j}^{(s)}(f)(x)|^q dx \right]^{\frac{1}{q}} \\ & \lesssim \sum_{j=1}^m \frac{\lambda_j |B_j|}{\|\mathbf{1}_{B_j}\|_X} \left[\int_{B_j} |T_{(\widetilde{K})} \left(\left[f - P_{2B_j}^{(s)}(f) \right] \mathbf{1}_{2B_j} \right)(x)|^q dx \right]^{\frac{1}{q}} \\ & \quad + \sum_{j=1}^m \frac{\lambda_j |B_j|}{\|\mathbf{1}_{B_j}\|_X} \left[\int_{B_j} |E_{B_j}(x)|^q dx \right]^{\frac{1}{q}} \\ & =: \mathbf{I}_1 + \mathbf{I}_2, \end{aligned} \tag{3.13}$$

where E_{B_j} is the same as in (3.6) with \widetilde{B} therein replaced by B_j here.

Now, we separately estimate \mathbf{I}_1 and \mathbf{I}_2 . Indeed, by Remarks 3.3 and 3.9, we find that

$$\mathbf{I}_1 \lesssim \sum_{j=1}^m \frac{\lambda_j |2B_j|}{\|\mathbf{1}_{2B_j}\|_X} \left[\int_{2B_j} |f(x) - P_{2B_j}^{(s)}(f)(x)|^q dx \right]^{\frac{1}{q}}. \tag{3.14}$$

This is a desired estimate of \mathbf{I}_1 . Besides, for any $j \in \{1, \dots, m\}$ and $k \in \mathbb{N}$, let $\lambda_{j,k} := \frac{\lambda_j \|\mathbf{1}_{2^k B_j}\|_X}{\|\mathbf{1}_{B_j}\|_X}$. Then, from both (3.8) and Lemma 3.15 with $\lambda := s + \delta$, we deduce that, for any given $r \in (0, p_-)$,

$$\begin{aligned} \mathbf{I}_2 & \lesssim \sum_{j=1}^m \frac{\lambda_j |B_j|}{\|\mathbf{1}_{B_j}\|_X} r^{s+\delta} \int_{\mathbb{R}^n \setminus 2B_j} \frac{|f(y) - P_{2B_j}^{(s)}(f)(y)|}{|y - z_j|^{n+d+\delta}} dy \\ & \lesssim \sum_{j=1}^m \frac{\lambda_j |B_j|}{\|\mathbf{1}_{B_j}\|_X} \sum_{k \in \mathbb{N}} \frac{1}{2^{k(s+\delta)}} \left[\int_{2^{k+1} B_j} |f(y) - P_{2^{k+1} B_j}^{(s)}(f)(y)|^q dy \right]^{\frac{1}{q}} \\ & \sim \sum_{k \in \mathbb{N}} 2^{k(-n-s-\delta)} \sum_{j=1}^m \frac{\lambda_{j,k} |2^{k+1} B_j|}{\|\mathbf{1}_{2^{k+1} B_j}\|_X} \\ & \quad \times \left[\int_{2^{k+1} B_j} |f(y) - P_{2^{k+1} B_j}^{(s)}(f)(y)|^q dy \right]^{\frac{1}{q}}. \end{aligned} \tag{3.15}$$

This is a desired estimate of I_2 .

Moreover, by both Remark 3.9 and the definition of $\lambda_{i,k}$ with both $i \in \{1, \dots, m\}$ and $k \in \mathbb{N}$, we conclude that, for any given $r \in (0, \min\{d, p_-\})$ and for any $k \in \mathbb{N}$,

$$\begin{aligned} \left\| \left\{ \sum_{i=1}^m \left[\frac{\lambda_{i,k}}{\|\mathbf{1}_{2^k B_i}\|_X} \right]^d \mathbf{1}_{2^k B_i} \right\} \right\|_X^{\frac{1}{d}} &\lesssim 2^{\frac{kn}{r}} \left\| \left\{ \sum_{i=1}^m \left[\frac{\lambda_{i,k}}{\|\mathbf{1}_{2^k B_i}\|_X} \right]^d \mathbf{1}_{B_i} \right\} \right\|_X^{\frac{1}{d}} \\ &\sim 2^{\frac{kn}{r}} \left\| \left\{ \sum_{i=1}^m \left[\frac{\lambda_i}{\|\mathbf{1}_{B_i}\|_X} \right]^d \mathbf{1}_{B_i} \right\} \right\|_X^{\frac{1}{d}}. \end{aligned}$$

From this, (3.13), (3.14), (3.15), and $s \in (\frac{n}{\min\{p_-, d\}} - n - \delta, \infty)$ which further implies that $\min\{d, p_-\} \in (\frac{n}{n+s+\delta}, \infty)$, we deduce that, for any given $r \in (\frac{n}{n+s+\delta}, \min\{d, p_-\})$,

$$\begin{aligned} &\left\| \left\{ \sum_{i=1}^m \left[\frac{\lambda_i}{\|\mathbf{1}_{B_i}\|_X} \right]^d \mathbf{1}_{B_i} \right\} \right\|_X^{\frac{1}{d}} \left\| \sum_{j=1}^m \frac{\lambda_j |B_j|}{\|\mathbf{1}_{B_j}\|_X} \left[\frac{1}{|B_j|} \int_{B_j} |f(x) - P_{B_j}^{(s)}(f)(x)|^q dx \right]^{\frac{1}{q}} \right\|_X^{-1} \\ &\lesssim \left\| \left\{ \sum_{i=1}^m \left[\frac{\lambda_i}{\|\mathbf{1}_{2B_i}\|_X} \right]^d \mathbf{1}_{2B_i} \right\} \right\|_X^{\frac{1}{d}} \left\| \sum_{j=1}^m \frac{\lambda_j |2B_j|}{\|\mathbf{1}_{2B_j}\|_X} \left[\int_{2B_j} |f(x) - P_{2B_j}^{(s)}(f)(x)|^q dx \right]^{\frac{1}{q}} \right\|_X^{-1} \\ &\quad + \sum_{k \in \mathbb{N}} 2^{k(-n-s-\delta+\frac{n}{r})} \left\| \left\{ \sum_{j \in \mathbb{N}} \left[\frac{\lambda_j}{\|\mathbf{1}_{2^k B_j}\|_X} \right]^d \mathbf{1}_{2^k B_j} \right\} \right\|_X^{\frac{1}{d}} \left\| \sum_{j=1}^m \frac{\lambda_j |2^{k+1} B_j|}{\|\mathbf{1}_{2^{k+1} B_j}\|_X} \left[\int_{2^{k+1} B_j} |f(y) - P_{2^{k+1} B_j}^{(s)}(f)(y)|^q dy \right]^{\frac{1}{q}} \right\|_X^{-1} \\ &\lesssim \|f\|_{\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)}. \end{aligned}$$

This, together with the definition of $\|\cdot\|_{\mathcal{L}_{X,q,s,d}(\mathbb{R}^n)}$, then further implies the desired conclusion, which completes the proof of the sufficiency and hence Theorem 3.13. \square

3.2 Relations between \widetilde{T} and T

In this subsection, we show that \widetilde{T} is just the adjoint operator of T . To this end, we first recall the concept of the Hardy space $H_X(\mathbb{R}^n)$ associated with X , which was originally studied in [50].

In what follows, denote by $\mathcal{S}(\mathbb{R}^n)$ the space of all Schwartz functions equipped with the topology determined by a well-known countable family of norms, and by $\mathcal{S}'(\mathbb{R}^n)$ its topological dual space equipped with the weak-* topology. For any $N \in \mathbb{N}$ and $\phi \in \mathcal{S}(\mathbb{R}^n)$, let

$$p_N(\phi) := \sum_{\alpha \in \mathbb{Z}_+^n, |\alpha| \leq N} \sup_{x \in \mathbb{R}^n} (1 + |x|)^{N+n} |\partial^\alpha \phi(x)|$$

and

$$\mathcal{F}_N(\mathbb{R}^n) := \{\phi \in \mathcal{S}(\mathbb{R}^n) : p_N(\phi) \in [0, 1]\},$$

where, for any $\alpha := (\alpha_1, \dots, \alpha_n) \in \mathbb{Z}_+^n$, $|\alpha| := \alpha_1 + \dots + \alpha_n$ and $\partial^\alpha := (\frac{\partial}{\partial x_1})^{\alpha_1} \dots (\frac{\partial}{\partial x_n})^{\alpha_n}$. Moreover, for any $r \in \mathbb{R}$, we denote by $\lfloor r \rfloor$ (resp., $\lceil r \rceil$) the *maximal* (resp., *minimal*) integer not greater (resp., less) than r .

Definition 3.16. Let X be a ball quasi-Banach function space and $N \in \mathbb{N}$ be sufficiently large. Then the *Hardy space* $H_X(\mathbb{R}^n)$ is defined to be the set of all the $f \in \mathcal{S}'(\mathbb{R}^n)$ such that

$$\|f\|_{H_X(\mathbb{R}^n)} := \|\mathcal{M}_N(f)\|_X < \infty,$$

where the *non-tangential grand maximal function* $\mathcal{M}_N(f)$ of $f \in \mathcal{S}'(\mathbb{R}^n)$ is defined by setting, for any $x \in \mathbb{R}^n$,

$$\mathcal{M}_N(f)(x) := \sup \{|f * \phi_t(y)| : \phi \in \mathcal{F}_N(\mathbb{R}^n), t \in (0, \infty), |x - y| < t\}.$$

Remark 3.17. Let all the symbols be the same as in Definition 3.16. Assume that there exists an $r \in (0, \infty)$ such that the Hardy–Littlewood maximal operator \mathcal{M} is bounded on $X^{\frac{1}{r}}$. If $N \in [\lfloor \frac{r}{r-1} \rfloor + 1, \infty) \cap \mathbb{N}$, then, by [50, Theorem 3.1], we find that the Hardy space $H_X(\mathbb{R}^n)$ is independent of the choice of N .

To obtain the desired result, we still need to recall some basic concepts. In what follows, for any $\theta \in (0, \infty)$, the *powered Hardy–Littlewood maximal operator* $\mathcal{M}^{(\theta)}$ is defined by setting, for any measurable function f and $x \in \mathbb{R}^n$,

$$\mathcal{M}^{(\theta)}(f)(x) := \left\{ \mathcal{M}(|f|^\theta)(x) \right\}^{\frac{1}{\theta}}.$$

Moreover, the associate space X' of any given ball Banach function space X is defined as follows (see [6, Chapter 1, Section 2] or [50, p. 9]).

Definition 3.18. For any given ball quasi-Banach function space X , its *associate space* (also called the *Köthe dual space*) X' is defined by setting

$$X' := \{f \in \mathcal{M}(\mathbb{R}^n) : \|f\|_{X'} < \infty\},$$

where, for any $f \in X'$,

$$\|f\|_{X'} := \sup \{\|fg\|_{L^1(\mathbb{R}^n)} : g \in X, \|g\|_X = 1\}$$

and $\|\cdot\|_{X'}$ is called the *associate norm* of $\|\cdot\|_X$.

Next, we recall the following mild assumption about the boundedness of the powered Hardy–Littlewood maximal operator on ball quasi-Banach function spaces (see, for instance, [50, (2.9)]), which is used later in this article.

Assumption 3.19. Let X be a ball quasi-Banach function space. Assume that there exists an $r_0 \in (0, \infty)$ and a $p_0 \in (r_0, \infty)$ such that X^{1/r_0} is a ball Banach function space and there exists a positive constant C such that, for any $f \in (X^{1/r_0})'$,

$$\|\mathcal{M}^{((p_0/r_0)')} (f)\|_{(X^{1/r_0})'} \leq C \|f\|_{(X^{1/r_0})'}.$$

Now, we present the following definitions of both the (X, q, s) -atom and the finite atomic Hardy space $H_{\text{fin}}^{X,q,s,d}(\mathbb{R}^n)$ which are just, respectively, [50, Definition 3.5] and [67, Definition 1.9].

Definition 3.20. Let X be a ball quasi-Banach function space, $q \in (1, \infty]$, and $s \in \mathbb{Z}_+$. Then a measurable function a on \mathbb{R}^n is called an (X, q, s) -atom if there exists a ball $B \in \mathbb{B}(\mathbb{R}^n)$ such that

- (i) $\text{supp}(a) := \{x \in \mathbb{R}^n : a(x) \neq 0\} \subset B$;
- (ii) $\|a\|_{L^q(\mathbb{R}^n)} \leq \frac{|B|^{\frac{1}{q}}}{\|\mathbf{1}_B\|_X}$;
- (iii) $\int_{\mathbb{R}^n} a(x)x^\gamma dx = 0$ for any $\gamma := (\gamma_1, \dots, \gamma_n) \in \mathbb{Z}_+^n$ with $|\gamma| \leq s$.

Definition 3.21. Let both X and p_- satisfy Assumption 3.8. Assume that $r_0 \in (0, \min\{1, p_-\})$ and $p_0 \in (r_0, \infty)$ satisfy Assumption 3.19. Let $s \in [\lfloor n(\frac{1}{\min\{1, p_-}} - 1) \rfloor, \infty) \cap \mathbb{Z}_+$, $d \in (0, r_0]$, and $q \in (\max\{1, p_0\}, \infty]$. The *finite atomic Hardy space* $H_{\text{fin}}^{X,q,s,d}(\mathbb{R}^n)$, associated with X , is defined to be the set of all finite linear combinations of (X, q, s) -atoms. The quasi-norm $\|\cdot\|_{H_{\text{fin}}^{X,q,s,d}(\mathbb{R}^n)}$ in $H_{\text{fin}}^{X,q,s,d}(\mathbb{R}^n)$ is defined by setting, for any $f \in H_{\text{fin}}^{X,q,s,d}(\mathbb{R}^n)$,

$$\|f\|_{H_{\text{fin}}^{X,q,s,d}(\mathbb{R}^n)} := \inf \left\{ \left\| \left[\sum_{j=1}^N \left(\frac{\lambda_j}{\|\mathbf{1}_{B_j}\|_X} \right)^d \mathbf{1}_{B_j} \right]^{\frac{1}{d}} \right\|_X \right\},$$

where the infimum is taken over all finite linear combinations of (X, q, s) -atoms of f , namely, $N \in \mathbb{N}$, $f = \sum_{j=1}^N \lambda_j a_j$, $\{\lambda_j\}_{j=1}^N \subset [0, \infty)$, and $\{a_j\}_{j=1}^N$ being (X, q, s) -atoms supported, respectively, in the balls $\{B_j\}_{j=1}^N \subset \mathbb{B}(\mathbb{R}^n)$.

Recall that X is said to have an *absolutely continuous quasi-norm* if, for any $f \in X$ and any measurable subsets $\{E_j\}_{j \in \mathbb{N}} \subset \mathbb{R}^n$ with $E_{j+1} \subset E_j$ for any $j \in \mathbb{N}$ and $\bigcap_{j \in \mathbb{N}} E_j = \emptyset$, $\|f \mathbf{1}_{E_j}\|_X \downarrow 0$ as $j \rightarrow \infty$. The following lemma is just [69, Theorem 3.14].

Lemma 3.22. *Let X , d , q , and s be the same as in Definition 3.21, and let X have an absolutely continuous quasi-norm. Then the dual space of $H_X(\mathbb{R}^n)$, denoted by $(H_X(\mathbb{R}^n))^*$, is $\mathcal{L}_{X,q',s,d}(\mathbb{R}^n)$ with $\frac{1}{q} + \frac{1}{q'} = 1$ in the following sense:*

- (i) *Let $g \in \mathcal{L}_{X,q',s,d}(\mathbb{R}^n)$. Then the linear functional*

$$L_g : f \rightarrow \langle L_g, f \rangle := \int_{\mathbb{R}^n} f(x)g(x) dx, \quad (3.16)$$

initially defined for any $f \in H_{\text{fin}}^{X,q,s,d}(\mathbb{R}^n)$, has a bounded extension to $H_X(\mathbb{R}^n)$.

- (ii) *Conversely, any continuous linear functional on $H_X(\mathbb{R}^n)$ arises the same as in (3.16) with a unique $g \in \mathcal{L}_{X,q',s,d}(\mathbb{R}^n)$.*

Moreover, $\|g\|_{\mathcal{L}_{X,q',s,d}(\mathbb{R}^n)} \sim \|L_g\|_{(H_X(\mathbb{R}^n))^}$ with the positive equivalence constants independent of both g and L_g .*

The following proposition is just a slight modification of [61, Theorem 3.11], which shows that T has a unique extension on $H_X(\mathbb{R}^n)$; we omit the details here.

Proposition 3.23. *Let both X and p_- satisfy Assumptions 3.8 and 3.19. Let $\delta \in (0, 1]$, $s \in (\frac{n}{p_-} - n - \delta, \infty)$, and T be the same as in Definition 3.4. Further assume that X has an absolutely continuous quasi-norm. Then T has a unique extension on $H_X(\mathbb{R}^n)$, still denoted by T , namely, there exists a positive constant C such that, for any $f \in H_X(\mathbb{R}^n)$,*

$$\|T(f)\|_{H_X(\mathbb{R}^n)} \leq C\|f\|_{H_X(\mathbb{R}^n)}.$$

Next, we present the main result of this subsection.

Theorem 3.24. *Let X , p_- , q , and d be the same as in Definition 3.21. Let $\delta \in (0, 1]$,*

$$s \in \left(\frac{n}{d} - n - \delta, \infty\right) \cap \left[\left\lceil n \left(\frac{1}{\min\{1, p_-\}} - 1\right) \right\rceil, \infty\right) \cap \mathbb{Z}_+,$$

T be the same as in Proposition 3.23, and \widetilde{T} the same as in Remark 3.7. Further assume that X has an absolutely continuous quasi-norm. Then \widetilde{T} is the adjoint operator of T in the following sense: for any $g \in \mathcal{L}_{X, q', s, d}(\mathbb{R}^n)$ and $f \in H_X(\mathbb{R}^n)$,

$$\langle L_g, T(f) \rangle = \langle L_{\widetilde{T}(g)}, f \rangle,$$

where L_g is the same as in (3.16), $L_{\widetilde{T}(g)}$ the same as in (3.16) with g replaced by $\widetilde{T}(g)$, and $\frac{1}{q} + \frac{1}{q'} = 1$.

To prove this theorem, we first establish three technique lemmas. We begin with showing that, for any (X, q, s) -atom a , $T(a)$ is a harmless constant multiple of an (X, q, s, τ) -molecule if and only if T has the vanishing moments up to order s . To this end, we first recall the following definition of (X, q, s, τ) -molecules, which is just [50, Definition 3.8].

Definition 3.25. Let X be a ball quasi-Banach function space, $q \in (1, \infty]$, $s \in \mathbb{Z}_+$, and $\tau \in (0, \infty)$. Then a measurable function m on \mathbb{R}^n is called an (X, q, s, τ) -molecule centered at a ball $B \in \mathbb{B}(\mathbb{R}^n)$ if

- (i) for any $j \in \mathbb{Z}_+$,

$$\|m \mathbf{1}_{L_j}\|_{L^q(\mathbb{R}^n)} \leq 2^{-j\tau} \frac{|B|^{\frac{1}{q}}}{\|\mathbf{1}_B\|_X},$$

where $L_0 := B$ and, for any $j \in \mathbb{N}$, $L_j := 2^j B \setminus 2^{j-1} B$;

- (ii) $\int_{\mathbb{R}^n} m(x) x^\gamma dx = 0$ for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \leq s$.

Lemma 3.26. *Let both X and p_- satisfy Assumption 3.8. Let $s \in \mathbb{Z}_+$, $\delta \in (0, 1]$, T be the same as in Definition 3.1, $q \in (1, \infty)$, $\frac{1}{q} + \frac{1}{q'} = 1$, and $\tau \in (0, \frac{n}{q} + s + \delta]$. Then the following two statements are equivalent:*

- (i) *there exists a positive constant C such that, for any (X, q, s) -atom a supported in the ball $B \in \mathbb{B}(\mathbb{R}^n)$, $\frac{T(a)}{C}$ is an (X, q, s, τ) -molecule centered at $2B$.*
- (ii) *T has the vanishing moments up to order s .*

Proof. We first show (i) \implies (ii). To this end, let $A \in L^q(\mathbb{R}^n)$ be supported in a ball $B \in \mathbb{B}(\mathbb{R}^n)$ and satisfy that, for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \leq s$, $\int_{\mathbb{R}^n} A(x)x^\gamma dx = 0$. Without loss of generality, we may assume that A satisfies that $\|A\|_{L^q(\mathbb{R}^n)} > 0$. Then it is easy to show that $\tilde{A} := \frac{|B|^{\frac{1}{q}}A}{\|\mathbf{1}_B\|_X \|A\|_{L^q(\mathbb{R}^n)}}$ is an (X, q, s) -atom supported in B . By this, (i) of the present lemma, and [37, (2.17)], we conclude that, for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \leq s$,

$$\int_{B(x_0, r_0)} \tilde{A}(x) \tilde{T}(y^\gamma)(x) dx = \int_{\mathbb{R}^n} T(\tilde{A})(x) x^\gamma dx = 0.$$

Applying this and an argument similar to that used in the proof of [37, Proposition 2.16], we find that, for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \leq s$, $\tilde{T}(y^\gamma) \in \mathcal{P}_s(\mathbb{R}^n)$ after changing values on a set of measure zero, which, combined with Proposition 3.14, further implies that T has the vanishing moments up to order s . This finishes the proof that (i) \implies (ii).

Next, we show (ii) \implies (i). Let a be an (X, q, s) -atom supported in the ball $B \in \mathbb{B}(\mathbb{R}^n)$. We prove that there exists a positive constant C , independent of a , such that $\frac{T(a)}{C}$ is an (X, q, s, τ) -molecule centered at $2B$. Indeed, using Remark 3.3, the size condition of a , and (3.12), we find that

$$\|T(a)\mathbf{1}_{2B}\|_{L^q(\mathbb{R}^n)} \lesssim \|a\|_{L^q(B)} \lesssim \frac{|B|^{\frac{1}{q}}}{\|\mathbf{1}_B\|_X} \sim \frac{|2B|^{\frac{1}{q}}}{\|\mathbf{1}_{2B}\|_X} \quad (3.17)$$

with the implicit positive constants independent of a . Moreover, similarly to the estimation of $T(a)$ in [37, p. 48], we find that, for any $x \notin 2B$,

$$|T(a)(x)| \lesssim \frac{r^{\frac{n}{q}+s+\delta}}{|z-x|^{n+s+\delta}} \|a\|_{L^q(B)},$$

which, together with both the size condition of a and $\tau \in (0, \frac{n}{q} + s + \delta]$, further implies that, for any $j \in \mathbb{N}$ and $x \in 2^{j+1}B \setminus 2^jB$,

$$|T(a)(x)| \lesssim 2^{-j(\frac{n}{q}+s+\delta)} (2^j r)^{-\frac{n}{q}} \frac{|B|^{\frac{1}{q}}}{\|\mathbf{1}_B\|_X} \lesssim 2^{-j\tau} |2^{j+1}B \setminus 2^jB|^{-\frac{1}{q}} \frac{|B|^{\frac{1}{q}}}{\|\mathbf{1}_B\|_X}.$$

From this and (3.12), we infer that, for any $j \in \mathbb{N}$,

$$\|T(a)\mathbf{1}_{2^{j+1}B \setminus 2^jB}\|_{L^q(\mathbb{R}^n)} \lesssim 2^{-j\tau} \frac{|B|^{\frac{1}{q}}}{\|\mathbf{1}_B\|_X} \sim 2^{-j\tau} \frac{|2B|^{\frac{1}{q}}}{\|\mathbf{1}_{2B}\|_X}.$$

This, combined with both (3.17) and (ii) of the present lemma, then finishes the proof that (ii) \implies (i) and hence the proof of Lemma 3.26. \square

The following lemma is just [10, Lemma 3.30].

Lemma 3.27. *Let X , p_- , d , q , and s be the same as in Definition 3.21. Let*

$$\tau \in \left(n \left[\frac{1}{p_-} - \frac{1}{q} \right], \infty \right) \cap \left(\frac{n}{q'} + s, \infty \right),$$

where $\frac{1}{q} + \frac{1}{q'} = 1$. Further assume that X has an absolutely continuous quasi-norm. Then, for any (X, q, s, τ) -molecule m and any $g \in \mathcal{L}_{X, q', s, d}(\mathbb{R}^n)$,

$$\langle L_g, m \rangle = \int_{\mathbb{R}^n} m(x)g(x) dx,$$

where L_g is the same as in (3.16).

The proof of the following lemma is a slight modification of [37, Lemma 3.19] and we only sketch some important steps.

Lemma 3.28. *Let X , p_- , d , and q be the same as in Definition 3.21. Let $\delta \in (0, 1]$,*

$$s \in \left(\frac{n}{d} - n - \delta, \infty \right) \cap \left[\left[n \left(\frac{1}{\min\{1, p_-\}} - 1 \right) \right], \infty \right) \cap \mathbb{Z}_+,$$

T be the same as in Definition 3.4, and \tilde{T} the same as in Remark 3.7. Further assume that X has an absolutely continuous quasi-norm. Then, for any $g \in \mathcal{L}_{X, q', s, d}(\mathbb{R}^n)$ and any (X, q, s) -atom a ,

$$\langle L_g, T(a) \rangle = \langle L_{\tilde{T}(g)}, a \rangle,$$

where L_g is the same as in (3.16), $L_{\tilde{T}(g)}$ the same as in (3.16) with g replaced by $\tilde{T}(g)$, and $\frac{1}{q} + \frac{1}{q'} = 1$.

Proof. By the ranges of both d and s , we have $d \in (\frac{n}{n+s+\delta}, \infty)$ and $p_- \in (\frac{n}{n+s+\delta}, \infty)$. Thus, we can choose a $\tau \in (n[\frac{1}{\min\{1, p_-\}} - \frac{1}{q}], \frac{n}{q'} + s + \delta] \cap (\frac{n}{q'} + s, \frac{n}{q'} + s + \delta]$. Then, using Lemma 3.26, we conclude that $T(a)$ is a harmless constant multiple of an (X, q, s, τ) -molecule. By this and Lemma 3.27, we find that, for any $g \in \mathcal{L}_{X, q', s, d}(\mathbb{R}^n)$ and any (X, q, s) -atom a ,

$$\langle L_g, T(a) \rangle = \int_{\mathbb{R}^n} g(x)T(a)(x) dx. \quad (3.18)$$

Besides, using both $\min\{d, p_-\} \in (\frac{n}{n+s+\delta}, \infty)$ and Theorem 3.12, we obtain $\tilde{T}(g) \in \mathcal{L}_{X, q', s, d}(\mathbb{R}^n)$, which, combined with Lemma 3.22, further implies that, for any (X, q, s) -atom a ,

$$\langle L_{\tilde{T}(g)}, a \rangle = \int_{\mathbb{R}^n} \tilde{T}(g)(x)a(x) dx. \quad (3.19)$$

Moreover, by Lemma 3.10 with $\lambda := n + s + \delta$, we conclude that, for any $g \in \mathcal{L}_{X, q', s, d}(\mathbb{R}^n)$ and any ball $B := B(x_0, r_0) \in \mathbb{B}(\mathbb{R}^n)$ with $x_0 \in \mathbb{R}^n$ and $r_0 \in (0, \infty)$,

$$\int_{\mathbb{R}^n \setminus B} \frac{|g(y) - P_B^{(s)}(g)(y)|}{|x_0 - y|^{n+s+\delta}} dy < \infty. \quad (3.20)$$

Using (3.20) and repeating the proof of [37, Lemma 3.19], we conclude that

$$\int_{\mathbb{R}^n} g(x)T(a)(x) dx = \int_{\mathbb{R}^n} \tilde{T}(g)(x)a(x) dx,$$

which, together with both (3.18) and (3.19), further implies that

$$\langle L_g, T(a) \rangle = \int_{\mathbb{R}^n} T(a)g(x) dx = \int_{\mathbb{R}^n} \tilde{T}(g)(x)a(x) dx = \langle L_{\tilde{T}(g)}, a \rangle.$$

This finishes the proof of Lemma 3.28. \square

Now, we show Theorem 3.24.

Proof of Theorem 3.24. To show the present theorem, let $f \in H_X(\mathbb{R}^n)$. Since X has an absolutely continuous quasi-norm, using this and [50, Remark 3.12], we find that $H_{\text{fin}}^{X, \infty, s, d}(\mathbb{R}^n)$ is dense in $H_X(\mathbb{R}^n)$. Thus, there exists a sequence $\{f_k\}_{k \in \mathbb{N}} \subset H_{\text{fin}}^{X, \infty, s, d}(\mathbb{R}^n)$ such that $f_k \rightarrow f$ in $H_X(\mathbb{R}^n)$. Besides, by the definition of $H_{\text{fin}}^{X, \infty, s, d}(\mathbb{R}^n)$, we conclude that, for any $k \in \mathbb{N}$, there exists an $m_k \in \mathbb{N}$, a sequence $\{a_j^{(k)}\}_{j=1}^{m_k}$ of (X, ∞, s) -atoms, and $\{\lambda_j^{(k)}\}_{j=1}^{m_k} \subset [0, \infty)$ such that

$$f_k = \sum_{j=1}^{m_k} \lambda_j^{(k)} a_j^{(k)}. \quad (3.21)$$

Moreover, combining this and Proposition 3.23, we have $T(f_k) \rightarrow T(f)$ in $H_X(\mathbb{R}^n)$ as $k \rightarrow \infty$. From this, Lemma 3.22, (3.21), and Lemmas 3.28 and 3.13, we deduce that, for any $g \in \mathcal{L}_{X, q', s, d}(\mathbb{R}^n)$,

$$\begin{aligned} \langle L_g, T(f) \rangle &= \lim_{k \rightarrow \infty} \langle L_g, T(f_k) \rangle = \lim_{k \rightarrow \infty} \sum_{j=1}^{m_k} \lambda_j^{(k)} \langle L_g, T(a_j^{(k)}) \rangle \\ &= \lim_{k \rightarrow \infty} \sum_{j=1}^{m_k} \lambda_j^{(k)} \langle L_{\tilde{T}(g)}, a_j^{(k)} \rangle = \lim_{k \rightarrow \infty} \langle L_{\tilde{T}(g)}, f_k \rangle = \langle L_{\tilde{T}(g)}, f \rangle. \end{aligned}$$

This finishes the proof of Theorem 3.24. \square

4 Applications

In this section, we apply all the above main results to eight concrete examples of ball quasi-Banach function spaces, namely, weighted Lebesgue spaces (Subsection 4.1), variable Lebesgue spaces (Subsection 4.2), Orlicz spaces (Subsection 4.3), Orlicz-slice spaces (Subsection 4.4), Morrey spaces (Subsection 4.5), mixed-norm Lebesgue spaces (Subsection 4.6), local generalized Herz spaces (Subsection 4.7), and mixed Herz spaces (Subsection 4.8). Moreover, to the best of our knowledge, all these results are new. These applications explicitly indicate the generality and the practicability of the main results of this article and more applications to new function spaces are obviously possible.

4.1 Weighted Lebesgue spaces

In this subsection, we apply Theorems 3.12, 3.13, and 3.24 to weighted Lebesgue spaces. We first present the definitions of both Muckenhoupt weights and weighted Lebesgue spaces (see, for instance, [23, Definitions 7.1.2 and 7.1.3]).

Definition 4.1. Let $p \in [1, \infty)$ and w be a nonnegative locally integrable function on \mathbb{R}^n . Then w is called an $A_p(\mathbb{R}^n)$ weight, denoted by $w \in A_p(\mathbb{R}^n)$, if, when $p \in (1, \infty)$,

$$[w]_{A_p(\mathbb{R}^n)} := \sup_{B \in \mathbb{B}(\mathbb{R}^n)} \left[\frac{1}{|B|} \int_B w(x) dx \right] \left\{ \frac{1}{|B|} \int_B [w(x)]^{-\frac{1}{p-1}} dx \right\}^{p-1} < \infty,$$

and

$$[w]_{A_1(\mathbb{R}^n)} := \sup_{B \in \mathbb{B}(\mathbb{R}^n)} \left[\frac{1}{|B|} \int_B w(x) dx \right] \left\{ \operatorname{ess\,sup}_{x \in \mathbb{R}^n} [w(x)]^{-1} \right\} < \infty.$$

Moreover, the class $A_\infty(\mathbb{R}^n)$ is defined by setting

$$A_\infty(\mathbb{R}^n) := \bigcup_{p \in [1, \infty)} A_p(\mathbb{R}^n).$$

Definition 4.2. Let $p \in (0, \infty)$ and $w \in A_\infty(\mathbb{R}^n)$. The weighted Lebesgue space $L_w^p(\mathbb{R}^n)$ is defined to be the set of all the measurable functions f on \mathbb{R}^n such that

$$\|f\|_{L_w^p(\mathbb{R}^n)} = \left[\int_{\mathbb{R}^n} |f(x)|^p w(x) dx \right]^{\frac{1}{p}} < \infty.$$

Remark 4.3. It is worth pointing out that, for any $p \in (0, \infty)$ and $w \in A_\infty(\mathbb{R}^n)$, $L_w^p(\mathbb{R}^n)$ is a ball quasi-Banach space but it may not be a quasi-Banach function space (see, for instance, [50, Section 7.1]).

Using Theorems 3.12, 3.13, and 3.24, we have the following conclusion.

Theorem 4.4. Let $p \in (0, \infty)$ and $w \in A_\infty(\mathbb{R}^n)$. Then both Theorems 3.12 and 3.13 with both $X := L_w^p(\mathbb{R}^n)$ and $p_- := \frac{p}{q_w}$ hold true; moreover, Theorem 3.24 with $X := L_w^p(\mathbb{R}^n)$, $p_- := \frac{p}{q_w}$, $q \in (\max\{1, \frac{p}{q_w}, p\}, \infty)$, $d \in (0, \min\{1, \frac{p}{q_w}\})$, and

$$s \in \left(\frac{n}{d} - n - \delta, \infty \right) \cap \left[\left[n \left(\frac{1}{\min\{1, p_-\}} - 1 \right) \right], \infty \right) \cap \mathbb{Z}_+$$

holds true.

Proof. It is easy to show that $L_w^p(\mathbb{R}^n)$ is a ball quasi-Banach space with an absolutely continuous quasi-norm. Moreover, from a claim in [8, p. 28], we deduce that both Assumptions 3.8 and 3.19 with $X := L_w^p(\mathbb{R}^n)$, $p_- := \frac{p}{q_w}$, $r_0 \in (d, \frac{p}{q_w})$, and $p_0 \in (\max\{\frac{p}{q_w}, p\}, q)$ hold true. Thus, all the assumptions of Theorems 3.12, 3.13, and 3.24 with $X := L_w^p(\mathbb{R}^n)$ are satisfied. Then, by Theorems 3.12, 3.13, and 3.24 with $X := L_w^p(\mathbb{R}^n)$, we obtain the desired conclusions, which completes the proof of Theorem 4.4. \square

Remark 4.5. (i) To the best of our knowledge, Theorem 4.4 is new.

(ii) Moreover, using [49, Theorem 246], we find that, for any $\alpha \in (0, \infty)$ and $q \in [1, \infty]$, $C_{\alpha, q, \lfloor \alpha n \rfloor}(\mathbb{R}^n) = \dot{C}^{\alpha n}(\mathbb{R}^n)$ with equivalent norms, where $\dot{C}^{\alpha n}(\mathbb{R}^n)$ is the homogeneous Hölder–Zygmund space (see, for instance, [49, Section 1.6.2]). Let $q \in (1, \infty)$, $s \in \mathbb{Z}_+$, $\alpha \in (0, \infty)$, $w \equiv 1$, and $X := L_w^{\frac{1}{\alpha+1}}(\mathbb{R}^n) = L^{\frac{1}{\alpha+1}}(\mathbb{R}^n)$. From Remark 2.5, we infer that, under these assumptions, $\mathcal{L}_{X, q, s}(\mathbb{R}^n)$ coincides with the Campanato space $C_{\alpha, q, s}(\mathbb{R}^n)$. Then, by Theorem 4.4, we conclude that, under these assumptions, the conclusions of Theorem 3.12 holds true with $X := L^{\frac{1}{\alpha+1}}(\mathbb{R}^n)$. This further implies that, under these assumptions, Theorem 3.12 with $X := L^{\frac{1}{\alpha+1}}(\mathbb{R}^n)$ coincides with [40, Theorem 4.21]. From these observations, we infer the following conclusion: For any $\alpha \in (0, \infty)$, if letting $\delta \in (\alpha n - \lfloor \alpha n \rfloor, 1]$ and \tilde{T} be the same as in Theorem 3.12, then \tilde{T} is bounded on $\dot{C}^{\alpha n}(\mathbb{R}^n)$ if and only if, for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \leq \lfloor \alpha n \rfloor$, $T^*(x^\gamma) = 0$, which is the known best possible result on the boundedness of Calderón–Zygmund operators on homogeneous Hölder–Zygmund spaces.

4.2 Variable Lebesgue spaces

In this subsection, we apply Theorems 3.12, 3.13, and 3.24 to variable Lebesgue spaces. We now recall the concept of variable Lebesgue spaces (see, for instance, [15, p. 4]). Moreover, we refer the reader to [15, 19, 39] for more studies on variable Lebesgue spaces.

Definition 4.6. Let $p(\cdot) : \mathbb{R}^n \rightarrow [0, \infty)$ be a measurable function. The *variable Lebesgue space* $L^{p(\cdot)}(\mathbb{R}^n)$ is defined to be the set of all the measurable functions f on \mathbb{R}^n such that

$$\|f\|_{L^{p(\cdot)}(\mathbb{R}^n)} := \inf \left\{ \lambda \in (0, \infty) : \int_{\mathbb{R}^n} \left[\frac{|f(x)|}{\lambda} \right]^{p(x)} dx \leq 1 \right\} < \infty.$$

Moreover, $p(\cdot)$ is said to be *globally log-Hölder continuous* if there exists a $p_\infty \in \mathbb{R}$ such that, for any $x, y \in \mathbb{R}^n$,

$$|p(x) - p(y)| \lesssim \frac{1}{\log(e + 1/|x - y|)} \text{ and } |p(x) - p_\infty| \lesssim \frac{1}{\log(e + |x|)},$$

where the implicit positive constants are independent of both x and y .

Remark 4.7. Let $p(\cdot) : \mathbb{R}^n \rightarrow [0, \infty)$ be a measurable function. It is easy to show that $L^{p(\cdot)}(\mathbb{R}^n)$ is a ball quasi-Banach space but it may not be a quasi-Banach function space (see, for instance, [50, Section 7.1]).

The following theorem is a corollary of Theorems 3.12, 3.13, and 3.24.

Theorem 4.8. Let $p(\cdot) : \mathbb{R}^n \rightarrow (0, \infty)$ be a globally log-Hölder continuous function satisfying $0 < \text{ess inf}_{x \in \mathbb{R}^n} p(x) \leq \text{ess sup}_{x \in \mathbb{R}^n} p(x) < \infty$. Then both Theorems 3.12 and 3.13 with both $X := L^{p(\cdot)}(\mathbb{R}^n)$ and $p_- := \text{ess inf}_{x \in \mathbb{R}^n} p(x)$ hold true; moreover, Theorem 3.24 with $X := L^{\vec{p}}(\mathbb{R}^n)$, $p_- := \text{ess inf}_{x \in \mathbb{R}^n} p(x)$, $q \in (\max\{1, \text{ess sup}_{x \in \mathbb{R}^n} p(x)\}, \infty)$, $d \in (0, \min\{1, \text{ess inf}_{x \in \mathbb{R}^n} p(x)\})$, and

$$s \in \left(\frac{n}{d} - n - \delta, \infty \right) \cap \left[\left[n \left(\frac{1}{\min\{1, p_-\}} - 1 \right) \right], \infty \right) \cap \mathbb{Z}_+$$

holds true.

Proof. It is easy to show that $L^{p(\cdot)}(\mathbb{R}^n)$ is a ball quasi-Banach space with an absolutely continuous quasi-norm. Moreover, using a claim in [69, p.36], we find that both Assumptions 3.8 and 3.19 with $X := L^{p(\cdot)}(\mathbb{R}^n)$, $p_- := \operatorname{ess\,inf}_{x \in \mathbb{R}^n} p(x)$, $r_0 \in (d, \operatorname{ess\,inf}_{x \in \mathbb{R}^n} p(x))$, and $p_0 \in (\operatorname{ess\,sup}_{x \in \mathbb{R}^n} p(x), q)$ hold true. Thus, all the assumptions of Theorems 3.12, 3.13, and 3.24 with $X := L^{p(\cdot)}(\mathbb{R}^n)$ are satisfied. Then, from Theorems 3.12, 3.13, and 3.24 with $X := L^{p(\cdot)}(\mathbb{R}^n)$, we deduce the desired conclusions, which completes the proof of Theorem 4.8. \square

Remark 4.9. To the best of our knowledge, Theorem 4.8 is new.

4.3 Orlicz spaces

In this subsection, we apply all the main results of this article to Orlicz spaces. Recall that there exist many operators that are not bounded on Lebesgue spaces $L^p(\mathbb{R}^n)$ with $p \in [1, \infty)$, especially on both $L^1(\mathbb{R}^n)$ and $L^\infty(\mathbb{R}^n)$. The Orlicz space was introduced to cover the failure of the boundedness of some integral operators. We refer the reader to [2, 28, 34, 53] for more studies on this.

A non-decreasing function $\Phi : [0, \infty) \rightarrow [0, \infty)$ is called an *Orlicz function* if it satisfies $\Phi(0) = 0$, $\Phi(t) > 0$ whenever $t \in (0, \infty)$, and $\lim_{t \rightarrow \infty} \Phi(t) = \infty$. An Orlicz function Φ is said to be of *lower* (resp., *upper*) *type* p with $p \in \mathbb{R}$ if there exists a positive constant $C_{(p)}$, depending on p , such that, for any $t \in [0, \infty)$ and $s \in (0, 1)$ [resp., $s \in [1, \infty)$],

$$\Phi(st) \leq C_{(p)} s^p \Phi(t).$$

In addition, an Orlicz function Φ is said to be of *positive lower* (resp., *upper*) *type* if it is of lower (resp., upper) type p for some $p \in (0, \infty)$.

Definition 4.10. Let Φ be an Orlicz function with lower type $p_\Phi^- \in \mathbb{R}$ and upper type $p_\Phi^+ \in \mathbb{R}$. The *Orlicz space* $L^\Phi(\mathbb{R}^n)$ is defined to be the set of all the measurable functions f on \mathbb{R}^n such that

$$\|f\|_{L^\Phi(\mathbb{R}^n)} := \inf \left\{ \lambda \in (0, \infty) : \int_{\mathbb{R}^n} \Phi \left(\frac{|f(x)|}{\lambda} \right) dx \leq 1 \right\} < \infty.$$

Remark 4.11. Let Φ be an Orlicz function on \mathbb{R}^n with positive lower type $p_\Phi^- \in (0, \infty)$ and positive upper type $p_\Phi^+ \in (0, \infty)$. It has been pointed out in [50, Section 7.6] that $L^\Phi(\mathbb{R}^n)$ is a quasi-Banach function space and hence a ball quasi-Banach function space.

Using Theorems 3.12, 3.13, and 3.24, we have the following conclusion.

Theorem 4.12. *Let Φ be an Orlicz function with positive lower type p_Φ^- and positive upper type p_Φ^+ satisfying $0 < p_\Phi^- \leq p_\Phi^+ < \infty$. Then both Theorems 3.12 and 3.13 with both $X := L^\Phi(\mathbb{R}^n)$ and $p_- := p_\Phi^-$ hold true; moreover, Theorem 3.24 with $X := L^\Phi(\mathbb{R}^n)$, $p_- := p_\Phi^-$, $q \in (\max\{1, p_\Phi^+\}, \infty)$, $d \in (0, \min\{1, p_\Phi^-\})$, and*

$$s \in \left(\frac{n}{d} - n - \delta, \infty \right) \cap \left[\left\lceil n \left(\frac{1}{\min\{1, p_\Phi^-\}} - 1 \right) \right\rceil, \infty \right) \cap \mathbb{Z}_+$$

holds true.

Proof. Similarly to the proof of [71, Lemma 4.5], we find that $L^\Phi(\mathbb{R}^n)$ is a ball quasi-Banach space with an absolutely continuous quasi-norm. Moreover, by the claim in [69, p. 38], we conclude that both Assumptions 3.8 and 3.19 with $X := L^\Phi(\mathbb{R}^n)$, $p_- := p_\Phi^-$, $r_0 \in (d, p_\Phi^-)$, and $p_0 \in (p_\Phi^+, q)$ hold true. Thus, all the assumptions of Theorems 3.12, 3.13, and 3.24 with $X := L^\Phi(\mathbb{R}^n)$ are satisfied. Then, using Theorems 3.12, 3.13, and 3.24 with $X := L^\Phi(\mathbb{R}^n)$, we obtain the desired conclusions, which completes the proof of Theorem 4.12. \square

Remark 4.13. To the best of our knowledge, Theorem 4.12 is new.

4.4 Orlicz-Slice spaces

Recall that the Orlicz-slice space was introduced by Zhang et al. [71], which is a generalization of the slice spaces proposed by Auscher and Mourgoglou [3] and Auscher and Prisuelos-Arribas [4]. We refer the reader to [70] for more studies on the local Orlicz-slice space. To be precise, we present the definition of the Orlicz-slice space as follows.

Definition 4.14. Let $t, r \in (0, \infty)$ and Φ be an Orlicz function on \mathbb{R}^n with positive lower type p_Φ^- and positive upper type p_Φ^+ . The *Orlicz-slice space* $(E_\Phi^r)_t(\mathbb{R}^n)$ is defined to be the set of all the measurable functions f on \mathbb{R}^n such that

$$\|f\|_{(E_\Phi^r)_t(\mathbb{R}^n)} := \left\{ \int_{\mathbb{R}^n} \left[\frac{\|f \mathbf{1}_{B(x,t)}\|_{L^\Phi(\mathbb{R}^n)}}{\|\mathbf{1}_{B(x,t)}\|_{L^\Phi(\mathbb{R}^n)}} \right]^r dx \right\}^{\frac{1}{r}} < \infty.$$

Remark 4.15. Let $t, r \in (0, \infty)$ and Φ be an Orlicz function on \mathbb{R}^n with $0 < p_\Phi^- \leq p_\Phi^+ < \infty$. By [71, Lemma 2.28], we find that $(E_\Phi^r)_t(\mathbb{R}^n)$ is a ball quasi-Banach space, but it may not be a quasi-Banach function space (see, for instance, [72, Remark 7.43(i)]).

The following theorem is a corollary of Theorems 3.12, 3.13, and 3.24.

Theorem 4.16. Let $t, r \in (0, \infty)$ and Φ be an Orlicz function with positive lower type p_Φ^- and positive upper type p_Φ^+ satisfying $0 < p_\Phi^- \leq p_\Phi^+ < \infty$. Then Theorems 3.12 and 3.13 with $X := (E_\Phi^r)_t(\mathbb{R}^n)$ and $p_- := \min\{r, p_\Phi^-\}$ hold true; moreover, Theorem 3.24 with $X := (E_\Phi^r)_t(\mathbb{R}^n)$, $p_- := \min\{r, p_\Phi^-\}$, $q \in (\max\{1, r, p_\Phi^+\}, \infty)$, $d \in (0, \min\{1, r, p_\Phi^-\})$, and

$$s \in \left(\frac{n}{d} - n - \delta, \infty \right) \cap \left[\left[n \left(\frac{1}{\min\{1, p_\Phi^-\}} - 1 \right) \right], \infty \right) \cap \mathbb{Z}_+$$

holds true.

Proof. Similarly to the proof of [71, Lemma 4.5], we find that $L^\Phi(\mathbb{R}^n)$ is a ball quasi-Banach space with an absolutely continuous quasi-norm. Moreover, by a claim in [69, p. 39], we conclude that both Assumptions 3.8 and 3.19 with $X := (E_\Phi^r)_t(\mathbb{R}^n)$, $p_- := \min\{r, p_\Phi^-\}$, $r_0 \in (d, \min\{r, p_\Phi^-\})$, and $p_0 \in (\max\{r, p_\Phi^+\}, q)$ hold true. Thus, all the assumptions of Theorems 3.12, 3.13, and 3.24 with $X := (E_\Phi^r)_t(\mathbb{R}^n)$ are satisfied. Then, using Theorems 3.12, 3.13, and 3.24 with $X := (E_\Phi^r)_t(\mathbb{R}^n)$, we obtain the desired conclusions, which completes the proof of Theorem 4.16. \square

Remark 4.17. To the best of our knowledge, Theorem 4.16 is new.

4.5 Morrey spaces

Recall that Morrey [44] introduced the Morrey space $M_r^p(\mathbb{R}^n)$ with $0 < r \leq p < \infty$ to study the regularity of solutions to certain equations. The Morrey space has many applications in the theory of elliptic partial differential equations, potential theory, and harmonic analysis (see, for instance, [11, 49, 59]).

Definition 4.18. Let $0 < r \leq p < \infty$. The Morrey space $M_r^p(\mathbb{R}^n)$ is defined to be the set of all the measurable functions f on \mathbb{R}^n such that

$$\|f\|_{M_r^p(\mathbb{R}^n)} := \sup_{B \in \mathcal{B}(\mathbb{R}^n)} |B|^{\frac{1}{p} - \frac{1}{r}} \|f\|_{L^r(B)} < \infty.$$

Remark 4.19. Let $0 < r \leq p < \infty$. By Definition 4.18, we easily find that $M_r^p(\mathbb{R}^n)$ is a ball quasi-Banach function space. However, it has been pointed out in [50, p. 87] that $M_r^p(\mathbb{R}^n)$ may not be a quasi-Banach function space. Moreover, $M_r^p(\mathbb{R}^n)$ does not have an absolutely continuous quasi-norm unless $r = p$, namely, $M_r^p(\mathbb{R}^n) = L^r(\mathbb{R}^n)$.

The following theorem is a corollary of both Theorems 3.12 and 3.13.

Theorem 4.20. Let $0 < r \leq p < \frac{n}{\alpha}$. Then Theorems 3.12 and 3.13 with both $X := M_r^p(\mathbb{R}^n)$ and $p_- := r$ hold true.

Proof. By Definition 4.18, we easily find that $M_r^p(\mathbb{R}^n)$ is a ball quasi-Banach function space. In addition, by [58, Lemma 2.5] (see also [72, Lemma 7.2]), we conclude that $M_r^p(\mathbb{R}^n)$ satisfies Assumption 3.8 with both $X := M_r^p(\mathbb{R}^n)$ and $p_- := r$. Thus, all the assumptions of Theorems 3.12 and 3.13 with $X := M_r^p(\mathbb{R}^n)$ are satisfied. Then, using Theorems 3.12 and 3.13 with $X := M_r^p(\mathbb{R}^n)$, we obtain the desired conclusions, which completes the proof of Theorem 4.20. \square

Remark 4.21. (i) Let $0 < r \leq p < \infty$. As Remark 4.19 point out that the Morrey space $M_r^p(\mathbb{R}^n)$ does not have an absolutely continuous quasi-norm unless $r = p$, Theorem 3.24 can not be applied to $M_r^p(\mathbb{R}^n)$.

(ii) To the best of our knowledge, Theorem 4.20 is new.

(iii) It is worth pointing out that there exist many studies about important operators on Morrey-type space; we refer the reader to [18, 24, 25, 51, 52, 68].

4.6 Mixed-norm Lebesgue spaces

The mixed-norm Lebesgue space $L^{\vec{p}}(\mathbb{R}^n)$ was studied by Benedek and Panzone [5] in 1961, which can be traced back to Hörmander [29]. We refer the reader to [12, 13, 14, 22, 30, 31, 33] for more studies on mixed-norm type spaces.

Definition 4.22. Let $\vec{p} := (p_1, \dots, p_n) \in (0, \infty]^n$. The mixed-norm Lebesgue space $L^{\vec{p}}(\mathbb{R}^n)$ is defined to be the set of all the measurable functions f on \mathbb{R}^n such that

$$\|f\|_{L^{\vec{p}}(\mathbb{R}^n)} := \left\{ \int_{\mathbb{R}} \cdots \left[\int_{\mathbb{R}} |f(x_1, \dots, x_n)|^{p_1} dx_1 \right]^{\frac{p_2}{p_1}} \cdots dx_n \right\}^{\frac{1}{p_n}} < \infty$$

with the usual modifications made when $p_i = \infty$ for some $i \in \{1, \dots, n\}$. Moreover, let

$$p_- := \min\{p_1, \dots, p_n\} \text{ and } p_+ := \max\{p_1, \dots, p_n\}. \quad (4.1)$$

Remark 4.23. Let $\vec{p} \in (0, \infty)^n$. From Definition 4.22, we easily deduce that $L^{\vec{p}}(\mathbb{R}^n)$ is a ball quasi-Banach space. However, as was pointed out in [72, Remark 7.21], $L^{\vec{p}}(\mathbb{R}^n)$ may not be a quasi-Banach function space.

The following theorem is a corollary of Theorems 3.12, 3.13, and 3.24.

Theorem 4.24. *Let $\vec{p} := (p_1, \dots, p_n) \in (0, \infty)^n$. Then Theorems 3.12, 3.13, and 3.24 with both $X := L^{\vec{p}}(\mathbb{R}^n)$ and $p_- := \min\{p_1, \dots, p_n\}$ hold true.*

Proof. Let all the symbols be the same as in the present theorem, and let p_+ be the same as in (4.1). By Definition 4.22, we easily find that $L^{\vec{p}}(\mathbb{R}^n)$ is a ball quasi-Banach space with an absolutely continuous quasi-norm. In addition, from [30, Lemma 3.7], we deduce that Assumption 3.8 with both $X := L^{\vec{p}}(\mathbb{R}^n)$ and p_- in (4.1) holds true. In addition, using both the dual theorem of $L^q(\mathbb{R}^n)$ (see [5, p. 304, Theorem 1.a]) and [30, Lemma 3.5] (see also [33, Lemma 4.3]), we conclude that Assumption 3.19 with any $r_0 \in (0, p_-)$ and $p_0 \in (p_+, \infty)$ also holds true. Thus, all the assumptions of Theorems 3.12, 3.13, and 3.24 with $X := L^{\vec{p}}(\mathbb{R}^n)$ are satisfied. Then, using Theorems 3.12, 3.13, and 3.24 with $X := L^{\vec{p}}(\mathbb{R}^n)$, we obtain the desired conclusions, which completes the proof of Theorem 4.24. \square

Remark 4.25. To the best of our knowledge, Theorem 4.24 is new.

4.7 Local generalized Herz spaces

The local generalized Herz space was originally introduced by Rafeiro and Samko [48], which is the generalization of the classical homogeneous Herz space and connects with the generalized Morrey type space. Very recently, Li et al. [41] developed a complete real-variable theory of Hardy spaces associated with the local generalized Herz space. We now present the concepts of both the function class $M(\mathbb{R}_+)$ and the local generalized Herz space $\mathcal{K}_{\omega, \mathbf{0}}^{p, q}(\mathbb{R}^n)$ (see [48, Definitions 2.1 and 2.2] and also [41, Definitions 1.1.1 and 1.2.1]).

Definition 4.26. Let $\mathbb{R}_+ := (0, \infty)$. The function class $M(\mathbb{R}_+)$ is defined to be the set of all the positive functions ω on \mathbb{R}_+ such that, for any $0 < \delta < N < \infty$,

$$0 < \inf_{t \in (\delta, N)} \omega(t) \leq \sup_{t \in (\delta, N)} \omega(t) < \infty$$

and there exist four constants $\alpha_0, \beta_0, \alpha_\infty, \beta_\infty \in \mathbb{R}$ such that

- (i) for any $t \in (0, 1]$, $\omega(t)t^{-\alpha_0}$ is almost increasing and $\omega(t)t^{-\beta_0}$ is almost decreasing;
- (ii) for any $t \in [1, \infty)$, $\omega(t)t^{-\alpha_\infty}$ is almost increasing and $\omega(t)t^{-\beta_\infty}$ is almost decreasing.

Definition 4.27. Let $p, q \in (0, \infty)$ and $\omega \in M(\mathbb{R}_+)$. The local generalized Herz space $\dot{\mathcal{K}}_{\omega, \mathbf{0}}^{p, q}(\mathbb{R}^n)$ is defined to be the set of all the measurable functions f on \mathbb{R}^n such that

$$\|f\|_{\dot{\mathcal{K}}_{\omega, \mathbf{0}}^{p, q}(\mathbb{R}^n)} := \left\{ \sum_{k \in \mathbb{Z}} [\omega(2^k)]^q \|f \mathbf{1}_{B(\mathbf{0}, 2^k) \setminus B(\mathbf{0}, 2^{k-1})}\|_{L^p(\mathbb{R}^n)}^q \right\}^{\frac{1}{q}}$$

is finite.

Remark 4.28. Let all the symbols be the same as in Definition 4.27. It has been proved in [41, Theorem 1.2.42] that $\dot{\mathcal{K}}_{\omega, \mathbf{0}}^{p, r}(\mathbb{R}^n)$ is a ball quasi-Banach space. However, as was pointed out in [9, Remark 4.15], $\dot{\mathcal{K}}_{\omega, \mathbf{0}}^{p, q}(\mathbb{R}^n)$ may not be a quasi-Banach function space.

Definition 4.29. Let ω be a positive function on \mathbb{R}_+ . Then the *Matuszewska-Orlicz indices* $m_0(\omega)$, $M_0(\omega)$, $m_\infty(\omega)$, and $M_\infty(\omega)$ of ω are defined, respectively, by setting, for any $h \in (0, \infty)$,

$$m_0(\omega) := \sup_{t \in (0, 1)} \frac{\ln(\overline{\lim}_{h \rightarrow 0^+} \frac{\omega(ht)}{\omega(h)})}{\ln t}, \quad M_0(\omega) := \inf_{t \in (0, 1)} \frac{\ln(\underline{\lim}_{h \rightarrow 0^+} \frac{\omega(ht)}{\omega(h)})}{\ln t},$$

$$m_\infty(\omega) := \sup_{t \in (1, \infty)} \frac{\ln(\underline{\lim}_{h \rightarrow \infty} \frac{\omega(ht)}{\omega(h)})}{\ln t},$$

and

$$M_\infty(\omega) := \inf_{t \in (1, \infty)} \frac{\ln(\overline{\lim}_{h \rightarrow \infty} \frac{\omega(ht)}{\omega(h)})}{\ln t}.$$

The following theorem is a corollary of Theorems 3.12, 3.13, and 3.24.

Theorem 4.30. Let $p, r \in (0, \infty)$ and $\omega \in M(\mathbb{R}_+)$. Then Theorems 3.12, 3.13, and 3.24 with both $X := \dot{\mathcal{K}}_{\omega, \mathbf{0}}^{p, r}(\mathbb{R}^n)$ and

$$p_- := \min \left\{ p, \frac{n}{\max\{M_0(\omega), M_\infty(\omega)\} + n/p} \right\}$$

hold true.

Proof. Let

$$r_0 \in (0, \min\{1, p_-, r\})$$

and

$$p_0 \in \left(\max \left\{ p, \frac{n}{\min\{m_0(\omega), m_\infty(\omega)\} + n/p} \right\}, \infty \right].$$

From [41, Theorems 1.2.42 and 1.4.1], we deduce that $\dot{\mathcal{K}}_{\omega, \mathbf{0}}^{p, r}(\mathbb{R}^n)$ is a ball quasi-Banach space with an absolutely continuous quasi-norm. To prove the required conclusions, it suffices to show that $\dot{\mathcal{K}}_{\omega, \mathbf{0}}^{p, r}(\mathbb{R}^n)$ satisfies all the assumptions of Theorems 3.12, 3.13, and 3.24 with $X := \dot{\mathcal{K}}_{\omega, \mathbf{0}}^{p, r}(\mathbb{R}^n)$. Indeed, by [41, Lemma 4.3.10], we conclude that Assumption 3.8 with $X := \dot{\mathcal{K}}_{\omega, \mathbf{0}}^{p, r}(\mathbb{R}^n)$ holds true. Moreover, from [41, Lemma 1.8.6], we deduce that Assumption 3.19 with $X := \dot{\mathcal{K}}_{\omega, \mathbf{0}}^{p, r}(\mathbb{R}^n)$ also holds true. Thus, all the assumptions of Theorems 3.12, 3.13, and 3.24 with $X := \dot{\mathcal{K}}_{\omega, \mathbf{0}}^{p, r}(\mathbb{R}^n)$ are satisfied. Then, using Theorems 3.12, 3.13, and 3.24 with $X := \dot{\mathcal{K}}_{\omega, \mathbf{0}}^{p, r}(\mathbb{R}^n)$, we obtain the desired conclusions, which completes the proof of Theorem 4.30. \square

Remark 4.31. To the best of our knowledge, Theorem 4.30 is new.

4.8 Mixed Herz spaces

We now present the following definition of mixed Herz spaces, which is just [73, Definition 2.3].

Definition 4.32. Let $\vec{p} := (p_1, \dots, p_n), \vec{q} := (q_1, \dots, q_n) \in (0, \infty]^n$, $\vec{\alpha} := (\alpha_1, \dots, \alpha_n) \in \mathbb{R}^n$, and $R_{k_i} := (-2^{k_i}, 2^{k_i}) \setminus (-2^{k_i-1}, 2^{k_i-1})$ for any $k_i \in \mathbb{Z}$ and $i \in \{1, \dots, n\}$. The mixed Herz space $\dot{E}_{\vec{q}}^{\vec{\alpha}, \vec{p}}(\mathbb{R}^n)$ is defined to be the set of all the functions $f \in \mathcal{M}(\mathbb{R}^n)$ such that

$$\begin{aligned} \|f\|_{\dot{E}_{\vec{q}}^{\vec{\alpha}, \vec{p}}(\mathbb{R}^n)} &:= \left\{ \sum_{k_n \in \mathbb{Z}} 2^{k_n p_n \alpha_n} \left[\int_{R_{k_n}} \dots \left\{ \sum_{k_1 \in \mathbb{Z}} 2^{k_1 p_1 \alpha_1} \right. \right. \right. \\ &\quad \times \left. \left. \left. \int_{R_{k_1}} |f(x_1, \dots, x_n)|^{q_1} dx_1 \right]^{\frac{p_1}{q_1}} \right]^{\frac{q_2}{p_1}} \dots dx_n \right]^{\frac{p_n}{q_n}} \right]^{\frac{1}{p_n}} \\ &=: \left\| \dots \|f\|_{\dot{K}_{q_1}^{\alpha_1, p_1}(\mathbb{R})} \dots \right\|_{\dot{K}_{q_n}^{\alpha_n, p_n}(\mathbb{R})} < \infty \end{aligned}$$

with the usual modifications made when $p_i = \infty$ or $q_j = \infty$ for some $i, j \in \{1, \dots, n\}$, where $\left\| \dots \|f\|_{\dot{K}_{q_1}^{\alpha_1, p_1}(\mathbb{R})} \dots \right\|_{\dot{K}_{q_n}^{\alpha_n, p_n}(\mathbb{R})}$ denotes the norm obtained after taking successively the $\dot{K}_{q_1}^{\alpha_1, p_1}(\mathbb{R})$ -norm to x_1, \dots , the $\dot{K}_{q_{n-1}}^{\alpha_{n-1}, p_{n-1}}(\mathbb{R})$ -norm to x_{n-1} , and the $\dot{K}_{q_n}^{\alpha_n, p_n}(\mathbb{R})$ -norm to x_n .

Recall that, to study the Lebesgue points of functions in mixed-norm Lebesgue spaces, Huang et al. [32] introduced a special case of the mixed Herz space and, later, Zhao et al. [73] generalized it to the above case. Also, both the dual theorem and the Riesz–Thorin interpolation theorem on the above mixed Herz space have been fully studied in [73].

Remark 4.33. Let $\vec{p} := (p_1, \dots, p_n), \vec{q} := (q_1, \dots, q_n) \in (0, \infty]^n$ and $\vec{\alpha} := (\alpha_1, \dots, \alpha_n) \in \mathbb{R}^n$. By [73, Proposition 2.22], we conclude that $\dot{E}_{\vec{q}}^{\vec{\alpha}, \vec{p}}(\mathbb{R}^n)$ is a ball quasi-Banach space if and only if, for any $i \in \{1, \dots, n\}$, $\alpha_i \in (-\frac{1}{q_i}, \infty)$. However, as was pointed out in [9, Remark 4.22], $\dot{E}_{\vec{q}}^{\vec{\alpha}, \vec{p}}(\mathbb{R}^n)$ may not be a quasi-Banach function space.

The following theorem is a corollary of Theorems 3.12, 3.13, and 3.24.

Theorem 4.34. Let $\vec{p} := (p_1, \dots, p_n), \vec{q} := (q_1, \dots, q_n) \in (0, \infty]^n$ and $\vec{\alpha} := (\alpha_1, \dots, \alpha_n) \in \mathbb{R}^n$. Then Theorems 3.12, 3.13, and 3.24 with both $X := \dot{E}_{\vec{q}}^{\vec{\alpha}, \vec{p}}(\mathbb{R}^n)$ and

$$p_- := \min \left\{ p_1, \dots, p_n, q_1, \dots, q_n, \left(\alpha_1 + \frac{1}{q_1} \right)^{-1}, \dots, \left(\alpha_n + \frac{1}{q_n} \right)^{-1} \right\} \quad (4.2)$$

hold true.

Proof. Let

$$r_0 \in (0, \min\{1, p_-\})$$

and

$$p_0 \in \left(\max \left\{ p_1, \dots, p_n, q_1, \dots, q_n, \left(\alpha_1 + \frac{1}{q_1} \right)^{-1}, \dots, \left(\alpha_n + \frac{1}{q_n} \right)^{-1} \right\}, \infty \right).$$

Using [73, Propositions 2.8 and 2.22], we conclude that $\dot{E}_{\vec{q}}^{\vec{\alpha}, \vec{p}}(\mathbb{R}^n)$ is a ball quasi-Banach space with an absolutely continuous quasi-norm. By both [73, Lemma 5.3(i)] and its proof, we conclude that Assumption 3.8 with both $X := \dot{E}_{\vec{q}}^{\vec{\alpha}, \vec{p}}(\mathbb{R}^n)$ and p_- in (4.2) holds true. In addition, using both [73, Lemma 5.3(ii)] and its proof, we find that Assumption 3.19 with $X := \dot{E}_{\vec{q}}^{\vec{\alpha}, \vec{p}}(\mathbb{R}^n)$ also holds true. Thus, all the assumptions of Theorems 3.12, 3.13, and 3.24 with $X := \dot{E}_{\vec{q}}^{\vec{\alpha}, \vec{p}}(\mathbb{R}^n)$ are satisfied. Then, using Theorems 3.12, 3.13, and 3.24 with $X := \dot{E}_{\vec{q}}^{\vec{\alpha}, \vec{p}}(\mathbb{R}^n)$, we obtain the desired conclusions, which completes the proof of Theorem 4.34. \square

Remark 4.35. To the best of our knowledge, Theorem 4.34 is new.

Acknowledgements The authors would like to thank both referees for their very carefully reading and several valuable remarks which definitely improve the quality of this article.

Data Availability Statement Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

Compliance with ethical standards

Conflict of interest The author declares that they have no conflict interests.

References

- [1] R. Arai and E. Nakai, Compact commutators of Calderón–Zygmund and generalized fractional integral operators with a function in generalized Campanato spaces on generalized Morrey spaces, *Tokyo J. Math.* 42 (2019), 471–496.
- [2] R. Arai, E. Nakai and Y. Sawano, Generalized fractional integral operators on Orlicz–Hardy spaces, *Math. Nachr.* 294 (2021), 224–235.
- [3] P. Auscher and M. Mourgoglou, Representation and uniqueness for boundary value elliptic problems via first order systems, *Rev. Mat. Iberoam.* 35 (2019), 241–315.
- [4] P. Auscher and C. Prisuelos-Arribas, Tent space boundedness via extrapolation, *Math. Z.* 286 (2017), 1575–1604.
- [5] A. Benedek and R. Panzone, The space L^p , with mixed norm, *Duke Math. J.* 28 (1961), 301–324.
- [6] C. Bennett and R. Sharpley, *Interpolation of Operators*, Pure and Applied Mathematics 129, Academic Press, Boston, MA, 1988.
- [7] S. Campanato, Proprietà di una famiglia di spazi funzionali, *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (3)* 18 (1964), 137–160.

- [8] D.-C. Chang, S. Wang, D. Yang and Y. Zhang, Littlewood–Paley characterizations of Hardy-type spaces associated with ball quasi-Banach function spaces, *Complex Anal. Oper. Theory* 14 (2020), Paper No. 40, 33 pp.
- [9] Y. Chen, H. Jia and D. Yang, Boundedness of fractional integrals on Hardy spaces associated with ball quasi-Banach function spaces, Submitted.
- [10] Y. Chen, H. Jia and D. Yang, Boundedness of fractional integrals on ball Campanato-type function spaces, Submitted.
- [11] F. Chiarenza and M. Frasca, Morrey spaces and Hardy–Littlewood maximal function, *Rend. Mat. Appl.* (7) 7 (1987), 273-279.
- [12] G. Cleanthous, A. G. Georgiadis and M. Nielsen, Anisotropic mixed-norm Hardy spaces, *J. Geom. Anal.* 27 (2017), 2758-2787.
- [13] G. Cleanthous, A. G. Georgiadis and M. Nielsen, Discrete decomposition of homogeneous mixed-norm Besov spaces, in: *Functional Analysis, Harmonic Analysis, and Image Processing: A Collection of Papers in Honor of Björn Jawerth*, 167-184, *Contemp. Math.* 693, Amer. Math. Soc., Providence, RI, 2017.
- [14] G. Cleanthous, A. G. Georgiadis and M. Nielsen, Molecular decomposition of anisotropic homogeneous mixed-norm spaces with applications to the boundedness of operators, *Appl. Comput. Harmon. Anal.* 47 (2019), 447-480.
- [15] D. V. Cruz-Uribe and A. Fiorenza, *Variable Lebesgue spaces. Foundations and harmonic analysis*, *Applied and Numerical Harmonic Analysis*, Birkhäuser/Springer, Heidelberg, 2013.
- [16] F. Dai, L. Grafakos, Z. Pan, D. Yang, W. Yuan and Y. Zhang, The Bourgain–Brezis–Mironescu formula on ball Banach function spaces, *Math. Ann.* (to appear).
- [17] R. del Campo, A. Fernández, F. Mayoral and F. Naranjo, Orlicz spaces associated to a quasi-Banach function space: applications to vector measures and interpolation, *Collect. Math.* 72 (2021), 481-499.
- [18] F. Deringoz, V. S. Guliyev, E. Nakai, Y. Sawano and M. Shi, Generalized fractional maximal and integral operators on Orlicz and generalized Orlicz–Morrey spaces of the third kind, *Positivity* 23 (2019), 727-757.
- [19] L. Diening, P. Hästö and S. Roudenko, Function spaces of variable smoothness and integrability, *J. Funct. Anal.* 256 (2009), 1731-1768.
- [20] J. Duoandikoetxea, *Fourier Analysis*, *Graduate Studies in Mathematics* 29, American Mathematical Society, Providence, RI, 2001.
- [21] C. Fefferman and E. M. Stein, H^p spaces of several variables, *Acta Math.* 129 (1972), 137-193.
- [22] A. G. Georgiadis and M. Nielsen, Pseudodifferential operators on mixed-norm Besov and Triebel–Lizorkin spaces, *Math. Nachr.* 289 (2016), 2019-2036.
- [23] L. Grafakos, *Classical Fourier Analysis*, Third edition, *Graduate Texts in Mathematics* 249, Springer, New York, 2014.
- [24] D. I. Hakim, Y. Sawano and T. Shimomura, Boundedness of generalized fractional integral operators from the Morrey space $L_{1,\phi}(X;\mu)$ to the Campanato space $\mathcal{L}_{1,\psi}(X;\mu)$ over non-doubling measure spaces, *Azerb. J. Math.* 6 (2016), 117-127.
- [25] K.-P. Ho, Fractional integral operators with homogeneous kernels on Morrey spaces with variable exponents, *J. Math. Soc. Japan* 69 (2017), 1059-1077.

- [26] K.-P. Ho, Integral operators on BMO and Campanato spaces, *Indag. Math. (N.S.)* 30 (2019), 1023-1035.
- [27] K.-P. Ho, Erdélyi-Kober fractional integral operators on ball Banach function spaces, *Rend. Semin. Mat. Univ. Padova* 145 (2021), 93-106.
- [28] K.-P. Ho, Fractional integral operators on Orlicz slice Hardy spaces, *Fract. Calc. Appl. Anal.* 25 (2022), 1294-1305.
- [29] L. Hörmander, Estimates for translation invariant operators in L^p spaces, *Acta Math.* 104 (1960), 93-140.
- [30] L. Huang, J. Liu, D. Yang and W. Yuan, Atomic and Littlewood–Paley characterizations of anisotropic mixed-norm Hardy spaces and their applications, *J. Geom. Anal.* 29 (2019), 1991-2067.
- [31] L. Huang, J. Liu, D. Yang and W. Yuan, Dual spaces of anisotropic mixed-norm Hardy spaces, *Proc. Amer. Math. Soc.* 147 (2019), 1201-1215.
- [32] L. Huang, F. Weisz, D. Yang and W. Yuan, Summability of Fourier transforms on mixed-norm Lebesgue spaces via associated Herz spaces, *Anal. Appl. (Singap.)* (2021), <http://doi.org/10.1142/S0219530521500135>.
- [33] L. Huang and D. Yang, On function spaces with mixed norms—a survey, *J. Math. Study* 54 (2021), 262-336.
- [34] T. Iwaniec and A. Verde, A study of Jacobians in Hardy–Orlicz spaces, *Proc. Roy. Soc. Edinburgh Sect. A* 129 (1999), 539-570.
- [35] M. Izuki and Y. Sawano, Characterization of BMO via ball Banach function spaces, *Vestn. St.-Peterbg. Univ. Mat. Mekh. Astron.* 4(62) (2017), 78-86.
- [36] H. Jia, J. Tao, D. Yang, W. Yuan and Y. Zhang, Special John–Nirenberg–Campanato spaces via congruent cubes, *Sci. China Math.* 65 (2022), 359-420.
- [37] H. Jia, J. Tao, D. Yang, W. Yuan and Y. Zhang, Boundedness of Calderón–Zygmund operators on special John–Nirenberg–Campanato and Hardy-type spaces via congruent cubes, *Anal. Math. Phys.* 12 (2022), Paper No. 15, 56 pp.
- [38] F. John and L. Nirenberg, On functions of bounded mean oscillation, *Comm. Pure Appl. Math.* 14 (1961), 415-426.
- [39] O. Kováčik and J. Rákosník, On spaces $L^p(x)$ and $W^{k,p(x)}$, *Czechoslovak Math. J.* 41 (116) (1991), 592-618.
- [40] S. Kislyakov and N. Kruglyak, Extremal Problems in Interpolation Theory, Whitney–Besicovitch Coverings, and Singular Integrals, *Instytut Matematyczny Polskiej Akademii Nauk. Monografie Matematyczne (New Series) 74*, Birkhäuser/Springer Basel AG, Basel, 2013.
- [41] Y. Li, D. Yang and L. Huang, Real-variable theory of Hardy spaces associated with generalized Herz spaces of Rafeiro and Samko, *Lecture Notes in Mathematics 2320*, Springer, Cham, 2022.
- [42] R. A. Macías and C. Segovia, Singular integrals on generalized Lipschitz and Hardy spaces. *Studia Math.* 65 (1979), 55-75.
- [43] Y. Meyer and R. R. Coifman, *Wavelets. Calderón–Zygmund and Multilinear Operators*, Cambridge Studies in Advanced Mathematics 48, Cambridge University Press, Cambridge, 1997.

- [44] C. B. Morrey, On the solutions of quasi-linear elliptic partial differential equations, *Trans. Amer. Math. Soc.* 43 (1938), 126-166.
- [45] E. Nakai, Singular and fractional integral operators on Campanato spaces with variable growth conditions, *Rev. Mat. Complut.* 23 (2010), 355-381.
- [46] E. Nakai, Singular and fractional integral operators on preduals of Campanato spaces with variable growth condition, *Sci. China Math.* 60 (2017), 2219-2240.
- [47] E. Nakai and G. Sadasue, Martingale Morrey–Campanato spaces and fractional integrals, *J. Funct. Spaces Appl.* 2012, Art. ID 673929, 29 pp.
- [48] H. Rafeiro and S. Samko, Herz spaces meet Morrey type spaces and complementary Morrey type spaces, *J. Fourier Anal. Appl.* 26 (2020), Paper No. 74, 14 pp.
- [49] Y. Sawano, G. Di Fazio and D. Hakim, *Morrey Spaces—Introduction and Applications to Integral Operators and PDE’s*, Vol I, *Monographs and Research Notes in Mathematics*, CRC Press, Boca Raton, FL, 2020.
- [50] Y. Sawano, K.-P. Ho, D. Yang and S. Yang, Hardy spaces for ball quasi-Banach function spaces, *Dissertationes Math.* 525 (2017), 1-102.
- [51] Y. Sawano and T. Shimomura, Boundedness of the generalized fractional integral operators on generalized Morrey spaces over metric measure spaces, *Z. Anal. Anwend.* 36 (2017), 159-190.
- [52] Y. Sawano, S. Sugano and H. Tanaka, A note on generalized fractional integral operators on generalized Morrey spaces, *Bound. Value Probl.* 2009, Art. ID 835865, 18 pp.
- [53] C. F. Serra, Molecular characterization of Hardy–Orlicz spaces, *Rev. Un. Mat. Argentina* 40 (1996), 203-217.
- [54] E. M. Stein, *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, Princeton Mathematical Series 43, *Monographs in Harmonic Analysis, III*, Princeton University Press, Princeton, NJ, 1993.
- [55] J. Sun, D. Yang and W. Yuan, Weak Hardy spaces associated with ball quasi-Banach function spaces on spaces of homogeneous type: decompositions, real interpolation, and Calderón–Zygmund operators, *J. Geom. Anal.* 32 (2022), Paper No. 191, 85 pp.
- [56] J. Sun, D. Yang and W. Yuan, Molecular characterization of weak Hardy spaces associated with ball quasi-Banach function spaces on spaces of homogeneous type with its application to Littelwood–Paley function characterization, *Forum Math.* (2022), <https://doi.org/10.1515/forum-2022-0074>.
- [57] M. H. Taibleson and G. Weiss, The molecular characterization of certain Hardy spaces, Representation theorems for Hardy spaces, in: *Astérisque* 77, 67-149, Soc. Math. France, Paris, 1980.
- [58] L. Tang and J. Xu, Some properties of Morrey type Besov–Triebel spaces, *Math. Nachr.* 278 (2005), 904-917.
- [59] J. Tao, Da. Yang and Do. Yang, Boundedness and compactness characterizations of Cauchy integral commutators on Morrey spaces, *Math. Methods Appl. Sci.* 42 (2019), 1631-1651.
- [60] J. Tao, D. Yang, W. Yuan and Y. Zhang, Compactness characterizations of commutators on ball Banach function spaces, *Potential Analysis* (2021), <https://doi.org/10.1007/s11118-021-09953-w>.
- [61] F. Wang, D. Yang and S. Yang, Applications of Hardy spaces associated with ball quasi-Banach function spaces, *Results Math.* 75 (2020), Paper No. 26, 58 pp.

- [62] S. Wang, D. Yang, W. Yuan and Y. Zhang, Weak Hardy-type spaces associated with ball quasi-Banach function spaces II: Littlewood–Paley characterizations and real interpolation, *J. Geom. Anal.* 31 (2021), 631-696.
- [63] S. Yamaguchi and E. Nakai, Generalized fractional integral operators on Campanato spaces and their bi-preduals, *Math. J. Ibaraki Univ.* 53 (2021), 17-34.
- [64] S. Yamaguchi and E. Nakai, Compactness of commutators of integral operators with functions in Campanato spaces on Orlicz–Morrey spaces, *J. Fourier Anal. Appl.* 28 (2022), Paper No. 33, 32 pp.
- [65] X. Yan, Z. He, D. Yang and W. Yuan, Hardy spaces associated with ball quasi-Banach function spaces on spaces of homogeneous type: Characterizations of maximal functions, decompositions, and dual spaces, *Math. Nachr.* (2022), <http://doi.org/10.1002/mana.202100432>.
- [66] X. Yan, Z. He, D. Yang and W. Yuan, Hardy spaces associated with ball quasi-Banach function spaces on spaces of homogeneous type: Littlewood–Paley characterizations with applications to boundedness of Calderón–Zygmund Operators, *Acta Math. Sin. (Engl. Ser.)* 38 (2022), 1133-1184.
- [67] X. Yan, D. Yang and W. Yuan, Intrinsic square function characterizations of Hardy spaces associated with ball quasi-Banach function spaces, *Front. Math. China* 15 (2020), 769-806.
- [68] T.-L. Yee and K.-P. Ho, Fractional integral operators with homogeneous kernels on generalized Lorentz–Morrey spaces, *J. Math. Inequal.* 15 (2021), 17-30.
- [69] Y. Zhang, L. Huang, D. Yang and W. Yuan, New ball Campanato-type function spaces and their applications, *J. Geom. Anal.* 32 (2022), Paper No. 99, 42 pp.
- [70] Y. Zhang, D. Yang, W. Yuan, Real-variable characterizations of local Orlicz-slice Hardy spaces with application to bilinear decompositions, *Commun. Contemp. Math.* 24 (2022), 2150004, 35 pp.
- [71] Y. Zhang, D. Yang, W. Yuan and S. Wang, Real-variable characterizations of Orlicz-slice Hardy spaces, *Anal. Appl. (Singap.)* 17 (2019), 597-664.
- [72] Y. Zhang, D. Yang, W. Yuan and S. Wang, Weak Hardy-type spaces associated with ball quasi-Banach function spaces I: Decompositions with applications to boundedness of Calderón–Zygmund operators, *Sci. China Math.* 64 (2021), 2007-2064.
- [73] Y. Zhao, D. Yang and Y. Zhang, Mixed-norm Herz spaces and their applications in related Hardy spaces, Submitted.

Yiqun Chen, Hongchao Jia and Dachun Yang (Corresponding author)

Laboratory of Mathematics and Complex Systems (Ministry of Education of China), School of Mathematical Sciences, Beijing Normal University, Beijing 100875, The People's Republic of China

E-mails: yiqunchen@mail.bnu.edu.cn (Y. Chen)

hcjia@mail.bnu.edu.cn (H. Jia)

dcyang@bnu.edu.cn (D. Yang)