

Wasserstein Complexity of Quantum Circuits

Lu Li,^{1,*} Kaifeng Bu,^{2,†} Dax Enshan Koh,^{3,‡} Arthur Jaffe,^{2,§} and Seth Lloyd^{4,¶}

¹*Department of Mathematics, Zhejiang Sci-Tech University, Hangzhou, Zhejiang 310018, China*

²*Department of Physics, Harvard University, Cambridge, Massachusetts 02138, USA*

³*Institute of High Performance Computing, Agency for Science, Technology and Research (A*STAR),*

1 Fusionopolis Way, #16-16 Connexis, Singapore 138632, Singapore

⁴*Department of Mechanical Engineering, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

Given a unitary transformation, what is the size of the smallest quantum circuit that implements it? This quantity, known as the quantum circuit complexity, is a fundamental property of quantum evolutions that has widespread applications in many fields, including quantum computation, quantum field theory, and black hole physics. In this letter, we obtain a new lower bound for the quantum circuit complexity in terms of a novel complexity measure that we propose for quantum circuits, which we call the quantum Wasserstein complexity. Our proposed measure is based on the quantum Wasserstein distance of order one (also called the quantum earth mover’s distance), a metric on the space of quantum states. We also prove several fundamental and important properties of our new complexity measure, which stand to be of independent interest. Finally, we show that our new measure also provides a lower bound for the experimental cost of implementing quantum circuits, which implies a quantum limit on converting quantum resources to computational resources. Our results provide novel applications of the quantum Wasserstein distance and pave the way for a deeper understanding of the resources needed to implement a quantum computation.

I. INTRODUCTION

How many elementary quantum gates does it take to synthesize a desired unitary transformation U ? Computing this number—called the quantum circuit complexity of U —is one of the holy grails of quantum computation, especially noisy intermediate-scale quantum computation where quantum resources are scarce [1, 2]. By using fewer elementary gates, a quantum algorithm stands to take less time to produce its output; this in turn increases its prospect for achieving the highly sought goal of quantum-computational advantage [3, 4].

In the search for optimal quantum circuits that use as few elementary gates as possible, one has taken various approaches to quantum circuit synthesis, including: reinforcement learning [5], quantum Karnaugh maps [6], and ZX-calculus [7]. While these methods usually succeed at finding smaller circuits, they seldom achieve optimal circuit solutions [8]. Computing the quantum circuit complexity of unitaries is generally hard, and no efficient algorithms are known [9, 10].

Unexpected connections have been found between the notion of quantum circuit complexity and high energy physics. A solution to the wormhole-growth paradox has been proposed, which asks: in the anti-de Sitter/conformal field theory (AdS/CFT) correspondence, which quantity in the CFT is dual to the wormhole volume [11]? Stanford and Susskind conjecture that the answer is the quantum circuit complexity of the boundary state [12]. In follow-up work by Brown and Susskind [13], they conjecture that the quantum circuit complexity grows linearly for an exponentially long time. This was subsequently formalized and proved by Haferkamp et

al. [14]. (See also [15] for two short proofs of this result.)

Nielsen pioneered a geometric approach to find upper and lower bounds for the quantum circuit complexity [16]. In a seminal series of papers, Nielsen and collaborators developed various geometric notions related to quantum computation and proved that the quantum circuit complexity of a unitary, up to polynomial factors and technical caveats, is equal to the circuit cost, defined as the length of the shortest path between two points in some curved Riemannian geometry [16–19].

A recent fruitful approach to bound Nielsen’s circuit cost is to relate it to the quantum resources in a quantum circuit that refer to ingredients in the circuit responsible for a quantum speedup. A famous example of a quantum resource is nonstabilizerness, also known as magic. The Gottesman-Knill theorem proves that stabilizer circuits are efficiently classically simulable [20], but when supplemented with magic states, such circuits become hard to simulate [21–24]. Hence, magic could be thought of as a resource for a quantum speedup [25–33]. In recent works, the circuit cost was shown to be bounded below by the circuit’s entangling power [34], magic [35], and sensitivity [35].

In this letter, we study the circuit complexity of quantum circuits using the so-called *Wasserstein distance*. Classically, the Wasserstein distance is a metric on the space of probability distributions on a metric space that traces its origins to the works of Kantorovich [36], Vaserstein [37], and others. The classical Wasserstein distance has numerous applications, including optimal transport [36, 38, 39], image retrieval in computer vision [40], and Wasserstein generative adversarial networks (WGANs) in classical machine learning [41–43].

What is an appropriate quantum generalization of the classical Wasserstein distance that is useful for quantum computation? To answer this question, several quantum generalizations of the Wasserstein distance have been proposed. These include the quantum W_2 distances, pioneered by Carlen, Maas, Datta, Rouzé, Junge, and others, based on a Riemannian met-

* lili93@zju.edu.cn

† kfbu@fas.harvard.edu

‡ dax_koh@ihpc.a-star.edu.sg

§ arthur_jaffe@harvard.edu

¶ slloyd@mit.edu

ric on the manifold of quantum states [44–50] that generalize the classical Wasserstein distance of order 2. These W_2 distances have been shown to be related to the entropy and Fisher information, two important concepts in quantum information theory [48]. In addition, several candidates have been proposed that generalize the Wasserstein distance of order 1 [51–53]. In this letter, we focus on the quantum Wasserstein distance of order 1, proposed by De Palma, Marvian, Trevisan, and Lloyd [53], which can be regarded as a quantum version of the Hamming distance on n -qudit systems. This new version of the quantum Wasserstein distance has numerous applications in quantum information and quantum computation, such as quantum machine learning (where it is commonly referred to as the *quantum earth mover's distance*) [54], quantum concentration inequalities [55], quantum differential privacy [56], characterizing limitations of variational quantum algorithms [57], etc. However, little is hitherto known about the connection between the quantum Wasserstein distance and the quantum circuit complexity.

In this work, we introduce a complexity measure for quantum circuits, called the quantum Wasserstein complexity, which we define as the maximal distance between the input and output states of the circuit, measured using the quantum Wasserstein distance of order 1. We show several useful properties of this complexity measure for quantum channels, such as subadditivity under concatenation, superadditivity under tensor product (or additivity for correlation enhanced complexity measures). Most importantly, we show a connection between the quantum Wasserstein complexity and the circuit complexity, where we show that the quantum Wasserstein complexity provides a lower bound for the circuit cost. Finally, we show that the quantum Wasserstein complexity also provides a lower bound on the experimental cost to implement quantum circuits.

II. MAIN RESULTS

Given the n -qudit system $\mathcal{H} = (\mathbb{C}^d)^{\otimes n}$, let us define $O(\mathcal{H})$ to be the set of traceless, Hermitian operators on \mathcal{H} , i.e., $O(\mathcal{H}) = \{A \in \mathcal{L}(\mathcal{H}) : \text{Tr}[A] = 0, A^\dagger = A\}$, where $\mathcal{L}(\mathcal{H})$ denotes the set of linear operators on \mathcal{H} . Also, let $D(\mathcal{H})$ denote the set of density operators on \mathcal{H} . The *quantum Wasserstein distance of order 1* [53] between two quantum states $\rho, \sigma \in D(\mathcal{H})$ is defined as:

$$\|\rho - \sigma\|_{W_1} := \frac{1}{2} \min_{\{X_i\}} \left\{ \sum_i \|X_i\|_1 : \rho - \sigma = \sum_i X_i, \right. \\ \left. X_i \in O(\mathcal{H}), \text{Tr}_i[X_i] = 0 \forall i \right\}, \quad (1)$$

where $\text{Tr}_i[\cdot]$ denotes the partial trace over the i -th subsystem, and $\|X\|_1 = \text{Tr}[\sqrt{X^\dagger X}]$ denotes the trace norm of X . This distance is also called the *quantum W_1 distance* or the *quantum earth mover's distance*. Using the quantum Wasserstein distance lets one distinguish between two quantum states that

differ locally; this is different from their global distinguishability. One example to show this distinction comes from considering the n -qubit states $|0\rangle^{\otimes n}$, $|1\rangle|0\rangle^{\otimes n-1}$ and $|1\rangle^{\otimes n}$. The trace distances [58] between $|0\rangle^{\otimes n}$ and $|1\rangle|0\rangle^{\otimes n-1}$ and between $|0\rangle^{\otimes n}$ and $|1\rangle^{\otimes n}$ are both equal to 1; hence, using the trace distance, we cannot distinguish which of $|1\rangle|0\rangle^{\otimes n-1}$ and $|1\rangle^{\otimes n}$ is further away from $|0\rangle^{\otimes n}$. However, the quantum W_1 distance between $|0\rangle^{\otimes n}$ and $|1\rangle|0\rangle^{\otimes n-1}$ is equal to 1, while the quantum W_1 distance between $|0\rangle^{\otimes n}$ and $|1\rangle^{\otimes n}$ is equal to n ; hence in quantum W_1 distance, $|1\rangle^{\otimes n}$ is further away from $|0\rangle^{\otimes n}$ than $|1\rangle|0\rangle^{\otimes n-1}$.

Another example comes from considering the trace distance between $|0\rangle^{\otimes n}$ and the Cat state $|\text{Cat}_a\rangle = a|0\rangle^{\otimes n} + \sqrt{1-|a|^2}|1\rangle^{\otimes n}$ with $0 < |a| < 1$. The trace distance is $\sqrt{1-|a|^2}$, which is very small when a is close to 1. However, the experimental resource, e.g., the number of gates, to transform $|0\rangle^{\otimes n}$ to $|\text{Cat}_a\rangle$ is proportional to the size of the system n , independent of how close a is to 1. On the other hand, the quantum W_1 distance between $|0\rangle^{\otimes n}$ and $|\text{Cat}_a\rangle$ is $\Omega((1-|a|^2)n)$. This motivates us to consider the circuit complexity in terms of the quantum W_1 distance.

Definition 1. Given an n -qudit quantum channel $\Lambda : D(\mathcal{H}) \rightarrow D(\mathcal{H})$, the *quantum Wasserstein complexity* $C_{W_1}(\Lambda)$ is the maximal distance between the input state and output state in quantum W_1 distance,

$$C_{W_1}(\Lambda) := \max_{\rho \in D(\mathcal{H})} \|\rho - \Lambda(\rho)\|_{W_1}. \quad (2)$$

By the convexity of the quantum W_1 distance, we need only to take the maximization over all pure states,

$$C_{W_1}(\Lambda) = \max_{|\psi\rangle \in \mathcal{H}} \|\|\psi\rangle\langle\psi| - \Lambda(|\psi\rangle\langle\psi|)\|_{W_1}. \quad (3)$$

To demonstrate applications of the quantum Wasserstein complexity, let us first study its basic properties.

Proposition 2. The quantum Wasserstein complexity $C_{W_1}(\Lambda)$ satisfies the following properties:

- (1) *Faithfulness:* $C_{W_1}(\Lambda) = 0$ if and only if Λ is the identity map;
- (2) *Convexity:* $C_{W_1}(\sum_i p_i \Lambda_i) \leq \sum_i p_i C_{W_1}(\Lambda_i)$, where $p_i \geq 0$ and $\sum_i p_i = 1$;
- (3) *Subadditivity under concatenation:* $C_{W_1}(\Lambda_1 \circ \Lambda_2) \leq C_{W_1}(\Lambda_1) + C_{W_1}(\Lambda_2)$;
- (4) $C_{W_1}(\Lambda_1 \otimes \Lambda_2) \leq C_{W_1}(\Lambda_1 \otimes I) + C_{W_1}(\Lambda_2 \otimes I)$;
- (5) *Superadditivity under tensorization:* $C_{W_1}(\Lambda_1 \otimes \Lambda_2) \geq C_{W_1}(\Lambda_1) + C_{W_1}(\Lambda_2)$;
- (6) For a unitary channel U , $C_{W_1}(U) = C_{W_1}(U^\dagger)$;
- (7) If the quantum channel Λ acts nontrivially on a k -qudit subsystem, then we have $C_{W_1}(\Lambda) \leq k$.

We prove Proposition 2 in Appendix B. Note that there exists a quantum channel Λ , such that the inequality (5) holds strictly. For example, for the single-qudit depolarizing channel $D_p(\cdot) = p(\cdot) + (1-p)\text{Tr}[\cdot]I/d$, it holds that $C_{W_1}(D_p) = (1-p)(1-1/d)$ and $C_{W_1}(D_p \otimes I) = (1-p)(1-1/d^2)$. Hence, the inequality in (5) holds strictly. See more examples in Table I.

The effects of the environment on quantum systems are usually taken into consideration, which means the ancilla qudits should be considered.

Definition 3. Given an n -qudit quantum channel $\Lambda : D(\mathcal{H}) \rightarrow D(\mathcal{H})$, the correlation-assisted quantum Wasserstein complexity AC_{W_1} for an n -qudit quantum channel Λ is defined as follows

$$AC_{W_1}(\Lambda) := \sup_{m \in \mathbb{Z}_{\geq 0}} C_{W_1}(\Lambda \otimes I_m), \quad (4)$$

where $\mathbb{Z}_{\geq 0}$ denotes the set of nonnegative integers, and I_m denotes the m -qubit identity operator.

Note that $C_{W_1}(\Lambda \otimes I_m) \geq C_{W_1}(\Lambda \otimes I_{m-1}), \forall m \in \mathbb{Z}_{\geq 0}$ and $C_{W_1}(\Lambda) \leq n$. Hence, AC_{W_1} is well-defined.

Proposition 4. The correlation-assisted Wasserstein complexity $AC_{W_1}(\Lambda)$ satisfies the following properties:

- (1) *Faithfulness:* $AC_{W_1}(\Lambda) = 0$ if and only if Λ is the identity map;
- (2) *Convexity:* $AC_{W_1}(\sum_i p_i \Lambda_i) \leq \sum_i p_i AC_{W_1}(\Lambda_i)$, where $p_i \geq 0$ and $\sum_i p_i = 1$;
- (3) *Subadditivity under concatenation:* $AC_{W_1}(\Lambda_1 \circ \Lambda_2) \leq AC_{W_1}(\Lambda_1) + AC_{W_1}(\Lambda_2)$;
- (4) *Additivity under tensorization:* $AC_{W_1}(\Lambda_1 \otimes \Lambda_2) = AC_{W_1}(\Lambda_1) + AC_{W_1}(\Lambda_2)$.

The proof of Proposition 4 is provided in Appendix B. Note that, unlike C_{W_1} , AC_{W_1} is additive under tensorization. Based on the definition, it is easy to see that $AC_{W_1}(\Lambda) \geq C_{W_1}(\Lambda)$. Also, there exists some quantum channel Λ , e.g., the 1-qudit depolarizing channel, such that $AC_{W_1}(\Lambda) > C_{W_1}(\Lambda)$. (See the examples in Table I).

After studying the basic properties of C_{W_1} and AC_{W_1} , let us consider an application of the quantum Wasserstein complexity to the study of the circuit complexity of quantum circuits. The circuit complexity of a unitary operator U is defined as the minimum number of basic gates needed to generate U [59–61] (See Fig. 1). Here, we consider the circuit cost of quantum circuits, which was introduced by Nielsen et al. [16–18] as a geodesic distance from the identity operator to the target unitary with respect to a given metric, and was shown to provide a useful lower bound for the quantum circuit complexity.

Here, the circuit cost of a unitary $U \in \text{SU}(d^n)$ with respect to traceless Hermitian operators h_1, \dots, h_m , supported on 2 qudits and normalized as $\|h_i\|_\infty = 1$, is defined to be

$$\text{Cost}(U) := \inf \int_0^1 \sum_{j=1}^m |r_j(s)| ds, \quad (5)$$




FIG. 1. Diagram showing a quantum circuit composed of single-qubit and two-qubit basic gates. Quantum circuit complexity quantifies the minimum number of basic gates needed to generate a target unitary.




FIG. 2. The circuit cost is the geodesic distance from the identity operator I to the target unitary U with respect to some metric, as illustrated by the length of the curve connecting I and U in the diagram.

where the infimum in (5) is taken over all continuous functions $r_j : [0, 1] \rightarrow \mathbb{R}$ that satisfy $H(s) = \sum_{j=1}^m r_j(s) h_j$ and $U = \mathcal{P} \exp \left(-i \int_0^1 H(s) ds \right)$, where \mathcal{P} denotes the path-ordering operator. (See Fig. 2)

For an n -qudit Hamiltonian acting nontrivially on a k -qudit subsystem, there is a simple upper bound on the total change of the quantum Wasserstein complexity through unitary evolution.

Lemma 5. Given an n -qudit system with a k -qudit Hamiltonian H , the change of the quantum Wasserstein complexity for the unitary evolution $U_t = \exp(-iHt)$ with any time interval Δt is bounded as follows

$$C_{W_1}(U_{t+\Delta t}) - C_{W_1}(U_t) \leq k. \quad (6)$$

Proof. This comes from the fact that $C_{W_1}(U_{t+\Delta t}) = C_{W_1}(U_{\Delta t} \circ U_t) \leq C_{W_1}(U_t) + C_{W_1}(U_{\Delta t})$, and $C_{W_1}(U_{\Delta t}) \leq k$. \square

To quantify the change of the quantum Wasserstein complexity in an infinitesimally small time interval, we need to

quantum channel Λ	$C_{W_1}(\Lambda)$	$AC_{W_1}(\Lambda)$
Single-qudit depolarizing channel D_p	$(1-p)(1-1/d)$	$(1-p)(1-1/d^2)$
Tensor product of depolarizing channels $D_p^{\otimes n}$	$\geq n(1-p)(1-1/d)$ $\leq n(1-p)(1-1/d^2)$	$n(1-p)(1-1/d^2)$
Tensor product of Hadamard gates $H^{\otimes n}$	n	n
CNOT gates $\prod_i \text{CNOT}_{i,i+1}$	$\Theta(n)$	$\Theta(n)$

TABLE I. We calculate (or estimate) the quantum Wasserstein complexity C_{W_1} and AC_{W_1} for several examples (See details in Appendix C). Here, $f(n) = \Theta(n)$ means that there exist constants c_1, c_2 such that $c_1 n \leq f(n) \leq c_2 n$ for any n .

introduce the quantum Wasserstein rate in the unitary dynamics generated by a Hamiltonian H , which will be useful for the connection between the quantum Wasserstein complexity and the circuit complexity. Given an n -qudit Hamiltonian H and an n -qudit pure state $|\psi\rangle$, the quantum Wasserstein rate of the unitary $U_t = \exp(-iHt)$ on the state $|\psi\rangle$ is defined as

$$R_{W_1}(\psi, H) = \lim_{\Delta t \rightarrow 0} \left\| \frac{|\psi\rangle\langle\psi| - U_{\Delta t}|\psi\rangle\langle\psi|U_{\Delta t}^\dagger}{\Delta t} \right\|_{W_1}. \quad (7)$$

Theorem 6 (Small incremental quantum Wasserstein complexity). *Given an n -qudit system with the Hamiltonian H acting on a k -qudit subsystem, and an n -qudit state $|\psi\rangle$, one has*

$$R_{W_1}(\psi, H) \leq 2\sqrt{2}k \|H\|_\infty, \quad (8)$$

where $\|\cdot\|_\infty$ denotes the operator norm.

We prove Theorem 6 in Appendix D. The small incremental quantum Wasserstein complexity is a key point to show the connection between circuit cost and the quantum Wasserstein complexity, for which we have the following relationship.

Theorem 7. *Given a unitary U , the circuit cost $\text{Cost}(U)$ is lower bounded by the quantum Wasserstein complexity $C_{W_1}(U)$ as*

$$\text{Cost}(U) \geq 4\sqrt{2}C_{W_1}(U). \quad (9)$$

We prove Theorem 7 in Appendix D. This relation can be generalized to the correlation-assisted quantum Wasserstein complexity AC_{W_1} defined in (4).

Corollary 8. *Under the assumptions of Theorem 7, we have*

$$\text{Cost}(U) \geq 4\sqrt{2}AC_{W_1}(U). \quad (10)$$

Besides the circuit cost, another important complexity measure of quantum circuits is the experimental cost to implement the circuit. Let us consider an n -qubit quantum circuit with a sequence of quantum gates $U = \prod_l U_l$, $U_l = \exp(-iH_l T_l)$, i.e., we run the gate U_1 first, then U_2 , and so on. T_l is the runtime of the l -th gate, and for each l , the time-independent Hamiltonian H_l acting on k_l qubits has the following spectral decomposition: $H_l = \sum_{i=1}^{2^{k_l}} h_i |h_i\rangle\langle h_i|$, $h_i \geq h_{i-1}, \forall i$.

Girolami and Anzà [62] proposed using the seminorm $E_l := (h_{2^{k_l}} - h_1)/2$ defined by the Hamiltonian H_l [63] to define the experimental cost of implementing the gate U_l and the circuit U in terms of physical resources as

$$\mathcal{R}_{U_l} := k_l E_l T_l, \quad \mathcal{R}_U := \sum_l \mathcal{R}_{U_l}. \quad (11)$$

Previously, the experimental cost of the quantum circuits was investigated using a weighted version of the Bures distance [62]. Here we find a new connection between the experimental cost \mathcal{R}_U and the quantum Wasserstein complexity $C_{W_1}(U)$ as follows:

Theorem 9. *The experimental cost $\mathcal{R}(U)$ of an n -qubit quantum circuit $U = \prod_l U_l$ is bounded from below by the quantum Wasserstein complexity $C_{W_1}(U)$ as*

$$\mathcal{R}(U) \geq \frac{1}{2}C_{W_1}(U). \quad (12)$$

The proof of Theorem 9 is presented in Appendix D. A similar argument also works for AC_{W_1} defined in (4).

Corollary 10. *Under the same conditions as in Theorem 9, we have*

$$\mathcal{R}(U) \geq \frac{1}{2}AC_{W_1}(U). \quad (13)$$

Example 1. Consider the CNOT gate $\exp(iH\pi/2)$, where $H = \frac{1}{2}(I - Z_1 - X_2 + Z_1 X_2)$. The experimental cost of CNOT is $\mathcal{R}(\text{CNOT}) = \pi$, and thus the quantum circuit $U = \prod_i \text{CNOT}_{i,i+1}$, which can be used to generate the cat state $(|0\rangle^{\otimes n} + |1\rangle^{\otimes n})/\sqrt{2}$, has the experimental cost $\mathcal{R}(U) = \pi n$. The quantum Wasserstein complexity is $C_{W_1}(U) = \Theta(n)$. (See Table I.) Thus, the quantum circuit $U = \prod_i \text{CNOT}_{i,i+1}$ has an experimental cost which is equivalent to the quantum Wasserstein complexity, up to some constant factor. Also relation (12) holds.

III. CONCLUSION

We introduce a new measure of quantum complexity, which we call *quantum Wasserstein complexity*. We provide a connection between this complexity and the circuit cost. This provides a useful lower bound for the circuit complexity of the shallow (or constant-depth) quantum circuits, in terms of the quantum Wasserstein complexity. We show that the quantum Wasserstein complexity provides a lower bound for the experimental cost of implementing quantum circuits.

Our results provide an application and operational interpretation of the quantum Wasserstein distance for shallow quantum circuits. This raises the interesting question: can one find a better bound for the circuit complexity of deep quantum circuits based on the quantum Wasserstein distance? To compute the quantum Wasserstein complexity, one needs to maximize

over all pure states, making it difficult to compute. So one naturally asks: can one discover an efficient way to approximate the quantum Wasserstein complexity? It is also interesting to study quantum Wasserstein distances of order other than 1, and how they relate to circuit complexity in quantum computation.

ACKNOWLEDGMENTS

We thank Michael Freedman, Chi-Ning Chou and Xun Gao for discussion of quantum circuit complexity; we thank Mar-

ius Junge, Weichen Gu and Hao Zhong for discussion of the quantum Wasserstein distance. This work was supported in part by the ARO Grant W911NF-19-1-0302 and the ARO MURI Grant W911NF-20-1-0082.

-
- [1] John Preskill, “Quantum Computing in the NISQ era and beyond,” *Quantum* **2**, 79 (2018).
- [2] Kishor Bharti, Alba Cervera-Lierta, Thi Ha Kyaw, Tobias Haug, Sumner Alperin-Lea, Abhinav Anand, Matthias Degroote, Hermanni Heimonen, Jakob S. Kottmann, Tim Menke, Wai-Keong Mok, Sukin Sim, Leong-Chuan Kwek, and Alán Aspuru-Guzik, “Noisy intermediate-scale quantum algorithms,” *Rev. Mod. Phys.* **94**, 015004 (2022).
- [3] Aram W Harrow and Ashley Montanaro, “Quantum computational supremacy,” *Nature* **549**, 203–209 (2017).
- [4] Alexander M. Dalzell, Aram W. Harrow, Dax Enshan Koh, and Rolando L. La Placa, “How many qubits are needed for quantum computational supremacy?” *Quantum* **4**, 264 (2020).
- [5] Thomas Fösel, Murphy Yuezhen Niu, Florian Marquardt, and Li Li, “Quantum circuit optimization with deep reinforcement learning,” *arXiv preprint arXiv:2103.07585* (2021).
- [6] J-H Bae, Paul M Alsing, Doyeol Ahn, and Warner A Miller, “Quantum circuit optimization using quantum Karnaugh map,” *Scientific reports* **10**, 1–8 (2020).
- [7] Ross Duncan, Aleks Kissinger, Simon Perdrix, and John van de Wetering, “Graph-theoretic Simplification of Quantum Circuits with the ZX-calculus,” *Quantum* **4**, 279 (2020).
- [8] Alternatively, one could aim to minimize the number of a particular type of gate in the circuit. For this goal, algorithms that achieve optimality—called resource-optimal circuit synthesis or re-synthesis algorithms [64]—include those that minimize the T-count [64–68] or CNOT-count [69] of Clifford+T circuits.
- [9] Nai-Hui Chia, Chi-Ning Chou, Jiayu Zhang, and Ruizhe Zhang, “Quantum meets the minimum circuit size problem,” *arXiv preprint arXiv:2108.03171* (2021).
- [10] Adi Botea, Akihiro Kishimoto, and Radu Marinescu, “On the complexity of quantum circuit compilation,” in *International Symposium on Combinatorial Search*, Vol. 9 (2018) pp. 138–142.
- [11] Leonard Susskind, “Computational complexity and black hole horizons,” *Fortschritte der Physik* **64**, 24–43 (2016).
- [12] Douglas Stanford and Leonard Susskind, “Complexity and shock wave geometries,” *Phys. Rev. D* **90**, 126007 (2014).
- [13] Adam R. Brown and Leonard Susskind, “Second law of quantum complexity,” *Phys. Rev. D* **97**, 086015 (2018).
- [14] Jonas Haferkamp, Philippe Faist, Naga BT Kothakonda, Jens Eisert, and Nicole Yunger Halpern, “Linear growth of quantum circuit complexity,” *Nature Physics* **18**, 528–532 (2022).
- [15] Zhi Li, “Short proofs of linear growth of quantum circuit complexity,” *arXiv preprint arXiv:2205.05668* (2022).
- [16] Michael A Nielsen, “A geometric approach to quantum circuit lower bounds,” *Quantum Information & Computation* **6**, 213–262 (2006).
- [17] Michael A. Nielsen, Mark R. Dowling, Mile Gu, and Andrew C. Doherty, “Quantum computation as geometry,” *Science* **311**, 1133–1135 (2006).
- [18] Michael A. Nielsen, Mark R. Dowling, Mile Gu, and Andrew C. Doherty, “Optimal control, geometry, and quantum computing,” *Phys. Rev. A* **73**, 062323 (2006).
- [19] Mark R Dowling and Michael A Nielsen, “The geometry of quantum computation,” *Quantum Information & Computation* **8**, 861–899 (2008).
- [20] D Gottesman, “The Heisenberg representation of quantum computers,” in *Proc. XXII International Colloquium on Group Theoretical Methods in Physics, 1998* (1998) pp. 32–43.
- [21] Richard Jozsa and Maarten Van den Nest, “Classical simulation complexity of extended Clifford circuits,” *Quantum Information & Computation* **14**, 633–648 (2014).
- [22] Dax Enshan Koh, “Further extensions of Clifford circuits and their classical simulation complexities,” *Quantum Information & Computation* **17**, 0262–0282 (2017).
- [23] Adam Bouland, Joseph F. Fitzsimons, and Dax Enshan Koh, “Complexity Classification of Conjugated Clifford Circuits,” in *33rd Computational Complexity Conference (CCC 2018)*, Leibniz International Proceedings in Informatics (LIPIcs), Vol. 102, edited by Rocco A. Servedio (Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik, Dagstuhl, Germany, 2018) pp. 21:1–21:25.
- [24] Mithuna Yoganathan, Richard Jozsa, and Sergii Strelchuk, “Quantum advantage of unitary Clifford circuits with magic state inputs,” *Proceedings of the Royal Society A* **475**, 20180427 (2019).
- [25] Sergey Bravyi, Graeme Smith, and John A. Smolin, “Trading classical and quantum computational resources,” *Phys. Rev. X* **6**, 021043 (2016).
- [26] Sergey Bravyi, Dan Browne, Pádraic Calpin, Earl Campbell, David Gosset, and Mark Howard, “Simulation of quantum circuits by low-rank stabilizer decompositions,” *Quantum* **3**, 181 (2019).
- [27] Mark Howard and Earl Campbell, “Application of a resource theory for magic states to fault-tolerant quantum computing,” *Phys. Rev. Lett.* **118**, 090501 (2017).
- [28] James R. Seddon, Bartosz Regula, Hakop Pashayan, Yingkai Ouyang, and Earl T. Campbell, “Quantifying quantum speedups: Improved classical simulation from tighter magic monotones,” *PRX Quantum* **2**, 010345 (2021).
- [29] James R. Seddon and Earl T. Campbell, “Quantifying magic

- for multi-qubit operations,” *Proc. R. Soc. A.* **475** (2019), doi.org/10.1098/rspa.2019.0251.
- [30] Xin Wang, Mark M Wilde, and Yuan Su, “Quantifying the magic of quantum channels,” *New Journal of Physics* **21**, 103002 (2019).
- [31] Dax Enshan Koh, Mark D Penney, and Robert W Spekkens, “Computing quopit Clifford circuit amplitudes by the sum-over-paths technique,” *Quantum Information & Computation* **17**, 1081–1095 (2017).
- [32] Kaifeng Bu and Dax Enshan Koh, “Efficient classical simulation of Clifford circuits with nonstabilizer input states,” *Phys. Rev. Lett.* **123**, 170502 (2019).
- [33] Kaifeng Bu and Dax Enshan Koh, “Classical simulation of quantum circuits by half Gauss sums,” *Commun. Math. Phys.* **390**, 471–500 (2022).
- [34] J. Eisert, “Entangling power and quantum circuit complexity,” *Phys. Rev. Lett.* **127**, 020501 (2021).
- [35] Kaifeng Bu, Roy J Garcia, Arthur Jaffe, Dax Enshan Koh, and Lu Li, “Complexity of quantum circuits via sensitivity, magic, and coherence,” *arXiv preprint arXiv:2204.12051* (2022).
- [36] Leonid V Kantorovich, “Mathematical methods of organizing and planning production,” *Management Science* **6**, 366–422 (1960).
- [37] Leonid Nisonovich Vaserstein, “Markov processes over denumerable products of spaces, describing large systems of automata,” *Problemy Peredachi Informatsii* **5**, 64–72 (1969).
- [38] Cédric Villani, *Optimal Transport* (Springer, Berlin, Germany, 2009).
- [39] Gabriel Peyré, Marco Cuturi, *et al.*, “Computational optimal transport: With applications to data science,” *Foundations and Trends® in Machine Learning* **11**, 355–607 (2019).
- [40] Yossi Rubner, Carlo Tomasi, and Leonidas J Guibas, “The earth mover’s distance as a metric for image retrieval,” *International Journal of Computer Vision* **40**, 99–121 (2000).
- [41] Martin Arjovsky, Soumith Chintala, and Léon Bottou, “Wasserstein generative adversarial networks,” in *Proceedings of the 34th International Conference on Machine Learning*, Proceedings of Machine Learning Research, Vol. 70, edited by Doina Precup and Yee Whye Teh (PMLR, 2017) pp. 214–223.
- [42] Ian Goodfellow, Jean Pouget-Abadie, Mehdi Mirza, Bing Xu, David Warde-Farley, Sherjil Ozair, Aaron Courville, and Yoshua Bengio, “Generative adversarial nets,” in *Advances in Neural Information Processing Systems*, Vol. 27, edited by Z. Ghahramani, M. Welling, C. Cortes, N. Lawrence, and K.Q. Weinberger (Curran Associates, Inc., 2014).
- [43] Ishaan Gulrajani, Faruk Ahmed, Martin Arjovsky, Vincent Dumoulin, and Aaron C Courville, “Improved training of Wasserstein GANs,” in *Advances in Neural Information Processing Systems*, Vol. 30, edited by I. Guyon, U. Von Luxburg, S. Bengio, H. Wallach, R. Fergus, S. Vishwanathan, and R. Garnett (Curran Associates, Inc., 2017).
- [44] Eric A Carlen and Jan Maas, “An analog of the 2-Wasserstein metric in non-commutative probability under which the fermionic Fokker–Planck equation is gradient flow for the entropy,” *Communications in Mathematical Physics* **331**, 887–926 (2014).
- [45] Eric A Carlen and Jan Maas, “Gradient flow and entropy inequalities for quantum Markov semigroups with detailed balance,” *Journal of Functional Analysis* **273**, 1810–1869 (2017).
- [46] Eric A Carlen and Jan Maas, “Non-commutative calculus, optimal transport and functional inequalities in dissipative quantum systems,” *Journal of Statistical Physics* **178**, 319–378 (2020).
- [47] Cambyse Rouzé and Nilanjana Datta, “Concentration of quantum states from quantum functional and transportation cost inequalities,” *Journal of Mathematical Physics* **60**, 012202 (2019).
- [48] Nilanjana Datta and Cambyse Rouzé, “Relating relative entropy, optimal transport and Fisher information: a quantum HWI inequality,” in *Annales Henri Poincaré*, Vol. 21 (Springer, 2020) pp. 2115–2150.
- [49] Li Gao, Marius Junge, and Nicholas LaRacuente, “Fisher Information and Logarithmic Sobolev Inequality for Matrix-Valued Functions,” in *Annales Henri Poincaré*, Vol. 21 (Springer, 2020) pp. 3409–3478.
- [50] Tan Van Vu and Yoshihiko Hasegawa, “Geometrical bounds of the irreversibility in Markovian systems,” *Phys. Rev. Lett.* **126**, 010601 (2021).
- [51] Yongxin Chen, Tryphon T. Georgiou, Lipeng Ning, and Allen Tannenbaum, “Matricial Wasserstein-1 distance,” *IEEE Control Systems Letters* **1**, 14–19 (2017).
- [52] Ernest K Ryu, Yongxin Chen, Wuchen Li, and Stanley Osher, “Vector and matrix optimal mass transport: theory, algorithm, and applications,” *SIAM Journal on Scientific Computing* **40**, A3675–A3698 (2018).
- [53] Giacomo De Palma, Milad Marvian, Dario Trevisan, and Seth Lloyd, “The quantum Wasserstein distance of order 1,” *IEEE Transactions on Information Theory* **67**, 6627–6643 (2021).
- [54] Bobak Toussi Kiani, Giacomo De Palma, Milad Marvian, Zi-Wen Liu, and Seth Lloyd, “Learning quantum data with the quantum earth mover’s distance,” *Quantum Science and Technology* **7**, 045002 (2022).
- [55] Giacomo De Palma and Cambyse Rouzé, “Quantum concentration inequalities,” *Annales Henri Poincaré* (2022), [10.1007/s00023-022-01181-1](https://doi.org/10.1007/s00023-022-01181-1).
- [56] Christoph Hirche, Cambyse Rouzé, and Daniel Stilck França, “Quantum differential privacy: An information theory perspective,” *arXiv preprint arXiv:2202.10717* (2022).
- [57] Giacomo De Palma, Milad Marvian, Cambyse Rouzé, and Daniel Stilck França, “Limitations of variational quantum algorithms: a quantum optimal transport approach,” *arXiv preprint arXiv:2204.03455* (2022).
- [58] Here, the trace distance d is defined as one-half the metric induced by the trace norm, i.e. $d : (x, y) \mapsto \frac{1}{2} \|x - y\|_1$.
- [59] Michael A. Nielsen and Isaac L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, 2010).
- [60] A. Kitaev, A. Shen, and M. Vyalyi, *Classical and quantum computation* (American Mathematical Society, 2002).
- [61] Scott Aaronson, “The complexity of quantum states and transformations: from quantum money to black holes,” *arXiv preprint arXiv:1607.05256* (2016).
- [62] Davide Girolami and Fabio Anzà, “Quantifying the difference between many-body quantum states,” *Phys. Rev. Lett.* **126**, 170502 (2021).
- [63] Sergio Boixo, Steven T. Flammia, Carlton M. Caves, and JM Geremia, “Generalized limits for single-parameter quantum estimation,” *Phys. Rev. Lett.* **98**, 090401 (2007).
- [64] Michele Mosca and Priyanka Mukhopadhyay, “A polynomial time and space heuristic algorithm for T-count,” *Quantum Science and Technology* **7**, 015003 (2021).
- [65] David Gosset, Vadym Kliuchnikov, Michele Mosca, and Vincent Russo, “An algorithm for the T-count,” *Quantum Information & Computation* **14**, 1261–1276 (2014).
- [66] Matthew Amy, Dmitri Maslov, and Michele Mosca, “Polynomial-time T-depth optimization of Clifford+T circuits via matroid partitioning,” *IEEE Transactions on Computer-Aided Design of Integrated Circuits and Systems* **33**, 1476–1489 (2014).

- [67] Matthew Amy and Michele Mosca, “T-count optimization and Reed–Muller codes,” *IEEE Transactions on Information Theory* **65**, 4771–4784 (2019).
- [68] Vlad Gheorghiu, Michele Mosca, and Priyanka Mukhopadhyay, “T-count and T-depth of any multi-qubit unitary,” *arXiv preprint arXiv:2110.10292* (2021).
- [69] Vlad Gheorghiu, Sarah Meng Li, Michele Mosca, and Priyanka Mukhopadhyay, “Reducing the CNOT count for Clifford+T circuits on NISQ architectures,” *arXiv preprint arXiv:2011.12191* (2020).
- [70] Sebastian Deffner, “Geometric quantum speed limits: a case for Wigner phase space,” *New Journal of Physics* **19**, 103018 (2017).

Appendix A: Basic properties of quantum Wasserstein norm of order 1

We state three basic properties of the quantum Wasserstein norm of order 1 [53], which is defined for all traceless, Hermitian operators A :

$$\|A\|_{W_1} := \frac{1}{2} \min_{\{X_i\}} \left\{ \sum_i \|X_i\|_1 : A = \sum_i X_i, X_i \in \mathcal{O}(\mathcal{H}), \text{Tr}_i[X_i] = 0 \forall i \right\}.$$

Lemma 11 (De Palma et al. [53]). *(Tensorization)* Given an n -qudit traceless, Hermitian operator A , and a subset $S \subset [n]$ with complement S^c ,

$$\|A\|_{W_1} \geq \|\text{Tr}_S[A]\|_{W_1} + \|\text{Tr}_{S^c}[A]\|_{W_1}. \quad (\text{A1})$$

Lemma 12 (De Palma et al. [53]). *Given an n -qudit traceless, Hermitian operator A and a subset $S \subset [n]$ with $\text{Tr}_S[A] = 0$,*

$$\|A\|_{W_1} \leq |S| \frac{d^2 - 1}{d^2} \|A\|_1. \quad (\text{A2})$$

Hence, for any n -qudit quantum channel Λ acting nontrivially on k -qudit subsystems,

$$\|\rho - \Lambda(\rho)\|_{W_1} \leq k. \quad (\text{A3})$$

Lemma 13 (De Palma et al. [53]). *Given an n -qudit traceless, Hermitian operator A , we have*

$$\frac{1}{2} \|A\|_1 \leq \|A\|_{W_1} \leq \frac{n}{2} \|A\|_1. \quad (\text{A4})$$

Moreover, if there exists some i such that $\text{Tr}_i[A] = 0$, then we have

$$\|A\|_{W_1} = \frac{1}{2} \|A\|_1. \quad (\text{A5})$$

Appendix B: Basic properties of quantum Wasserstein complexity

Proposition 14 (Restatement of Proposition 2). *The quantum Wasserstein complexity C_{W_1} satisfies the following properties:*

- (1) *Faithfulness:* $C_{W_1}(\Lambda) = 0$ if and only if Λ is the identity map;
- (2) *Convexity:* $C_{W_1}(\sum_i p_i \Lambda_i) \leq \sum_i p_i C_{W_1}(\Lambda_i)$, where $p_i \geq 0$ and $\sum_i p_i = 1$;
- (3) *Subadditivity under concatenation:* $C_{W_1}(\Lambda_1 \circ \Lambda_2) \leq C_{W_1}(\Lambda_1) + C_{W_1}(\Lambda_2)$;
- (4) $C_{W_1}(\Lambda_1 \otimes \Lambda_2) \leq C_{W_1}(\Lambda_1 \otimes I) + C_{W_1}(\Lambda_2 \otimes I)$;
- (5) *Superadditivity under tensorization:* $C_{W_1}(\Lambda_1 \otimes \Lambda_2) \geq C_{W_1}(\Lambda_1) + C_{W_1}(\Lambda_2)$;
- (6) *For a unitary channel U ,* $C_{W_1}(U) = C_{W_1}(U^\dagger)$;
- (7) *If Λ acts nontrivially on a k -qudit subsystem, then we have $C_{W_1}(\Lambda) \leq k$.*

Proof. Property (1) follows directly from the faithfulness of the quantum W_1 norm, properties (2)–(4) follow directly from the triangle inequality of the quantum W_1 norm, and property (5) follows from the tensorization of W_1 norm (see Lemma 11) as follows

$$\|\rho - \Lambda_1 \otimes \Lambda_2(\rho)\|_{W_1} \geq \|\rho_1 - \Lambda_1(\rho_1)\|_{W_1} + \|\rho_2 - \Lambda_2(\rho_2)\|_{W_1},$$

where ρ_i (for $i = 1, 2$) denotes the corresponding reduced state. Property (6) follows directly from the definition of the quantum Wasserstein complexity. Finally, Property (7) follows from Lemma 12. \square

Proposition 15 (Restatement of Proposition 4). *The correlation-assisted Wasserstein complexity $\text{AC}_{W_1}(\Lambda)$ satisfies the following properties:*

- (1) *Faithfulness:* $\text{AC}_{W_1}(\Lambda) = 0$ if and only if Λ is the identity map;

(2) *Convexity*: $\text{AC}_{W_1}(\sum_i p_i \Lambda_i) \leq \sum_i p_i \text{AC}_{W_1}(\Lambda_i)$, where $p_i \geq 0$ and $\sum_i p_i = 1$;

(3) *Subadditivity under concatenation*: $\text{AC}_{W_1}(\Lambda_1 \circ \Lambda_2) \leq \text{AC}_{W_1}(\Lambda_1) + \text{AC}_{W_1}(\Lambda_2)$;

(4) *Additivity under tensorization*: $\text{AC}_{W_1}(\Lambda_1 \otimes \Lambda_2) = \text{AC}_{W_1}(\Lambda_1) + \text{AC}_{W_1}(\Lambda_2)$.

Proof. The property (1) follows from the faithfulness of the quantum W_1 distance. Properties (2) and (3) follow from the triangle inequality of the quantum W_1 distance. Hence, it suffices for us to prove the property (4). Since

$$C_{W_1}(\Lambda_1 \otimes \Lambda_2) \leq C_{W_1}(\Lambda_1 \otimes I) + C_{W_1}(\Lambda_2 \otimes I), \quad (\text{B1})$$

by the definition of EW_1 , we have

$$\text{AC}_{W_1}(\Lambda_1 \otimes \Lambda_2) \leq \text{AC}_{W_1}(\Lambda_1) + \text{AC}_{W_1}(\Lambda_2). \quad (\text{B2})$$

Besides, since

$$C_{W_1}(\Lambda_1 \otimes \Lambda_2) \geq C_{W_1}(\Lambda_1) + C_{W_1}(\Lambda_2), \quad (\text{B3})$$

we have

$$\text{AC}_{W_1}(\Lambda_1 \otimes \Lambda_2) \geq \text{AC}_{W_1}(\Lambda_1) + \text{AC}_{W_1}(\Lambda_2). \quad (\text{B4})$$

□

Appendix C: Estimation of C_{W_1} and AC_{W_1} : interesting examples

Let us start with the simplest case, where U is a single-qudit unitary.

Claim 16. For a 1-qudit unitary U , we have

$$C_{W_1}(U) = \text{AC}_{W_1}(U). \quad (\text{C1})$$

Proof. For any 1-qudit unitary and pure state $|\psi\rangle$, we have

$$\| |\psi\rangle\langle\psi| - U|\psi\rangle\langle\psi|U^\dagger \|_{W_1} = \frac{1}{2} \| |\psi\rangle\langle\psi| - U|\psi\rangle\langle\psi|U^\dagger \|_1 = \sqrt{1 - |\langle\psi|U|\psi\rangle|^2}. \quad (\text{C2})$$

Let $U = \sum_j \exp(i\theta_j) |a_j\rangle\langle a_j|$ be the eigenvalue decomposition of U . Then, the state $|\psi\rangle$ can be written as $|\psi\rangle = \sum_j c_j |a_j\rangle$, from which it follows that $|\langle\psi|U|\psi\rangle|^2 = |\sum_j |c_j|^2 \exp(i\theta_j)|^2 = |\sum_j d_j \exp(i\theta_j)|^2$, where $d_j = |c_j|^2$. Hence

$$\begin{aligned} C_{W_1}(U) &= \max_{\psi} \| |\psi\rangle\langle\psi| - U|\psi\rangle\langle\psi| \|_{W_1} = \max_{\psi} \sqrt{1 - |\langle\psi|U|\psi\rangle|^2} \\ &= \sqrt{1 - \min_{\psi} |\langle\psi|U|\psi\rangle|^2} = \sqrt{1 - \min_{d_j: \sum_j d_j=1} \left| \sum_j d_j \exp(i\theta_j) \right|^2}. \end{aligned}$$

Let us consider $U \otimes I_m$ for any $m \geq 1$. For any $(m+1)$ -qudit state $|\psi\rangle = \sum_j \sqrt{\lambda_j} |\psi_j\rangle |j\rangle$, with $|j\rangle$ being a basis on m -qudit systems and $\sum_j \lambda_j = 1$, we have

$$\langle\psi|U \otimes I|\psi\rangle = \sum_j \lambda_j \langle\psi_j|U|\psi_j\rangle,$$

where for each j , $|\psi_j\rangle = \sum_k \sqrt{d_{jk}} e^{i\theta_{jk}} |a_k\rangle$, and

$$\langle\psi_j|U|\psi_j\rangle = \sum_k d_{jk} \exp(i\theta_k).$$

Hence

$$\langle\psi|U \otimes I|\psi\rangle = \sum_{j,k} \lambda_j d_{jk} \exp(i\theta_k) = \sum_k f_k \exp(i\theta_k),$$

where $f_k = \sum_j \lambda_j d_{jk}$, and $\sum_k f_k = 1$. Hence, we have the following expression for $\text{AC}_{W_1}(U)$:

$$\text{AC}_{W_1}(U) = \sup_m \max_{\psi} \| |\psi\rangle\langle\psi| - U \otimes I_m |\psi\rangle\langle\psi|U^\dagger \otimes I_m \|_{W_1} = \sqrt{1 - \min_{f_k: \sum_k f_k=1} \left| \sum_k f_k \exp(i\theta_k) \right|^2},$$

which implies that $AC_{W_1}(U) = C_{W_1}(U)$. □

Example 1: Depolarizing channel

Let us consider the single-qudit depolarizing channel, for which we have the following statement.

Claim 17. For the 1-qudit depolarizing channel $D_p(\cdot) = p(\cdot) + (1-p)\text{Tr}[\cdot]I/d$, we have

$$C_{W_1}(D_p) = (1-p)(1-1/d), \quad (C3)$$

and

$$AC_{W_1}(D_p) = (1-p)(1-1/d^2). \quad (C4)$$

Proof. For any 1-qudit state, we have

$$\| |\psi\rangle\langle\psi| - D_p(|\psi\rangle\langle\psi|) \|_{W_1} = \frac{1}{2} \| |\psi\rangle\langle\psi| - D_p(|\psi\rangle\langle\psi|) \|_1 = (1-p)(1-1/d).$$

Hence, we have

$$C_{W_1}(D_p) = (1-p)(1-1/d).$$

Now, let us take a 2-qudit state $|\psi\rangle = \frac{1}{\sqrt{d}} \sum_j |j\rangle |j\rangle$ and the quantum channel $D_p \otimes I$. Then,

$$\| |\psi\rangle\langle\psi| - D_p \otimes I(|\psi\rangle\langle\psi|) \|_{W_1} = \frac{1}{2} \| |\psi\rangle\langle\psi| - D_p \otimes I(|\psi\rangle\langle\psi|) \|_1 = (1-p)(1-1/d^2).$$

This implies that

$$AC_{W_1}(D_p) \geq (1-p)(1-1/d^2).$$

Besides, let us prove that for any integer $m \geq 0$,

$$C_{W_1}(D_p \otimes I_m) \leq (1-p)(1-1/d^2).$$

For any $(m+1)$ -qudit state $|\psi\rangle$, we have

$$\begin{aligned} & \| |\psi\rangle\langle\psi| - D_p \otimes I_m(|\psi\rangle\langle\psi|) \|_{W_1} \\ &= (1-p) \left\| |\psi\rangle\langle\psi| - \frac{I}{d} \otimes \text{Tr}_1[|\psi\rangle\langle\psi|] \right\|_{W_1} \\ &= (1-p) \left\| |\psi\rangle\langle\psi| - \frac{1}{d^2} \sum_{s,t \in \mathbb{Z}_d} X_1^s Z_1^t |\psi\rangle\langle\psi| Z_1^{-t} X_1^{-s} \right\|_{W_1} \\ &= (1-p) \frac{d^2-1}{d^2} \left\| \mathbb{E}_{(s,t) \neq (0,0)} (|\psi\rangle\langle\psi| - X_1^s Z_1^t |\psi\rangle\langle\psi| Z_1^{-t} X_1^{-s}) \right\|_{W_1} \\ &\leq (1-p) \frac{d^2-1}{d^2} \mathbb{E}_{(s,t) \neq (0,0)} \left\| |\psi\rangle\langle\psi| - X_1^s Z_1^t |\psi\rangle\langle\psi| Z_1^{-t} X_1^{-s} \right\|_{W_1} \\ &= (1-p) \frac{d^2-1}{d^2} \frac{1}{2} \left\| |\psi\rangle\langle\psi| - X_1^s Z_1^t |\psi\rangle\langle\psi| Z_1^{-t} X_1^{-s} \right\|_1 \\ &\leq (1-p) \frac{d^2-1}{d^2}, \end{aligned}$$

where the third line comes from the fact that

$$\frac{1}{d^2} \sum_{(s,t) \in \mathbb{Z}_d \times \mathbb{Z}_d} X^s Z^t (\cdot) Z^{-t} X^{-s} = \text{Tr}[\cdot] \frac{I}{d},$$

the fifth line comes from the convexity of the quantum W_1 distance, and the last line comes from the fact that $\text{Tr}_1[|\psi\rangle\langle\psi| - X_1^s Z_1^t |\psi\rangle\langle\psi| Z_1^{-t} X_1^{-s}] = 0$, and thus $\| |\psi\rangle\langle\psi| - X_1^s Z_1^t |\psi\rangle\langle\psi| Z_1^{-t} X_1^{-s} \|_{W_1} = \frac{1}{2} \| |\psi\rangle\langle\psi| - X_1^s Z_1^t |\psi\rangle\langle\psi| Z_1^{-t} X_1^{-s} \|_1$. Thus, we have

$$AC_{W_1}(D_p) \leq (1-p)(1-1/d^2).$$

□

Corollary 18. For the n -fold tensor product of the depolarizing channel $D_p^{\otimes n}$, we have

$$n(1-p)(1-1/d) \leq C_{W_1}(D_p^{\otimes n}) \leq n(1-p)(1-1/d^2) \quad (C5)$$

and

$$AC_{W_1}(D_p^{\otimes n}) = n(1-p)(1-1/d^2). \quad (C6)$$

Example 2: n -fold tensor product of Hadamard gates

Let us consider the n -fold tensor product of Hadamard gates, i.e., $H^{\otimes n}$. The quantum Wasserstein complexity of $H^{\otimes n}$ is

$$C_{W_1}(H^{\otimes n}) = n. \quad (C7)$$

Proof of (C7). The inequality $C_{W_1}(H^{\otimes n}) \leq n$ holds because $\|\rho - \sigma\|_{W_1} \leq \frac{n}{2} \|\rho - \sigma\|_1 \leq n$, which follows from Lemma 12. To prove that $C_{W_1}(H^{\otimes n}) \geq n$, we use the fact that for the Hadamard gate H , there exists some pure state $|\phi\rangle$ such that $H|\phi\rangle \perp |\phi\rangle$. Hence,

$$\begin{aligned} C_{W_1}(H^{\otimes n}) &\geq \|(H|\phi\rangle\langle\phi|H)^{\otimes n} - |\phi\rangle\langle\phi|^{\otimes n}\|_{W_1} \\ &= \sum_i \|H|\phi\rangle\langle\phi|H - |\phi\rangle\langle\phi|\|_{W_1} \\ &= \frac{1}{2} \sum_i \|H|\phi\rangle\langle\phi|H - |\phi\rangle\langle\phi|\|_1 \\ &= n, \end{aligned}$$

where the first line follows from the definition of C_{W_1} ; the second line follows from the tensorization of the quantum W_1 distance, i.e., $\|\rho_1 \otimes \rho_2 - \sigma_1 \otimes \sigma_2\|_{W_1} = \|\rho_1 - \sigma_1\|_{W_1} + \|\rho_2 - \sigma_2\|_{W_1}$ [53]; the third line follows from the fact that $\|\rho - \sigma\|_{W_1} = \frac{1}{2} \|\rho_1 - \sigma_1\|_1$ for 1-qudit states ρ, σ [53]; and the last line follows from the fact that $\|\rho_1 - \sigma_1\|_1 = 1$ as $H|\phi\rangle \perp |\phi\rangle$. \square

Besides, as $C_{W_1}(H^{\otimes n} \otimes I_m) \leq n$, we also have

$$AC_{W_1}(H^{\otimes n}) = n. \quad (C8)$$

Example 3: CNOT gates

Let us consider the n -qubit quantum circuit $\prod_i \text{CNOT}_{i,i+1}$ (See Fig. 3). Then we have

$$C_{W_1}\left(\prod_i \text{CNOT}_{i,i+1}\right) = \Theta(n), \quad (C9)$$

where $f(n) = \Theta(n)$ means that there exist constants c_1, c_2 such that $c_1 n \leq f(n) \leq c_2 n$ for any n . (1) The upper bound $C_{W_1}(\prod_i \text{CNOT}_{i,i+1}) \leq n$ holds for the same reason as the one in the above example. (2) Next, we show that $C_{W_1}(\prod_i \text{CNOT}_{i,i+1}) \geq n/2$. To this end, let us take the input state to be $|+\rangle \otimes |0\rangle^{n-1}$. Then, the output state is $\prod_i \text{CNOT}_{i,i+1} |+\rangle \otimes |0\rangle^{n-1} = \frac{1}{\sqrt{2}}(|0\rangle^n + |1\rangle^n)$, which we denote as $|\text{Cat}_{1/2}\rangle$. Hence

$$\begin{aligned} C_{W_1}\left(\prod_i \text{CNOT}_{i,i+1}\right) &\geq \||+\rangle\langle+| \otimes |0\rangle\langle 0|^{n-1} - |\text{Cat}_{1/2}\rangle\langle\text{Cat}_{1/2}|\|_{W_1} \\ &\geq \frac{1}{2} \sum_i \|\rho_i - \sigma_i\|_1 = n/2, \end{aligned}$$

where ρ_i (or σ_i) is the reduced state of $|+\rangle\langle+| \otimes |0\rangle\langle 0|^{n-1}$ (or $|\text{Cat}_{1/2}\rangle\langle\text{Cat}_{1/2}|$) on the i -th position, and the second inequality comes from the tensorization of the quantum W_1 distance.

Hence, we also have

$$AC_{W_1}\left(\prod_i \text{CNOT}_{i,i+1}\right) = \Theta(n). \quad (C10)$$




FIG. 3. A circuit diagram for the circuit of cascading CNOT gates $\prod_i \text{CNOT}_{i,i+1}$.

Appendix D: Application in quantum circuit complexity and experimental cost

Lemma 19. *Given an n -qudit system with a Hamiltonian H acting on a k -qudit subsystem, and an n -qudit state $|\psi\rangle$, for the unitary $U_{\Delta t} = \exp(-iH\Delta t)$, one has*

$$\left\| |\psi\rangle\langle\psi| - U_{\Delta t} |\psi\rangle\langle\psi| U_{\Delta t}^\dagger \right\|_{W_1} \leq 2\sqrt{2k} \frac{d^2 - 1}{d^2} \|H\|_\infty |\Delta t| e^{\|H\|_\infty |\Delta t|}, \quad (\text{D1})$$

which implies that

$$R_{W_1}(\psi, H) \leq 2\sqrt{2} \|H\|_\infty. \quad (\text{D2})$$

Proof. Since H acts on a k -qudit subsystem, there exists a subsystem S for which $\text{Tr}_S [|\psi\rangle\langle\psi| - U_{\Delta t} |\psi\rangle\langle\psi| U_{\Delta t}^\dagger] = 0$. Hence, by Lemma 12, we have

$$\left\| |\psi\rangle\langle\psi| - U_{\Delta t} |\psi\rangle\langle\psi| U_{\Delta t}^\dagger \right\|_{W_1} \leq k \frac{d^2 - 1}{d^2} \left\| |\psi\rangle\langle\psi| - U_{\Delta t} |\psi\rangle\langle\psi| U_{\Delta t}^\dagger \right\|_1 = 2k \frac{d^2 - 1}{d^2} \sqrt{1 - |\langle\psi| U_{\Delta t} |\psi\rangle|^2},$$

Let us define $|\psi_{\Delta t}\rangle = U_{\Delta t} |\psi\rangle$. Then, the Taylor expansion of $|\psi_{\Delta t}\rangle\langle\psi_{\Delta t}|$ is

$$|\psi_{\Delta t}\rangle\langle\psi_{\Delta t}| = |\psi\rangle\langle\psi| + i\Delta t [H, |\psi\rangle\langle\psi|] + \frac{(i\Delta t)^2}{2!} [H, [H, |\psi\rangle\langle\psi|]] + \dots$$

where $[A, B] = AB - BA$ denotes the commutator between A and B . Then

$$\begin{aligned} 1 - |\langle\psi| U_{\Delta t} |\psi\rangle|^2 &= 1 - \left[\text{Tr} [|\psi\rangle\langle\psi|^2] + i\Delta t \text{Tr} [|\psi\rangle\langle\psi| [H, |\psi\rangle\langle\psi|]] + \frac{(i\Delta t)^2}{2!} \text{Tr} [|\psi\rangle\langle\psi| [H, [H, |\psi\rangle\langle\psi|]]] + \dots \right] \\ &= -\frac{(i\Delta t)^2}{2!} \text{Tr} [|\psi\rangle\langle\psi| [H, [H, |\psi\rangle\langle\psi|]]] + \dots \end{aligned}$$

where $\text{Tr} [|\psi\rangle\langle\psi| [H, |\psi\rangle\langle\psi|]] = 0$. Let us define ad_H as $\text{ad}_H(A) := [H, A]$. Then, the above formula can be rewritten as

$$1 - |\langle\psi| U_{\Delta t} |\psi\rangle|^2 = -\sum_{k \geq 2} \frac{(i\Delta t)^k}{k!} \text{Tr} [|\psi\rangle\langle\psi| \text{ad}_H^k(|\psi\rangle\langle\psi|)].$$

For each term $\text{Tr} [|\psi\rangle\langle\psi| \text{ad}_H^k(|\psi\rangle\langle\psi|)]$, by Hölder's inequality, we have

$$\left| \text{Tr} [|\psi\rangle\langle\psi| \text{ad}_H^k(|\psi\rangle\langle\psi|)] \right| \leq \left\| \text{ad}_H^k(|\psi\rangle\langle\psi|) \right\|_\infty \leq \|H\|_\infty^k 2^k,$$

which implies that

$$|1 - |\langle\psi| U_{\Delta t} |\psi\rangle|^2| \leq 2 \|H\|_\infty^2 |\Delta t|^2 \exp(2 \|H\|_\infty |\Delta t|). \quad (\text{D3})$$

Hence, we have

$$\left\| |\psi\rangle\langle\psi| - U_{\Delta t} |\psi\rangle\langle\psi| U_{\Delta t}^\dagger \right\|_{W_1} \leq 2\sqrt{2k} \frac{d^2 - 1}{d^2} \|H\|_\infty |\Delta t| e^{\|H\|_\infty |\Delta t|}. \quad (\text{D4})$$

□

Theorem 20. Given a unitary U , the circuit cost $\text{Cost}(U)$ is lower bounded in terms of $C_{W_1}(U)$ as follows

$$\text{Cost}(U) \geq 4\sqrt{2}C_{W_1}(U). \quad (\text{D5})$$

Proof. First, let us take a Trotter decomposition of U such that for arbitrarily small $\varepsilon > 0$,

$$\|U - V_N\|_\infty \leq \varepsilon,$$

where V_N is defined as follows

$$V_N := \prod_{t=1}^N W_t,$$

$$W_t := \exp\left(-\frac{i}{N} \sum_{j=1}^m r_j \left(\frac{t}{N}\right) h_j\right).$$

and

$$W_t = \lim_{l \rightarrow \infty} W_t^{(l)},$$

$$W_t^{(l)} := \left(W_{t,1}^{1/l} \cdots W_{t,m}^{1/l}\right)^l,$$

$$W_{t,j} := \exp\left(-\frac{i}{N} r_j \left(\frac{t}{N}\right) h_j\right).$$

where $\|h_j\|_\infty \leq 1$ for any j . Let us define $|\psi_t\rangle = W_t |\psi_{t-1}\rangle$ with $|\psi_0\rangle = |\psi\rangle$, then by the triangle inequality of the quantum W_1 distance, we have

$$\left\| |\psi\rangle\langle\psi| - V_N |\psi\rangle\langle\psi| V_N^\dagger \right\|_{W_1} \leq \sum_{t=1}^N \left\| |\psi_{t-1}\rangle\langle\psi_{t-1}| - |\psi_t\rangle\langle\psi_t| \right\|_{W_1}. \quad (\text{D6})$$

For each $\left\| |\psi_{t-1}\rangle\langle\psi_{t-1}| - |\psi_t\rangle\langle\psi_t| \right\|_{W_1}$, we have

$$\begin{aligned} \left\| |\psi_{t-1}\rangle\langle\psi_{t-1}| - |\psi_t\rangle\langle\psi_t| \right\|_{W_1} &= \left\| |\psi_{t-1}\rangle\langle\psi_{t-1}| - W_t |\psi_{t-1}\rangle\langle\psi_{t-1}| W_t^\dagger \right\|_{W_1} \\ &= \lim_{l \rightarrow \infty} \left\| |\psi_{t-1}\rangle\langle\psi_{t-1}| - W_t^{(l)} |\psi_{t-1}\rangle\langle\psi_{t-1}| W_t^{(l)\dagger} \right\|_{W_1} \\ &\leq \lim_{l \rightarrow \infty} l \sum_j \frac{4\sqrt{2}}{Nl} \left| r_j \left(\frac{t}{N}\right) \right| \exp\left(\frac{1}{Nl} \left| r_j \left(\frac{t}{N}\right) \right|\right) \\ &= \frac{4\sqrt{2}}{N} \sum_j \left| r_j \left(\frac{t}{N}\right) \right|, \end{aligned}$$

where the last inequality comes from the triangle inequality of quantum W_1 distance and Lemma 19 by taking $\Delta t = \frac{1}{l}$. Therefore,

$$\left\| |\psi\rangle\langle\psi| - V_N |\psi\rangle\langle\psi| V_N^\dagger \right\|_{W_1} \leq \frac{4\sqrt{2}}{N} \sum_{t=1}^N \sum_{j=1}^m \left| r_j \left(\frac{t}{N}\right) \right|. \quad (\text{D7})$$

Since the circuit cost can be expressed as

$$\text{Cost}(U) = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{t=1}^N \sum_{j=1}^m \left| r_j \left(\frac{t}{N}\right) \right|,$$

we have

$$\text{Cost}(U) \geq 4\sqrt{2}C_{W_1}(U). \quad (\text{D8})$$

□

Note that Lemma 19 also holds for $(n+m)$ -qudit systems, and hence the statement also holds for AC_{W_1} . Therefore, the following corollary follows immediately.

Corollary 21. Given a unitary U , the circuit cost $\text{Cost}(U)$ is lower bounded in terms of $\text{AC}_{W_1}(U)$ as follows:

$$\text{Cost}(U) \geq 4\sqrt{2}\text{AC}_{W_1}(U). \quad (\text{D9})$$

Theorem 22. *The experimental cost \mathcal{R}_U of an n -qubit quantum circuit $U = \prod_l U_l$ is lower bounded in terms of $C_{W_1}(U)$ as follows:*

$$\mathcal{R}(U) \geq \frac{1}{2} C_{W_1}(U). \quad (\text{D10})$$

Proof. For any pure n -qubit quantum state $|\psi\rangle$, let us define

$$|\psi_{l+1}\rangle := U_l |\psi_l\rangle, \quad \text{with } |\psi_1\rangle = |\psi\rangle. \quad (\text{D11})$$

And we also define

$$|\psi_{l,t_l}\rangle := e^{iH_l t_l} |\psi_l\rangle, \quad (\text{D12})$$

as the intermediate state at time $t_l \in [0, T_l]$ while implementing U_l with $|\psi_{l,0}\rangle = |\psi_l\rangle$, $\psi_{l,T_l} = |\psi_{l+1}\rangle$.

Since U_l acts on only k_l qubits, by Lemma 12, we have

$$\| |\psi_{l+1}\rangle\langle\psi_{l+1}| - |\psi_l\rangle\langle\psi_l| \|_{W_1} \leq k_l \| |\psi_{l+1}\rangle\langle\psi_{l+1}| - |\psi_l\rangle\langle\psi_l| \|_1. \quad (\text{D13})$$

Besides, it was shown in [70] that the quantum speed limit for the trace norm is bounded by the trace norm of the derivative of the state, that is,

$$\| |\psi_{l+1}\rangle\langle\psi_{l+1}| - |\psi_l\rangle\langle\psi_l| \|_1 \leq \int_0^{T_l} \| |d|\psi_{l,t_l}\rangle\langle\psi_{l,t_l}|/dt_l \|_1 dt_l, \quad (\text{D14})$$

where $d|\psi_{l,t_l}\rangle\langle\psi_{l,t_l}|/dt_l = d e^{iH_l t_l} |\psi_l\rangle\langle\psi_l| e^{-iH_l t_l} / dt_l = i[H_l, |\psi_{l,t_l}\rangle\langle\psi_{l,t_l}|]$. The trace norm of the derivative of the state under unitary evolution $e^{iH_l t_l}$ is bounded above by the square root of the variance of the Hamiltonian H_l , i.e.,

$$\| |d|\psi_{l,t_l}\rangle\langle\psi_{l,t_l}|/dt_l \|_1 \leq 2\sqrt{\text{Var}_{\psi_{l,t_l}}[H_l]}, \quad (\text{D15})$$

where

$$\text{Var}_{\psi_{l,t_l}}[H_l] = \text{Tr}[H_l^2 |\psi_{l,t_l}\rangle\langle\psi_{l,t_l}|] - \text{Tr}[H_l |\psi_{l,t_l}\rangle\langle\psi_{l,t_l}|]^2. \quad (\text{D16})$$

Note that $\text{Var}_{\psi_l}[H_l] \leq E_l^2$ with $E_l = (h_{2^{k_l}} - h_1)/2$, and so we have

$$\| |\psi_{l+1}\rangle\langle\psi_{l+1}| - |\psi_l\rangle\langle\psi_l| \|_1 \leq 2E_l T_l. \quad (\text{D17})$$

Hence,

$$\| |\psi_{l+1}\rangle\langle\psi_{l+1}| - |\psi_l\rangle\langle\psi_l| \|_{W_1} \leq 2k_l E_l T_l. \quad (\text{D18})$$

Therefore, for the quantum circuit $U = \prod_l U_l$ with any input state $|\psi\rangle$,

$$\begin{aligned} \| |\psi\rangle\langle\psi| - U |\psi\rangle\langle\psi| U^\dagger \|_{W_1} &\leq \sum_l \| |\psi_{l+1}\rangle\langle\psi_{l+1}| - |\psi_l\rangle\langle\psi_l| \|_{W_1} \\ &\leq \sum_l k_l \| |\psi_{l+1}\rangle\langle\psi_{l+1}| - |\psi_l\rangle\langle\psi_l| \|_1 \\ &\leq 2 \sum_l k_l E_l T_l = 2\mathcal{R}(U), \end{aligned}$$

where the first inequality comes from the triangle inequality, the second inequality comes from (D13), and the third inequality comes from (D17). \square

The proof given for the above theorem also works for the case where we have m qubits as ancillas. Hence, the following corollary follows immediately.

Corollary 23. *The experimental cost of an n -qubit quantum circuit $\mathcal{R}(U)$ is lower bounded in terms of $\text{AC}_{W_1}(U)$ as follows:*

$$\mathcal{R}(U) \geq \frac{1}{2} \text{AC}_{W_1}(U). \quad (\text{D19})$$