

Probing R -parity violation in B -meson decays to a baryon and a light neutralino

Claudio O. Dib,^a Juan Carlos Helo,^{b,c} Valery E. Lyubovitskij,^{d,a,c} Nicolás A. Neill,^f
Abner Soffer,^g Zeren Simon Wang^{h,i}

^a*Departamento de Física and CCTVal, Universidad Técnica Federico Santa María, Valparaíso 2340000, Chile*

^b*Departamento de Física, Facultad de Ciencias, Universidad de La Serena, Avenida Cisternas 1200, La Serena, Chile*

^c*Millennium Institute for Subatomic Physics at the High Energy Frontier (SAPHIR), Fernández Concha 700, Santiago, Chile*

^d*Institut für Theoretische Physik, Universität Tübingen, Kepler Center for Astro and Particle Physics, Auf der Morgenstelle 14, D-72076 Tübingen, Germany*

^f*Departamento de Ingeniería Eléctrica-Electrónica, Universidad de Tarapacá, Arica 1010069, Chile*

^g*School of Physics and Astronomy, Tel Aviv University, Tel Aviv 69978, Israel*

^h*Department of Physics, National Tsing Hua University, Hsinchu 300, Taiwan*

ⁱ*Center for Theory and Computation, National Tsing Hua University, Hsinchu 300, Taiwan*

E-mail: claudio.dib@usm.cl, jchelo@userena.cl, valeri.lyubovitskij@uni-tuebingen.de, naneill@outlook.com, asoffer@tau.ac.il, wzs@mx.nthu.edu.tw

ABSTRACT: We propose a search for B meson decays to a proton or a hyperon plus missing energy at the Belle II experiment to probe supersymmetry with a GeV-scale lightest neutralino $\tilde{\chi}_1^0$ and R -parity violation (RPV). We perform analytic computations of the signal branching fractions in the framework of effective field theory, with a single nonzero RPV operator $\lambda''_{1j3} \bar{U}_1^c \bar{D}_j^c \bar{D}_3^c$, where $j = 1, 2$. The hadronic form factors are calculated using an SU(3) phenomenological Lagrangian approach. Since the decay of the neutralino is kinematically and CKM suppressed in this theoretical scenario, it decays outside the detector and appears experimentally only as missing energy. We detail the analysis techniques at the experimental level and estimate the background in the $B^+ \rightarrow p \tilde{\chi}_1^0$ search using published results for $B^+ \rightarrow K^+ \nu \bar{\nu}$. Our final sensitivity plots are shown for both λ''_{113} versus the squark mass $m_{\tilde{q}}$ and $\lambda''_{113}/m_{\tilde{q}}^2$ versus the neutralino mass $m_{\tilde{\chi}_1^0}$. We find that the search at Belle II could probe $\lambda''_{113}/m_{\tilde{q}}^2$ down to the order of 10^{-8} GeV^{-2} in the kinematically allowed $m_{\tilde{\chi}_1^0}$ range. We also obtain current limits on $B^0 \rightarrow \Lambda^0 \tilde{\chi}_1^0$ by recasting an existing search, and comment about searches for $B^+ \rightarrow \Sigma^+ \tilde{\chi}_1^0$ and $B^0 \rightarrow \Sigma^0 \tilde{\chi}_1^0$. In closing, we briefly discuss potential searches at the LHCb and BESIII experiments.

Contents

1	Introduction	1
2	Theoretical framework: effective Lagrangian and neutralino production and decay	3
3	Experimental sensitivity estimation	7
3.1	Analysis technique	7
3.2	Background and sensitivity estimate for $\text{Br}(B^+ \rightarrow p\tilde{\chi}_1^0)$	9
3.3	Sensitivity estimate for $\text{Br}(B^0 \rightarrow \Lambda^0\tilde{\chi}_1^0)$, $\text{Br}(B^+ \rightarrow \Sigma^+\tilde{\chi}_1^0)$, $\text{Br}(B^0 \rightarrow \Sigma^0\tilde{\chi}_1^0)$	10
4	Results for the RPV parameters	11
5	Discussion and Conclusions	12
A	Transition Form Factors	14

1 Introduction

Supersymmetry (SUSY) [1, 2] has been one of the most sought candidates for physics beyond the Standard Model, in particular because it offers an elegant solution to the hierarchy problem [3, 4] in the Higgs sector of the Standard Model (SM). Although no evidence of SUSY has been found so far, the LHC explored wide regions of the SUSY parameter space and has established tight mass bounds on TeV-scale SUSY. The minimal supersymmetric extension of the Standard Model (MSSM) in its general form contains lepton-number violating (LNV) and baryon-number violating (BNV) terms, which can induce undesirable proton decay. One can impose by hand a simple Z_2 symmetry, called R -parity, which forbids these terms. Under R -parity, all SM particles are even, while all their superpartners are odd. If R -parity is conserved, not only is the proton stable but so is also the lightest supersymmetric particle (LSP). Nevertheless, in SUSY with R -parity violation (RPV), one can still prevent proton decay by imposing certain conditions¹. RPV-SUSY models are thus acceptable and, moreover, provide a rich phenomenology that is still only loosely constrained by colliders (see e.g. Refs. [6–8] for reviews of these models).

The LHC has established strong lower bounds on the masses of heavy SUSY particles, especially color-charged superpartners [9–13]. Color-neutral superpartners, however, are not so strictly bounded. The lightest neutralino $\tilde{\chi}_1^0$ can be as light as a GeV or even lighter [14–23], in which case its production in decays of τ lepton or heavy mesons is

¹See Ref. [5] for a study on the updated bounds on RPV couplings from nucleon decays.

kinematically allowed. Such a light neutralino should be dominantly Bino according to current bounds [21–25]. In addition, it has to decay, e.g., via RPV couplings so as to avoid overclosing the Universe [26–29]. If the RPV couplings are tiny, such light neutralinos can be long-lived so that they appear as displaced signatures or even missing energy in collider experiments. In this context, in a previous work [30] we proposed searching for such a neutralino in τ lepton decays at Belle II. Refs. [31–34] have estimated the sensitivities of various LHC far detectors, and Ref. [35] investigated the sensitivity of neutrino detectors to atmospheric light neutralinos.

In this work we consider a light neutralino produced in B -meson decays through the superpotential RPV terms $\lambda''_{ijk}\bar{U}_i\bar{D}_j\bar{D}_k$, where U and D are the up-type and down-type right-handed supermultiplet fields, λ''_{ijk} are dimensionless couplings, and i, j, k are generation indices in the quark-mass basis. Such interaction terms not only violate R -parity but also baryon number. However, proton decay does not occur in this scenario, as long as no LNV RPV terms are allowed and the neutralino is not lighter than the proton. We focus on the case in which only λ''_{113} or λ''_{123} is non-zero and there is no squark mixing. The decay of such a light neutralino is suppressed, so that it appears as missing energy at the detector level [25]. We propose to probe λ''_{113} via the decay mode $B^+ \rightarrow p\tilde{\chi}_1^0$, and to study λ''_{123} via the decays $B^0 \rightarrow \Lambda^0\tilde{\chi}_1^0$, $B^+ \rightarrow \Sigma^+\tilde{\chi}_1^0$, and $B^0 \rightarrow \Sigma^0\tilde{\chi}_1^0$. We generically refer to all four modes as $B \rightarrow \mathcal{B}\tilde{\chi}_1^0$, where \mathcal{B} indicates one of the above four baryons. These decays can be searched for at an $e^+e^- \rightarrow B\bar{B}$ “ B -factory” facility, namely, the currently running Belle II experiment [36, 37] or the BABAR and Belle experiments, which are no longer collecting data. While these decays violate baryon number, strictly speaking it is not possible to verify experimentally that the neutralino, which escapes detection, does not carry baryon number.

The superpotential terms $\lambda''_{ijk}\bar{U}_i\bar{D}_j\bar{D}_k$ can also contribute to baryon-antibaryon oscillations ($n - \bar{n}$) [38, 39] and di-nucleon decay (e.g., $NN \rightarrow \pi\pi$, $NN \rightarrow KK$) [40]. However, in the scenario studied here, with no squark mixing, the contributions of λ''_{113} and λ''_{123} to these processes are suppressed by two weak insertions, yielding limits on λ''_{113} and λ''_{123} that are much weaker than current LHC bounds and weaker still than the sensitivity of the studies we propose here. Following Eq. (23) in Ref. [40], we can estimate the order of magnitude of the baryon anti-baryon transition amplitude $\delta_{B\bar{B}}$ in terms of the RPV couplings:

$$\delta_{B\bar{B}} \sim \frac{\kappa^2 m_{\tilde{\chi}_1^0}}{m_B^2 - m_{\tilde{\chi}_1^0}^2} \left(\frac{\lambda''_{ijk} g_{1R}^{\tilde{q}}}{m_{\tilde{q}}^2} \right)^2, \quad (1.1)$$

where $\kappa \sim 10^{-2} \text{ GeV}^3$ [41]. From the LHCb bound on the $\Xi_b^0 - \bar{\Xi}_b^0$ oscillation rate, namely $\omega < 0.08 \text{ ps}^{-1}$, we extract the bound $\lambda''_{123}/m_{\tilde{q}}^2 \lesssim 4 \times 10^{-4} \text{ GeV}^{-2}$, where we have taken $m_{\tilde{\chi}_1^0} = 2.5 \text{ GeV}$ in Eq. (1.1). Similarly, the bounds on $\delta_{B\bar{B}}$ from dinucleon decay listed in Table I of Ref. [40] can be translated into bounds on the RPV couplings. The bounds $\delta_{(udb)^2} \lesssim 10^{-13}$ and $\delta_{(usb)^2} \lesssim 10^{-10}$ translate into $\lambda''_{113}/m_{\tilde{q}}^2 \lesssim 6 \times 10^{-4} \text{ GeV}^{-2}$ and $\lambda''_{123}/m_{\tilde{q}}^2 \lesssim 2 \times 10^{-2} \text{ GeV}^{-2}$, respectively, again taking $m_{\tilde{\chi}_1^0} = 2.5 \text{ GeV}$ in Eq. (1.1).

Decays of a B meson to a baryon and missing energy are motivated also in the context of B -mesogenesis models [42, 43]. Signatures of directly detectable or escaping long-lived

particles at B factories have also been discussed in Refs. [30, 44–55]

This paper is organized as follows: section 2 contains the theoretical framework of the RPV scenario. This includes the effective Lagrangian with the RPV terms and the calculation of the matrix elements, which we estimate using an effective BNV Lagrangian. In section 3 we explain the experimental analysis technique, estimate the background in the search for $B^+ \rightarrow p\tilde{\chi}_1^0$, and determine the sensitivity of a Belle II measurement of the branching fraction $\text{Br}(B^+ \rightarrow p\tilde{\chi}_1^0)$. We also interpret preliminary BABAR results [56] in terms of $B^0 \rightarrow \Lambda^0\tilde{\chi}_1^0$ and compare the sensitivity of this mode to those of $B^0 \rightarrow \Sigma^0\tilde{\chi}_1^0$ and $B^+ \rightarrow \Sigma^+\tilde{\chi}_1^0$. Our main physics results are provided in section 4, where we extract current limits on λ''_{123} and estimate the Belle II sensitivity to λ''_{113} . Finally, section 5 contains our summary, as well as a brief discussion of related measurements that we propose to perform at the LHCb and BESIII experiments. We have included an Appendix with details of the form factor calculation for the hadronic transitions.

2 Theoretical framework: effective Lagrangian and neutralino production and decay

In this section we describe our theoretical framework. We start with an effective Lagrangian needed for the description of $B \rightarrow \mathcal{B}\tilde{\chi}_1^0$ decays at the partonic level. Then we define the effective four-fermion operators that generate the matrix elements for the transition hadronic form factors. Finally, we discuss the computation of the decay rates $\Gamma(B \rightarrow \mathcal{B}\tilde{\chi}_1^0)$ and present the contributions arising from the different intermediate squarks as a function of the neutralino mass.

The effective Lagrangian \mathcal{L}_{eff} for the description of $B \rightarrow \mathcal{B}\tilde{\chi}_1^0$ at the partonic level contains two parts: an R -parity violating part, \mathcal{L}_{RPV} , with trilinear couplings of two quarks and one squark, and an R -parity conserving (RPC) part describing the couplings of the neutralino $\tilde{\chi}_1^0$ with a quark and an squark [57–60]:

$$\begin{aligned} \mathcal{L}_{\text{eff}} &= \mathcal{L}_{\text{RPV}} + \mathcal{L}_{\text{RPC}}, \\ \mathcal{L}_{\text{RPV}} &= \lambda''_d \epsilon_{abc} \left(\tilde{u}_{R,a}^* \bar{d}_{R,b} b_{R,c}^C + \tilde{d}_{R,a}^* \bar{u}_{R,b} b_{R,c}^C + \tilde{b}_{R,a}^* \bar{u}_{R,b} d_{R,c}^C \right) + (d \rightarrow s) \\ &\quad + \text{h.c.} + \dots, \\ \mathcal{L}_{\text{RPC}} &= - \sum_{q=u,d,s,b} g_{1R}^{\tilde{q}} (\bar{q}_{R,a} P_L \tilde{\chi}_1^0) \tilde{q}_{R,a} + \text{h.c.} + \dots \end{aligned} \quad (2.1)$$

Here $(d \rightarrow s)$ indicates replacement of the d quark by the s quark, $\lambda''_d = \lambda''_{113}$ and $\lambda''_s = \lambda''_{123}$ are the RPV couplings, a, b, c are color indices, ϵ_{abc} is the color three-dimensional Levi-Civita tensor, $P_L = (1 - \gamma^5)/2$ is the chiral projector, and the upper index C denotes charge conjugation. Since we take $\tilde{\chi}_1^0$ to be a pure Bino, the coupling $g_{1R}^{\tilde{q}}$ is given by

$$g_{1R}^{\tilde{q}} = -\sqrt{2} g_W e_q \tan \theta_W, \quad (2.2)$$

where θ_W is the Weinberg angle with $\tan \theta_W \simeq 0.54840$, $g_W = e/\sin \theta_W \simeq 0.62977$ is the SU(2) weak coupling, and e_q is the quark electric charge, i.e. $e_u = 2/3$ and $e_d = e_s = e_b = -1/3$. In the last line of Eq. (2.1) we explicitly write only the terms with right-chiral fields,

since we consider vanishing squark mixing, and only such fields are involved in the RPV interactions considered here.




Figure 1. Parton-level diagrams for the decays $B \rightarrow \mathcal{B}\tilde{\chi}_1^0$ (where \mathcal{B} is one of the baryons $p, \Lambda, \Sigma^+, \Sigma^0$) through the RPV couplings λ''_{113} and λ''_{123} , respectively.

The Lagrangian of Eq. (2.1) generates the parton-level diagrams shown in Fig. 1 for the decays $B \rightarrow \mathcal{B}\tilde{\chi}_1^0$, as long as the decays are kinematically allowed². To simplify the discussion, we first focus on the $B^+ \rightarrow p\tilde{\chi}_1^0$ process, i.e., the diagrams that have a d quark in the final state.

For simplicity, we assume the squark masses to be degenerate and no squark mixing. In the limit that the squark mass $m_{\tilde{q}}$ is large, the squark propagators are approximately $1/m_{\tilde{q}}^2$. Therefore, the d -quark diagrams in Fig. 1 correspond to the four-fermion effective Lagrangian

$$\mathcal{L}_d^{BNV} = \mathcal{L}_d^{bud\tilde{\chi}_1^0} + \mathcal{L}_d^{udb\tilde{\chi}_1^0} + \mathcal{L}_d^{dub\tilde{\chi}_1^0}, \quad (2.3)$$

where

$$\mathcal{L}_d^{q_1q_2q_3\tilde{\chi}_1^0} = \mathcal{O}_d^{q_1q_2q_3} \tilde{\chi}_1^0 + \text{h.c.}, \quad (2.4)$$

the effective operators are defined as

$$\mathcal{O}_d^{q_1q_2q_3} = g_d^{\tilde{q}_1R} \mathcal{O}_{q_1q_2q_3}^{LL}, \quad (2.5)$$

and we have used the definitions

$$g_d^{\tilde{q}R} = \frac{g_{1R}^{\tilde{q}} \lambda_d''}{m_{\tilde{q}}^2}, \quad \mathcal{O}_{q_1q_2q_3}^{LL} = \varepsilon_{abc} (\bar{q}_{3,c} P_L C \bar{q}_{2,b}) \bar{q}_{1,a} P_L, \quad (2.6)$$

where $C = i\gamma^0\gamma^2$ is the charge conjugation matrix.

Following Refs. [61–65], we define the invariant matrix element for the $B^+ \rightarrow p\tilde{\chi}_1^0$ decay as

$$\mathcal{M} = \bar{u}_p(p', s') \left[W_0^{LL}(q^2) + \frac{\not{q}}{m_{\tilde{\chi}_1^0}} W_1^{LL}(q^2) \right] P_L v_{\tilde{\chi}_1^0}(q, s), \quad (2.7)$$

where $\bar{u}_p(p', s')$ and $v_{\tilde{\chi}_1^0}(q, s)$ are the spinors of the final-state proton and neutralino, respectively; p, p' , and $q = p - p'$ are the momenta of B^+ meson, proton, and neutralino,

²While the decay $B^0 \rightarrow n\tilde{\chi}_1^0$ can also take place, we do not consider it due to the experimental difficulty involved with neutron reconstruction.

respectively; m_B , m_p , and $m_{\tilde{\chi}_1^0}$ are the masses of these particles; and $W_0^{LL}(q^2)$ and $W_1^{LL}(q^2)$ are form factors that parametrize the hadronic matrix elements of the BNV three-quark operator

$$\mathcal{O}^{LL} = g_d^{\tilde{b}R} \mathcal{O}_{bud}^{LL} + g_d^{\tilde{u}R} \mathcal{O}_{udb}^{LL} + g_d^{\tilde{d}R} \mathcal{O}_{dub}^{LL} \quad (2.8)$$

between the proton and B^+ meson states:

$$\langle p | \mathcal{O}^{LL} | B^+ \rangle = \bar{u}_p(p', s') \left[W_0^{LL}(q^2) + \frac{\not{q}}{m_{\tilde{\chi}_1^0}} W_1^{LL}(q^2) \right] P_L. \quad (2.9)$$

The form factors $W_0^{LL}(q^2)$ and $W_1^{LL}(q^2)$ can be calculated using an SU(3) phenomenological Lagrangian (see Refs. [66] and [61]-[65]) applied to the BNV matrix elements involving a proton and light pseudoscalar mesons (π, K, η) and extending it to the bottom sector. The contribution to the form factors of each of the \tilde{b} , \tilde{u} , and \tilde{d} squarks is the sum of a *direct* contribution from the $B^+ \rightarrow p \tilde{\chi}_1^0$ process and a *pole* contribution induced by the bottom baryon resonances Λ_b^0 and Σ_b^0 , leading to the virtual two-step process $B^+ \rightarrow p + [\Lambda_b^0 / \Sigma_b^0 \rightarrow \tilde{\chi}_1^0]$.

Here, we estimate our matrix elements by extrapolating the strange flavor in $\langle p | \mathcal{O}^{LL} | K^+ \rangle$ to the bottom flavor. In Refs. [67–71] detailed discussion of possibilities to extend the phenomenological Lagrangians from the SU(3) sector to the heavy quark sector have been provided. Derived effective Lagrangians involving light and heavy mesons and baryons have been successfully applied to different tasks. More concretely, one replaces the decay constant of the Kaon by that of the B meson, $f_K \rightarrow f_B$, and the masses of the K^+ meson and the Λ^0 and Σ^0 hyperons by the corresponding masses of B^+ , Λ_b^0 and Σ_b^0 . Our analytical results for the corresponding form factors are shown in Appendix A. We also demonstrate in the appendix that the same results for the transition form factors can be obtained using an effective phenomenological BNV Lagrangian constructed in terms of the fields of the B^+ , proton, Λ_b^0 , and Σ_b^0 , which contains some adjustable couplings fixed from matching of matrix elements $\langle p | \mathcal{O}^{LX} | B^+ \rangle$ in the two approaches.

The $B^+ \rightarrow p \tilde{\chi}_1^0$ decay width is calculated according to the formula

$$\begin{aligned} \Gamma(B^+ \rightarrow p \tilde{\chi}_1^0) &= \frac{\lambda^{1/2}(m_B^2, m_p^2, m_{\tilde{\chi}_1^0}^2)}{16\pi m_B^3} \sum_{\text{pol.}} |\mathcal{M}|^2, \\ \sum_{\text{pol.}} |\mathcal{M}|^2 &= \left(m_B^2 - m_p^2 - m_{\tilde{\chi}_1^0}^2 \right) (\mathcal{A} + \mathcal{B})^2 - 2 \left(m_B^2 - (m_p + m_{\tilde{\chi}_1^0})^2 \right) \mathcal{A} \cdot \mathcal{B} \\ &= \left(m_B^2 - (m_p + m_{\tilde{\chi}_1^0})^2 \right) (\mathcal{A}^2 + \mathcal{B}^2) + 2m_p m_{\tilde{\chi}_1^0} (\mathcal{A} + \mathcal{B})^2, \end{aligned} \quad (2.10)$$

where

$$\mathcal{A} = W_0^{LL}(m_{\tilde{\chi}_1^0}^2), \quad \mathcal{B} = W_1^{LL}(m_{\tilde{\chi}_1^0}^2), \quad (2.11)$$

and

$$\lambda(x, y, z) = x^2 + y^2 + z^2 - 2xy - 2xz - 2yz \quad (2.12)$$

is the kinematical triangle Källén function. The last line in Eq. (2.10) shows that the matrix element squared $\sum_{pol} |\mathcal{M}|^2$ is manifestly positive because of the kinematical constraint $m_B > m_p + m_{\tilde{\chi}_1^0}$.

We estimate the rate of the decay $B^+ \rightarrow \Sigma^+ \tilde{\chi}_1^0$ in the same way, replacing the d quark by the s quark. Specifically: $\lambda_d'' \rightarrow \lambda_s''$, $p(duu) \rightarrow \Sigma^+(suu)$, and the intermediate baryons Λ_b^0 and Σ_b^0 are replaced by the Ξ_b^0 and $\Xi_b'^0$, which have a (bus) quark-flavor content with an antisymmetric light spin-0 diquark $[us]_0$ and a symmetric light spin-1 diquark $\{us\}_1$, respectively. We also replace the couplings α and β that define the matrix elements of the three-quark operators between the proton and vacuum state, by the respective couplings that define the matrix elements for the case of the Σ^+ baryon. See the discussion in Appendix A for more details, as well as for the form-factor calculation for $B^0 \rightarrow \Lambda^0 \tilde{\chi}_1^0$ and $B^0 \rightarrow \Sigma^0 \tilde{\chi}_1^0$.




Figure 2. The branching fractions of $B^+ \rightarrow p \tilde{\chi}_1^0$, $B^+ \rightarrow \Sigma^+ \tilde{\chi}_1^0$, $B^0 \rightarrow \Sigma^0 \tilde{\chi}_1^0$, and $B^0 \rightarrow \Lambda^0 \tilde{\chi}_1^0$ normalized by $G^2 = \lambda_{d/s}''^2 \times (1 \text{ TeV}/\tilde{m}_q)^4$, as functions of the neutralino mass.

In Fig. 2 we compare the four branching fractions of interest, normalized by $G^2 = \lambda_{d/s}''^2 \times (1 \text{ TeV}/\tilde{m}_q)^4$, as functions of $m_{\tilde{\chi}_1^0}$. One can see that $\text{Br}(B^0 \rightarrow \Lambda^0 \tilde{\chi}_1^0)$ is significantly suppressed relative to $\text{Br}(B^+ \rightarrow p \tilde{\chi}_1^0)$. This is explained by the compensation of the relative \tilde{b}^* , \tilde{u} , and \tilde{s} contributions in $\text{Br}(B^0 \rightarrow \Lambda^0 \tilde{\chi}_1^0)$, whose magnitudes are shown in Fig. 3. In particular, at small values of $m_{\tilde{\chi}_1^0}$, the ratio $R_1 = \text{Br}(B^0 \rightarrow \Lambda^0 \tilde{\chi}_1^0)/\text{Br}(B^+ \rightarrow p \tilde{\chi}_1^0)$ is dominated by the direct contributions to W_0^{LL} (see Eq. (A.17)), and is given by

$$R_1 \simeq \frac{3}{8} \left[\frac{e_u + e_d - 2e_s}{e_u - e_s} \right]^2 = \frac{1}{24}. \quad (2.13)$$

In the case of the Σ production modes, the corresponding ratio $R_2 = \text{Br}(B^0 \rightarrow \Sigma^0 \tilde{\chi}_1^0)/\text{Br}(B^+ \rightarrow \Sigma^+ \tilde{\chi}_1^0) \simeq 1/2$ is explained by the isotopic flavor factor $1/\sqrt{2}$.

Before moving on to the experimental aspects of the proposed search, we briefly discuss the decay of the lightest neutralino. For the considered RPV couplings, the neutralino mass range of interest, and the absence of squark mixing, the neutralino decay is kinematically suppressed by 3 off-shell propagators: a squark, a bottom quark, and a W boson, and is also CKM suppressed. This renders the lifetime of the lightest neutralino so long that it appears only as missing energy in the detector.




Figure 3. Individual squark contributions to the branching fractions shown in Fig. 2.

3 Experimental sensitivity estimation

The proposed search for $B^+ \rightarrow p\tilde{\chi}_1^0$ is experimentally very similar to those of $B^+ \rightarrow K^+\nu\bar{\nu}$ and $B^+ \rightarrow \pi^+\nu\bar{\nu}$. In all these cases, the signal involves a B decay to a single charged particle and unobserved, “missing” particles. We describe the basic analysis techniques in Sec. 3.1 and use previous searches for $B^+ \rightarrow K^+\nu\bar{\nu}$ to estimate the sensitivity for $B^+ \rightarrow p\tilde{\chi}_1^0$ in Sec. 3.2. In Sec. 3.3 we discuss the relative sensitivities of the modes $B^0 \rightarrow \Lambda^0\tilde{\chi}_1^0$, $B^+ \rightarrow \Sigma^+\tilde{\chi}_1^0$, and $B^0 \rightarrow \Sigma^0\tilde{\chi}_1^0$.

3.1 Analysis technique

Searches for $B^+ \rightarrow K^+\nu\bar{\nu}$ and $B^+ \rightarrow \pi^+\nu\bar{\nu}$ have been published by the CLEO [72], BABAR [73–75], Belle [76–78], and Belle II [79] collaborations. Using $e^+e^- \rightarrow B^+B^-$ data collected at the $\Upsilon(4S)$ resonance, these searches were performed by reconstructing, or “tagging”, the decays of one of the B mesons, referred to as the tag B . A single kaon is the only visible particle associated with the signal B candidate, which is assumed

to undergo the signal decay of interest. Tagging is performed via one of three methods: hadronic, semileptonic, or inclusive. Hadronic tagging has the highest signal purity and lowest efficiency, and the opposite is true for inclusive tagging. In what follows we briefly describe the three tagging methods for use in the decay $B^+ \rightarrow p\tilde{\chi}_1^0$.

In hadronic tagging, one attempts to reconstruct the tag B in up to thousands of decay modes involving the Cabibbo-favored quark-level processes $b \rightarrow c\bar{u}d$ and $b \rightarrow c\bar{c}s$. Background is suppressed with selections based on the variables $\Delta E = E_B - \sqrt{s}/2$ and $M_{bc} = \sqrt{s/4 - p_B^2}$, where E_B and p_B are the measured energy and momentum of the tag B in the center-of-mass (CM) frame of the e^+e^- collision, and \sqrt{s} is the CM energy. Signal events peak at $\Delta E = 0$ and $M_{bc} = m_B$, while background events are more broadly distributed. Hadronic tagging provides also the squared invariant mass of the unobserved neutralino candidate, the so-called missing mass

$$m_{\text{miss}}^2 = (p_{ee} - p_{\text{tag}} - p_p)^2. \quad (3.1)$$

Here p_{ee} is the known 4-momentum of the beam particles [80], and p_{tag} and p_p are the measured 4-momenta of the tag B and of the proton, respectively. The presence of a signal would appear as a peak in the m_{miss} distribution centered at $m_{\text{miss}} = m_{\tilde{\chi}_1^0}$.

In semileptonic tagging, the tag B is reconstructed in $B^- \rightarrow D^{(*)}\ell^- \bar{\nu}$ decays, where ℓ is an electron or muon. Because of the unobserved neutrino, the values of ΔE and M_{bc} are not physically meaningful. Rather, one applies the constraint $E_B = \sqrt{s}/2$ to calculating the variable

$$\cos\theta_{B-D^{(*)}\ell} = \frac{E_{D^{(*)}\ell} s - M_B^2 - m_{D^{(*)}\ell}^2}{2p_{D^{(*)}\ell}\sqrt{s/4 - M_B^2}}, \quad (3.2)$$

where $E_{D^{(*)}\ell}$, $p_{D^{(*)}\ell}$, and $m_{D^{(*)}\ell}$ are the CM-frame energy, momentum, and invariant mass of the $D^{(*)}\ell^-$ system. For signal events reconstructed without measurement errors, $\cos\theta_{B-D^{(*)}\ell}$ is the cosine of the angle between the true momenta of the tag B and the $D^{(*)}\ell$ system. Therefore, signal events are distributed mostly in the range $-1 < \cos\theta_{B-D^{(*)}\ell} < 1$, with tails arising from the finite measurement resolution and the spread in \sqrt{s} , while background events reach greater negative values. The unobserved neutrino makes it impossible to determine p_{tag} . Nonetheless, the neutralino mass can be calculated with the assumption that the two B mesons are at rest in the CM frame, i.e., $\vec{p}_{\text{tag}} = 0$ in Eq. (3.2). The resulting m_{miss} signal peak is significantly broader than for hadronic tagging.

Inclusive tagging is similar to semileptonic tagging in that the tag B decay products are not all observed. This may arise from particles that are outside the angular or momentum acceptance of the detector, poor resolution and identification capability for K_L mesons and neutrons, particles with overlapping detector signatures, accelerator background, electronic noise, and limitations of the detector or the reconstruction algorithms. In Ref. [79], the variable used to discriminate between signal and background was the output of a multivariate algorithm, which was trained with simulated events. Previous uses of inclusive tagging, beginning with Ref. [81], used kinematic variables such as M_{bc} for background discrimination.

In all tagging methods, one would require that the event contain only one charged-particle track that is not associated with the tag B . For $B^+ \rightarrow p\tilde{\chi}_1^0$, this track should be identified as a proton using the detector’s hadron-identification capabilities. In addition, the energy deposition E_{extra} in calorimeter clusters that are not associated with the proton or the tag- B decay products must be small in order to suppress background from neutrons and K_L^0 mesons. As in $B^+ \rightarrow K^+\nu\bar{\nu}$ searches, this requirement can be applied directly on E_{extra} , or by including this variable as input to a multivariate algorithm.

In both $B^+ \rightarrow K^+\nu\bar{\nu}$ and other studies of rare B decays with final-state neutrinos [82] the three methods have similar sensitivities for a given data set. Therefore, all can be used for the search of $B^+ \rightarrow p\tilde{\chi}_1^0$, yielding similar sensitivities. In what follows, we use hadronic tagging for our sensitivity estimation.

3.2 Background and sensitivity estimate for $\text{Br}(B^+ \rightarrow p\tilde{\chi}_1^0)$

We estimate the m_{miss} -dependent background in the hadronic-tagging analysis from Fig. 8 of Ref. [75], which shows the m_{miss} distribution in the BABAR $B^+ \rightarrow K^+\nu\bar{\nu}$ search, performed with a data sample of 0.471×10^9 $B\bar{B}$ pairs. The background estimate is 0.2 events per 100-MeV-wide bin for $1 < m_{\text{miss}} < 2.5$ GeV, thereby rising linearly to 3.5 events per bin at $m_{\text{miss}} = 4.2$ GeV. We scale the background estimate to the Belle II data set by multiplying it by the ratio $R_L = N_{BB}(\text{Belle II})/N_{BB}(\text{BABAR}) = 55 \times 10^9/0.471 \times 10^9$ of the numbers of $e^+e^- \rightarrow B\bar{B}$ events in the final Belle II data set and in Ref. [75], respectively.

The estimated background is reduced relative to that in $B^+ \rightarrow K^+\nu\bar{\nu}$ by the fact that the proton production rate in B^+ decays is about $R_{p/K} \approx 1/16$ of K^+ production. Specifically, $\text{Br}(B^+ \rightarrow K^+ + \text{anything}) = (66 \pm 5)\%$ [83]. While this was measured by ARGUS [84] and CLEO [85] for a nearly equal admixture of B^+ and B^0 , it is not expected to depend significantly on the isospin state of the B meson. Therefore, we take this branching fraction to apply to the B^+ . By contrast, $\text{Br}(B \rightarrow \bar{p}/p + \text{anything}) = (8.0 \pm 0.4)\%$ [83], which is about 8 times smaller. Furthermore, correlating the charge of the proton with the charge of the tag B will reduce the background by a factor of two, given that such a correlation was not performed in the $\text{Br}(B \rightarrow \bar{p}/p + \text{anything})$ measurement. Baryon production is expected to be similarly suppressed in the non- $B\bar{B}$ background, specifically $e^+e^- \rightarrow q\bar{q}$, where $q = u, d, s, c$ is a light quark. Furthermore, this background is reportedly smaller than the $B\bar{B}$ background in the $B^+ \rightarrow K^+\nu\bar{\nu}$ searches [46, 78, 79]. Thus, multiplication by $R_{p/K}$ gives the expected background yield N_b^{100} for the $B^+ \rightarrow p\tilde{\chi}_1^0$ search at Belle II in each 100-MeV-wide m_{miss} bin. From this, we calculate the number of background events in a m_{miss} bin with a width of 2-standard-deviation,

$$N_b = N_b^{100} \frac{2\sigma(m_{\text{miss}})}{100 \text{ MeV}}. \quad (3.3)$$

The m_{miss} resolution $\sigma(m_{\text{miss}})$ is dominated by the resolution on the transverse momentum p_T of the proton for most of the m_{miss} range. For the Belle experiment, this has been measured to be [80]

$$\left(\frac{\sigma(p_T)}{p_T}\right)^2 = \left(0.0019 \frac{p_T}{\text{GeV}}\right)^2 + \left(0.003 \frac{1}{\beta}\right)^2, \quad (3.4)$$

where β is the charged-particle velocity, which we calculate on average assuming a uniform angular distribution for the protons. We ignore the fact that the p_T resolution is expected to be somewhat better for Belle II, which has a larger drift chamber. A smaller contribution to $\sigma(m_{\text{miss}})$ arises from the spread of the collider center-of-mass energy, which we take to be 5 MeV. The value of $\sigma(m_{\text{miss}})$ as a function of $m_{\tilde{\chi}_1^0}$ is shown in Fig. 4.




Figure 4. The missing-mass resolution $\sigma(m_{\text{miss}})$ (left) and the expected Belle II 90% confidence-level upper limit on the branching fraction $\text{Br}(B^+ \rightarrow p\tilde{\chi}_1^0)$ as functions of the neutralino mass $m_{\tilde{\chi}_1^0}$. The upper limit decreases with the decreasing value of $\sigma(m_{\text{miss}})$ but increases for $m_{\tilde{\chi}_1^0} > 2.5$ GeV due to the increasing background. The kink at 2.5 GeV results from the simple parameterization of the background governed by the limited data available in Ref. [75].

The 90% confidence-level (CL) upper limit on the $B^+ \rightarrow p\tilde{\chi}_1^0$ branching fraction at Belle II, assuming the absence of a signal, is calculated from

$$\text{Br}(B^+ \rightarrow p\tilde{\chi}_1^0) < \frac{1.64\sqrt{N_b}}{N_{BB}(\text{Belle II}) \epsilon}, \quad (3.5)$$

where $\epsilon \approx 95 \times 10^{-5}$ is the total reconstruction efficiency [75].³ We check this branching-fraction sensitivity estimate against an inclusive-tagging study of $B^+ \rightarrow K^+a$, where a is an axion-like particle that is long lived enough to always escape the detector [46]. Fig. 10 of Ref. [46] gives the 90% CL limit $\text{Br}(B^+ \rightarrow K^+a) < 10^{-7}$. Scaling this by $\sqrt{R_{p/K}}$ yields $\text{Br}(B^+ \rightarrow p\tilde{\chi}_1^0) < 2.5 \times 10^{-8}$, in agreement with our hadronic-tagging-based estimate.

3.3 Sensitivity estimate for $\text{Br}(B^0 \rightarrow \Lambda^0\tilde{\chi}_1^0)$, $\text{Br}(B^+ \rightarrow \Sigma^+\tilde{\chi}_1^0)$, $\text{Br}(B^0 \rightarrow \Sigma^0\tilde{\chi}_1^0)$

The Belle collaboration has published a search for $B^0 \rightarrow \Lambda^0\psi_D$, where ψ_D is an invisible dark-matter particle [86], obtaining limits on the branching fraction in 500 MeV-wide steps of m_{miss} . A preliminary result from BABAR [56] shows tighter limits in the range 193 steps in $1 < m_{\text{miss}} < 4.2$ GeV. In Sec. 4 we reinterpret the BABAR search in terms of $B^0 \rightarrow \Lambda^0\tilde{\chi}_1^0$ and extract limits on λ''_{123} .

Considering the $B^0 \rightarrow \Sigma^0\tilde{\chi}_1^0$ signal, we note that the dominant decay mode of the Σ^0 is $\Sigma^0 \rightarrow \Lambda^0\gamma$. Since the $\Sigma^0 - \Lambda^0$ mass difference is only 77 MeV, the photon is soft, so its reconstruction incurs lower efficiency and higher background than for $B^0 \rightarrow \Lambda^0\tilde{\chi}_1^0$

³The efficiency is taken from Table VI of Ref. [75], where it is given multiplied by $\text{Br}(B^+ \rightarrow J/\psi K^+)$.

reconstruction. The efficiency loss may be avoided by foregoing the photon reconstruction altogether. However, this results in an additional contribution of order the photon energy to the m_{miss} resolution. Furthermore, while the production rate of Σ baryons in B -meson decays has not been measured, one can assume that it is similar to that of Λ^0 . As a result, background from true Σ baryons is not smaller than that arising from true Λ baryons. Overall, although $B^0 \rightarrow \Sigma^0 \tilde{\chi}_1^0$ appears theoretically more promising than $B^0 \rightarrow \Lambda^0 \tilde{\chi}_1^0$ in Fig. 2, these experimental difficulties mean that it does not offer an advantage over $B^0 \rightarrow \Lambda^0 \tilde{\chi}_1^0$.

In the search for $B^+ \rightarrow \Sigma^+ \tilde{\chi}_1^0$, one must reconstruct the Σ^+ in the final state $p\pi^0$, which has a branching fraction of 51.6% [83]. About 48.3% of the Σ^+ decays are to $n\pi^+$. However, since the neutron interacts only hadronically, the reconstruction of this mode is much less practical. The efficiency for reconstruction of the soft π^0 is in the low tens of percent [87]. This approximately offsets the advantage of the $B^+ \rightarrow \Sigma^+ \tilde{\chi}_1^0$ mode seen in Fig. 2.

4 Results for the RPV parameters

We proceed to calculate the sensitivity to the trilinear RPV couplings from the experimental sensitivity to the decay branching fractions of B mesons into a charged baryon and a light neutralino. As stated above, we take only one RPV coupling to be non-vanishing at a time, either λ''_{113} or λ''_{123} , and assume degenerate squark masses and no squark mixing.

In Fig. 5 we show the limits on $\lambda''_{123}/m_{\tilde{q}}^2$ vs. $m_{\tilde{\chi}_1^0}$ that we extract from the BABAR search for $B^0 \rightarrow \Lambda^0 \psi_D$ [56]. These limits are around $\lambda''_{123} < 1 \times 10^{-6}/\text{GeV}^{-2}$, with fluctuations that reflect the BABAR results. The BABAR search is effectively background-free in $m_{\tilde{\chi}_1^0}$ regions in which the m_{miss} difference between background events is significantly larger than the experimental m_{miss} resolution. With more than 100 times more data, the Belle II search will probably not have a background-free region, unless further analysis improvements are implemented. Therefore, the Belle II sensitivity to λ''_{123} is naively expected to be better than that of BABAR by about $1/\sqrt{R_L}$ for most regions of $m_{\tilde{\chi}_1^0}$. Related to this, as a result of the nonvanishing background, the Belle II sensitivity is expected to become poorer as the m_{miss} resolution decreases for low values of $m_{\tilde{\chi}_1^0}$.

Fig. 6 shows the expected sensitivity of the search for $B^+ \rightarrow p\tilde{\chi}_1^0$ at Belle II with a sample of $N_{BB} = 55 \times 10^9 B\bar{B}$ pairs. The sensitivity is shown in the λ''_{113} vs. $m_{\tilde{q}}$ plane for the neutralino-mass values $m_{\tilde{\chi}_1^0} = (1, 2.4, 4)$ GeV, and corresponds to 90% confidence-level upper limits. Fig. 6 also compares our limits with current squark-mass limits, $m_{\tilde{q}} \gtrsim 1.85$ TeV for 8 mass-degenerate first- and second-generation squarks and $m_{\tilde{q}} \gtrsim 1.3$ TeV for a single squark belonging to the first or second generation. These limits were obtained by the ATLAS collaboration from a search for $pp \rightarrow \tilde{q}\tilde{q}, \tilde{q} \rightarrow q\tilde{\chi}_1^0$ in a signature of jets plus missing transverse momentum in a data sample with an integrated luminosity of 139 fb^{-1} [88]. The squark mass limits assume $\text{Br}(\tilde{q} \rightarrow q\tilde{\chi}_1^0) = 1$. In our theoretical scenario this is a good approximation only for small values of λ''_{113} , for which the competing decay channel $\tilde{q} \rightarrow qq$ is suppressed. As λ''_{113} increases, the ATLAS $m_{\tilde{q}}$ limits weaken relative to the values shown in Fig. 6. As mentioned in Sec. 1, current bounds on λ''_{113} and λ''_{123} arising from di-nucleon




Figure 5. Limits on λ''_{123} as a function of $m_{\tilde{\chi}_1^0}$ from the preliminary BABAR limits on $\text{Br}(B^0 \rightarrow \Lambda^0 \psi_D)$ [56] reinterpreted in terms of $B^0 \rightarrow \Lambda^0 \tilde{\chi}_1^0$.

and $n - \bar{n}$ oscillation are much weaker than bounds shown in Fig. 6 and are hence not included in the figure.

Fig. 7 shows our estimated Belle II sensitivity in terms of bounds on $\lambda''_{113}/m_{\tilde{q}}^2$ vs. the neutralino mass. The bound is between 1.5 and $2 \times 10^{-8} \text{ GeV}^{-2}$ for most values of $m_{\tilde{\chi}_1^0}$.

5 Discussion and Conclusions

A GeV-scale, Bino-like lightest neutralino $\tilde{\chi}_1^0$ is allowed by current bounds, provided it decays, e.g., via R -parity-violating (RPV) interactions, and is heavier than the proton, so as not to mediate rapid proton decay. In this work we consider such a light $\tilde{\chi}_1^0$ in the minimal realization of RPV supersymmetry. We propose to probe this scenario with a search for B meson decays into $\tilde{\chi}_1^0$ plus a baryon, specifically p , Λ , Σ^+ , or Σ^0 . The decays are mediated by the RPV operators $\lambda''_{ijk} \bar{U}_i \bar{D}_j \bar{D}_k$. We study the case where only λ''_{113} or λ''_{123} is nonzero in the absence of squark mixing and with the simplifying assumption of mass-degenerate squarks. We compute the corresponding B -meson decay rates in the framework of an effective phenomenological BNV Lagrangian. We note that the lightest neutralino in our benchmark scenarios is so long-lived that it appears as missing energy at the detector level.

Experimentally, the presence of the $\tilde{\chi}_1^0$ and its mass can be determined from the missing mass at an e^+e^- B factory, namely, BABAR, Belle, or Belle II. We interpret a BABAR search for B^0 decay into a Λ baryon plus missing energy [56] to obtain current limits on $\lambda''_{123}/m_{\tilde{q}}^2$ around $1 \times 10^{-6}/\text{GeV}^2$, where $m_{\tilde{q}}$ is the mediating squark mass. Relative to this search, we discuss the sensitivities of the modes $B^+ \rightarrow \Sigma^+ \tilde{\chi}_1^0$ and $B^0 \rightarrow \Sigma^0 \tilde{\chi}_1^0$, as well as subtleties of the extrapolation to the final integrated luminosity of Belle II. Furthermore, we estimate the background level in a future analysis of $B^+ \rightarrow p \tilde{\chi}_1^0$ and the resulting sensitivity to the branching fraction of this decay. From this we obtain the sensitivity of




Figure 6. The estimated Belle II sensitivity in terms of upper bound on λ''_{113} vs. $m_{\tilde{q}}$ for different values of the neutralino mass $m_{\tilde{\chi}_1^0}$, extracted from the proposed search for $B^+ \rightarrow p\tilde{\chi}_1^0$ assuming the sensitivities given in Sec. 3.2. We also plot current collider limits on the scalar squark masses under the hypotheses of eight mass-degenerate light-flavor squarks ($m_{\tilde{q}} \gtrsim 1850$ GeV) and only one light-flavor squark state ($m_{\tilde{q}} \gtrsim 1300$ GeV) [88].




Figure 7. Upper bounds on $\lambda''_{113}/m_{\tilde{q}}^2$ as a function of the neutralino mass, extracted from the process $B^+ \rightarrow p\tilde{\chi}_1^0$, assuming the sensitivity given in Sec. 3.2.

Belle II in terms of the upper limit on $\lambda''_{113}/m_{\tilde{q}}^2$, which is between 1.5 and 2×10^{-8} GeV^{-2} for most kinematically allowed values of $m_{\tilde{\chi}_1^0}$. For the currently allowed squark mass of about

1850 GeV (1300 GeV) in the case of eight (one) degenerate squark states, this corresponds to bounds on λ''_{113} as low as 0.05 (0.025). The bounds are relevant also for multi-TeV squarks beyond the reach of direct detection at LHC.

Lastly, we note that similar studies can be performed with other processes, exploiting the capabilities of other experiments. At LHCb, one can use the decay $B^+ \rightarrow p\pi^+\pi^-\tilde{\chi}_1^0$, with the B^+ produced in the decay $B_{s2}^{*0} \rightarrow B^+K^-$, which has already been used to study semileptonic B decays [89]. Owing to the unobserved neutralino, the 4-momenta of the $\tilde{\chi}_1^0$, B^+ , and B_{s2}^{*0} are unknown. However, they can be determined up to a 2-fold ambiguity from the 12 constraints of the signal decay chain: 4-momentum conservation in the B_{s2}^{*0} and B^+ decays, the known B_{s2}^{*0} and B^+ masses, and the flight direction of the B^+ determined from the position of the $p\pi^+\pi^-$ vertex. Solution of the constraint equations results in two values of the neutralino mass, m_{\pm} , similarly to Refs. [30, 45]. Signal events peak more significantly than background in the m_{\pm} distribution, except for expected peaks around masses of (mostly charmed) baryons that decay to only neutral particles. The feasibility of this approach depends, among other factors, on the m_{\pm} measurement resolution. The decay chains $B_{s2}^{*+} \rightarrow B^0K^+$, $B^0 \rightarrow p\pi^-\tilde{\chi}_1^0$ and $\Sigma_b^+ \rightarrow \Lambda_b\pi^+$, $\Lambda_b \rightarrow \pi^+\pi^-\tilde{\chi}_1^0$ may be used in a similar way. One may also probe λ''_{212} in the decays $D^+ \rightarrow \bar{\Sigma}^-\tilde{\chi}_1^0$ and $D^0 \rightarrow \bar{\Lambda}^0\tilde{\chi}_1^0$ for a smaller range of the neutralino mass. Specifically, the D can be produced in $\psi(3770) \rightarrow D\bar{D}$ at BESIII, with full reconstruction of the \bar{D} in a manner identical to the Belle II analysis we propose here.

Acknowledgements:

We thank Florian Domingo and Martin Hirsch for useful discussions, and Dexu Lin and the BABAR collaboration for help in the interpretation of Ref. [56]. Z. S. W. is supported by the Ministry of Science and Technology (MoST) of Taiwan with grant number MoST-110-2811-M-007-542-MY3. C. O. D. acknowledges support from FONDECYT (Chile) Grant No. 1210131 and ANID (Chile) PIA/APOYO AFB 180002. J. C. H. acknowledges support from grant FONDECYT (Chile) No.1201673 and ANID – Programa Milenio - code ICN2019.044. V. E. L. is supported by BMBF (Germany) “Verbundprojekt 05P2021 (ErUM-FSP T01) - Run 3 von ALICE am LHC: Perturbative Berechnungen von Wirkungsquerschnitten für ALICE” (Förderkennzeichen: 05P21VTCAA), by ANID PIA/APOYO AFB180002 (Chile), by FONDECYT (Chile) under Grant No. 1191103, and by ANID–Millennium Program–ICN2019.044 (Chile). N. A. N. was supported by ANID (Chile) under the grant ANID REC Convocatoria Nacional Subvención a Instalación en la Academia Convocatoria Año 2020, PAI77200092. A. S. is supported by grants from the Israel Science Foundation, the United State-Israel Binational Science Fund, and the Tel Aviv University Center for Artificial Intelligence and Data Science.

A Transition Form Factors

Here, we list the results for the form factors of the $B^+ \rightarrow p(\Sigma^+)\tilde{\chi}_1^0$ and $B^+ \rightarrow \Lambda^0(\Sigma^0)\tilde{\chi}_1^0$ transitions and discuss how to extract the results for the couplings of light baryons with

corresponding three-quark currents. First, we specify the nucleon couplings α and β , which define the matrix elements of three-quark operators between nucleon and vacuum (they are essential ingredient of our calculation of the transition form factors):

$$\langle p | \mathcal{O}_{uud}^{LR} | 0 \rangle = \bar{u}_p(p', s') \alpha P_R, \quad \langle p | \mathcal{O}_{uud}^{LL} | 0 \rangle = -\bar{u}_p(p', s') \beta P_L, \quad (\text{A.1})$$

where

$$\begin{aligned} \mathcal{O}_{q_1 q_2 q_3}^{XY} &= \varepsilon^{a_1 a_2 a_3} (\bar{q}_3^{a_3} P_X \bar{q}_2^{a_2}) \bar{q}_1^{a_1} P_Y, \\ \bar{\mathcal{O}}_{q_1 q_2 q_3}^{XY} &= \varepsilon^{a_1 a_2 a_3} P_Y q_1^{a_1} (q_2^{a_2} P_X q_3^{a_3}), \end{aligned} \quad (\text{A.2})$$

and $X, Y = L, R$. In our calculations we deal only with $\mathcal{O}_{q_1 q_2 q_3}^{LL}$ operators. Therefore, we need only β couplings and form factors $W_{0,1}^{LL}(q^2)$. In addition, for completeness, we also discuss and show results for the case of the $\mathcal{O}_{q_1 q_2 q_3}^{LR}$ operators, i.e. for α couplings and $W_{0,1}^{LR}(q^2)$ form factors.

The nucleon couplings α and β have been calculated using various QCD-motivated approaches and lattice QCD (see, e.g., detailed compilation of the predictions in Table 5 of Ref. [63] and recent lattice results in Ref. [65]). One can see that results of QCD models and lattice QCD for the α coupling range from -0.003 to -0.03 and from -0.006 to -0.03 , respectively. In our analysis, we take recent lattice result $\alpha = -0.01257 \simeq -0.0126$ of the USQCD Collaboration [65] (continuum extrapolation) as central value and take into account its variation as: $\alpha = -0.0126_{-0.0096}^{+0.0174}$. In case of the β coupling mainly lattice QCD results are available, which are distributed from 0.01 to 0.0127 . Therefore, for the β coupling we will use the following predictions with lattice QCD approaches, of the central value from a recent paper of the USQCD Collaboration [65] and take into account error occurring in other lattices calculations [63, 65]: $\beta = 0.0108 \pm 0.007$.

For the case of the transition $B^+ \rightarrow \Sigma^+ \bar{\chi}_1^0$ we need the Σ^+ coupling β_{Σ^+} defining the matrix element $\langle \Sigma^+ | \mathcal{O}_{uus}^{LL} | 0 \rangle$ (for completeness we also calculate the coupling α_{Σ^+} defining the matrix element $\langle \Sigma^+ | \mathcal{O}_{uus}^{LR} | 0 \rangle$):

$$\langle \Sigma^+ | \mathcal{O}_{uus}^{LR} | 0 \rangle = \bar{u}_{\Sigma^+}(p', s') \alpha_{\Sigma^+} P_R, \quad \langle \Sigma^+ | \mathcal{O}_{uus}^{LL} | 0 \rangle = -\bar{u}_{\Sigma^+}(p', s') \beta_{\Sigma^+} P_L. \quad (\text{A.3})$$

The couplings α_{Σ^+} and β_{Σ^+} can be fixed using results for the so-called vector λ_V and tensor λ_T couplings estimated in Ref. [90] using QCD sum rules:

$$\langle 0 | (suu)_V | \Sigma^+ \rangle = \lambda_{\Sigma^+}^V u_{\Sigma^+}(p, s), \quad \langle 0 | (suu)_T | \Sigma^+ \rangle = \lambda_{\Sigma^+}^T u_{\Sigma^+}(p, s), \quad (\text{A.4})$$

where

$$\begin{aligned} (q_1 q_2 q_3)_V &= \varepsilon^{a_1 a_2 a_3} \gamma^\mu \gamma^5 q_1^{a_1} (q_2^{a_2} C \gamma_\mu q_3^{a_3}), \\ (q_1 q_2 q_3)_T &= \varepsilon^{a_1 a_2 a_3} \sigma^{\mu\nu} \gamma^5 q_1^{a_1} (q_2^{a_2} C \sigma^{\mu\nu} q_3^{a_3}) \end{aligned} \quad (\text{A.5})$$

are the vector V and tensor T three-quark currents (operators). Also we will need for further calculations the pseudoscalar P and scalar S three-quark operators:

$$\begin{aligned} (q_1 q_2 q_3)_P &= \varepsilon^{a_1 a_2 a_3} q_1^{a_1} (q_2^{a_2} C \gamma^5 q_3^{a_3}), \\ (q_1 q_2 q_3)_S &= \varepsilon^{a_1 a_2 a_3} \gamma^5 q_1^{a_1} (q_2^{a_2} C q_3^{a_3}). \end{aligned} \quad (\text{A.6})$$

Using Fierz transformation relating V and T currents defined in Eq. (A.5) we can relate them with pseudoscalar P and scalar S currents and then with the currents with specific chirality occurring in Eq. (A.3). In particular, for three-quark operators with flavor content of the Σ^+ baryon we have the following Fierz identities:

$$(suu)_V = 2[(uus)_P - (uus)_S], \quad (suu)_T = 4[(uus)_P + (uus)_S]. \quad (\text{A.7})$$

Next using

$$(uus)_P = \overline{\mathcal{O}}_{uus}^{RR} - \overline{\mathcal{O}}_{uud}^{LL} + \overline{\mathcal{O}}_{uud}^{RL} - \overline{\mathcal{O}}_{uud}^{LR}, \quad (\text{A.8})$$

$$(uus)_S = \overline{\mathcal{O}}_{uus}^{RR} - \overline{\mathcal{O}}_{uud}^{LL} - \overline{\mathcal{O}}_{uud}^{RL} + \overline{\mathcal{O}}_{uud}^{LR}, \quad (\text{A.9})$$

we can express the matrix elements of V and T currents through the matrix elements of the $\overline{\mathcal{O}}_{uus}^{RR}$ operators and, therefore, relate then the pair of the couplings $\lambda_{\Sigma^+}^V, \lambda_{\Sigma^+}^T$ with couplings α_{Σ^+} and β_{Σ^+} :

$$\begin{aligned} \langle 0|(suu)_V|\Sigma^+\rangle &= 4 \left[\langle 0|\overline{\mathcal{O}}_{uus}^{RL}|\Sigma^+\rangle - \langle 0|\overline{\mathcal{O}}_{uus}^{LR}|\Sigma^+\rangle \right] \\ &= 4\alpha_{\Sigma^+} u_{\Sigma^+}(p, s), \end{aligned} \quad (\text{A.10})$$

$$\begin{aligned} \langle 0|(suu)_T|\Sigma^+\rangle &= 8 \left[\langle 0|\overline{\mathcal{O}}_{uus}^{RR}|\Sigma^+\rangle - \langle 0|\overline{\mathcal{O}}_{uus}^{LL}|\Sigma^+\rangle \right] \\ &= -8\beta_{\Sigma^+} u_{\Sigma^+}(p, s). \end{aligned} \quad (\text{A.11})$$

Matching Eqs. (A.4) and (A.10) we arrive at the required relations:

$$\alpha_{\Sigma^+} = \frac{1}{4}\lambda_{\Sigma^+}^V, \quad \beta_{\Sigma^+} = -\frac{1}{8}\lambda_{\Sigma^+}^T. \quad (\text{A.12})$$

Taking $\lambda_{\Sigma^+}^{V,T}$ couplings from Ref. [90] with the sign conventions as in Ref. [63] we get:

$$\alpha_{\Sigma^+} = -(0.743 \pm 0.030) \times 10^{-2} \text{ GeV}^3, \quad \beta_{\Sigma^+} = (0.654 \pm 0.030) \times 10^{-2} \text{ GeV}^3. \quad (\text{A.13})$$

For the transitions $B^0 \rightarrow \Lambda^0 \tilde{\chi}_1^0$ and $B^0 \rightarrow \Sigma^0 \tilde{\chi}_1^0$ we need the Λ^0 and Σ^0 couplings β_{Λ^0} and β_{Σ^0} defining the following matrix elements

$$\langle \Lambda^0 | \mathcal{O}_{sud}^{LL} | 0 \rangle = -\bar{u}_p(p', s') \beta_{\Lambda^0} P_L, \quad (\text{A.14})$$

$$\langle \Sigma^0 | 1/\sqrt{2} [\mathcal{O}_{uds}^{LL} + \mathcal{O}_{dus}^{LL}] | 0 \rangle = -\bar{u}_p(p', s') \beta_{\Sigma^0} P_L. \quad (\text{A.15})$$

The β_{Λ^0} coupling was calculated in Ref. [90] using QCD sum rules:

$$\beta_{\Lambda^0} = (0.926 \pm 0.056) \times 10^{-2} \text{ GeV}^3. \quad (\text{A.16})$$

As for β_{Σ^0} we accept isospin-symmetry relation: $\beta_{\Sigma^0} = \beta_{\Sigma^+}$.

As we stressed above, the form factors $W_0^{LX}(q^2)$ and $W_1^{LX}(q^2)$ defining matrix elements $\langle p | \mathcal{O}^{LX} | B^+ \rangle$ can be calculated using SU(5) extension of the SU(3) version of the phenomenological Lagrangian proposed and used in Refs. [66] and [61–65]. At the leading order form factors $W_0^{LX}(q^2)$ and $W_1^{LX}(q^2)$ are contributed by direct BNV couplings

of proton and pseudoscalar mesons and pole contributions because of exchange of baryon resonances. In particular, using results for the BNV matrix elements $\langle p | \mathcal{O}^{LX} | K^+ \rangle$ and making replacements $f_K \rightarrow f_B$, $m_{\Lambda^0} \rightarrow m_{\Lambda_b}$, and $m_{\Sigma} \rightarrow m_{\Sigma_b}$, one obtains:

$$\begin{aligned}
W_0^{LL}(q^2) = & -\frac{\beta}{f_B} \left[g^{\tilde{b}R} \left(1 + \frac{D+3F}{3} \frac{m_{\Lambda_b} m_p + q^2}{m_{\Lambda_b}^2 - q^2} \right) \right. \\
& + g^{\tilde{u}R} \left(-1 - \frac{D+3F}{6} \frac{m_{\Lambda_b} m_p + q^2}{m_{\Lambda_b}^2 - q^2} + \frac{D-F}{2} \frac{m_{\Sigma_b} m_p + q^2}{m_{\Sigma_b}^2 - q^2} \right) \\
& \left. + g^{\tilde{d}R} \left(\frac{D+3F}{6} \frac{m_{\Lambda_b} m_p + q^2}{m_{\Lambda_b}^2 - q^2} + \frac{D-F}{2} \frac{m_{\Sigma_b} m_p + q^2}{m_{\Sigma_b}^2 - q^2} \right) \right], \quad (\text{A.17})
\end{aligned}$$

$$\begin{aligned}
W_1^{LL}(q^2) = & -\frac{\beta}{f_B} m_{\tilde{\chi}_1^0} \left[g^{\tilde{b}R} \frac{D+3F}{3} \frac{m_{\Lambda_b} + m_p}{m_{\Lambda_b}^2 - q^2} \right. \\
& + g^{\tilde{u}R} \left(-\frac{D+3F}{6} \frac{m_{\Lambda_b} + m_p}{m_{\Lambda_b}^2 - q^2} + \frac{D-F}{2} \frac{m_{\Sigma_b} + m_p}{m_{\Sigma_b}^2 - q^2} \right) \\
& \left. + g^{\tilde{d}R} \left(\frac{D+3F}{6} \frac{m_{\Lambda_b} + m_p}{m_{\Lambda_b}^2 - q^2} + \frac{D-F}{2} \frac{m_{\Sigma_b} + m_p}{m_{\Sigma_b}^2 - q^2} \right) \right], \quad (\text{A.18})
\end{aligned}$$

$$\begin{aligned}
W_0^{LR}(q^2) = & -\frac{\alpha}{f_B} \left[g^{\tilde{b}L} \left(1 + \frac{D+3F}{3} \frac{m_{\Lambda_b} m_p + q^2}{m_{\Lambda_b}^2 - q^2} \right) \right. \\
& + g^{\tilde{u}L} \left(-1 - \frac{D+3F}{6} \frac{m_{\Lambda_b} m_p + q^2}{m_{\Lambda_b}^2 - q^2} + \frac{D-F}{2} \frac{m_{\Sigma_b} m_p + q^2}{m_{\Sigma_b}^2 - q^2} \right) \\
& \left. + g^{\tilde{d}L} \left(\frac{D+3F}{6} \frac{m_{\Lambda_b} m_p + q^2}{m_{\Lambda_b}^2 - q^2} + \frac{D-F}{2} \frac{m_{\Sigma_b} m_p + q^2}{m_{\Sigma_b}^2 - q^2} \right) \right], \quad (\text{A.19})
\end{aligned}$$

$$\begin{aligned}
W_1^{LR}(q^2) = & -\frac{\alpha}{f_B} m_{\tilde{\chi}_1^0} \left[g^{\tilde{b}L} \frac{D+3F}{3} \frac{m_{\Lambda_b} + m_p}{m_{\Lambda_b}^2 - q^2} \right. \\
& + g^{\tilde{u}L} \left(-\frac{D+3F}{6} \frac{m_{\Lambda_b} + m_p}{m_{\Lambda_b}^2 - q^2} + \frac{D-F}{2} \frac{m_{\Sigma_b} + m_p}{m_{\Sigma_b}^2 - q^2} \right) \\
& \left. + g^{\tilde{d}L} \left(\frac{D+3F}{6} \frac{m_{\Lambda_b} + m_p}{m_{\Lambda_b}^2 - q^2} + \frac{D-F}{2} \frac{m_{\Sigma_b} + m_p}{m_{\Sigma_b}^2 - q^2} \right) \right], \quad (\text{A.20})
\end{aligned}$$

where $D = 0.8$, $F = 0.47$, $f_B = 0.192$ GeV, $m_{\Lambda_b} = 5.6196$ GeV, $m_{\Sigma_b} = 5.81056$ GeV, $m_{B^+} = 5.27934$ GeV, and $m_p = 0.93827$ GeV. In case of the $B^+ \rightarrow \Sigma^+ + \tilde{\chi}_1^0$ transition we replace the masses of the baryons in the expressions for the form factors (see Eqs. (A.17) and (A.18)) and kinematical formula for the decay rate (Eq. (2.10)) accordingly: $m_{\Lambda_b} \rightarrow m_{\Xi_b} = 5.7919$ GeV, $m_{\Sigma_b} \rightarrow m_{\Xi_b^+} = 5.93502$ GeV, and $m_p \rightarrow m_{\Sigma^+} = 1.18937$ GeV.

Note that direct BNV couplings of proton/ Σ^+ and B^+ meson occur on the partonic level only in the case of the subprocesses induced by the \tilde{b} and \tilde{u} squarks and can not occur

in the case of the \tilde{d} squark for the proton mode and \tilde{s} for the Σ^+ mode. In particular, direct BNV coupling of proton and Σ^+ with B^+ means that the latter transits into two three-quark currents. One current corresponds to the proton current J^P or Σ^+ hyperon current J^{Σ^+} and another one to the BNV operator J^{BNV} . The BNV operator has a form of scalar diquark $[q_1 q_2]_0$ coupled to the third quark q_3 . We have three possibilities for the J^{BNV} (for simplicity we drop color indices):

$$J_{bud}^{\text{BNV}} = \bar{b}[\bar{u}\bar{d}]_0, \quad J_{udb}^{\text{BNV}} = \bar{u}[\bar{d}\bar{b}]_0, \quad J_{dub}^{\text{BNV}} = \bar{d}[\bar{u}\bar{b}]_0, \quad (\text{A.21})$$

for the proton mode and

$$J_{bus}^{\text{BNV}} = \bar{b}[\bar{u}\bar{s}]_0, \quad J_{usb}^{\text{BNV}} = \bar{u}[\bar{s}\bar{b}]_0, \quad J_{sub}^{\text{BNV}} = \bar{s}[\bar{u}\bar{b}]_0, \quad (\text{A.22})$$

for the Σ^+ mode. It means that the B^+ meson having spin $J = 0$ can decay into light baryons having scalar quark configuration. In our case, proton and Σ^+ hyperon could be formed as quark-scalar diquark bound state:

$$J^P = u[ud]_0 \quad J^{\Sigma^+} = u[us]_0. \quad (\text{A.23})$$

Therefore, the following direct transitions are allowed: (1) B^+ meson annihilates into u and \bar{b} quarks, which further play the role of third quarks in light baryon and BNV current. In this case, the pair of scalar diquark $[ud]_0$ and antidiquark $[\bar{u}\bar{d}]_0$ is produced from vacuum; (2) B^+ meson annihilates into the pair of scalar diquark $[ud]_0$ and antidiquark $[\bar{b}\bar{d}]_0$. The allowed direct modes are listed as:

Direct transition induced by the \tilde{b} squark

$$B^+(u\bar{b}) \rightarrow J^P + J_{bud}^{\text{BNV}}, \quad B^+(u\bar{b}) \rightarrow J^{\Sigma^+} + J_{bus}^{\text{BNV}}. \quad (\text{A.24})$$

Direct transition induced by the \tilde{u} squark

$$B^+(u\bar{b}) \rightarrow J^P + J_{udb}^{\text{BNV}}, \quad B^+(u\bar{b}) \rightarrow J^{\Sigma^+} + J_{usb}^{\text{BNV}}. \quad (\text{A.25})$$

We have no direct transitions induced by the \tilde{d} squark for the proton mode and by the \tilde{s} squark for the Σ^+ mode, because it requires production of the pair of scalar diquark contributing to the J^{BNV} current and vector diquark ($d\{uu\}_1$ or $s\{uu\}_1$) contributing to the current of light baryon, which is suppressed by spin conservation.

Now we turn to discussion of the allowed pole transitions induced by bottom baryon resonances. We have two types of bottom baryon resonances with quark content of the BNV operators: baryons Λ_b and Ξ_b with antisymmetric light quark configuration, whose three-quark currents could have a form of light spin-0 diquark $[ud]_0$ and $[us]_0$, respectively, coupled to the b quark

$$J^{\Lambda_b} = b[ud]_0, \quad J^{\Xi_b} = b[us]_0, \quad (\text{A.26})$$

or vector diquarks composed of light quark u , d , or s and bottom quark b :

$$J^{\Lambda_b} = u\{db\}_1 - d\{ub\}_1, \quad J^{\Xi_b} = u\{sb\}_1 - s\{ub\}_1, \quad (\text{A.27})$$

and baryons $\Sigma_b = b\{ud\}_1$ and $\Xi'_b = b\{us\}_1$ with symmetric light quark configuration, whose three-quark currents could have a form of light spin-1 diquark $\{ud\}_1$ and $\{us\}_1$, respectively coupled to the b quark

$$J^{\Sigma_b} = b\{ud\}_1, \quad J^{\Xi'_b} = b\{us\}_1, \quad (\text{A.28})$$

or scalar diquarks composed of light quark u , d , or s and bottom quark b :

$$J^{\Sigma_b} = u[db]_0 + d[ub]_0, \quad J^{\Xi'_b} = u[sb]_0 + s[ub]_0. \quad (\text{A.29})$$

Therefore, pole transitions with Λ_b and Ξ_b baryon resonances are allowed for all three squarks \tilde{b} , \tilde{u} , and \tilde{d} in the case of the proton mode and \tilde{b} , \tilde{u} , and \tilde{s} in the case of the Σ^+ mode.

On the partonic level $B^+ \rightarrow p\Lambda_b$ and $B^+ \rightarrow \Sigma^+\Xi_b$ transitions are induced by the quark transitions, respectively:

$$B^+(u\bar{b}) \rightarrow p(u[ud]_0) + \bar{\Lambda}_b^0(\bar{b}[\bar{u}\bar{d}]_0), \quad B^+(u\bar{b}) \rightarrow \Sigma^+(u[us]_0) + \bar{\Xi}_b^0(\bar{b}[\bar{u}\bar{s}]_0). \quad (\text{A.30})$$

Next, the Λ_b or Σ_b converses into J^{BNV} with emission of the neutralino. In particular, the transitions induced by the squark \tilde{b} exchange are

$$\bar{b}[\bar{u}\bar{d}]_0 \rightarrow \bar{b}[\bar{u}\bar{d}]_0, \quad \bar{b}[\bar{u}\bar{s}]_0 \rightarrow \bar{b}[\bar{u}\bar{s}]_0. \quad (\text{A.31})$$

Transitions induced by the squarks \tilde{u} and $\tilde{d}(s)$ are

$$\bar{u}[\bar{d}\bar{b}]_0 \rightarrow \bar{u}[\bar{d}\bar{b}]_0, \quad \bar{u}[\bar{s}\bar{b}]_0 \rightarrow \bar{u}[\bar{s}\bar{b}]_0, \quad (\text{A.32})$$

and

$$\bar{d}[\bar{u}\bar{b}]_0 \rightarrow \bar{s}[\bar{u}\bar{b}]_0, \quad \bar{s}[\bar{u}\bar{b}]_0 \rightarrow \bar{s}[\bar{u}\bar{b}]_0. \quad (\text{A.33})$$

One can explain relative contributions of all three channels to the hadronic form factors. The \tilde{u} and $\tilde{d}(s)$ channel contributions are generated by the Fierz transformation of the Λ_b and Ξ_b currents $J^{\Lambda_b} = b[ud]_0$ and $J^{\Xi_b} = b[us]_0$ allowing to interchange $b \leftrightarrow u$ and $b \leftrightarrow d(s)$. It leads to the currents

$$J^{\Lambda_b} = -\frac{1}{2}u[db]_0 + \dots = \frac{1}{2}d[ub]_0 + \dots, \quad (\text{A.34})$$

$$J^{\Xi_b} = -\frac{1}{2}u[sb]_0 + \dots = \frac{1}{2}s[ub]_0 + \dots. \quad (\text{A.35})$$

From Eqs. (A.31)-(A.35) one can see that relative contributions of the \tilde{b} , \tilde{u} , and $\tilde{d}(s)$ squark channels are $2 : (-1) : 1$. Note, that the contributions of \tilde{u} and $\tilde{d}(s)$ squarks are equal on magnitude and of opposite sign as a result of antisymmetric configuration of (u, d) and (u, s) quarks in the Λ_b and Ξ_b baryons, respectively (see Eq. (A.26)).

From Eqs. (A.28) and (A.29), we conclude that in the case of the pole contributions induced by the bottom baryons Σ_b and Ξ'_b with symmetric configurations of light quarks, the squark \tilde{b} mode is absent because spin-1 diquark $[ud(s)]_1$ cannot transit to the spin-0

diquark $[ud(s)]_0$. On the other hand, the \tilde{u} and $\tilde{d}(s)$ squark modes are allowed and they have exactly the same contributions because of the symmetric configuration of light quarks in the Σ_b and Ξ'_b baryons.

By analogy, we consider, in addition, decay modes of the neutral B^0 mesons $B^0 \rightarrow \Lambda^0 \tilde{\chi}_1^0$ and $B^0 \rightarrow \Sigma^0 \tilde{\chi}_1^0$. In particular, the corresponding hadronic form factors defining these decay read:

$$\begin{aligned}
W_0^{LL}(q^2) = & -\sqrt{\frac{3}{2}} \frac{\beta_{\Lambda^0}}{f_B} \left[g^{\tilde{b}R} \left(-\frac{1}{2} - \frac{D+3F}{6} \frac{m_{\Xi_b} m_{\Lambda^0} + q^2}{m_{\Xi_b}^2 - q^2} \right) \right. \\
& + g^{\tilde{u}R} \left(-\frac{1}{2} + \frac{D+3F}{12} \frac{m_{\Xi_b} m_{\Lambda^0} + q^2}{m_{\Xi_b}^2 - q^2} + \frac{3(D-F)}{4} \frac{m_{\Xi'_b} m_{\Lambda^0} + q^2}{m_{\Xi'_b}^2 - q^2} \right) \\
& \left. + g^{\tilde{s}R} \left(-1 - \frac{D+3F}{12} \frac{m_{\Xi_b} m_{\Lambda^0} + q^2}{m_{\Xi_b}^2 - q^2} + \frac{3(D-F)}{4} \frac{m_{\Xi'_b} m_{\Lambda^0} + q^2}{m_{\Xi'_b}^2 - q^2} \right) \right], \quad (\text{A.36})
\end{aligned}$$

$$\begin{aligned}
W_1^{LL}(q^2) = & -\sqrt{\frac{3}{2}} \frac{\beta_{\Lambda^0}}{f_B} m_{\tilde{\chi}_1^0} \left[-g^{\tilde{b}R} \frac{D+3F}{6} \frac{m_{\Xi_b} + m_{\Lambda^0}}{m_{\Xi_b}^2 - q^2} \right. \\
& + g^{\tilde{u}R} \left(\frac{D+3F}{12} \frac{m_{\Xi_b} + m_{\Lambda^0}}{m_{\Xi_b}^2 - q^2} + \frac{3(D-F)}{4} \frac{m_{\Xi'_b} + m_{\Lambda^0}}{m_{\Xi'_b}^2 - q^2} \right) \\
& \left. + g^{\tilde{s}R} \left(-\frac{D+3F}{12} \frac{m_{\Xi_b} + m_{\Lambda^0}}{m_{\Xi_b}^2 - q^2} + \frac{3(D-F)}{4} \frac{m_{\Xi'_b} + m_{\Lambda^0}}{m_{\Xi'_b}^2 - q^2} \right) \right], \quad (\text{A.37})
\end{aligned}$$

for the Λ^0 mode and

$$\begin{aligned}
W_0^{LL}(q^2) = & -\sqrt{\frac{1}{2}} \frac{\beta_{\Sigma}}{f_B} \left[g^{\tilde{b}R} \left(1 + \frac{D+3F}{3} \frac{m_{\Xi_b} m_{\Sigma^0} + q^2}{m_{\Xi_b}^2 - q^2} \right) \right. \\
& + g^{\tilde{u}R} \left(-1 - \frac{D+3F}{6} \frac{m_{\Xi_b} m_{\Sigma^0} + q^2}{m_{\Xi_b}^2 - q^2} + \frac{D-F}{2} \frac{m_{\Xi'_b} m_p + q^2}{m_{\Xi'_b}^2 - q^2} \right) \\
& \left. + g^{\tilde{s}R} \left(\frac{D+3F}{6} \frac{m_{\Xi_b} m_{\Sigma^0} + q^2}{m_{\Xi_b}^2 - q^2} + \frac{D-F}{2} \frac{m_{\Xi'_b} m_{\Sigma^0} + q^2}{m_{\Xi'_b}^2 - q^2} \right) \right], \quad (\text{A.38})
\end{aligned}$$

$$\begin{aligned}
W_1^{LL}(q^2) = & -\sqrt{\frac{1}{2}} \frac{\beta_{\Sigma}}{f_B} m_{\tilde{\chi}_1^0} \left[g^{\tilde{b}R} \frac{D+3F}{3} \frac{m_{\Xi_b} + m_{\Sigma^0}}{m_{\Xi_b}^2 - q^2} \right. \\
& + g^{\tilde{u}R} \left(-\frac{D+3F}{6} \frac{m_{\Xi_b} + m_{\Sigma^0}}{m_{\Xi_b}^2 - q^2} + \frac{D-F}{2} \frac{m_{\Xi'_b} + m_{\Sigma^0}}{m_{\Xi'_b}^2 - q^2} \right) \\
& \left. + g^{\tilde{s}R} \left(\frac{D+3F}{6} \frac{m_{\Xi_b} + m_{\Sigma^0}}{m_{\Xi_b}^2 - q^2} + \frac{D-F}{2} \frac{m_{\Xi'_b} + m_{\Sigma^0}}{m_{\Xi'_b}^2 - q^2} \right) \right], \quad (\text{A.39})
\end{aligned}$$

for the Σ^0 mode. For the masses of the Λ^0 and Σ^0 , we take the central values from the Particle Data Group [91]: $m_{\Lambda^0} = 1.115683$ GeV and $m_{\Sigma^0} = 1.192642$ GeV.

References

- [1] H. P. Nilles, *Supersymmetry, Supergravity and Particle Physics*, *Phys. Rept.* **110** (1984) 1.
- [2] S. P. Martin, *A Supersymmetry primer*, *Adv. Ser. Direct. High Energy Phys.* **21** (2010) 1, [[hep-ph/9709356](#)].
- [3] E. Gildener, *Gauge Symmetry Hierarchies*, *Phys. Rev. D* **14** (1976) 1667.
- [4] M. Veltman, *The Infrared - Ultraviolet Connection*, *Acta Phys. Polon. B* **12** (1981) 437.
- [5] N. Chamoun, F. Domingo, and H. K. Dreiner, *Nucleon decay in the R-parity violating MSSM*, *Phys. Rev. D* **104** (2021), no. 1 015020, [[arXiv:2012.11623](#)].
- [6] H. K. Dreiner, *An Introduction to explicit R-parity violation*, *Adv. Ser. Direct. High Energy Phys.* **21** (2010) 565, [[hep-ph/9707435](#)].
- [7] R. Barbier et al., *R-parity violating supersymmetry*, *Phys. Rept.* **420** (2005) 1, [[hep-ph/0406039](#)].
- [8] R. N. Mohapatra, *Supersymmetry and R-parity: an Overview*, *Phys. Scripta* **90** (2015) 088004, [[arXiv:1503.06478](#)].
- [9] **ATLAS** Collaboration, M. Aaboud et al., *Search for photonic signatures of gauge-mediated supersymmetry in 13 TeV pp collisions with the ATLAS detector*, *Phys. Rev. D* **97** (2018), no. 9 092006, [[arXiv:1802.03158](#)].
- [10] **CMS** Collaboration, A. M. Sirunyan et al., *Search for gauge-mediated supersymmetry in events with at least one photon and missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV*, *Phys. Lett. B* **780** (2018) 118–143, [[arXiv:1711.08008](#)].
- [11] **CMS** Collaboration, A. M. Sirunyan et al., *Search for supersymmetry in final states with photons and missing transverse momentum in proton-proton collisions at 13 TeV*, *JHEP* **06** (2019) 143, [[arXiv:1903.07070](#)].
- [12] **CMS** Collaboration, A. M. Sirunyan et al., *Search for supersymmetry in proton-proton collisions at 13 TeV in final states with jets and missing transverse momentum*, *JHEP* **10** (2019) 244, [[arXiv:1908.04722](#)].
- [13] **ATLAS** Collaboration, G. Aad et al., *Search for new phenomena in final states with large jet multiplicities and missing transverse momentum using $\sqrt{s} = 13$ TeV proton-proton collisions recorded by ATLAS in Run 2 of the LHC*, *JHEP* **10** (2020) 062, [[arXiv:2008.06032](#)].
- [14] D. Choudhury and S. Sarkar, *A Supersymmetric resolution of the KARMEN anomaly*, *Phys. Lett. B* **374** (1996) 87–92, [[hep-ph/9511357](#)].
- [15] D. Choudhury, H. K. Dreiner, P. Richardson, and S. Sarkar, *A Supersymmetric solution to the KARMEN time anomaly*, *Phys. Rev. D* **61** (2000) 095009, [[hep-ph/9911365](#)].
- [16] G. Belanger, F. Boudjema, A. Pukhov, and S. Rosier-Lees, *A Lower limit on the neutralino mass in the MSSM with nonuniversal gaugino masses*, in *10th International Conference on Supersymmetry and Unification of Fundamental Interactions (SUSY02)*, pp. 919–924, 12, 2002. [[hep-ph/0212227](#)].
- [17] A. Bottino, N. Fornengo, and S. Scopel, *Light relic neutralinos*, *Phys. Rev. D* **67** (2003) 063519, [[hep-ph/0212379](#)].
- [18] G. Belanger, F. Boudjema, A. Cottrant, A. Pukhov, and S. Rosier-Lees, *Lower limit on the neutralino mass in the general MSSM*, *JHEP* **03** (2004) 012, [[hep-ph/0310037](#)].

- [19] D. Albornoz Vasquez, G. Belanger, C. Boehm, A. Pukhov, and J. Silk, *Can neutralinos in the MSSM and NMSSM scenarios still be light?*, *Phys. Rev. D* **82** (2010) 115027, [[arXiv:1009.4380](#)].
- [20] L. Calibbi, J. M. Lindert, T. Ota, and Y. Takahashi, *Cornering light Neutralino Dark Matter at the LHC*, *JHEP* **10** (2013) 132, [[arXiv:1307.4119](#)].
- [21] I. Gogoladze, J. D. Lykken, C. Macesanu, and S. Nandi, *Implications of a Massless Neutralino for Neutrino Physics*, *Phys. Rev. D* **68** (2003) 073004, [[hep-ph/0211391](#)].
- [22] H. K. Dreiner, S. Heinemeyer, O. Kittel, U. Langenfeld, A. M. Weber, and G. Weiglein, *Mass Bounds on a Very Light Neutralino*, *Eur. Phys. J. C* **62** (2009) 547–572, [[arXiv:0901.3485](#)].
- [23] J. de Vries, H. K. Dreiner, and D. Schmeier, *R-Parity Violation and Light Neutralinos at SHiP and the LHC*, *Phys. Rev. D* **94** (2016), no. 3 035006, [[arXiv:1511.07436](#)].
- [24] H. Dreiner, C. Hanhart, U. Langenfeld, and D. R. Phillips, *Supernovae and light neutralinos: SN1987A bounds on supersymmetry revisited*, *Phys. Rev. D* **68** (2003) 055004, [[hep-ph/0304289](#)].
- [25] F. Domingo and H. K. Dreiner, *Decays of a bino-like particle in the low-mass regime*, [arXiv:2205.08141](#).
- [26] D. Hooper and T. Plehn, *Supersymmetric dark matter: How light can the LSP be?*, *Phys. Lett. B* **562** (2003) 18, [[hep-ph/0212226](#)].
- [27] A. Bottino, N. Fornengo, and S. Scopel, *Phenomenology of light neutralinos in view of recent results at the CERN Large Hadron Collider*, *Phys. Rev. D* **85** (2012) 095013, [[arXiv:1112.5666](#)].
- [28] G. Bélanger, G. Drieu La Rochelle, B. Dumont, R. M. Godbole, S. Kraml, and S. Kulkarni, *LHC constraints on light neutralino dark matter in the MSSM*, *Phys. Lett. B* **726** (2013) 773, [[arXiv:1308.3735](#)].
- [29] P. Bechtle et al., *Killing the cMSSM softly*, *Eur. Phys. J. C* **76** (2016), no. 2 96, [[arXiv:1508.05951](#)].
- [30] S. Dey, C. O. Dib, J. Carlos Helo, M. Nayak, N. A. Neill, A. Soffer, and Z. S. Wang, *Long-lived light neutralinos at Belle II*, *JHEP* **02** (2021) 211, [[arXiv:2012.00438](#)].
- [31] D. Dercks, J. De Vries, H. K. Dreiner, and Z. S. Wang, *R-parity Violation and Light Neutralinos at CODEX-b, FASER, and MATHUSLA*, *Phys. Rev. D* **99** (2019), no. 5 055039, [[arXiv:1810.03617](#)].
- [32] D. Dercks, H. K. Dreiner, M. Hirsch, and Z. S. Wang, *Long-Lived Fermions at AL3X*, *Phys. Rev. D* **99** (2019), no. 5 055020, [[arXiv:1811.01995](#)].
- [33] H. K. Dreiner, J. Y. Günther, and Z. S. Wang, *R-parity Violation and Light Neutralinos at ANUBIS and MAPP*, [arXiv:2008.07539](#).
- [34] H. K. Dreiner, D. Köhler, S. Nangia, and Z. S. Wang, *Searching for a Single Photon from Lightest Neutralino Decays in R-parity-violating Supersymmetry at FASER*, [arXiv:2207.05100](#).
- [35] P. Candia, G. Cottin, A. Méndez, and V. Muñoz, *Searching for light long-lived neutralinos at Super-Kamiokande*, *Phys. Rev. D* **104** (2021), no. 5 055024, [[arXiv:2107.02804](#)].
- [36] **Belle-II** Collaboration, T. Abe et al., *Belle II Technical Design Report*, [arXiv:1011.0352](#).

- [37] **Belle-II** Collaboration, W. Altmannshofer et al., *The Belle II Physics Book*, *PTEP* **2019** (2019), no. 12 123C01, [[arXiv:1808.10567](#)]. [Erratum: *PTEP* 2020, 029201 (2020)].
- [38] L. Calibbi, G. Ferretti, D. Milstead, C. Petersson, and R. Pöttgen, *Baryon number violation in supersymmetry: $n - \bar{n}$ oscillations as a probe beyond the LHC*, *JHEP* **05** (2016) 144, [[arXiv:1602.04821](#)]. [Erratum: *JHEP* 10, 195 (2017)].
- [39] D. McKeen and A. E. Nelson, *CP Violating Baryon Oscillations*, *Phys. Rev. D* **94** (2016), no. 7 076002, [[arXiv:1512.05359](#)].
- [40] K. Aitken, D. McKeen, T. Neder, and A. E. Nelson, *Baryogenesis from Oscillations of Charmed or Beautiful Baryons*, *Phys. Rev. D* **96** (2017), no. 7 075009, [[arXiv:1708.01259](#)].
- [41] M. I. Buchoff and M. Wagman, *Perturbative Renormalization of Neutron-Antineutron Operators*, *Phys. Rev. D* **93** (2016), no. 1 016005, [[arXiv:1506.00647](#)]. [Erratum: *Phys.Rev.D* 98, 079901 (2018)].
- [42] G. Elor, M. Escudero, and A. Nelson, *Baryogenesis and Dark Matter from B Mesons*, *Phys. Rev. D* **99** (2019), no. 3 035031, [[arXiv:1810.00880](#)].
- [43] G. Alonso-Álvarez, G. Elor, and M. Escudero, *Collider signals of baryogenesis and dark matter from B mesons: A roadmap to discovery*, *Phys. Rev. D* **104** (2021), no. 3 035028, [[arXiv:2101.02706](#)].
- [44] A. Filimonova, R. Schäfer, and S. Westhoff, *Probing dark sectors with long-lived particles at BELLE II*, *Phys. Rev. D* **101** (2020), no. 9 095006, [[arXiv:1911.03490](#)].
- [45] C. Dib, J. Helo, M. Nayak, N. Neill, A. Soffer, and J. Zamora-Saa, *Searching for a sterile neutrino that mixes predominantly with ν_τ at B factories*, *Phys. Rev. D* **101** (2020), no. 9 093003, [[arXiv:1908.09719](#)].
- [46] T. Ferber, A. Filimonova, R. Schäfer, and S. Westhoff, *Displaced or invisible? ALPs from B decays at Belle II*, [arXiv:2201.06580](#).
- [47] E. Bertholet, S. Chakraborty, V. Loladze, T. Okui, A. Soffer, and K. Tobioka, *Heavy QCD axion at Belle II: Displaced and prompt signals*, *Phys. Rev. D* **105** (2022), no. 7 L071701, [[arXiv:2108.10331](#)].
- [48] K. Cheung, A. Soffer, Z. S. Wang, and Y.-H. Wu, *Probing charged lepton flavor violation with axion-like particles at Belle II*, *JHEP* **11** (2021) 218, [[arXiv:2108.11094](#)].
- [49] C. S. Kim, Y. Kwon, D. Lee, S. Oh, and D. Sahoo, *Probing sterile neutrinos in $B(D)$ meson decays at Belle II (BESIII)*, *Eur. Phys. J. C* **80** (2020), no. 8 730, [[arXiv:1908.00376](#)].
- [50] D. W. Kang, P. Ko, and C.-T. Lu, *Exploring properties of long-lived particles in inelastic dark matter models at Belle II*, *JHEP* **04** (2021) 269, [[arXiv:2101.02503](#)].
- [51] M. Acevedo, A. Blackburn, N. Blinov, B. Shuve, and M. Stone, *Multi-track displaced vertices at B-factories*, *JHEP* **09** (2021) 154, [[arXiv:2105.12744](#)].
- [52] S. Dreyer et al., *Physics reach of a long-lived particle detector at Belle II*, [arXiv:2105.12962](#).
- [53] M. Duerr, T. Ferber, C. Hearty, F. Kahlhoefer, K. Schmidt-Hoberg, and P. Tunney, *Invisible and displaced dark matter signatures at Belle II*, *JHEP* **02** (2020) 039, [[arXiv:1911.03176](#)].
- [54] M. Duerr, T. Ferber, C. Garcia-Cely, C. Hearty, and K. Schmidt-Hoberg, *Long-lived Dark Higgs and Inelastic Dark Matter at Belle II*, *JHEP* **04** (2021) 146, [[arXiv:2012.08595](#)].
- [55] X. Chen, Z. Hu, Y. Wu, and K. Yi, *Search for dark photon and dark matter signatures around electron-positron colliders*, *Phys. Lett. B* **814** (2021) 136076, [[arXiv:2001.04382](#)].

- [56] **BABAR** Collaboration, D. Lin and S. M. (for the collaboration), *Search for Baryogenesis and Dark Matter in B-meson Decays at BABAR, Poster presented at ICHEP 2022* (2022).
- [57] S. Weinberg, *Supersymmetry at Ordinary Energies. 1. Masses and Conservation Laws*, *Phys. Rev. D* **26** (1982) 287.
- [58] L. J. Hall and M. Suzuki, *Explicit R-Parity Breaking in Supersymmetric Models*, *Nucl. Phys. B* **231** (1984) 419.
- [59] A. Faessler, T. Gutsche, J. C. Helo, S. Kovalenko, and V. E. Lyubovitskij, *On the possible resolution of the B-meson decay polarization anomaly in R-parity violating SUSY*, *Phys. Rev. D* **75** (2007) 074029, [[hep-ph/0702020](#)].
- [60] C. S. Li, Z. Li, R. J. Oakes, and L. L. Yang, *Threshold Resummation Effects in the Associated Production of Chargino and Neutralino at Hadron Colliders*, *Phys. Rev. D* **77** (2008) 034010, [[arXiv:0707.3952](#)].
- [61] **JLQCD** Collaboration, S. Aoki et al., *Nucleon decay matrix elements from lattice QCD*, *Phys. Rev. D* **62** (2000) 014506, [[hep-lat/9911026](#)].
- [62] Y. Aoki, C. Dawson, J. Noaki, and A. Soni, *Proton decay matrix elements with domain-wall fermions*, *Phys. Rev. D* **75** (2007) 014507, [[hep-lat/0607002](#)].
- [63] **RBC-UKQCD** Collaboration, Y. Aoki, P. Boyle, P. Cooney, L. Del Debbio, R. Kenway, C. M. Maynard, A. Soni, and R. Tweedie, *Proton lifetime bounds from chirally symmetric lattice QCD*, *Phys. Rev. D* **78** (2008) 054505, [[arXiv:0806.1031](#)].
- [64] Y. Aoki, E. Shintani, and A. Soni, *Proton decay matrix elements on the lattice*, *Phys. Rev. D* **89** (2014), no. 1 014505, [[arXiv:1304.7424](#)].
- [65] J.-S. Yoo, Y. Aoki, P. Boyle, T. Izubuchi, A. Soni, and S. Syritsyn, *Proton decay matrix elements on the lattice at physical pion mass*, *Phys. Rev. D* **105** (2022), no. 7 074501, [[arXiv:2111.01608](#)].
- [66] M. B. Gavela, S. F. King, C. T. Sachrajda, G. Martinelli, M. L. Paciello, and B. Taglienti, *Lattice Computation of Proton Decay Amplitudes*, *Nucl. Phys. B* **312** (1989) 269.
- [67] S. Okubo, *SU(4), SU(8) Mass Formulas and Weak Interactions*, *Phys. Rev. D* **11** (1975) 3261.
- [68] W. Liu, C. M. Ko, and Z. W. Lin, *Cross-section for charmonium absorption by nucleons*, *Phys. Rev. C* **65** (2002) 015203.
- [69] Y. Dong, A. Faessler, and V. E. Lyubovitskij, *Description of heavy exotic resonances as molecular states using phenomenological Lagrangians*, *Prog. Part. Nucl. Phys.* **94** (2017) 282.
- [70] Y. Dong, A. Faessler, T. Gutsche, and V. E. Lyubovitskij, *Strong two-body decays of the $\Lambda_c(2940)^+$ in a hadronic molecule picture*, *Phys. Rev. D* **81** (2010) 014006, [[arXiv:0910.1204](#)].
- [71] Y. Dong, A. Faessler, T. Gutsche, and V. E. Lyubovitskij, *Charmed baryon $\Sigma_c(2800)$ as a ND hadronic molecule*, *Phys. Rev. D* **81** (2010) 074011, [[arXiv:1002.0218](#)].
- [72] **CLEO** Collaboration, T. E. Browder et al., *A Search for $B \rightarrow \tau\nu$* , *Phys. Rev. Lett.* **86** (2001) 2950, [[hep-ex/0007057](#)].
- [73] **BaBar** Collaboration, B. Aubert et al., *A search for the decay $B^+ \rightarrow K^+\nu\bar{\nu}$* , *Phys. Rev. Lett.* **94** (2005) 101801, [[hep-ex/0411061](#)].

- [74] **BaBar** Collaboration, P. del Amo Sanchez et al., *Search for the Rare Decay $B \rightarrow K\nu\bar{\nu}$* , *Phys. Rev. D* **82** (2010) 112002, [[arXiv:1009.1529](#)].
- [75] **BaBar** Collaboration, J. P. Lees et al., *Search for $B \rightarrow K^{(*)}\nu\bar{\nu}$ and invisible quarkonium decays*, *Phys. Rev. D* **87** (2013), no. 11 112005, [[arXiv:1303.7465](#)].
- [76] **Belle** Collaboration, K. F. Chen et al., *Search for $B \rightarrow h^{(*)}\nu\bar{\nu}$ decays at Belle*, *Phys. Rev. Lett.* **99** (2007) 221802, [[arXiv:0707.0138](#)].
- [77] **Belle** Collaboration, O. Lutz et al., *Search for $B \rightarrow h^{(*)}\nu\bar{\nu}$ with the full Belle $\Upsilon(4S)$ data sample*, *Phys. Rev. D* **87** (2013), no. 11 111103, [[arXiv:1303.3719](#)].
- [78] **Belle** Collaboration, J. Grygier et al., *Search for $B \rightarrow h\nu\bar{\nu}$ decays with semileptonic tagging at Belle*, *Phys. Rev. D* **96** (2017), no. 9 091101, [[arXiv:1702.03224](#)]. [Addendum: *Phys.Rev.D* 97, 099902 (2018)].
- [79] **Belle-II** Collaboration, F. Abudinén et al., *Search for $B \rightarrow K + \nu\nu$ Decays Using an Inclusive Tagging Method at Belle II*, *Phys. Rev. Lett.* **127** (2021), no. 18 181802, [[arXiv:2104.12624](#)].
- [80] **BaBar, Belle** Collaboration, A. J. Bevan et al., *The Physics of the B Factories*, *Eur. Phys. J. C* **74** (2014) 3026, [[arXiv:1406.6311](#)].
- [81] **CLEO** Collaboration, J. P. Alexander et al., *First measurement of the $B \rightarrow \pi$ lepton neutrino and $B \rightarrow \rho$ (omega) lepton neutrino branching fractions*, *Phys. Rev. Lett.* **77** (1996) 5000.
- [82] A. Soffer, *B-meson decays into final states with a τ lepton*, *Mod. Phys. Lett. A* **29** (2014), no. 07 1430007, [[arXiv:1401.7947](#)].
- [83] **Particle Data Group** Collaboration, P. Zyla et al., *Review of Particle Physics*, *PTEP* **2020** (2020), no. 8 083C01.
- [84] **ARGUS** Collaboration, H. Albrecht et al., *Kaons in flavor tagged B decays*, *Z. Phys. C* **62** (1994) 371–382.
- [85] **CLEO** Collaboration, M. S. Alam et al., *Branching Ratios of B Meson to K^+ , K^- and K^0/\bar{K}^0* , *Phys. Rev. Lett.* **58** (1987) 1814.
- [86] **Belle** Collaboration, C. Hadjivasiliou et al., *Search for B^0 meson decays into Λ and missing energy with a hadronic tagging method at Belle*, *Phys. Rev. D* **105** (2022), no. 5 L051101, [[arXiv:2110.14086](#)].
- [87] S. Dey, *private communication*, .
- [88] **ATLAS** Collaboration, G. Aad et al., *Search for squarks and gluinos in final states with jets and missing transverse momentum using 139 fb^{-1} of $\sqrt{s} = 13 \text{ TeV}$ pp collision data with the ATLAS detector*, *JHEP* **02** (2021) 143, [[arXiv:2010.14293](#)].
- [89] **LHCb** Collaboration, R. Aaij et al., *Measurement of the relative $B^- \rightarrow D^0/D^{*0}/D^{**0}\mu^-\bar{\nu}_\mu$ branching fractions using B^- mesons from \bar{B}_{s2}^{*0} decays*, *Phys. Rev. D* **99** (2019), no. 9 092009, [[arXiv:1807.10722](#)].
- [90] Y.-L. Liu and M.-Q. Huang, *Distribution amplitudes of Σ and Λ and their electromagnetic form factors*, *Nucl. Phys. A* **821** (2009) 80, [[arXiv:0811.1812](#)].
- [91] **Particle Data Group** Collaboration, P. A. Zyla et al., *Review of Particle Physics*, *PTEP* **2020** (2020), no. 8 083C01.