FERMI ISOSPECTRALITY OF DISCRETE PERIODIC SCHRÖDINGER OPERATORS WITH SEPARABLE POTENTIALS ON \mathbb{Z}^2

WENCAI LIU

ABSTRACT. Let $\Gamma = q_1 \mathbb{Z} \oplus q_2 \mathbb{Z}$ with $q_1 \in \mathbb{Z}_+$ and $q_2 \in \mathbb{Z}_+$. Let $\Delta + X$ be the discrete periodic Schrödinger operator on \mathbb{Z}^2 , where Δ is the discrete Laplacian and $X : \mathbb{Z}^2 \to \mathbb{C}$ is Γ -periodic. In this paper, we develop tools from complex analysis to study the isospectrality of discrete periodic Schrödinger operators. We prove that if two Γ -periodic potentials X and Y are Fermi isospectral and both $X = X_1 \oplus X_2$ and $Y = Y_1 \oplus Y_2$ are separable functions, then, up to a constant, one dimensional potentials X_j and Y_j are Floquet isospectral, j = 1, 2. This allows us to prove that for any non-constant separable real-valued Γ -periodic potential, the Fermi variety $F_{\lambda}(V)/\mathbb{Z}^2$ is irreducible for any $\lambda \in \mathbb{C}$, which partially confirms a conjecture of Gieseker, Knörrer and Trubowitz in the early 1990s.

1. INTRODUCTION AND MAIN RESULTS

Given $q_j \in \mathbb{Z}_+$, $j = 1, 2, \cdots, d$, let $\Gamma = q_1 \mathbb{Z} \oplus q_2 \mathbb{Z} \oplus \cdots \oplus q_d \mathbb{Z}$. We say that a function $V : \mathbb{Z}^d \to \mathbb{C}$ is Γ -periodic (or just periodic) if for any $\gamma \in \Gamma$ and $n \in \mathbb{Z}^d$, $V(n + \gamma) = V(n)$. For $n = (n_1, n_2, \cdots, n_d) \in \mathbb{Z}^d$, denote by $||n||_1 = \sum_{j=1}^d |n_j|$. Let Δ be the discrete Laplacian on lattices \mathbb{Z}^d , namely

$$(\Delta u)(n) = \sum_{n' \in \mathbb{Z}^d, ||n'-n||_1 = 1} u(n').$$

In the following, we always assume that q_j , $j = 1, 2, \dots, d$, are pairwise coprime and V is Γ -periodic.

Key words and phrases. Fermi variety, irreducibility, Rouche's theorem, Fermi isospectrality, Floquet isospectrality, periodic Schrödinger operator, separable potentials.

²⁰²⁰ Mathematics Subject Classification. Primary: 58J53. Secondary: 35J10, 47A75.

In this article we are interested in the isospectrality problem and irreducibility of Fermi varieties of discrete periodic Schrödinger operators $\Delta + V$. We refer readers to two survey articles [21, 27] for background and recent developments about the two topics.

Let $\{\mathbf{e}_j\}, j = 1, 2, \cdots d$, be the standard basis in \mathbb{Z}^d :

$$\mathbf{e}_1 = (1, 0, \cdots, 0), \mathbf{e}_2 = (0, 1, 0, \cdots, 0), \cdots, \mathbf{e}_d = (0, 0, \cdots, 0, 1).$$

Definition 1. The Bloch variety B(V) of $\Delta + V$ consists of all pairs $(k, \lambda) \in \mathbb{C}^{d+1}$ for which there exists a non-zero solution of the equation

(1)
$$(\Delta u)(n) + V(n)u(n) = \lambda u(n), n \in \mathbb{Z}^d,$$

satisfying the so called Floquet-Bloch boundary condition

(2)
$$u(n+q_j e_j) = e^{2\pi i k_j} u(n), j = 1, 2, \cdots, d, \text{ and } n \in \mathbb{Z}^d,$$

where $k = (k_1, k_2, \cdots, k_d) \in \mathbb{C}^d$.

Given $\lambda \in \mathbb{C}$, the Fermi surface (variety) $F_{\lambda}(V)$ is defined as the level set of the Bloch variety:

$$F_{\lambda}(V) = \{k : (k, \lambda) \in B(V)\}.$$

We call $k = (k_1, k_2, \dots, k_d)$ that appears in (2) quasi-momentum. One can see that both Fermi and Bloch varieties are analytic sets, in fact algebraic sets after changing variables [21, 26, 27].

Our first interest is the isospectrality problems.

Let $D_V(k)$ be the periodic operator $\Delta + V$ with the Floquet-Bloch boundary condition (2) (see Section 2 for the precise description of $D_V(k)$). Two Γ -periodic potentials X and Y are called Floquet isospectral if

(3)
$$\sigma(D_X(k)) = \sigma(D_Y(k)), \text{ for any } k \in \mathbb{R}^d$$

Two Γ -periodic potentials X and Y are called isospectral if

(4)
$$\sigma(D_X(k)) = \sigma(D_Y(k)) \text{ with } k = 0.$$

Understanding when two periodic potentials X and Y are Floquet isospectral or isospectral is a fascinating subject and has been extensively studied [5–7, 13–15, 17–19, 21, 28, 31].

In [25], the author introduced a new type of isospectrality: Fermi isospectrality.

Definition 2. [25] Let X and Y be two Γ -periodic functions. We say X and Y are Fermi isospectral if $F_{\lambda_0}(X) = F_{\lambda_0}(Y)$ for some $\lambda_0 \in \mathbb{C}$.

It is not difficult to see that two periodic functions X and Y are Floquet isospectral if and only if Bloch varieties of X and Y are the same

 $\mathbf{2}$

(or Fermi varieties of X and Y are the same for every $\lambda \in \mathbb{C}$) [25]. So Fermi isospectrality is a "hyperplane" version of Floquet isospectrality.

In [25], the author proved several rigidity theorems of discrete periodic Schrödinger operators about separable functions. We say that a function V on \mathbb{Z}^d is (d_1, d_2, \dots, d_r) separable (or simply separable, denote it by $V = \bigoplus_{j=1}^r V_j$), where $\sum_{j=1}^r d_j = d$ with $r \ge 2$, if there exist functions V_j on \mathbb{Z}^{d_j} , $j = 1, 2, \dots, r$, such that for any $(n_1, n_2, \dots, n_d) \in \mathbb{Z}^d$,

$$V(n_1, n_2, \cdots, n_d) = V_1(n_1, \cdots, n_{d_1}) + V_2(n_{d_1+1}, n_{d_1+2}, \cdots, n_{d_1+d_2}) + \cdots + V_r(n_{d_1+d_2+\cdots+d_{r-1}+1}, \cdots, n_{d_1+d_2+\cdots+d_r}).$$

One of rigidity theorems in [25] states

Theorem 1.1. [25] Let $d \ge 3$. Assume that two separable Γ -periodic potentials $X = \bigoplus_{j=1}^{r} X_j$ and $Y = \bigoplus_{j=1}^{r} Y_j$ are Fermi isospectral. Then, up to a constant, lower dimensional decompositions V_j and Y_j are Floquet isospectral, $j = 1, 2, \dots, r$.

In the present work, we prove that the statement in Theorem 1.1 holds for dimension d = 2. Namely,

Theorem 1.2. Let d = 2. Assume that two Γ -periodic potentials Xand Y are Fermi isospectral and both $X = X_1 \oplus X_2$ and $Y = Y_1 \oplus Y_2$ are separable. Then, up to a constant, one dimensional functions X_j and Y_j are Floquet isospectral, j = 1, 2.

Remark 1. In Theorems 1.1 and 1.2, potentials are allowed to be complex-valued.

Our second interest of this paper is the irreducibility of Fermi varieties. Irreducibility of Fermi varieties (also Bloch varieties) and related applications such as embedded eigenvalues and spectral band edges have seen continuous progress in the past 30 years [1–4, 9–12, 20, 22–24, 29, 30].

Recently, the author introduced an algebraic method and provided more general proofs of irreducibility of Fermi varieties [26].

Denote by [V] the average of V over one periodicity cell, namely

$$[V] = \frac{1}{q_1 q_2 \cdots q_d} \sum_{\substack{1 \le n_j \le q_j \\ 1 \le j \le d}} V(n_1, n_2, \cdots, n_d).$$

Theorem 1.3. [26] For any $d \geq 3$, the Fermi variety $F_{\lambda}(V)/\mathbb{Z}^d$ is irreducible for any $\lambda \in \mathbb{C}$. For d = 2, the Fermi variety $F_{\lambda}(V)/\mathbb{Z}^2$ is irreducible for any $\lambda \in \mathbb{C}$ except maybe for $\lambda = [V]$ and $F_{[V]}(V)/\mathbb{Z}^2$ has at most two irreducible components.

Before [26], the irreducibility of Fermi varieties at all energy levels for d = 3 and at all energy levels but finitely many λ for d = 2 was proved in [2, 12] by an different approach (compactification).

Let d = 2. When the potential V is a constant function, direct computation (e.g., see [27]) implies that $F_{[V]}(V)/\mathbb{Z}^2$ has exactly two irreducible components. When the complex-valued functions are allowed, there exist non-constant complex valued functions V such that the Fermi variety is reducible at the energy level [V] (e.g. [8]).

However, for real-valued potentials, people believe the constant potential is the only case that the Fermi variety $F_{\lambda}(V)/\mathbb{Z}^2$ is reducible at some energy level, which has been formulated as a conjecture by Gieseker, Knörrer and Trubowitz in the early 1990s [12].

Conjecture 1: [12, p.43] Assume that V is a non-constant realvalued periodic potential on \mathbb{Z}^2 . Then the Fermi variety $F_{\lambda}(V)/\mathbb{Z}^2$ is irreducible for any $\lambda \in \mathbb{C}$.

Theorem 1.2 allows us to confirm the Conjecture 1 for separable potentials.

Theorem 1.4. Assume that V is a non-constant separable realvalued periodic potential on \mathbb{Z}^2 . Then the Fermi variety $F_{\lambda}(V)/\mathbb{Z}^2$ is irreducible for any $\lambda \in \mathbb{C}$.

The irreducibility of Fermi variety and Fermi isospectrality of discrete periodic Schrödinger operators (dimension $d \geq 3$) are well understood in two recent papers [25, 26]. Besides Theorem 1.1, there are other Fermi isospectrality results in [25] for dimension d > 3. However, approaches in [25] can not be extended to dimension d = 2 since there are not enough free variables available. For irreducibility results of the Fermi variety in Theorem 1.3, the proof for d = 2 is more difficult than that for d > 3. For continuous periodic Schrödinger operators, Bättig, Knörrer and Trubowitz [3] proved the irreducibility of Fermi varieties and a rigidity theorem of separable functions in dimension three. However, the proof in [3] does not work for dimension d = 2. For discrete periodic Schrödinger operators on \mathbb{Z}^d with $d \geq 3$, the Fermi variety $F_{\lambda}(V)/\mathbb{Z}^d$ for any complex-valued potential is irreducible at any energy level λ (see Theorem 1.3). For d = 2, there are many complexvalued potentials V such that the Fermi variety $F_{\lambda}(V)/\mathbb{Z}^2$ has two irreducible components at the average energy level [V] [8].

Finally, we want to comment that dimension two is the transition of Fermi isospectrality problems of periodic Shcrödinger operators. For d = 1, it does not make sense to study Fermi isospectrality since for any periodic potential V, $F_{\lambda_0}(V)$ contains at most two points. For d = 2 and any periodic potential Y, all periodic potentials X such that

X and Y are Fermi isospectral at λ_0 (namely $F_{\lambda_0}(X) = F_{\lambda_0}(Y)$) is an algebraic set with at least one dimension [8]. For $d \geq 3$ and any periodic potential Y, all periodic potentials X such that X and Y are Fermi isospectral at λ_0 could be an algebraic set with zero dimension [8].

All evidence above seems to indicate that when d = 2, problems related to Fermi varieties are special (often more challenging).

In this paper, we present a novel approach to study the Fermi isospectrality of discrete periodic Schrödinger operators. As in [25], we focus on the study of a family of Laurent polynomials whose zero sets are Fermi varieties after changing variables. Our strategy is to develop tools from complex analysis to study the eigenvalue problems of (1) and (2) (or (6) and (7)) with complexified quasi-momenta. One needs to relabel spectral band functions of one dimensional periodic Schrödinger operators based on asymptotics of eigenvalues and show that those functions are analytic with respect to quasi-momenta in an appropriate domain. Applying Rouche's Theorem, one sees that for any two one-dimensional q-periodic potentials with the same average, there exist q choices of quasi-momenta such that for those quasieigenvalues of two potentials only differ by a momenta, labelled (same) constant. This enables us to show that separable components of Fermi isospectrality potentials with respect to one coordinate are Floquet isospectral and hence remaining separable components are Floquet isospectral as well.

The rest of this paper is organized as follows. In Section 2, we recall some basics for Fermi varieties. In Section 3, we study one dimensional periodic Schrödinger operators. Section 4 is devoted to proving Theorems 1.2 and 1.4.

2. Basics of Fermi varieties

Let $\mathbb{C}^{\star} = \mathbb{C} \setminus \{0\}$ and $z = (z_1, z_2, \cdots, z_d)$. For any $z \in (\mathbb{C}^{\star})^d$, consider the equation

(6)
$$(\Delta + V)u = \lambda u$$

with the boundary condition

(7)
$$u(n+q_j\mathbf{e}_j) = z_j u(n), j = 1, 2, \cdots, d, \text{ and } n \in \mathbb{Z}^d,$$

Introduce a fundamental domain W for Γ :

$$W = \{ n = (n_1, n_2, \cdots, n_d) \in \mathbb{Z}^d : 0 \le n_j \le q_j - 1, j = 1, 2, \cdots, d \}.$$

By writing out $\Delta + V$ as acting on the $Q = q_1 q_2 \cdots q_d$ dimensional space $\{u(n), n \in W\}$, the equation (6) with boundary condition (7)

((1) and (2)) translates into the eigenvalue problem for a $Q \times Q$ matrix $\mathcal{D}_V(z)$ $(D_V(k))$.

Let

(8)
$$\mathcal{P}_V(z,\lambda) = \det(\mathcal{D}_V(z) - \lambda I), P_V(k,\lambda) = \det(D_V(k) - \lambda I).$$

We remark that $\mathcal{D}_V(z)$ and $D_V(k)$ ($\mathcal{P}_V(z,\lambda)$ and $\mathcal{P}_V(k,\lambda)$) are the same under the relations $z_j = e^{2\pi i k_j}, j = 1, 2, \cdots, d$.

Example 1: When d = 1, the equation $(\Delta + V)u = \lambda u$ with the Floquet-Bloch boundary condition $u(n + q) = zu(n), z \in \mathbb{C}^*$, can be reduced to an eigenvalue problem of a $q \times q$ matrix:

(9)
$$\mathcal{D}_{V}(z) = \begin{pmatrix} V(1) & 1 & 0 & \cdots & 0 & z^{-1} \\ 1 & V(2) & 1 & \cdots & 0 & 0 \\ 0 & 1 & V(3) & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & V(q-1) & 1 \\ z & 0 & 0 & \cdots & 1 & V(q) \end{pmatrix}.$$

We have the following **Basic Facts**:

(1) Fermi variety is given by

(10)
$$F_{\lambda}(V) = \{k \in \mathbb{C}^d : P_V(k,\lambda) = 0\}.$$

(2) Two periodic functions X and Y are Floquet isospectral if and only if

(11)
$$\mathcal{P}_X(z,\lambda) = \mathcal{P}_Y(z,\lambda).$$

3. One dimensional discrete periodic Schrödinger operators

In this section, we study one dimensional discrete periodic Schrödinger operators $\Delta + V$:

$$((\Delta + V)u)(n) = u(n+1) + u(n-1) + V(n)u(n), n \in \mathbb{Z},$$

where V is a periodic function on \mathbb{Z} , namely, $V(n+q) = V(n), n \in \mathbb{Z}$ for some positive integer q.

In the following, we say z is large if |z| is large.

By (9) in Example 1, $\mathcal{P}_V(z,\lambda) - (-1)^{q+1}z - (-1)^{q+1}z^{-1}$ is independent of variable z. So let $\hat{\mathcal{P}}_V(\lambda)$ be such that

(12)
$$\mathcal{P}_V(z,\lambda) = \hat{\mathcal{P}}_V(\lambda) + (-1)^{q+1}z + (-1)^{q+1}z^{-1}.$$

By (9), one has that (recall that $[V] = \frac{1}{q} (\sum_{j=1}^{q} V(j))$)

(13) $\hat{\mathcal{P}}_V(z,\lambda) = (-1)^q \lambda^q - (-1)^q q[V] \lambda^{q-1} + \text{lower order terms of } \lambda.$

 $\mathbf{6}$

 $\overline{7}$

Fixing $z \in \mathbb{C}$, solve the algebraic equation

(14)
$$\mathcal{P}_V(z^q,\lambda) = 0.$$

By (12) and (13), there exist solutions $\lambda_V^l(z)$ of equation (14), $l = 0, 1, 2, \cdots, q-1$ such that $\lambda^l(z)$ is analytic in $\Omega = \{z \in \mathbb{C} : |z| > R\}$ with large R (the largeness only depends on the potential V). Moreover, $\lambda_V^l(z), l = 0, 1, 2, \cdots, q-1$ have the following representations in Laurent series,

(15)
$$\lambda_V^l(z) = e^{2\pi \frac{l}{q}i} z + [V] + \sum_{k=1}^{\infty} \frac{a_k(V)}{z^k},$$

where the coefficient $a_k(V)$ depends on V.

Clearly, for any large $z, \lambda_V^l(z), l = 0, 1, 2, \dots, q-1$, are eigenvalues of $\mathcal{D}(z^q)$, and hence

(16)
$$\mathcal{P}_V(z^q,\lambda) = \det(\mathcal{D}(z^q) - \lambda I) = \prod_{l=0}^{q-1} (\lambda_V^l(z) - \lambda).$$

Lemma 3.1. Assume $[V] = [\tilde{V}]$, and V and \tilde{V} are not Floquet isospectral. Then there exist sufficiently large R > 0 and small $\epsilon > 0$ such that for any $\eta \in \mathbb{C}$ with $0 < |\eta| < \epsilon$ and $l = 0, 1, 2, \cdots, q-1$, the equation $\lambda_V^l(z) = \lambda_{\tilde{V}}^l(z) + \eta$ has at least one solution in $\{z \in \mathbb{C} : |z| \ge R\}$.

Proof. Fix any $l \in \{0, 1, \dots, q-1\}$. By (15), one has that for any large z,

(17)
$$\lambda_V^l(z) = e^{2\pi \frac{l}{q}i} z + [V] + \sum_{k=1}^{\infty} \frac{a_k(V)}{z^k},$$

and

(18)
$$\lambda_{\tilde{V}}^{l}(z) = e^{2\pi \frac{l}{q}i}z + [\tilde{V}] + \sum_{k=1}^{\infty} \frac{a_{k}(\tilde{V})}{z^{k}},$$

If $\lambda_V^l(z) = \lambda_{\tilde{V}}^l(z)$ for any large z, then V and \tilde{V} must be Flqouet isospectral, which contradicts the assumption. So functions $\lambda_V^l(z)$ and $\lambda_{\tilde{V}}^l(z)$ are not identical.

When $a_1(V) \neq a_1(\tilde{V})$, let $k_0 = 1$. Otherwise, let $k_0 \in \mathbb{Z}_+$ be such that $a_k(V) = a_k(\tilde{V})$ for any $k < k_0$ and $a_{k_0}(V) \neq a_{k_0}(\tilde{V})$. Consider a ball $B_{\epsilon_1} = \{z \in \mathbb{C} : |z| \leq \epsilon_1\}$ with a small $\epsilon_1 > 0$. Define an analytic function f(z) in B_{ϵ_1} by

(19)
$$f(z) = \sum_{k=k_0}^{\infty} (a_k(V)z^k - a_k(\tilde{V})z^k).$$

Then for any large z,

(20)
$$f(z^{-1}) = \lambda_V^l(z) - \lambda_{\tilde{V}}^l(z).$$

Let

$$f_1(z) = a_k(V)z^{k_0} - a_k(\tilde{V})z^{k_0},$$

and

$$f_2(z) = \sum_{k=k_0+1}^{\infty} (a_k(V)z^k - a_k(\tilde{V})z^k)$$

For any $z \in \partial B_{\epsilon_1} = \{z \in \mathbb{C} : |z| = \epsilon_1\}$, one has that

(21)
$$|f_1(z)| \ge |f_2(z)| + \frac{|a_{k_0}(V) - a_{k_0}(\tilde{V})|}{2} \epsilon_1^{k_0}.$$

Let ϵ be sufficiently small (depending on ϵ_1). Choose any η with $0 < |\eta| \le \epsilon$. By (21), one has that for any $z \in \partial B_{\epsilon_1}$,

(22)
$$|f_1(z) - \eta| > |f_2(z)|.$$

By Rouche's theorem, $f_1(z) - \eta = 0$ and $f(z) - \eta = f_1(z) + f_2(z) - \eta = 0$ have the same number of zeros (counting multiplicity) in $\{z \in \mathbb{C} : |z| < \epsilon_1\}$. This particularly implies that $f(z) = \eta$ has at least one non-zero solution in $\{z \in \mathbb{C} : |z| < \epsilon_1\}$. Now Lemma 3.1 follows from (20).

4. Proof of Theorems 1.2 and 1.4

Lemma 4.1. [25, Lemma 2.3] Assume $F_{\lambda_0}(X) = F_{\lambda_0}(Y)$. Then for any $z \in (\mathbb{C}^*)^2$,

(23)
$$\mathcal{P}_X(z,\lambda_0) = \mathcal{P}_Y(z,\lambda_0).$$

Lemma 4.2. Assume that separable functions $X = X_1 \oplus X_2$ and $Y = Y_1 \oplus Y_2$ are Fermi isospectral. Assume that X_2 and Y_2 are Floquet isospectral. Then X_1 and Y_1 are Floquet isospectral.

Proof. Recall that $z = (z_1, z_2)$. By the assumption that $F_{\lambda_0}(X) = F_{\lambda_0}(Y)$ and Lemma 4.1, one has that

$$\mathcal{P}_X(z,\lambda_0) = \mathcal{P}_Y(z,\lambda_0),$$

and hence

(24)
$$\mathcal{P}_X(z_1, z_2^{q_2}, \lambda_0) = \mathcal{P}_Y(z_1, z_2^{q_2}, \lambda_0).$$

Using the fact that both X and Y are separable, one has that for any large z_2 ,

$$\mathcal{P}_X(z_1, z_2^{q_2}, \lambda_0) = \prod_{l=0}^{q_2-1} \mathcal{P}_{X_1}(z_1, -\lambda_{X_2}^l(z_2) + \lambda_0)$$

(25)
$$= \prod_{l=0}^{q_2-1} (\hat{\mathcal{P}}_{X_1}(-\lambda_{X_2}^l(z_2) + \lambda_0) + (-1)^{q_1+1}z_1 + (-1)^{q_1+1}z_1^{-1}),$$

and

$$\mathcal{P}_{Y}(z_{1}, z_{2}^{q_{2}}, \lambda_{0}) = \prod_{l=0}^{q_{2}-1} \mathcal{P}_{Y_{1}}(z_{1}, -\lambda_{Y_{2}}^{l}(z_{2}) + \lambda_{0})$$

$$(26) \qquad = \prod_{l=0}^{q_{2}-1} (\hat{\mathcal{P}}_{Y_{1}}(-\lambda_{Y_{2}}^{l}(z_{2}) + \lambda_{0}) + (-1)^{q_{1}+1}z_{1} + (-1)^{q_{1}+1}z_{1}^{-1}).$$

By (15), (24), (25), (26), and the unique factorization theorem (using $(-1)^{q_1+1}z_1 + (-1)^{q_1+1}z_1^{-1}$ as a variable), one has that for any $l = 0, 1, \dots, q_2 - 1$,

(27)
$$\hat{\mathcal{P}}_{X_1}(-\lambda_{X_2}^l(z_2)+\lambda_0) = \hat{\mathcal{P}}_{Y_1}(-\lambda_{Y_2}^l(z_2)+\lambda_0),$$

and

(28)
$$\mathcal{P}_{X_1}(z_1, -\lambda_{X_2}^l(z_2) + \lambda_0) = \mathcal{P}_{Y_1}(z_1, -\lambda_{Y_2}^l(z_2) + \lambda_0).$$

Since X_2 and Y_2 are Floquet isospectral, we know that for any large z_2 ,

(29)
$$\lambda_{X_2}^l(z_2) = \lambda_{Y_2}^l(z_2), l = 0, 1, 2, \cdots, q_2 - 1.$$

By (28) and (29), one has that for any $z_1 \in \mathbb{C}^*$ and $\lambda \in \mathbb{C}$,

(30)
$$\mathcal{P}_{X_1}(z_1,\lambda) = \mathcal{P}_{Y_1}(z_1,\lambda)$$

By (30) and basic fact (2) appearing at the end of Section 2, we conclude that X_1 and Y_1 are Floquet isospectral.

Proof of Theorem 1.2. Without loss of generality, assume that $q_2 > q_1$ and $[X_2] = [Y_2] = 0$. If X_2 and Y_2 are Floquet isospectral, then Theorem 1.2 follows from Lemma 4.2. So we assume X_2 and Y_2 are not Floquet isospectral.

Applying Lemma 3.1 with $V = X_2$, $\tilde{V} = Y_2$ and $q = q_2$, there exist η and large x_l , $l = 0, 1, 2, \dots, q_2 - 1$, such that

(31)
$$\lambda_{X_2}^l(x_l) = \lambda_{Y_2}^l(x_l) + \eta.$$

By (27) and (31), we have that for any $l = 0, 1, 2, \dots, q_2 - 1$,

(32)
$$\hat{\mathcal{P}}_{X_1}(-\lambda_{Y_2}^l(x_l) - \eta + \lambda_0) = \hat{\mathcal{P}}_{Y_1}(-\lambda_{Y_2}^l(x_l) + \lambda_0).$$

Since both $\hat{\mathcal{P}}_{X_1}(\lambda - \eta)$ and $\hat{\mathcal{P}}_{Y_1}(\lambda)$ are polynomials of λ with degree q_1 , by (32) and the fact that $q_2 > q_1$, one has that

(33)
$$\hat{\mathcal{P}}_{X_1}(\lambda - \eta) = \hat{\mathcal{P}}_{X_1}(\lambda), \lambda \in \mathbb{C}.$$

This implies that X_1 and Y_1 are Floquet isospectral up to a constant (by letting $\eta \to 0$ in (33), we can indeed show that X_1 and Y_1 are Floquet isospectral. This is because we have already shifted the constant by setting $[X_2] = [Y_2] = 0$). Now Theorem 1.2 follows from Lemma 4.2 (exchange X_1 and X_2 , and Y_1 and Y_2).

Denote by **0** the zero function on \mathbb{Z}^2 .

Proof of Theorem 1.4. Assume that $F_{\lambda}(V)$ is reducible at some $\lambda = \lambda_0$. By Remark 4 in [26] (also Theorem 2.5 in [25]), $\lambda_0 = [V]$ and $\mathcal{P}_V(z,\lambda_0) = \mathcal{P}_0(z,0)$. Therefore $\mathcal{P}_{V-[V]}(z,0) = \mathcal{P}_0(z,0)$. By Theorem 1.2, we have that V and a constant potential are Floquet isospectral. Therefore, Ambarzumian-type theorem (e.g. [16]) implies V is constant. This contradicts the assumption.

Acknowledgments

This research was supported by NSF DMS-2000345 and DMS-2052572.

References

- D. Bättig. A directional compactification of the complex Fermi surface and isospectrality. In Séminaire sur les Équations aux Dérivées Partielles, 1989–1990, pages Exp. No. IV, 11. École Polytech., Palaiseau, 1990.
- [2] D. Bättig. A toroidal compactification of the Fermi surface for the discrete Schrödinger operator. *Comment. Math. Helv.*, 67(1):1–16, 1992.
- [3] D. Bättig, H. Knörrer, and E. Trubowitz. A directional compactification of the complex Fermi surface. *Compositio Math.*, 79(2):205– 229, 1991.
- [4] N. Do, P. Kuchment, and F. Sottile. Generic properties of dispersion relations for discrete periodic operators. J. Math. Phys., 61(10):103502, 19, 2020.
- [5] G. Eskin. Inverse spectral problem for the Schrödinger equation with periodic vector potential. *Comm. Math. Phys.*, 125(2):263– 300, 1989.

- [6] G. Eskin, J. Ralston, and E. Trubowitz. On isospectral periodic potentials in Rⁿ. Comm. Pure Appl. Math., 37(5):647–676, 1984.
- [7] G. Eskin, J. Ralston, and E. Trubowitz. On isospectral periodic potentials in Rⁿ. II. Comm. Pure Appl. Math., 37(6):715–753, 1984.
- [8] M. Faust, W. Liu, and F. Sottile. In preparation. 2022.
- [9] M. Faust and F. Sottile. Critical points of discrete periodic operators. arXiv preprint arXiv:2206.13649, 2022.
- [10] J. Fillman, W. Liu, and R. Matos. Irreducibility of the bloch variety for finite-range Schrödinger operators. J. Funct. Anal. to appear.
- [11] L. Fisher, W. Li, and S. P. Shipman. Reducible Fermi surface for multi-layer quantum graphs including stacked graphene. *Comm. Math. Phys.*, 385(3):1499–1534, 2021.
- [12] D. Gieseker, H. Knörrer, and E. Trubowitz. The geometry of algebraic Fermi curves, volume 14 of Perspectives in Mathematics. Academic Press, Inc., Boston, MA, 1993.
- [13] C. S. Gordon and T. Kappeler. On isospectral potentials on tori. Duke Math. J., 63(1):217–233, 1991.
- [14] C. S. Gordon and T. Kappeler. On isospectral potentials on flat tori. II. Comm. Partial Differential Equations, 20(3-4):709–728, 1995.
- [15] V. Guillemin. Inverse spectral results on two-dimensional tori. J. Amer. Math. Soc., 3(2):375–387, 1990.
- [16] B. Hatinoğlu, J. Eakins, W. Frendreiss, L. Lamb, S. Manage, and A. Puente. Ambarzumian-type problems for discrete Schrödinger operators. *Complex Anal. Oper. Theory*, 15(8):Paper No. 118, 13, 2021.
- [17] T. Kappeler. On isospectral periodic potentials on a discrete lattice. I. Duke Math. J., 57(1):135–150, 1988.
- [18] T. Kappeler. On isospectral potentials on a discrete lattice. II. Adv. in Appl. Math., 9(4):428–438, 1988.
- [19] T. Kappeler. Isospectral potentials on a discrete lattice. III. Trans. Amer. Math. Soc., 314(2):815–824, 1989.
- [20] H. Knörrer and E. Trubowitz. A directional compactification of the complex Bloch variety. *Comment. Math. Helv.*, 65(1):114–149, 1990.
- [21] P. Kuchment. An overview of periodic elliptic operators. Bull. Amer. Math. Soc. (N.S.), 53(3):343–414, 2016.
- [22] P. Kuchment and B. Vainberg. On absence of embedded eigenvalues for Schrödinger operators with perturbed periodic potentials. *Comm. Partial Differential Equations*, 25(9-10):1809–1826, 2000.

- [23] P. Kuchment and B. Vainberg. On the structure of eigenfunctions corresponding to embedded eigenvalues of locally perturbed periodic graph operators. *Comm. Math. Phys.*, 268(3):673–686, 2006.
- [24] W. Li and S. P. Shipman. Irreducibility of the Fermi surface for planar periodic graph operators. *Lett. Math. Phys.*, 110(9):2543– 2572, 2020.
- [25] W. Liu. Fermi isospectrality for discrete periodic Schrödinger operators. *Comm. Pure Appl. Math. to appear.*
- [26] W. Liu. Irreducibility of the Fermi variety for discrete periodic Schrödinger operators and embedded eigenvalues. *Geom. Funct. Anal.*, 32(1):1–30, 2022.
- [27] W. Liu. Topics on Fermi varieties of discrete periodic Schrödinger operators. J. Math. Phys., 63(2):Paper No. 023503, 13, 2022.
- [28] H. P. McKean and E. Trubowitz. Hill's operator and hyperelliptic function theory in the presence of infinitely many branch points. *Comm. Pure Appl. Math.*, 29(2):143–226, 1976.
- [29] S. P. Shipman. Eigenfunctions of unbounded support for embedded eigenvalues of locally perturbed periodic graph operators. *Comm. Math. Phys.*, 332(2):605–626, 2014.
- [30] S. P. Shipman. Reducible Fermi surfaces for non-symmetric bilayer quantum-graph operators. J. Spectr. Theory, 10(1):33–72, 2020.
- [31] A. Waters. Isospectral periodic torii in dimension 2. Ann. Inst. H. Poincaré Anal. Non Linéaire, 32(6):1173–1188, 2015.

(W. Liu) Department of Mathematics, Texas A&M University, College Station, TX 77843-3368, USA

Email address: liuwencai1226@gmail.com; wencail@tamu.edu