Every graph with no \mathcal{K}_8^{-4} minor is 7-colorable

Michael Lafferty^{*} Zi-Xia Song[†]

August 22, 2022

Dedicated to the memory of Robin Thomas on his 60th birthday

Abstract

Hadwiger's Conjecture from 1943 states that every graph with no K_t minor is (t-1)colorable; it remains wide open for all $t \geq 7$. For positive integers t and s, let \mathcal{K}_t^{-s} denote the family of graphs obtained from the complete graph K_t by removing s edges. We say that a graph G has no \mathcal{K}_t^{-s} minor if it has no H minor for every $H \in \mathcal{K}_t^{-s}$. Jakobsen in 1971 proved that every graph with no \mathcal{K}_7^{-2} minor is 6-colorable. In this paper we consider the next step and prove that every graph with no \mathcal{K}_8^{-4} minor is 7-colorable. Our result implies that H-Hadwiger's Conjecture, suggested by Paul Seymour in 2017, is true for every graph H on eight vertices such that the complement of H has maximum degree at least four, a perfect matching, a triangle and a cycle of length four. Our proof utilizes an extremal function for \mathcal{K}_8^{-4} minors obtained in this paper, generalized Kempe chains of contraction-critical graphs by Rolek and the second author, and the method for finding K_7 minors from three different K_5 subgraphs by Kawarabayashi and Toft; this method was first developed by Robertson, Seymour and Thomas in 1993 to prove Hadwiger's Conjecture for t = 6.

1 Introduction

All graphs in this paper are finite and simple. For a graph G we use |G|, e(G), $\delta(G)$, $\Delta(G)$, $\alpha(G)$, $\alpha(G)$, $\chi(G)$ to denote the number of vertices, number of edges, minimum degree, maximum degree, independence number, and chromatic number of G, respectively. The complement of G is denoted by \overline{G} . For any positive integer k, we write [k] for the set $\{1, \ldots, k\}$. A graph H is a minor of a graph G if H can be obtained from a subgraph of G by contracting edges. We write $G \succeq H$ if H is a minor of G. In those circumstances we also say that G has an H minor. For positive integers t, s, we use \mathcal{K}_t^{-s} to denote the family of graphs obtained from the complete graph K_t by deleting s edges. When $s \leq 2$, we use K_t^- to denote the unique graph in \mathcal{K}_t^{-1} ; and $K_t^=$ and $K_t^<$ to denote the graphs obtained from K_t by deleting two independent edges and two adjacent edges, respectively. Note that $\mathcal{K}_t^{-2} = \{K_t^=, K_t^<\}$. We say that a graph G has no \mathcal{K}_t^{-s} if G has a \mathcal{K}_t^{-s} minor.

^{*}Department of Mathematics, University of Central Florida, Orlando, FL 32816, USA. Supported by NSF grant DMS-2153945. Email: michaelmlafferty@Knights.ucf.edu.

[†]Department of Mathematics, University of Central Florida, Orlando, FL 32816, USA. Supported by NSF grant DMS-2153945. Email: Zixia.Song@ucf.edu.

Our work is motivated by Hadwiger's Conjecture [11], which is perhaps the most famous conjecture in graph theory.

Conjecture 1.1 (Hadwiger's Conjecture [11]). Every graph with no K_t minor is (t-1)-colorable.

Conjecture 1.1 is trivially true for $t \leq 3$, and reasonably easy for t = 4, as shown independently by Hadwiger [11] and Dirac [6]. However, for $t \ge 5$, Hadwiger's Conjecture implies the Four Color Theorem [2, 3]. Wagner [35] proved that the case t = 5 of Hadwiger's Conjecture is, in fact, equivalent to the Four Color Theorem, and the same was shown for t = 6 by Robertson, Seymour and Thomas [31]. Despite receiving considerable attention over the years, Hadwiger's Conjecture remains wide open for all $t \geq 7$, and is considered among the most important problems in graph theory and has motivated numerous developments in graph coloring and graph minor theory. Proving that graphs with no K_7 minor are 6-colorable is thus the first case of Hadwiger's Conjecture that is still open. It is not even known yet whether every graph with no K_7 minor is 7colorable; Rolek, Thomas and the second author [30] proved that every 8-contraction-critical graph with no K_7 minor has at most one vertex of degree eight. Jakobsen [13] in 1971 proved that every graph with no K_7 minor is 9-colorable. Kawarabayashi and Toft [15] in 2005 proved that every graph with no K_7 or $K_{4,4}$ minor is 6-colorable. Recently, Albar and Gonçalves [1] proved that every graph with no K_7 minor is 8-colorable, and every graph with no K_8 minor is 10-colorable. Their proofs are computer-assisted; Rolek and the second author [29] then gave computer-free proofs of their results, and further proved that every graph with no K_9 minor is 12-colorable, and every graph with no K_t minor is (2t-6)-colorable for all $t \ge 10$ if Conjecture 5.1 in [29] holds.

Until very recently the best known upper bound on the chromatic number of graphs with no K_t minor is $O(t(\log t)^{1/2})$, obtained independently by Kostochka [17, 18] and Thomason [34], while Norin, Postle and the second author [26] improved the frightening $(\log t)^{1/2}$ term to $(\log t)^{1/4}$. The current record is $O(t \log \log t)$ due to Delcourt and Postle [10]. Kühn and Osthus [22] proved that Hadwiger's Conjecture is true for C_4 -free graphs of sufficiently large chromatic number, and for all graphs of girth at least 19. We refer the reader to recent surveys [5, 16, 32] for further background on Hadwiger's Conjecture.

Given the difficulty of Hadwiger's Conjecture, Paul Seymour in 2017 suggested the study of the following H-Hadwiger's Conjecture.

Conjecture 1.2 (*H*-Hadwiger's Conjecture). For every graph *H* on *t* vertices, every graph with no *H* minor is (t-1)-colorable.

It is worth noting that Wagner [36] in 1960 initiated the study of this type of problem and proved that every graph with no K_5^- minor is 4-colorable; Dirac [9] in 1964 proved that every graph with no K_6^- minor is 5-colorable; and Jakobsen [13] in 1971 proved that every graph with no K_7^- minor is 7-colorable. Thus proving graphs with no K_7^- minor is 6-colorable remains open. Rolek and the second author [29] in 2017 proved that every graph with no K_8^- minor is 9-colorable. Very recently, Norin and Seymour [27] proved that every graph on *n* vertices with independence number two has an *H* minor, where *H* is a graph with $\lceil n/2 \rceil$ vertices and at least 0.98688 $\cdot {\binom{|H|}{2}} - o(n^2)$ edges. Woodall [37] studied a special case of *H*-Hadwiger's Conjecture in 2001 by excluding the complete bipartite graph $K_{s,t}$ minor and made the following conjecture.

Conjecture 1.3 (Woodall [37]). Every graph with no $K_{s,t}$ minor is (s + t - 1)-colorable.

In the same paper Woodall confirmed Conjecture 1.3 for $s \in \{1, 2\}$ without using the extremal function for $K_{s,t}$ minors. The s = 2 case also follows from the extremal function for $K_{2,t}$ minors by Myers [25], and Chudnovsky, Reed and Seymour [4]; the s = 3 case when $t \ge 6300$ follows from the extremal function for $K_{3,t}$ minors by Kostochka and Prince [20]. Asymptomatic bounds for the chromatic number of graphs with no $K_{s,t}$ minor, when t is sufficiently larger than s, follow from the extremal functions for $K_{s,t}$ minors by Kühn and Osthus [23], and Kostochka and Prince [19]; in particular, the extremal function by Kostochka and Prince implies that every graph with no $K_{s,t}$ minor is (3s + t - 1)-colorable when t is sufficiently larger than s. Years later Kostochka [21] proved that Conjecture 1.3 is true for $t > C(s \log s)^3$.

Dirac in 1964 began the study of a variation of *H*-Hadwiger's Conjecture in [8] by excluding more than one forbidden minor simultaneously; he proved that every graph with no \mathcal{K}_t^{-2} minor is (t-1)-colorable for each $t \in \{5, 6\}$. Jakobsen [12] in 1971 proved that every graph with no \mathcal{K}_7^{-2} minor is 6-colorable; this implies that *H*-Hadwiger's Conjecture is true for all graphs *H* on seven vertices such that $\Delta(\overline{H}) \geq 2$ and \overline{H} has a matching of size two. Recently, Rolek and the second author [29] proved that every graph with no \mathcal{K}_8^{-2} minor is 8-colorable, and Rolek [28] later proved that every graph with no \mathcal{K}_9^{-2} minor is 10-colorable. Proving that every graph with no \mathcal{K}_8^{-2} minor is 7-colorable is still open. We state the result of Jakobsen [12] below.

Theorem 1.4 (Jakobsen [12]). Every graph with no \mathcal{K}_7^{-2} minor is 6-colorable. In particular, H-Hadwiger's Conjecture is true for all graphs H on seven vertices such that $\Delta(\overline{H}) \geq 2$ and \overline{H} has a matching of size two.

The purpose of this paper is to consider the next step and prove the following main result.

Theorem 1.5. Every graph with no \mathcal{K}_8^{-4} minor is 7-colorable.

There are 11 ways to delete four edges from K_8 , including four edges that form a perfect matching and four edges that are incident with the same vertex. Let H be a graph on eight vertices such that $\Delta(\overline{H}) \geq 4$, and \overline{H} has a perfect matching, a triangle and a cycle of length four. It is simple to check that every graph with no H minor has no \mathcal{K}_8^{-4} minor. Combining this with Theorem 1.5 leads to the observation that H-Hadwiger's Conjecture is true for all such graphs H. In particular, H-Hadwiger's Conjecture is true for five graphs H obtained from K_8 by deleting eight edges, where \overline{H} is given in Figure 1.

Corollary 1.6. *H*-Hadwiger's Conjecture is true for all graphs *H* on eight vertices such that $\Delta(\overline{H}) \geq 4$, and \overline{H} has a perfect matching, a triangle and a cycle of length four.

Our proof of Theorem 1.5 utilizes an extremal function for \mathcal{K}_8^{-4} minors (see Theorem 1.7), generalized Kempe chains of contraction-critical graphs obtained by Rolek and the second author [29]




Figure 1: Five non-isomorphic \overline{H} .

(see Lemma 2.4), and the method for finding K_7 minors from three different K_5 subgraphs by Kawarabayashi and Toft [15] (see Theorem 2.7).

Theorem 1.7. Every graph on $n \ge 8$ vertices with at least 4.5n - 12 edges has a \mathcal{K}_8^{-4} minor.

Theorem 1.7 is best possible in the sense that every $(K_{2,2,2,2}, 4)$ -cockade (see the definition of an (H, k)-cockade in [33]) on n vertices has 4.5n - 12 edges but no \mathcal{K}_8^{-3} minor.

This paper is organized as follows. In the next section, we introduce the necessary definitions and collect several tools which we will need later on. In Section 3, we prove Theorem 1.5. In Section 4, we prove Theorem 1.7 using more involved arguments.

2 Notation and tools

Let G be a graph. If x, y are adjacent vertices of G, then we denote by G/xy the graph obtained from G by contracting the edge xy and deleting all resulting parallel edges. We simply write G/e if e = xy. If u, v are distinct nonadjacent vertices of G, then by G + uv we denote the graph obtained from G by adding an edge with ends u and v. If u, v are adjacent or equal, then we define G + uvto be G. Similarly, if $M \subseteq E(G) \cup E(\overline{G})$, then by G + M we denote the graph obtained from G by adding all the edges of M to G. Every edge in \overline{G} is called a missing edge of G. For a vertex $x \in V(G)$, we will use N(x) to denote the set of vertices in G which are adjacent to x. We define $N[x] = N(x) \cup \{x\}$. The degree of x is denoted by $d_G(x)$ or simply d(x). If $A, B \subseteq V(G)$ are disjoint, we say that A is complete to B if each vertex in A is adjacent to all vertices in B, and A is anticomplete to B if no vertex in A is adjacent to any vertex in B. If $A = \{a\}$, we simply say a is complete to B or a is anticomplete to B. We use e(A, B) to denote the number of edges between A and B in G. The subgraph of G induced by A, denoted by G[A], is the graph with vertex set A and edge set $\{xy \in E(G) \mid x, y \in A\}$. We denote by $B \setminus A$ the set B - A, and $G \setminus A$ the subgraph of G induced on $V(G) \setminus A$, respectively. If $A = \{a\}$, we simply write $B \setminus a$ and $G \setminus a$, respectively. An (A, B)-path in G, when A and B are not necessarily disjoint, is a path with one end in A and the other in B such that all its internal vertices lie in $G \setminus (A \cup B)$. We simply say an (a, B)-path if $A = \{a\}$. It is worth noting that each vertex in $A \cap B$ is an (A, B)-path. For a positive integer k, a k-vertex is a vertex of degree k, and a k-clique is a set of k pairwise adjacent vertices. Let \mathcal{F} be a family of graphs. A graph G is \mathcal{F} -free if it has no subgraph isomorphic to H for every $H \in \mathcal{F}$. We simply say G is H-free if $\mathcal{F} = \{H\}$. The join G + H (resp. union $G \cup H$) of two vertex-disjoint graphs G and H is the graph having vertex set $V(G) \cup V(H)$ and edge set $E(G) \cup E(H) \cup \{xy \mid x \in V(G), y \in V(H)\}$ (resp. $E(G) \cup E(H)$). We use the convention "A :=" to mean that A is defined to be the right-hand side of the relation. Finally, if H is a connected subgraph of a graph G and $y \in V(H)$, we simply say that we contract $H \setminus y$ onto y when we contract H to a single vertex, that is, contract all the edges of H.

To prove Theorem 1.5, we need to investigate the basic properties of contraction-critical graphs. For a positive integer k, a graph G is k-contraction-critical if $\chi(G) = k$ and every proper minor of G is (k-1)-colorable. Lemma 2.1 is a result of Dirac [7].

Lemma 2.1 (Dirac [7]). Let G be a k-contraction-critical graph. Then for each $v \in V(G)$,

$$\alpha(G[N(v)]) \le d(v) - k + 2.$$

We will make use of the following results of Mader [24]. It seems very difficult to improve Theorem 2.3; it remains open whether every k-contraction-critical graph is 8-connected for all $k \ge 8$. Let z_1, z_2, z_3, z_4 be four vertices in a graph G. We say that G contains a K_4 minor rooted at z_1, z_2, z_3, z_4 if there exist $Z_1, Z_2, Z_3, Z_4 \subseteq V(G)$ such that $z_i \in Z_i$ and $G[Z_i]$ is connected for all $i \in [4]$, and Z_i and Z_j are disjoint and there is an edge between Z_i and Z_j in G for $1 \le i < j \le 4$.

Theorem 2.2 (Mader [24]). For all $i \in [4]$, let $Z_i \subseteq V(G)$ with $z_i \in Z_i$ such that $Z_i \cap Z_j = \emptyset$ for $1 \leq i < j \leq 4$, and $\alpha(G[\{z_1, z_2, z_3, z_4\}] \leq 2$. If there exists a (z_i, z_j) -path consisting only of vertices from $Z_i \cup Z_j$ for $1 \leq i < j \leq 4$, then G has a K_4 minor rooted at $\{z_1, z_2, z_3, z_4\}$.

Theorem 2.3 (Mader [24]). For all $k \ge 7$, every k-contraction-critical graph is 7-connected.

Lemma 2.4 on contraction-critical graphs turns out to be very powerful, as the existence of pairwise vertex-disjoint paths is guaranteed without using the connectivity of such graphs. The proof of Lemma 2.4 uses Kempe chains. We recall the proof here as it will be needed in the proof of Lemma 2.5. Recall that every edge in \overline{H} is a missing edge of a graph H.

Lemma 2.4 (Rolek and Song [29]). Let G be any k-contraction-critical graph. Let $x \in V(G)$ be a vertex of degree k+s with $\alpha(G[N(x)]) = s+2$ and let $S \subset N(x)$ with |S| = s+2 be any independent set, where $k \ge 4$ and $s \ge 0$ are integers. Let M be a set of missing edges of $G[N(x) \setminus S]$. Then there exists a collection $\{P_{uv} \mid uv \in M\}$ of paths in G such that for each $uv \in M$, P_{uv} has ends u, v and all its internal vertices in $G \setminus N[x]$. Moreover, if vertices u, v, w, z with $uv, wz \in M$ are distinct, then the paths P_{uv} and P_{wz} are vertex-disjoint.

Proof. Let G, x, S and M be given as in the statement. Let H be obtained from G by contracting $S \cup \{x\}$ to a single vertex, say w. Then H is (k-1)-colorable. Let $c: V(H) \to \{1, 2, \ldots, k-1\}$ be a proper (k-1)-coloring of H. We may assume that c(w) = 1. Then each of the colors $2, \ldots, k-1$ must appear in $G[N(x) \setminus S]$, else we could assign x the missing color and all vertices in S the color 1 to obtain a proper (k-1)-coloring of G, a contradiction. Since $|N(x) \setminus S| = k-2$, we have $c(u) \neq c(v)$ for any two distinct vertices u, v in $G[N(x) \setminus S]$. We next claim that for each $uv \in M$ there must exist a path between u and v with its internal vertices in $G \setminus N[x]$. Suppose not. Let

 H^* be the subgraph of H induced by the vertices colored c(u) or c(v) under the coloring c. Then $V(H^*) \cap N(x) = \{u, v\}$. Notice that u and v must belong to different components of H^* as there is no path between u and v with its internal vertices in $G \setminus N[x]$. By switching the colors on the component of H^* containing u, we obtain a (k-1)-coloring of H with the color c(u) missing on $G[N(x) \setminus S]$, a contradiction. This proves that there must exist a path P_{uv} in H^* with ends u, v and all its internal vertices in $H^* \setminus N[x]$ for each $uv \in M$. Clearly, for any $uv, wz \in M$ with u, v, w, z distinct, the paths P_{uv}, P_{wz} are vertex-disjoint, because no two vertices of u, v, w, z are colored the same under the coloring c.

We shall make frequent use of the following lemma in the proof of Theorem 1.5.

Lemma 2.5. Let G be any k-contraction-critical graph. Let $x \in V(G)$ be a vertex of degree k+s with $\alpha(G[N(x)]) = s+2$ and let $S \subset N(x)$ with |S| = s+2 be any independent set, where $k \ge 4$ and $s \ge 0$ are integers. If $M = \{x_1y_1, x_1y_2, x_2y_1, x_2y_2, a_1b_{11}, \ldots, a_1b_{1r_1}, \ldots, a_mb_{m1}, \ldots, a_mb_{mr_m}\}$ is a set of missing edges of $G[N(x) \setminus S]$, where the vertices $x_1, x_2, y_1, y_2, a_1, \ldots, a_m, b_{11}, \ldots, b_{mr_m} \in N(x) \setminus S$ are all distinct, and for all $1 \le i \le m$, $a_ib_{i1}, \ldots, a_ib_{ir_i}$ are r_i missing edges with a_i as a common end, and $x_1x_2, y_1y_2 \in E(G)$, then $G \succcurlyeq G[N[x]] + M$.

Proof. By Lemma 2.4, there exists a collection $\{P_{uv} \mid uv \in M\}$ of paths in G such that for each $uv \in M$, P_{uv} has ends u, v and all its internal vertices in $G \setminus N[x]$. In particular, for any $1 \leq i < j \leq m$, the paths $P_{a_ib_{i1}}, \ldots, P_{a_ib_{ir_i}}$ are vertex-disjoint from the paths $P_{a_jb_{j1}}, \ldots, P_{a_jb_{jr_j}}$ and $P_{x_1y_1}, P_{x_1y_2}, P_{x_2y_1}, P_{x_2y_2}; P_{x_1y_1}$ and $P_{x_2y_2}$ are vertex-disjoint; $P_{x_1y_2}$ and $P_{x_2y_1}$ are vertex-disjoint but each of $P_{x_1y_2}$ and $P_{x_2y_1}$ is not necessarily vertex-disjoint from $P_{x_1y_1}$ and $P_{x_2y_2}$. By contracting each of $P_{a_ib_{i\ell}} \setminus b_{i\ell}$ onto a_i for all $i \in [m]$ and $\ell \in [r_i]$, we see that $G \succeq G[N[x]] + (M \setminus \{x_1y_1, x_1y_2, x_2y_1, x_2y_2\})$. We next apply Theorem 2.2 to show that $G \succeq G[N[x]] + M$.

Let $z_1 = x_1, z_2 = x_2, z_3 = y_1, z_4 = y_2$. From the proof of Lemma 2.4, each of $P_{x_1y_1}, P_{x_1y_2}, P_{x_2y_1}, P_{x_2y_2}$ is a Kempe chain, and vertices of z_1, z_2, z_3, z_4 are colored differently under the coloring c, where c is given in the proof of Lemma 2.4. For each $i \in [4]$, let Z_i be the set of vertices v of $G[V(P_{x_1y_1}) \cup V(P_{x_1y_2}) \cup V(P_{x_2y_1}) \cup V(P_{x_2y_2})]$ such that $c(v) = c(z_i)$. Then $z_i \in Z_i$ and $Z_i \cap Z_j = \emptyset$ for $1 \leq i < j \leq 4$. Note that $\alpha(G[\{z_1, z_2, z_3, z_4\}] \leq 2$ because $z_1z_2, z_3z_4 \in E(G)$; the (z_1, z_2) -path has only one edge z_1z_2 , the (z_3, z_4) -path has only one edge z_3z_4 , and the (z_i, z_j) -path for each $i \in [2]$ and $j \in \{3, 4\}$ is $P_{x_iy_j}$. By Theorem 2.2, we see that $G[V(P_{x_1y_1}) \cup V(P_{x_1y_2}) \cup V(P_{x_2y_1}) \cup V(P_{x_2y_2})]$ has a K_4 minor rooted at z_1, z_2, z_3, z_4 . It follows that $G \succcurlyeq G[N[x]] + M$, as desired.

Remark 2.6. Lemma 2.5 can be applied when

 $M = \{x_1y_1, x_1y_2, x_2y_1, x_2y_2, a_1b_{11}, \dots, a_1b_{1r_1}, \dots, a_mb_{m1}, \dots, a_mb_{mr_m}\}$

is a subset of edges and missing edges of $G[N(x) \setminus S]$, where $x_1, x_2, y_1, y_2, a_1, \ldots, a_m, b_{11}, \ldots, b_{mr_m} \in N(x) \setminus S$ are all distinct, and $x_1x_2, y_1y_2 \in E(G)$. Under those circumstances, it suffices to apply Lemma 2.5 to M^* , where $M^* = \{e \in M \mid e \text{ is a missing edge of } G[N(x) \setminus S])\}$. It is straightforward to see that $G \succcurlyeq G[N[x]] + M$.




Figure 2: The nine possibilities for three 5-cliques in Theorem 2.7.

Finally we need a tool to find a desired \mathcal{K}_8^{-4} minor through three different 5-cliques in 7connected graphs. This method was first introduced by Robertson, Seymour and Thomas [31] to prove Hadwiger's Conjecture for t = 6: they found a desired \mathcal{K}_6 minor via three different 4-cliques in 6-connected non-apex graphs. This method was later extended by Kawarabayashi and Toft [15] to find a desired \mathcal{K}_7 minor via three different 5-cliques in 7-connected graphs.

Theorem 2.7 (Kawarabayashi and Toft [15]). Let G be a 7-connected graph such that $|G| \ge 19$. If G contains three different 5-cliques, say L_1, L_2, L_3 , such that $|L_1 \cup L_2 \cup L_3| \ge 12$, that is, they fit into one of the nine configurations depicted in Figure 2, then $G \succcurlyeq K_7$. In particular, G has seven pairwise vertex-disjoint "good paths", where a "good path" is an (L_i, L_j) -path in G with $i \ne j$.

It is worth noting that Theorem 2.7 corresponds to [15, Lemma 5], where the existence of such seven "good paths" follows from the proof of [15, Lemma 5]. Theorem 2.7 implies the following:

Corollary 2.8. Let G be a 7-connected graph such that $|G| \ge 19$. If G contains three different 5-cliques, say L_1, L_2, L_3 , such that $|L_1 \cup L_2 \cup L_3| \ge 12$, that is, they fit into one of the nine configurations depicted in Figure 2, then $G \succcurlyeq \mathcal{K}_8^{-4}$. Moreover, if $L_1 \cap L_2 \cap L_3 = \emptyset$, then $G \succcurlyeq \mathcal{K}_8^{-3}$.

Proof. By Theorem 2.7, G has seven pairwise vertex-disjoint "good paths". By choosing such seven

"good paths", say Q_1, \ldots, Q_7 , with $|V(Q_1) \cup \ldots \cup V(Q_7)|$ minimum, we may assume that no internal vertex of each Q_i lies in $L_1 \cup L_2 \cup L_3$. Moreover, if $v \in L_i \cap L_j$ for some $i \neq j$, then v does not belong to any "good path" on at least two vertices. It is then straightforward to verify that each possibility of L_1, L_2, L_3 , given in Figure 2(a-g), together with Q_1, \ldots, Q_7 , yields a \mathcal{K}_8^{-3} minor; and each possibility of L_1, L_2, L_3 , given in Figure 2(h,i), together with Q_1, \ldots, Q_7 , yields a \mathcal{K}_8^{-4} minor, as desired.




Figure 3: The graph H_8 .

Finally we need a lemma of Rolek, Thomas and the second author.

Lemma 2.9 (Rolek, Song and Thomas [30]). Let H be a graph with |H| = 8 and $\alpha(H) = 2$. Then H contains K_4 or H_8 as a subgraph, where H_8 is depicted in Figure 3.

3 Coloring graphs with no \mathcal{K}_8^{-4} minor

We begin this section with two lemmas. Lemma 3.1 below follows directly from Lemma 2.5. We give a proof here without using Theorem 2.2 (its proof was written in German).




Figure 4: Applying Lemma 2.4 to N(x) with $S = \{w_1, w_6\}$ and $M = \{w_2w_5, w_3w_8, w_4w_7, w_3w_7\}$.

Lemma 3.1. Let G be an 8-contraction-critical graph. If there exists $x \in V(G)$ such that G[N(x)] is isomorphic to H_8 , then $G \succeq \mathcal{K}_8^{-3}$.

Proof. Suppose $G[N(x)] = H_8$ for some $x \in V(G)$. Let w_1, \ldots, w_8 be the vertices of N(x) as depicted in Figure 4. Note that $|H_8| = 8$ and $\alpha(H_8) = 2$. By Lemma 2.4 applied to N(x)with $S = \{w_1, w_6\}$ and $M = \{w_2w_5, w_3w_8, w_4w_7, w_3w_7\}$, there exist pairwise vertex-disjoint paths Q_1, Q_2, Q_3 and another path Q_4 such that Q_1, Q_2, Q_3, Q_4 have ends in $\{w_2, w_5\}$, $\{w_3, w_8\}$, $\{w_4, w_7\}$ and $\{w_3, w_7\}$, respectively, and all their internal vertices are in $G \setminus N[x]$. Note that Q_4 and Q_2 may have more than w_3 in common, and Q_4 and Q_3 may have more than w_7 in common; see Figure 4. Let $u \in V(Q_4) \cap V(Q_2)$ such that the (w_7, u) -subpath Q'_4 of Q_4 and Q_2 have exactly u in common; and let $v \in V(Q'_4) \cap V(Q_3)$ such that the (u, v)-subpath Q''_4 of Q'_4 and Q_3 have exactly v in common. Let Q'''_4 denote the (w_7, v) -subpath of Q'_4 . By contracting $Q_1 \setminus w_5$ onto w_2 , both $Q_2 \setminus w_8$ and $Q''_4 \setminus v$ onto w_3 , and both $Q_3 \setminus w_4$ and Q'''_4 onto w_7 , we see that $H_8 + M$ has a \mathcal{K}_7^{-3} minor after contracting the edge $w_6 w_8$. Hence $G \geq G[N[x]] + M \geq \mathcal{K}_8^{-3}$, as desired.

Lemma 3.2. Let G be an 8-contraction-critical graph. If G has a 6-clique A and a 5-clique B such that $B \not\subseteq A$, then $G \succcurlyeq \mathcal{K}_8^{-3}$.

Proof. Let $A := \{a_1, \ldots, a_6\}$ and $B := \{b_1, \ldots, b_5\}$. Let $t := |A \cap B|$. Then $0 \le t \le 4$ because $B \not\subseteq A$. We may further assume that $b_i = a_i$ for each $i \in [t]$ when $t \ne 0$. Assume first t = 4. Then $G[A \cup B]$ contains $K_7^<$ as a subgraph. By Theorem 2.3, G is 7-connected. We obtain a $K_8^<$ minor by contracting a component of $G \setminus (A \cup B)$ to a single vertex, as desired. We may assume that $0 \le t \le 3$. Then there exist 5 - t pairwise vertex-disjoint paths Q_{t+1}, \ldots, Q_5 between $A \setminus B$ and $B \setminus A$ in $G \setminus (A \cap B)$. We may assume that Q_i has ends a_i, b_i for each $i \in \{t + 1, \ldots, 5\}$. Then $G \setminus \{a_1, \ldots, a_5\}$ is connected, so there must exist a path Q with one end a_6 and the other in $V(Q_{t+1} \setminus a_{t+1}) \cup \cdots \cup V(Q_5 \setminus a_5)$, say in $V(Q_5 \setminus a_5)$. We may assume that Q is vertex-disjoint from Q_{t+1}, \ldots, Q_4 by choosing Q to be a shortest such path. Now contracting both $Q_5 \setminus a_5$ and $Q \setminus a_6$ onto $b_5, Q_4 \setminus a_4$ onto b_4 , and all the edges of Q_j for each $j \in \{t+1, \ldots, 3\}$ when $t \le 2$, we see that $G \succcurlyeq G[A \cup B] + \{a_{t+1}b_{t+1}, \ldots, a_5b_5, a_6b_5\} \succcurlyeq K_8^{-3}$.

We are now ready to prove Theorem 1.5, which we restate for convenience.

Theorem 1.5. Every graph with no \mathcal{K}_8^{-4} minor is 7-colorable.

Proof. Suppose the assertion is false. Let G be a graph with no \mathcal{K}_8^{-4} minor such that $\chi(G) \geq 8$. We may choose such a graph G so that it is 8-contraction-critical. Then $\delta(G) \geq 7$, G is 7-connected by Theorem 2.3, and $\delta(G) \leq 8$ by Theorem 1.7. Let $x \in V(G)$ be of minimum degree. Since G is 8-contraction-critical and has no \mathcal{K}_8^{-4} minor, by Lemma 2.1 applied to G[N(x)], we see that

(1)
$$\delta(G) = 8$$
 and $\alpha(G[N(x)]) = 2$.

(2)
$$G$$
 is \mathcal{K}_7^{-4} -free.

Proof. Suppose G contains an $H \in \mathcal{K}_7^{-4}$ as a subgraph. Since G is 7-connected, we obtain a \mathcal{K}_8^{-4} minor of G by contracting a component of $G \setminus V(H)$ to a single vertex, a contradiction.

(3) G[N(x)] contains a 4-clique.

Proof. Suppose G[N(x)] is K_4 -free. By Lemma 2.9, G[N(x)] contains H_8 as a subgraph. Let the vertices of H_8 be labeled as in Figure 4. Then $w_1w_6 \notin E(G)$ because G[N(x)] is K_4 -free. We consider the worst scenario that $G[N(x)] = H_8$. By Lemma 3.1, $G \succeq \mathcal{K}_8^{-3}$, a contradiction.

(4) $n_8 \ge 25$, where n_8 denotes the number of vertices of degree eight in G.

Proof. Suppose $|n_8| \le 24$. Then $e(G) \ge (8n_8 + 9(|G| - n_8))/2 \ge (9|G| - 24)/2$. By Theorem 1.7, $G \succcurlyeq \mathcal{K}_8^{-4}$, a contradiction. □

(5) G is K_6 -free.

Proof. Suppose G has a 6-clique A. By (4), $n_8 \ge 25$. Let $y \in V(G) \setminus A$ be an 8-vertex in G. By (3), N[y] has a 5-clique B such that $y \in B$. Then $B \not\subseteq A$. By Lemma 3.2, $G \succcurlyeq \mathcal{K}_8^{-3}$, a contradiction. \Box

(6) G[N(x)] has two disjoint 4-cliques.

Proof. Suppose *G*[*N*(*x*)] does not have two disjoint 4-cliques. By (5), *G*[*N*(*x*)] is *K*₅-free. Then $\delta(G[N(x)]) \geq 3$ because $\alpha(G[N(x)]) = 2$. We claim that $\delta(G[N(x)]) \geq 4$. Suppose not. Let $y \in N(x)$ such that *y* has exactly three neighbors $y_1, y_2, y_3 \in N(x)$. Let $Y := \{y, y_1, y_2, y_3\}$ and $Z := \{z_1, z_2, z_3, z_4\} = N(x) \setminus Y$. Then *Z* is a 4-clique in *G* and *Y* is not a clique. We may assume that $y_1y_2 \notin E(G)$. Note that $e(y_i, Z) \geq 1$ for each $i \in [2]$. We may assume that $y_1z_1 \in E(G)$. Suppose y_1 is anticomplete to $\{z_2, z_3, z_4\}$. Then $y_1y_3 \in E(G)$ and y_2 must be complete to $\{z_2, z_3, z_4\}$. By Lemma 2.5 applied to *N*(*x*) with $S = \{y, z_4\}$ and $M = \{y_1z_2, y_1z_3, y_3z_2, y_3z_3, y_2z_1\}$, we obtain a \mathcal{K}_8^{-3} minor from G[N[x]] + M after contracting the edge yy_2 , a contradiction. Thus $e(y_1, Z) \geq 2$. Similarly, $e(y_2, Z) \geq 2$. Note that $e(\{y_1, y_2\}, Z) \leq 5$, else $G[Z \cup \{x, y_1, y_2\}]$ is not \mathcal{K}_7^{-3} -free, contrary to (2). We may assume that $e(y_1, Z) = 2$ and $y_1z_2 \in E(G)$. Then y_1 is anticomplete to $\{z_3, z_4\}$. By Lemma 2.5 applied to $\{z_3, z_4\}$. By Lemma 2.5 applied to $\{z_3, z_4\}$. By Lemma 2.5 applied to $\{z_3, z_4\}$. We obtain a \mathcal{K}_8^{-3} minor from G[N[x]] + M after contracting the edge yy_2 , a contradiction. Thus $e(y_1, Z) \geq 2$. Similarly, $e(y_2, Z) \geq 2$. Note that $e(\{y_1, Z_2\}, Z_2) \leq 5$, else $G[Z \cup \{x, y_1, y_2\}]$ is not \mathcal{K}_7^{-3} -free, contrary to (2). We may assume that $e(y_1, Z) = 2$ and $y_1z_2 \in E(G)$. Then y_1 is anticomplete to $\{z_3, z_4\}$. By Lemma 2.5 applied to N(x) with $S = \{y, z_1\}$ and $M = \{y_1z_3, y_1z_4, y_3z_3, y_3z_4, y_2z_2\}$, we obtain a \mathcal{K}_8^{-4} minor from G[N[x]] + M after contracting the edge yy_1 , a contradiction. This proves that $\delta(G[N(x)]) \geq 4$, as claimed.

By (3), G[N(x)] has a 4-clique A. Let $N(x) = \{a_1, \ldots, a_4, b_1, \ldots, b_4\}$, where $A = \{a_1, \ldots, a_4\}$ is a 4-clique and $B := \{b_1, \ldots, b_4\}$ is not. We may assume that $b_1b_2 \notin E(G)$. Since G[N(x)] is K_5 -free and $\alpha(G[N(x)]) = 2$, we may further assume that $b_1a_1, b_2a_2 \notin E(G)$. Then $b_1a_2, b_2a_1 \in E(G)$. Note that $e(b_i, A) \ge 2$ for each $i \in [2]$ because $\delta(G[N(x)]) \ge 4$. On the other hand, $e(\{b_1, b_2\}, A) \le 5$, else $G[A \cup \{x, b_1, b_2\}]$ is not \mathcal{K}_7^{-3} -free, contrary to (2). We may assume that $e(b_1, A) = 2$ and $2 \le e(b_2, A) \le 3$. We may further assume that $b_1a_3, b_2b_3 \in E(G)$. Then $b_1a_4 \notin E(G)$. It follows that $b_2a_4 \in E(G)$ and b_1 is complete to $\{b_3, b_4\}$. By Lemma 2.5 applied to N(x) with $S = \{b_1, a_4\}$ and $M = \{b_2a_2, b_2a_3, b_3a_2, b_3a_3, b_4a_1\}$, we obtain a \mathcal{K}_8^{-4} minor from G[N[x]] + M after contracting the edge b_1b_4 , a contradiction.

To complete the proof, recall that x is an 8-vertex in G. Since $n_8 \ge 25$ by (4), let $y \in V(G) \setminus N[x]$ such that y is an 8-vertex in G. By (6), let A, B be disjoint 4-cliques of G[N(x)] and let C be a 4-clique of G[N(y)]. Let $L_1 := A \cup \{x\}$, $L_2 := B \cup \{x\}$ and $L_3 := C \cup \{y\}$. Then $L_1 \cap L_2 = \{x\}$ and $L_1 \cap L_2 \cap L_3 = \emptyset$. By Corollary 2.8, L_1, L_2, L_3 are not as depicted in Figure 2(c,d,f,g) because G has no \mathcal{K}_8^{-4} minor. We may assume that $|L_1 \cap L_3| \ge |L_2 \cap L_3|$. Then $|L_1 \cap L_3| \ge 2$, else L_1, L_2, L_3 are as depicted in Figure 2(c,d,g).

Let $L_1 := \{x, x_1, \ldots, x_4\}$, $L_2 := \{x, y_1, \ldots, y_4\}$ and $L_3 := \{y, z_1, \ldots, z_4\}$. Suppose first $|L_1 \cap L_3| = 4$. We may assume that $x_i = z_i$ for each $i \in [4]$. Note that $G[L_1 \cup L_3] = K_6^-$. By Menger's Theorem, there exist four pairwise vertex-disjoint $(\{z_1, \ldots, z_4\}, \{y_1, \ldots, y_4\})$ -paths, say Q_1, \ldots, Q_4 , in $G \setminus \{x, y\}$. We may assume that Q_j has ends y_j, z_j for each $j \in [4]$. Let Q be a shortest path in $G \setminus \{x, y, y_1, y_2, y_3, z_4\}$ with one end y_4 and the other in $V(Q_1 \setminus y_1) \cup V(Q_2 \setminus y_2) \cup V(Q_3 \setminus y_3)$. We may assume that Q is vertex-disjoint from Q_1 and Q_2 but contains a vertex on $Q_3 \setminus y_3$. Let $u \in V(Q) \cap V(Q_3)$ and $v \in V(Q) \cap V(Q_4)$ such that the (u, v)-subpath of Q has no internal vertex in $V(Q_3) \cup V(Q_4)$. Let Q_3^* be the (u, y_3) -subpath of Q_3 and Q_4^* be the (v, z_4) -subpath of Q_4 . Then $G \succcurlyeq G[L_1 \cup L_2 \cup L_3] + \{y_1z_1, y_2z_2, y_3z_3, y_4z_3, y_4z_4\} \succcurlyeq K_8^{-4}$ by first contracting all the edges of Q_1, Q_2 , then the (y_4, v) -subpath of Q_4 onto $y_4, Q_4^* \setminus v$ onto $z_4, Q_3^* \setminus u$ onto y_3 , and both $Q \setminus v$ and the (u, z_3) -subpath of Q_3 onto z_3 , a contradiction. Suppose next $|L_1 \cap L_3| = 3$. Then $G[L_1 \cup L_3]$ is not K_7^{-4} -free, contrary to (2). Thus $|L_1 \cap L_3| \leq 2$. Note that $|L_2 \cap L_3| \neq 0$, else L_1, L_2, L_3 are as depicted in Figure 2(f). Thus $|L_1 \cap L_3| = 2$ and $1 \leq |L_2 \cap L_3| \leq 2$. We may assume that $x_3 = z_1, x_4 = z_2$.

We first consider the case $|L_2 \cap L_3| = 2$. We may assume that $y_3 = z_3, y_4 = z_4$. Then there exist two vertex-disjoint $(\{x_1, x_2\}, \{y_1, y_2\})$ -paths, say Q_1, Q_2 , in $G \setminus \{x, x_3, x_4, y_3, y_4\}$. We may assume that Q_j has ends x_j, y_j for each $j \in [2]$. We may further assume $y \notin V(Q_1)$. Let Q_2^* be the (y_2, y) subpath of Q_2 when $y \in V(Q_2)$. Then $G \succcurlyeq G[L_1 \cup L_2 \cup L_3] + \{x_1y_1, x_2y_2\} \succcurlyeq \mathcal{K}_8^{-3}$ by contracting all the edges of Q_1 and Q_2 when $y \notin V(Q_2)$; and $G \succcurlyeq G[L_1 \cup L_2 \cup L_3] + \{x_1y_1, x_2y, y_2y\} \succcurlyeq \mathcal{K}_8^{-2}$ by first contracting all the edges of Q_1 , then $Q_2^* \setminus y$ onto y_2 , and the (y, x_2) -subpath of Q_2 onto x_2 when $y \in V(Q_2)$, a contradiction.

It remains to consider the case $|L_2 \cap L_3| = 1$. We may assume that $z_3 = y_4$. Then there exist three vertex-disjoint $(\{x_1, x_2, z_4, y\}, \{y_1, y_2, y_3\})$ -paths, say Q_1, Q_2, Q_3 , in $G \setminus \{x, x_3, x_4, y_4\}$. We may assume that Q_1 has ends $\{x_1, y_1\}$. Note that $L_3 \setminus (L_1 \cup L_2) = \{y, z_4\}$. By symmetry, we may further assume that Q_3 has ends y_3, z_4 . Then Q_2 has ends y_2, x_2 , or y_2, y . By contracting all the edges of Q_1, Q_2 and Q_3 , we see that $G \succcurlyeq G[L_1 \cup L_2 \cup L_3] + \{x_1y_1, x_2y_2, y_3z_4\} \succcurlyeq \mathcal{K}_8^{-3}$ in the former case, and $G \succcurlyeq G[L_1 \cup L_2 \cup L_3] + \{x_1y_1, y_2y, y_3z_4\} \succcurlyeq \mathcal{K}_8^{-3}$ in the latter case, a contradiction.

This completes the proof of Theorem 1.5.

4 An extremal function for \mathcal{K}_8^{-4} minors

Throughout this section, if G is a graph and K is a subgraph of G, then by N(K) we denote the set of vertices of $V(G) \setminus V(K)$ that are adjacent to a vertex of K. If $V(K) = \{x\}$, then N(K) = N(x). It can be easily checked that for each vertex $x \in V(G)$, if K is a component of $G \setminus N[x]$, then N(K)is a minimal separating set of G.

We first give a brief outline of the proof of Theorem 1.7. We follow the main ideas in [33].

Suppose for a contradiction that G is a counterexample to Theorem 1.7 with as few vertices as possible. Since deletion or contraction of edges does not produce smaller counterexamples, it follows easily that G has minimum degree at least five, and every edge of G belongs to at least four triangles. With some effort it can be shown that G is 5-connected, and has at most one 5-vertex but no 6-vertex and no 7-vertex. As e(G) = 4.5|G| - 12, we see that G has an 8-vertex. Fix such a vertex x. We then show that x is not adjacent to a 5-vertex in G. If $G \setminus N[x]$ has a component K such that $M \subseteq N(K)$, where M is the set of all vertices of N(x) that are not adjacent to every other vertex of N(x), then we can find a vertex $y \in N(x)$ such that $G[N(x) \setminus y] \succeq \mathcal{K}_6^{-4}$, and so $G \succeq \mathcal{K}_8^{-4}$ by contracting the connected graph $G[V(K) \cup \{y\}]$ to a single vertex. Thus we may assume that for no 8-vertex x such a component exists. In particular, $G \setminus N[x]$ is disconnected. In the next step we prove that there is no component K of $G \setminus N[x]$ with $|K| \ge 2$ such that $d_G(v) \ge 9$ for all vertices $v \in V(K)$, except possibly one. In the last step, we select an 8-vertex $x \in V(G)$ to minimize the size of a component K of $G \setminus N[x]$ with $|K| \ge 2$. It follows easily that K does not have a vertex that is an 8-vertex in G.

We next prove two lemmas that will be needed in the proof of Theorem 1.7.

Lemma 4.1. Let H be a graph with |H| = 8 and $\delta(H) \ge 4$. Then there exists $x \in V(H)$ such that $H \setminus x$ has a \mathcal{K}_6^{-4} minor.

Proof. We may assume that $\delta(H) = 4$ and every edge is incident with a 4-vertex in H. Suppose $H \setminus x$ has no \mathcal{K}_6^{-4} minor for every x in H. Suppose first that H has a 4-clique, say $A := \{a_1, \ldots, a_4\}$. Let $B := V(H) \setminus A$. By the minimality of e(H), we may assume that a_i is a 4-vertex in H for each $i \in [3]$. Let $b_i \in B$ be the unique neighbor of a_i in B for each $i \in [3]$. Note that b_1, b_2, b_3 are not necessarily distinct. We claim that for all $a \in A$ and $b \in B$, if $ab \in E(H)$, then a and b have at least one common neighbor. Suppose not. We may assume that $a \neq a_1$. Let $H^* := H \setminus a_1$. Then $e(H^*/ab) = e(H^*) - 1 = (e(H) - 4) - 1 \ge (16 - 4) - 1 = 11 = e(K_6) - 4$. Thus $H \setminus a_1$ has a \mathcal{K}_6^{-4} minor, a contradiction. This implies that $b_i a_4 \in E(H)$ for each $i \in [3]$. Note that $e(b, A) \ge 1$ for each $b \in B$. It follows that a_4 is complete to B and so $d(a_4) = 7$. By the minimality of e(H), we see that each vertex in B is a 4-vertex in H, which is impossible. This proves that H is K_4 -free. Suppose next $\alpha(H) = 2$. By Lemma 2.9, H contains H_8 as a subgraph. Let the vertices of H be labeled as in Figure 3. It is simple to check that $H_8 \setminus w_2$ has a \mathcal{K}_6^{-4} minor after contracting the edge w_1w_3 . Thus H is K_4 -free and $\alpha(H) \ge 3$.

It is straightforward to check that $H = K_{4,4}$ when $\alpha(H) = 4$; and $K_{4,4} \setminus x$ has a \mathcal{K}_6^{-4} minor for every x in $K_{4,4}$. Thus $\alpha(H) = 3$. Let $S := \{x_1, x_2, x_3\}$ be an independent set of H and let $V(H) \setminus S := \{y_1, \ldots, y_5\}$. Since $d(x_i) \ge 4$ for each $i \in [3]$, we may assume that $\{y_1, y_2\}$ is complete to S. Suppose y_3 is complete to S. Since $K_{3,4}$ has a \mathcal{K}_6^{-4} minor, we see that y_4 is neither complete to S nor complete to $\{y_1, y_2, y_3\}$. We may assume that $y_4x_1 \notin E(H)$ and $y_4y_1 \notin E(H)$. Then $H \setminus y_5$ has a \mathcal{K}_6^{-4} minor after contracting the edge x_1y_1 , a contradiction. This proves that no y_j is complete to S for each $j \in \{3, 4, 5\}$. It follows that $d(x_i) = 4$, and $e(y_j, S) = 2$ for each $i \in [3]$ and $j \in \{3, 4, 5\}$; in addition, we may assume that y_3 is complete to $\{x_1, x_2\}, y_4$ is complete to $\{x_2, x_3\}$ and y_5 is complete to $\{x_1, x_3\}$. If $H[\{y_3, y_4, y_5\}] = K_3$, then $H \setminus y_2$ has a \mathcal{K}_6^{-4} minor after contracting the edge x_1y_1 , a contradiction. Thus we may assume that $y_3y_5 \notin E(G)$ and $y_3y_2 \in E(G)$. Then either $y_3y_4 \in E(G)$ or $y_3y_1 \in E(G)$. Thus $H \setminus y_5$ has a \mathcal{K}_6^{-4} minor after contracting the edge x_3y_4 , a contradiction.

Lemma 4.2 follows from the proof of Lemma 16 of Jørgensen [14]. We recall the proof here for convenience.

Lemma 4.2 (Jørgensen [14]). Let G be a 4-connected graph and let $S \subseteq V(G)$ be a separating set of four vertices. Let G_1 and G_2 be proper subgraphs of G so that $G_1 \cup G_2 = G$ and $G_1 \cap G_2 = G[S]$. Let d_1 be the maximum number of edges that can be added to G_2 by contracting edges of G with at least one end in G_1 . If $|G_1| \ge 6$, then

$$e(G[S]) + d_1 \ge 5.$$

Proof. Let $x \in V(G_1) \setminus S$. Then there exist four pairwise internally vertex-disjoint (x, S)-paths, say Q_1, \ldots, Q_4 , in G_1 . For each $i \in [4]$, let s_i be the vertex in $V(Q_i) \cap S$. If all four of these paths have length one, then, since $|G_1| \ge 6$, we may choose a vertex $y \in V(G_1) \setminus (S \cup \{x\})$. Then there are at least three internally vertex-disjoint (y, S)-paths in $G_1 \setminus \{x\}$. Contracting some of these paths results in S having at least five edges, as desired.

We may now assume that Q_1 has length at least two. Since $\{x, s_1\}$ is not a separating set, there is a path Q from a vertex on $Q_1 \setminus \{x, s_1\}$ to a vertex on $Q_j \setminus x$ for some $j \in \{2, 3, 4\}$, so that only the ends of Q belong to $V(Q_1) \cup V(Q_2) \cup V(Q_3) \cup V(Q_4)$. We may assume that j = 2. Since $\{x, s_1, s_2\}$ does not separate the graph, there is a path R from $V(Q_1 \setminus \{x, s_1\}) \cup V(Q_2 \setminus s_2) \cup V(Q)$ to $Q_\ell \setminus x$ for some $\ell \in \{3, 4\}$, so that only the ends of R belong to $V(Q_1) \cup V(Q_2) \cup V(Q_3) \cup V(Q_4) \cup V(Q)$. The result follows from the existence of these paths.

We are now ready to prove Theorem 1.7, which we restate for convenience.

Theorem 1.7. Every graph on $n \ge 8$ vertices with at least 4.5n - 12 edges has a \mathcal{K}_8^{-4} minor.

Proof. Suppose the assertion is false. Let G be a graph on $n \ge 8$ vertices with $e(G) \ge 4.5n - 12$ and, subject to this, n is minimum. We may assume that $e(G) = \lfloor 4.5n - 12 \rfloor$. It is simple to check that $G \succcurlyeq \mathcal{K}_8^{-4}$ for each $n \in \{8, 9\}$. Thus $n \ge 10$. We next prove several claims.

Claim 1. $\delta(G) \ge 5$.

Proof. Suppose $\delta(G) \leq 4$. Let $x \in V(G)$ with $d(x) \leq 4$. Then

$$e(G \setminus x) = e(G) - d(x) \ge \lfloor 4.5n - 12 \rfloor - 4 > \lfloor 4.5 | G \setminus x | - 12 \rfloor.$$

Thus $G \setminus x$ has a \mathcal{K}_8^{-4} minor by the minimality of G, a contradiction.

Claim 2. Every edge in G belongs to at least 4 triangles. Moreover, if $x \in V(G)$ is a 5-vertex, then $G[N[x]] = K_6$.

Proof. Suppose there exists $e \in E(G)$ such that e belongs to at most three triangles. Then

$$e(G/e) \ge \lceil 4.5n - 12 \rceil - 4 > \lceil 4.5 | G/e | - 12 \rceil.$$

Thus $G/e \geq \mathcal{K}_8^{-4}$ by the minimality of G, a contradiction. This implies that $G[N[x]] = K_6$ for each 5-vertex x in G.

Claim 3. No two 5-vertices in G are adjacent.

Proof. Suppose there exist two distinct 5 vertices, say x, y, in G such that $xy \in E(G)$. Then $|G \setminus \{x, y\}| = n - 2 \ge 8$ and

$$e(G \setminus \{x, y\}) = e(G) - 9 = \lceil 4.5(n-2) - 12 \rceil.$$

Thus $G \setminus \{x, y\}$ has a \mathcal{K}_8^{-4} by the minimality of G, a contradiction.

Let S be a minimal separating set of vertices in G, and let G_1 and G_2 be proper subgraphs of G so that $G = G_1 \cup G_2$ and $G_1 \cap G_2 = G[S]$. For each $i \in [2]$, let d_i be the maximum number of edges that can be added to G_{3-i} by contracting edges of G with at least one end in G_i . More precisely, let d_i be the largest integer so that G_i contains pairwise disjoint sets of vertices V_1, \ldots, V_p so that $G_i[V_j]$ is connected, $|S \cap V_j| = 1$ for $1 \leq j \leq p := |S|$, and so that the graph obtained from G_i by contracting each of $G_i[V_1], \ldots, G_i[V_p]$ to a single vertex and deleting $V(G) \setminus \bigcup_{j=1}^p V_j$ has $e(G[S]) + d_i$ edges. It follows from the minimality of G that for each $i \in [2]$,

$$e(G_i) + d_{3-i} < 4.5|G_1| - 12 \text{ if } |G_i| \ge 8.$$
 (*)

Claim 4. If $|G_i| = 7$ for some $i \in [2]$, then $|S| \le 5$, G_i contains K_7^- or $K_7^=$ as a spanning subgraph, and each missing edge of G_i lies in G[S].

Proof. Suppose, say, $|G_1| = 7$. Let C be a component of $G_2 \setminus S$. We first prove that $G_1 \setminus S$ is connected. Suppose not. By Claims 1, 2 and 3, $G_1 \setminus S$ must contain two nonadjacent 5-vertices in G with $G[S] = K_5$. Thus $G_1 = K_7^-$ and so $G \succeq \mathcal{K}_8^{-3}$ by contracting C to a single vertex, a contradiction. Thus $G_1 \setminus S$ is connected, and so $G_1 \setminus S$ has least one 6-vertex, say x, in G. Suppose next |S| = 6. Then G[S] = G[N(x)] contains $K_{2,2,2}$ as a spanning subgraph by Claim 2. But then $G \succeq \mathcal{K}_8^{-4}$ by contracting C to a single vertex, a contradiction. This proves that $|S| \leq 5$ and so $|G_1 \setminus S| \geq 2$.

We next show that no vertex in $G_1 \setminus S$ is a 5-vertex in G. Suppose not. Let $y \in V(G_1 \setminus S)$ be a 5-vertex in G. Then $xy \in E(G)$ and $G_1 = K_7^-$ by Claim 2. Note that G[S] is a complete subgraph because $G[N(y)] = K_5$ and G[N(x)] contains $K_{2,2,2}$ as a subgraph by Claim 2. Then $|S| \leq 3$, else $G \succcurlyeq \mathcal{K}_8^{-4}$ by contracting C to a single vertex. Suppose $|G_2| \geq 8$. Then

$$e(G_2) = e(G) - e(G_1) + e(G[S])$$

= $\lceil 4.5n - 12 \rceil - 20 + {|S| \choose 2}$
= $\lceil (4.5 \times (n - (7 - |S|)) - 12) + 4.5 \times (7 - |S|) \rceil - 20 + {|S| \choose 2}$
 $\ge \lceil 4.5|G_2| - 12 \rceil$

because $|S| \leq 3$. By the minimality of G, we see that $G_2 \succeq \mathcal{K}_8^{-4}$, a contradiction. Thus $|G_2| \leq 7$. If $|G_2| = 6$, then $G_2 = K_6$ and n = 7 + 6 - |S| = 13 - |S|; but then

$$\lceil 4.5 \times (13 - |S|) - 12 \rceil = e(G) = e(G_1) + e(G_2) - e(G[S]) = 20 + 15 - \binom{|S|}{2},$$

which is impossible because $|S| \leq 3$. Thus $|G_2| = 7$ and so n = 7 + 7 - |S| = 14 - |S|. Using a similar argument for G_1 , we see that $G_2 \setminus S$ is connected and so $G_2 = K_7$ or K_7^- . Note that $G_2 = K_7^-$ when |S| = 3, else $G \succcurlyeq \mathcal{K}_8^{-4}$ by contracting $G_1 \setminus S$ to a single vertex. But then

$$\lceil 4.5 \times (14 - |S|) - 12 \rceil = e(G) = e(G_1) + e(G_2) - e(G[S]) \le 20 + (21 - \max\{0, |S| - 2\}) - {|S| \choose 2},$$

which is impossible because $|S| \leq 3$. This proves that no vertex in $G_1 \setminus S$ is a 5-vertex in G. It then follows that each vertex in $G_1 \setminus S$ is a 6-vertex in G. By Claim 2 and the fact that $|S| \leq 5$, we see that G[S] is isomorphic to $K_{|S|}$, $K_{|S|}^-$, or $K_{|S|}^=$, and every vertex in $V(G_1 \setminus S)$ is adjacent to all the other vertices in G_1 . Thus G_1 contains K_7^- or $K_7^=$ as a spanning subgraph such that all its missing edges lie in G[S].

Claim 5. Neither G_1 nor G_2 has exactly seven vertices.

Proof. Suppose not, say $|G_1| = 7$. By Claim 4, $|S| \leq 5$, G_1 contains K_7^- or $K_7^=$ as a spanning subgraph, and each missing edge of G_1 lies in G[S]. We next prove that $|G_2| \geq 8$. Suppose $|G_2| \leq 7$. Note that $|G_2| \geq 6$ by Claim 1. Suppose $|G_2| = 6$. Then $G_2 = K_6$ and $G[S] = K_5$. But then $n = |G_1| + |G_2| - |S| = 7 + 6 - 5 < 10$, a contradiction. Thus $|G_2| = 7$ and so n = 14 - |S|. Since $n \geq 10$, we see that $|S| \leq 4$. By Claim 4, G_2 contains K_7^- or $K_7^=$ as a spanning subgraph, and each missing edge of G_2 lies in G[S]. Suppose |S| = 4. Then $G[S] = K_4^-$, else $G \succeq \mathcal{K}_8^{-4}$ by contracting $G_2 \setminus S$ to a single vertex. Let $y \in V(G_2) \setminus S$ and $z \in S$ such that z is incident with a missing edge in G[S]. It is simple to check that $G \succeq \mathcal{K}_8^{-4}$ by contracting the edge yz and $G_2 \setminus (S \cup \{y\})$ to a single vertex, a contradiction. Thus $|S| \leq 3$. As noted in the proof of Claim 4, we see that $G_1 = G_2 = K_7^-$ when |S| = 3. But then

$$\lceil 4.5 \times (14 - |S|) - 12 \rceil = e(G) = e(G_1) + e(G_2) - e(G[S]) \le 2 \times (21 - \max\{0, |S| - 2\}) - {|S| \choose 2},$$

which is impossible because $|S| \leq 3$. This proves that $|G_2| \geq 8$.

Recall that G_1 contains K_7^- or $K_7^=$ as a spanning subgraph, and each missing edge of G_1 lies in G[S]. It follows that $e(G[S]) + d_1 = {|S| \choose 2}$. Suppose $G_1 = K_7^=$. Then $4 \le |S| \le 5$. But then

$$e(G_2) + d_1 = e(G) - e(G_1) + e(G[S]) + d_1$$

= $\lceil 4.5n - 12 \rceil - 19 + {|S| \choose 2}$
= $\lceil (4.5 \times (n - (7 - |S|)) - 12) + 4.5 \times (7 - |S|) \rceil - 19 + {|S| \choose 2}$
 $\ge \lceil 4.5 |G_2| - 12 \rceil,$

contrary to (*) because $4 \leq |S| \leq 5$ and $|G_2| \geq 8$. Thus $G_1 = K_7^-$ or K_7 . Note that $|S| \leq 3$, else $G \geq \mathcal{K}_8^{-4}$ by contracting a component of $G_2 \setminus S$ to a single vertex. But then

$$e(G_2) + d_1 = e(G) - e(G_1) + e(G[S]) + d_1$$

$$\geq \lceil 4.5n - 12 \rceil - 21 + \binom{|S|}{2}$$

$$= \lceil (4.5 \times (n - (7 - |S|)) - 12) + 4.5 \times (7 - |S|) \rceil - 21 + \binom{|S|}{2}$$

$$\geq \lceil 4.5 |G_2| - 12 \rceil,$$

contrary to (*) because $|S| \leq 3$ and $|G_2| \geq 8$. This proves Claim 5.

Observe that, if $|G_1| \ge 8$ and $|G_2| \ge 8$, then by (*), we have

$$4.5n - 12 \le e(G) = e(G_1) + e(G_2) - e(G[S])$$

$$< (4.5|G_1| - 12 - d_2) + (4.5|G_2| - 12 - d_1) - e(G[S])$$

$$= 4.5(n + |S|) - 24 - d_1 - d_2 - e(G[S]).$$

It follows that

$$9|S| > 24 + 2(d_1 + d_2 + e(G[S]))$$
 if $|G_1| \ge 8$ and $|G_2| \ge 8$. (**)

Claim 6. G is 5-connected.

Proof. Suppose G is not 5-connected. Let S be a minimal separating set of G, and G_1, G_2, d_1, d_2 be defined as above. Then $|G_1| \neq 6$, else by Claim 2, we have $G_1 = K_6$ and so $G[S] = K_5$, a contradiction. Similarly, $|G_2| \neq 6$. By Claim 5, $|G_1| \geq 8$ and $|G_2| \geq 8$. By (**), $|S| \geq 3$. If |S| = 3, then either $e(G[S]) \geq 2$ or $\min\{d_1, d_2\} \geq 1$; in either case, $d_1 + d_2 + e(G[S]) \geq 2$, contrary to (**). Therefore, |S| = 4 and G is 4-connected. By Lemma 4.2, $e(G[S]) + d_1 \geq 5$. Note that $d_2 \geq 1$ when S is not a 4-clique, and e(G[S]) = 6 when S is a 4-clique. In either case, we have $d_1 + d_2 + e(G[S]) \geq 6$, contrary to (**).

Claim 7. If there exists $x \in S$ such that $S \setminus x$ is a clique, then $G[S] = K_5$.

Proof. Suppose $S \setminus x$ is a clique but $G[S] \neq K_5$. Let G_1 and G_2 be as above. Then $|G_1| \neq 6$, else $G_1 = K_6$ and $G[S] = K_5$. Similarly, $|G_2| \neq 6$. By Claim 5, $|G_1| \geq 8$ and $|G_2| \geq 8$. By Claim 6, $|S| \geq 5$. If S contains a 6-clique, then $G \succcurlyeq K_8^-$ by contracting a component of $G_1 \setminus S$ and a component of $G_2 \setminus S$ to two distinct vertices, a contradiction. Thus $5 \leq |S| \leq 6$ and S is not a clique. Then $\delta(G[S]) = d_{G[S]}(x) \leq |S| - 2$. Since $S \setminus x$ is a clique, we see that

$$d_1 = d_2 = |S| - 1 - d_{G[S]}(x) = |S| - 1 - \delta(G[S]).$$

It follows that

$$e(G[S]) = {|S| - 1 \choose 2} + d_{G[S]}(x) = {|S| - 1 \choose 2} + \delta(G[S]).$$

This, together with (**), implies that

$$\begin{aligned} 9|S| &> 24 + 2(d_1 + d_2) + 2e(G[S]) \\ &= 24 + 4|S| - 4 - 4\delta(G[S]) + (|S| - 1)(|S| - 2) + 2\delta(G[S]) \\ &= 20 + 4|S| + |S|^2 - 3|S| + 2 - 2\delta(G[S]) \\ &\geq 20 + |S|^2 + |S| + 2 - 2(|S| - 2) \\ &= |S|^2 - |S| + 26, \end{aligned}$$

which is impossible because $5 \le |S| \le 6$.

Claim 8. No vertex in G is a 6-vertex.

Proof. Suppose to the contrary that G has a 6-vertex, say x. By Claim 2, G[N(x)] contains $K_{2,2,2}$ as a spanning subgraph. Let C be a component of $G \setminus N[x]$. By Claim 6, $|N(C)| \ge 5$. If N(C) = N(x)or G[N(x)] contains K_6^{\pm} as a spanning subgraph, then $G \succcurlyeq \mathcal{K}_8^{-4}$ by contracting C to a single vertex. Thus $|N(C) \cap N(x)| = 5$ for every component C of $G \setminus N[x]$ and $G[N(x)] = K_{2,2,2}$. By Claim 2, no vertex in N(x) is a 5-vertex in G. Thus $G \setminus N[x]$ is disconnected. Let $C' \ne C$ be another component of $G \setminus N[x]$. By Claim 7, $G[N(C)] = K_5^{\pm}$ and $G[N(C')] = K_5^{\pm}$. Let $y \in N(x) \cap N(C)$ such that y is incident with a missing edge of G[N(C)]. Then $G \succcurlyeq \mathcal{K}_8^{-4}$ by contracting C onto y and C' to a single vertex, a contradiction.

Claim 9. No vertex in G is a 7-vertex.

Proof. Suppose G has a 7-vertex, say x. By Claim 2, $\delta(G[N(x)]) \geq 4$, and so $\overline{G}[N(x)]$ is the disjoint union of paths and cycles. By Claim 9, no vertex in N(x) is a 6-vertex in G. Suppose there exists $y \in N(x)$ such that y is a 5-vertex in G. Then $N[y] \subseteq N[x]$ and $G[N[y]] = K_6$. Thus $G[N[x]] \succeq \mathcal{K}_8^{-4}$ because $\delta(G[N(x)]) \ge 4$, a contradiction. This proves that no vertex in N(x) is a 5-vertex or 6-vertex in G. Let $\mathcal{F} := \{C_7, C_6 \cup K_1, C_5 \cup K_2, C_4 \cup K_3\}$. Then $\overline{G}[N(x)]$ is a spanning subgraph of some member in \mathcal{F} because $\overline{G}[N(x)]$ is the disjoint union of paths and cycles. We only consider the worst scenario that $\overline{G}[N(x)]$ is isomorphic to some graph in \mathcal{F} . It is straightforward to check that every graph in \mathcal{F} has a \mathcal{K}_6^{-3} minor. Let C be a component of $G \setminus N[x]$. By Claim 6, $|N(C)| \ge 5$. If N(C) = N(x), then $G \succcurlyeq \mathcal{K}_8^{-4}$ by contracting C to a single vertex. Thus $N(C) \neq N(x)$ for every component C of $G \setminus N[x]$. We claim that $G \setminus N[x]$ is disconnected. Suppose not. Then $C = G \setminus N[x]$ and $\overline{G}[N(x)] = K_1 + C_6$. Note that every vertex on C_6 belongs to N(C). Thus $G \succeq \mathcal{K}_8^{-4}$ by contracting $G \setminus N[x]$ onto a vertex of C_6 , a contradiction. This proves that $G \setminus N[x]$ is disconnected, as claimed. Let C and C' be two distinct components of $G \setminus N[x]$ such that N(C) contains the most number of missing edges of G[N(x)]. It is straightforward to check that either G[N(C)] must contain two adjacent missing edges, say x_1x_2, x_1x_3 , where $x_1, x_2, x_3 \in N(x)$ are distinct, or $\overline{G}[N(x)] \in \{K_1 \cup C_6, K_2 \cup C_5\}$ and $G[N(C)] = G[N(C')] = K_5^{=}$. In the former case, since $|N(C')| \ge 5$, by Claim 7, we see that $G[N(C') \setminus x_1]$ must contain a missing edge, say e. By contracting C onto x_1 and C' onto an end of the edge e, it is straightforward to check that $G[N[x]] + \{x_1x_2, x_1x_3, e\} \geq \mathcal{K}_8^{-4}$, a contradiction. In the latter case, let y_1y_2 be a missing edge

in G[N(C)] and y_3y_4 be a missing edge in G[N(C')] such that y_1y_2, y_3y_4 are two different missing edges in G[N[x]]. We may assume that $y_1 \neq y_3$. Then $G[N[x]] + \{y_1y_2, y_3y_4\} \succeq \mathcal{K}_8^{-4}$ by contracting C onto y_1 and C' onto y_3 , a contradiction.

Claim 10. G is \mathcal{K}_7^{-2} -free.

Proof. Suppose G has a subgraph H such that $H \in \mathcal{K}_7^{-2}$. Since G is 5-connected, we obtain a \mathcal{K}_8^{-4} minor in G by contracting a component of $G \setminus V(H)$ to a single vertex, a contradiction.

Claim 11. G has at most one 5-vertex.

Proof. Suppose to the contrary that G has two distinct 5-vertices, say x and y. By Claims 2 and 3, $xy \notin E(G)$ and N[x] and N[y] are 6-cliques. Suppose $N(x) \neq N(y)$. By Claim 6 and Menger's Theorem, there exist five pairwise internally vertex-disjoint (x, y)-paths Q_1, \ldots, Q_5 . We may assume that Q_1 has at least four vertices with $V(Q_1) \cap N(x) = \{x_1\}$ and $V(Q_1) \cap N(y) = \{y_1\}$. Let Q_1^* be the (x_1, y_1) -subpath of Q_1 . Then $G \succcurlyeq \mathcal{K}_8^{-4}$ by contracting all the edges of $Q_1^* \setminus y_1$, $Q_2 \setminus \{x, y\}, \ldots, Q_5 \setminus \{x, y\}$, a contradiction. Thus N(x) = N(y). Then $G[N[x] \cup N[y]] = K_7^-$, contrary to Claim 10.

Claim 12. No 8-vertex is adjacent to a 5-vertex in G.

Proof. Suppose to the contrary that there exists $xy \in E(G)$ such that d(x) = 8 and d(y) = 5. By Claim 2, $G[N[y]] = K_6$, $N[y] \subseteq N[x]$ and $\delta(G[(N(x)]) \ge 4$. Let $A := N(x) \setminus N[y]$. Then |A| = 3 because $xy \in E(G)$. Let $A := \{a_1, a_2, a_3\}$. Suppose $G[A] \ne K_3$, say $a_1a_2 \notin E(G)$. Then either $e(\{a_1, a_2\}, N(x) \setminus A) \ge 7$, or $e(\{a_1, a_2\}, N(x) \setminus A) = 6$ and a_3 is complete to $\{a_1, a_2\}$, because $\delta(G[(N(x)]]) \ge 4$. Then $G[N[x] \setminus a_3] \succcurlyeq K_8^{-4}$ in the former case, and $G[N[x]]/a_1a_3 \succcurlyeq K_8^{-4}$ in the latter case, a contradiction. This proves that $G[A] = K_3$. Then $e(a_i, N(x) \setminus A) \ge 2$ for each $i \in [3]$. If $e(a_j, N(x) \setminus A) \ge 3$ for some $j \in [3]$, then $G[N[y] \cup \{a_j\}]$ is not K_7^{-2} -free, contrary to Claim 10. Thus $e(a_i, N(x) \setminus A) = 2$ for each $i \in [3]$. Next if there exists $z \in N(x) \cap N(y)$ such that z has exactly one neighbor, say a_1 , in A, then $G[N[x]]/za_1 \succcurlyeq K_8^{-4}$, a contradiction. Since $e(A, N(x) \setminus A) = 6$ and $|N(x) \cap N(y)| = 4$, it follows that there exists $z^* \in N(x) \cap N(y)$ such that z^* is anticomplete to A in G. By Claim 11, z^* is not a 5-vertex in G. Let C be a component of $G \setminus N[x]$ such that $z^* \in N(C)$. Note that $y \notin N(C)$ and $N(x) \cap N(y)$ is a 4-clique. By Claim 7, N(C) must contain at least two vertices, say a_1, a_2 , in A. By contracting C onto z^* , we see that $G[N[x] \setminus a_3] + \{z^*a_1, z^*a_2\} \models K_8^{-4}$, a contradiction. □

Claim 13. Let $x \in V(G)$ be an 8-vertex in G, and let M be the set of vertices of N(x) not adjacent to all other vertices of N(x). Then there is no component K of $G \setminus N[x]$ such that $M \subseteq N(K)$. In particular, $G \setminus N[x]$ is disconnected.

Proof. Suppose such a component K exists. Then every vertex in M has a neighbor in K because $M \subseteq N(K)$. By Lemma 4.1, there exists $y \in N(x)$ such that $G[N(x)] \setminus y$ has a \mathcal{K}_6^{-4} minor. If N[y] = N[x], then $G[N[x]] \succeq \mathcal{K}_8^{-4}$, a contradiction. Thus $y \in M \subseteq N(K)$. By contracting K onto y, we obtain a \mathcal{K}_8^{-4} minor in G, a contradiction. This proves that no such component K exists. By

Claims 8, 9 and 12, every vertex in M has degree at least eight, and thus every vertex in M has a neighbor in $G \setminus N[x]$. It follows that $G \setminus N[x]$ is disconnected.

Claim 14. Let $x \in V(G)$ be an 8-vertex in G. Then $G \setminus N[x]$ has at most one component C with |C| = 1. Moreover, if C is a component of $G \setminus N[x]$ such that |C| = 1, then the sole vertex in C is a 5-vertex in G.

Proof. Let C be a component of $G \setminus N[x]$ with |C| = 1. Let y be the only vertex in C. Suppose y is not a 5-vertex in G. Then $d(y) \ge 8$ by Claims 8 and 9, and so N(C) = N(x), contrary to Claim 13. Thus y is a 5-vertex in G. By Claim 11, $G \setminus N[x]$ has at most one such component C.

Claim 15. Let $x \in V(G)$ be an 8-vertex in G. Then there is no component C of $G \setminus N[x]$ such that $|C| \geq 2$ and for every vertex $y \in V(C)$, either $d_G(y) = 5$ or $d_G(y) \geq 9$.

Proof. Suppose such a component C exists. By Claim 11, C has at most one 5-vertex in G. Let $y \in V(C)$ such that $d_G(y) \ge 9$. Then $|V(C) \cup N(C)| \ge 10$. By Claim 13, $N(C) \ne N(x)$. Thus $|C| \ge |N[y] \setminus N(C)| \ge 10 - 7 = 3$. Let $G_1 := G \setminus C$ and $G_2 := G[C \cup N(C)]$. Note that $|G_2| \ge 10$ and N(C) is a minimal separating set of G. Let d_1 be defined as in the paragraph prior to Claim 4. Let $z \in N(C)$ such that $d_{G[N(C)]}(z) = \delta(G[N(C)])$. Let $d := d_{G[N(C)]}(z)$. By contracting $G_1 \setminus N(C)$ onto z, we see that $d_1 \ge |N(C)| - d - 1$. By (*),

$$e(G_2) < 4.5(|C| + |N(C)|) - 12 - (|N(C)| - d - 1) = 4.5|C| + 3.5|N(C)| + d - 11.$$
 (a)

Now let $t = e_G(C, N(C))$. Then $e(G_2) = e(C) + t + e(G[N(C)])$. Note that $2e(C) \ge 9(|C|-1) + 5 - t$ and $2e(G[N(C)]) \ge d|N(C)|$, so we have

$$2e(G_2) = 2e(C) + 2t + 2e(G[N(C)]) \ge 9|C| - 4 + t + d|N(C)|.$$
 (b)

Combining (a) and (b) yields

$$9|C| + 7|N(C)| + 2d - 22 > 2e(G_2) \ge 9|C| - 4 + t + d|N(C)$$

and so

$$-t > d(|N(C)| - 2) - 7|N(C)| + 18.$$
 (c)

Note that $\delta(G[N(x)]) \ge 4$ by Claim 2, and N(C) is a subset of N(x), so

$$d = \delta(G[N(C)]) \ge 4 - (8 - |N(C)|) = |N(C)| - 4.$$

This, together with (c), implies that

$$-t > (|N(C)| - 4) (|N(C)| - 2) - 7|N(C)| + 18$$

= $|N(C)|^2 - 13|N(C)| + 26$
= $(|N(C)| - \frac{13}{2})^2 - \frac{65}{4},$

so $-t \geq -16$. But then

$$|C|(|C|-1) \ge 2e(C) \ge 9(|C|-1) + 5 - t \ge 9|C| - 20,$$

which implies that $|C| \ge 8$ because $|C| \ge 3$. Thus $G \succcurlyeq C \succcurlyeq \mathcal{K}_8^{-4}$, contrary to the choice of G. \Box

To complete the proof, since $e(G) = \lceil 4.5n - 12 \rceil$, we have $\delta(G) \leq 8$. By Claims 3, 8 and 9, let x be an 8-vertex in G. By Claim 14, let C be a component of $G \setminus N_G[x]$ with $|C| \geq 2$ and, subject to that, x and C are chosen so that |C| is minimized. By Claims 11 and 15, C contains an 8-vertex, say y, in G. Note that $N_G(x) \neq N_G(y)$ because $|C| \geq 2$. Let K be the component of $G \setminus N_G[y]$ containing x. Then $|K| \geq 2$ because $N_G(x) \neq N_G(y)$. Note that $N_G(x) \cap N_G(y) \subseteq N(K)$, and every vertex in $N_G(x) \setminus N_G(y)$ belongs to K. Let M be the set of vertices of $N_G(y)$ not adjacent to all other vertices of $N_G(y)$. We claim that $M \subseteq N(K)$. Suppose not. Let $z \in M \setminus N(K)$. Then $z \notin N_G(x)$, else $z \in N(K)$ because $x \in V(K)$. It follows that $z \in V(C)$. By Claim 12, we have $d_G(z) \geq 8$; in addition, $N_G(z)$ has no 5-vertex in G if $d_G(z) = 8$, and at least one vertex in $N_G(z) \setminus N_G(y)$ is not a 5-vertex in G if $d_G(z) \geq 9$, due to Claim 11. In either case, let z' be a neighbor of z in $G \setminus N_G[y]$ such that $d_G(z') \geq 8$. Note that $z' \in (N_G(x) \setminus N_G(y)) \cup V(C)$. Suppose $z' \notin V(K)$. Then $z' \in V(C)$ because every vertex in $N_G(x) \setminus N_G(y)$ belongs to K. Let C' be the component of $G \setminus N_G[y]$ that contains z'. Then $|C'| \geq 2$ by Claim 14, and C' is a proper subset of C, contrary to our choice of x and C. This proves that $z' \in V(K)$, and so $z \in N(K)$, contrary to the choice of z. Thus K is a component of $G \setminus N_G[y]$ such that $M \subseteq N(K)$, contrary to Claim 13.

This completes the proof of Theorem 1.7.

References

[1] Boris Albar and Daniel Gonçalves. On triangles in K_r -minor free graphs. J. Graph Theory, 88(1):154–173, 2018.

- [2] K. Appel and W. Haken. Every planar map is four colorable. I. Discharging. Illinois J. Math., 21(3):429–490, 1977.
- [3] K. Appel, W. Haken, and J. Koch. Every planar map is four colorable. II. Reducibility. *Illinois J. Math.*, 21(3):491–567, 1977.
- [4] Maria Chudnovsky, Bruce Reed, and Paul Seymour. The edge-density for $K_{2,t}$ minors. J. Combin. Theory Ser. B, 101(1):18–46, 2011.
- [5] Kathie Cameron and Kristina Vuvsković. Hadwiger's conjecture for some hereditary classes of graphs: a survey. Bulletin of the European Association for Theoretical Computer Science, 131, 2020.
- [6] G. A. Dirac. A property of 4-chromatic graphs and some remarks on critical graphs. J. London Math. Soc., 27:85–92, 1952.
- [7] G. A. Dirac. Trennende Knotenpunktmengen und Reduzibilität abstrakter Graphen mit Anwendung auf das Vierfarbenproblem. J. Reine Agew. Math., 204:116–131, 1960.
- [8] G. A. Dirac. Homomorphism theorems for graphs. Math. Ann., 153:69–80, 1964.
- [9] G. A. Dirac. On the structure of 5- and 6-chromatic abstract graphs. J. Reine Angew. Math., 214(215):43-52, 1964.

- [10] Michelle Delcourt and Luke Postle. Reducing Linear Hadwiger's Conjecture to coloring small graphs. arXiv:2108.01633.
- [11] H. Hadwiger. Über eine Klassifikation der Streckenkomplexe. Vierteljschr. Naturforsch. Ges. Zürich, 88:133–142, 1943.
- [12] I. T. Jakobsen. A homomorphism theorem with an application to the conjecture of Hadwiger. Studia Sci. Math. Hungar., 6:151–160, 1971.
- [13] I. T. Jakobsen. Weakening of the conjecture of Hadwiger for 8- and 9-chromatic graphs. Aarhus Universitet, Matematisk Institut Preprint Series, 22:1–17, 1971.
- [14] Leif K. Jørgensen. Vertex partitions of $K_{4,4}$ -minor free graphs. Graphs Combin., 17(2):265–274, 2001.
- [15] Ken-ichii Kawarabayashi and Bjarne Toft. Any 7-chromatic graph has K_7 or $K_{4,4}$ as a minor. Combinatorica, 25:327–353, 2005.
- [16] Ken-ichi Kawarabayashi. Hadwiger's conjecture. In *Topics in chromatic graph theory*, volume 156 of *Encyclopedia Math. Appl.*, pages 73–93. Cambridge Univ. Press, Cambridge, 2015.
- [17] Alexandr V. Kostochka. The minimum Hadwiger number for graphs with a given mean degree of vertices. *Metody Diskret. Analiz.*, (38):37–58, 1982.
- [18] Alexandr V. Kostochka. Lower bound of the Hadwiger number of graphs by their average degree. Combinatorica, 4(4):307–316, 1984.
- [19] Alexandr V. Kostochka and Noah Prince. On $K_{s,t}$ -minors in graphs with given average degree. Discrete Math., 308(19):4435–4445, 2008.
- [20] Alexandr V. Kostochka and Noah Prince. Dense graphs have $K_{3,t}$ minors. Discrete Mathematics, 310(20):2637-2654, 2010.
- [21] A. V. Kostochka. $K_{s,t}$ minors in (s+t)-chromatic graphs, II. J. Graph Theory, 75(4):377–386, 2014.
- [22] Daniela Kühn and Deryk Osthus. Minors in graphs of large girth. Random Structures Algorithms, 22(2):213–225, 2003.
- [23] Daniela Kühn and Deryk Osthus. Forcing unbalanced complete bipartite minors. European Journal of Combinatorics, 26(1):75–81, 2005.
- [24] W. Mader. Uber trennende Eckenmengen in homomorphiekritischen Graphen. Math. Ann., 175:243–252, 1968.
- [25] Joseph Samuel Myers. The extremal function for unbalanced bipartite minors. Discrete mathematics, 271(1-3):209–222, 2003.

- [26] Sergey Norin, Luke Postle, and Zi-Xia Song. Breaking the degeneracy barrier for coloring graphs with no K_t minor. arXiv:12206.001862.
- [27] Sergey Norin and Paul Seymour. Dense minors of graphs with independence number two. arXiv:1910.09378v2.
- [28] Martin Rolek. Graphs with no $K_9^{=}$ minor are 10-colorable. J. Graph Theory, 93(4):560–565, 2020.
- [29] Martin Rolek and Zi-Xia Song. Coloring graphs with forbidden minors. J. Combin. Theory Ser. B, 127:14–31, 2017.
- [30] Martin Rolek, Zi-Xia Song, and Robin Thomas. Properties of 8-contraction-critical graphs with no K_7 minor. arXiv:2208.07335.
- [31] Neil Robertson, Paul Seymour, and Robin Thomas. Hadwiger's conjecture for K_6 -free graphs. Combinatorica, 13(3):279–361, 1993.
- [32] P. Seymour. Hadwiger's Conjecture, chapter 13, pages 417–437. Springer, Cham, 2016. In: Open Problems in Mathematics (edited by J. Nash Jr. and M. Rassias).
- [33] Zi-Xia Song and Robin Thomas. The extremal function for K_9 minors. J. Combin. Theory Ser. B, 96(2):240–252, 2006.
- [34] Andrew Thomason. An extremal function for contractions of graphs. Math. Proc. Cambridge Philos. Soc., 95(2):261–265, 1984.
- [35] K. Wagner. Über eine Eigenschaft der ebenen Komplexe. Mathematische Annalen, 114:570– 590, 1937.
- [36] K. Wagner. Bemerkungen zu Hadwiger's Vermutung. Mathematische Annalen, 141:433–451, 1960.
- [37] Douglas R. Woodall. List colourings of graphs. In Surveys in combinatorics, 2001 (Sussex), volume 288 of London Math. Soc. Lecture Note Ser., pages 269–301. Cambridge Univ. Press, Cambridge, 2001.