arXiv:2208.07361v1 [cond-mat.str-el] 15 Aug 2022

A toy model for categorical charges

CLEMENT DELCAMP*

Max Planck Institute for the Physics of Complex Systems, Nöthnitzer Straße 38, 01187 Dresden, Germany

We consider a higher gauge topological model in three spatial dimensions whose input datum is a 2-group encoding the mixing of a 0-form \mathbb{Z}_2 - and 1-form \mathbb{Z}_3 -symmetry. We study the excitation content of the theory on the symmetry-preserving boundary. We show that boundary operators are organised into the fusion 2-category of 2-representations of the 2-group. These can be interpreted as categorical charges for an effective boundary model that inherits a global 2-group symmetry from the bulk topological order. Interestingly, we find that certain simple 2-representations are physically interpreted as composites of intrinsic excitations and condensation defects.

SEC. I | Introduction

In modern parlance, ordinary (global) symmetry operations leaving a physical system invariant, are referred to as ∂ -form symmetries. These are generated by operators supported on one-codimensional submanifolds of spacetime and labelled by elements of a group so that fusion of symmetry operators obeys the multiplication rule of the group. Crucially, symmetry operators are topological so that correlation functions are insensitive to deformations of the underlying submanifold that preserve its topology, unless it passes through charged operators, i.e. operators supported on zero-dimensional manifolds that transform non-trivially in a representation of the group. A special role is then played by irreducible representations as they decompose the action of the group and provide meaningful sets of quantum numbers.

In recent years, the concept of symmetry has been generalised in various ways. A first generalisation is obtained by relaxing the requirement that operators are supported on one-codimensional submanifolds. This results in the notion of q-form symmetry that admits symmetry operators for every q-codimensional submanifolds of spacetime [1]. Symmetry operators are still labelled by group elements, but for q > 1 the corresponding group must now be *abelian* [2]. Furthermore, the topological nature of the symmetry operators requires the charged operators to be *q*-dimensional. These higher-form global symmetries have received widespread attention in high-energy physics [3-15] and condensed matter theory [16-24]. A further generalisation consists in relaxing the *invertibility* condition of symmetry operators so that fusion rules are encoded into abstractly higher mathematical structures such as fusion categories [25]. These so-called categorical symmetries have also been under scrutiny as of late [26–39]. One concrete application of these generalised notions of symmetry is the broadening of Landau's theory of symmetry breaking so as to include phases of matter previously believed not to be accommodated by it, e.g. topological phases [40–43].

Interestingly, higher-form symmetries of various degrees can be combined in non-trivial ways so that they do not factorise. A prototypical example is the mixing of 0- and 1-form symmetries, whereby the 0-form symmetry acts on the 1-form charged operators. The associated groups combine into a categorification of the notion of group referred to as a 2-group [44]. Much progress has been made towards understanding the resulting 2-group global symmetries and the corresponding anomalies [45– 50]. Nevertheless, the corresponding *higher representation theory* in connection with the study of charged operators remains somewhat elusive.

There is indeed a generalised notion of representation that is suitable to 2-groups. These so-called 2representations are defined as 2-functors from the 2group—thought as a one-object 2-groupoid—to the 2category of 2-vector spaces [51–53], categorifying the definition of a group (1-)representation as a functor from the group—thought as a one-object (1-)groupoid—to the category of vector spaces. Since every group is in particular a 2-group, it makes sense to compute its 2representations. This special kind of 2-representations has been studied in detail in the past [54–56], and was recently shown to naturally arise in the context of the electromagnetic duality of higher-dimensional symmetric quantum models [57–59]. Perhaps surprisingly, the 2-category of 2-representations of a group contains more information than its lower-categorical counterpart, encoding in particular so-called *condensation defects* that arise for instance when gauging higher-form symmetries along submanifolds [60–66].

In this work, we consider an effective model with a specific global 2-group symmetry and elucidate in this context the physical interpretation of the corresponding 2representations. Our approach exploits a correspondence between symmetry-preserving gapped phases in (2+1)dand Neumann boundary conditions of (3+1)d topological models satisfying gauged versions of the symmetries [34]. A non-anomalous global 2-group symmetry can indeed be gauged by coupling it to 1- and 2-form connections that interact in a non-trivial way. The resulting theory is typically referred to as a higher gauge theory [67–69]. Practically, we study an exactly solvable model with a higher gauge theory interpretation that provides a lat-

^{*} clement.delcamp@ugent.be


tice Hamiltonian realisation of Yetter's homotopy 2-type topological quantum field theory [70]. The input datum is a strict 2-group encoding the mixing of a 0-form \mathbb{Z}_2 symmetry and 1-form \mathbb{Z}_3 -symmetry. This model yields bulk charge and flux composite *intrinsic* excitations that are either point-like or loop-like. The Neumann gapped boundary condition is then obtained by condensing all the magnetic excitations, and the resulting boundary operators can be identified with charged operators for the effective symmetric model. We explain how these *categorical charges* and their fusion statistics are organised into the monoidal bicategory of 2-representations of the 2-group. Interestingly, certain simple 2-representations are interpreted as composites of intrinsic excitations and condensation defects.

SEC. II | Categorical charges

We propose in this section a toy model for categorical charges. These charges arise as boundary topological excitations of a lattice Hamiltonian possessing a higher gauge theory interpretation.

II.A. Lattice Hamiltonian

Let us consider an oriented three-dimensional manifold \mathcal{M} with a non-empty boundary $\partial \mathcal{M}$. For now we assume that the boundary has a single connected component. We further suppose that \mathcal{M} can be endowed with a *cubic* cellulation \mathcal{M}_{\Box} whose vertices, edges, plaquettes and cubes are notated via v, e, p and c, respectively:



where bold lines correspond to edges along the boundary $\partial \mathcal{M}$. We require parallel edges to be all oriented in the positive direction according to the frame of reference depicted above, and denote by s(e) and t(e) the source and target vertices of the directed edge $e \subset \mathcal{M}_{\Box}$, respectively. This in turn induces an orientation for every plaquette $p \subset \mathcal{M}_{\Box}$, and canonically assigns to it a *basepoint* bp(p) defined as the unique vertex $v \subset \partial p$ such that $s(e_1) = v = s(e_2)$ for $e_1, e_2 \subset \partial p$.

Qubit and qutrit degrees of freedom are assigned to edges and plaquettes of \mathcal{M}_{\Box} , respectively. We identify such an assignment with a choice of *colouring* of the edges and plaquettes by \mathbb{Z}_2 and \mathbb{Z}_3 group variables, respectively, so that the microscopic Hilbert space is provided by $(\bigotimes_{\mathbf{e}} \mathbb{C}[\mathbb{Z}_2]) \otimes (\bigotimes_{\mathbf{p}} \mathbb{C}[\mathbb{Z}_3])$. We choose the presentations $\mathbb{Z}_2 = (\{+1, -1\}, \times) \ni g$ and $\mathbb{Z}_3 = (\{0, 1, 2\}, +) \ni h$, where the addition is modulo 3. Given a colouring of the cellulation we notate its restriction to an edge \mathbf{e} or a plaquette \mathbf{p} via $|g_{\mathbf{e}}\rangle$ and $|h_{\mathbf{p}}\rangle$, respectively, both regarded as elements of the microscopic Hilbert space. We identify by convention a qubit in the 'up' state $|\uparrow\rangle$ with $|+1\rangle$ or in the 'down' state $|\downarrow\rangle$ with $|-1\rangle$.

Qubit and qutrit degrees of freedom are governed by a lattice Hamiltonian obtained as a sum of mutually *commuting* local operators split in four families. Let us begin by introducing some definitions. We require the following operators:

$$\sigma_X = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_Z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

such that $\sigma_X \sigma_Z = -\sigma_Z \sigma_X$ and

$$\Sigma_X = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \quad \Sigma_Z = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & \bar{\omega} \end{pmatrix}, \quad \Gamma = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix},$$

with $\omega = \exp(\frac{2\pi i}{3})$ such that $\Sigma_Z^3 = \Sigma_X^3 = \operatorname{id}, \Sigma_Z \Sigma_X = \omega \Sigma_X \Sigma_Z, \Sigma_Z^\dagger \Sigma_X = \bar{\omega} \Sigma_X \Sigma_Z^\dagger, \Gamma \Sigma_Z \Gamma = \Sigma_Z^\dagger$ and $\Gamma \Sigma_X \Gamma = \Sigma_X^\dagger$. We also need *controlled gates* of the form

$$c\Sigma_X : \mathbb{C}^2 \otimes \mathbb{C}^3 \to \mathbb{C}^2 \otimes \mathbb{C}^3 : |g,h\rangle \mapsto (\mathrm{id} \otimes \Sigma_X^g)|g,h\rangle ,$$
(1)

where the qubit plays the role of the control. In practice we shall apply this gate between a qutrit assigned to a plaquette \mathbf{p} and a qubit assigned to an edge \mathbf{e} that may or may not be in the boundary of \mathbf{p} . We similarly define a controlled gate $c\Sigma_Z$. It immediately follows from the definitions that the following commutation relations are satisfied:


$$c\Sigma_X(\mathrm{id}\otimes\Gamma) = (\mathrm{id}\otimes\Gamma)c\Sigma_X^{\dagger},$$

$$c\Sigma_X(\sigma_X\otimes\mathrm{id}) = (\sigma_X\otimes\mathrm{id})c\Sigma_X^{\dagger},$$

$$c\Sigma_X(\sigma_X\otimes\Gamma) = (\sigma_X\otimes\Gamma)c\Sigma_X.$$

(2)

Similar commutation relations hold for $c\Sigma_Z$. Moreover, we have $c\Sigma_Z c\Sigma_X = \omega c\Sigma_X c\Sigma_Z$, and generalisation thereof. We graphically represent such controlled gates by means of a dotted line connecting a control qubit identified by 'c' and a target qutrit, e.g.



for any $g \in \mathbb{Z}_2$ and $h \in \mathbb{Z}_3$, with $[g+h] \equiv (g+h)$ mod 3. We shall also consider generalisations of these

controlled gates with more than one control qubits, e.g. $cc\Sigma_X : |g_1, g_2, h\rangle \mapsto (id \otimes id \otimes \Sigma_X^{g_1g_2})|g_1, g_2, h\rangle$, depicted as before via dotted lines connecting all the control qubits to the single qutrit.

We are now ready to write the lattice Hamiltonian, which we decompose as $\mathbb{H} = \mathbb{H}_{int(\mathcal{M})} + \mathbb{H}_{\partial\mathcal{M}}$, where $\mathbb{H}_{int(\mathcal{M})}$ only contains terms that act solely on qubit and qutrit degrees of freedom in the interior $int(\mathcal{M})$ of \mathcal{M} , and $\mathbb{H}_{\partial\mathcal{M}}$ only contains terms involving boundary degrees of freedom. Since we are ultimately interested in the boundary excitations, we shall only write down explicitly the component $\mathbb{H}_{\partial\mathcal{M}}$. Graphically, we have

where $g_{\mathbf{p}} := \prod_{\mathbf{e} \subset \partial \mathbf{p}} g_{\mathbf{e}}$. We can immediately check that $\mathbb{H}_{\partial \mathcal{M}}$ is Hermitian. As mentioned earlier, we distinguish four families of local operators, which act on neighbourhoods of the plaquettes, vertices, cubes and edges of the cubic lattice \mathcal{M}_{\Box} , respectively. We interpret these operators as enforcing *magnetic* and *electric* stabiliser conditions for the qubit and qutrit degrees of freedom, respectively. Notice, however, that except for the plaquette operators, all operators involve both qubit and qutrit degrees of freedom. We provide additional comments regarding the physical interpretations of these various terms in sec. II B.

Let us confirm that all the operators entering the definition of $\mathbb{H}_{\partial \mathcal{M}}$ are mutually commuting. First of all, the plaquette operators readily commute with all operators. Vertex operators commute with any cube operator in virtue of $(\sigma_X \otimes \Gamma) c \Sigma_Z^g = c \Sigma_Z^g (\sigma_X \otimes \Gamma)$ and with the sum of any edge operator and its Hermitian conjugate either in virtue of $(\sigma_X \otimes \Gamma) c \Sigma_X^g = c \Sigma_X^g (\sigma_X \otimes \Gamma)$ or $(\sigma_X \otimes id) c \Sigma_X^g =$ $c \Sigma_X^{-g} (\sigma_X \otimes id)$ together with $\Gamma \Sigma_X^g \Gamma = \Sigma_X^{-g}$. Finally, the commutation between cube and edge operators is guaranteed by $\Sigma_X^{g_1} \Sigma_Z^{g_2} = \omega^{g_1 g_2} \Sigma_Z^{g_2} \Sigma_X^{g_1}$ as well as the fact that cube operators vanish whenever the plaquette stabiliser conditions are not satisfied. Let us emphasise that this last requirement is crucial whenever the commutation relation between an edge and a plaquette operator involves four controlled gates whose control qubits are associated with boundary edges of the same plaquette. All the operators commute and thus the model is exactly solvable. Note finally that when discarding the qutrit degrees of freedom, the model reduces to the usual (3+1)d toric code [71, 72].

II.B. Higher gauge theory interpretation

Let us now reveal the higher gauge theory interpretation of the Hamiltonian we consider. Loosely speaking, a higher gauge theory is a gauge theory for which the group is replaced by a *categorification* of the notion of group referred to as a 2-group. We focus in this manuscript on strict 2-groups defined as strict monoidal categories whose objects and morphisms are invertible [44, 67]. Crucially strict 2-groups can be equivalently described in terms of *crossed modules* [73]. A crossed module \mathbb{G} is a quadruple $(G, H, \triangleright, \partial)$ consisting of two groups G and H, a group action $\triangleright : G \times H \to H$ by automorphisms, and a group homomorphism $\partial: G \to H$, which are subject to the so-called *Peiffer* identities. Our attention is on the so-called *dihedral* 2-group $\mathbb{G}(2,3) \equiv (\mathbb{Z}_2,\mathbb{Z}_3,\mathbb{1}_{\mathbb{Z}_2}:\mathbb{Z}_3)$ $\{+1\}, \triangleright$, where $g \triangleright h = h^g$ for any $g \in \mathbb{Z}_2$ and $h \in \mathbb{Z}_3$, so that $\mathbb{Z}_2 \ltimes \mathbb{Z}_3$ is isomorphic to the dihedral group \mathbb{D}_6 of order 6. The Peiffer identities being trivially satisfied for $\mathbb{G}(2,3)$, we need not writing them down explicitly. We can conveniently rewrite the controlled gates in terms of the action \triangleright as

$$c\Sigma_{X}^{k}:|g,h\rangle\mapsto|g,[h+(g\triangleright k)]\rangle$$

$$c\Sigma_{Z}^{k}:|g,h\rangle\mapsto(\omega^{h})^{g\triangleright k}|g,h\rangle,$$
(4)

for any $k \in \mathbb{Z}_3$.

Assigning group variables $g_{e} \in \mathbb{Z}_{2}$ to every edge $\mathbf{e} \subset \mathcal{M}_{\Box}$ defines an (ordinary) 1-form \mathbb{Z}_2 -connection, whereas assigning group variables g_{p} to every plaquette $p \subset \mathcal{M}_{\Box}$ defines a 2-form $\mathbb{Z}_3\text{-connection}.$ Together with the 1-flatness condition encoded into the plaquette operators, these assignments define a so-called 2-connection on the strict 2-group $\mathbb{G}(2,3)$. The cube operators then enforce a 2-flatness constraint on this 2-connection. Such a gauge field configuration is subject to two kinds of gauge transformations: On the one hand, we have ordinary (0form) \mathbb{Z}_2 -gauge transformations whose actions on the 2connection are encoded into the vertex operators. On the other hand, there are 1-form \mathbb{Z}_3 -gauge transformations acting on the 2-connection via the edge operators. Note that 0-form and 1-form gauge transformations typically do not commute, but commutation of the operators in the definition of the Hamiltonian is achieved by considering an edge operator together with its Hermitian conjugate.

A general gauge transformation can then be defined as a 0-form transformation preceded by a 1-form transformation. Composition of such general gauge transformations is then provided by group multiplications in \mathbb{D}_6 . The model we consider is an example of higher gauge models of topological phases considered in ref. [74, 75].

Let us remark that the higher gauge model with input datum $\mathbb{G}(2,3)$ admits by no means a unique choice of gapped boundary condition. The boundary Hamiltonian $\mathbb{H}_{\partial \mathcal{M}}$ we focus on in this manuscript encodes the socalled 'smooth' (or Neumann) boundary condition. This boundary condition is obtained by condensing all the magnetic excitations, i.e. violations of 1- and 2-flatness constraints, so that only electric excitations, i.e. violations of 0- and 1-form gauge invariance, live at the boundary. The remainder of this manuscript is dedicated to the study of these electric excitations.


II.C. Boundary excitations

Our model yields four types of bulk excitations obtained by violating stabiliser conditions associated with each family of Hamiltonian operators. In light of the higher gauge theory interpretation of the model, we shall refer to the corresponding four types of *elementary* bulk excitations as *1-flux*, *0-charge*, *2-flux* and *1-charge*, respectively. We know from general considerations that 0-charges and 2-fluxes should be zero-dimensional objects, whereas 1-fluxes and 1-charges should be onedimensional.

We focus in this manuscript on topological boundary excitations for the boundary condition encoded into our choice of Hamiltonian $\mathbb{H}_{\partial \mathcal{M}}$. As evoked above, this is the boundary condition that condenses 1- and 2-fluxes. To precise this statement, we first need to clarify the notion of 'condensation' within our context. For twodimensional topological models, condensing a topological excitation amounts to making it *local*—in the sense that it can be created by a local operator. However, this convention becomes ambiguous in higher dimensions. Consider for instance the (3+1)d toric code that hosts pointlike (0-)charges and loop-like (1-)fluxes created at the boundary of string- and membrane-like topological operators, respectively. The 'rough' (or Dirichlet) boundary condition condenses electric excitations since a charge brought to the boundary can be annihilated by a local boundary operator. What about the 'smooth' (or Neumann) boundary condition? When bringing a loop-like flux to the boundary, it cannot be annihilated by a local operator but by a genuine line-like operator, i.e. an operator that does *not* depend on the topology of a choice of bounded surface. Equivalently, a loop-like flux excitation can be created at the boundary with a genuine line-like operator but it cannot be detected as we cannot link it to a topological charge, and thus it is identified with the vacuum. Going back to the higher gauge model of interest, we could show that the boundary condition

under consideration condenses the magnetic excitations as 1-fluxes can be annihilated at the boundary by genuine line-like operators and 2-fluxes by local operators. This also means that 1-fluxes are defined up to genuine line-like operators. We shall exploit this property in the following but for now let us explicitly identify the boundary (electric) operators.


Firstly, we denote by 1 the trivial (string-like) 1-charge and by 1_1 the trivial (point-like) 0-charge. As suggested by the notation, we shall think of a trivial 0-charge 1_1 as being attached to a trivial 1-charge 1, or rather as being a domain wall $1 \rightarrow 1$ between two trivial 1-charges. The first non-trivial excitation is the 0-charge e. Akin to the (3+1)d toric code, pairs of 0-charges e can be created at the endpoints of a string of σ_Z operators, e.g.



As before, we think of a 0-charge e as being a domain wall $e : \mathbb{1} \to \mathbb{1}$ between two trivial 1-charges 1. From these 0-charges descends another type of object, namely one-dimensional defects that condense 0-charges e. These condensation defects do not constitute excitations of the Hamiltonian per se, but rather deformations of the model itself. Physically, such a defect is interpreted as a onedimensional gas of 0-charges e that spontaneously breaks the effective \mathbb{Z}_2 -symmetry [60, 61, 64, 65]. Microscopically, it can be realised as follows: Given a (1-)path ℓ along the edges of the cubic lattice, we discard the vertex operators associated with the vertices $\mathsf{v} \subset \ell$ along the path, and add the energy term

$$-\sum_{\mathbf{e}\subset\ell}-\sigma_{Z}-, \qquad (6)$$

which effectively projects the qubits along the path ℓ onto the spin- \uparrow subspace. With respect to the original Hamiltonian, this defect amounts to creating at every edge along the path an equal-weight superposition of the vacuum and the excited state. We can then readily show that as we bring a 0-charge e in contact with this defect, it cannot be distinguished from the vacuum anymore. Henceforth, we denote this defect by \mathcal{E}_1 and depict it as a wiggly red line \frown . Importantly, this string-like defect does not have to be closed, i.e. there exist domain walls between the trivial 1-charge 1 and the condensation defect \mathcal{E}_1 . We notate via $x^{\vdash} : \mathbb{1} \to \mathcal{E}_1$ and $x^{\dashv} : \mathcal{E}_1 \to \mathbb{1}$ the (unique) domain walls associated with each endpoint of the string. Furthermore, we distinguish two types of domain walls between two condensation defects \mathcal{E}_1 . Indeed, in addition to the trivial domain wall $1_{\mathcal{E}_1} : \mathcal{E}_1 \to \mathcal{E}_1$, there is a domain wall $n : \mathcal{E}_1 \to \mathcal{E}_1$ that corresponds to violating the 1-flatness stabiliser conditions associated with the three plaquettes meeting at an edge $\mathbf{e} \subset \ell$ along the path. We graphically summarise these various objects as




where the blue-shaded plaquettes are those for which the 1-flatness/plaquette stabiliser conditions are violated. Note however that interpreting this excitation strictly as a 1-flux requires some care as the cube operators are defined to vanish whenever the 1-flatness is not everywhere satisfied around the cube (see [76] for a discussion on the interplay between 1- and 2-fluxes). The domain wall nis distinct from the trivial one $1_{\mathcal{E}_1}$ since we cannot annihilate the resulting magnetic excitation with a local or genuine line-like operator without creating additional excitations or altering the condensation defect itself. Indeed, the only local operation that could annihilate this magnetic excitation would amount to modifying the defect by projecting the qubit at the intersection of the blue-shaded plaquettes onto the spin- \downarrow subspace. This provides an alternative interpretation for the domain wall n as

$$\mathcal{E}_1$$
, (8)

where $\rightarrow \rightarrow$ indicates an energy term in the Hamiltonian effectively projecting the corresponding qubit onto the spin- \downarrow subspace. We exploit both interpretations in the following.

The defects identified so far coincide with those found on the smooth boundary of the (3+1)d toric code by Kong et al. in [64]. Let us now study another type of electric excitations, which is obtained by violating the 1-form gauge invariance. Consider inserting a Σ_Z operator on a given plaquette **p**. Such an operator insertion certainly breaks the edge stabiliser conditions at every $\mathbf{e} \subset \partial \mathbf{p}$. However, these violations do not correspond to elementary excitations, as witnessed by the fact that the commutator of Σ_Z with two of the edge operators along $\partial \mathbf{p}$ depends on the \mathbb{Z}_2 -colouring of two edges $\mathbf{e}_1, \mathbf{e}_2 \subset \partial \mathbf{p}$, as well as the fact that it does not commute with the vertex operator associated with bp(**p**). This is remedied by projecting these boundary qubits onto a specific state. More precisely, we construct the 1-charge \mathcal{E}_{ω} by combining a Σ_Z operator with a condensation defect running along the boundary of p so as to project the corresponding qubits onto the spin- \uparrow subspace, i.e



Replacing Σ_Z by Σ_Z^{\dagger} yields another 1-charge denoted by $\mathcal{E}_{\bar{\omega}}$. How about defining larger excitations? Given a 2path \mathcal{P} , acting with $(\Sigma_Z^k)_{\mathsf{p}}$, with $k \in \mathbb{Z}_3$, at every $\mathsf{p} \subset \mathcal{P}$ would not yield the expected excitations. Indeed, given two adjoining plaquettes p_1 and p_2 , the edge stabiliser operator acting at $e = p_1 \cap p_2$ may or may not commute with $(\Sigma_Z^k)_{\mathbf{P}_1} \otimes (\Sigma_Z^k)_{\mathbf{P}_2}$ depending on the \mathbb{Z}_2 -colouring of the edges bounding these plaquettes. More precisely, we are looking for a topological membrane operator that only creates a loop-like excitation at its boundary. We must therefore adapt this naive attempt in order to account for the dependence of the edge stabiliser operators on the qubits living on the 2-path. Assigning a basepoint $bp(\mathcal{P})$ to the 2-path \mathcal{P} and denoting by $\ell_{\mathcal{P}}^{p}$: bp $(\mathcal{P}) \to bp(p)$ a choice of 1-path from $bp(\mathcal{P})$ to the basepoint of any plaquette $\mathbf{p} \subset \mathcal{P}$, we consider instead the operator¹

$$\prod_{\mathbf{p}\subset\mathcal{P}}\prod_{\mathbf{e}\subset\ell_{\mathcal{P}}^{\mathbf{p}}}(\mathbf{c}\Sigma_{Z}^{k})_{\mathbf{e},\mathbf{p}}=\prod_{\mathbf{p}\subset\mathcal{P}}(\Sigma_{Z}^{k})_{\mathbf{p}}^{\prod_{\mathbf{e}\subset\ell_{\mathcal{P}}^{\mathbf{p}}}(\sigma_{Z})_{\mathbf{e}}},\qquad(10)$$


with $k \in \mathbb{Z}_3$. Concretely, this operator acts on a state associated with a plaquette **p** as

$$\begin{aligned} |g_{\mathbf{e}_{1}},\ldots,g_{\mathbf{e}_{|\ell_{\mathcal{P}}^{\mathbf{p}}|}},h_{\mathbf{p}}\rangle &\mapsto (\Sigma_{Z}^{k})_{\mathbf{p}}^{\prod_{\mathbf{e}\subset \ell_{\mathcal{P}}^{\mathbf{p}}}g_{\mathbf{e}}}|g_{\mathbf{e}_{1}},\ldots,g_{\mathbf{e}_{|\ell_{\mathcal{P}}^{\mathbf{p}}|}},h_{\mathbf{p}}\rangle \\ &= (\Sigma_{Z}^{g_{\ell_{\mathcal{P}}^{\mathbf{p}}}\triangleright k})_{\mathbf{p}}|g_{\mathbf{e}_{1}},\ldots,g_{\mathbf{e}_{|\ell_{\mathcal{P}}^{\mathbf{p}}|}},h_{\mathbf{p}}\rangle \\ &= (\omega^{h_{\mathbf{p}}})^{(g_{\ell_{\mathcal{P}}^{\mathbf{p}}}\triangleright k)}|g_{\mathbf{e}_{1}},\ldots,g_{\mathbf{e}_{|\ell_{\mathcal{P}}^{\mathbf{p}}|}},h_{\mathbf{p}}\rangle\,,\end{aligned}$$


where $\mathbf{e}_1, \ldots, \mathbf{e}_{|\ell_{\mathcal{P}}^p|}$ are the constitutive edges of the path $\ell_{\mathcal{P}}^p$ and $g_{\ell_{\mathcal{P}}^p} = \prod_{\mathbf{e} \subset \ell_{\mathcal{P}}^p} g_{\mathbf{e}}$. As long as the 1-flatness stabiliser conditions are everywhere satisfied, this operator is insensitive to the choices of 1-paths $\mathrm{bp}(\mathcal{P}) \to \mathrm{bp}(\mathbf{p})$ for every $\mathbf{p} \subset \mathcal{P}$. In practice, we shall focus on rectangular membrane-like operators for convenience, in which case we can always choose without loss of generality the basepoint $\mathrm{bp}(\mathcal{P})$ of the 2-path \mathcal{P} to be the basepoint of the bottom left plaquette. Combined with condensation defects running along the boundary $\partial \mathcal{P}$ of the 2-path, the operator (10) creates a loop-like 1-charge \mathcal{E}_{ω^k} . For

¹ Analogous operators for more general higher gauge models were constructed in [77, 78].


instance, we have




where we exploited the fact that qubits along the condensation defect are projected onto the spin- \uparrow subspace to simplify the operators. Let us confirm that this defect excitation yields a well-defined 1-charge. In particular, we need to check that all the stabiliser conditions in the interior of \mathcal{P} are fulfilled. Typically, we cannot readily apply the commutation relations between the $c\Sigma_X$ and $c\Sigma_Z$ gates as they are defined with respect to different control qubits. This is remedied by exploiting the fact that, as long as the 1-flatness stabiliser conditions are satisfied, we can modify the 1-paths $\ell_{\mathcal{P}}^{\mathcal{P}}$ with respect to which the operator is defined. For instance, we have



Since operators of the form




always commute, this guarantees that the membrane operator defined in eq. (11) commutes with every edge operator in the interior of \mathcal{P} . Similarly, it follows from the commutation relations given in eq. (2) that operators of the form




always commute, which in turn guarantees that the membrane operator defined in eq. (11) commutes with every

vertex operator in the interior of \mathcal{P} . More generally, the interiors of the membrane-like operators defined according to eq. (10) are 0- and 1-form gauge invariant.

Naturally, when picking $k = 0 \in \mathbb{Z}_3$, the 1-charge \mathcal{E}_{ω^k} reduces to the condensation defect \mathcal{E}_1 previously defined. Notice however that since they appear at the boundary of topological membrane-like operators—in contrast to the condensation defect \mathcal{E}_1 —the 1-charges \mathcal{E}_{ω} and $\mathcal{E}_{\bar{\omega}}$ necessarily have the topology of a circle, i.e. there is no domain wall between them and the trivial 1-charge 1, nor there are domain walls between $\mathcal{E}_{\omega/\bar{\omega}}$ and \mathcal{E}_1 . Moreover, there are only trivial domain walls between two 1-charges \mathcal{E}_{ω} or two 1-charges $\mathcal{E}_{\bar{\omega}}$, namely $1_{\mathcal{E}_{\omega}} : \mathcal{E}_{\omega} \to \mathcal{E}_{\omega}$ and $1_{\mathcal{E}_{\bar{\omega}}} : \mathcal{E}_{\bar{\omega}} \to \mathcal{E}_{\bar{\omega}}$. But there are non-trivial domain walls $n_{\omega} : \mathcal{E}_{\omega} \to \mathcal{E}_{\bar{\omega}}$ and $n_{\bar{\omega}} : \mathcal{E}_{\bar{\omega}} \to \mathcal{E}_{\omega}$. These are constructed as follows: Consider a rectangular 2-path \mathcal{P} , halve it for instance vertically, and apply operators (10)so as to define a 1-charge \mathcal{E}_{ω} on the left-hand side and a 1-charge $\mathcal{E}_{\bar{\omega}}$ on the right-hand side. The domain wall itself then amounts to altering the condensation defect along $\partial \mathcal{P}$ by projecting the interfacial qubits on the side of the basepoint $bp(\mathcal{P})$ onto the spin- \downarrow subspace. For instance, we have



where we recall that the wiggly purple lines \longrightarrow indicate a qubit that is projected onto the spin- \downarrow subspace. The domain wall $n_{\bar{\omega}} : \mathcal{E}_{\bar{\omega}} \to \mathcal{E}_{\omega}$ is defined similarly. Let us confirm that these domain walls are such that the resulting defects are well-defined 1-charges. More specifically, we need to confirm that only stabiliser conditions along the boundary of the 2-paths are violated. Consider for instance the two adjoining plaquettes in the middle of the operator considered in eq. (15). In order to compute the commutation relation with the interstitial edge operator, we must first modify the 1-path $\ell_{\mathcal{P}}^{\mathsf{p}}$ associated with the plaquette on the left-hand side. The operator is modified as follows:



We can then invoke the commutation of operators of the form (13) to confirm that the edge stabiliser condition

is indeed fulfilled. Similarly, we can confirm that all the edge stabiliser conditions in the interior of \mathcal{P} are satisfied as well as all the vertex stabiliser constraints.


This concludes the enumeration of the boundary excitations of the higher gauge model. We summarise our findings so far in the table below:

In sec. III we shall recover these 1-charges and the domain walls between them as the simple 2-representations of the 2-group $\mathbb{G}(2,3)$ and simple 1-intertwiners between them, respectively.


As a final remark, let us comment on the physical interpretation of these defects as categorical charges. By definition, a ground state of the higher gauge model satis fies all the stabiliser conditions, i.e. it is in the +1eigenspace of all the operators. In particular, it is invariant under 0- and 1-form gauge transformations, as generated by the vertex and edge operators, respectively. At the boundary, these ground state constraints enforce global 0- and 1-form symmetry conditions, respectively. The corresponding symmetry operators can be obtained by composing vertex and edge operators of the bulk Hamiltonian. In virtue of the commutation relation between vertex and edge operators, 0-form symmetry operators act on 1-form symmetry operators, resulting in a global symmetry encoded into the 2-group $\mathbb{G}(2,3)$. By definition such symmetry operators act trivially on the quasi-two-dimensional system associated with our choice of gapped boundary condition. In other words, this is the symmetry-preserving gapped boundary condition. In that context, we can interpret the boundary excitations derived in this section as categorical charges with respect to this global 2-group symmetry that is enforced by the bulk topological order.

II.D. Excitation statistics


Let us now compute the fusion rules of the point-like domain wall excitations on the one hand and the fusion of the 1-charges on the other hand.² We begin with domain walls of the trivial 1-charge 1. Trivially, we have $l_1 \circ l_1 \simeq l_1$ and $l_1 \circ e \simeq e \simeq e \circ l_1$. Moreover, in virtue of $\sigma_Z^2 = \text{id}$, we have $e \circ e \simeq l_1$. Let us pursue with domain walls of the 1-charge \mathcal{E}_1 . The only non-trivial fusion rule is associated with $n \circ n$. Recall that a domain wall n may be interpreted as a violation of the 1-flatness stabiliser conditions surrounding an edge along the condensation defect. Bringing two such domain walls next to one another, the resulting excitation can be annihilated with a local operator:



and thus $n \circ n \simeq 1_{\mathcal{E}_1}$. The remaining fusion rules involving the 1-charge \mathcal{E}_1 are $x^{\vdash} \circ x^{\dashv}$ and $x^{\dashv} \circ x^{\vdash}$. First of all, remark that, by construction, the output of these fusion processes must be domains walls $\mathbb{1} \to \mathbb{1}$ and $\mathcal{E}_1 \to \mathcal{E}_1$, respectively. Focusing on the first scenario, let us bring together the domain walls x^{\dashv} and x^{\vdash} as follows:




The qubit located in the middle of the horizontal edge in-between the domain walls x^{\dashv} and x^{\vdash} can either be \uparrow or \downarrow . This gives rise to a decomposition of the Hilbert space:



In the first configuration, the qubit being projected onto the spin- \uparrow subspace, the two condensation defects get connected. In the second configuration, we can think of the qubit being projected onto the spin- \downarrow subspace as an alternation of the condensation defect as in eq. (8), or equivalently as the outcome of acting with a σ_X operator on a short condensation defect \mathcal{E}_1 effectively causing a violation of the 1-flatness stabiliser conditions associated


² Although we refer to the 'fusion' of domain walls, it is usually not encoded into a monoidal structure but rather a composition rule.

with the surrounding plaquettes, i.e.



Notice however that for this interpretation to be selfconsistent, it is required to apply the same mechanism somewhere else on the condensation defect, or along a different condensation defect, as the resulting wave function would not be in the ground state subspace otherwise. Putting everything together, we find $x^{\dashv} \circ x^{\vdash} \simeq 1_{\mathcal{E}_1} \oplus n$. Conversely, bringing together x^{\vdash} and x^{\dashv} yields a short condensation defect \mathcal{E}_1 that amounts to the local operator creating a pair of non-elementary point-like electric excitations decomposing as $1_{\mathbb{I}} \oplus e$, and therefore $x^{\vdash} \circ x^{\dashv} \simeq 1_{\mathbb{I}} \oplus e$.

The fusion rules computed so far coincide with those found on the smooth boundary of the (3+1)d toric code as computed in ref. [64]. Let us now consider the fusion of domain walls involving the 1-charges \mathcal{E}_{ω} and $\mathcal{E}_{\bar{\omega}}$. The only non-trivial fusion rules are those associated with $n_{\omega} \circ n_{\bar{\omega}}$ and $n_{\bar{\omega}} \circ n_{\omega}$. Given the obvious symmetry, it is sufficient to focus on the former. This situation arises when a 'thin' membrane-like operator creating a 1-charge $\mathcal{E}_{\bar{\omega}}$ is inserted between two membrane-like operators creating two 1-charges \mathcal{E}_{ω} , e.g.




Acting with vertex operators at the vertices where the domain walls meet has the effect of flipping the qubits projected onto the spin- \downarrow subspace, thus annihilating the domain walls, while turning the 1-charge $\mathcal{E}_{\bar{\omega}}$ into \mathcal{E}_{ω} without introducing any magnetic excitations. This follows in particular from $\Gamma \Sigma_Z^{\dagger} = \Sigma_Z \Gamma$ as well as $(\sigma_X \otimes \mathrm{id}) c \Sigma_Z^{\dagger} = c \Sigma_Z (\sigma_X \otimes \mathrm{id})$. Therefore, the result is a single 1-charge \mathcal{E}_{ω} . Therefore, we have $n_{\omega} \circ n_{\bar{\omega}} \simeq 1_{\mathcal{E}_{\omega}}$, and by symmetry we also find $n_{\bar{\omega}} \circ n_{\omega} = 1_{\mathcal{E}_{\bar{\omega}}}$. Note that, together, these fusions rules imply in particular that the 1-charges \mathcal{E}_{ω} are isomorphic. We summarise the non-trivial fusion

rules of domain walls we found below:


Let us now consider the fusion of the 1-charges themselves. First of all, we trivially have

$$\mathbb{1} \boxtimes \mathcal{E}_{\eta} \cong \mathcal{E}_{\eta} \cong \mathcal{E}_{\eta} \boxtimes \mathbb{1} , \qquad (23)$$

where $\eta \in \{1, \omega, \bar{\omega}\}$ labels a character of \mathbb{Z}_3 . Non-trivial fusion rules are those of the form $\mathcal{E}_{\eta} \boxtimes \mathcal{E}_{\mu}$, where $\eta, \mu \in \{1, \omega, \bar{\omega}\}$. For concreteness, let us focus on $\mathcal{E}_{\omega} \boxtimes \mathcal{E}_{\omega}$. Microscopically, such a fusion amounts to bringing the looplike defects very close to one another and consider the resulting system as a single (typically non-elementary) excitation. As such this fusion does not strictly take place on the boundary, rather we should think of one of the 1-charges as coming from the bulk, e.g.




The task is now to determine which excitation this system corresponds to. The qubit degrees of freedom between the two 1-charges can a priori be \uparrow or \downarrow , but it follows from the plaquette stabiliser conditions being satisfied that these interstitial qubits must all be in the same state. The previous configuration is thus equivalent to




On the one hand, we can think of qubits projected onto the spin- \uparrow subspace as corresponding to the presence of condensation defects \mathcal{E}_1 , and can thus be represented as wiggly red lines. On the other hand, we can think of the qubits projected onto the spin- \downarrow subspace as the result of acting with σ_X operators on qubits in the \uparrow state,

causing violations of 1-flatness stabiliser conditions associated with the surrounding plaquettes. But recall that topological defects are defined up to genuine line-like operators. It turns out the resulting magnetic excitations may be annihilated by acting with genuine line-like operators on the boundary. In order to appreciate the fact that the required operators are indeed genuinely line-like, we need to consider larger 1-charges following the pattern described in eq. (11). Given a defect configuration of 1charges and magnetic excitations, we consider genuine line-like operators of the form



Although the σ_X operators are sufficient to annihilate the 1-fluxes arising from the violations of the 1-flatness stabiliser conditions, the resulting configuration would not commute with some of the cube operators—giving rise to 2-fluxes—if not for the addition of the Γ operators. It follows from the commutation relations given in eq. (2) that the effect of such an operator—in addition to annihilating the magnetic excitations—is to turn the $c\Sigma_Z$ operators at the front into $c\Sigma_Z^{\dagger}$ operators. Applying a smaller version of this operator, we thus find that the configuration depicted in eq. (24) ends up being equivalent to



Since $\Sigma_Z^2 = \Sigma_Z^{\dagger}$, the first term in this decomposition amounts to a single 'thick' 1-charge $\mathcal{E}_{\bar{\omega}}$, whereas the second term amounts to a single 1-charge \mathcal{E}_1 . Putting everything together, we find the fusion rules $\mathcal{E}_{\omega} \boxtimes \mathcal{E}_{\omega} \cong \mathcal{E}_{\bar{\omega}} \boxplus \mathcal{E}_1$, and more generally

$$\mathcal{E}_{\eta} \boxtimes \mathcal{E}_{\mu} \cong \mathcal{E}_{\eta\mu} \boxplus \mathcal{E}_{\eta\bar{\mu}} , \qquad (26)$$

for any $\eta, \mu \in \{1, \omega, \bar{\omega}\}$, where we think of \mathcal{E}_{μ} as being the 1-charge at the front. We recover in particular the fusion rules $\mathcal{E}_1 \boxtimes \mathcal{E}_1 \cong \mathcal{E}_1 \boxplus \mathcal{E}_1$ of \mathbb{Z}_2 condensation defects [61, 64, 66].

Let us conclude this enumeration of the boundary excitations and their statistics with a couple of representation theoretic comments: We have already mentioned that a short string-like condensation defect \mathcal{E}_1 spanning over a single edge of the lattice amounts to the local operator creating a superposition $1_{\mathbb{1}} \oplus e$ of pairs of 0charges. In the same vein, the *tube algebra* analysis carried out in [75, 79] predicts that a *small* loop-like electric excitation, to which a point-like electric excitation may be attached, are labelled by irreducible representations of $\mathbb{D}_6 \simeq \mathbb{Z}_2 \ltimes \mathbb{Z}_3$. Recall that \mathbb{D}_6 possesses three irreducible representations, namely $\{\underline{0}, \underline{1}, \underline{2}\}$ referred to as the trivial, sign and standard representation. In that context, we can identify the 1-charges \mathcal{E}_1 , \mathcal{E}_{ω} and $\mathcal{E}_{\bar{\omega}}$ studied in this section as being labelled by the induced representations $\operatorname{Ind}_{\mathbb{Z}_3}^{\mathbb{D}_6}(1) \simeq \underline{0} \oplus \underline{1}$, $\operatorname{Ind}_{\mathbb{Z}_3}^{\mathbb{D}_6}(\omega) \simeq \underline{2}$ and $\operatorname{Ind}_{\mathbb{Z}_3}^{\mathbb{D}_6}(\bar{\omega}) \simeq \underline{2}$, respectively. Notice furthermore that this identification is compatible with the fusion rules (26) in virtue of $\underline{2} \otimes \underline{2} \simeq (\underline{0} \oplus \underline{1}) \oplus \underline{2}$.

SEC. III | Categorical representations

We show in this section how the categorical charges defined above as well as their statistics are organised into the monoidal bicategory of 2-representations of the 2-group $\mathbb{G}(2,3)$.

III.A. Preliminaries

In order to introduce the notion of 2-group 2representation employed in this manuscript, let us first review the category theoretical definition of a group representation. We invite the reader to consult [80] for basic definitions. Every finite group G can be thought as a one-object groupoid referred to as the delooping BG of G. It is the category with unique object \bullet and hom-set $\operatorname{Hom}(\mathsf{B} G) = G \ni \bullet \xrightarrow{g} \bullet$ such that the composition rule is provided by multiplication in G, i.e.

•
$$\xrightarrow{g_1}$$
 • $\xrightarrow{g_2}$ • = • $\xrightarrow{g_1g_2}$ • , $\forall g_1, g_2 \in G$. (27)

A representation of G is then defined as a *functor* from BG to Vec, where Vec is the category of complex vector spaces and linear maps:

$$\rho: BG \to \text{Vec}$$

: • $\mapsto \rho(\bullet) =: V$, (28)
:• $\xrightarrow{g} \bullet \mapsto V \xrightarrow{\rho(g)} V \in \text{End}_{\mathbb{C}}(V)$

i.e. it assigns a vector space V to the unique object • in BG, and a linear map $\rho(g)$ in $\operatorname{End}_{\mathbb{C}}(V)$ to every morphism $\bullet \xrightarrow{g} \bullet$ such that $\rho(g_1g_2) = \rho(g_1) \circ \rho(g_2)$ for any $g_1, g_2 \in G$. In the same vein, given two representations ρ and σ , an intertwiner $\vartheta : \rho \to \sigma$ between them is defined as a *natural transformation* between the corresponding functors, i.e. the assignment of a morphism $\vartheta_{\bullet} : \rho(\bullet) \to \sigma(\bullet)$ to the unique object \bullet in BG such that $\rho(g) \circ \vartheta_{\bullet} = \vartheta_{\bullet} \circ \sigma(g)$ for every $g \in G$. Putting everything together we obtain that the category $\operatorname{Rep}(G)$ of representations and intertwiners can be defined as the *functor category* $\operatorname{Fun}(\mathsf{B}G, \mathsf{Vec})$.

Let us now go up one level of abstraction so as to enter the realm of 2-categories, where, in addition to objects and (1-)morphisms, there are 2-morphisms between 1-morphisms. Given two objects $A, B \in Ob(\mathcal{C})$ in a 2-category \mathcal{C} , we denote by $\mathsf{Hom}_{\mathcal{C}}(A, B)$ the category of morphisms (hom-category) from A to B. As customary, objects and morphisms in hom-categories shall be referred to as 1- and 2-morphisms in \mathcal{C} , respectively. Given $A, B, C \in Ob(\mathcal{C})$, the horizontal composition functor of 1- and 2-morphisms is notated via \circ : $\operatorname{Hom}_{\mathcal{C}}(A,B) \times \operatorname{Hom}_{\mathcal{C}}(B,C) \to \operatorname{Hom}_{\mathcal{C}}(A,C)$, and the corresponding identity 1- and 2-morphisms via 1_A and 1_{1_A} , respectively. The vertical composition of 2-morphisms within a hom-category $\operatorname{Hom}_{\mathcal{C}}(A, B)$ is denoted by \cdot and the corresponding identity 1-morphisms by 1_f for any $f: A \to B$. Furthermore, we recall that a *bicategory* is a 2-category for which the associativity of the horizontal composition is weakened by a natural 2-isomorphisms referred to as the 1-associator.

Given a 2-group \mathbb{G} associated with a crossed module of the form $(G, H, \mathbb{1}_G : H \to {\mathbb{1}_G}, \triangleright)$ with H abelian,³ its delooping BG is a 2-groupoid consisting of a single object • with hom-category $\operatorname{Hom}_{BG}(\bullet, \bullet) = \mathbb{G}$ such that horizontal and vertical compositions are provided by the monoidal structure in \mathbb{G} and the composition of (1-)morphisms in \mathbb{G} , respectively [67]. More concretely, 1-morphisms are of the form $\bullet \xrightarrow{g} \bullet$, with $g \in G$, and compose via group multiplication in G, whereas 2-morphisms are of the form

•
$$\forall g \in G, h \in H$$
. (29)

Vertical composition of 2-morphisms is defined via group multiplication in H as

with $h := h_1 + h_2$ for any $g \in G$ and $h_1, h_2 \in H$, whereas horizontal composition of 2-morphisms is provided by group multiplication in $G \ltimes H$ as

$$\bullet \underbrace{ \left(\begin{array}{c} g_1 \\ h_1 \\ g_1 \end{array}\right)}_{g_1} \bullet \underbrace{ \left(\begin{array}{c} g_2 \\ h_2 \\ g_2 \end{array}\right)}_{g_2} \bullet = \bullet \underbrace{ \left(\begin{array}{c} g_1 g_2 \\ g_1 g_2 \\ g_1 g_2 \end{array}\right)}_{g_1 g_2} \bullet$$
(31)

with $\tilde{h} := h_1 + (g_1 \triangleright h_2)$ for any $g_1, g_2 \in G$ and $h_1, h_2 \in H$.

In order to introduce a notion of 2-representations, we require a *categorification* of the notion of vector spaces. We denote by 2Vec the 2-category of finite semi-simple linear categories, linear functors and natural transformations. As customary, we shall refer to objects in 2Vec as 2-vector spaces. This bicategory can be further equipped with a monoidal structure via the Deligne tensor product \boxtimes of abelian categories. Note that 2Vec has a unique simple object, namely Vec.

Given the above, we define the bicategory $2\text{Rep}(\mathbb{G})$ of 2-representations, 1-interwiners and 2-intertwiners of a 2-group \mathbb{G} as the bicategory $2\text{Fun}(\mathbb{BG}, 2\text{Vec})$ of *pseudofunctors*, *pseudonatural transformations* and *modifications*. Instead of providing the general definition of these notions, let us immediately unpack the bicategory $2\text{Fun}(\mathbb{BG}, 2\text{Vec})$.

A 2-representation ρ of \mathbb{G} is a map

$$\rho: \quad B\mathbb{G} \quad \to 2 \operatorname{Vec}$$

$$: \quad \bullet \quad \mapsto \quad \rho(\bullet) =: \mathcal{V}$$

$$: \quad \bullet \stackrel{g}{\to} \bullet \quad \mapsto \quad \mathcal{V} \stackrel{\rho(g)}{\longrightarrow} \mathcal{V} \in \operatorname{End}(\mathcal{V}) \quad , \quad (32)$$

$$: \quad \bullet \stackrel{g}{\longrightarrow} \bullet \quad \mapsto \quad \mathcal{V} \stackrel{\rho(g)}{\longrightarrow} \stackrel{\rho(g)}{\longrightarrow} \mathcal{V} \in \operatorname{End}_{\operatorname{End}(\mathcal{V})}(\rho(g))$$

where \mathcal{V} is a finite linear semisimple (1-)category, $\rho(g)$ are endofunctors of \mathcal{V} and $\rho(h) : \rho(g) \Rightarrow \rho(g)$ are natural transformations. Furthermore, ρ is required to strictly preserve vertical and horizontal compositions of 2-morphisms as well as to preserve composition of 1-morphisms up to natural isomorphisms. In other words, we have

$$\rho(h_1) \cdot \rho(h_2) = \rho(h_1 \cdot h_2) \tag{33}$$

$$\rho(h_1) \circ \rho(h_2) = \rho(h_1 \circ h_2),$$
(34)

for any $h_1: g_1 \Rightarrow g_1, h_2: g_2 \Rightarrow g_2$ labelled by $h_1, h_2 \in H$, and ρ assigns to every pair of 1-morphisms labelled by $g_1, g_2 \in G$ a natural 2-isomorphism

$$\rho_{g_1,g_2}:\rho(g_1)\circ\rho(g_2) \xrightarrow{\sim} \rho(g_1g_2), \qquad (35)$$

which is required to fulfil

$$[1_{\rho(g_1)} \circ \rho_{g_2,g_3}] \cdot \rho_{g_1,g_2g_3} = [\rho_{g_1,g_2} \circ 1_{\rho(g_3)}] \cdot \rho_{g_1g_2,g_3} , \quad (36)$$
for any $g_1, g_2, g_3 \in G$.

³ Crossed modules with arbitrary group homomorphisms $\partial: G \to H$ with H non-abelian can be treated in the exact same way, but since we ultimately focus on $\mathbb{G}(2,3)$ for which ∂ is trivial, we have no need dealing with the extra technicalities.

Given two 2-representations ρ and σ of \mathbb{G} , a 1intertwiner $\vartheta : \rho \to \sigma$ between them consists of an object ϑ_{\bullet} in Fun $(\rho(\bullet), \sigma(\bullet))$ together with a collection of natural 2-isomorphisms ϑ_{-} defined via

$$\vartheta_g:\rho(g)\circ\vartheta_{\bullet} \xrightarrow{\sim} \vartheta_{\bullet}\circ\sigma(g)\,,\quad g\in G\,. \tag{37}$$

such that $\vartheta_{\mathbb{1}_G} = 1_{\vartheta_{\bullet}}$. This collection of 2-isomorphisms is subject to a coherence relation ensuring compatibility with the horizontal composition of 1-morphisms in \mathbb{C} , namely

$$\begin{aligned} & [\rho_{g_1,g_2} \circ 1_{\vartheta_{\bullet}}] \cdot \vartheta_{g_1g_2} \\ &= [1_{\rho(g_1)} \circ \vartheta_{g_2}] \cdot [\vartheta_{g_1} \circ 1_{\sigma(g_2)}] \cdot [1_{\vartheta_{\bullet}} \circ \sigma_{g_1,g_2}] \end{aligned}$$
(38)

for any $g_1, g_2 \in G$. Moreover, the naturality condition stipulates that

$$[\rho(h) \circ 1_{\vartheta_{\bullet}}] \cdot \vartheta_g = \vartheta_g \cdot [1_{\vartheta_{\bullet}} \circ \sigma(h)]$$
(39)

for any 2-morphism $h: g \Rightarrow g$ in \mathbb{G} .

Finally, given two 1-intertwiners ϑ , $\tilde{\vartheta} : \rho \to \sigma$ between two 2-representations ρ and σ of \mathbb{G} , a 2-intertwiner Θ : $\vartheta \Rightarrow \tilde{\vartheta}$ between them consists of a natural transformation $\Theta_{\bullet} : \vartheta_{\bullet} \Rightarrow \tilde{\vartheta}_{\bullet}$ in $\mathsf{Fun}(\rho(\bullet), \sigma(\bullet))$ such that

$$\vartheta_g \cdot [\Theta_{\bullet} \circ 1_{\sigma(g)}] = [1_{\rho(g)} \circ \Theta_{\bullet}] \cdot \tilde{\vartheta}_g \tag{40}$$

is fulfilled for any $g \in G$.

The definitions above are still somewhat terse so let us unfold them further before specialising to $\mathbb{G}(2,3)$. Let us begin with 2-representations. A finite semisimple 1category \mathcal{V} together with endofunctors $\rho(g): \mathcal{V} \to \mathcal{V}$, for every $g \in G$, and natural 2-isomorphisms $\rho_{g_1,g_2}: \rho(g_1) \circ$ $\rho(g_2) \Rightarrow \rho(g_1g_2)$ in 2Vec satisfying the coherence relation given in eq. (35) define a right *module* category over the category Vec_G of *G*-graded vector spaces. Recall that simple objects in Vec_G are the one-dimensional vector spaces \mathbb{C}_g , for every $g \in G$, such that $\mathbb{C}_{g_1} \otimes \mathbb{C}_{g_2} \simeq \mathbb{C}_{g_2}$ and $\operatorname{Hom}_{\operatorname{Vec}_G}(\mathbb{C}_{g_1}, \mathbb{C}_{g_2}) = \delta_{g_1,g_2}\mathbb{C}$. Introducing the notations $\triangleleft: \mathcal{V} \times \operatorname{Vec}_G \to \mathcal{V}$, whereby $M \triangleleft \mathbb{C}_g := \rho(g)(M)$ for every $M \in \operatorname{Ob}(\mathcal{V})$, and $\alpha^{\triangleleft}_{M,\mathbb{C}_{g_1},\mathbb{C}_{g_2}} := (\rho_{g_1,g_2})_M$, it follows from the 2-cocycle condition satisfied by ρ_{g_1,g_2} that

$$\begin{aligned} \alpha^{\triangleleft}_{M \triangleleft \mathbb{C}_{g_1}, \mathbb{C}_{g_2}, \mathbb{C}_{g_3}} \circ \alpha^{\triangleleft}_{M, \mathbb{C}_{g_1}, \mathbb{C}_{g_2g_3}} \\ &= (\alpha^{\triangleleft}_{M, \mathbb{C}_{g_1}, \mathbb{C}_{g_2}} \triangleleft \mathbb{1}_{\mathbb{C}_{g_3}}) \circ \alpha^{\triangleleft}_{M, \mathbb{C}_{g_1g_2}, \mathbb{C}_{g_3}} \circ \mathbb{1}_{M \triangleleft \mathbb{C}_{g_1g_2g_3}} \end{aligned}$$
(41)

holds for all $g_1, g_2, g_3 \in G$ and $M \in \operatorname{Ob}(\mathcal{V})$, and thus $\alpha^{\triangleleft} : (-\triangleleft -) \triangleleft - \rightarrow - \triangleleft (- \otimes -)$ defines a module associator for \mathcal{V} with respect to \triangleleft . Putting everything together, the triple $(\mathcal{V}, \triangleleft, \alpha^{\triangleleft})$ does define a right module category over Vec_G . In virtue of $\rho(h)$ defining a natural transformation between $\rho(g)$ and itself, together with the compatibility condition eq. (33), we find that ρ further assigns to every object $M \in \operatorname{Ob}(\mathcal{V})$ a representation $\rho(-)_M : H \to \operatorname{End}_{\mathcal{V}}(\rho(g)(M))$ of H on $\rho(g)(M) \in \operatorname{Ob}(\mathcal{V})$

for any $g \in G$, and to every 1-morphism in \mathcal{V} an intertwiner between the corresponding representations. The compatibility condition (34) finally requires that

$$\rho(g \triangleright -)_M = \rho(-)_{\rho(g)(M)} \tag{42}$$

for every $g \in G$ and $M \in Ob(\mathcal{V})$.

Let us now examine further 1-intertwiners at the light of this interpretation of 2-representations. Given two 2-representations ρ and σ of \mathbb{G} with underlying Vec_{G} module categories $(\mathcal{V}, \triangleleft, \alpha^{\triangleleft})$ and $(\mathcal{W}, \triangleleft, \alpha^{\triangleleft})$, respectively, an object ϑ_{\bullet} in $\operatorname{Fun}(\mathcal{V}, \mathcal{W})$ together with a collection of natural 2-isomorphisms $\vartheta_g : \rho(g) \circ \vartheta_{\bullet} \Rightarrow \vartheta_{\bullet} \circ \sigma(g)$ satisfying eq. (38) defines a Vec_G -module functor from \mathcal{V} to \mathcal{W} . Introducing the notation, $\omega_{M,\mathbb{C}_g} := (\vartheta_g)_M$, it follows from eq. (38) that

$$\vartheta_{\bullet}(\alpha_{M,\mathbb{C}_{g_{1}},\mathbb{C}_{g_{2}}}^{\triangleleft}) \circ \omega_{M,\mathbb{C}_{g_{1}g_{2}}}$$

$$= \omega_{M \triangleleft \mathbb{C}_{g_{1}},\mathbb{C}_{g_{2}}} \circ (\omega_{M,\mathbb{C}_{g_{1}}} \triangleleft 1_{\mathbb{C}_{g_{2}}}) \circ \alpha_{\vartheta_{\bullet}(M),\mathbb{C}_{g_{1}},\mathbb{C}_{g_{2}}}^{\triangleleft}$$

$$(43)$$

holds for every $g_1, g_2 \in G$ and $M \in Ob(\mathcal{V})$, and thus the pair $(\vartheta_{\bullet}, \omega)$ with $\omega : \vartheta_{\bullet}(\neg \triangleleft \neg) \rightarrow \vartheta_{\bullet}(\neg) \triangleleft \neg$ does define a Vec_G-module functor. The category of such module functors over Vec_G is denoted by Fun_{Vec_G}(\mathcal{V}, \mathcal{W}). The naturality condition (39) further stipulates that

$$\vartheta_{\bullet}(\rho(h)_M) \circ \omega_{M,\mathbb{C}_g} = \omega_{M,\mathbb{C}_g} \circ \sigma(h)_{\vartheta_{\bullet}(M)}$$
(44)

must hold for every $h: g \Rightarrow g$ and $M \in Ob(\mathcal{V})$, and thus not all objects in $\mathsf{Fun}_{\mathsf{Vec}_G}(\mathcal{V}, \mathcal{W})$ necessarily provide a 1intertwiner between the corresponding 2-representations.

Before computing explicitly the bicategory $2\text{Rep}(\mathbb{G}(2,3))$, let us remark that in the limiting case where we forget about the group \mathbb{Z}_3 , which amounts to treating the group \mathbb{Z}_2 as a 2-group, the construction presented in this section yields the monoidal bicategory $2\text{Rep}(\mathbb{Z}_2)$ studied by Kong et al. in [64] in the context of the (3+1)d toric code.

III.B. 2-group 2-representations

Let us now apply the definitions spelt out above to the case of $\mathbb{G}(2,3)$ beginning with the computation of all the (simple) 2-representations.⁴ Note that a similar derivation was carried out by Barrett and Mackaay in [52] but their definition of $2\text{Rep}(\mathbb{G}(2,3))$ differs a little it from the one employed in this manuscript.

We established earlier that a 2-representation of $\mathbb{G}(2,3)$ is labelled—amongst other things—by a choice of right module category over $\mathsf{Vec}_{\mathbb{Z}_2}$. There are only two indecomposable $\mathsf{Vec}_{\mathbb{Z}_2}$ -module categories, namely $\mathsf{Vec}_{\mathbb{Z}_2}$ itself via the monoidal product in $\mathsf{Vec}_{\mathbb{Z}_2}$ and Vec via the forgetful functor $\mathsf{Vec}_{\mathbb{Z}_2} \to \mathsf{Vec}$ [54]. Notice that both module

⁴ We are referring here to the notion of semi-simplicity in the sense of Douglas and Reutter [81].

categories have module associators α^{\triangleleft} given by the identity morphisms. Let us treat these two cases separately:

* Let ρ be a 2-representation of $\mathbb{G}(2,3)$ whose underlying $\operatorname{Vec}_{\mathbb{Z}_2}$ -module category ($\mathcal{V} := \rho(\bullet), \triangleleft, \alpha^{\triangleleft}$) is Vec. The 2-representation ρ assigns to the unique simple object in Vec, namely \mathbb{C} , a representation $\rho(-)_{\mathbb{C}} : \mathbb{Z}_3 \to$ $\operatorname{End}_{\mathcal{V}}(\rho(g)(\mathbb{C}))$ for any $g \in \mathbb{Z}_2$. Since $\rho(g)(\mathbb{C}) \simeq \mathbb{C}$, $\rho(-)_{\mathbb{C}}$ amounts to a *character* on \mathbb{C} . This one-dimensional representation of \mathbb{Z}_3 is further required to fulfil $\rho(g \triangleright -)_{\mathbb{C}} =$ $\rho(-)_{\mathbb{C}}$ for every $g \in \mathbb{Z}_2$, forcing $\rho(-)_{\mathbb{C}}$ to be the trivial representation of \mathbb{Z}_3 . We notate the resulting 2representation of $\mathbb{G}(2,3)$ via 1, which was referred to as the trivial 1-charge in the previous section.

* Let ρ be a 2-representation of $\mathbb{G}(2,3)$ whose underlying $\operatorname{Vec}_{\mathbb{Z}_2}$ -module category $(\mathcal{V}, \triangleleft, \alpha^{\triangleleft})$ is $\operatorname{Vec}_{\mathbb{Z}_2}$. The 2-representation ρ assigns to the simple objects \mathbb{C}_{+1} and \mathbb{C}_{-1} in $\mathsf{Vec}_{\mathbb{Z}_2}$ representations $\rho(-)_{\mathbb{C}_{+1}}$: $\mathbb{Z}_3 \to$ $\operatorname{End}_{\mathcal{V}}(\rho(g)(\mathbb{C}_{\pm 1}))$ for any $g \in \mathbb{Z}_2$. Since $\rho(g)(\mathbb{C}_{\pm 1}) =$ $\mathbb{C}_{\pm 1} \triangleleft \mathbb{C}_g \simeq \mathbb{C}_{\pm g}$ and $\operatorname{End}_{\mathcal{V}}(\mathbb{C}_{\pm g}) \simeq \mathbb{C}, \ \rho(-)_{\mathbb{C}_{\pm 1}}$ are \mathbb{C} -valued characters of \mathbb{Z}_3 . These characters are further required to fulfill eq. (42), i.e. $\rho(g \triangleright -)_{\mathbb{C}_{+1}} =$ $\rho(-)_{\mathbb{C}_{\pm 1} \triangleleft \mathbb{C}_g} = \rho(-)_{\mathbb{C}_{\pm g}}$. Choosing $g = -1 \in \mathbb{Z}_2$, this imposes that $\rho(h^{-1})_{\mathbb{C}_{+1}} = \rho(h)_{\mathbb{C}_{-1}}$ for every $h \in \mathbb{Z}_3$ and thus $\rho(-)_{\mathbb{C}_{-1}} = \rho(-)_{\mathbb{C}_{+1}}^{-1}$. We thus obtain three distinct 2-representations labelled by a complex character of \mathbb{Z}_3 . These correspond to the non-trivial 1-charges $\mathcal{E}_1, \mathcal{E}_{\omega}$ and $\mathcal{E}_{\bar{\omega}}$ derived in the previous section. Notice that condition (42) precisely translates the fact that we required $c\Sigma_Z$ gates in order to construct membrane operators creating 1-charges along their boundaries.

III.C. 1-intertwiners

We pursue our analysis with the derivation of the 1intertwiners between the four 2-representations defined above:

1 → 1: A 1-intertwiner between the 2-representation 1 and itself is specified—amongst other things—by a choice of Vec_{Z2}-module endofunctors of Vec, i.e. an object in Fun_{VecZ2} (Vec, Vec). But we can easily show that the category Fun_{VecZ2} (Vec, Vec) is equivalent as a monoidal category to Rep(Z₂) [80]. Such a Vec_{Z2}-module functor is further required to fulfil eq. (44), but it turns out to be trivial in that case given that $\rho(-)_{\mathbb{C}}$ corresponds to the trivial representation of Z₃. We thus identify two simple 1-intertwiners 1 → 1 denoted by 1₁ and *e*, associated with the two simple objects in Rep(Z₂). These correspond to the 0-charges constructed in the previous section, thought as domain walls between two trivial 1charges.

 $\mathcal{E}_1 \to \mathcal{E}_1$: Such a 1-intertwiner is now specified by a choice of $\mathsf{Vec}_{\mathbb{Z}_2}$ -module endofunctors of $\mathsf{Vec}_{\mathbb{Z}_2}$, whereby $\mathsf{Fun}_{\mathsf{Vec}_{\mathbb{Z}_2}}(\mathsf{Vec}_{\mathbb{Z}_2},\mathsf{Vec}_{\mathbb{Z}_2}) \cong \mathsf{Vec}_{\mathbb{Z}_2}$. Similarly to the previous scenario, since the 2-representation \mathcal{E}_1 assigns the

trivial character of \mathbb{Z}_3 to both simple objects in $\mathsf{Vec}_{\mathbb{Z}_2}$, the naturality condition (44) happens to be trivially satisfied. We thus identify two simple 1-intertwiners $\mathcal{E}_1 \to \mathcal{E}_1$ denotes by $1_{\mathcal{E}_1}$ and n, associated with the two simple objects in $\mathsf{Vec}_{\mathbb{Z}_2}$. These correspond to the two types of domain walls between condensation defects constructed in the previous section.

 $\mathcal{E}_{\omega/\bar{\omega}} \to \mathcal{E}_{\omega/\bar{\omega}}$: As for the previous case, such a 1intertwiner is specified by an object in $\operatorname{Vec}_{\mathbb{Z}_2}$ thought as a module endofunctor of $\operatorname{Vec}_{\mathbb{Z}_2}$ over itself. But the naturality condition (44) is not trivial anymore. There are two simple objects in $\operatorname{Vec}_{\mathbb{Z}_2}$, namely \mathbb{C}_{+1} and \mathbb{C}_{-1} , and the corresponding module functors are given by $\mathbb{C}_{+1}\otimes-$ and $\mathbb{C}_{-1}\otimes-$, respectively, with the isomorphisms ω being identity morphisms. Focusing on $\mathbb{C}_{-1}\otimes -$, the naturality condition stipulates that we must have $\mathbb{1}_{\mathbb{C}_{-1}}\otimes\rho(h)_{\mathbb{C}_{\pm 1}}=\rho(h)_{\mathbb{C}_{-1}\otimes\mathbb{C}_{\pm 1}}$, i.e. $\rho(h)_{\mathbb{C}_{\pm 1}}=\rho(h)_{\mathbb{C}_{\mp 1}}$, which is not satisfied whenever $h\neq 0\in\mathbb{Z}_3$ given that $\rho(h)_{\mathbb{C}_{\mp 1}}=\rho(h^{-1})_{\mathbb{C}_{\pm 1}}$. On other hand, the naturality condition is always satisfied for $\mathbb{C}_{+1}\otimes-$. It follows that there is a unique simple 1-intertwiner $\mathcal{E}_{\omega/\bar{\omega}}\to \mathcal{E}_{\omega/\bar{\omega}}$, which we notate via $\mathbb{1}_{\mathcal{E}_{w}/\bar{\omega}}$.

 $\mathcal{E}_{\omega/\bar{\omega}} \to \mathcal{E}_{\bar{\omega}/\omega}$: This case is very similar to the previous one with the difference that the module endofunctor $\mathbb{C}_{+1} \otimes -$ over itself is the one that does not satisfy the naturality condition. Indeed, the naturality condition stipulates that $1_{\mathbb{C}_{+1}} \otimes \rho(h)_{\mathbb{C}_{\pm 1}} = \sigma(h)_{\mathbb{C}_{\pm 1}}$ with $\sigma(h)_{\mathbb{C}_{\pm 1}} = \rho(h)_{\mathbb{C}_{\pm 1}}^{-1}$. On the other hand, the naturality condition is always satisfied for $\mathbb{C}_{-1} \otimes -$. It follows that there is a unique simple 1-intertwiner $\mathcal{E}_{\omega/\bar{\omega}} \to \mathcal{E}_{\omega/\bar{\omega}}$, which we notate via $n_{\omega/\bar{\omega}}$.

 $\mathcal{E}_1 \to \mathbb{1}$: This scenario is different than the previous ones in that the $\mathsf{Vec}_{\mathbb{Z}_2}$ -module categories underlying the 2-representations \mathcal{E}_1 and $\mathbb{1}$ differ so that such a 1-interviner is now specified by a choice of objects in $\operatorname{Fun}_{\operatorname{Vec}_{\mathbb{Z}_2}}(\operatorname{Vec}_{\mathbb{Z}_2},\operatorname{Vec})$. But the category $\operatorname{\mathsf{Fun}}_{\operatorname{\mathsf{Vec}}_{\mathbb{Z}_2}}(\operatorname{\mathsf{Vec}}_{\mathbb{Z}_2},\operatorname{\mathsf{Vec}})$ is equivalent to $\operatorname{\mathsf{Vec}}$, whereby the unique object in Vec amounts to the forgetful functor $\mathsf{Vec}_{\mathbb{Z}_2} \to \mathsf{Vec}$. The naturality condition further stipulates that $\rho(h)_{\mathbb{C}_{+1}} = \sigma(h)_{\mathbb{C}}$ for every $h \in \mathbb{Z}_3$, which is always satisfied since both \mathcal{E}_1 and $\mathbb{1}$ are associated with the trivial character of \mathbb{Z}_3 . We denote this unique 1intertwiner $\mathcal{E}_1 \to \mathbb{1}$ by x^{\dashv} . By transposition, we find a unique 1-intertwiner $\mathbb{1} \to \mathcal{E}_1$, which we denote by x^{\vdash} . These provide the domain walls between the condensation defect and the vacuum found in the previous section.

It follows immediately from the previous derivations that there are no 1-intertwiners of the form $\mathcal{E}_1 \to \mathcal{E}_{\omega/\bar{\omega}}$, $\mathbb{1} \to \mathcal{E}_{\omega/\bar{\omega}}$, or transpositions thereof, as trivial and nontrivial characters of \mathbb{Z}_3 are incompatible preventing the naturality condition (44) from being satisfied. This is consistent with the absence of domain walls between the corresponding 1-charges witnessed in the previous section and concludes our analysis of the 1-intertwiners.

III.D. Composition rule

We established above that the domain walls derived in the previous section are provided by the 1-intertwiners in $2\text{Rep}(\mathbb{G}(2,3))$. We shall now confirm that the fusion of these domain walls in encoded into the composition rule of the corresponding 1-intertwiners. Generally speaking, given two 1-intertwiners $\vartheta : \rho \to \sigma$ and $\tilde{\vartheta} : \sigma \to \varsigma$, their horizontal composition is defined as the 1-intertwiner $\vartheta \circ \tilde{\vartheta}$ such that

$$\begin{aligned} (\vartheta \circ \tilde{\vartheta})_{\bullet} &:= \vartheta_{\bullet} \circ \tilde{\vartheta}_{\bullet} \in \mathsf{Fun}(\rho(\bullet),\varsigma(\bullet)) \\ (\vartheta \circ \tilde{\vartheta})_g &:= (\vartheta_g \circ 1_{\tilde{\vartheta}_{\bullet}}) \cdot (1_{\vartheta_{\bullet}} \circ \tilde{\vartheta}_g) \end{aligned}$$
(45)

for every $g \in G$. This simply expresses the composition of the $\operatorname{Vec}_{\mathbb{Z}_2}$ -module functors underlying the 1-intertwiners. In particular, the defining isomorphism associated with the module functor $(\vartheta \circ \vartheta)_{\bullet}$ is given by $((\vartheta \circ \tilde{\vartheta})_g)_M :=$ $\tilde{\vartheta}_{\bullet}(\omega_{M,\mathbb{C}_g}) \circ \tilde{\omega}_{\vartheta \bullet (M),\mathbb{C}_g}$ for any $M \in \operatorname{Ob}(\rho(\bullet))$ and $g \in \mathbb{Z}_2$.

* We established above that simple 1-intertwiners $1 \to 1$ are in one-to-one correspondence with simple objects in $\operatorname{Fun}_{\operatorname{Vec}_{\mathbb{Z}_2}}(\operatorname{Vec},\operatorname{Vec}) \cong \operatorname{Rep}(\mathbb{Z}_2)$. It follows from the definition of $((\vartheta \circ \tilde{\vartheta})_g)_M$ given above that (horizontal) composition of these 1-intertwiners is provided by the monoidal structure in $\operatorname{Rep}(\mathbb{Z}_2)$, and thus we have $e \circ e \simeq 1_1$. Similarly composition of 1-intertwiners of the form $\mathcal{E}_1 \to \mathcal{E}_1$ is provided by the monoidal structure in $\operatorname{Vec}_{\mathbb{Z}_2}$ and thus $n \circ n \simeq 1_{\mathcal{E}_1}$. These composition rules are compatible with the fusion rules of the corresponding domain walls derived in the previous section.

* Analogously to the previous cases, the 1-intertwiners $x^{\dashv} : \mathcal{E}_1 \to \mathbb{1}$ and $x^{\vdash} : \mathbb{1} \to \mathcal{E}_1$ are provided by the unique simple objects in $\operatorname{Fun}_{\operatorname{Vec}_{\mathbb{Z}_2}}(\operatorname{Vec}_{\mathbb{Z}_2}, \operatorname{Vec}) \cong \operatorname{Vec}$ and $\operatorname{Fun}_{\operatorname{Vec}_{\mathbb{Z}_2}}(\operatorname{Vec}_{\mathbb{Z}_2}, \operatorname{Vec}) \cong \operatorname{Vec}$ and $x^{\vdash} \circ x^{\dashv}$ and $x^{\dashv} \circ x^{\vdash}$ of the 1-intertwiners are then dictated by the compositions of the corresponding $\operatorname{Vec}_{\mathbb{Z}_2}$ module functors [54]. By definition $x^{\vdash} \circ x^{\dashv}$ is an object in $\operatorname{Fun}_{\operatorname{Vec}_{\mathbb{Z}_2}}(\operatorname{Vec}, \operatorname{Vec}) \cong \operatorname{Rep}(\mathbb{Z}_2)$ and can be shown to correspond to the *regular* representation of \mathbb{Z}_2 . It follows that $x^{\vdash} \circ x^{\dashv} \simeq 1_{\mathbb{I}} \oplus e$. Similarly, we find that $x^{\dashv} \circ x^{\vdash} \circ 1_{\mathcal{E}_1} \oplus n$ as an object in $\operatorname{Fun}_{\operatorname{Vec}_{\mathbb{Z}_2}}(\operatorname{Vec}_{\mathbb{Z}_2}, \operatorname{Vec}_{\mathbb{Z}_2}) \cong \operatorname{Vec}_{\mathbb{Z}_2}$.

* Let us finally consider the composite $n_{\omega} \circ n_{\bar{\omega}}$. Since the composition of 1-intertwiners boils down to the composition of the corresponding module functors, we already know that we must have $n_{\omega} \circ n_{\bar{\omega}} \simeq 1_{\mathcal{E}_{\omega}}$ as it is the unique 1-intertwiner $\mathcal{E}_{\omega} \to \mathcal{E}_{\omega}$, but let us check explicitly how the composition behaves with respect to the naturality condition (44). Recall that both n_{ω} and $n_{\bar{\omega}}$ corresponds to $\mathbb{C}_{-1} \otimes -$ as the unique simple $\operatorname{Vec}_{\mathbb{Z}_2}$ -module endofunctor of $\operatorname{Vec}_{\mathbb{Z}_2}$ satisfying the naturality condition. It follows from eq. (45) that $n_{\omega} \circ n_{\bar{\omega}}$ corresponds to $\mathbb{C}_{+1} \otimes -$ and that the naturality condition for 1-intertwiners $\mathcal{E}_{\omega} \to \mathcal{E}_{\omega}$ is satisfied as $\rho(h)_{\mathbb{C}_{\pm 1}} = \rho(h)_{\mathbb{C}_{-1} \otimes (\mathbb{C}_{-1} \otimes \mathbb{C}_{\pm 1})}$ for every $h \in \mathbb{Z}_3$.

III.E. Monoidal structure

We close our dictionary between boundary excitations of the higher gauge model and the bicategory of 2representations of $\mathbb{G}(2,3)$ with the recovery of the fusion of the 1-charges from the monoidal structure in $2\text{Rep}(\mathbb{G}(2,3))$.

Recall that the simple objects in $2\operatorname{Rep}(\mathbb{G}(2,3))$ associated with the 1-charges \mathcal{E}_1 , \mathcal{E}_{ω} and $\mathcal{E}_{\overline{\omega}}$ are specified by the $\operatorname{Vec}_{\mathbb{Z}_2}$ -module category $\operatorname{Vec}_{\mathbb{Z}_2}$ and the characters of \mathbb{Z}_3 labelled by 1, ω and $\overline{\omega}$, respectively. A (right) $\operatorname{Vec}_{\mathbb{Z}_2}$ -module structure can be readily defined on the product $\operatorname{Vec}_{\mathbb{Z}_2} \boxtimes \operatorname{Vec}_{\mathbb{Z}_2}$ via $(M \boxtimes N) \triangleleft \mathbb{C}_g := (M \triangleleft \mathbb{C}_g) \boxtimes (N \triangleleft \mathbb{C}_g) = (M \otimes \mathbb{C}_g) \boxtimes (N \otimes \mathbb{C}_g)$ for any $M, N \in \operatorname{Ob}(\operatorname{Vec}_{\mathbb{Z}_2})$ and $g \in G$ [82]. The resulting module category decomposes as $\operatorname{Vec}_{\mathbb{Z}_2} \boxplus \operatorname{Vec}_{\mathbb{Z}_2}$, where the first copy of $\operatorname{Vec}_{\mathbb{Z}_2}$ has simple objects $\mathbb{C}_{\pm 1} \boxtimes \mathbb{C}_{\pm 1}$ and the latter has simple objects $\mathbb{C}_{\pm 1} \boxtimes \mathbb{C}_{\mp 1}$. This already provides the fusion rule $\mathcal{E}_1 \boxtimes \mathcal{E}_1 \cong \mathcal{E}_1 \boxplus \mathcal{E}_1$. It remains to understand how this tensor structure interacts with non-trivial characters of \mathbb{Z}_3 .

Consider the 2-representations $\rho \equiv \mathcal{E}_{\eta}$ and $\sigma \equiv \mathcal{E}_{\mu}$, where $\eta, \mu \in \{1, \omega, \bar{\omega}\}$ label characters of \mathbb{Z}_3 . Following the derivation above, we are looking for 2-representations ς_1 and ς_2 such that $\rho \boxtimes \sigma \cong \varsigma_1 \boxplus \varsigma_2$ with $\varsigma_1(\bullet) = \varsigma_2(\bullet) =$ $\operatorname{Vec}_{\mathbb{Z}_2}$ as $\operatorname{Vec}_{\mathbb{Z}_2}$ -module categories. Moreover, we established that simple objects in $\varsigma_1(\bullet)$ are given by $\mathbb{C}_{\pm 1} \boxtimes \mathbb{C}_{\pm 1}$, whereas simple objects in $\varsigma_2(\bullet)$ are given by $\mathbb{C}_{\pm 1} \boxtimes \mathbb{C}_{\pm 1}$. The representations of \mathbb{Z}_3 associated with ς_1 and ς_2 are then defined via $\varsigma_1(-)_{\mathbb{C}_{\pm 1} \boxtimes \mathbb{C}_{\pm 1}} := \rho(-)_{\mathbb{C}_{\pm 1}} \otimes \sigma(-)_{\mathbb{C}_{\pm 1}}$ and $\varsigma_2(-)_{\mathbb{C}_{\pm 1} \boxtimes \mathbb{C}_{\mp 1}} := \rho(-)_{\mathbb{C}_{\pm 1}} \otimes \sigma(-)_{\mathbb{C}_{\pm 1}}$, respectively. Let us check that these representations are well-defined. In particular, they must be compatible with the action of $\operatorname{Vec}_{\mathbb{Z}_2}$ on $\varsigma_1(\bullet)$ and $\varsigma_2(\bullet)$. Recall that we have $\rho(g \triangleright -)_{\mathbb{C}_{\pm 1}} = \rho(-)_{\mathbb{C}_{\pm g}}$ and $\sigma(g \triangleright -)_{\mathbb{C}_{\pm 1}} = \sigma(-)_{\mathbb{C}_{\pm g}}$ for every $g \in G$. It follows that we have

$$\varsigma_{1}(g \triangleright -)_{\mathbb{C}_{\pm 1} \boxtimes \mathbb{C}_{\pm 1}} = \rho(g \triangleright -)_{\mathbb{C}_{\pm 1}} \otimes \sigma(g \triangleright -)_{\mathbb{C}_{\pm 1}} \qquad (46)$$
$$= \rho(-)_{\mathbb{C}_{\pm 1} \otimes \mathbb{C}_{g}} \otimes \sigma(-)_{\mathbb{C}_{\pm 1} \otimes \mathbb{C}_{g}}$$
$$= \varsigma_{1}(-)_{(\mathbb{C}_{\pm 1} \otimes \mathbb{C}_{g}) \boxtimes (\mathbb{C}_{\pm 1} \otimes \mathbb{C}_{g})}$$
$$= \varsigma_{1}(-)_{(\mathbb{C}_{\pm 1} \boxtimes \mathbb{C}_{\pm 1}) \triangleleft \mathbb{C}_{g}}$$

for every $g \in G$, as requested, and similarly for ς_2 . Recall finally that while $\varsigma_1(-)_{\mathbb{C}_{+1}}$ corresponds to the character $\eta, \varsigma_1(-)_{\mathbb{C}_{-1}}$ corresponds to $\bar{\eta}$, and similarly for ς_2 with respect to μ . Putting everything together, we recover the fusion rules $\mathcal{E}_{\eta} \boxtimes \mathcal{E}_{\mu} \cong \mathcal{E}_{\eta\mu} \boxplus \mathcal{E}_{\eta\bar{\mu}}$, for any $\eta, \mu \in$ $\{1, \omega, \bar{\omega}\}$. This concludes our analysis of the bicategory $2\mathsf{Rep}(\mathbb{G}(2, 3))$.

SEC. IV | **Discussion**

Relying on the paradigm that global symmetry operators in any quantum model correspond to topological defects, we explored in this manuscript the (categorical) charges associated with a simple case of higher group symmetry. More precisely, we studied the excitation content of the symmetry-preserving boundary of a higher gauge topological model mixing 0-form \mathbb{Z}_{2} - and 1-form \mathbb{Z}_{3} -gauge symmetries.

Partially guided by the physical interpretation of the non-trivial 2-representation of the (1-)group \mathbb{Z}_2 as a string-like condensation defect in the (3+1)d toric code [61, 64], we proposed an interpretation of the simple 2representations of the 2-group $\mathbb{G}(2,3)$ as composites of condensation defects and violations of 1-form gauge invariance in the higher gauge model. Although we focused on $\mathbb{G}(2,3)$ for concreteness, we could have dealt with the dihedral 2-group $\mathbb{G}(2, n)$ for an arbitrary n with no additional effort, and with the more general case discussed in sec. III almost as easily. We expect the case of an arbitrary 2-group to be treatable invoking similar techniques, however the interpretation we offered of 2-representations and 1-intertwiners in terms of module categories and module category functors would have to be adapted. That being said, we expect the interplay between condensation defects—gases of 0-charges that spontaneously break the effective 0-form symmetry along one-dimensional submanifolds-and 1-charges to persist in the most general case.

Acknowledgements: The author is grateful to Joe Huxford for answering questions about manuscript [77], as well as Alex Bullivant and Apoorv Tiwari for numerous discussions on closely related topics.

Note added: After completion of this work, the author was made aware of the manuscripts [83] and [84] that might contain partially overlapping results.

References

- D. Gaiotto, A. Kapustin, N. Seiberg, and B. Willett, Generalized Global Symmetries, JHEP 02, 172, arXiv:1412.5148 [hep-th].
- [2] B. Eckman and P. J. Hilton, Structure maps in group theory, Fundamenta Mathematicae 50, 207 (1961).
- [3] A. Kapustin and N. Seiberg, Coupling a QFT to a TQFT and Duality, JHEP 04, 001, arXiv:1401.0740 [hep-th].
- [4] E. Sharpe, Notes on generalized global symmetries in QFT, Fortsch. Phys. 63, 659 (2015), arXiv:1508.04770 [hep-th].
- [5] D. Gaiotto, A. Kapustin, Z. Komargodski, and N. Seiberg, Theta, Time Reversal, and Temperature, JHEP 05, 091, arXiv:1703.00501 [hep-th].
- [6] D. Gaiotto, Z. Komargodski, and N. Seiberg, Timereversal breaking in QCD₄, walls, and dualities in 2 + 1 dimensions, JHEP **01**, 110, arXiv:1708.06806 [hep-th].
- [7] L. Bhardwaj and Y. Tachikawa, On finite symmetries and their gauging in two dimensions, JHEP 03, 189, arXiv:1704.02330 [hep-th].
- [8] Y. Tachikawa, On gauging finite subgroups, SciPost Phys. 8, 015 (2020), arXiv:1712.09542 [hep-th].

- [9] D. M. Hofman and N. Iqbal, Goldstone modes and photonization for higher form symmetries, SciPost Phys. 6, 006 (2019), arXiv:1802.09512 [hep-th].
- [10] P.-S. Hsin, H. T. Lam, and N. Seiberg, Comments on One-Form Global Symmetries and Their Gauging in 3d and 4d, SciPost Phys. 6, 039 (2019), arXiv:1812.04716 [hep-th].
- [11] C. Delcamp and A. Tiwari, On 2-form gauge models of topological phases, JHEP 05, 064, arXiv:1901.02249 [hep-th].
- [12] N. Sogabe and N. Yamamoto, Triangle Anomalies and Nonrelativistic Nambu-Goldstone Modes of Generalized Global Symmetries, Phys. Rev. D 99, 125003 (2019), arXiv:1903.02846 [hep-th].
- [13] N. Seiberg and S.-H. Shao, Exotic Symmetries, Duality, and Fractons in 2+1-Dimensional Quantum Field Theory, SciPost Phys. 10, 027 (2021), arXiv:2003.10466 [cond-mat.str-el].
- [14] N. Seiberg and S.-H. Shao, Exotic \mathbb{Z}_N symmetries, duality, and fractons in 3+1-dimensional quantum field theory, SciPost Phys. 10, 003 (2021), arXiv:2004.06115 [cond-mat.str-el].
- [15] N. Iqbal and J. McGreevy, Mean string field theory: Landau-Ginzburg theory for 1-form symmetries (2021), arXiv:2106.12610 [hep-th].
- [16] Z. Nussinov and G. Ortiz, Sufficient symmetry conditions for Topological Quantum Order, Proc. Nat. Acad. Sci. 106, 16944 (2009), arXiv:cond-mat/0605316.
- [17] Z. Nussinov and G. Ortiz, A symmetry principle for topological quantum order, Annals Phys. **324**, 977 (2009), arXiv:cond-mat/0702377.
- [18] X.-G. Wen, Emergent anomalous higher symmetries from topological order and from dynamical electromagnetic field in condensed matter systems, Phys. Rev. B 99, 205139 (2019), arXiv:1812.02517 [cond-mat.str-el].
- [19] E. Lake, Higher-form symmetries and spontaneous symmetry breaking (2018), arXiv:1802.07747 [hep-th].
- [20] J. Zhao, Z. Yan, M. Cheng, and Z. Y. Meng, Higher-form symmetry breaking at Ising transitions, Phys. Rev. Res. 3, 033024 (2021), arXiv:2011.12543 [cond-mat.str-el].
- [21] N. Iqbal and J. McGreevy, Toward a 3d Ising model with a weakly-coupled string theory dual, SciPost Phys. 9, 019 (2020), arXiv:2003.04349 [hep-th].
- [22] B. C. Rayhaun and D. J. Williamson, Higher-Form Subsystem Symmetry Breaking: Subdimensional Criticality and Fracton Phase Transitions (2021), arXiv:2112.12735 [cond-mat.str-el].
- [23] Z. Bi, E. Lake, and T. Senthil, Landau ordering phase transitions beyond the Landau paradigm, Phys. Rev. Res. 2, 023031 (2020), arXiv:1910.12856 [cond-mat.strel].
- [24] J. McGreevy, Generalized Symmetries in Condensed Matter (2022), arXiv:2204.03045 [cond-mat.str-el].
- [25] P. Etingof, D. Nikshych, and V. Ostrik, On fusion categories (2002), arXiv:math/0203060 [math.QA].
- [26] A. Feiguin, S. Trebst, A. W. W. Ludwig, M. Troyer, A. Kitaev, Z. Wang, and M. H. Freedman, Interacting anyons in topological quantum liquids: The golden chain, Phys. Rev. Lett. 98, 160409 (2007).
- [27] S. Trebst, E. Ardonne, A. Feiguin, D. A. Huse, A. W. W. Ludwig, and M. Troyer, Collective states of interacting fibonacci anyons, Phys. Rev. Lett. **101**, 050401 (2008).
- [28] C. Gils, E. Ardonne, S. Trebst, A. W. W. Ludwig, M. Troyer, and Z. Wang, Collective states of interact-

- [29] A. Kapustin and M. Tikhonov, Abelian duality, walls and boundary conditions in diverse dimensions, JHEP 11, 006, arXiv:0904.0840 [hep-th].
- [30] A. Kapustin and N. Saulina, Surface operators in 3d Topological Field Theory and 2d Rational Conformal Field Theory (2010), arXiv:1012.0911 [hep-th].
- [31] E. Ardonne, J. Gukelberger, A. W. W. Ludwig, S. Trebst, and M. Troyer, Microscopic models of interacting yang-lee anyons, New Journal of Physics 13, 045006 (2011).
- [32] C. Gils, E. Ardonne, S. Trebst, D. A. Huse, A. W. W. Ludwig, M. Troyer, and Z. Wang, Anyonic quantum spin chains: Spin-1 generalizations and topological stability, Phys. Rev. B 87, 235120 (2013).
- [33] M. Buican and A. Gromov, Anyonic Chains, Topological Defects, and Conformal Field Theory, Commun. Math. Phys. 356, 1017 (2017), arXiv:1701.02800 [hep-th].
- [34] R. Thorngren and Y. Wang, Fusion Category Symmetry I: Anomaly In-Flow and Gapped Phases (2019), arXiv:1912.02817 [hep-th].
- [35] T.-C. Huang and Y.-H. Lin, Topological Field Theory with Haagerup Symmetry (2021), arXiv:2102.05664 [hepth].
- [36] R. Thorngren and Y. Wang, Fusion Category Symmetry II: Categoriosities at c = 1 and Beyond (2021), arXiv:2106.12577 [hep-th].
- [37] J. Kaidi, K. Ohmori, and Y. Zheng, Kramers-Wannierlike duality defects in (3+1)d gauge theories (2021), arXiv:2111.01141 [hep-th].
- [38] Y. Choi, C. Cordova, P.-S. Hsin, H. T. Lam, and S.-H. Shao, Non-Invertible Duality Defects in 3+1 Dimensions (2021), arXiv:2111.01139 [hep-th].
- [39] N. Tantivasadakarn, R. Thorngren, A. Vishwanath, and R. Verresen, Long-range entanglement from measuring symmetry-protected topological phases (2021), arXiv:2112.01519 [cond-mat.str-el].
- [40] F. J. Wegner, Duality in Generalized Ising Models and Phase Transitions without Local Order Parameters, Journal of Mathematical Physics 12, 2259 (1971), https://doi.org/10.1063/1.1665530.
- [41] R. B. Laughlin, Anomalous quantum hall effect: An incompressible quantum fluid with fractionally charged excitations, Phys. Rev. Lett. 50, 1395 (1983).
- [42] X. G. Wen, Topological Order in Rigid States, Int. J. Mod. Phys. B 4, 239 (1990).
- [43] X. Chen, Z.-C. Gu, and X.-G. Wen, Local unitary transformation, long-range quantum entanglement, wave function renormalization, and topological order, Phys. Rev. B 82, 155138 (2010).
- [44] Hoàng Xuân Sính, Gr-catégories, Ph.D. thesis (1973).
- [45] A. Kapustin and R. Thorngren, Higher symmetry and gapped phases of gauge theories (2013), arXiv:1309.4721 [hep-th].
- [46] C. Delcamp and A. Tiwari, From gauge to higher gauge models of topological phases, JHEP 10, 049, arXiv:1802.10104 [cond-mat.str-el].
- [47] C. Córdova, T. T. Dumitrescu, and K. Intriligator, Exploring 2-Group Global Symmetries, JHEP 02, 184, arXiv:1802.04790 [hep-th].
- [48] F. Benini, C. Córdova, and P.-S. Hsin, On 2-Group Global Symmetries and their Anomalies, JHEP 03, 118, arXiv:1803.09336 [hep-th].

- [49] P.-S. Hsin and A. Turzillo, Symmetry-enriched quantum spin liquids in (3 + 1)d, JHEP 09, 022, arXiv:1904.11550 [cond-mat.str-el].
- [50] N. Iqbal and N. Poovuttikul, 2-group global symmetries, hydrodynamics and holography (2020), arXiv:2010.00320 [hep-th].
- [51] L. Crane and D. N. Yetter, Measurable categories and 2-groups (2003), arXiv:math/0305176.
- [52] J. W. Barrett and M. Mackaay, Categorical representations of categorical groups (2004), arXiv:math/0407463.
- [53] J. Elgueta, Representation theory of 2-groups on finite dimensional 2-vector spaces (2004), arXiv:math/0408120.
- [54] V. Ostrik, Module categories over the Drinfeld double of a finite group, arXiv Mathematics e-prints, math/0202130 (2002), arXiv:math/0202130 [math.QA].
- [55] N. Ganter and M. Kapranov, Representation and character theory in 2-categories, arXiv Mathematics e-prints , math/0602510 (2006), arXiv:math/0602510 [math.KT].
- [56] B. Bartlett, On unitary 2-representations of finite groups and topological quantum field theory (2009), arXiv:0901.3975 [math.QA].
- [57] W. Ji and X.-G. Wen, Categorical symmetry and noninvertible anomaly in symmetry-breaking and topological phase transitions, Phys. Rev. Research 2, 033417 (2020).
- [58] L. Kong, T. Lan, X.-G. Wen, Z.-H. Zhang, and H. Zheng, Algebraic higher symmetry and categorical symmetry: A holographic and entanglement view of symmetry, Phys. Rev. Research 2, 043086 (2020).
- [59] C. Delcamp, Tensor network approach to electromagnetic duality in (3+1)d topological gauge models (2021), arXiv:2112.08324 [cond-mat.str-el].
- [60] L. Kong and X.-G. Wen, Braided fusion categories, gravitational anomalies, and the mathematical framework for topological orders in any dimensions (2014), arXiv:1405.5858 [cond-mat.str-el].
- [61] D. V. Else and C. Nayak, Cheshire charge in (3+1)dimensional topological phases, Phys. Rev. B 96, 045136 (2017).
- [62] D. Gaiotto and T. Johnson-Freyd, Condensations in higher categories (2019), arXiv:1905.09566 [math.CT].
- [63] L. Kong, Y. Tian, and S. Zhou, The center of monoidal 2categories in 3+1D Dijkgraaf-Witten theory, Adv. Math. 360, 106928 (2020), arXiv:1905.04644 [math.QA].
- [64] L. Kong, Y. Tian, and Z.-H. Zhang, Defects in the 3dimensional toric code model form a braided fusion 2category, JHEP 12, 078, arXiv:2009.06564 [cond-mat.strel].
- [65] T. Johnson-Freyd, (3+1)D topological orders with only a Z₂-charged particle (2020), arXiv:2011.11165 [math.QA].
- [66] K. Roumpedakis, S. Seifnashri, and S.-H. Shao, Higher Gauging and Non-invertible Condensation Defects (2022), arXiv:2204.02407 [hep-th].
- [67] J. C. Baez and A. D. Lauda, Higher-Dimensional Algebra V: 2-Groups, arXiv Mathematics e-prints, math/0307200 (2003), arXiv:math/0307200 [math.QA].
- [68] J. C. Baez and U. Schreiber, Higher Gauge Theory, arXiv Mathematics e-prints , math/0511710 (2005), arXiv:math/0511710 [math.DG].
- [69] J. C. Baez and J. Huerta, An Invitation to Higher Gauge Theory, Gen. Rel. Grav. 43, 2335 (2011), arXiv:1003.4485 [hep-th].
- [70] D. N. Yetter, TQFT's from homotopy 2 types, J. Knot Theor. Ramifications 2, 113 (1993).
- [71] A. Kitaev, Fault-tolerant quantum computation by

anyons, Annals of Physics 303, 2 (2003).

- [72] A. Hamma, P. Zanardi, and X.-G. Wen, String and membrane condensation on three-dimensional lattices, Phys. Rev. B 72, 035307 (2005).
- [73] J. H. C. Whitehead, Combinatorial homotopy. I, Bulletin of the American Mathematical Society 55, 213 (1949).
- [74] A. Bullivant, M. Calçada, Z. Kádár, P. Martin, and J. a. F. Martins, Topological phases from higher gauge symmetry in 3 + 1 dimensions, Phys. Rev. B 95, 155118 (2017).
- [75] A. Bullivant and C. Delcamp, Excitations in strict 2group higher gauge models of topological phases, JHEP 01, 107, arXiv:1909.07937 [cond-mat.str-el].
- [76] J. Huxford and S. H. Simon, Excitations in the Higher Lattice Gauge Theory Model for Topological Phases III: the 3+1d Case (2022), arXiv:2206.09941 [cond-mat.strel].
- [77] J. Huxford and S. H. Simon, Excitations in the Higher Lattice Gauge Theory Model for Topological Phases I: Overview (2022), arXiv:2202.08294 [cond-mat.str-el].
- [78] J. Huxford and S. H. Simon, Excitations in the Higher

Lattice Gauge Theory Model for Topological Phases II: the 2+1d Case (2022), arXiv:2204.05341 [cond-mat.str-el].

- [79] A. Bullivant and C. Delcamp, Tube algebras, excitations statistics and compactification in gauge models of topological phases, JHEP 10, 216, arXiv:1905.08673 [condmat.str-el].
- [80] P. Etingof, S. Gelaki, D. Nikshych, and V. Ostrik, *Tensor categories*, Vol. 205 (American Mathematical Soc., 2016).
- [81] C. L. Douglas and D. J. Reutter, Fusion 2-categories and a state-sum invariant for 4-manifolds (2018), arXiv:1812.11933 [math.QA].
- [82] J. Greenough, Monoidal 2-structure of bimodule categories, Journal of Algebra 324, 1818 (2010).
- [83] T. Bartsch, M. Bullimore, A. E. V. Ferrari, and J. Pearson, Non-invertible Symmetries and Higher Representation Theory I (2022), arXiv:2208.05993 [hep-th].
- [84] L. Bhardwaj, S. Schafer-Nameki, and J. Wu, Universal Non-Invertible Symmetries (2022), arXiv:2208.05973 [hep-th].