

Exclusive doubly charged Higgs boson pair production in pp collisions at the LHC

Laura DUARTE,^{1,*} Victor P. GONÇALVES,^{2,3,4,†} Daniel E. MARTINS,^{4,‡} Têssio B. de MELO,^{4,§} and Farinaldo S. QUEIROZ^{1,5,6,¶}

¹*International Institute of Physics, Universidade Federal do Rio Grande do Norte, Campus Universitario, Lagoa Nova, Natal-RN 59078-970, Brazil*

²*Institut für Theoretische Physik, Westfälische Wilhelms-Universität Münster, Wilhelm-Klemm-Straße 9, D-48149 Münster, Germany*

³*Institute of Modern Physics, Chinese Academy of Sciences, Lanzhou 730000, China*

⁴*Institute of Physics and Mathematics, Federal University of Pelotas, Postal Code 354, 96010-900, Pelotas, RS, Brazil*

⁵*Departamento de Física, Universidade Federal do Rio Grande do Norte, 59078-970, Natal, RN, Brazil*

⁶*Millennium Institute for SubAtomic Physics at the High-energy frontIeR, SAPHIR, Chile*

The type II seesaw model predicts the existence of a doubly charged Higgs boson ($H^{\pm\pm}$), which can be produced through the Drell - Yan process and photon – induced interactions at hadronic colliders. In recent years, such reactions have been largely investigated in inclusive processes, where both incident protons breakup and a large number of particles is produced in addition to the doubly charged Higgs pair. In this paper, we investigate, for the first time, the possibility of searching for $H^{\pm\pm}$ in exclusive processes, which are characterized by two rapidity gaps and two intact very forward protons in final state. We estimate the associated cross section making use of the exclusive nature of the final state, together with the use of timing information provided by forward proton detectors. Moreover, the background is estimated considering distinct amounts of pile-up. Our results indicate that the exclusive doubly charged Higgs pair production is a promising way to search for signatures of the type II seesaw and to obtain lower mass bounds on $H^{\pm\pm}$.

PACS numbers:

Keywords: Doubly charged Higgs, exclusive processes, proton-proton collisions

*Electronic address: l.duarte@unesp.br

†Electronic address: barros@ufpel.edu.br

‡Electronic address: dan.ernani@gmail.com

§Electronic address: tessiomelo@gmail.com

¶Electronic address: farinaldo.queiroz@ufrn.br

I. INTRODUCTION

Many beyond Standard Model (SM) scenarios predict the existence of a scalar particle carrying two units of electric charge, especially those related to neutrino masses. One of the most popular scenarios featuring this kind of particle is the type II seesaw mechanism [1–4], in which a $SU(2)_L$ scalar triplet $\Delta = (H^{++}, H^+, H^0)$ accounts for the tiny neutrino masses after its neutral component develop a vacuum expectation value (VEV). Accompanying the neutral scalar inside the triplet, there is the doubly charged scalar H^{++} (along with a singly charged scalar as well). Doubly charged scalars are also present in left-right symmetric models [5–7] (which were, in fact, the first models to propose the existence of these particles), coming from either a $SU(2)_L$ or a $SU(2)_R$ triplet. They can also arise from other scalar multiplet representations in models such as radiative seesaw models [8, 9], 3-3-1 models [12–15], Georgi-Machacek [16–18] and little Higgs models [19–21].

Several searches for doubly charged scalar particles have been performed recently, specially in high-energy colliders [22–29]. From the non-observation of any excess compared to the SM background, limits on the production cross section of the H^{++} have been placed, which can be translated into bounds on the H^{++} mass, depending on the assumptions about its decays modes. Since in many scenarios neutrino masses are generated via the coupling between the SM lepton doublet and the scalar multiplet that hosts the H^{++} , a reasonable assumption is that H^{++} is able to decay into SM leptons. In particular, the decay into same-sign dileptons provides a smoking gun signature for a doubly charged particle and, for this reason, this signature has been used in most of the experimental analysis. The most stringent limits to date come from the resonant searches by the ATLAS Collaboration obtained with 36.1 fb^{-1} data collected at 13 TeV [29]. Assuming that H^{++} decays 100% into $e^\pm e^\pm$, the lower limit on its mass is around 770 GeV, at 95% C.L.

The current lower limit on the mass of the doubly charged Higgs in hadronic colliders have been mainly obtained by considering that the $H^{++}H^{--}$ pair can be produced via the Drell - Yan (DY) process as well as by photon - photon and gluon - gluon interactions. As demonstrated e.g. in Ref. [30], photon - induced processes are suppressed by about an order of magnitude with respect to the DY production, and the gluon fusion is subdominant compared to the $\gamma\gamma$ fusion. The DY production is an inclusive process, characterized by the fragmentation of both incident protons and a large number of particles is produced in addition to the doubly charged Higgs pair. On the other hand, in $\gamma\gamma$ interactions, the processes can be classified as elastic, semi - elastic or inelastic, depending if both, only one or none of the incident protons remain intact (For previous studies see, e.g. Refs. [31, 32]). As a consequence, the elastic events, represented in Fig. 1, although subdominant with respect to the other channels, has a final state characterized by the presence of two regions devoid of hadronic activity, called rapidity gaps, separating the intact very forward protons from the central massive




FIG. 1: Diagram for the exclusive doubly charged Higgs pair production in pp collisions at the LHC.

object. Such very clean final state is a characteristic of exclusive processes, which are generated by the exchange of color singlet objects (e.g. a photons or a Pomeron) [33]. Such distinct topology of exclusive processes was not explored in previous studies and will be considered, for the first time, in the present paper. In our analysis we will consider that these events can be collected using the forward proton detectors (FPD) such as the ATLAS Forward Proton detector (AFP) [34, 35] and Precision Proton Spectrometer (CT-PPS) [36] that are installed symmetrically around the interaction point at a distance of roughly 210 m from the interaction point. In particular, we will assume that $m(H^{\pm\pm}) \geq 350$ GeV and that the doubly charged Higgs decays into same - signal dileptons. The large invariant mass of the produced system implies that the intact protons in the final state can be tagged by FPDs. Consequently, such events can, in principle, be separated and be used to searching for the doubly charged Higgs and/or impose limits on its mass. In our analysis we will also consider the presence of pile-up events, which is a reality in current and future LHC runs, and arise due to the presence of extra pp interactions per bunch crossing in high luminosity pp collisions at the LHC. These extra interactions generates additional tracks increase the background stemming from the inelastic doubly charged Higgs pair produced in a different primary vertex and, in general, destroy the signature associated to two rapidity gaps. Following Refs. [37, 38], such background will be suppressed by considering the time-of-flight (ToF) detectors [39–42]. As we will demonstrate below, the exclusive $H^{++}H^{--}$ production is a promising alternative to searching for signals of the type II seesaw mechanism (For related searches of BSM physics in exclusive processes see, e.g., Refs. [39, 43–51]).

This paper is organized as follows. In the next section we present a brief review of the formalism for the doubly charged Higgs interactions and for the treatment of the $H^{\pm\pm}$ pair production in photon-induced interactions in pp collisions. In Section III we discuss the backgrounds considered in our

analysis as well as the exclusivity cuts. In addition, we present our predictions for the invariant mass, transverse momentum and acoplanarity distributions as well as for the total cross sections for the $H^{\pm\pm}$ pair production in $\gamma\gamma$ interactions. Finally, in Section IV we summarize our main conclusions.

II. FORMALISM

In this section we describe the relevant H^{++} interactions for its production and decay at the LHC. Since doubly charged scalar bosons appear in several extensions of the SM [8, 16, 52, 53], here we will first focus on the case of the Higgs Triplet Model, which is a minimal realization of the type II seesaw mechanism, and later extend to more general H^{++} particles, keeping some of the key features from this prototypical case.

In the Higgs Triplet Model, the SM scalar sector is extended by adding a weak scalar triplet Δ ,

$$\Delta(1, 2, 1) = \begin{pmatrix} H^+/\sqrt{2} & H^{++} \\ H^0 & -H^+/\sqrt{2} \end{pmatrix}, \quad (1)$$

where the numbers in parentheses denote the representation under the $SU(3)_C \times SU(2)_L \times U(1)_Y$ gauge group of the SM ¹. The triplet Δ interacts with the SM leptons via the coupling,

$$\mathcal{L}_Y \supset -y\bar{L}^c i\sigma^2 \Delta L + h.c. \quad (2)$$

where y is the matrix of Yukawa coupling constants, $L = (\nu_L \ e_L)^T$ is the SM lepton doublet, σ^2 is the second Pauli matrix and the superscript c denotes charge conjugation. By means of this interaction, neutrino masses are generated after H^0 develops a VEV v_Δ . As the masses will be proportional to this VEV, it is clear that v_Δ is required to be small in order to account for the tiny neutrino masses.

The H^{++} in the Higgs Triplet Model further couples to gauge bosons and other scalars at tree level. The coupling to quarks, however, is forbidden (note that the $Y = 1$ hypercharge that allows the coupling of Δ with the leptons in Eq. (2), automatically forbids a coupling to the quark doublets). Thus, the H^{++} has basically two decays modes: $H^{\pm\pm} \rightarrow \ell_i^\pm \ell_j^\pm$ and $H^{\pm\pm} \rightarrow W^\pm W^\pm$. The widths for these two body decays are given by [54]:

$$\Gamma(H^{\pm\pm} \rightarrow \ell_i^\pm \ell_j^\pm) = \frac{m_{H^{\pm\pm}}}{8\pi(1 + \delta_{ij})} \frac{|\mathcal{M}_\nu^{ij}|^2}{v_\Delta^2}, \quad (3)$$

$$\Gamma(H^{\pm\pm} \rightarrow W^\pm W^\pm) = \frac{g^4 v_\Delta^2}{8\pi m_{H^{\pm\pm}}} \sqrt{1 - \frac{4m_W^2}{m_{H^{\pm\pm}}^2}} \left[2 + \left(\frac{m_{H^{\pm\pm}}^2}{2m_W^2} - 1 \right)^2 \right], \quad (4)$$

¹ We are using the convention $Q = T_3 + Y$ for the electric charge and hypercharge.

where δ_{ij} is the Kronecker delta, \mathcal{M}_{ν}^{ij} are the elements of the neutrino mass matrix and $\ell_i^\pm = e^\pm, \mu^\pm, \tau^\pm$. From these expressions, it is clear that the branching ratios for H^{++} decays are very sensitive to the triplet VEV v_Δ . Most of the searches so far have been focused in the same sign dilepton decays $H^{\pm\pm} \rightarrow \ell_i^\pm \ell_j^\pm$, which provides a clean distinctive signature. This amounts to assume that $v_\Delta \leq 10^{-4}$ GeV, so that this decay is dominant over the $H^{\pm\pm} \rightarrow W^\pm W^\pm$ channel. Although this channel is more difficult to analyze due to the presence of neutrinos in the final state, and the consequent larger SM backgrounds, searches in this channel have been done [28, 29]. It should be mentioned that, if kinematically allowed, H^{++} can also have cascade decays to scalar and vector bosons, such as $H^{\pm\pm} \rightarrow H^\pm W^\pm \rightarrow H^0 W^\pm W^\pm$. However, these decays are much more challenging to probe experimentally.

Doubly-charged Higgs bosons can be pair produced at the LHC via the Drell-Yan (DY) process $q\bar{q} \rightarrow \gamma^*/Z^* \rightarrow H^{\pm\pm} H^{\mp\mp}$. The trilinear and quartic gauge interactions relevant for the calculation of the pair production come from the kinetic Lagrangian for H^{++} , and can be written as,

$$\begin{aligned} \mathcal{L}_{\text{kin}} = & i \left[2eA_\mu + \frac{g}{c_W} (2 - Y - 2s_W^2) Z_\mu \right] H^{++} \partial^\mu H^{--} \\ & + \left[2eA_\mu + \frac{g}{c_W} (2 - Y - 2s_W^2) Z_\mu \right] \left[2eA^\mu + \frac{g}{c_W} (2 - Y - 2s_W^2) Z^\mu \right] H^{++} H^{--}, \end{aligned} \quad (5)$$

where $c_W = \cos \theta_W$, $s_W = \sin \theta_W$, with θ_W being the weak mixing angle. The H^{++} can also be produced in association with a W boson. However, in the Higgs Triplet Model this production rate is highly suppressed, since it is proportional to the factor (v_Δ^2/m_W^2) . Production of H^{++} in association with H^- can occur at significant rates via the process $u\bar{d} \rightarrow W^{+*} \rightarrow H^{++} H^-$. Nevertheless, the H^- decays into final states involving neutrinos, as a result its signatures are not very clean. For these reasons, in their searches for doubly charged Higgs bosons, the ATLAS and CMS collaborations have considered only the pair production via DY process. In the next section we shall discuss a new way to search for the H^{++} , employing elastic photon fusion production, combined with the tagging of the intact protons after the collision, which is complementary to the ATLAS and CMS searches and leads to excellent detection prospects.

In our analysis, we also consider doubly charged scalars arising from more general $SU(2)_L$ representations. We assume that most of the features of the H^{++} from the Higgs triplet model remain valid for these other cases, in particular concerning its production and decay modes. More specifically, our analysis includes:

- H^{++} from a $SU(2)_L$ singlet: $\phi = H^{++}$; ($T_3 = 0, Y = 2$);
- H^{++} from a $SU(2)_L$ doublet: $\phi = (H^{++} H^+)$; ($T_3 = 1/2, Y = 3/2$);
- H^{++} from a $SU(2)_L$ triplet: $\phi = (H^{++} H^+ H^0)$; ($T_3 = 1, Y = 1$).

These representations arise quite commonly in a variety of models. For instance, in some versions of the 3-3-1 model [14, 15], an scalar triplet emerges from an original $SU(3)_L$ scalar sextet, after this symmetry is spontaneously broken down to the SM gauge group. In the left-right symmetric model [10, 11], besides the H_L^{++} from the $SU(2)_L$ triplet Δ_L , there is also a companion H_R^{++} from the $SU(2)_R$ triplet Δ_R (which is however singlet under $SU(2)_L$). In radiative neutrino mass models [8, 9], the doubly charged Higgs boson is analogous to the singlet H_R^{++} of the left-right model. As the presence of singlets, doublets or triplets containing a doubly charged boson is very common in several neutrino mass studies [55–60], we will stick to these representations. It should be noted that other scalar multiplets, such as $SU(2)_L$ quadruplets or quintuplets, can also host a H^{++} . In these cases, however, the coupling of H^{++} to the leptons would require the presence of new vector-like fermions that mix with the SM leptons. For this reason, we will not consider these higher isospin representations here. It is important to emphasize that the gauge interactions for the DY pair production given in Eq. (5) remain valid for all the aforementioned representations (provided the appropriate hypercharge value is used), while the production via photon-photon fusion, discussed below, is insensitive to the multiplet representation the H^{++} originates from. In addition, we will assume that, similarly to the case of the Higgs Triplet Model in the regime of small v_Δ , the H^{++} decays dominantly into same-sign dileptons, also for the singlet and doublet cases.

A. Production via photon-photon fusion

As discussed in the Introduction, our focus in this paper is on the analysis of the doubly charged Higgs production by elastic $\gamma\gamma$ interactions, as represented in Fig. 1. The total cross section for these interactions can be factorized in terms of the equivalent flux of photons into the hadron projectiles and the photon-photon cross section. In particular, the $H^{++}H^{--}$ production in photon-photon interactions is described by

$$\sigma(h_1 h_2 \rightarrow h_1 \otimes H^{++} H^{--} \otimes h_2) = S_{abs}^2 \int dx_1 \int dx_2 \gamma_1^{el}(x_1) \cdot \gamma_2^{el}(x_2) \cdot \hat{\sigma}(\gamma\gamma \rightarrow H^{++} H^{--}), \quad (6)$$

where \otimes represents the presence of a rapidity gap in the final state, S_{abs}^2 is the absorptive factor (see below), x is the fraction of the hadron energy carried by the photon and $\gamma^{el}(x)$ is the elastic equivalent photon distribution of the hadron. The general expression for the elastic photon flux of the proton is given by [61]

$$\gamma^{el}(x) = -\frac{\alpha}{2\pi} \int_{-\infty}^{-\frac{m^2 x^2}{1-x}} \frac{dt}{t} \left\{ \left[2 \left(\frac{1}{x} - 1 \right) + \frac{2m^2 x}{t} \right] H_1(t) + x G_M^2(t) \right\}, \quad (7)$$

where $t = q^2$ is the momentum transfer squared of the photon,

$$H_1(t) \equiv \frac{G_E^2(t) + \tau G_M^2(t)}{1 + \tau} \quad (8)$$

with $\tau \equiv -t/m^2$, m being the nucleon mass, and where G_E and G_M are the Sachs elastic form factors. In our analysis, we will use the photon flux derived in Ref. [62], where an analytical expression is presented. Moreover, we will also assume that the absorptive factor S_{abs}^2 , which takes into account of additional soft interactions between incident protons which leads to an extra production of particles that destroy the rapidity gaps in the final state [63], is equal to the unity. Such assumption is a reasonable approximation since the contribution of the soft interactions is expected to be small in $\gamma\gamma$ interactions due to the long range of the electromagnetic interaction. In contrast, for exclusive processes mediated by a Pomeron, such factor can be $\mathcal{O}(10^{-2})$ and its precise value is a theme of intense debate (for reviews see Refs. [64–66]).

A comment is in order. In recent years, the determination of photon PDF in a global analysis was performed by different groups assuming distinct assumptions for, e.g., the initial conditions and the treatment of the higher order corrections [67–69]. In general, in these analyzes the authors have considered that the photon PDF is given by the elastic γ^{el} and the inelastic γ^{inel} contributions (For a more detailed discussion see, e.g. Refs. [70, 71]). The elastic photon distribution is associated to the probability that a proton emits a photon and remains intact and can be expressed in terms of the electric and magnetic form factors using the Equivalent Photon Approximation (EPA) method [61, 62], as in Eq. (7) above. On the other hand, the inelastic photon distribution γ^{inel} provides the probability for a photon emission from a proton in an inelastic interaction and can be estimated assuming that the photon is a constituent of the proton, along with quarks and gluons, with its contribution being derived by solving the DGLAP evolution equations modified by the inclusion of the QED parton splittings. The resulting predictions for γ^{inel} , and consequently for the photon PDF, are strongly dependent on the assumptions assumed in the global analysis, implying a large uncertainty on the predictions for the inclusive processes derived e.g. in Ref. [30]. In contrast, for exclusive (elastic) processes, the uncertainty on the predictions is strongly reduced since it depends only on γ^{el} , which is well - known.

III. RESULTS

In what follows we will present our results for the doubly charged Higgs production in exclusive processes considering pp collisions at $\sqrt{s} = 13$ TeV. In our analysis, we will take into account of the current detector acceptances and resolutions as well as the pile-up effects. We will assume that the $H^{++}H^{--}$ system decays leptonically, $H^{++}H^{--} \rightarrow e^+e^+e^-e^-$, both forward protons are tagged and pile-up is present. In particular, we will consider two different amounts for the pile-up, $\langle\mu\rangle = 5$ and 50. Moreover, our results will be derived by considering additional exclusivity cuts and utilizing the time-of-flight (TOF) detectors for suppressing the combinatorial background coming from pile-up. The signal is assumed to be the $pp \rightarrow p \otimes H^{++}H^{--} \otimes p \rightarrow p \otimes (e^+e^+)(e^-e^-) \otimes p$ process, which




FIG. 2: Predictions for the invariant mass distribution of the four lepton system (left panel) and for the transverse momentum distribution of the e^+e^+ pair (right panel) at the generation level considering $m(H^{\pm\pm}) = 350$ GeV and pp collisions at $\sqrt{s} = 13$ TeV. DY* indicates that the double tagging and time - of - flight suppression factor have been applied to the inclusive DY predictions.

will be generated using the MadGraph 5 [72, 73]. For the background we will consider the photon - induced processes $pp \rightarrow p \otimes e^+e^-e^+e^- \otimes p$ and $pp \rightarrow p \otimes ZZ \otimes p \rightarrow p \otimes e^+e^-e^+e^- \otimes p$, where the last process occurs via the box diagram. In addition, the inclusive $H^{++}H^{--}$ production via the DY production is also estimated, considering the probability of fake double tagging. All these backgrounds are also generated by MadGraph 5 [72]. As in Refs. [37, 38], each event is then properly mixed with such a number of pile-up interactions that corresponds to the studied luminosity scenario. Detector effects are incorporated and the pile-up mixing is done using Delphes 3.5 [74] with input cards with CMS detector specifications. In order to perform the separation of the signal from backgrounds, we first select the central system as in the inclusive processes and then we apply exclusivity criteria. We require both forward protons to be detected by Forward Proton Detectors (FPDs) and processed by Time-of-Flight (ToF) detectors. In our analysis one has assumed a TOF resolution of 10 ps.

Initially, we will focus on the $H^{\pm\pm}$ from the Higgs triplet model, in which $Y = 1$, and let's estimate the differential distributions at the generation level. In Fig. 2 (left panel) we present the associated predictions for the invariant mass distribution of the $e^+e^+e^-e^-$ system, denoted $m(4lep)$, considering $m(H^{\pm\pm}) = 350$ GeV. The signal and the main background contributions are shown. The results for the $pp \rightarrow p \otimes ZZ \otimes p \rightarrow p \otimes e^+e^-e^+e^- \otimes p$ process are not presented since they are 6 orders of magnitude smaller than the signal. The background associated to the inclusive DY contribution is evaluated by applying the the suppression factors related to the probability of fake double tagging and ToF in the forward detector. As expected, the four lepton production by the QED process $\gamma\gamma \rightarrow e^+e^-e^+e^-$ dominates at small invariant mass, below the production threshold of the doubly charged Higgs pair. On the other hand, for $m(4lep) > 2m(H^{\pm\pm})$, one has that this contribution is suppressed by two orders

p-p @ $\sqrt{s} = 13$ TeV	$\gamma\gamma \rightarrow H^{++}H^{--} \rightarrow (e^+e^+)(e^-e^-)$	DY[After FT and ToF]: $q\bar{q} \rightarrow H^{++}H^{--} \rightarrow (e^+e^+)(e^-e^-)$	$\gamma\gamma \rightarrow e^+e^-e^+e^-$	$\gamma\gamma \rightarrow ZZ \rightarrow (e^+e^-)(e^+e^-)$ [box]
$m(H^{\pm\pm}) = 350.0$ GeV, $\langle\mu\rangle = 5$				
Total Cross section (fb)	0.53	0.0023	0.09	$\lesssim 10^{-7}$
$p_T(e^+e^+, e^-e^-) > 50.0$ GeV	0.44	0.0023	$\lesssim 10^{-5}$	0.0
$ \eta < 2.5$	0.44	0.0023	$\lesssim 10^{-5}$	0.0
$m(H^{\pm\pm}) \pm 10$ GeV	0.34	0.0016	0.0	0.0
$m(H^{\pm\pm}) = 350.0$ GeV, $\langle\mu\rangle = 50$				
Total Cross section (fb)	0.53	0.29	0.09	$\lesssim 10^{-7}$
$p_T(e^+e^+, e^-e^-) > 50.0$ GeV	0.44	0.29	$\lesssim 10^{-5}$	0.0
$ \eta < 2.5$	0.43	0.29	$\lesssim 10^{-5}$	0.0
$m(H^{\pm\pm}) \pm 10$ GeV	0.32	0.20	0.0	0.0
$m(H^{\pm\pm}) = 850.0$ GeV, $\langle\mu\rangle = 5$				
Total Cross section (fb)	0.061	0.00013	0.09	$\lesssim 10^{-7}$
$p_T(e^+e^+, e^-e^-) > 50.0$ GeV	0.060	0.00013	$\lesssim 10^{-5}$	0.0
$ \eta < 2.5$	0.060	0.00012	$\lesssim 10^{-5}$	0.0
$m(H^{\pm\pm}) \pm 10$ GeV	0.046	0.00008	0.0	0.0
$m(H^{\pm\pm}) = 850.0$ GeV, $\langle\mu\rangle = 50$				
Total Cross section (fb)	0.061	0.016	0.09	$\lesssim 10^{-7}$
$p_T(e^+e^+, e^-e^-) > 50.0$ GeV	0.060	0.016	$\lesssim 10^{-5}$	0.0
$ \eta < 2.5$	0.060	0.016	$\lesssim 10^{-5}$	0.0
$m(H^{\pm\pm}) \pm 10$ GeV	0.046	0.010	0.0	0.0

TABLE I: Predictions for the total cross sections of the signal and background at the detector level considering two different values of the doubly charged Higgs mass and two scenarios for the pile-up.

of magnitude in comparison to the signal, but is still larger than the corrected DY background.

In Fig. 2 (right panel) we present our predictions for the transverse momentum distribution of the e^+e^+ system, denoted by $p_T(e^+e^+)$. One has that the four lepton production by the QED process $\gamma\gamma \rightarrow e^+e^-e^+e^-$ dominates at small $p_T(e^+e^+)$, but is strongly suppressed with the increasing of the transverse momentum. In particular, our results indicate that signal dominates for $p_T(e^+e^+) > 50$ GeV. Moreover, one has that in this kinematical range the signal is larger than the DY contribution by ≈ 2 orders of magnitude.

In Table I we present our predictions at the detector level, which have been derived assuming the following cuts: (a) $p_T(e^\pm e^\pm) > 50$ GeV; (b) that the leptons are in the rapidity range $|\eta| \leq 2.5$, and (c) that the invariant mass of the dilepton system $m(e^\pm e^\pm)$ is in the range $m(H^{\pm\pm}) \pm 10$ GeV. The results are presented for two values of the doubly charged Higgs and two scenarios for the pile-up. One has that the signal and corrected DY cross sections decrease for larger values of $m(H^{\pm\pm})$, with the inclusive DY production being the main background. For a small amount of the pile-up and $m(H^{\pm\pm}) = 350$ GeV, one has that signal is two orders of magnitude larger than the background, as expected from Fig. 2. Such conclusion is also valid for $m(H^{\pm\pm}) = 850$ GeV and $\langle\mu\rangle = 5$. In contrast, for a larger amount of pile-up, $\langle\mu\rangle = 50$, the difference between signal and background decreases and both cross




FIG. 3: Mass dependence of the statistical significance for the exclusive doubly charge Higgs production in pp collisions at $\sqrt{s} = 13$ TeV considering two scenarios for the pile-up configuration and integrated luminosity ($\langle \mu \rangle, L$). Results for the Higgs triplet model ($Y = 1$). The vertical black line indicates the upper value for $m(H^{\pm\pm})$ excluded by the analysis of inclusive processes.

sections become of the same order of magnitude, with the signal being still larger than the corrected DY prediction.

The previous results allows to estimate the statistical significance and the signal to background ratio, S/B . In order to estimate these quantities, we consider two luminosity scenarios in terms of $\langle \mu \rangle$ and \mathcal{L} , where $\langle \mu \rangle$ represents the average number of pile-up interactions per event and \mathcal{L} is the integrated luminosity. Moreover, we assume \mathcal{L} to be 10 and 300 fb^{-1} for $\langle \mu \rangle = 5$ and 50, respectively. In addition, the significance will be estimated using the formula based on Asimov data set [75], which is more reliable for $S \approx B$ and reduces to the S/\sqrt{B} ratio if $S \ll B$. The results are presented in Fig. 3 as a function of doubly charged Higgs mass. We restrict our study for the mass range $m(H^{\pm\pm}) \leq 1000$ GeV due to the current limitations of the forward detectors which are not able to measure final states with invariant mass larger than ≈ 2600 GeV. The results indicate that the significance is larger than 2σ (5σ) for $m(H^{\pm\pm}) \lesssim 850$ GeV and $\mathcal{L} = 10$ (300) fb^{-1} . Considering that values smaller than 780 GeV are already excluded by the experimental analysis of the doubly charged Higgs pair production in inclusive processes performed by the ATLAS Collaboration, one has that a future experimental analysis of the exclusive production at the HL - LHC will be able to improve the limits on $m(H^{\pm\pm})$ at the 5σ level. Moreover, our results indicate that for $m(H^{\pm\pm}) > 850$ GeV, the exclusive process is also an important alternative to search for the doubly charged Higgs and extend the current limits on its mass.




FIG. 4: Mass dependence of the statistical significance for the exclusive doubly charge Higgs production in pp collisions at $\sqrt{s} = 13$ TeV considering distinct $SU(2)_L$ representations. and different scenarios for the pile-up configuration and integrated luminosity ($\langle \mu \rangle, L$).

The previous results have been derived assuming that the $H^{\pm\pm}$ has the properties predicted by the Higgs triplet model, in which $Y = 1$. As discussed in the Section II, doubly charged scalars can also arise from more general $SU(2)_L$ representations. In our analysis, we also have performed a detailed analysis of singlet ($Y = 2$) and doublet ($Y = 3/2$) representations and estimated the associated significances. In Fig. 4 we present a comparison of significances derived for the distinct representations and different values of the integrated luminosity. One has that the singlet predictions are similar to those derived for the triplet case. In contrast, the significance associated to the doublet representation is larger in comparison to the singlet and triplet predictions. In particular, we predict values larger than 10σ for $m(H^{\pm\pm}) \lesssim 1300$ GeV. Such results are expected due to the larger suppression of the DY background in the Higgs doublet representation.

IV. SUMMARY

One of the challenge problems in Particle Physics today is the description of the neutrino masses and all neutrino oscillation data. One of the simplest ways to describe these data is provided by the type II seesaw model, which predicts the existence of seven physical Higgs bosons, two of them being doubly charged. Another feature of the type II seesaw model is that the Higgs triplet can be produced directly through gauge interactions with electroweak bosons. Such aspect has motivated the study of the doubly charged Higgs pair production through the Drell - Yan process, where the pair is produced via s - channel γ^*/Z exchange, or via the vector boson fusion processes $\gamma\gamma \rightarrow H^{++}H^{--}$ and $ZZ \rightarrow H^{++}H^{--}$. In recent years, several experimental studies in high - energy colliders have searched for doubly charged scalar particles and derived bounds on the $H^{\pm\pm}$ mass. Distinctly from

previous studies, which focused on inclusive reactions, in this paper we have proposed the study of exclusive processes as an alternative to searching for the doubly charged Higgs. As demonstrated in our analysis, although the associated cross sections are smaller than those predicted for the inclusive reactions, the very clean final state and possibility of tagging of the intact protons in the final states and the reduction of the impact of the pile-up using the time - of - flight detectors, makes feasible the study of the exclusive $H^{++}H^{--}$ production in pp collisions at the LHC.

Acknowledgments

V.P.G. and D.E.M. acknowledge useful discussions with M. Tasevsky and G. Gil da Silveira. This work was partially supported by INCT-FNA (Process No. 464898/2014-5) and Simons Foundation (Award Number:884966, AF). V.P.G. was partially supported by the CAS President's International Fellowship Initiative (Grant No. 2021VMA0019) and by CNPq, CAPES and FAPERGS. T.B.M. acknowledges CNPq for the financial support (grant no. 164968/2020-2). D.E.M. was partially supported by CNPq (grant no.164609/2020-2 and 152473/2022-0). We also thank FAPESP grant 2021/01089-1, ICTP-SAIFR FAPESP grant 2016/01343-7, CNPq grant 408295/2021-0, Serrapilheira Foundation grant Serra-1912-31613, and ANID-Programa Milenio-code *ICN2019_044*.

-
- [1] J. Schechter and J. W. F. Valle, Phys. Rev. D **22**, 2227 (1980).
 - [2] T. P. Cheng and L. F. Li, Phys. Rev. D **22**, 2860 (1980).
 - [3] A. Arhrib, R. Benbrik, M. Chabab, G. Moultaqa, M. C. Peyranere, L. Rahili and J. Ramadan, Phys. Rev. D **84**, 095005 (2011).
 - [4] S. Ashanujjaman and K. Ghosh, JHEP **03**, 195 (2022).
 - [5] J. C. Pati and A. Salam, Phys. Rev. D **10**, 275-289 (1974) [erratum: Phys. Rev. D **11**, 703-703 (1975)].
 - [6] R. N. Mohapatra and J. C. Pati, Phys. Rev. D **11**, 2558 (1975).
 - [7] G. Senjanovic and R. N. Mohapatra, Phys. Rev. D **12**, 1502 (1975).
 - [8] A. Zee, Nucl. Phys. B **264**, 99-110 (1986).
 - [9] K. S. Babu, Phys. Lett. B **203**, 132-136 (1988).
 - [10] D. Borah and A. Dasgupta, JHEP **01** (2017), 072
 - [11] P. S. Bhupal Dev and Y. Zhang, JHEP **10** (2018), 199
 - [12] J. E. Cieza Montalvo, N. V. Cortez, J. Sa Borges and M. D. Tonasse, Nucl. Phys. B **756**, 1-15 (2006) [erratum: Nucl. Phys. B **796**, 422-423 (2008)].
 - [13] A. Alves, E. Ramirez Barreto, A. G. Dias, C. A. de S.Pires, F. S. Queiroz and P. S. Rodrigues da Silva, Phys. Rev. D **84**, 115004 (2011).
 - [14] A. C. B. Machado, P. Pasquini and V. Pleitez, [arXiv:1810.02817 [hep-ph]].

- [15] M. M. Ferreira, T. B. de Melo, S. Kovalenko, P. R. D. Pinheiro and F. S. Queiroz, *Eur. Phys. J. C* **79**, no.11, 955 (2019).
- [16] H. Georgi and M. Machacek, *Nucl. Phys. B* **262**, 463-477 (1985).
- [17] M. S. Chanowitz and M. Golden, *Phys. Lett. B* **165**, 105-108 (1985).
- [18] C. W. Chiang, G. Cottin and O. Eberhardt, *Phys. Rev. D* **99**, no.1, 015001 (2019).
- [19] N. Arkani-Hamed, A. G. Cohen, E. Katz, A. E. Nelson, T. Gregoire and J. G. Wacker, *JHEP* **08**, 021 (2002).
- [20] N. Arkani-Hamed, A. G. Cohen, E. Katz and A. E. Nelson, *JHEP* **07**, 034 (2002).
- [21] A. Hektor, M. Kadastik, M. Muntel, M. Raidal and L. Rebane, *Nucl. Phys. B* **787**, 198-210 (2007).
- [22] G. Aad *et al.* [ATLAS], *Eur. Phys. J. C* **72**, 2244 (2012).
- [23] S. Chatrchyan *et al.* [CMS], *Eur. Phys. J. C* **72**, 2189 (2012).
- [24] G. Aad *et al.* [ATLAS], *JHEP* **03**, 041 (2015).
- [25] V. Khachatryan *et al.* [CMS], *Phys. Rev. Lett.* **114**, no.5, 051801 (2015).
- [26] M. Aaboud *et al.* [ATLAS], *Eur. Phys. J. C* **78**, no.3, 199 (2018).
- [27] A. M. Sirunyan *et al.* [CMS], *Phys. Rev. Lett.* **120**, no.8, 081801 (2018).
- [28] M. Aaboud *et al.* [ATLAS], *Eur. Phys. J. C* **79**, no.1, 58 (2019).
- [29] G. Aad *et al.* [ATLAS], *JHEP* **06**, 146 (2021).
- [30] B. Fuks, M. Nemevšek and R. Ruiz, *Phys. Rev. D* **101**, no.7, 075022 (2020).
- [31] T. Han, B. Mukhopadhyaya, Z. Si and K. Wang, *Phys. Rev. D* **76**, 075013 (2007).
- [32] K. S. Babu and S. Jana, *Phys. Rev. D* **95**, no.5, 055020 (2017).
- [33] A. Hebecker, *Phys. Rept.* **331**, 1 (2000); L. Schoeffel, *Prog. Part. Nucl. Phys.* **65**, 9 (2010); M. G. Albrow, T. D. Coughlin and J. R. Forshaw, *Prog. Part. Nucl. Phys.* **65**, 149 (2010).
- [34] L. Adamczyk *et al.*, CERN-LHCC-2015-009, ATLAS-TDR-024.
- [35] M. Tasevsky [ATLAS Collaboration], *AIP Conf. Proc.* **1654**, 090001 (2015).
- [36] M. Albrow *et al.* [CMS and TOTEM Collaborations], CERN-LHCC-2014-021, TOTEM-TDR-003, CMS-TDR-13.
- [37] V. P. Gonçalves, D. E. Martins, M. S. Rangel and M. Tasevsky, *Phys. Rev. D* **102**, no.7, 074014 (2020).
- [38] D. E. Martins, M. Tasevsky and V. P. Goncalves, *Phys. Rev. D* **105**, no.11, 114002 (2022).
- [39] L. A. Harland-Lang, V. A. Khoze, M. G. Ryskin and M. Tasevsky, *JHEP* **1904**, 010 (2019).
- [40] M. Tasevsky, *Int. J. Mod. Phys. A* **29**, 1446012 (2014).
- [41] K. Cerny, M. Tasevsky, T. Sykora and R. Zlebcik, *JINST* **16** (2021) no.01, P01030.
- [42] G. Aad *et al.* [ATLAS], ATL-FWD-PUB-2021-002.
- [43] J. Ohnemus, T. F. Walsh and P. M. Zerwas, *Phys. Lett. B* **328**, 369-373 (1994)
- [44] L. A. Harland-Lang, C. H. Kom, K. Sakurai and W. J. Stirling, *Eur. Phys. J. C* **72**, 1969 (2012)
- [45] S. Fichtel, G. von Gersdorff, O. Kepka, B. Lenzi, C. Royon and M. Saimpert, *Phys. Rev. D* **89**, 114004 (2014).
- [46] V. P. Goncalves, W. K. Sauter and M. Thiel, *Phys. Rev. D* **89**, no.7, 076003 (2014).
- [47] V. P. Goncalves and W. K. Sauter, *Phys. Rev. D* **91**, 035004 (2015).
- [48] P. Lebiedowicz and A. Szczurek, *Phys. Rev. D* **91**, 095008 (2015).

- [49] P. Lebiedowicz, M. Luszczak, R. Pasechnik and A. Szczurek, Phys. Rev. D **94**, no.1, 015023 (2016).
- [50] C. Baldenegro, S. Fichet, G. von Gersdorff and C. Royon, JHEP **06**, 142 (2017).
- [51] V. P. Goncalves and W. K. Sauter, Phys. Lett. B **811**, 135981 (2020).
- [52] L. Basso, B. Fuks, M. E. Krauss and W. Porod, JHEP **07**, 147 (2015).
- [53] K. Kumericki, I. Picek and B. Radovic, Phys. Rev. D **86**, 013006 (2012).
- [54] A. Melfo, M. Nemevsek, F. Nesti, G. Senjanovic and Y. Zhang, Phys. Rev. D **85**, 055018 (2012).
- [55] A. Arhrib, B. Bajc, D. K. Ghosh, T. Han, G. Y. Huang, I. Puljak and G. Senjanovic, Phys. Rev. D **82**, 053004 (2010)
- [56] P. Bandyopadhyay, S. Choubey and M. Mitra, JHEP **10**, 012 (2009)
- [57] T. Toma and A. Vicente, JHEP **01**, 160 (2014)
- [58] A. Vicente and C. E. Yaguna, JHEP **02**, 144 (2015)
- [59] S. Fraser, E. Ma and O. Popov, Phys. Lett. B **737**, 280-282 (2014)
- [60] A. Biswas, D. Borah and D. Nanda, JHEP **12**, 109 (2019)
- [61] B. A. Kniehl, Phys. Lett. B **254**, 267 (1991).
- [62] V. M. Budnev, I. F. Ginzburg, G. V. Meledin and V. G. Serbo, Phys. Rept. **15**, 181 (1975).
- [63] J. D. Bjorken, Phys. Rev. D **47**, 101 (1993).
- [64] V. A. Khoze, A. D. Martin and M. G. Ryskin, Int. J. Mod. Phys. A **30** no.08, 1542004 (2015).
- [65] E. Gotsman, E. Levin and U. Maor, Int. J. Mod. Phys. A **30**, no. 08, 1542005 (2015).
- [66] C. O. Rasmussen and T. Sjöstrand, JHEP **1602**, 142 (2016).
- [67] A. V. Manohar, P. Nason, G. P. Salam and G. Zanderighi, JHEP **12**, 046 (2017).
- [68] V. Bertone *et al.* [NNPDF], SciPost Phys. **5**, no.1, 008 (2018).
- [69] L. A. Harland-Lang, A. D. Martin, R. Nathvani and R. S. Thorne, Eur. Phys. J. C **79**, no.10, 811 (2019).
- [70] G. G. da Silveira and V. P. Goncalves, Phys. Rev. D **92**, no.1, 014013 (2015).
- [71] G. G. da Silveira, V. P. Goncalves and G. G. Vargas Veronez, Phys. Rev. D **103**, no.7, 074021 (2021).
- [72] J. Alwall, M. Herquet, F. Maltoni *et al.*, JHEP **06**, 128 (2011).
- [73] N. D. Christensen and C. Duhr, Comput. Phys. Commun. **180**, 1614-1641 (2009)
- [74] J. de Favereau *et al.* [DELPHES 3], JHEP **02**, 057 (2014).
- [75] G. Cowan, K. Cranmer, E. Gross and O. Vitells, Eur. Phys. J. C **71** (2011), 1554 [erratum: Eur. Phys. J. C **73** (2013), 2501].