

Investigation of magnetic moment of $P_{cs}(4338)$ and $P_{cs}(4459)$ molecular pentaquarks

Ulaş Özdem^{1,*}

¹*Health Services Vocational School of Higher Education,
Istanbul Aydin University, Sefakoy-Kucukcekmece, 34295 Istanbul, Turkey*

In this study, we discuss the magnetic moment of $P_{cs}(4338)$ and $P_{cs}(4459)$ hidden-charmed molecular pentaquarks, which are closely related to their substructures. The magnetic moments of these states are calculated with the help of the light-cone sum rules method with quantum numbers $I(J^P) = 0(1/2^-)$ and $I(J^P) = 0(3/2^-)$ for $P_{cs}(4338)$ and $P_{cs}(4459)$, respectively. Our prediction for the magnetic moment $\mu_{P_{cs}} = 0.34 \pm 0.12 \mu_N$ for the $P_{cs}(4338)$ state and $\mu_{P_{cs}} = -1.67 \pm 0.58 \mu_N$ for the $P_{cs}(4459)$ state. As a byproduct, the magnetic moments of the isospin-1 partners of these states have been also obtained. The magnetic moment are obtained as $\mu_{P_{cs}} = 0.63 \pm 0.21 \mu_N$ and $\mu_{P_{cs}} = -3.33 \pm 1.04 \mu_N$ for the isospin-1 partners of the $P_{cs}(4338)$ and $P_{cs}(4459)$ states, respectively. Our results regarding the magnetic moments of these molecular pentaquark states are compared with the results in the literature.

Keywords: Hidden-charm pentaquarks, magnetic moment, molecular picture, light-cone sum rules

I. INTRODUCTION

The discovery and investigation of hadron states with quantum numbers that go beyond the traditional hadron structure is one of the most attractive platforms of hadron spectroscopy due to the insight that these states can ensure information on confinement, gluonic degrees of freedom and, on QCD in the GeV scale. Since the discovery of the $X(3872)$ in 2003 by Belle Collaboration, many unconventional states were discovered in the past two decades. The different elucidations also have appeared together with conventional hadrons, loosely bound molecules, compact tetraquarks or pentaquarks, kinematic effects, glueballs, hybrids and so on. Though there is still a long way to precisely figure out how the strong interaction binds these quarks and antiquarks together, this subject has become one of the most attractive research subjects in hadron physics. For the recent experimental and theoretical progresses on the unconventional states see, for instance, Refs. [1–16].

In 2015, two pentaquark states, $P_c(4380)$ and $P_c(4450)$, were observed by the LHCb collaboration in the $J/\psi p$ invariant mass distribution [17]. The LHCb Collaboration then updated its previous analysis and announced that it had observed three new pentaquark states, $P_c(4312)$, $P_c(4440)$, and $P_c(4457)$ [18]. In 2020, the LHCb Collaboration announced a pentaquark state, $P_{cs}(4459)$, in the invariant mass spectrum of $J/\psi\Lambda$ in the $\Xi_b^0 \rightarrow J/\psi \Lambda K^-$ decay [19]. The measured mass and width are $4458.8 \pm 2.7_{-1.1}^{+4.7}$ MeV and $17.3 \pm 6.5_{-5.7}^{+8.0}$ MeV respectively. Since this pentaquark state is observed in the invariant mass distribution $J/\psi\Lambda$, the quark content is assumed to be $udsc\bar{c}$; However, the spin-parity quantum numbers have not yet been clarified. In 2021, an evidence for a new pentaquark structure, $P_c(4337)$, was found by the LHCb Collaboration in $B_s \rightarrow J/\psi p\bar{p}$ decays [20] and

spin-parity quantum numbers of this state were predicted with $J^P = 1/2^+$ [21]. Recently, the LHCb collaboration observed an evidence for a new structure $P_{cs}(4338)$ in the $J/\psi\Lambda$ mass distribution in the $B^- \rightarrow J/\psi\Lambda^- p$ decays [22]. The measured mass and width are $4338.2 \pm 0.7 \pm 0.4$ MeV and $7.0 \pm 1.2 \pm 1.3$ MeV respectively and the amplitude analysis prefers the $J^P = 1/2^-$ spin-parity quantum numbers and excludes the possibility of $J^P = 1/2^+$ at 90% confidence level. As regards to its decay products, one can conclude that this newly reported state consist of $udsc\bar{c}$ quarks. After the experimental discovery was announced, several studies were conducted to understand the spectroscopic properties of this state [23–28].

Studies in the literature for currently experimentally observed pentaquark states indicate that spectroscopic analyzes may not be sufficient to detect the properties of such states. In addition to the spectroscopic properties of these states, investigation of their strong, weak and electromagnetic decays may contribute to the understanding of the nature of these states. In the literature, there are few studies on the calculation of electromagnetic multipole moments and transitions magnetic moments of pentaquark states by means of different configurations [29–39]. In this study, we study the magnetic moment of $P_{cs}(4338)$ and $P_{cs}(4459)$ pentaquark states (hereafter we will show these states as P_{cs}^1 and P_{cs}^2 , respectively) making use of the light-cone sum rule formalism in the molecular picture. The magnetic moments belong to the non-perturbative domain of QCD, and to achieve their calculations we need non-perturbative methods. QCD sum rule is one of the effective techniques that consider the non-perturbative effects. One of the modifications of the traditional QCD sum rules technique is the light-cone version where operator product expansion is applied over the twist of operators instead of the dimensions of operators [40–42].

This paper is organized in the following manner: In Sect. II, we will briefly describe the formalism of the light-cone sum rules method, which is used to obtain the magnetic moments of the P_{cs}^1 pentaquark in the first step

* ulasozdem@aydin.edu.tr

and then the P_{cs}^2 pentaquark. Sect. III provides the details of numerical computations to get the numerical results. Sect. IV is reserved for summary and concluding remarks. In Appendix A, we will present explicit ex-

pressions of the magnetic moment of the P_{cs}^1 pentaquark state. The explicit expressions of the photon distribution amplitudes are also given in Appendix B.

II. THE LIGHT-CONE SUM RULES FOR THE PENTAQUARK MOLECULAR STATES

A. Formalism of the P_{cs}^1 state

The correlation function, which is essential for magnetic moment calculations, is written in the following form:

$$\Pi(p, q) = i \int d^4x e^{ip \cdot x} \langle 0 | T \left\{ J^{P_{cs}^1}(x) \bar{J}^{P_{cs}^1}(0) \right\} | 0 \rangle_\gamma, \quad (1)$$

where sub-indices γ is the external electromagnetic field. Here, $J(x)$ represents the interpolating current of the state considered, which in our case P_{cs}^1 pentaquark state, and it is required for further calculations. This interpolating current is written with isospin and spin-parity $I(J^P) = 0(1/2^-)$ as follows [23]:

$$J^{P_{cs}^1}(x) = \frac{1}{\sqrt{2}} \left\{ | \bar{D}^0 \Xi_c^0 \rangle - | \bar{D}^- \Xi_c^+ \rangle \right\} = \frac{1}{\sqrt{2}} \left\{ [\bar{c}^d(x) i \gamma_5 u^d(x)] [\varepsilon^{abc} d^{aT}(x) C \gamma_5 s^b(x) c^c(x)] \right. \\ \left. - [\bar{c}^d(x) i \gamma_5 d^d(x)] [\varepsilon^{abc} u^{aT}(x) C \gamma_5 s^b(x) c^c(x)] \right\}, \quad (2)$$

where a, b, c and d being color indices and the C is the charge conjugation operator.

To obtain the hadronic representation of the correlation function, we insert a full set of intermediate states P_{cs}^1 with the same quantum numbers as the interpolation currents into the correlation function. As a result, we get the following expression

$$\Pi^{Had}(p, q) = \frac{\langle 0 | J(x) | P_{cs}(p, s) \rangle \langle P_{cs}(p, s) | P_{cs}(p+q, s) \rangle_\gamma \langle P_{cs}(p+q, s) | \bar{J}(0) | 0 \rangle}{[p^2 - m_{P_{cs}}^2]} + \dots \quad (3)$$

The matrix element $\langle P_{cs}(p, s) | P_{cs}(p+q, s) \rangle_\gamma$ in Eq. (3) can be defined in relation to form factors as follows:

$$\langle P_{cs}(p, s) | P_{cs}(p+q, s) \rangle_\gamma = \varepsilon^\mu \bar{u}(p, s) \left[[F_1(q^2) + F_2(q^2)] \gamma_\mu + F_2(q^2) \frac{(2p+q)_\mu}{2m_{P_{cs}}} \right] u(p+q, s). \quad (4)$$

After the Eq. (4) is substituted in the Eq. (3) and some calculations are made, the following result is obtained for the hadronic side,

$$\Pi^{Had}(p, q) = \lambda_{P_{cs}}^2 \gamma_5 \frac{(\not{p} + m_{P_{cs}})}{[p^2 - m_{P_{cs}}^2]} \varepsilon^\mu \left[[F_1(q^2) + F_2(q^2)] \gamma_\mu + F_2(q^2) \frac{(2p+q)_\mu}{2m_{P_{cs}}} \right] \gamma_5 \frac{(\not{p} + \not{q} + m_{P_{cs}})}{[(p+q)^2 - m_{P_{cs}}^2]}. \quad (5)$$

The value of form factors $F_1(q^2)$ and $F_2(q^2)$ give us the magnetic form factor $F_M(q^2)$ at different q^2 :

$$F_M(q^2) = F_1(q^2) + F_2(q^2). \quad (6)$$

For real photon, i.e. $q^2 = 0$, magnetic form factor $F_M(q^2 = 0)$ is proportional to the magnetic moment $\mu_{P_{cs}}$:

$$\mu_{P_{cs}} = \frac{e}{2m_{P_{cs}}} F_M(q^2 = 0). \quad (7)$$

The second representation of the correlation function, the QCD side, is achieved employing the interpolating currents explicitly into the correlation function. Then, the proper quark fields are contracted with the help of Wick's theorem and the demanded outcomes are achieved. As a result of above procedures, the QCD representation of the correlation function is obtained as follows:

$$\Pi^{QCD}(p, q) = -\frac{1}{2} \varepsilon^{abc} \varepsilon^{a'b'c'} \int d^4x e^{ip \cdot x} \langle 0 | \left\{ \text{Tr} \left[\gamma_5 S_u^{dd'}(x) \gamma_5 S_c^{d'd}(-x) \right] \text{Tr} \left[\gamma_5 S_s^{bb'}(x) \gamma_5 \tilde{S}_d^{aa'}(x) \right] \right. \\ \left. - \text{Tr} \left[\gamma_5 S_u^{da'}(x) \gamma_5 \tilde{S}_s^{bb'}(x) \gamma_5 S_d^{ad'}(x) \gamma_5 S_c^{d'd}(-x) \right] - \text{Tr} \left[\gamma_5 S_d^{da'}(x) \gamma_5 \tilde{S}_s^{bb'}(x) \gamma_5 S_u^{ad'}(x) \gamma_5 S_c^{d'd}(-x) \right] \right. \\ \left. + \text{Tr} \left[\gamma_5 S_d^{dd'}(x) \gamma_5 S_c^{d'd}(-x) \right] \text{Tr} \left[\gamma_5 S_s^{bb'}(x) \gamma_5 \tilde{S}_u^{aa'}(x) \right] \right\} | 0 \rangle_\gamma, \quad (8)$$

where $\tilde{S}_{c(q)}^{ij}(x) = C S_{c(q)}^{ijT}(x) C$ and, $S_c(x)$ and $S_q(x)$ are the charm and light quark propagators, respectively. The expressions of these propagators are given as [43, 44]

$$S_q(x) = \frac{1}{2\pi x^2} \left(i \frac{\not{x}}{x^2} - \frac{m_q}{2} \right) - \frac{\langle \bar{q}q \rangle}{12} \left(1 - i \frac{m_q \not{x}}{4} \right) - \frac{\langle \bar{q}q \rangle}{192} m_0^2 x^2 \left(1 - i \frac{m_q \not{x}}{6} \right) - \frac{ig_s}{32\pi^2 x^2} G^{\mu\nu}(x) \left[\not{x} \sigma_{\mu\nu} + \sigma_{\mu\nu} \not{x} \right], \quad (9)$$

$$S_c(x) = \frac{m_c^2}{4\pi^2} \left[\frac{K_1(m_c \sqrt{-x^2})}{\sqrt{-x^2}} + i \frac{\not{x} K_2(m_c \sqrt{-x^2})}{(\sqrt{-x^2})^2} \right] - \frac{g_s m_c}{16\pi^2} \int_0^1 dv G^{\mu\nu}(vx) \left[(\sigma_{\mu\nu} \not{x} + \not{x} \sigma_{\mu\nu}) \frac{K_1(m_c \sqrt{-x^2})}{\sqrt{-x^2}} + 2\sigma_{\mu\nu} K_0(m_c \sqrt{-x^2}) \right], \quad (10)$$

where $\langle \bar{q}q \rangle$, $G^{\mu\nu}$, v , and K_i 's are light-quark condensate, the gluon field strength tensor, line variable and modified Bessel functions of the second kind, respectively. The first term of the light and heavy quark propagators correspond to perturbative or free part and the rest belong to the interacting parts.

The correlation function given in Eq. (8) receives both perturbative, i.e., when photon interacts perturbatively with quark propagators, and nonperturbative, i.e., photon interacts with light quarks at large distance, contributions. In the first part, the propagator of the quark interacting with the photon perturbatively is replaced by

$$S^{free}(x) \rightarrow \int d^4y S^{free}(x-z) \mathcal{A}(z) S^{free}(z), \quad (11)$$

and the remaining propagators in Eq. (8) are substituted with the full quark propagators including the perturbative and nonperturbative parts. Here we use $A_\mu(z) = -\frac{1}{2} F_{\mu\nu}(z) z^\nu$ where the electromagnetic field strength tensor is written as $F_{\mu\nu}(z) = -i(\varepsilon_\mu q_\nu - \varepsilon_\nu q_\mu) e^{iq \cdot z}$. The total perturbative contribution is achieved by performing the replacement mentioned above for the perturbatively interacting quark propagator with the photon and making use of the replacement of the remaining propagators by their free parts.

In the next part, one of the light quark propagators in Eq. (9), defining the photon emission at large distances, is substitute by

$$S_{\alpha\beta}^{ab}(x) \rightarrow -\frac{1}{4} [\bar{q}^a(x) \Gamma_i q^b(x)] (\Gamma_i)_{\alpha\beta}, \quad (12)$$

and the rest propagators are substituted with the full quark propagators. Here, Γ_i represent the full set of Dirac matrices. Once Eq. (12) is plugged into Eq. (8), there appear matrix elements of $\langle \gamma(q) | \bar{q}(x) \Gamma_i q(0) | 0 \rangle$ and $\langle \gamma(q) | \bar{q}(x) \Gamma_i G_{\alpha\beta} q(0) | 0 \rangle$ kinds, representing the nonperturbative contributions. To calculate the nonperturbative contributions, we need these matrix elements which are parameterized in terms of photon wave functions with definite twists. The explicit expressions of the photon distribution amplitudes (DAs) are presented in Ref. [45]. The QCD side of the correlation function can be acquired in terms of quark-gluon parameters as well as the DAs of the photon using Eqs. (8)–(12) and after performing the Fourier transformation to remove the calculations to the momentum space.

As a final step, the \not{x} Lorentz structure is chosen from both representations and the coefficients of the Lorentz structure are matched from both the hadronic and QCD representations. To eliminate the effects of the continuum and higher states, Borel transformation and continuum subtraction are performed. After these procedures, we obtain the demanded light-cone sum rules for the magnetic moments:

$$\mu_{P_{cs}} \lambda_{P_{cs}}^2 m_{P_{cs}} = e \frac{m_{P_{cs}}^2}{M^2} \Delta_1^{QCD}(M^2, s_0). \quad (13)$$

The analytical expressions obtained for the $\Delta_1^{QCD}(M^2, s_0)$ function are given in the Appendix A.

B. Formalism of the P_{cs}^2 state

For the magnetic moment of P_{cs}^2 state required correlation function is given as

$$\Pi_{\mu\nu}(p, q) = i \int d^4x e^{ip \cdot x} \langle 0 | T \left\{ J_\mu^{P_{cs}^2}(x) \bar{J}_\nu^{P_{cs}^2}(0) \right\} | 0 \rangle_\gamma. \quad (14)$$

The interpolating current used for P_{cs}^2 pentaquark state with isospin and spin-parity $I(J^P) = 0(3/2^-)$ is as follows [23]:

$$J_{\mu}^{P_{cs}^2}(x) = \frac{1}{\sqrt{2}} \left\{ |\bar{D}^{*0}\Xi_c^0\rangle - |\bar{D}^{*-}\Xi_c^+\rangle \right\} = \frac{1}{\sqrt{2}} \left\{ [\bar{c}^d(x)\gamma_{\mu}u^d(x)] [\varepsilon^{abc}d^{aT}(x)C\gamma_5s^b(x)c^c(x)] \right. \\ \left. - [\bar{c}^d(x)\gamma_{\mu}d^d(x)] [\varepsilon^{abc}u^{aT}(x)C\gamma_5s^b(x)c^c(x)] \right\}. \quad (15)$$

The correlation function obtained depending on the hadron parameters is written as,

$$\Pi_{\mu\nu}^{Had}(p, q) = \frac{\langle 0 | J_{\mu}^{P_{cs}^2}(x) | P_{cs}^2(p, s) \rangle \langle P_{cs}^2(p, s) | P_{cs}^2(p+q, s) \rangle_{\gamma}}{[p^2 - m_{P_{cs}^2}^2]} \frac{\langle P_{cs}^2(p+q, s) | \bar{J}_{\nu}^{P_{cs}^2}(0) | 0 \rangle}{[(p+q)^2 - m_{P_{cs}^2}^2]} + \dots \quad (16)$$

The matrix elements of the interpolating current between the vacuum and the P_{cs}^2 pentaquark are defined as

$$\langle 0 | J_{\mu}^{P_{cs}^2}(x) | P_{cs}^2(p, s) \rangle = \lambda_{P_{cs}^2} u_{\mu}(p, s), \\ \langle P_{cs}^2(p+q, s) | \bar{J}_{\nu}^{P_{cs}^2}(0) | 0 \rangle = \lambda_{P_{cs}^2} \bar{u}_{\nu}(p+q, s), \quad (17)$$

where the $u_{\mu}(p, s)$, $u_{\nu}(p+q, s)$ and $\lambda_{P_{cs}^2}$ are the spinors and residue doubly-heavy P_{cs}^2 pentaquark states, respectively. Summation over spins of P_{cs}^2 pentaquark state is performed as:

$$\sum_s u_{\mu}(p, s) \bar{u}_{\nu}(p, s) = -(\not{p} + m_{P_{cs}^2}) \left[g_{\mu\nu} - \frac{1}{3} \gamma_{\mu} \gamma_{\nu} - \frac{2p_{\mu}p_{\nu}}{3m_{P_{cs}^2}^2} + \frac{p_{\mu}\gamma_{\nu} - p_{\nu}\gamma_{\mu}}{3m_{P_{cs}^2}} \right]. \quad (18)$$

The transition matrix element $\langle P_{cs}^2(p) | P_{cs}^2(p+q) \rangle_{\gamma}$ entering Eq. (16) can be written as follows [46–49]:

$$\langle P_{cs}^2(p, s) | P_{cs}^2(p+q, s) \rangle_{\gamma} = -e \bar{u}_{\mu}(p, s) \left[F_1(q^2) g_{\mu\nu} \not{\epsilon} - \frac{1}{2m_{P_{cs}^2}} \left[F_2(q^2) g_{\mu\nu} + F_4(q^2) \frac{q_{\mu}q_{\nu}}{(2m_{P_{cs}^2})^2} \right] \not{\epsilon} \not{q} \right. \\ \left. + F_3(q^2) \frac{1}{(2m_{P_{cs}^2})^2} q_{\mu}q_{\nu} \not{\epsilon} \right] u_{\nu}(p+q, s). \quad (19)$$

where F_i 's are the Lorentz invariant form factors.

In principle, using the Eqs. (16)–(19), we can get the final expression of the hadronic side of the correlation function, however in doing so we encounter two problems: not all Lorentz structures are independent, and the correlation function can also receive contributions from spin-1/2 particles that need to be eliminated. In fact, the matrix element of the current J_{μ} between spin-1/2 pentaquarks and vacuum is nonzero and is determined as

$$\langle 0 | J_{\mu}(0) | B(p, s = 1/2) \rangle = (Ap_{\mu} + B\gamma_{\mu})u(p, s = 1/2). \quad (20)$$

As is seen the unwanted spin-1/2 contributions are proportional to γ_{μ} and p_{μ} . By multiplying both sides with γ^{μ} and using the condition $\gamma^{\mu}J_{\mu} = 0$ one can determine the constant A in terms of B. To remove the spin-1/2 pollutions and obtain only independent structures in the correlation function, we apply the ordering for Dirac matrices as $\gamma_{\mu}\not{p}\not{\epsilon}\not{q}\gamma_{\nu}$ and eliminate terms with γ_{μ} at the beginning, γ_{ν} at the end and those proportional to p_{μ} and p_{ν} [50]. Consequently, employing Eqs. (16)–(20) the hadronic side take the form,

$$\Pi_{\mu\nu}^{Had}(p, q) = \frac{\lambda_{P_{cs}^2}^2}{[(p+q)^2 - m_{P_{cs}^2}^2][p^2 - m_{P_{cs}^2}^2]} \left[g_{\mu\nu} \not{p}\not{\epsilon}\not{q} F_1(q^2) - m_{P_{cs}^2} g_{\mu\nu} \not{\epsilon}\not{q} F_2(q^2) - \frac{F_3(q^2)}{4m_{P_{cs}^2}} q_{\mu}q_{\nu} \not{\epsilon}\not{q} \right. \\ \left. - \frac{F_4(q^2)}{4m_{P_{cs}^2}^3} (\varepsilon \cdot p) q_{\mu}q_{\nu} \not{p}\not{q} + \dots \right]. \quad (21)$$

The final expression of the hadronic representation for the chosen Lorentz structures is obtained as follows:

$$\Pi_{\mu\nu}^{Had}(p, q) = \Pi_1^{Had} g_{\mu\nu} \not{p}\not{\epsilon}\not{q} + \Pi_2^{Had} g_{\mu\nu} \not{\epsilon}\not{q} + \dots, \quad (22)$$

where Π_1^{Had} and Π_2^{Had} are functions of the form factors $F_1(q^2)$ and $F_2(q^2)$, respectively; and other independent structures and form factors are denoted by dots.

The magnetic form factor, $G_M(q^2)$, is characterized with respect to the form factors $F_i(q^2)$ as follows [46–49]:

$$G_M(q^2) = [F_1(q^2) + F_2(q^2)](1 + \frac{4}{5}\tau) - \frac{2}{5}[F_3(q^2) + F_4(q^2)]\tau(1 + \tau), \quad (23)$$

where $\tau = -\frac{q^2}{4m_{P_{cs}^2}}$. At $q^2 = 0$, the magnetic moment is obtained with respect to the functions $F_1(0)$ and $F_2(0)$ form factors as:

$$G_M(0) = F_1(0) + F_2(0). \quad (24)$$

The magnetic moment of the spin-3/2 state, $(\mu_{P_{cs}^2})$, is described as follows,

$$\mu_{P_{cs}^2} = \frac{e}{2m_{P_{cs}^2}}G_M(0). \quad (25)$$

When the above procedures are applied, the analysis with regards to hadronic parameters, which is the first step of the light-cone sum rule analysis, is completed.

The next step in the light-cone sum rule analysis is to obtain the correlation function in relation to the quark-gluon parameters as well as the photon DAs. Doing again the steps in the previous subsection yields the following result:

$$\begin{aligned} \Pi_{\mu\nu}^{QCD}(p, q) = & \frac{i}{2}\varepsilon^{abc}\varepsilon^{a'b'c'} \int d^4x e^{ip\cdot x} \langle 0 | \left\{ \text{Tr} \left[\gamma_\mu S_u^{dd'}(x) \gamma_\nu S_c^{d'd}(-x) \right] \text{Tr} \left[\gamma_5 S_s^{bb'}(x) \gamma_5 \tilde{S}_d^{aa'}(x) \right] \right. \\ & - \text{Tr} \left[\gamma_\mu S_u^{da'}(x) \gamma_5 \tilde{S}_s^{bb'}(x) \gamma_5 S_d^{ad'}(x) \gamma_\nu S_c^{d'd}(-x) \right] - \text{Tr} \left[\gamma_\mu S_d^{da'}(x) \gamma_5 \tilde{S}_s^{bb'}(x) \gamma_5 S_u^{ad'}(x) \gamma_\nu S_c^{d'd}(-x) \right] \\ & \left. + \text{Tr} \left[\gamma_\mu S_d^{dd'}(x) \gamma_\nu S_c^{d'd}(-x) \right] \text{Tr} \left[\gamma_5 S_s^{bb'}(x) \gamma_5 \tilde{S}_u^{aa'}(x) \right] \right\} | 0 \rangle_\gamma, \end{aligned} \quad (26)$$

As a result, the QCD representation of the correlation function in relation to the chosen Lorentz structures is obtained as:

$$\Pi_{\mu\nu}^{QCD}(p, q) = \Pi_1^{QCD} g_{\mu\nu} \not{p} \not{q} + \Pi_2^{QCD} g_{\mu\nu} \not{q} + \dots \quad (27)$$

where Π_1^{QCD} and Π_2^{QCD} are functions of the QCD degrees of freedom and photon DAs parameters. The analytical expressions obtained for Π_i^{QCD} functions are not given in the text because they are very lengthy.

The correlation function has been obtained in connection with both hadronic and quark-gluon parameters. For magnetic moment analysis, the QCD and hadronic representations of the correlation function are equated using the quark-hadron duality approach. For the form factor F_1 and F_2 , we obtain the light-cone sum rules by matching the coefficients of the Lorentz structures $g_{\mu\nu} \not{p} \not{q}$ and $g_{\mu\nu} \not{q}$, respectively.

$$\Pi_{\mu\nu}^{Had}(p, q) = \Pi_{\mu\nu}^{QCD}(p, q). \quad (28)$$

Analytical results are obtained for both P_{cs}^1 and P_{cs}^2 pentaquark states. The next step would be to perform numerical calculations for these pentaquark states. We should also mentioned that the magnetic moments of the isospin-1 partners of the P_{cs}^1 and P_{cs}^2 states have also been calculated. To do this, we substitute the "+" sign instead of the "-" sign in Eqs. (2) and (15). For the sake of brevity, only the analytical results of the isospin-0 P_{cs}^1 and P_{cs}^2 states have been presented.

III. NUMERICAL ANALYSIS AND DISCUSSION

The light cone sum rule for the magnetic moment of the P_{cs} states contains many input parameters whose numerical values we need. The values of these parameters we use in the analysis are given as follows: $m_u = m_d = 0$, $m_s = 93.4_{-3.4}^{+8.6}$ MeV, $m_c = 1.27 \pm 0.02$ GeV [51], $m_{P_{cs}^1} = 4338.2 \pm 0.7 \pm 0.4$ MeV, $m_{P_{cs}^2} = 4458.8 \pm 2.9_{-1.1}^{+4.7}$ MeV, $f_{3\gamma} = -0.0039$ GeV² [45], $\langle \bar{s}s \rangle = 0.8 \langle \bar{u}u \rangle$ GeV³ with $\langle \bar{u}u \rangle = \langle \bar{d}d \rangle = (-0.24 \pm 0.01)^3$ GeV³ [52],

$m_0^2 = 0.8 \pm 0.1$ GeV² [52], $\langle g_s^2 G^2 \rangle = 0.88$ GeV⁴ [5], $\lambda_{P_{cs}^1} = (1.43_{-0.18}^{+0.19}) \times 10^{-3}$ GeV⁶ and $\lambda_{P_{cs}^2} = (1.69_{-0.21}^{+0.22}) \times 10^{-3}$ GeV⁶ [23]. The photon DAs and explicit expressions of the wave functions used in these DAs and their input parameters are given in Appendix B.

The light-cone sum rules for magnetic moments of these pentaquarks are obtained as a function of the Borel mass parameter M^2 and continuum threshold s_0 . To get a reliable light-cone sum rules result, one should define appropriate working regions for these two helping parameters. The operator product expansion (OPE) con-




FIG. 1. The magnetic moments of isospin-0 ($[I, I_3] = [0, 0]$) pentaquark states versus M^2 at three different values of s_0 ; (a) for P_{cs}^1 state and (b) for P_{cs}^2 state (in unit of μ_N).

vergence and pole dominance (PC) limitations are widely used to define the working regions of these two helping parameters. Bearing in mind these limitations, the following working regions are obtained for these two helping parameters as a result of the numerical analysis,

$$\begin{aligned} 5.0 \text{ GeV}^2 &\leq M^2 \leq 7.0 \text{ GeV}^2, \\ 22.5 \text{ GeV}^2 &\leq s_0 \leq 24.5 \text{ GeV}^2, \end{aligned} \quad (29)$$

for the P_{cs}^1 state and

$$\begin{aligned} 5.0 \text{ GeV}^2 &\leq M^2 \leq 7.0 \text{ GeV}^2, \\ 23.5 \text{ GeV}^2 &\leq s_0 \leq 25.5 \text{ GeV}^2, \end{aligned} \quad (30)$$

for the P_{cs}^2 state.

Using the above working regions for the M^2 and s_0 , the PC changes in the intervals $60\% \geq \text{PC} \geq 33\%$. At

$M_{max}^2 = 7.0 \text{ GeV}^2$, the PC is equal to 33%, whilst at $M_{min}^2 = 5.0 \text{ GeV}^2$, it is equal to 60%. In the standard analysis of QCD sum rules, the PC should be larger than 50% for baryons and mesons. In the case of multi-quark states, it turns out to be as $\text{PC} > 20\%$. When we investigate the OPE convergence, we have acquired that the contribution of the higher dimensional terms in OPE is less than $\sim 2\%$, thus the convergence of the sum rules is ensured. As is seen the chosen working regions for M^2 and s_0 fulfill the requirements of the light-cone sum rules technique. For completeness, in Fig. 1, we also depict the dependence of the magnetic moment of P_{cs}^1 and P_{cs}^2 states, on the Borel mass parameter, M^2 at various values of s_0 . As you can see from this figure, the change of magnetic moments with respect to M^2 is reasonably stable. Though the change is high compared to s_0 , this variation stays within the error limits of the light-cone sum rules method.

Now that we have determined the numerical values of all the input parameters, we can start performing the numerical analysis. Our final results for magnetic moments are given as:

$$\begin{aligned} \mu_{P_{cs}^1} &= 0.34 \pm 0.12 \mu_N, \\ \mu_{P_{cs}^2} &= -1.67 \pm 0.58 \mu_N. \end{aligned} \quad (31)$$

The errors in the magnetic moment results given above are due to the input parameters, auxiliary parameters such as s_0 and M^2 , as well as the input parameters used in the photon DAs.

In Refs. [35, 36], the magnetic moments of the molecular pentaquark states have been investigated by means of the constituent quark model based on different combinations of the flavour-spin wave functions of the component hadrons. In Ref. [36], the magnetic moment of the P_{cs}^1 state is obtained as $\mu_{P_{cs}^1} = 0.377 \mu_N$. For the P_{cs}^2 state, while $\mu_{P_{cs}^2} = 0.465 \mu_N$ was found for magnetic moment in Ref. [35], $\mu_{P_{cs}^2} = -0.231 \mu_N$ result was obtained in Ref. [36]. Our result for the magnetic moment of state P_{cs}^1 seems to be consistent with the result obtained in Ref. [36]. For the P_{cs}^2 pentaquark state, there are large discrepancy among results not only the magnitude but also by the sign.

Our final remark is that the magnetic moment of the isospin-1 ($[I, I_3] = [1, 0]$) partners of these pentaquark states have also been obtained. The results are given as

$$\begin{aligned}\mu_{P_{cs}^1} &= 0.63 \pm 0.21 \mu_N, \\ \mu_{P_{cs}^2} &= -3.33 \pm 1.04 \mu_N.\end{aligned}\quad (32)$$

When these results are examined, it is seen that the difference between the magnetic moment values of isospin-0, ($[I, I_3] = [0, 0]$), and isospin-1, ($[I, I_3] = [1, 0]$), pentaquark states is almost twice. In Ref. [36], the magnetic moments of the isospin-1 partners of these pentaquark states have been calculated and the results are obtained as: $\mu_{P_{cs}^1} = 0.377 \mu_N$ and $\mu_{P_{cs}^2} = 0.465 \mu_N$. It is seen that the obtained results are not consistent with each other.

It appears from these results that different approaches lead to quite different estimates for the P_{cs}^1 and P_{cs}^2 magnetic moments that can be used to distinguish these models. Obviously, it seems that more study is needed to understand the current situation. Hopefully, our study may attract the lattice QCD and experiment plans in future.

IV. SUMMARY AND CONCLUDING REMARKS

Searching for multi-quark states is an interesting topic in hadron physics. The recently reported pentaquark state, the hidden-charmed strange $P_{cs}(4338)$, added a new member to the pentaquark family. The mass and width of this state were measured as $4338.2 \pm 0.7 \pm 0.4$ MeV and $7.0 \pm 1.2 \pm 1.3$ MeV respectively. Inspired

by this, in this study, we discuss the magnetic moment of $P_{cs}(4338)$ and $P_{cs}(4459)$ hidden-charmed molecular pentaquarks, which are closely related to their substructures. The magnetic moments of these states are calculated with the help of the light-cone sum rules method with quantum numbers $I(J^P) = 0(1/2^-)$ and $I(J^P) = 0(3/2^-)$. Our prediction for the magnetic moment $\mu_{P_{cs}} = 0.34 \pm 0.12 \mu_N$ for the $P_{cs}(4338)$ state and $\mu_{P_{cs}} = -1.67 \pm 0.58 \mu_N$ for the $P_{cs}(4459)$ state. As a byproduct, the magnetic moments of the isospin-1 partners of these states have been also obtained. The magnetic moment are obtained as $\mu_{P_{cs}} = 0.63 \pm 0.21 \mu_N$ and $\mu_{P_{cs}} = -3.33 \pm 1.04 \mu_N$ for the isospin-1 partners of the $P_{cs}(4338)$ and $P_{cs}(4459)$ states, respectively. Our results regarding the magnetic moments of these molecular pentaquark states are compared with the results in the literature.

It is important to examine features such states theoretically in different perspectives not only to ensure some insights into the future experiments but also to better figure out the inner structure of these pentaquark states. The magnetic moments of hadronic molecular states play an important role in understanding their internal structure. We hope that our current research, together with existing studies in the literature, can inspire our colleagues to focus on the electromagnetic properties of hadronic molecular pentaquark states in the future. With these efforts, our information of the features of hadronic molecular pentaquark states will turn into more abundant. We are looking forward to future experimental progress along these directions.

Appendix A: Explicit expression for $\Delta_1^{QCD}(M^2, s_0)$

In this Appendix, we present the explicit expressions of the function $\Delta_1^{QCD}(M^2, s_0)$ for the magnetic moment of the P_{cs}^1 state entering into the sum rule.

$$\begin{aligned}\Delta_1^{QCD}(M^2, s_0) &= \frac{m_c P_1 P_2}{679477248\pi^5} \left[2e_c \left\{ -6m_c \left(5I[0, 3, 1, 0] + 2I[0, 3, 1, 1] - 7I[0, 3, 1, 2] - 2(5I[0, 3, 2, 0] + I[0, 3, 2, 1]) \right. \right. \right. \\ &\quad + 5I[0, 3, 3, 0] \Big) + 3m_0^2 \left(18m_c (I[0, 2, 1, 0] - I[0, 2, 1, 1] - I[0, 2, 2, 0]) - m_s (I[0, 2, 1, 0] - 2I[0, 2, 1, 1] \right. \\ &\quad + I[0, 2, 1, 2] - 2I[0, 2, 2, 0] + 2I[0, 2, 2, 1] + I[0, 2, 3, 0] - 2I[1, 1, 1, 0] - 4I[1, 1, 1, 1] + 2I[1, 1, 1, 2] \\ &\quad - 4I[1, 1, 2, 0] + 4I[1, 1, 2, 1] + 2I[1, 1, 3, 0]) \Big) - 4m_s \left(31I[0, 3, 1, 0] - 63I[0, 3, 1, 1] + 33I[0, 3, 1, 2] \right. \\ &\quad - I[0, 3, 1, 3] - 62I[0, 3, 2, 0] + 64I[0, 3, 2, 1] - 2I[0, 3, 2, 2] + 31I[0, 3, 3, 0] - I[0, 3, 3, 1] + 3I[1, 2, 1, 1] \\ &\quad \left. \left. \left. - 6I[1, 2, 1, 2] + 3I[1, 2, 1, 3] - 6I[1, 2, 2, 1] + 6I[1, 2, 2, 2] + 6I[1, 2, 3, 1] \right) \right\} \right]\end{aligned}$$

$$\begin{aligned}
& - e_d \left\{ 3m_0^2 \left(2m_s (I[0, 2, 1, 0] - 2I[0, 2, 1, 1] + I[0, 2, 1, 2] - 2I[0, 2, 2, 0] + 2I[0, 2, 2, 1] + I[0, 2, 3, 0] \right. \right. \\
& - 2I[1, 1, 1, 0] - 4I[1, 1, 1, 1] + 2I[1, 1, 1, 2] - 4I[1, 1, 2, 0] + 4I[1, 1, 2, 1] + 2I[1, 1, 3, 0]) \\
& + m_c (9I[0, 2, 1, 0] - 18I[0, 2, 1, 1] + 9I[0, 2, 1, 2] - 14I[0, 2, 2, 0] + 14I[0, 2, 2, 1] + 5I[0, 2, 3, 0] \\
& - 10I[1, 1, 1, 0] - 28I[1, 1, 1, 1] + 18I[1, 1, 1, 2] - 20I[1, 1, 2, 0] + 28I[1, 1, 2, 1] + 10I[1, 1, 3, 0]) \left. \right) \\
& + 4m_c \left(5I[0, 3, 1, 0] - 19I[0, 3, 1, 1] + 21I[0, 3, 1, 2] - 7I[0, 3, 1, 3] - 10I[0, 3, 2, 0] + 24I[0, 3, 2, 1] \right. \\
& - 12I[0, 3, 2, 2] + 5I[0, 3, 3, 0] - 5I[0, 3, 3, 1] + 15I[1, 2, 1, 1] - 36I[1, 2, 1, 2] + 21I[1, 2, 1, 3] \\
& - 30I[1, 2, 2, 1] + 36I[1, 2, 2, 2] + 15I[1, 2, 3, 1] \left. \right) + 8m_s \left(I[0, 3, 1, 0] - 3I[0, 3, 1, 1] + 3I[0, 3, 1, 2] \right. \\
& - I[0, 3, 1, 3] - 2I[0, 3, 2, 0] + 4I[0, 3, 2, 1] - 2I[0, 3, 2, 2] + I[0, 3, 3, 0] - I[0, 3, 3, 1] + 3I[1, 2, 1, 1] \\
& - 6I[1, 2, 1, 2] + 3I[1, 2, 1, 3] - 6I[1, 2, 2, 1] + 6I[1, 2, 2, 2] + 3I[1, 2, 3, 1] \left. \right) \left. \right\} \\
& - 2e_s m_c \left\{ 3m_0^2 \left(I[0, 2, 1, 0] - 2I[0, 2, 1, 1] + I[0, 2, 1, 2] - 6I[0, 2, 2, 0] + 6I[0, 2, 2, 1] + 5I[0, 2, 3, 0] \right. \right. \\
& - 10I[1, 1, 1, 0] - 12I[1, 1, 1, 1] + I[1, 1, 1, 2] - 20I[1, 1, 2, 0] + 12I[1, 1, 2, 1] + 10I[1, 1, 3, 0]) \\
& + 4 \left(5I[0, 3, 1, 0] - 11I[0, 3, 1, 1] + 9I[0, 3, 1, 2] - 3I[0, 3, 1, 3] - 10I[0, 3, 2, 0] + 16I[0, 3, 2, 1] \right. \\
& - 8I[0, 3, 2, 2] + 5I[0, 3, 3, 0] - 5I[0, 3, 3, 1] + 15I[1, 2, 1, 1] - 24I[1, 2, 1, 2] + 9I[1, 2, 1, 3] \\
& \left. \left. - 30I[1, 2, 2, 1] + 24I[1, 2, 2, 2] + 10I[1, 2, 3, 1] \right) \right\} \\
& - e_u \left\{ 3m_0^2 \left(2m_s (I[0, 2, 1, 0] - 2I[0, 2, 1, 1] + I[0, 2, 1, 2] - 2I[0, 2, 2, 0] + 2I[0, 2, 2, 1] + I[0, 2, 3, 0] \right. \right. \\
& - 2I[1, 1, 1, 0] - 4I[1, 1, 1, 1] + 2I[1, 1, 1, 2] - 4I[1, 1, 2, 0] + 4I[1, 1, 2, 1] + 2I[1, 1, 3, 0]) \\
& + m_c (9I[0, 2, 1, 0] - 18I[0, 2, 1, 1] + 9I[0, 2, 1, 2] - 14I[0, 2, 2, 0] + 14I[0, 2, 2, 1] + 5I[0, 2, 3, 0] \\
& - 10I[1, 1, 1, 0] - 28I[1, 1, 1, 1] + 18I[1, 1, 1, 2] - 20I[1, 1, 2, 0] + 28I[1, 1, 2, 1] + 10I[1, 1, 3, 0]) \left. \right) \\
& + 4m_c \left(5I[0, 3, 1, 0] - 19I[0, 3, 1, 1] + 21I[0, 3, 1, 2] - 7I[0, 3, 1, 3] - 10I[0, 3, 2, 0] + 24I[0, 3, 2, 1] \right. \\
& - 12I[0, 3, 2, 2] + 5I[0, 3, 3, 0] - 5I[0, 3, 3, 1] + 15I[1, 2, 1, 1] - 36I[1, 2, 1, 2] + 21I[1, 2, 1, 3] \\
& - 30I[1, 2, 2, 1] + 36I[1, 2, 2, 2] + 15I[1, 2, 3, 1] \left. \right) + 8m_s \left(I[0, 3, 1, 0] - 3I[0, 3, 1, 1] + 3I[0, 3, 1, 2] - I[0, 3, 1, 3] \right. \\
& - 2I[0, 3, 2, 0] + 4I[0, 3, 2, 1] - 2I[0, 3, 2, 2] + I[0, 3, 3, 0] - I[0, 3, 3, 1] + 3I[1, 2, 1, 1] - 6I[1, 2, 1, 2] + 3I[1, 2, 1, 3] \\
& \left. \left. - 6I[1, 2, 2, 1] + 6I[1, 2, 2, 2] + 3I[1, 2, 3, 1] \right) \right\} \\
& + \frac{m_c P_1 P_3}{679477248 \pi^5} \left[2e_c \left\{ m_0^2 \left(6m_c (5I[0, 2, 1, 0] - 4I[0, 2, 1, 1] - I[0, 2, 1, 2] - 5I[0, 2, 2, 0] - I[0, 2, 2, 1] - 2I[1, 1, 1, 1] \right. \right. \right. \\
& + 2I[1, 1, 1, 2] + 2I[1, 1, 2, 1]) + m_s (I[0, 2, 1, 0] - 2I[0, 2, 1, 1] + I[0, 2, 1, 2] - 2I[0, 2, 2, 0] + 2I[0, 2, 2, 1] + I[0, 2, 3, 0] \\
& - 2I[1, 1, 1, 0] - 4I[1, 1, 1, 1] + 2I[1, 1, 1, 2] - 4I[1, 1, 2, 0] + 4I[1, 1, 2, 1] + 2I[1, 1, 3, 0]) \left. \right) + 2m_c \left(3I[0, 3, 1, 0] \right. \\
& - 26I[0, 3, 1, 1] + 21I[0, 3, 1, 2] + 2I[0, 3, 1, 3] - 6I[0, 3, 2, 0] + 26I[0, 3, 2, 1] + 2I[0, 3, 2, 2] + 3I[0, 3, 3, 0] + 6I[1, 2, 1, 2] \\
& - 6I[1, 2, 1, 3] - 6I[1, 2, 2, 2] \left. \right) + 2m_s \left(31I[0, 3, 1, 0] - 63I[0, 3, 1, 1] + 33I[0, 3, 1, 2] - I[0, 3, 1, 3] - 62I[0, 3, 2, 0] \right. \\
& + 64I[0, 3, 2, 1] - 2I[0, 3, 2, 2] + 31I[0, 3, 3, 0] - I[0, 3, 3, 1] + 3I[1, 2, 1, 1] \\
& \left. \left. - 6I[1, 2, 1, 2] + 3I[1, 2, 1, 3] - 6I[1, 2, 2, 1] + 6I[1, 2, 2, 2] + 3I[1, 2, 3, 1] \right) \right\} \\
\end{aligned}$$

$$\begin{aligned}
& - (e_d + e_u) \left\{ m_0^2 \left(-2m_s (I[0, 2, 1, 0] - 2I[0, 2, 1, 1] + I[0, 2, 1, 2] - 2I[0, 2, 2, 0] + 2I[0, 2, 2, 1] + I[0, 2, 3, 0] \right. \right. \\
& - 2I[1, 1, 1, 0] - 4I[1, 1, 1, 1] + 2I[1, 1, 1, 2] - 4I[1, 1, 2, 0] + 4I[1, 1, 2, 1] + 2I[1, 1, 3, 0]) + 3m_c (5I[0, 2, 1, 0] \\
& - 10I[0, 2, 1, 1] + 5I[0, 2, 1, 2] - 6I[0, 2, 2, 0] + 6I[0, 2, 2, 1] + I[0, 2, 3, 0] - 2(I[1, 1, 1, 0] - 6I[1, 1, 1, 1] + 5I[1, 1, 1, 2] \\
& - 2I[1, 1, 2, 0] + 6I[1, 1, 2, 1] + I[1, 1, 3, 0])) + m_s \left(-I[0, 3, 1, 0] + 3I[0, 3, 1, 1] - 3I[0, 3, 1, 2] + I[0, 3, 1, 3] \right. \\
& - I[0, 3, 2, 0] + 2I[0, 3, 2, 1] - I[0, 3, 2, 2] + 5I[0, 3, 3, 0] - 5I[0, 3, 3, 1] - 3I[0, 3, 4, 0] - 15I[1, 2, 1, 1] + 3I[1, 2, 1, 2] \\
& + 3I[1, 2, 1, 3] + 30I[1, 2, 2, 1] - 3I[1, 2, 2, 2] - 15I[1, 2, 3, 1]) + 4m_c \left(I[0, 3, 1, 0] - 7I[0, 3, 1, 1] + 9I[0, 3, 1, 2] \right. \\
& - 3I[0, 3, 1, 3] - 2I[0, 3, 2, 0] + 8I[0, 3, 2, 1] - 4I[0, 3, 2, 2] + I[0, 3, 3, 0] - I[0, 3, 3, 1] + 3I[1, 2, 1, 1] - 12I[1, 2, 1, 2] \\
& \left. \left. + 9I[1, 2, 1, 3] - 6I[1, 2, 2, 1] + 12I[1, 2, 2, 2] + 3I[1, 2, 3, 1] \right) \right\} \\
& - \frac{5e_c m_c P_2 P_3}{2359296\pi^3} \left[4m_c m_s \left(I[0, 3, 1, 0] - 2I[0, 3, 1, 1] + I[0, 3, 1, 2] - 2I[0, 3, 2, 0] + 2I[0, 3, 2, 1] + I[0, 3, 3, 0] \right) \right. \\
& + 3 \left(-I[0, 4, 1, 0] + 3I[0, 4, 1, 1] - 3I[0, 4, 1, 2] + I[0, 4, 1, 3] + 3I[0, 4, 2, 0] - 6I[0, 4, 2, 1] + 3I[0, 4, 2, 2] - 3I[0, 4, 3, 0] \right. \\
& \left. \left. + 3I[0, 4, 3, 1] + I[0, 4, 4, 0] \right) \right] \\
& + \frac{m_c P_1}{27179089920\pi^7} \left[-e_c \left(-108I[0, 5, 1, 1] - 498I[0, 5, 1, 2] + 673I[0, 5, 1, 3] - 284I[0, 5, 1, 4] + I[0, 5, 1, 5] \right. \right. \\
& - 324I[0, 5, 2, 1] + 996I[0, 5, 2, 2] - 674I[0, 5, 2, 3] + 2I[0, 5, 2, 4] + 324I[0, 5, 3, 1] - 498I[0, 5, 3, 2] + I[0, 5, 3, 3] \\
& - 108I[0, 5, 4, 1] + 10m_c m_s (9I[0, 4, 1, 1] - 48I[0, 4, 1, 2] + 37I[0, 4, 1, 3] + 2I[0, 4, 1, 4] - 18I[0, 4, 2, 1] + 48I[0, 4, 2, 2] \\
& + 2I[0, 4, 2, 3] + 9I[0, 4, 3, 1] + 8I[1, 3, 1, 3] - 8I[1, 3, 1, 4] - 8I[1, 3, 2, 3]) - 5(I[1, 4, 1, 3] - 2I[1, 4, 1, 4] + I[1, 4, 1, 5] \\
& - 2I[1, 4, 2, 3] + 2I[1, 4, 2, 4] + I[1, 4, 3, 3]) \left. \right) \\
& + e_s \left(90I[0, 5, 1, 1] - 315I[0, 5, 1, 2] + 419I[0, 5, 1, 3] - 235I[0, 5, 1, 4] + 47I[0, 5, 1, 5] - 270I[0, 5, 2, 1] \right. \\
& + 687I[0, 5, 2, 2] - 556I[0, 5, 2, 3] + 139I[0, 5, 2, 4] + 270I[0, 5, 3, 1] - 411I[0, 5, 3, 2] \\
& + 137I[0, 5, 3, 3] - 5(18I[0, 5, 4, 1] - 9I[0, 5, 4, 2] - 45I[1, 4, 1, 2] + 137I[1, 4, 1, 3] - 139I[1, 4, 1, 4] + 47I[1, 4, 1, 5] \\
& \left. \left. + 135I[1, 4, 2, 2] - 274I[1, 4, 2, 3] + 139I[1, 4, 2, 4] - 135I[1, 4, 3, 2] + 137I[1, 4, 3, 3] - 45I[1, 4, 4, 2]) \right) \right) \\
& + e_d \left(54I[0, 5, 1, 1] - 183I[0, 5, 1, 2] + 229I[0, 5, 1, 3] - 125I[0, 5, 1, 4] + 25I[0, 5, 1, 5] - 162I[0, 5, 2, 1] + 393I[0, 5, 2, 2] \right. \\
& - 308I[0, 5, 2, 3] + 77I[0, 5, 2, 4] + 162I[0, 5, 3, 1] - 237I[0, 5, 3, 2] + 79I[0, 5, 3, 3] - 54I[0, 5, 4, 1] + 27I[0, 5, 4, 2] \\
& + 5m_c m_s (6I[0, 4, 1, 1] - 27I[0, 4, 1, 2] + 28I[0, 4, 1, 3] - 7I[0, 4, 1, 4] - 12I[0, 4, 2, 1] + 30I[0, 4, 2, 2] - 10I[0, 4, 2, 3] \\
& + 6I[0, 4, 3, 1] - 3I[0, 4, 3, 2] + 12I[1, 3, 1, 2] - 40I[1, 3, 1, 3] + 28I[1, 3, 1, 4] - 24I[1, 3, 2, 2] + 40I[1, 3, 2, 3] \\
& + 12I[1, 3, 3, 2]) + 5(27I[1, 4, 1, 2] - 79I[1, 4, 1, 3] + 77I[1, 4, 1, 4] - 25I[1, 4, 1, 5] - 81I[1, 4, 2, 2] + 158I[1, 4, 2, 3] \\
& \left. \left. - 77I[1, 4, 2, 4] + 81I[1, 4, 3, 2] - 79I[1, 4, 3, 3] - 27I[1, 4, 4, 2]) \right) \right) \\
& + e_u \left(54I[0, 5, 1, 1] - (189 - 6I)I[0, 5, 1, 2] + 229I[0, 5, 1, 3] - 125I[0, 5, 1, 4] + 25I[0, 5, 1, 5] - 162I[0, 5, 2, 1] \right. \\
& + 393I[0, 5, 2, 2] - 308I[0, 5, 2, 3] + 77I[0, 5, 2, 4] + 162I[0, 5, 3, 1] - 237I[0, 5, 3, 2] + 79I[0, 5, 3, 3] - 54I[0, 5, 4, 1] \\
& + 27I[0, 5, 4, 2] + 5m_c m_s (6I[0, 4, 1, 1] - 27I[0, 4, 1, 2] + 28I[0, 4, 1, 3] - 7I[0, 4, 1, 4] - 12I[0, 4, 2, 1] + 30I[0, 4, 2, 2] \\
& - 10I[0, 4, 2, 3] + 6I[0, 4, 3, 1] - 3I[0, 4, 3, 2] + 12I[1, 3, 1, 2] - 40I[1, 3, 1, 3] + 28I[1, 3, 1, 4] - 24I[1, 3, 2, 2] \\
& + 40I[1, 3, 2, 3] + 12I[1, 3, 3, 2]) + 5(27I[1, 4, 1, 2] - 79I[1, 4, 1, 3] + 77I[1, 4, 1, 4] - 25I[1, 4, 1, 5] - 81I[1, 4, 2, 2] \\
& \left. \left. + 158I[1, 4, 2, 3] - 77I[1, 4, 2, 4] + 81I[1, 4, 3, 2] - 79I[1, 4, 3, 3] + 27I[1, 4, 4, 2]) \right) \right) \right]
\end{aligned}$$

$$\begin{aligned}
& - \frac{e_c m_c P_2}{37748736\pi^5} \left[2m_c \left(15m_0^2 \left(I[0, 4, 1, 1] - 2I[0, 4, 1, 2] + I[0, 4, 1, 3] - 2I[0, 4, 2, 1] + 2I[0, 4, 2, 2] + I[0, 4, 3, 1] \right) \right. \right. \\
& + 6 \left(I[0, 5, 1, 2] - 2I[0, 5, 1, 3] + I[0, 5, 1, 4] - 2I[0, 5, 2, 2] + 2I[0, 5, 2, 3] + I[0, 5, 3, 2] \right) \left. \right) - 3 \left(15m_0^2 m_s \left(I[0, 4, 1, 0] \right. \right. \\
& - 3I[0, 4, 1, 1] + 3I[0, 4, 1, 2] - I[0, 4, 1, 3] - 3 \left(I[0, 4, 2, 0] - 2I[0, 4, 2, 1] + I[0, 4, 2, 2] - I[0, 4, 3, 0] + I[0, 4, 3, 1] \right) \\
& - I[0, 4, 4, 0] \left. \right) + 12m_s \left(-I[0, 5, 1, 1] + 3I[0, 5, 1, 2] - 3I[0, 5, 1, 3] + I[0, 5, 1, 4] + 3 \left(I[0, 5, 2, 1] - 2I[0, 5, 2, 2] \right. \right. \\
& \left. \left. + I[0, 5, 2, 3] - I[0, 5, 3, 1] + I[0, 5, 3, 2] \right) + I[0, 5, 4, 1] \right) \left. \right) \left. \right] \\
& - \frac{e_c m_c P_3}{125829120\pi^5} \left[5m_0^2 \left(2m_c \left(I[0, 4, 1, 1] - 2I[0, 4, 1, 2] + I[0, 4, 1, 3] - 2I[0, 4, 2, 1] + 2I[0, 4, 2, 2] + I[0, 4, 3, 1] \right) \right. \right. \\
& + 11m_s \left(I[0, 4, 1, 0] - 3I[0, 4, 1, 1] + 3I[0, 4, 1, 2] - I[0, 4, 1, 3] - 3 \left(I[0, 4, 2, 0] - 2I[0, 4, 2, 1] + I[0, 4, 2, 2] - I[0, 4, 3, 0] \right. \right. \\
& \left. \left. + I[0, 4, 3, 1] \right) - I[0, 4, 4, 0] \right) \left. \right) - 4m_c \left(I[0, 5, 1, 2] - 2I[0, 5, 1, 3] + I[0, 5, 1, 4] - 2I[0, 5, 2, 2] + 2I[0, 5, 2, 3] \right. \\
& \left. \left. + I[0, 5, 3, 2] \right) - 66m_s \left(I[0, 5, 1, 1] - 3I[0, 5, 1, 2] + 3I[0, 5, 1, 3] - I[0, 5, 1, 4] - 3 \left(I[0, 5, 2, 1] - 2I[0, 5, 2, 2] + I[0, 5, 2, 3] \right. \right. \\
& \left. \left. - I[0, 5, 3, 1] + I[0, 5, 3, 2] \right) - I[0, 5, 4, 1] \right) \left. \right] \\
& - \frac{e_c m_c^2 m_s}{754974720\pi^7} \left[I[0, 6, 1, 3] - 2I[0, 6, 1, 4] + I[0, 6, 1, 5] - 2I[0, 6, 2, 3] + 2I[0, 6, 2, 4] + I[0, 6, 3, 3] \right] \\
& + \frac{11e_c m_c}{3523215360\pi^7} \left[I[0, 7, 1, 3] - 3I[0, 7, 1, 4] + 3I[0, 7, 1, 5] - I[0, 7, 1, 6] - 3I[0, 7, 2, 3] + 6I[0, 7, 2, 4] - 3I[0, 7, 2, 5] \right. \\
& \left. + 3I[0, 7, 3, 3] - 3I[0, 7, 3, 4] - I[0, 7, 4, 3] \right] \\
& + \frac{f_{3\gamma} m_c P_1}{86973087744\pi^5} \left[64 \left(ed \left(-4m_c m_s \left(I[0, 3, 1, 0] - 7I[0, 3, 1, 1] + 9I[0, 3, 1, 2] - 3I[0, 3, 1, 3] - 2I[0, 3, 2, 0] \right. \right. \right. \right. \\
& + 8I[0, 3, 2, 1] - 4I[0, 3, 2, 2] + I[0, 3, 3, 0] - I[0, 3, 3, 1] \left. \right) + 9I[0, 4, 1, 0] - 38I[0, 4, 1, 1] + 59I[0, 4, 1, 2] \\
& - 40I[0, 4, 1, 3] + 10I[0, 4, 1, 4] - 27I[0, 4, 2, 0] + 85I[0, 4, 2, 1] - 87I[0, 4, 2, 2] + 29I[0, 4, 2, 3] + 27I[0, 4, 3, 0] \\
& - 56I[0, 4, 3, 1] + 28I[0, 4, 3, 2] - 9I[0, 4, 4, 0] + 9I[0, 4, 4, 1] \left. \right) + e_u \left(-4m_c m_s \left(I[0, 3, 1, 0] - 7I[0, 3, 1, 1] + 9I[0, 3, 1, 2] \right. \right. \\
& - 3I[0, 3, 1, 3] - 2I[0, 3, 2, 0] + 8I[0, 3, 2, 1] - 4I[0, 3, 2, 2] + I[0, 3, 3, 0] - I[0, 3, 3, 1] \left. \right) + 9I[0, 4, 1, 0] - 38I[0, 4, 1, 1] \\
& + 59I[0, 4, 1, 2] - 40I[0, 4, 1, 3] + 10I[0, 4, 1, 4] - 27I[0, 4, 2, 0] + 85I[0, 4, 2, 1] - 87I[0, 4, 2, 2] + 29I[0, 4, 2, 3] \\
& + 27I[0, 4, 3, 0] - 56I[0, 4, 3, 1] + 28I[0, 4, 3, 2] - 9I[0, 4, 4, 0] + 9I[0, 4, 4, 1] \left. \right) - e_s \left(21I[0, 4, 1, 0] - 82I[0, 4, 1, 1] \right. \\
& + 121I[0, 4, 1, 2] - 80I[0, 4, 1, 3] + 20I[0, 4, 1, 4] - 63I[0, 4, 2, 0] + 185I[0, 4, 2, 1] - 183I[0, 4, 2, 2] + 61I[0, 4, 2, 3] \\
& + 63I[0, 4, 3, 0] - 124I[0, 4, 3, 1] + 62I[0, 4, 3, 2] - 21I[0, 4, 4, 0] + 21I[0, 4, 4, 1] \left. \right) \right) \varphi^a[u_0] \\
& + I_2[\mathcal{V}] \left(-45(11e_d + 24e_s)I[0, 4, 3, 0] + e_u \left(736m_c m_s I[0, 3, 2, 0] - 459I[0, 4, 3, 0] - 30I[0, 4, 4, 0] \right) \right. \\
& \left. - 60e_s I[0, 4, 4, 0] \right) \left. \right]
\end{aligned}$$

$$\begin{aligned}
& - \frac{m_c P_2}{377487360\pi^5} \left[10f_{3\gamma}\pi^2 I_2[\mathcal{V}] \left(20e_s m_c (m_0^2 I[0, 3, 3, 0] - I[0, 4, 3, 0]) + e_u (10m_0^2 m_c I[0, 3, 3, 0] - 10m_c I[0, 4, 3, 0] \right. \right. \\
& - 3m_s I[0, 4, 4, 0]) + 2e_d (5m_0^2 m_c I[0, 3, 3, 0] - 5m_c I[0, 4, 3, 0] + 9m_s I[0, 4, 4, 0]) \left. \left. \right) + 3I_4[\mathcal{S}] \left(5e_d m_c I[0, 5, 3, 0] \right. \right. \\
& - e_u m_c I[0, 5, 3, 0] + 21e_d m_s I[0, 5, 4, 0] + 21e_u m_s I[0, 5, 4, 0] \left. \left. \right) \right] \\
& + \frac{m_c P_3}{377487360\pi^5} \left[5f_{3\gamma}\pi^2 I_2[\mathcal{V}] \left(33e_d m_s I[0, 4, 4, 0] + e_u (4m_0^2 m_c I[0, 3, 3, 0] - 4m_c I[0, 4, 3, 0] + 15m_s I[0, 4, 4, 0]) \right) \right. \\
& \left. - 3e_s m_c I_4[\mathcal{S}] I[0, 5, 3, 0] \right] \\
& \left. \frac{f_{3\gamma} m_c}{1006632960\pi^5} \left[I_2[\mathcal{V}] \left(11(e_d + e_s) I[0, 6, 4, 0] + e_u (4m_c m_s I[0, 5, 3, 0] + 5I[0, 6, 4, 0]) \right) \right] \right], \tag{A1}
\end{aligned}$$

where $P_1 = \langle g_s^2 G^2 \rangle$ is gluon condensate, $P_2 = \langle \bar{q}q \rangle$ stands for u/d quark condensate and $P_3 = \langle \bar{s}s \rangle$ denotes s quark condensate. We should also remark that in the Eq. (A1), for simplicity we have only given the terms that give significant contributions to the numerical values of the magnetic moment and neglected to give many higher dimensional operators though they have been considered in the numerical calculations. The $I[n, m, l, k]$, $I_1[\mathcal{F}]$, $I_2[\mathcal{F}]$, $I_3[\mathcal{F}]$, $I_4[\mathcal{F}]$, $I_5[\mathcal{F}]$, and $I_6[\mathcal{F}]$ functions are defined as:

$$\begin{aligned}
I[n, m, l, k] &= \int_{4m_Q^2}^{s_0} ds \int_0^1 dt \int_0^1 dw e^{-s/M^2} s^n (s - 4m_Q^2)^m t^l w^k, \\
I_1[\mathcal{F}] &= \int D\alpha_i \int_0^1 dv \mathcal{A}(\alpha_{\bar{q}}, \alpha_q, \alpha_g) \delta'(\alpha_q + \bar{v}\alpha_g - u_0), \\
I_2[\mathcal{F}] &= \int D\alpha_i \int_0^1 dv \mathcal{A}(\alpha_{\bar{q}}, \alpha_q, \alpha_g) \delta'(\alpha_{\bar{q}} + v\alpha_g - u_0), \\
I_3[\mathcal{F}] &= \int D\alpha_i \int_0^1 dv \mathcal{A}(\alpha_{\bar{q}}, \alpha_q, \alpha_g) \delta(\alpha_q + \bar{v}\alpha_g - u_0), \\
I_4[\mathcal{F}] &= \int D\alpha_i \int_0^1 dv \mathcal{A}(\alpha_{\bar{q}}, \alpha_q, \alpha_g) \delta(\alpha_{\bar{q}} + v\alpha_g - u_0), \\
I_5[\mathcal{F}] &= \int_0^1 du A(u) \delta'(u - u_0), \\
I_6[\mathcal{F}] &= \int_0^1 du A(u),
\end{aligned}$$

where \mathcal{F} denotes the corresponding photon DAs.

Appendix B: Distribution amplitudes of the photon

In the present appendix, the matrix elements $\langle \gamma(q) | \bar{q}(x) \Gamma_i q(0) | 0 \rangle$ and $\langle \gamma(q) | \bar{q}(x) \Gamma_i G_{\mu\nu} q(0) | 0 \rangle$ associated with the photon DAs are presented as follows [45],

$$\begin{aligned}
\langle \gamma(q) | \bar{q}(x) \gamma_\mu q(0) | 0 \rangle &= e_q f_{3\gamma} \left(\varepsilon_\mu - q_\mu \frac{\varepsilon x}{qx} \right) \int_0^1 du e^{i\bar{u}qx} \psi^v(u) \\
\langle \gamma(q) | \bar{q}(x) \gamma_\mu \gamma_5 q(0) | 0 \rangle &= -\frac{1}{4} e_q f_{3\gamma} \epsilon_{\mu\nu\alpha\beta} \varepsilon^\nu q^\alpha x^\beta \int_0^1 du e^{i\bar{u}qx} \psi^a(u) \\
\langle \gamma(q) | \bar{q}(x) \sigma_{\mu\nu} q(0) | 0 \rangle &= -ie_q \langle \bar{q}q \rangle (\varepsilon_\mu q_\nu - \varepsilon_\nu q_\mu) \int_0^1 du e^{i\bar{u}qx} \left(\chi \varphi_\gamma(u) + \frac{x^2}{16} \mathbb{A}(u) \right) \\
&\quad - \frac{i}{2(qx)} e_q \bar{q}q \left[x_\nu \left(\varepsilon_\mu - q_\mu \frac{\varepsilon x}{qx} \right) - x_\mu \left(\varepsilon_\nu - q_\nu \frac{\varepsilon x}{qx} \right) \right] \int_0^1 du e^{i\bar{u}qx} h_\gamma(u) \\
\langle \gamma(q) | \bar{q}(x) g_s G_{\mu\nu}(vx) q(0) | 0 \rangle &= -ie_q \langle \bar{q}q \rangle (\varepsilon_\mu q_\nu - \varepsilon_\nu q_\mu) \int \mathcal{D}\alpha_i e^{i(\alpha_{\bar{q}} + v\alpha_g)qx} \mathcal{S}(\alpha_i) \\
\langle \gamma(q) | \bar{q}(x) g_s \tilde{G}_{\mu\nu}(vx) i\gamma_5 q(0) | 0 \rangle &= -ie_q \langle \bar{q}q \rangle (\varepsilon_\mu q_\nu - \varepsilon_\nu q_\mu) \int \mathcal{D}\alpha_i e^{i(\alpha_{\bar{q}} + v\alpha_g)qx} \tilde{\mathcal{S}}(\alpha_i) \\
\langle \gamma(q) | \bar{q}(x) g_s \tilde{G}_{\mu\nu}(vx) \gamma_\alpha \gamma_5 q(0) | 0 \rangle &= e_q f_{3\gamma} q_\alpha (\varepsilon_\mu q_\nu - \varepsilon_\nu q_\mu) \int \mathcal{D}\alpha_i e^{i(\alpha_{\bar{q}} + v\alpha_g)qx} \mathcal{A}(\alpha_i) \\
\langle \gamma(q) | \bar{q}(x) g_s G_{\mu\nu}(vx) i\gamma_\alpha q(0) | 0 \rangle &= e_q f_{3\gamma} q_\alpha (\varepsilon_\mu q_\nu - \varepsilon_\nu q_\mu) \int \mathcal{D}\alpha_i e^{i(\alpha_{\bar{q}} + v\alpha_g)qx} \mathcal{V}(\alpha_i) \\
\langle \gamma(q) | \bar{q}(x) \sigma_{\alpha\beta} g_s G_{\mu\nu}(vx) q(0) | 0 \rangle &= e_q \langle \bar{q}q \rangle \left\{ \left[\left(\varepsilon_\mu - q_\mu \frac{\varepsilon x}{qx} \right) \left(g_{\alpha\nu} - \frac{1}{qx} (q_\alpha x_\nu + q_\nu x_\alpha) \right) q_\beta \right. \right. \\
&\quad - \left(\varepsilon_\mu - q_\mu \frac{\varepsilon x}{qx} \right) \left(g_{\beta\nu} - \frac{1}{qx} (q_\beta x_\nu + q_\nu x_\beta) \right) q_\alpha - \left(\varepsilon_\nu - q_\nu \frac{\varepsilon x}{qx} \right) \left(g_{\alpha\mu} - \frac{1}{qx} (q_\alpha x_\mu + q_\mu x_\alpha) \right) q_\beta \\
&\quad + \left. \left(\varepsilon_\nu - q_\nu \frac{\varepsilon x}{qx} \right) \left(g_{\beta\mu} - \frac{1}{qx} (q_\beta x_\mu + q_\mu x_\beta) \right) q_\alpha \right] \int \mathcal{D}\alpha_i e^{i(\alpha_{\bar{q}} + v\alpha_g)qx} \mathcal{T}_1(\alpha_i) \\
&\quad + \left[\left(\varepsilon_\alpha - q_\alpha \frac{\varepsilon x}{qx} \right) \left(g_{\mu\beta} - \frac{1}{qx} (q_\mu x_\beta + q_\beta x_\mu) \right) q_\nu \right. \\
&\quad - \left(\varepsilon_\alpha - q_\alpha \frac{\varepsilon x}{qx} \right) \left(g_{\nu\beta} - \frac{1}{qx} (q_\nu x_\beta + q_\beta x_\nu) \right) q_\mu \\
&\quad - \left. \left(\varepsilon_\beta - q_\beta \frac{\varepsilon x}{qx} \right) \left(g_{\mu\alpha} - \frac{1}{qx} (q_\mu x_\alpha + q_\alpha x_\mu) \right) q_\nu \right. \\
&\quad + \left. \left(\varepsilon_\beta - q_\beta \frac{\varepsilon x}{qx} \right) \left(g_{\nu\alpha} - \frac{1}{qx} (q_\nu x_\alpha + q_\alpha x_\nu) \right) q_\mu \right] \int \mathcal{D}\alpha_i e^{i(\alpha_{\bar{q}} + v\alpha_g)qx} \mathcal{T}_2(\alpha_i) \\
&\quad + \frac{1}{qx} (q_\mu x_\nu - q_\nu x_\mu) (\varepsilon_\alpha q_\beta - \varepsilon_\beta q_\alpha) \int \mathcal{D}\alpha_i e^{i(\alpha_{\bar{q}} + v\alpha_g)qx} \mathcal{T}_3(\alpha_i) \\
&\quad + \left. \frac{1}{qx} (q_\alpha x_\beta - q_\beta x_\alpha) (\varepsilon_\mu q_\nu - \varepsilon_\nu q_\mu) \int \mathcal{D}\alpha_i e^{i(\alpha_{\bar{q}} + v\alpha_g)qx} \mathcal{T}_4(\alpha_i) \right\} ,
\end{aligned}$$

where $\varphi_\gamma(u)$ is the DA of leading twist-2, $\psi^v(u)$, $\psi^a(u)$, $\mathcal{A}(\alpha_i)$ and $\mathcal{V}(\alpha_i)$, are the twist-3 amplitudes, and $h_\gamma(u)$, $\mathbb{A}(u)$, $\mathcal{S}(\alpha_i)$, $\tilde{\mathcal{S}}(\alpha_i)$, $\mathcal{T}_1(\alpha_i)$, $\mathcal{T}_2(\alpha_i)$, $\mathcal{T}_3(\alpha_i)$ and $\mathcal{T}_4(\alpha_i)$ are the twist-4 photon DAs. The measure $\mathcal{D}\alpha_i$ is defined as

$$\int \mathcal{D}\alpha_i = \int_0^1 d\alpha_{\bar{q}} \int_0^1 d\alpha_q \int_0^1 d\alpha_g \delta(1 - \alpha_{\bar{q}} - \alpha_q - \alpha_g) .$$

The expressions of the DAs that entering into the matrix elements above are described as follows:

$$\begin{aligned}
\varphi_\gamma(u) &= 6u\bar{u} \left(1 + \varphi_2(\mu) C_2^{\frac{3}{2}}(u - \bar{u}) \right), \\
\psi^v(u) &= 3 \left(3(2u - 1)^2 - 1 \right) + \frac{3}{64} (15w_\gamma^V - 5w_\gamma^A) (3 - 30(2u - 1)^2 + 35(2u - 1)^4), \\
\psi^a(u) &= (1 - (2u - 1)^2) (5(2u - 1)^2 - 1) \frac{5}{2} \left(1 + \frac{9}{16} w_\gamma^V - \frac{3}{16} w_\gamma^A \right), \\
h_\gamma(u) &= -10 (1 + 2\kappa^+) C_2^{\frac{1}{2}}(u - \bar{u}), \\
\mathbb{A}(u) &= 40u^2\bar{u}^2 (3\kappa - \kappa^+ + 1) + 8(\zeta_2^+ - 3\zeta_2) [u\bar{u}(2 + 13u\bar{u}) \\
&\quad + 2u^3(10 - 15u + 6u^2) \ln(u) + 2\bar{u}^3(10 - 15\bar{u} + 6\bar{u}^2) \ln(\bar{u})], \\
\mathcal{A}(\alpha_i) &= 360\alpha_q\alpha_{\bar{q}}\alpha_g^2 \left(1 + w_\gamma^A \frac{1}{2} (7\alpha_g - 3) \right), \\
\mathcal{V}(\alpha_i) &= 540w_\gamma^V (\alpha_q - \alpha_{\bar{q}}) \alpha_q \alpha_{\bar{q}} \alpha_g^2, \\
\mathcal{T}_1(\alpha_i) &= -120(3\zeta_2 + \zeta_2^+) (\alpha_{\bar{q}} - \alpha_q) \alpha_{\bar{q}} \alpha_q \alpha_g, \\
\mathcal{T}_2(\alpha_i) &= 30\alpha_g^2 (\alpha_{\bar{q}} - \alpha_q) ((\kappa - \kappa^+) + (\zeta_1 - \zeta_1^+) (1 - 2\alpha_g) + \zeta_2(3 - 4\alpha_g)), \\
\mathcal{T}_3(\alpha_i) &= -120(3\zeta_2 - \zeta_2^+) (\alpha_{\bar{q}} - \alpha_q) \alpha_{\bar{q}} \alpha_q \alpha_g, \\
\mathcal{T}_4(\alpha_i) &= 30\alpha_g^2 (\alpha_{\bar{q}} - \alpha_q) ((\kappa + \kappa^+) + (\zeta_1 + \zeta_1^+) (1 - 2\alpha_g) + \zeta_2(3 - 4\alpha_g)), \\
\mathcal{S}(\alpha_i) &= 30\alpha_g^2 \{ (\kappa + \kappa^+) (1 - \alpha_g) + (\zeta_1 + \zeta_1^+) (1 - \alpha_g) (1 - 2\alpha_g) + \zeta_2 [3(\alpha_{\bar{q}} - \alpha_q)^2 - \alpha_g(1 - \alpha_g)] \}, \\
\tilde{\mathcal{S}}(\alpha_i) &= -30\alpha_g^2 \{ (\kappa - \kappa^+) (1 - \alpha_g) + (\zeta_1 - \zeta_1^+) (1 - \alpha_g) (1 - 2\alpha_g) + \zeta_2 [3(\alpha_{\bar{q}} - \alpha_q)^2 - \alpha_g(1 - \alpha_g)] \}.
\end{aligned}$$

Numerical values of parameters used in DAs are: $\varphi_2(1 \text{ GeV}) = 0$, $w_\gamma^V = 3.8 \pm 1.8$, $w_\gamma^A = -2.1 \pm 1.0$, $\kappa = 0.2$, $\kappa^+ = 0$, $\zeta_1 = 0.4$, $\zeta_2 = 0.3$.

-
- [1] A. Esposito, A. L. Guerrieri, F. Piccinini, A. Pilloni, A. D. Polosa, Four-Quark Hadrons: an Updated Review, *Int. J. Mod. Phys. A* 30 (2015) 1530002. [arXiv:1411.5997](#), [doi:10.1142/S0217751X15300021](#).
- [2] A. Esposito, A. Pilloni, A. D. Polosa, Multiquark Resonances, *Phys. Rept.* 668 (2017) 1–97. [arXiv:1611.07920](#), [doi:10.1016/j.physrep.2016.11.002](#).
- [3] S. L. Olsen, T. Skwarnicki, D. Zieminska, Nonstandard heavy mesons and baryons: Experimental evidence, *Rev. Mod. Phys.* 90 (1) (2018) 015003. [arXiv:1708.04012](#), [doi:10.1103/RevModPhys.90.015003](#).
- [4] R. F. Lebed, R. E. Mitchell, E. S. Swanson, Heavy-Quark QCD Exotica, *Prog. Part. Nucl. Phys.* 93 (2017) 143–194. [arXiv:1610.04528](#), [doi:10.1016/j.pnnp.2016.11.003](#).
- [5] M. Nielsen, F. S. Navarra, S. H. Lee, New Charmonium States in QCD Sum Rules: A Concise Review, *Phys. Rept.* 497 (2010) 41–83. [arXiv:0911.1958](#), [doi:10.1016/j.physrep.2010.07.005](#).
- [6] N. Brambilla, S. Eidelman, C. Hanhart, A. Nefediev, C.-P. Shen, C. E. Thomas, A. Vairo, C.-Z. Yuan, The *XYZ* states: experimental and theoretical status and perspectives, *Phys. Rept.* 873 (2020) 1–154. [arXiv:1907.07583](#), [doi:10.1016/j.physrep.2020.05.001](#).
- [7] S. Agaev, K. Azizi, H. Sundu, Four-quark exotic mesons, *Turk. J. Phys.* 44 (2) (2020) 95–173. [arXiv:2004.12079](#), [doi:10.3906/fiz-2003-15](#).
- [8] H.-X. Chen, W. Chen, X. Liu, S.-L. Zhu, The hidden-charm pentaquark and tetraquark states, *Phys. Rept.* 639 (2016) 1–121. [arXiv:1601.02092](#), [doi:10.1016/j.physrep.2016.05.004](#).
- [9] A. Ali, J. S. Lange, S. Stone, Exotics: Heavy Pentaquarks and Tetraquarks, *Prog. Part. Nucl. Phys.* 97 (2017) 123–198. [arXiv:1706.00610](#), [doi:10.1016/j.pnnp.2017.08.003](#).
- [10] F.-K. Guo, C. Hanhart, U.-G. Meißner, Q. Wang, Q. Zhao, B.-S. Zou, Hadronic molecules, *Rev. Mod. Phys.* 90 (1) (2018) 015004, [Erratum: *Rev. Mod. Phys.* 94, 029901 (2022)]. [arXiv:1705.00141](#), [doi:10.1103/RevModPhys.90.015004](#).
- [11] Y.-R. Liu, H.-X. Chen, W. Chen, X. Liu, S.-L. Zhu, Pentaquark and Tetraquark states, *Prog. Part. Nucl. Phys.* 107 (2019) 237–320. [arXiv:1903.11976](#), [doi:10.1016/j.pnnp.2019.04.003](#).
- [12] G. Yang, J. Ping, J. Segovia, Tetra- and penta-quark structures in the constituent quark model, *Symmetry* 12 (11) (2020) 1869. [arXiv:2009.00238](#), [doi:10.3390/sym12111869](#).
- [13] X.-K. Dong, F.-K. Guo, B.-S. Zou, A survey of heavy-antiheavy hadronic molecules, *Progr. Phys.* 41 (2021) 65–93. [arXiv:2101.01021](#), [doi:10.13725/j.cnki.pip.2021.02.001](#).
- [14] X.-K. Dong, F.-K. Guo, B.-S. Zou, A survey of heavy-heavy hadronic molecules, *Commun. Theor. Phys.* 73 (12) (2021) 125201. [arXiv:2108.02673](#), [doi:10.1088/](#)

- 1572-9494/ac27a2.
- [15] L. Meng, B. Wang, G.-J. Wang, S.-L. Zhu, Chiral perturbation theory for heavy hadrons and chiral effective field theory for heavy hadronic molecules (4 2022). [arXiv:2204.08716](#).
- [16] H.-X. Chen, W. Chen, X. Liu, Y.-R. Liu, S.-L. Zhu, An updated review of the new hadron states (4 2022). [arXiv:2204.02649](#).
- [17] R. Aaij, et al., Observation of $J/\psi p$ Resonances Consistent with Pentaquark States in $\Lambda_b^0 \rightarrow J/\psi K^- p$ Decays, *Phys. Rev. Lett.* 115 (2015) 072001. [arXiv:1507.03414](#), [doi:10.1103/PhysRevLett.115.072001](#).
- [18] R. Aaij, et al., Observation of a narrow pentaquark state, $P_c(4312)^+$, and of two-peak structure of the $P_c(4450)^+$, *Phys. Rev. Lett.* 122 (22) (2019) 222001. [arXiv:1904.03947](#), [doi:10.1103/PhysRevLett.122.222001](#).
- [19] R. Aaij, et al., Evidence of a $J/\psi\Lambda$ structure and observation of excited Ξ^- states in the $\Xi_b^- \rightarrow J/\psi\Lambda K^-$ decay, *Sci. Bull.* 66 (2021) 1278–1287. [arXiv:2012.10380](#), [doi:10.1016/j.scib.2021.02.030](#).
- [20] R. Aaij, et al., Evidence for a new structure in the $J/\psi p$ and $J/\psi\bar{p}$ systems in $B_s^0 \rightarrow J/\psi p\bar{p}$ decays, *Phys. Rev. Lett.* 128 (6) (2022) 062001. [arXiv:2108.04720](#), [doi:10.1103/PhysRevLett.128.062001](#).
- [21] C.-W. Shen, D. Rönchen, U.-G. Meißner, B.-S. Zou, Exploratory study of possible resonances in heavy meson - heavy baryon coupled-channel interactions, *Chin. Phys. C* 42 (2) (2018) 023106. [arXiv:1710.03885](#), [doi:10.1088/1674-1137/42/2/023106](#).
- [22] Chen Chen and Elisabetta Spadaro on behalf of the LHCb Collaboration, New tetra- and pentaquarks at LHCb, LHC Seminar, <https://indico.cern.ch/event/1176505/>.
- [23] X.-W. Wang, Z.-G. Wang, Analysis of the $P_{cs}(4338)$ and related pentaquark molecular states via the QCD sum rules (7 2022). [arXiv:2207.06060](#).
- [24] M. Karliner, J. R. Rosner, New strange pentaquarks (7 2022). [arXiv:2207.07581](#).
- [25] F.-L. Wang, X. Liu, Emergence of molecular-type characteristic spectrum of hidden-charm pentaquark with strangeness embodied in the $P_{\psi_s}^\Lambda(4338)$ and $P_{cs}(4459)$ (7 2022). [arXiv:2207.10493](#).
- [26] M.-J. Yan, F.-Z. Peng, M. S. Sánchez, M. Pavon Valderama, The $P_{\psi_s}^\Lambda(4338)$ pentaquark and its partners in the molecular picture (7 2022). [arXiv:2207.11144](#).
- [27] L. Meng, B. Wang, S.-L. Zhu, The double thresholds distort the lineshapes of the $P_{\psi_s}^\Lambda(4338)^0$ resonance (8 2022). [arXiv:2208.03883](#).
- [28] T. J. Burns, E. S. Swanson, The LHCb state $P_{\psi_s}^\Lambda(4338)$ as a triangle singularity (8 2022). [arXiv:2208.05106](#).
- [29] G.-J. Wang, R. Chen, L. Ma, X. Liu, S.-L. Zhu, Magnetic moments of the hidden-charm pentaquark states, *Phys. Rev. D* 94 (9) (2016) 094018. [arXiv:1605.01337](#), [doi:10.1103/PhysRevD.94.094018](#).
- [30] U. Özdem, K. Azizi, Electromagnetic multipole moments of the $P_c^+(4380)$ pentaquark in light-cone QCD, *Eur. Phys. J. C* 78 (5) (2018) 379. [arXiv:1803.06831](#), [doi:10.1140/epjc/s10052-018-5873-2](#).
- [31] E. Ortiz-Pacheco, R. Bijker, C. Fernández-Ramírez, Hidden charm pentaquarks: mass spectrum, magnetic moments, and photocouplings, *J. Phys. G* 46 (6) (2019) 065104. [arXiv:1808.10512](#), [doi:10.1088/1361-6471/ab096d](#).
- [32] Y.-J. Xu, Y.-L. Liu, M.-Q. Huang, The magnetic moment of $P_c(4312)$ as a $\bar{D}\Sigma_c$ molecular state, *Eur. Phys. J. C* 81 (5) (2021) 421. [arXiv:2008.07937](#), [doi:10.1140/epjc/s10052-021-09211-8](#).
- [33] U. Özdem, Electromagnetic properties of the $P_c(4312)$ pentaquark state, *Chin. Phys. C* 45 (2) (2021) 023119. [doi:10.1088/1674-1137/abd01c](#).
- [34] U. Özdem, Magnetic dipole moments of the hidden-charm pentaquark states: $P_c(4440)$, $P_c(4457)$ and $P_{cs}(4459)$, *Eur. Phys. J. C* 81 (4) (2021) 277. [arXiv:2102.01996](#), [doi:10.1140/epjc/s10052-021-09070-3](#).
- [35] M.-W. Li, Z.-W. Liu, Z.-F. Sun, R. Chen, Magnetic moments and transition magnetic moments of P_c and P_{cs} states, *Phys. Rev. D* 104 (5) (2021) 054016. [arXiv:2106.15053](#), [doi:10.1103/PhysRevD.104.054016](#).
- [36] F. Gao, H.-S. Li, Magnetic moments of the hidden-charm strange pentaquark states (12 2021). [arXiv:2112.01823](#).
- [37] U. Özdem, Magnetic moments of the vector hidden-charm tetraquark states, *Phys. Rev. D* 105 (11) (2022) 114030. [arXiv:2206.05196](#), [doi:10.1103/PhysRevD.105.114030](#).
- [38] U. Özdem, Electromagnetic properties of doubly-heavy pentaquark states (1 2022). [arXiv:2201.00979](#).
- [39] H.-Y. Zhou, F.-L. Wang, Z.-W. Liu, X. Liu, Probing the electromagnetic properties of the $\Sigma_c^{(*)}D^{(*)}$ -type doubly charmed molecular pentaquarks (7 2022). [arXiv:2207.08660](#).
- [40] V. L. Chernyak, I. R. Zhitnitsky, B meson exclusive decays into baryons, *Nucl. Phys. B* 345 (1990) 137–172. [doi:10.1016/0550-3213\(90\)90612-H](#).
- [41] V. M. Braun, I. E. Filyanov, QCD Sum Rules in Exclusive Kinematics and Pion Wave Function, *Z. Phys. C* 44 (1989) 157. [doi:10.1007/BF01548594](#).
- [42] I. I. Balitsky, V. M. Braun, A. V. Kolesnichenko, Radiative Decay $\Sigma^+ \rightarrow p \gamma$ in Quantum Chromodynamics, *Nucl. Phys. B* 312 (1989) 509–550. [doi:10.1016/0550-3213\(89\)90570-1](#).
- [43] K.-C. Yang, W. Y. P. Hwang, E. M. Henley, L. S. Kisslinger, QCD sum rules and neutron proton mass difference, *Phys. Rev. D* 47 (1993) 3001–3012. [doi:10.1103/PhysRevD.47.3001](#).
- [44] V. M. Belyaev, B. Y. Blok, CHARMED BARYONS IN QUANTUM CHROMODYNAMICS, *Z. Phys. C* 30 (1986) 151. [doi:10.1007/BF01560689](#).
- [45] P. Ball, V. M. Braun, N. Kivel, Photon distribution amplitudes in QCD, *Nucl. Phys. B* 649 (2003) 263–296. [arXiv:hep-ph/0207307](#), [doi:10.1016/S0550-3213\(02\)01017-9](#).
- [46] H. J. Weber, H. Arenhovel, Isobar Configurations in Nuclei, *Phys. Rept.* 36 (1978) 277–348. [doi:10.1016/0370-1573\(78\)90187-4](#).
- [47] S. Nozawa, D. B. Leinweber, Electromagnetic formfactors of spin 3/2 baryons, *Phys. Rev. D* 42 (1990) 3567–3571. [doi:10.1103/PhysRevD.42.3567](#).
- [48] V. Pascalutsa, M. Vanderhaeghen, S. N. Yang, Electromagnetic excitation of the Delta(1232)-resonance, *Phys. Rept.* 437 (2007) 125–232. [arXiv:hep-ph/0609004](#), [doi:10.1016/j.physrep.2006.09.006](#).
- [49] G. Ramalho, M. T. Pena, F. Gross, Electric quadrupole and magnetic octupole moments of the Delta, *Phys. Lett. B* 678 (2009) 355–358. [arXiv:0902.4212](#), [doi:10.1016/j.physletb.2009.06.052](#).
- [50] V. M. Belyaev, B. L. Ioffe, Determination of the baryon mass and baryon resonances from the quantum-

- chromodynamics sum rule. Strange baryons, Sov. Phys. JETP 57 (1983) 716–721.
- [51] R. L. Workman, Review of Particle Physics, PTEP 2022 (2022) 083C01.
- [52] B. L. Ioffe, QCD at low energies, Prog. Part. Nucl. Phys. 56 (2006) 232–277. [arXiv:hep-ph/0502148](#), [doi:10.1016/j.ppnp.2005.05.001](#).