

Finite groups with some bounded codegrees

Dongfang Yang^a, Yu Zeng^b, Heng Lv^{a*}

^aSchool of Mathematics and Statistics, Southwest University, Chongqing, China

^bDepartment of Mathematics, Changshu Institute of Technology, Jiangsu, China

Abstract. For a character χ of a finite group G , the number $\text{cod}(\chi) := |G : \ker(\chi)|/\chi(1)$ is called the codegree of χ . In this paper, we give a solvability criterion for a finite group G depending on the minimum of the ratio $\chi(1)^2/\text{cod}(\chi)$, when χ varies among the irreducible characters of G .

Key words. Finite group, codegree, degree, simple group.

1 Introduction

Throughout all groups are finite and p is a prime number. Let G be a group. We denote by $\text{Irr}(G)$ the set of irreducible characters of G ; and use $\text{cd}(G)$ to denote the set of all irreducible character degrees of G . Let χ be a character of G . Qian, et al. [10] called the positive integer $\text{cod}(\chi) := |G : \ker(\chi)|/\chi(1)$ the codegree of the character χ and studied the connection between the structure of a group and the codegrees of its irreducible characters (see the details in [6], [11], [9]). In these papers, they studied the groups through considering the prime divisors of the codegrees of its irreducible characters. In [13], Yang et al. obtained that the group G is solvable if $\text{cod}(\chi) \leq p_\chi \cdot \chi(1)$ for each nonlinear irreducible character χ of a group G , where p_χ is the largest prime divisor of $|G : \ker(\chi)|$. Notice that the codegree is larger than its degree for any nonprincipal irreducible character of G (for any nonprincipal irreducible character χ of G , $\chi \in \text{Irr}(G/\ker(\chi))$, so that $|G/\ker(\chi)| > \chi(1)^2$ since $G/\ker(\chi) > 1$). So, it is natural to ask that what positive real number k such that

$$k \cdot \text{cod}(\chi) \leq \chi(1)^2, \quad (*)$$

for all nonlinear irreducible characters χ of G guarantees the solvability of G . In the next theorem, we give a sharp lower bound of k .

Theorem A. *Let G be a finite group with $k \cdot \text{cod}(\chi) \leq \chi(1)^2$ for every nonlinear irreducible character χ of G . If $k > a = \frac{2^9 \cdot 3^2 \cdot 19^2}{5 \cdot 7^3 \cdot 11 \cdot 31}$, then G is solvable. The real number a is best possible.*

*Corresponding author. Email: lvh529@163.com

2010 AMS Mathematics Subject Classification: 20C15

This research was supported by the National Natural Science Foundation of China (No.11971391, 12071376), the first and second authors are supported by the Chinese Scholarship Council.

The positive real number a in the above theorem is best possible. For instance, let $G \cong O'N$, and let $\theta \in \text{Irr}(G)$ be of minimal degree $10944 (= 2^6 \cdot 3^2 \cdot 19)$, we get

$$a \cdot |G| = 2^{18} \cdot 3^6 \cdot 19^3 = \theta(1)^3.$$

It forces $a \cdot |G| \leq \theta(1)^3$ for every nonlinear irreducible character θ of a nonabelian simple group G .

More generally, given a positive integer s , one could ask what positive real number k such that

$$k \cdot \text{cod}(\chi) \leq \chi(1)^s$$

for all nonlinear irreducible characters χ of G guarantees the solvability of G .

2 Proof

At first, we gather some lemmas before we actually prove the main theorem.

Lemma 2.1. *Suppose that N is a minimal normal nonabelian subgroup of a group G . Then there exists an irreducible character θ of N such that θ is extendible to G with $\theta(1) \geq 5$.*

Proof. See [7, Theorem 1.1]. □

We also need the following result regarding the characters of simple groups of Lie type.

Lemma 2.2. *Let S be a simple group of Lie type in characteristic p defined over a field of size q . Assume that $S \not\cong A_1(q), {}^2F_4(2)'$. Then there exist two irreducible characters θ_i , $i = 1, 2$, of S such that both θ_i extend to $\text{Aut}(S)$ with $1 < \theta_1(1) < \theta_2(1)$ and $\theta_2(1) = |S|_p$.*

Proof. See [12, Lemma 2.4]. □

Our proof will be based on the classification of finite simple groups. The main point of the classification is that if S is a nonabelian simple group, then S is either a simple group of Lie type, an alternating group of degree at least 5, one of 26 sporadic simple groups, or the Tits group. Often the Tits group, which denoted by ${}^2F_4(2)'$, is lumped in the groups of Lie type. Since the Tits group is in the Atlas [4], here we will consider this simple group separately.

For the simple groups of Lie type, there is a particular character called Steinberg character. If S is a simple group of Lie type with defining characteristic p and θ is the Steinberg character of S , then $\theta(1) = |S|_p$. By Lemma 2.1, if G has a normal subgroup S such that S is isomorphic to a finite simple group of Lie type, then the Steinberg character of S extends to G . Actually, the character θ_2 in Lemma 2.2 is the Steinberg character (see for instance [8, Proposition 3.1]).

For the alternating groups A_n , where $n \geq 5$, the automorphism group of A_n is S_n except $n = 6$. In [2, Theorem 3], it is proved that A_n , where $n \geq 6$, has a character of degree $\frac{n(n-3)}{2}$ which is extendible to $\text{Aut}(A_n)$.

For the sporadic simple groups, irreducible character degrees are available in the Atlas [4]. In Table 2, we list some character degrees of sporadic simple groups which will be relevant to our problems.

Lemma 2.3. *Let S be a simple group and $k > a = \frac{2^9 \cdot 3^2 \cdot 19^2}{5 \cdot 7^3 \cdot 11 \cdot 31}$. If S is not isomorphic to one of the groups in $\{A_1(q), G_2(q), {}^2G_2(q^2), Fi_{22}\}$, then S has some nontrivial irreducible character θ that is extendible to $\text{Aut}(S)$ and such that $k \cdot |S| > \theta(1)^3$.*

For a simple group of Lie type S , we know from [1, Lemma 2.6] that $|S| < \theta(1)^3$ for the Steinberg character θ (which is actually the θ_2 in Lemma 2.2). By Lemma 2.2, it is natural to check whether the character θ_1 of S satisfies $k \cdot |S| > \theta_1(1)^3$ or not. Based on the proof of [12, Lemma 2.4], Table 1 lists $\theta_1(1)$ for every simple group of Lie type in Lemma 2.3.

Proof of Lemma 2.3 Observe that $k > a = \frac{2^9 \cdot 3^2 \cdot 19^2}{5 \cdot 7^3 \cdot 11 \cdot 31} > \frac{5}{2}$.

Assume that S is an alternating group A_n , where $n \geq 5$. By [2, Theorem 3], S has some irreducible character θ of degree $\frac{n(n-3)}{2}$ that extends to $\text{Aut}(A_n)$. Furthermore, we have

$$\frac{k \cdot |A_n|}{\theta(1)^3} = \frac{4k \cdot (n-1)(n-2) \cdot (n-4)!}{n^2(n-3)^2}.$$

Note that $(n-1)(n-2) > n(n-3)$. If $n \geq 6$, then $4k \cdot (n-4)! > n(n-3)$ and hence $k \cdot |A_n| > \theta(1)^3$. If $n = 5$, A_5 has an irreducible character θ of degree 4 that is extendible to $\text{Aut}(A_5)$, and we have $k \cdot |A_5| > 2^3 \cdot 3 \cdot 5 > 4^3 = \theta(1)^3$, and we are done for the alternating groups A_n of degree at least 5.

Assume that S is a sporadic simple group other than Fi_{22} . By [2, Theorem 4], there exists some irreducible character $\theta \in \text{Irr}(S)$ of degree as in Table 2 that is extendible to $\text{Aut}(S)$. Also, it's easy to check $k \cdot |S| > \theta(1)^3$, we are also done for the sporadic simple groups but other than Fi_{22} .

Assume that $S \cong {}^2F_4(2)'$, by the Atlas [4], S has an irreducible character θ of degree 27 that is extendible to $\text{Aut}(S)$. Obviously, we also have $k \cdot |S| = k \cdot 2^{11} \cdot 3^3 \cdot 5^2 \cdot 13 > 27^3 = \theta(1)^3$.

Assume that S is a simple group of Lie type and $S \notin \{A_1(q), G_2(q), {}^2G_2(q^2)\}$. Take $\theta \in \text{Irr}(S)$ as the irreducible character θ_1 in Lemma 2.2, we have that θ is extendible to $\text{Aut}(S)$. Next we will check the stronger conclusion that $|S| > \theta(1)^3$ and hence the result follows immediately.

If $S \cong A_n(q)$, where $n \geq 2$, then

$$\begin{aligned} |A_n(q)| &= \frac{1}{(n+1, q-1)} q^{\frac{n(n+1)}{2}} (q^2 - 1)(q^3 - 1) \cdots (q^{n+1} - 1) \\ &> \frac{1}{q-1} q^{\frac{n(n+1)}{2}} q \cdot q^2 \cdots q^n = \frac{q^{n(n+1)}}{q-1}. \end{aligned}$$

By Table 1, choose $\theta \in \text{Irr}(S)$ with $\theta(1) = \frac{q(q^n-1)}{q-1} < \frac{q^{n+1}}{q-1}$, and we have

$$\frac{|S|}{\theta(1)^3} > \frac{q^{n(n+1)}(q-1)^3}{q^{3n+3}(q-1)} = q^{n^2-2n-3} \cdot (q-1)^2 = q^{(n-1)^2-4}(q-1)^2.$$

If $n \geq 3$, it is obvious that $(n-1)^2 - 4 \geq 0$. That means $|S| > \theta(1)^3$. If $n = q = 2$, then there exists a $\theta \in \text{Irr}(S)$ of degree 3 such that $|S| > \theta(1)^3$ by [4]. If $n = 2$ and $q \geq 3$, then since $|S| = |A_2(q)| \geq \frac{1}{q-1} q^3 (q^3 - 1)(q^2 - 1) = q^3 (q^3 - 1)(q + 1)$ and $\theta(1) = q(q + 1)$ we have

$$\frac{|S|}{\theta(1)^3} > \frac{q^3 (q^3 - 1)(q + 1)}{q^3 (q + 1)^3} = \frac{q^3 - 1}{(q + 1)^2} > 1.$$

This yields $|S| > \theta(1)^3$.

For every remaining simple group of Lie type, it is routine to check $|S| > \theta(1)^3$ for character θ of degree as in Table 1 (The similar calculation process is omitted here). Therefore, the proof is complete. $\square$

Let H be a subgroup of a finite group G , and let θ be an irreducible character of H . We use $\text{Irr}(G|\theta)$ to denote the set of irreducible characters of G lying over θ .

Lemma 2.4. *Let G be a group and N be its nonsolvable minimal normal subgroup. Suppose that $\text{cod}(\chi) \leq \chi(1)^2$ for each nonlinear irreducible character χ of G . Then N is nonabelian simple.*

Proof. Let $N = S_1 \times \cdots \times S_t$ where $S_i \cong S$ for a nonabelian simple group S . By the unique minimality of N , it follows that G acts transitively on $\{S_1, S_2, \dots, S_t\}$. Let $S = S_1$. Then $S \trianglelefteq N_G(S)$ and $t = |G : N_G(S)|$. By Lemma 2.1, S has some irreducible nonlinear character θ such that θ is extendible to $N_G(S)$. Take $\chi \in \text{Irr}(G|\theta)$. Then $\chi(1) \leq |G : N_G(S)| \cdot \theta(1) = t \cdot \theta(1)$.

Note that $\ker(\chi) = 1$. So, we conclude from the hypothesis that $|G| \leq \chi(1)^3$. Furthermore,

$$t^3 \cdot \theta(1)^3 \geq \chi(1)^3 \geq |G| \geq |G : N_G(S)| \cdot |N| = t \cdot |S|^t.$$

Since $\theta(1)^2 < |S|$, we have

$$t^2 \geq \frac{|S|^t}{\theta(1)^3} > |S|^{t-\frac{3}{2}}.$$

If $t \geq 3$, then since $t - \frac{3}{2} \geq \frac{t}{2}$ we have $t^2 > |S|^{t-\frac{3}{2}} \geq |S|^{\frac{t}{2}} = \sqrt{|S|}^t > 7^t$, a contradiction. Similarly, if $t = 2$, we have $4 > \sqrt{|S|} > 7$, which is also a contradiction. Thus $t = 1$. $\square$

Lemma 2.5. *If $q = p^f$ is a power of a prime p such that $q \equiv 3 \pmod{6}$, then $f < q^{\frac{1}{3}}$.*

Proof. If $f = 1$, we are done immediately. If $f = 2$, it follows from $q \geq 9$ that $2 \leq q^{\frac{1}{3}}$. Let function $T(x) = p^{\frac{x}{3}} - x$, where $x \geq 3$. Then the derivative function $T'(x) = \frac{\ln p}{3} \cdot p^{\frac{x}{3}} - 1 \geq \ln p - 1 \geq 0$. That means $T(f) \geq T(0) > 0$. That is, $f < p^{\frac{f}{3}} = q^{\frac{1}{3}}$. $\square$

Recall that $\Phi_n(x)$ is the n -th cyclotomic polynomial with indeterminate x . For a simple group of Lie type over finite field of q elements, we define $\Phi_n = \Phi_n(q)$. Next we will prove Theorem A.

Proof of the Theorem A. Suppose that G is a nonsolvable group of minimal order which satisfies the hypothesis of Theorem A.

We claim that G is not simple. Suppose not. If $G \notin \{A_1(q), G_2(q), {}^2G_2(q^2), Fi_{22}\}$, by Lemma 2.3, there exists some nonlinear irreducible character χ of G such that $k \cdot |G| > \chi(1)^3$. That is, $k \cdot \text{cod}(\chi) > \chi(1)^2$, which is contrary to the hypothesis. If G is isomorphic to one of the group in $\{A_1(q), G_2(q), {}^2G_2(q^2), Fi_{22}\}$, similarly, to get a contradiction, we just need to show that there also exists some nonlinear irreducible character θ of G such that $k \cdot |G| > \theta(1)^3$.

Suppose that $G \cong A_1(q)$. Let $\chi \in \text{Irr}(G)$ be of degree $q - 1$. Then since $k > \frac{5}{2}$ we have

$$k \cdot |G| > q(q^2 - 1) > (q - 1)^3 = \chi(1)^3.$$

Suppose that $G \cong G_2(q)$. Since $G_2(2) \cong U_3(3).2$ is not simple, we can assume that $q \geq 3$. Note that the character degrees of $G_2(q)$ are listed in the paper [3]. If $q \equiv 1 \pmod{6}$ or $q \equiv 4 \pmod{6}$, then $\Phi_2\Phi_6 \in \text{cd}(G)$. If $q \equiv 2 \pmod{6}$ or $q \equiv 5 \pmod{6}$, then $\Phi_1\Phi_3 \in \text{cd}(G)$. If $q \equiv 3 \pmod{6}$, then $\Phi_3\Phi_6 \in \text{cd}(G)$. It follows that

$$\frac{k \cdot |G|}{(\Phi_3\Phi_6)^3} = \frac{3q^6(q^6 - 1)(q^2 - 1)}{(q^2 + q + 1)^3(q^2 - q + 1)^3} > \frac{3(q^2 - 1)^4}{q^6 - 1} > 1.$$

Obviously, $\Phi_3\Phi_6 > \Phi_2\Phi_6 > \Phi_1\Phi_3$. So there always exists some character χ of G such that $k \cdot |G| > \chi(1)^3$.

Suppose that $G \cong {}^2G_2(q^2)$, where $q^2 = 3^{2m+1}$. Taking $\chi \in \text{Irr}(G)$ with degree Φ_{12} . Then

$$\frac{k \cdot |G|}{\Phi_{12}^3} \geq \frac{q^6(q^6 + 1)(q^2 - 1) \cdot (q^2 + 1)^3}{(q^4 + q^2 + 1)^3 \cdot (q^2 + 1)^3} \geq \frac{(q^2 - 1)(q^2 + 1)^3}{q^6 - 1} > 1.$$

Suppose that $G \cong Fi_{22}$. By the Atlas [4], it is easy to get that $k \cdot |G| > \chi(1)^3$ for some nonlinear $\chi \in \text{Irr}(G)$.

We next conclude the final contradiction. Note that G is not simple. Assume that G has distinct minimal normal subgroups, says N_1, N_2 . It follows from the minimality of G that G/N_i is solvable for $i = 1, 2$, and then $G \lesssim G/N_1 \times G/N_2$ is solvable, a contradiction. So we may assume that G has a unique minimal normal subgroup N ; and, similarly, G/N is solvable; and this means that N is also nonsolvable. Hence N is a direct product of isomorphic nonabelian simple groups. Since $\text{cod}(\chi) < k \cdot \text{cod}(\chi) \leq \chi(1)^2$ for every nonlinear irreducible character χ of G , it follows from Lemma 2.4 that the minimal normal subgroup N of G is simple. If $N \notin \{A_1(q), G_2(q), {}^2G_2(q^2), Fi_{22}\}$, then, by Lemma 2.3, N has some nontrivial irreducible character θ that is extendible to $\text{Aut}(N)$ (also to G) and $k \cdot |N| > \theta(1)^3$. Let $\chi \in \text{Irr}(G)$ be such that $\chi_N = \theta$. Then

$$k \cdot |G| > k \cdot |N| > \theta(1)^3 = \chi(1)^3.$$

Observe that $\ker(\chi) = 1$. This means $k \cdot \text{cod}(\chi) > \chi(1)^2$, a contradiction. Next we will also get a contradiction when N is isomorphic to one of the groups in $\{A_1(q), G_2(q), {}^2G_2(q^2), Fi_{22}\}$.

(1) Suppose that $N \cong A_1(q)$. Let $\theta \in \text{Irr}(N)$ be the Steinberg character. Then there exists $\chi \in \text{Irr}(G)$ such that $\chi_N = \theta$. Thus

$$k \cdot |G| \geq 2k \cdot |N| \geq kq(q^2 - 1) > q^3 = \theta(1)^3 = \chi(1)^3.$$

(2) Suppose that $N \cong G_2(q)$, where $q = p^f$. Then $|N| = q^6(q^6 - 1)(q^2 - 1)$.

If $q \not\equiv 3 \pmod{6}$, then, by the Atlas [4], we obtain $|G : N| \leq |\text{Out}(N)| \leq q$. Also, there exists some $\theta \in \text{Irr}(N)$ of degree at most $q^3 + 1$ (see for instance [3]). Let $\chi \in \text{Irr}(G|\theta)$. It follows that $\chi(1) \leq |G : N| \cdot \theta(1) \leq q(q^3 + 1)$. Hence

$$\frac{k \cdot |G|}{\chi(1)^3} \geq \frac{k \cdot q^6(q^6 - 1)(q^2 - 1)}{q^3(q^3 + 1)^3} = \frac{k \cdot q^3(q^3 - 1)(q^2 - 1)}{(q^3 + 1)^2}.$$

Note that $k > a > \frac{5}{2}$. Since $k \cdot q^3(q-1) > q^3 + 1$ and $(q^3 - 1)(q+1) > q^3 + 1$, then we have $k \cdot |G| > \chi(1)^3$.

If $q \equiv 3 \pmod{6}$, then $|G : N| \leq |\text{Out}(N)| \leq 2 \cdot f < 2 \cdot q^{\frac{1}{3}}$ where the last inequality follows from Lemma 2.5. Now let $\theta \in \text{Irr}(N)$ be of degree $(q^2 + q + 1)(q^2 - q + 1)$ and $\chi \in \text{Irr}(G|\theta)$. Then $\chi(1) < 2 \cdot q^{\frac{1}{3}}(q^2 + q + 1)(q^2 - q + 1)$. Thus

$$\frac{k \cdot |G|}{\chi(1)^3} \geq \frac{k \cdot q^6(q^6 - 1)(q^2 - 1)}{8q \cdot (q^2 + q + 1)^3(q^2 - q + 1)^3} = \frac{k \cdot q^5(q^2 - 1)^4}{8(q^6 - 1)^2} = \frac{k \cdot q^5(q + 1)^4(q - 1)^4}{8(q^6 - 1)^2}.$$

Since $q^5(q + 1) > q^6 - 1$ and $(q + 1)^3 > q^3 + 1$, it follows that

$$\frac{k \cdot |G|}{\chi(1)^3} \geq \frac{k(q + 1)^3(q - 1)^4}{8(q^6 - 1)} \geq \frac{k(q - 1)^4}{8(q^3 - 1)}.$$

Recall that $q \equiv 3 \pmod{6}$. If $q \geq 9$, it is easy to get $k \cdot |G| > \chi(1)^3$. If $q = 3$,

$$\frac{k \cdot |G|}{\chi(1)^3} \geq \frac{2k \cdot |N|}{\chi(1)^3} = \frac{k \cdot 2^4 \cdot 3^6(3^6 - 1)}{2^3 \cdot 3 \cdot 13^3 \cdot 7^3} = \frac{2k \cdot 3^5(3^6 - 1)}{13^3 \cdot 7^3} > 1.$$

(3) Suppose that $N \cong {}^2G_2(q^2)$, where $q^2 = 3^{2m+1}$. Since $q^2 = 3^{2m+1} \equiv 3 \pmod{6}$, it follows from Lemma 2.5 that $|G : N| \leq |\text{Out}(N)| = 2m + 1 < q^{\frac{2}{3}}$. Let $\theta \in \text{Irr}(N)$ be of degree Φ_{12} and $\chi \in \text{Irr}(G|\theta)$. Then $\chi(1) \leq |G : N| \cdot \theta(1) \leq q^{\frac{2}{3}} \cdot \Phi_{12}$. Thus

$$\frac{3 \cdot |G|}{\chi(1)^3} \geq \frac{q^6(q^6 + 1)(q^2 - 1) \cdot (q^2 + 1)^3}{q^2 \cdot (q^4 + q^2 + 1)^3 \cdot (q^2 + 1)^3} \geq \frac{(q^2 - 1)(q^2 + 1)^3}{q^2(q^6 - 1)} > 1.$$

(4) Suppose that $N \cong Fi_{22}$. By the Atlas [4], N has an irreducible character θ of degree 78. Let $\chi \in \text{Irr}(G|\theta)$. Then since $|\text{Out}(N)| = 2$ we get $\chi(1) \leq |G : N| \cdot \theta(1) = 156 = 2^2 \cdot 3 \cdot 13$. It follows that

$$k \cdot |G| \geq 2k \cdot |N| = k \cdot 2^{18} \cdot 3^9 \cdot 5^2 \cdot 7 \cdot 11 \cdot 13 > (2^2 \cdot 3 \cdot 13)^3 = \chi(1)^3.$$

Therefore we can find an irreducible character χ of G such that $k \cdot |G| > \chi(1)^3$ when $N \in \{A_1(q), G_2(q), {}^2G_2(q^2), Fi_{22}\}$. Since $[\chi_N, 1_N] = 0$, it follows from the minimality of N that $\ker(\chi) = 1$. So $k \cdot \text{cod}(\chi) > \chi(1)^2$, a contradiction. The proof is established. $\square$

Table 1: some degrees of some simple groups of Lie type

Group	sym.	Degree	Group	sym.	Degree
$A_n(q)(n \geq 2)$	$(1, n)$	$\frac{q^{n+1}-q}{q-1}$	$F_4(q)$	$\theta_{9,2}$	$q^2\Phi_3^2\Phi_6^2\Phi_{12}$
${}^2A_n(q)$	$(1, n)$	$\frac{q^{n+1}+(-1)^{n+1}q}{q+1}$	$E_6(q)$	$\theta_{6,1}$	$q\Phi_8\Phi_9$
$B_n(q)$	$\begin{pmatrix} 0 & 1 & n \\ & - & \end{pmatrix}$	$\frac{(q^n-1)(q^n-q)}{2(q+1)}$	${}^2E_6(q^2)$	$\theta_{2,4'}$	$q\Phi_8\Phi_{18}$
$C_n(q)$	$\begin{pmatrix} 0 & 1 & n \\ & - & \end{pmatrix}$	$\frac{(q^n-1)(q^n-q)}{2(q+1)}$	$E_7(q)$	$\theta_{7,1}$	$q\Phi_7\Phi_{12}\Phi_{14}$
$D_n(q)$	$\begin{pmatrix} n-1 \\ & 1 \end{pmatrix}$	$\frac{(q^n-1)(q^{n-1}+q)}{q^2-1}$	$E_8(q)$	$\theta_{8,1}$	$q\Phi_4^2\Phi_8\Phi_{12}\Phi_{20}\Phi_{24}$
${}^2D_n(q^2)$	$\begin{pmatrix} 1 & n-1 \\ & - \end{pmatrix}$	$\frac{(q^n+1)(q^{n-1}-q)}{q^2-1}$	${}^2B_2(q^2)$	${}^2B_2[a]$	$\frac{1}{\sqrt{2}}q\Phi_1\Phi_2$
${}^3D_4(q^3)$	$\theta_{1,3'}$	$q\Phi_{12}$	${}^2F_4(q^2)$	${}^2B_2[a]$	$\frac{1}{\sqrt{2}}q\Phi_1\Phi_2\Phi_4^2\Phi_6$

Table 2: degrees of the sporadic simple groups

Group	char.	Degree	Group	char.	Degree
M_{11}	χ_2	$10 = 2 \cdot 5$	$O'N$	χ_2	$10944 = 2^6 \cdot 3^2 \cdot 19$
M_{12}	χ_7	$54 = 2 \cdot 3^3$	Co_3	χ_2	23
J_1	χ_4	$76 = 2^2 \cdot 19$	Co_2	χ_2	23
M_{22}	χ_2	$21 = 3 \cdot 7$	Fi_{22}	χ_{56}	$1441792 = 2^{17} \cdot 11$
J_2	χ_6	$36 = 2^2 \cdot 3^2$	HN	χ_{10}	$16929 = 3^4 \cdot 11 \cdot 19$
M_{23}	χ_6	$22 = 2 \cdot 11$	Ly	χ_7	$120064 = 2^8 \cdot 7 \cdot 67$
HS	χ_2	$22 = 2 \cdot 11$	Th	χ_2	$248 = 2^3 \cdot 31$
J_3	χ_6	$324 = 2^3 \cdot 3^4$	Fi_{23}	χ_4	$5083 = 13 \cdot 17 \cdot 23$
M_{24}	χ_2	23	Co_1	χ_3	$299 = 13 \cdot 23$
M^cL	χ_2	$22 = 2 \cdot 11$	J_4	χ_2	$1333 = 31 \cdot 43$
He	χ_9	$1275 = 3 \cdot 5^2 \cdot 17$	Fi'_{24}	χ_2	$8671 = 23 \cdot 29 \cdot 13$
Ru	χ_5	$783 = 3^3 \cdot 29$	B	χ_2	$4371 = 3 \cdot 31 \cdot 47$
Suz	χ_2	$143 = 11 \cdot 13$	M	χ_2	$196883 = 59 \cdot 71 \cdot 47$

References

- [1] N. Ahanjideh, On Thompson's conjecture for some finite simple groups, *J. Algebra* 344(2011), 205-228.
- [2] M. Bianchi, D. Chillai, M. Lewis and E. Pacifici, Character degree graphs that are complete graphs, *Proc. Amer. Math. Soc.* 135(3) (2007), 671-676.
- [3] B. Chang and R. Ree, The characters of $G_2(q)$, *Symposia Mathematica*(Convegno di Gruppi e loro Rappresentazioni, INDAM,Rome) 13(1972), 395-413.
- [4] J. Conway, R. Curtis, S. Norton, R. Parker and R. Wilson, *Atlas of finite groups*, Oxford University Press, London, 1984.
- [5] I. Isaacs, *Character theory of finite groups*, Academic Press, New York, 1976.
- [6] D. Liang and G. Qian, Finite groups with coprime character degrees and codegrees, *J. Group Theory* 19 (2016), 763-776.
- [7] K. Magaard and H. Tong-Viet, Character degree sums in finite nonsolvable groups, *J. Group Theory* 14 (2011), 53-57.
- [8] G. Malle and A. Zalesski, Steinberg-like characters for finite simple groups, *J. Group Theory* 23(1) (2020), 25-78.
- [9] G. Qian, A character theoretic criterion for a p -closed group, *Israel J. Math.* 190 (2012), 401-412.
- [10] G. Qian, Y. Yang and H. Wei, Codegrees of irreducible characters in finite group, *J. Algebra* 312 (2007), 946-955.
- [11] G. Qian, A note on element orders and character codegrees, *Arch. Math.* 97 (2011), 99-103.
- [12] H. Tong-Viet, Symmetric groups are determined by their character degrees, *J. Algebra* 334 (2011), 275-284.
- [13] D. Yang, C. Gao and H. Lv, Finite groups with special codegrees, *Arch. Math.* 115 (2020), 121-130.