

Poisson approximation to the binomial distribution: extensions to the convergence of positive operators

Ana-Maria Acu^{*a}, Margareta Heilmann^b, Ioan Rasa^c, Andra Seserman^d

^aLucian Blaga University of Sibiu, Department of Mathematics and Informatics, Str. Dr. I. Ratiu, No.5-7, RO-550012 Sibiu, Romania, e-mail: anamaria.acu@ulbsibiu.ro, telephone number: +040766258465

^bSchool of Mathematics and Natural Sciences, University of Wuppertal, Gaußstraße 20, D-42119 Wuppertal, Germany, e-mail: heilmann@math.uni-wuppertal.de

^cTechnical University of Cluj-Napoca, Faculty of Automation and Computer Science, Department of Mathematics, Str. Memorandumului nr. 28, 400114 Cluj-Napoca, Romania e-mail: ioan.rasa@math.utcluj.ro

^dTechnical University of Cluj-Napoca, Faculty of Automation and Computer Science, Department of Mathematics, Str. Memorandumului nr. 28, 400114 Cluj-Napoca, Romania e-mail: campan_aandra@yahoo.com

Abstract

The idea behind Poisson approximation to the binomial distribution was used in [J. de la Cal, F. Luquin, J. Approx. Theory, 68(3), 1992, 322-329] and subsequent papers in order to establish the convergence of suitable sequences of positive linear operators. The proofs in these papers are given using probabilistic methods. We use similar methods, but in analytic terms. In this way we recover some known results and establish several new ones. In particular, we enlarge the list of the limit operators and give characterizations of them.

Keywords: Poisson approximation to binomial distribution; positive linear operators; convergence; limit operator.

2010 MSC: 41A36, 60E05, 60E10.

1. Introduction

Let $X_n(p_n)$ be a binomial random variable with parameters $n \in \mathbb{N}$ and $p_n \in [0, 1]$. Its characteristic function is

$$\begin{aligned} g_n(s) &= E(\exp(isX_n)) = \sum_{k=0}^n \binom{n}{k} p_n^k (1-p_n)^{n-k} \exp(isk) \\ &= (1-p_n + p_n e^{is})^n, s \in \mathbb{R}, \end{aligned}$$

where E stands for mathematical expectation and i is the imaginary unit.

Let $X(\lambda)$ be a Poisson variable with parameter $\lambda \geq 0$, having the characteristic function

$$h(s) = \sum_{k=0}^{\infty} e^{-\lambda} \frac{\lambda^k}{k!} \exp(isk) = e^{\lambda(e^{is}-1)}, s \in \mathbb{R}.$$

Suppose that $\lim_{n \rightarrow \infty} np_n = \lambda$. Then $\lim_{n \rightarrow \infty} g_n(s) = h(s)$, $s \in \mathbb{R}$, and hence $X_n(p_n)$ converges in law to $X(\lambda)$. This is the celebrated result about "Poisson approximation to the binomial distribution", see, e.g., [19], [36]. It implies

$$\lim_{n \rightarrow \infty} E\varphi(X_n(p_n)) = E\varphi(X(\lambda)), \quad \varphi \in C_b[0, \infty), \quad (1.1)$$

where $C_b[0, \infty)$ is the space of all real valued, continuous and bounded functions on $[0, \infty)$.

Now let $x \geq 0$, $m \in \mathbb{N}$, $p_n = \frac{x}{n}$, $\lambda = mx$. From (1.1) we deduce

$$\lim_{n \rightarrow \infty} E\varphi\left(X_{mn}\left(\frac{x}{n}\right)\right) = E\varphi(X(mx)), \quad \varphi \in C_b[0, \infty).$$

This can be written as

$$\lim_{n \rightarrow \infty} \sum_{k=0}^{mn} \binom{mn}{k} \left(\frac{x}{n}\right)^k \left(1 - \frac{x}{n}\right)^{mn-k} \varphi(k) = \sum_{k=0}^{\infty} e^{-mx} \frac{(mx)^k}{k!} \varphi(k). \quad (1.2)$$

Let $f \in C_b[0, \infty)$ and $\varphi \in C_b[0, \infty)$, $\varphi(t) = f\left(\frac{t}{m}\right)$, $t \geq 0$. Now (1.2) becomes

$$\lim_{n \rightarrow \infty} \sum_{k=0}^{mn} \binom{mn}{k} \left(\frac{x}{n}\right)^k \left(1 - \frac{x}{n}\right)^{mn-k} f\left(\frac{k}{m}\right) = \sum_{k=0}^{\infty} e^{-mx} \frac{(mx)^k}{k!} f\left(\frac{k}{m}\right).$$

In terms of Bernstein and Szász-Mirakjan operators this means that

$$\lim_{n \rightarrow \infty} B_{mn}\left(f\left(\frac{nt}{n}\right); \frac{x}{n}\right) = B_m^{[0]}(f(t); x), \quad f \in C_b[0, \infty). \quad (1.3)$$

This result and similar ones involving other sequences of operators, were obtained in [6, 7, 8, 13, 14, 15] using probabilistic methods and making references to several probability distributions. Rates of convergence in relations like (1.3) can be found in the same papers and the references therein.

In our paper we present a general result extending (1.3), using analytic terms and analytic methods. We apply it to many families of positive linear operators recovering some known results and establishing new ones.

In (1.3) a specific modification of the Bernstein operators, inspired by Poisson approximation to the binomial distribution, is presented. We will consider several other modifications corresponding to suitable families of operators. As limit operator the Szász-Mirakjan operator appears in (1.3). It will have the same role in several other examples. But other operators will also appear as limit operators and we will characterize some of them.

Let L_n and L be positive linear operators associated with random variables. In Section 2 we show that the family of the complex exponentials $t \in \mathbb{R} \rightarrow e^{ist}$, indexed by $s \in \mathbb{R}$, can be used as test family for the convergence of sequences L_n toward the operator L . This is presented in Theorem 2.1 which is the main result of our paper. We will apply Theorem 2.1 to several classical or new sequences of positive linear operators. Sections 3-9 are devoted to sequences for which the limit operator is the classical Szász-Mirakjan operator. The Gamma operator and the Weierstrass operator appear as the limit operator in Sections 10-11. Other limit operators are considered in Section 12. This section contains several characterizations of the limit operators.

The necessary definitions and notations will be presented in the next sections.

2. The complex exponentials as test functions for convergence of operators

Let $I \subseteq \mathbb{R}$ be an interval and $C_b(I)$ the space of all real-valued continuous and bounded functions on I . Let Z^x, Z_1^x, Z_2^x, \dots be I -valued random variables whose probability distributions depend on a real parameter $x \in I$. Suppose that for each $f \in C_b(I)$ the functions $x \rightarrow Ef(Z^x)$ and $x \rightarrow Ef(Z_n^x)$, $x \in I$, $n \geq 1$, are continuous on I .

Consider the positive linear operators

$$L_n : C_b(I) \rightarrow C_b(I), \quad n \geq 1, \quad \text{and} \quad L : C_b(I) \rightarrow C_b(I),$$

defined for $f \in C_b(I)$ and $x \in I$ by

$$L_n(f(t); x) := Ef(Z_n^x), \quad L(f(t); x) := Ef(Z^x). \quad (2.1)$$

For each $s \in \mathbb{R}$ we consider the function $t \rightarrow e^{ist}$, $t \in I$, and define

$$L_n(e^{ist}; x) = L_n(\cos(st); x) + iL_n(\sin(st); x)$$

and similarly for $L(e^{ist}; x)$.

Theorem 2.1. *Suppose that for each $s \in \mathbb{R}$ and $x \in I$,*

$$\lim_{n \rightarrow \infty} L_n(e^{ist}; x) = L(e^{ist}; x), \quad (2.2)$$

and for each $x \in I$ the function $s \rightarrow L(e^{ist}; x)$ is continuous on \mathbb{R} . Then

$$\lim_{n \rightarrow \infty} L_n(f(t); x) = L(f(t); x) \quad (2.3)$$

for all $f \in C_b(I)$ and $x \in I$.

Proof. Let $g_n^x(s)$ and $g^x(s)$ be the characteristic functions of Z_n^x and Z^x , $n \geq 1$, $x \in I$. Then, according to (2.1)

$$\begin{aligned} g_n^x(s) &:= E(e^{isZ_n^x}) = L_n(e^{ist}; x), \\ g^x(s) &:= E(e^{isZ^x}) = L(e^{ist}; x). \end{aligned}$$

Now (2.2) shows that

$$\lim_{n \rightarrow \infty} g_n^x(s) = g^x(s), \quad s \in \mathbb{R},$$

and by hypothesis g^x is continuous on \mathbb{R} .

Consequently we can apply the continuity theorem of Lévy (see [19], [36]) to conclude that the sequence $(Z_n^x)_{n \geq 1}$ converges in law to Z^x . Furthermore, this implies (see [19], [36])

$$\lim_{n \rightarrow \infty} Ef(Z_n^x) = Ef(Z^x), \quad (2.4)$$

and now (2.3) is a consequence of (2.1) and (2.4). \square

In the next sections we will be concerned with positive linear operators L_n and L for which it is easy to identify the corresponding random variables. See also [6, 7, 8, 13, 14, 15].

3. The Baskakov type operators $B_n^{[c]}$

Let $c \in \mathbb{R}$, $n \in \mathbb{N}$, $n > c$ for $c \geq 0$ and $-n/c \in \mathbb{N}$ for $c < 0$. Furthermore let $I_c = [0, \infty)$ for $c \geq 0$ and $I_c = [0, -1/c]$ for $c < 0$. Consider $f : I_c \rightarrow \mathbb{R}$ given in such a way that the corresponding integrals and series are convergent.

The Baskakov-type operators are defined as follows (see [11, 28, 27])

$$B_n^{[c]}(f; x) = \sum_{j=0}^{\infty} p_{n,j}^{[c]}(x) f\left(\frac{j}{n}\right),$$

where

$$p_{n,j}^{[c]}(x) = \begin{cases} \frac{n^j}{j!} x^j e^{-nx} & , c = 0, \\ \frac{n^{c,\bar{j}}}{j!} x^j (1+cx)^{-(\frac{n}{c}+j)} & , c \neq 0, \end{cases} \quad (3.1)$$

and $a^{c,\bar{j}} := \prod_{l=0}^{j-1} (a + cl)$, $a^{c,\bar{0}} := 1$.

For $c = -1$ one recovers the well known Bernstein operators

$$B_n^{[-1]}(f; x) := \sum_{j=0}^n f\left(\frac{j}{n}\right) p_{n,j}^{[-1]}(x), \text{ where } p_{n,j}^{[-1]}(x) := \binom{n}{j} x^j (1-x)^{n-j}, \quad x \in [0, 1].$$

The classical Baskakov operators are obtained for $c = 1$ and are defined as follows

$$B_n^{[1]}(f; x) := \sum_{j=0}^{\infty} f\left(\frac{j}{n}\right) p_{n,j}^{[1]}(x), \text{ where } p_{n,j}^{[1]}(x) := \binom{n+j-1}{j} \frac{x^j}{(1+x)^{n+j}}, \quad x \in [0, \infty).$$

The classical Szász-Mirakjan operators are Baskakov type operators with $c = 0$ defined by

$$B_n^{[0]}(f; x) := \sum_{j=0}^{\infty} p_{n,j}^{[0]}(x) f\left(\frac{j}{n}\right), \text{ where } p_{n,j}^{[0]}(x) = e^{-nx} \frac{(nx)^j}{j!}, \quad x \in [0, \infty). \quad (3.2)$$

Theorem 3.1. *Let $m \in \mathbb{N}$, $c \in \mathbb{R}$ and $x \in I_c$ be given. Then*

$$\lim_{n \rightarrow \infty} B_{mn}^{[c]} \left(f(nt); \frac{x}{n} \right) = B_m^{[0]}(f(t); x), \quad f \in C_b(I_c). \quad (3.3)$$

Proof. Consider first the case $c = 0$. Using elementary calculations we find

$$B_{mn}^{[0]} \left(f(nt); \frac{x}{n} \right) = B_m^{[0]}(f(t); x).$$

Now let $c \neq 0$. Then

$$\begin{aligned} B_n^{[c]}(e^{ist}; x) &= \sum_{j=0}^{\infty} \frac{n^{c,\bar{j}}}{j!} x^j (1+cx)^{-n/c-j} e^{isj/n} \\ &= (1+cx)^{-n/c} \sum_{j=0}^{\infty} \frac{n(n+c) \dots (n+(j-1)c)}{j!} \left(\frac{xe^{is/n}}{1+cx} \right)^j \\ &= (1+cx)^{-n/c} \sum_{j=0}^{\infty} \binom{n/c+j-1}{j} \left(\frac{cxe^{is/n}}{1+cx} \right)^j \\ &= (1+cx)^{-n/c} \left(1 - \frac{cxe^{is/n}}{1+cx} \right)^{-n/c}, \end{aligned}$$

and so

$$B_n^{[c]}(e^{ist}; x) = \left(1 - cx(e^{is/n} - 1)\right)^{-n/c}. \quad (3.4)$$

Define

$$L_n(f(t); x) := B_{mn}^{[c]} \left(f(nt); \frac{x}{n}\right), \quad (3.5)$$

$$L(f(t); x) := B_m^{[0]}(f(t); x). \quad (3.6)$$

From (3.4), (3.5), (3.6) we obtain

$$\begin{aligned} \lim_{n \rightarrow \infty} L_n(e^{ist}; x) &= \lim_{n \rightarrow \infty} B_{mn}^{[c]} \left(e^{isnt}; \frac{x}{n}\right) \\ &= \lim_{n \rightarrow \infty} \left(1 - c \frac{x}{n} (e^{is/m} - 1)\right)^{-mn/c} = e^{mx(e^{is/m} - 1)} \\ &= B_m^{[0]}(e^{ist}; x) = L(e^{ist}; x). \end{aligned}$$

Thus (2.2) is satisfied and (3.3) is a consequence of (2.3), (3.5) and (3.6). \square

4. The k th order Kantorovich modification of the Baskakov type operators

The k -th order Kantorovich modifications of the operators $B_n^{[c]}$ are defined by

$$B_n^{[c](k)} := D^k \circ B_n^{[c]} \circ I_k, \quad k \in \mathbb{N},$$

where D^k denotes the k -th order ordinary differential operator and

$$I_k f = f, \text{ if } k = 0 \text{ and } (I_k f)(x) = \int_0^x \frac{(x-t)^{k-1}}{(k-1)!} f(t) dt, \text{ if } k \in \mathbb{N}.$$

We have $B_n^{[c](k)}(f(t); x) := \frac{d^k}{dx^k} B_n^{[c]}((I_k f)(t); x)$. Using (3.4) we get

$$\begin{aligned} B_n^{[c](k)}(e^{ist}; x) &= \frac{1}{(is)^k} \frac{d^k}{dx^k} B_n^{[c]}(e^{ist}; x) \\ &= n^{c, \bar{k}} \left(\frac{e^{is/n} - 1}{is}\right)^k \left(1 - cx(e^{is/n} - 1)\right)^{-n/c-k}, \quad c \neq 0, \end{aligned} \quad (4.1)$$

$$B_n^{[c](k)}(e^{ist}; x) = \left(\frac{n(e^{is/n} - 1)}{is}\right)^k e^{nx(e^{is/n} - 1)}, \quad c = 0. \quad (4.2)$$

The image of the constant function $\mathbf{1}$ under $B_n^{[c](k)}$ is obtained with $s = 0$ and is the constant function $\frac{n^{c, \bar{k}}}{n^k} \mathbf{1}$. Therefore the operators

$$V_n^{[c](k)} := (n^k / n^{c, \bar{k}}) B_n^{[c](k)} \quad (4.3)$$

preserve the constant function $\mathbf{1}$.

Theorem 4.1. For $m \in \mathbb{N}$, $c \in \mathbb{R}$, $x \in I_c$, and $f \in C_b(I)$, we have

$$\lim_{n \rightarrow \infty} V_{mn}^{[c](k)} \left(f(nt); \frac{x}{n} \right) := V_m^{[0](k)} (f(t); x). \quad (4.4)$$

Proof. According to Theorem 2.1, it suffices to show that (4.4) holds for $f(t) := e^{ist}$, $s \in \mathbb{R}$.

Let $c \neq 0$. Using (4.3) and (4.1) we have

$$\begin{aligned} \lim_{n \rightarrow \infty} V_{mn}^{[c](k)} \left(e^{isnt}; \frac{x}{n} \right) &= \lim_{n \rightarrow \infty} \left(\frac{m}{is} (e^{is/m} - 1) \right)^k \left(1 - c \frac{x}{n} (e^{is/m} - 1) \right)^{-mn/c-k} \\ &= \left(\frac{m}{is} (e^{is/m} - 1) \right)^k e^{mx(e^{is/m} - 1)} \\ &= V_m^{[0](k)} (f(t); x). \end{aligned}$$

Using (4.3) and (4.2) it is easy to check that (4.4) holds for $f(t) = e^{ist}$, $s \in \mathbb{R}$, also for $c = 0$. Now (4.4) for $f \in C_b(I)$ is a consequence of Theorem 2.1. \square

Remark 4.1. Since $\lim_{n \rightarrow \infty} n^k/n^{c,k} = 1$, from (4.3) and (4.4) we derive

$$\lim_{n \rightarrow \infty} B_{mn}^{[c](k)} \left(f(nt); \frac{x}{n} \right) = B_m^{[c](k)} (f(t); x).$$

5. Discrete operators associated with $V_{n,\rho}^{[c](k)}$

Depending on a parameter $\rho \in \mathbb{R}^+$ the linking operators are given by

$$(B_{n,\rho}^{[c]} f)(x) = \sum_{j=0}^{\infty} F_{n,j}^{[c],\rho}(f) p_{n,j}^{[c]}(x), \quad f : I_c \longrightarrow \mathbb{R},$$

where

$$F_{n,j}^{[c],\rho}(f) = \begin{cases} f(0) & , \quad j = 0, c \in \mathbb{R}, \\ f\left(-\frac{1}{c}\right) & , \quad j = -\frac{n}{c}, c < 0, \\ \int_{I_c} \mu_{n,j}^{[c],\rho}(t) f(t) dt & , \quad \text{otherwise,} \end{cases}$$

with

$$\mu_{n,j}^{[c],\rho}(t) = \begin{cases} \frac{(-c)^{j\rho}}{B(j\rho, -(\frac{n}{c} + j)\rho)} t^{j\rho-1} (1+ct)^{-(\frac{n}{c}+j)\rho-1} & , \quad c < 0, \\ \frac{(n\rho)^{j\rho}}{\Gamma(j\rho)} t^{j\rho-1} e^{-n\rho t} & , \quad c = 0, \\ \frac{c^{j\rho}}{B(j\rho, \frac{n}{c}\rho + 1)} t^{j\rho-1} (1+ct)^{-(\frac{n}{c}+j)\rho-1} & , \quad c > 0. \end{cases}$$

By $B(x, y) = \int_0^1 t^{x-1} (1-t)^{y-1} dt$, $x, y > 0$ we denote Euler's Beta function and by Γ the Gamma function.

These operators represent a link between Baskakov type operators $B_n^{[c]}$ (for $\rho = \infty$) and the genuine Baskakov-Durrmeyer type operators (for $\rho = 1$). Convergence results concerning the linking operators can be found in [10, Section 5], [20, Th. 2.3], [24, Th. 1, Th. 2, Cor. 1] and [33, Th. 4].

Let $B_{n,\rho}^{[c](k)} := D^k \circ B_{n,\rho}^{[c]} \circ I_k$ be the k -th order Kantorovich modification of the operators $B_{n,\rho}^{[c]}$ and $V_{n,\rho}^{[c](k)} := \frac{(n\rho)^{c,\underline{k}}}{\rho^k n^{c,\overline{k}}} B_{n,\rho}^{[c](k)}$. For $\rho \in \mathbb{N}$ explicit representation are given in [25], [26] and [27].

The discrete operator associated with $V_{n,\rho}^{[c](k)}$ was introduced in [23] as follows

$$D_{n,\rho}^{[c](k)}(f(t), x) = \sum_{j=0}^{\infty} f\left(\frac{(2j+k)\rho+k}{2(n\rho-kc)}\right) p_{n+kc,j}^{[c]}(x).$$

Note that $D_{n,\rho}^{[c](k)}e_0 = e_0$. For $c \neq 0$ and $\rho n > kc$, we have

$$\begin{aligned} D_{n,\rho}^{[c](k)}(e^{ist}; x) &= e^{\frac{isk(\rho+1)}{2(n\rho-kc)}} (1+cx)^{-n/c-k} \sum_{j=0}^{\infty} \binom{n/c+k+j-1}{j} \left(\frac{cx e^{is\rho/(n\rho-kc)}}{1+cx}\right)^j \\ &= e^{\frac{isk(\rho+1)}{2(n\rho-kc)}} (1+cx)^{-n/c-k} \left(1 - \frac{cx e^{is\rho/(n\rho-kc)}}{1+cx}\right)^{-n/c-k}. \end{aligned}$$

Therefore,

$$D_{n,\rho}^{[c](k)}(e^{ist}; x) = e^{\frac{isk(\rho+1)}{2(n\rho-kc)}} \left(1 - cx(e^{is\rho/(n\rho-kc)} - 1)\right)^{-n/c-k}. \quad (5.1)$$

Similarly,

$$D_{n,\rho}^{[0](k)}(e^{ist}; x) = e^{\frac{isk(\rho+1)}{2n\rho} + nx(e^{is/n} - 1)}. \quad (5.2)$$

Theorem 5.1. *Let $m \in \mathbb{N}$, $c \in \mathbb{R}$, $x \in I_c$, $f \in C_b(I)$. Then*

$$\lim_{n \rightarrow \infty} D_{mn,\rho}^{[c](k)}\left(f(nt); \frac{x}{n}\right) = D_{m,\rho}^{[0](k)}(f(t); x). \quad (5.3)$$

Proof. Using (5.1) and (5.2) it is easy to check that (5.3) is valid for $f(t) = e^{ist}$, $s \in \mathbb{R}$. Therefore, (5.3) is a consequence of Theorem 2.1. \square

6. Other examples

6.1.

For a non-negative constant a , independent of n , in [32] it was introduced the operator

$$Q_n^a(f; x) = \sum_{k=0}^{\infty} q_{n,k}^a(x) f\left(\frac{k}{n}\right),$$

where

$$q_{n,k}^a(x) = e^{-\frac{ax}{1+x}} \frac{\sum_{i=0}^k \binom{k}{i} n^{1,\bar{i}} a^{k-i}}{k!} \frac{x^k}{(1+x)^{n+k}}.$$

For the operators Q_n^a we have (see [21, 1.10.2])

$$Q_n^a(e^{ist}; x) = (1+x - x e^{is/n})^{-n} \exp\left(\frac{ax}{1+x} (e^{is/n} - 1)\right)$$

Consequently,

$$\begin{aligned}\lim_{n \rightarrow \infty} Q_{mn}^a \left(e^{isnt}, \frac{x}{n} \right) &= \lim_{n \rightarrow \infty} \left(1 + x/n \left(1 - e^{is/m} \right) \right)^{-mn} e^{\frac{ax}{n+x} (e^{is/m} - 1)} \\ &= e^{mx(e^{is/m} - 1)} = B_m^{[0]}(e^{ist}; x).\end{aligned}$$

Using Theorem 2.1 we get for $f \in C_b[0, \infty)$, $m \geq 1$, $x \in [0, \infty)$,

$$\lim_{n \rightarrow \infty} Q_{mn}^a \left(f(nt); \frac{x}{n} \right) = B_m^{[0]}(f(t); x).$$

6.2.

Let $c \neq 0$ and

$$\begin{aligned}\sigma : I_c &\rightarrow I_{-c}, \quad \sigma(x) = \frac{x}{1 + cx}, \\ \psi : I_c &\rightarrow I_{-c}, \quad \psi(x) = \frac{x}{1 - cx}.\end{aligned}$$

Let $B_{n,1}^{[c](k)}$ be the k th order Kantorovich modifications of the genuine Baskakov-Durrmeyer type operators. In [22] the second author of this paper presented a method to relate the k th order Kantorovich modifications of Durrmeyer type variants of Bleimann-Butzer-Hahn operators for $c < 0$ and Meyer-König and Zeller operators for $c > 0$ with $B_{n,1}^{[c](k)}$ as follows

$$\overline{B}_{n,1}^{[c](k)}(f; x) := B_{n,1}^{[c](k)}(f \circ \sigma; \psi(x)).$$

For details see [22, Sec.6]. Using Theorem 2.1 we derive

$$\begin{aligned}\lim_{n \rightarrow \infty} \overline{B}_{mn,1}^{[c](k)} \left(f(nt); \frac{x}{n} \right) &= \lim_{n \rightarrow \infty} B_{mn,1}^{[c](k)} \left((f \circ \sigma)(nt); \frac{\psi(x)}{n} \right) \\ &= B_{m,1}^{[0](k)}((f \circ \sigma)(t); \psi(x)).\end{aligned}\tag{6.1}$$

7. The Bleimann-Butzer-Hahn operators

Let

$$H_n(f(t); x) := (1+x)^{-n} \sum_{k=0}^n \binom{n}{k} x^k f\left(\frac{k}{n-k+1}\right)\tag{7.1}$$

be the Bleimann-Butzer-Hahn operators [12]. Using the probabilistic approach it was proved in [13, Th. 2 a)] that

$$\lim_{n \rightarrow \infty} H_{mn} \left(f\left(\frac{nt}{1+t}\right); \frac{x}{n} \right) = B_m^{[0]}(f(t); x).\tag{7.2}$$

Here is an analytic proof based on Theorem 2.1. In fact, all that we have to prove is that (7.2) holds for $f(t) := e^{ist}$, $s \in \mathbb{R}$.

This immediately follows from

$$\begin{aligned}\lim_{n \rightarrow \infty} H_{mn} \left(e^{isnt/(1+t)}; \frac{x}{n} \right) &= \lim_{n \rightarrow \infty} \left(1 + \frac{x}{n} \right)^{-mn} \sum_{k=0}^{mn} \binom{mn}{k} \left(\frac{x}{n} \right)^k e^{isnk/(mn+1)} \\ &= \lim_{n \rightarrow \infty} \left(1 + \frac{x}{n} \right)^{-mn} \left(1 + \frac{x}{n} e^{isn/(mn+1)} \right)^{mn} \\ &= e^{mx(e^{is/m} - 1)} = B_m^{[0]}(e^{ist}; x).\end{aligned}$$

Example 7.1. It was proved in [6, Corollary 3] that

$$\lim_{n \rightarrow \infty} H_{mn} \left[f \left(\frac{(mn+1)t}{m(1+t)} \right); \frac{x}{n} \right] = B_m^{[0]}(f(t); x). \quad (7.3)$$

Here is the proof based on Theorem 2.1. We have to prove (7.3) for $f(t) = e^{ist}$, $s \in \mathbb{R}$.

$$\begin{aligned} & \lim_{n \rightarrow \infty} H_{mn} \left[\exp \left(is \frac{(mn+1)t}{m(1+t)} \right); \frac{x}{n} \right] \\ &= \lim_{n \rightarrow \infty} \left(1 + \frac{x}{n} \right)^{-mn} \sum_{k=0}^{mn} \binom{mn}{k} \left(\frac{x}{n} \right)^k \exp \left(is \frac{(mn+1) \frac{k}{mn-k+1}}{m \left(1 + \frac{k}{mn-k+1} \right)} \right) \\ &= \lim_{n \rightarrow \infty} \left(1 + \frac{x}{n} \right)^{-mn} \sum_{k=0}^{mn} \binom{mn}{k} \left(\frac{x}{n} \right)^k \left(e^{is/m} \right)^k \\ &= \lim_{n \rightarrow \infty} \left(1 + \frac{x}{n} \right)^{-mn} \left(1 + \frac{x}{n} e^{is/m} \right)^{mn} = e^{-mn} e^{mxe^{is/m}} \\ &= e^{mx(e^{is/m}-1)} = B_m^{[0]}(e^{ist}; x). \end{aligned}$$

This ends the proof.

Concerning this example see also [6, Remark 2, p.496], where it is proved that

$$\left| H_{mn} \left(f \left(\frac{(mn+1)t}{m(1+t)} \right); \frac{x}{n} \right) - H_{mn} \left(f \left(\frac{nt}{1+t} \right); \frac{x}{n} \right) \right| \leq 2\omega \left(f; \frac{x}{mn} \right),$$

ω being the first modulus of continuity.

8. Meyer-König and Zeller operators

The operators of Meyer-König and Zeller [31] in the slight modification of Cheney and Sharma [16] are defined for $f \in C[0, 1]$ as follows

$$M_n(f; x) = \begin{cases} \sum_{k=0}^{\infty} \binom{n+k}{k} x^k (1-x)^{n+1} f \left(\frac{k}{n+k} \right), & x \in [0, 1), \\ f(1), & x = 1. \end{cases} \quad (8.1)$$

8.1. Modified Meyer-König and Zeller operators converging to Szász-Mirakyan operator

Let us remark that

$$M_n \left(f \left(\frac{t}{1-t} \right); x \right) = (1-x)^{n+1} \sum_{k=0}^{\infty} f \left(\frac{k}{n} \right) \binom{n+k}{k} x^k,$$

and consequently. In particular

$$\begin{aligned} M_{mn} \left(f \left(\frac{nt}{1-t} \right); \frac{x}{n} \right) &= \left(1 - \frac{x}{n} \right)^{mn+1} \sum_{k=0}^{\infty} \binom{mn+k}{k} \left(\frac{x}{n} \right)^k f \left(\frac{n \frac{k}{mn+k}}{1 - \frac{k}{mn+k}} \right) \\ &= \left(1 - \frac{x}{n} \right)^{mn+1} \sum_{k=0}^{\infty} \binom{mn+k}{k} \left(\frac{x}{n} \right)^k f \left(\frac{k}{m} \right), \end{aligned}$$

and so,

$$\begin{aligned}\lim_{n \rightarrow \infty} M_{mn} \left(e^{is \frac{nt}{1-t}}; \frac{x}{n} \right) &= \lim_{n \rightarrow \infty} \left(1 - \frac{x}{n} \right)^{mn+1} \sum_{k=0}^{\infty} \binom{mn+k}{k} \left(\frac{x}{n} \right)^k \left(e^{is/m} \right)^k \\ &= \lim_{n \rightarrow \infty} \left(1 - \frac{x}{n} \right)^{mn+1} \frac{1}{\left(1 - \frac{x}{n} e^{is/m} \right)^{mn}} = e^{-mx} e^{mx e^{is/m}}.\end{aligned}$$

We see that

$$\lim_{n \rightarrow \infty} M_{mn} \left(e^{is \frac{nt}{1-t}}; \frac{x}{n} \right) = e^{mx(e^{is/m}-1)} = B_m^{[0]}(e^{ist}; x),$$

and therefore,

$$\lim_{n \rightarrow \infty} M_{mn} \left(f \left(\frac{nt}{1-t} \right); \frac{x}{n} \right) = B_m^{[0]}(f(t); x).$$

In a similar way it can be proved that

$$\lim_{n \rightarrow \infty} M_{mn} \left[f \left(\frac{(mn+1)t}{m(1-t)} \right); \frac{x}{n} \right] = B_m^{[0]}(f(t); x).$$

8.2. Modified Meyer-König and Zeller operators converging to Gamma operator

Here we consider another modification of Meyer-König and Zeller operators, namely

$$\begin{aligned}M_m \left[f \left(\frac{t}{n(1-t)} \right); \frac{nx}{1+nx} \right] &= \left(1 - \frac{nx}{1+nx} \right)^{m+1} \sum_{k=0}^{\infty} f \left(\frac{\frac{k}{m+k}}{n \left(1 - \frac{k}{m+k} \right)} \right) \binom{m+k}{k} \left(\frac{nx}{1+nx} \right)^k \\ &= \frac{1}{(1+nx)^{m+1}} \sum_{k=0}^{\infty} f \left(\frac{k}{mn} \right) \binom{m+k}{k} \left(\frac{nx}{1+nx} \right)^k.\end{aligned}$$

In particular,

$$\begin{aligned}\lim_{n \rightarrow \infty} M_m \left[e^{is \frac{t}{n(1-t)}}; \frac{nx}{1+nx} \right] &= \lim_{n \rightarrow \infty} \frac{1}{(1+nx)^{m+1}} \sum_{k=0}^{\infty} \binom{m+k}{k} \left(\frac{nx}{1+nx} e^{is/mn} \right)^k \\ &= \lim_{n \rightarrow \infty} \frac{1}{(1+nx)^{m+1}} \frac{1}{\left(1 - \frac{nx}{1+nx} e^{is/mn} \right)^{m+1}} \\ &= \lim_{n \rightarrow \infty} \frac{1}{[1+nx(1-e^{is/mn})]^{m+1}} = \frac{1}{\left(1 - \frac{isx}{m} \right)^{m+1}}.\end{aligned}$$

Using (10.2) we get

$$\lim_{n \rightarrow \infty} M_m \left[f \left(\frac{t}{n(1-t)} \right); \frac{nx}{1+nx} \right] = G_{m+1} \left(f(t); \frac{m+1}{m} x \right). \quad (8.2)$$

9. Convergence of Lototsky-Schnabl operators

Let $\lambda : [0, 1] \rightarrow [0, 1]$ be continuous, $\lambda(0) = 1$. For $f \in C[0, 1]$, $a \in [0, 1]$, $x \in [0, 1]$, we consider the function

$$f_{x,a}(t) := f(at + (1-a)x), \quad t \in [0, 1].$$

Using the Bernstein operators $(B_r^{[-1]})_{r \geq 1}$ one defines the Lototsky-Schnabl operators $A_{n,\lambda} : C[0, 1] \rightarrow C[0, 1]$,

$$A_{n,\lambda}(f(t); x) := \sum_{r=0}^n \binom{n}{r} \lambda(x)^r (1 - \lambda(x))^{n-r} B_r^{[-1]}(f_{x, \frac{x}{n}}(t); x).$$

For details see [8].

Theorem 9.1. *Given $f \in C_b[0, \infty)$, $x \in [0, \infty)$, $m \geq 1$, we have*

$$\lim_{n \rightarrow \infty} A_{mn,\lambda}\left(f(nt); \frac{x}{n}\right) = B_m^{[0]}(f(t); x), \quad (9.1)$$

where $B_m^{[0]}$ is the classical Szász-Mirakjan operator.

Proof. Let $L_n(f(t); x) := A_{mn,\lambda}\left(f(nt); \frac{x}{n}\right)$. Remark that

$$f_{\frac{x}{n}, \frac{r}{mn}}(nt) = f\left(\frac{rt}{m} + \left(1 - \frac{r}{mn}\right) \frac{x}{n}\right)$$

and

$$B_r^{[-1]}\left(f_{\frac{x}{n}, \frac{r}{mn}}(nt); \frac{x}{n}\right) = \sum_{k=0}^r \binom{r}{k} \left(\frac{x}{n}\right)^k \left(1 - \frac{x}{n}\right)^{r-k} f\left(\frac{k}{m} + \left(1 - \frac{r}{mn}\right) \frac{x}{n}\right).$$

With $f(t) := e^{ist}$ we have

$$B_r^{[-1]}\left(f_{\frac{x}{n}, \frac{r}{mn}}(nt); \frac{x}{n}\right) = \left(1 - \frac{x}{n} + \frac{x}{n} e^{is/m}\right)^r e^{is(1 - \frac{r}{mn}) \frac{x}{n}}.$$

Therefore,

$$\begin{aligned} L_n(e^{ist}; x) &= \sum_{r=0}^{mn} \binom{mn}{r} \lambda\left(\frac{x}{n}\right)^r \left(1 - \lambda\left(\frac{x}{n}\right)\right)^{mn-r} \left(1 - \frac{x}{n} + \frac{x}{n} e^{is/m}\right)^r e^{is(1 - \frac{r}{mn}) \frac{x}{n}} \\ &= e^{isx/n} \left[1 - \lambda\left(\frac{x}{n}\right) + \lambda\left(\frac{x}{n}\right) \left(1 + \frac{x}{n} (e^{is/m} - 1)\right)\right]^{mn} e^{-isx/mn^2}. \end{aligned}$$

By straightforward calculations we get

$$\lim_{n \rightarrow \infty} L_n(e^{ist}; x) = e^{mx(e^{is/m} - 1)} = B_m^{[0]}(e^{ist}; x).$$

Now Theorem 2.1 shows that

$$\lim_{n \rightarrow \infty} A_{mn,\lambda}\left(f(nt); \frac{x}{n}\right) = \lim_{n \rightarrow \infty} L_n(f(t); x) = B_m^{[0]}(f(t); x),$$

and this concludes the proof. \square

Remark 9.1. *An estimate of the rate of convergence in (9.1) can be found in [8, Theorem 4].*

10. Convergence toward Gamma type operators

Consider the Gamma operator

$$G_m(f(t); x) := \frac{x^{-m}}{(m-1)!} \int_0^\infty f\left(\frac{t}{m}\right) t^{m-1} e^{-t/x} du, \quad (10.1)$$

where $f \in C_b[0, \infty)$, $x \in [0, \infty)$. (If $x = 0$, $G_m(f(t); 0) = f(0)$).

It is not difficult to prove that

$$G_m(e^{ist}; x) = \left(1 - \frac{isx}{m}\right)^{-m}, \quad s \in \mathbb{R} \quad (10.2)$$

As in Section 3, it can be proved that

$$V_m^{[c](k)}(e^{ist}; x) = \left(\frac{m}{is}(e^{is/m} - 1)\right)^k \left(1 - cx(e^{is/m} - 1)\right)^{-\frac{m}{c}-k},$$

and so, for $c > 0$,

$$\begin{aligned} \lim_{n \rightarrow \infty} V_m^{[c](k)}(e^{ist/n}; nx) &= \lim_{n \rightarrow \infty} \left(\frac{mn}{is}(e^{\frac{is}{mn}} - 1)\right)^k \left(1 - cnx(e^{\frac{is}{mn}} - 1)\right)^{-\frac{m}{c}-k} \\ &= \left(1 - \frac{iscx}{m}\right)^{-\frac{m}{c}-k} = G_{\frac{m}{c}+k}\left(e^{ist}; \frac{m+ck}{m}x\right). \end{aligned}$$

Now from Theorem 2.1 we deduce

Theorem 10.1. *With the above notation we have*

$$\lim_{n \rightarrow \infty} V_m^{[c](k)}\left(f\left(\frac{t}{n}\right); nx\right) = G_{\frac{m}{c}+k}\left(f(t); \frac{m+ck}{m}x\right). \quad (10.3)$$

For $k = 0$ and $c = 1$, this is Theorem 2 d) in [13].

11. Convergence toward Weierstrass operator

Let $a > 0$. The Weierstrass operator is defined by (see [9])

$$W_a(f(t); x) := \frac{1}{\sqrt{2\pi a}} \int_{\mathbb{R}} f(t) e^{-\frac{(t-x)^2}{2a}} dt, \quad f \in C_b[0, \infty), x \geq 0.$$

The following results were briefly presented in [37, (24.25)] and [34]. Here we give detailed proofs based on Theorem 2.1.

Theorem 11.1. *Let $0 < u < 1$ be given, and let $b_n > 0$ such that $\lim_{n \rightarrow \infty} nb_n^2 = a$. Then*

$$\lim_{n \rightarrow \infty} B_n^{[-1]}\left(f\left(x + \frac{nb_n}{\sqrt{u(1-u)}}(t-u)\right); u\right) = W_a(f(t); x), \quad (11.1)$$

where $B_n^{[-1]}$, $n \geq 1$, are the classical Bernstein operators.

Proof. Let

$$L_n(f(t); x) := B_n^{[-1]} \left(f \left(x + \frac{nb_n}{\sqrt{u(1-u)}}(t-u) \right); u \right). \quad (11.2)$$

Then, by a straightforward calculation we get

$$\begin{aligned} L_n(e^{ist}; x) &:= B_n^{[-1]} \left(e^{is \left(x + \frac{nb_n}{\sqrt{u(1-u)}}(t-u) \right)}; u \right) \\ &= e^{isx} \sum_{j=0}^n \binom{n}{j} u^j (1-u)^{n-j} e^{is \frac{nb_n}{\sqrt{u(1-u)}} \left(\frac{j}{n} - u \right)} \\ &= e^{ist} \left[(1-u) e^{\frac{-isu b_n}{\sqrt{u(1-u)}}} + u e^{\frac{is(1-u)b_n}{\sqrt{u(1-u)}}} \right] n. \end{aligned}$$

Remark that

$$\lim_{n \rightarrow \infty} L_n(e^{ist}; x) = e^{isx} e^{\lim_{n \rightarrow \infty} \left(\frac{\frac{-isu b_n}{\sqrt{u(1-u)}} + u e^{\frac{is(1-u)b_n}{\sqrt{u(1-u)}}}}{b_n^2} n b_n^2 \right)}.$$

Since $nb_n^2 \rightarrow a$ we have $b_n \rightarrow 0$ and it readily follows that

$$\lim_{n \rightarrow \infty} L_n(e^{ist}; x) = e^{isx - \frac{1}{2}as^2}. \quad (11.3)$$

On the other hand, using the well-known characteristic function of a Gaussian random variable with mean x and variance a , we can write

$$W_a(e^{ist}; x) = \frac{1}{\sqrt{2\pi a}} \int_{\mathbb{R}} e^{ist} e^{-\frac{(t-x)^2}{2a}} dt = e^{isx - \frac{1}{2}as^2}. \quad (11.4)$$

From (11.3), (11.4) and Theorem 2.1 we infer that

$$\lim_{n \rightarrow \infty} L_n(f(t); x) = W_a(f(t); x). \quad (11.5)$$

Now (11.1) is a consequence of (11.2) and (11.5). \square

We consider again a sequence $b_n > 0$, $n \geq 1$, such that $\lim_{n \rightarrow \infty} nb_n^2 = a > 0$. The following Bernstein-Schnabl type operators were introduced in [35] and investigated in [1], [2], [37], [34], [3], [5]:

$$S_{n,b_n}(f(t); x) := \frac{1}{(2nb_n)^n} \int_{x-nb_n}^{x+nb_n} \cdots \int_{x-nb_n}^{x+nb_n} f \left(\frac{u_1 + \cdots + u_n}{n} \right) du_1 \cdots du_n.$$

Theorem 11.2. *Let $f \in C_b[0, \infty)$ and $x \geq 0$. Then*

$$\lim_{n \rightarrow \infty} S_{n,b_n}(f(t); x) = W_{a/3}(f(t); x). \quad (11.6)$$

Proof. Using Theorem 2.1 it suffices to show that

$$\lim_{n \rightarrow \infty} S_{n,b_n}(e^{ist}; x) = W_{a/3}(e^{ist}; x). \quad (11.7)$$

According to the definition we have

$$S_{n,b_n}(e^{ist}; x) = \frac{1}{(2nb_n)^n} \int_{x-nb_n}^{x+nb_n} \dots \int_{x-nb_n}^{x+nb_n} e^{is(u_1+\dots+u_n)/n} du_1 \dots du_n \quad (11.8)$$

$$= \frac{1}{(2nb_n)^n} \left(\int_{x-nb_n}^{x+nb_n} e^{isu/n} du \right)^n = e^{isx} \left(\frac{\sin sb_n}{sb_n} \right)^n. \quad (11.9)$$

Since $nb_n^2 \rightarrow a$ we have $b_n \rightarrow 0$ and therefore

$$\begin{aligned} \lim_{n \rightarrow \infty} S_{n,b_n}(e^{ist}; x) &= e^{isx} \exp \left(\lim_{n \rightarrow \infty} \left(n \frac{\sin sb_n - sb_n}{sb_n} \right) \right) \\ &= e^{isx} \exp \left(s^2 (nb_n^2) \frac{\sin sb_n - sb_n}{(sb_n)^3} \right). \end{aligned}$$

It follows that

$$\lim_{n \rightarrow \infty} S_{n,b_n}(e^{ist}; x) = e^{isx - \frac{1}{6}as^2}. \quad (11.10)$$

Now (11.7) is a consequence of (11.4) and (11.10). Thus (11.6) is proved. \square

12. The limit operator

Proposition 12.1. *Let $(L_n)_{n \geq 1}$ and $(T_m)_{m \geq 1}$ be two sequences of positive linear operators such that*

$$\lim_{n \rightarrow \infty} L_{mn}(f(nt); \frac{x}{n}) = T_m(f(t); x)$$

for all $m \geq 1$, $x \geq 0$, $f \in C_b[0, \infty)$. Then

$$T_{m\nu} \left(f(\nu t); \frac{x}{\nu} \right) = T_m(f(t); x), \quad (12.1)$$

for all $m \geq 1$, $\nu \geq 1$, $x \geq 0$, $f \in C_b[0, \infty)$.

Proof. Let $\nu \geq 1$, $m \geq 0$. Then

$$\begin{aligned} \lim_{n \rightarrow \infty} L_{m(n\nu)} \left(f((n\nu)t); \frac{x}{n\nu} \right) &= T_m(f(t); x), \\ \lim_{n \rightarrow \infty} L_{(m\nu)n} \left(f(n(\nu t)); \frac{x/\nu}{n} \right) &= T_{m\nu} \left(f(\nu t); \frac{x}{\nu} \right) \end{aligned}$$

and so we get (12.1). \square

Theorem 12.1. *Let $0 \leq a_0 < a_1 < \dots$, $\varphi_{m,k} \in C_b[0, \infty)$, $m \in \mathbb{N}$, $k \in \mathbb{N}_0$,*

$$T_m(f(t); x) := \sum_{k=0}^{\infty} \varphi_{m,k}(x) f\left(\frac{a_k}{m}\right), \quad f \in C_b[0, \infty), \quad x \geq 0.$$

Then (12.1) is equivalent to

$$\varphi_{m\nu,k}(x) = \varphi_{m,k}(\nu x), \quad m, \nu \in \mathbb{N}, \quad k \in \mathbb{N}_0, \quad x \geq 0, \quad (12.2)$$

and to

$$\varphi_{m,k}(x) = \varphi_{1,k}(mx), \quad m \in \mathbb{N}, \quad k \in \mathbb{N}_0, \quad x \geq 0. \quad (12.3)$$

Proof. $T_{m\nu} \left(f(\nu t); \frac{x}{\nu} \right) = \sum_{k=0}^{\infty} \varphi_{m\nu,k} \left(\frac{x}{\nu} \right) f \left(\frac{a_k}{m} \right)$, and it readily follows that (12.1) is equivalent to (12.2). Suppose that (12.3) holds true. Then

$$\varphi_{m\nu,k}(x) = \varphi_{1,k}(m\nu x) \text{ and } \varphi_{m,k}(\nu x) = \varphi_{1,k}(m\nu x).$$

This implies (12.2).

Finally, suppose that (12.2) is valid. Using it with $m = 1$ it follows that $\varphi_{\nu,k}(x) = \varphi_{1,k}(\nu x)$ for all $\nu \geq 1$, $k \in \mathbb{N}_0$, $x \geq 0$, and this (12.3). \square

The Jakimovski-Leviatan operators are generalization of Szász-Mirakjan operators (see [29], [4]). To define them we need a sequence of Appell polynomials $(A_n(x))_{n \geq 0}$. If they have the explicit representation

$$A_n(x) = \alpha_n + \binom{n}{1} \alpha_{n-1} x + \binom{n}{2} \alpha_{n-2} x^2 + \cdots + \alpha_0 x^n, \quad n = 0, 1, \dots, \quad (12.4)$$

one associates the power series

$$a(t) = \alpha_0 + \frac{t}{1!} \alpha_1 + \frac{t^2}{2!} \alpha_2 + \cdots + \frac{t^n}{n!} \alpha_n + \cdots, \quad \alpha_0 \neq 0, \quad (12.5)$$

called the generating function.

In fact, the sequence and its generating function are related by

$$a(t)e^{tx} = A_0(x) + \frac{t}{1!} A_1(x) + \frac{t^2}{2!} A_2(x) + \cdots + \frac{t^n}{n!} A_n(x) + \cdots \quad (12.6)$$

For details see, e.g., [18].

The Jakimovski-Leviatan type operators associated with $(A_n(x))_{n \geq 0}$ are defined by

$$\Psi_n f(x) := \frac{e^{-nx}}{a(1)} \sum_{k=0}^{\infty} \frac{1}{k!} A_k(nx) f \left(\frac{k}{n} \right), \quad n \geq 1, \quad (12.7)$$

where $f \in C[0, \infty)$ is a function for which the series is convergent for all $x \in [0, \infty)$.

Remark 12.1. *The Szász-Mirakjan and more generally the Jakimovski-Leviatan operators satisfy the equivalent conditions (12.1), (12.2), (12.3). The Szász-Mirakjan operator appeared as limit operator in previous examples. The next example presents an instance where a Jakimovski-Leviatan operator appears as limit operator.*

Example 12.1. *Let $p \in [0, \infty)$ and $a(t) := e^{pt}$, $t \in \mathbb{R}$. Using (12.6) we find the associated Appell polynomials $A_{k,p}(x) = (x+p)^k$, $k \geq 0$. The Jakimovski-Leviatan operators are*

$$\Psi_{n,p}(f(t); x) = e^{-nx-p} \sum_{k=0}^{\infty} \frac{(nx+p)^k}{k!} f \left(\frac{k}{n} \right), \quad f \in C_b[0, \infty).$$

Let us remark that $\Psi_{n,p}(f(t); x)$ are the Szász-Mirakjan-Schurer operators presented, e.g. in [9, p.338].

Let $B_n^{[1]}$ be the classical Baskakov operators, and

$$L_n(f(t); x) := B_{mn}^{[1]} \left(f(nt); \frac{x+p/m}{n} \right).$$

Then $\lim_{n \rightarrow \infty} L_n(e^{ist}; x) = e^{(mx+p)(e^{is/m}-1)} = \Psi_{m,p}(e^{ist}; x)$ and consequently

$$\lim_{n \rightarrow \infty} B_{mn}^{[1]} \left(f(nt); \frac{x+p/m}{n} \right) = \Psi_{m,p}(f(t); x). \quad (12.8)$$

Example 12.2. Consider the operator

$$L_m^*(f(t); x) := \frac{1}{\cosh(mx)} \sum_{k=0}^{\infty} \frac{(mx)^{2k}}{(2k)!} f\left(\frac{2k}{m}\right), \quad f \in C_b[0, \infty), \quad x \geq 0,$$

introduced by Leśniewicz and Rempulska [30]; see also [17]. It satisfies the equivalent conditions (12.1), (12.2), (12.3).

Now let $\tilde{B}_n(f(t); x) := \sum_{k=0}^n \binom{n}{k} (-x)^k (1-x)^{n-k} f\left(\frac{k}{n}\right)$, $f \in C[0, 1]$, $x \in [0, 1]$, and

$$L_n(f(t); x) := \frac{B_n^{[-1]}(f(t); x) + \tilde{B}_n(f(t); x)}{1 + (1-2x)^n},$$

where

$$L_{2p+1}(f(t); 1) := \lim_{x \rightarrow 1} L_{2p+1}(f(t); x) = f\left(\frac{2p}{2p+1}\right).$$

Then L_n reproduces the constant functions and

$$\lim_{n \rightarrow \infty} L_{mn} \left(e^{isnt}; \frac{x}{n} \right) = \frac{\cosh(mxe^{is/m})}{\cosh(mx)} = L_m^*(e^{ist}; x).$$

Consequently,

$$\lim_{n \rightarrow \infty} L_{mn} \left(f(nt); \frac{x}{n} \right) = L_m^*(f(t); x), \quad f \in C_b[0, \infty). \quad (12.9)$$

Proposition 12.2. Let $(L_m)_{m \geq 1}$ and $(Z_m)_{m \geq 1}$ be two sequences of positive linear operators such that

$$\lim_{n \rightarrow \infty} L_m \left(f\left(\frac{t}{n}\right); nx \right) = Z_m(f(t), x)$$

for all $m \geq 1$, $x \geq 0$, $f \in C_b[0, \infty)$. Then

$$Z_m \left(f\left(\frac{t}{\nu}\right); \nu x \right) = Z_m(f(t), x), \quad \nu \in \mathbb{N}, \quad x \geq 0. \quad (12.10)$$

Proof. We have

$$\lim_{n \rightarrow \infty} L_m \left(f\left(\frac{t/\nu}{n}\right); n(\nu x) \right) = Z_m \left(f\left(\frac{t}{\nu}\right); \nu x \right),$$

and

$$\lim_{n \rightarrow \infty} L_m \left(f\left(\frac{t}{n\nu}\right); (n\nu)x \right) = Z_m(f(t), x).$$

Then leads to (12.10). \square

Theorem 12.2. Let Z_m be an integral operator of the form

$$Z_m(f(t); x) := \int_0^\infty f(t)K_m(t, x)dt, \quad f \in C_b[0, \infty), \quad x \geq 0,$$

with a suitable continuous kernel $K_m(t, x)$. Then (12.10) is equivalent to

$$K_m(\nu t, \nu x) = \frac{1}{\nu} K_m(t, x), \quad \nu > 0, \quad t, x \geq 0. \quad (12.11)$$

Proof. We have

$$\begin{aligned} Z_m\left(f\left(\frac{t}{\nu}\right); \nu x\right) &= \int_0^\infty f\left(\frac{t}{\nu}\right) K_m(t, \nu x) dt \\ &= \nu \int_0^\infty f(t) K_m(\nu t, \nu x) dt. \end{aligned}$$

So (12.10) becomes

$$\int_0^\infty f(t) K_m(\nu t, \nu x) dt = \frac{1}{\nu} \int_0^\infty f(t) K_m(t, x) dt,$$

for all $f \in C_b[0, \infty)$, $\nu > 0$, $x > 0$. This is equivalent to (12.11) and the proof is finished. \square

Example 12.3. For the Gamma operator the kernel is

$$K_m(t, x) := \frac{x^{-m}}{(m-1)!} m^m t^{m-1} e^{-mt/x},$$

and it satisfies (12.11).

Example 12.4. Let R_m be an operator of the form

$$R_m(f(t); x) := \sum_{k=0}^\infty \varphi_{m,k}(x) \frac{\int_0^\infty \varphi_{m,k}(t) f(t) dt}{\int_0^\infty \varphi_{m,k}(t) dt}.$$

If the functions $\varphi_{m,k}$ satisfy (12.2), then the operators R_m satisfy (12.1).

Problem 12.1. Is the converse true?

Comment 1. Suppose that $(R_m)_{m \geq 1}$ satisfies (12.1) for all $m \geq 1$, $\nu \geq 1$, $x \geq 0$, $f \in C_b[0, \infty)$. With $m = 1$ we get $R_\nu\left(f(\nu t); \frac{x}{\nu}\right) = R_1(f(t); x)$. With an obvious change of notation we have $R_m(f(s); y) = R_1\left(f\left(\frac{s}{\nu}\right); \nu y\right)$. Even better,

$$R_m(f(t); x) = R_1\left(f\left(\frac{t}{m}\right); mx\right), \quad \text{for all } m \geq 1, \quad x \geq 0. \quad (12.12)$$

Therefore each R_m , $m \geq 2$, is completely determined by R_1 .

A natural hypothesis is $\int_0^\infty \varphi_{m,k}(x) dx = 1/m$, for all $m \geq 1$, $k \geq 0$. Then (12.12) becomes

$$\sum_{k=0}^\infty \varphi_{m,k}(x) \int_0^\infty \varphi_{m,k}(t) f(t) dt = \sum_{k=0}^\infty \varphi_{1,k}(mx) \int_0^\infty \varphi_{1,k}(mt) f(t) dt, \quad (12.13)$$

for all $m \geq 1$, $x \geq 0$, $f \in C_b[0, \infty)$.

In this context Problem 12.1 is reduced to: does (12.13) imply (12.3)?

Generally speaking, if a Durrmeyer operator is represented as

$$\sum_{k=0}^{\infty} u_k(x) \int_0^{\infty} u_k(t) f(t) dt = \sum_{k=0}^{\infty} v_k(x) \int_0^1 v_k(t) f(t) dt,$$

for all $x \geq 0$ and $f \in C_b[0, \infty)$, does it follow that $u_k = v_k$, $k \geq 0$?

Proposition 12.3. *The following two statements are equivalent*

$$(1) R_{m\nu} \left(f(\nu t); \frac{x}{\nu} \right) = R_m(f(t); x), \quad \forall m, \nu \geq 1, x \geq 0, f \in C_b[0, \infty).$$

$$(2) R_m(f(t); x) = R_1 \left(f \left(\frac{t}{m} \right); mx \right), \quad \forall m \geq 1, x \geq 0, f \in C_b[0, \infty).$$

Proof. (1) \Rightarrow (2). Take $m = 1$. Then $R_{\nu} \left(f(\nu t); \frac{x}{\nu} \right) = R_1(f(t); x)$. Set $s = \nu t$, $y = \frac{x}{\nu}$.

Then $R_{\nu}(f(s); y) = R_1 \left(f \left(\frac{s}{\nu} \right); \nu y \right)$, which is (2).

(2) \Rightarrow (1). Using (2) we have

$$\begin{aligned} R_{m\nu} \left(f(\nu t); \frac{x}{\nu} \right) &= R_1 \left(f \left(\frac{\nu t}{m\nu} \right); m\nu \frac{x}{\nu} \right) \\ &= R_1 \left(f \left(\frac{t}{m} \right); mx \right) = R_m(f(t); x), \end{aligned}$$

and (1) is proved. □

Example 12.5. *We try to determine a convolution operator $U(f(t); x) = \int_{\mathbb{R}} f(t) \varphi(x - t) dt$ satisfying (12.10) for all $\nu \in (0, \infty)$.*

$$U \left(f \left(\frac{t}{\nu} \right); \nu x \right) = \int_{\mathbb{R}} f \left(\frac{t}{\nu} \right) \varphi(\nu x - t) dt = \int_{\mathbb{R}} f(s) \varphi(\nu x - \nu s) \nu ds$$

and so $\varphi(\nu(x - s)) = \frac{1}{\nu} \varphi(x - s)$, i.e., $\varphi(\nu z) = \frac{1}{\nu} \varphi(z)$.

Taking $\varphi(z) := \frac{1}{\pi z}$ we see that Uf is the convolution of $f(t)$ and $\frac{1}{\pi t}$, i.e., the Hilbert transform of f . But, of course, this is not a positive linear operator.

We end this section with a list with the limit operators presented in the previous sections, together with an equation where they appear: $B_m^{[0]}$ (1.3), $V_m^{[0](k)}$ (4.4), $D_{m,\rho}^{[0](k)}$ (5.3), $V_m^{[0](k)} \left((f \circ \sigma)(t); \frac{\psi(x)}{n} \right)$ (6.1), $G_{\frac{m}{n}+k}$ (10.3), W_a (11.1), $W_{a/3}$ (11.6), $\Psi_{m,p}$ (12.8), L_m^* (12.9), G_{m+1} (8.2).

Funding. This work has been supported by the University of Wuppertal.

Conflicts of interest. The authors declare no competing financial interests.

Availability of data and material. No data were used to support this study.

References

- [1] U. Abel, M. Ivan, Asymptotic approximation with a sequence of positive linear operators, *J. Comput. Anal. Appl.* 3(4) (2001), 331-341.
- [2] U. Abel, M. Ivan, Asymptotic expansion of a sequence of divided differences with application to positive linear operators, *J. Comput. Anal. Appl.* 7(1) (2005), 89-101.
- [3] A.M. Acu, G. Başcanbaz-Tunca, I. Rasa, Information potential for some probability density functions, *Appl. Math. Comput.* 389 (2021), 125578.
- [4] A.M. Acu, I.C. Buscu, I. Rasa, A sequence of Appell polynomials and the associated Jakimovski-Leviatan operators. *Anal. Math. Phys.* 11, 88 (2021).
- [5] A.M. Acu, M. Dancs, M. Heilmann, V. Paşca, I. Raşa, Voronovskaya type results for special sequences of operators, *RACSAM* 116, 19 (2022).
- [6] J.A. Adell, J. de la Cal, Limiting properties of Inverse Beta and generalized Bleimann-Butzer-Hahn operators, *Math. Proc. Camb. Phil. Soc.* (1993), 114-489.
- [7] J.A. Adell, F.G. Badia, J. de la Cal, On the Iterates of Some Bernstein-Type Operators, *J. Math. Anal. Appl.* 209(2) (1997), 529-541.
- [8] J.A. Adell, J. de la Cal, I. Rasa, Lototsky-Schabl operators on the unit interval, *Rend. Circ. Mat. Palermo*, 48(3)(1999), 517-536
- [9] F. Altomare, M. Campiti, Korovkin-type Approximation Theory and its Applications, Series: De Gruyter Studies in Mathematics, 17, 1994.
- [10] K. Baumann, M. Heilmann, I. Raşa, Further results for kth order Kantorovich modification of linking Baskakov type operators. *Results Math.* 69(3) (2016), 297–315.
- [11] V.A. Baskakov, A sequence of linear positive operators in the space of continuous functions, *Dokl. Acad. Nauk. SSSR* 113 (1957), 249-251.
- [12] G. Bleimann, P.L. Butzer, L. Hahn, A Bernstein-type operator approximating continuous functions of the semi-axis, *Indag. Math.* 42 (1980), 255–262.
- [13] J. de la Cal, F. Luquin, A note on limiting properties of some Bernstein-type operators, *J. Approx. Theory*, 68(3) (1992), 322-329.
- [14] J. de la Cal, On Stancu-Mühlbach operators and some connected problems concerning probability distribution, *J. Aprox. Theory*, 74(1) (1993), 59-68.
- [15] J. de la Cal, F. Luquin, Approximation Szász and Gamma operators by Baskakov operators, *J. Math. Anal. Appl.*, 184 (1994), 585-593.

- [16] E.W. Cheney, A. Sharma, Bernstein power series, *Canad. J. Math.* 16(1964), 241-253.
- [17] A. Ciupa, A Voronovskaya-type theorem for a positive linear operator, *International Journal of Mathematics and Mathematical Sciences*, Volume 2006, Article ID 42368, 1–7.
- [18] F.A. Costabile, E. Longo, A determinantal approach to Appell polynomials, *J. Comput. Appl. Math.*, 234 (5) (2010), 1528-1542.
- [19] W. Feller, *An Introduction to Probability Theory and Its Applications*, Vol. I, 2nd ed., Wiley, New York, 1951.
- [20] H. Gonska, R. Păltănea, Simultaneous approximation by a class of Bernstein–Durrmeyer operators preserving linear functions. *Czechoslovak Math. J.* 60(135) (2010), 783–799.
- [21] V.Gupta, M.Th. Rassias, *Moments of Linear Positive Operators and Approximation*, Springer, Cham, 2019.
- [22] M Heilmann, Commutativity and spectral properties of genuine Baskakov-Durrmeyer type operators and their th order Kantorovich modification, *J. Numer. Anal. Approx. Theory*, 44 (4) (2015), 166–179.
- [23] M. Heilmann, F. Nasaireh, I. Raşa, I. Discrete operators associated with linking operators. *Anal. Math. Phys.* 10(72) (2020).
- [24] M. Heilmann, I. Raşa, k-th order Kantorovich modification of linking Baskakov type operators, in P.N. Agrawal et al. (ed.) *Recent Trends in Mathematical Analysis and its Applications*, Proceedings of the Conference ICRTMAA 2014, Roorkee, India, December 2014, *Proceedings in Mathematics & Statistics*, 143, 229–242 (2015).
- [25] M. Heilmann, I. Raşa, A nice representation for a link between Bernstein-Durrmeyer and Kantorovich operators, in *Proceedings of ICMC 2017*, Haldia, India, January 2017, (ed. D. Giri et. al.), *Springer Proceedings in Mathematics and Computing*, (2017), 312-320.
- [26] M. Heilmann, I. Raşa, A Nice Representation for a Link between Baskakov- and Szász-Mirakjan-Durrmeyer Operators and their Kantorovich Variants, *Results Math* (2019) 74: 9.
- [27] M. Heilmann, I. Raşa, Note on a proof for the representation of the kth order Kantorovich modification of linking Baskakov type operators, In: Deo N., Gupta V., Acu A., Agrawal P. (eds) *Mathematical Analysis I: Approximation Theory*. ICRAPAM 2018. Springer Proceedings in Mathematics & Statistics, vol 306. Springer, Singapore, 2020, 89-93.
- [28] M. Heilmann, I. Raşa, A nice representation for a link between Baskakov-and Szasz-Mirakjan-Durrmeyer operators and their Kantorovich variants, *Results Math.* (2019), 74:9.
- [29] A. Jakimovski, D. Leviatan, Generalized Szász operators for the approximation in the infinite interval. *Mathematica (Cluj)* 34(1969), 97-103.
- [30] M. Leśniewicz, L Rempulska, Approximation by some operators of the Szasz-Mirakjan type in exponential weight spaces, *Glasnik Matematički*, 32(52) (1997), 57-69.
- [31] W. Meyer-Konig and K. Zeller, Bernsteinsche Potenzreihen. *Studia Math.* 19 (1960), 89-94.

- [32] V. Miheşan, Uniform approximation with positive linear operators generated by generalized Baskakov method, *Automat. Comput. Appl. Math.* 7 (1) (1998), 34-37.
- [33] R. Păltănea, Simultaneous approximation by a class of Szász–Mirakjan operators. *J. Appl. Funct. Anal.* 9(3–4) (2014), 356–368.
- [34] D. Popa, I. Raşa, Steklov averages as positive linear operators, *Filomat* 30:5 (2016), 1195–1201.
- [35] I. Raşa, Korovkin approximation and parabolic functions, *Conf. Sem. Mat. Univ. Bari.* 236 (1991).
- [36] A. N. Shiriyayev, *Probability*, Springer, New York, 1984.
- [37] T. Vladislav, I. Raşa, *Analiză Numerică. Aproximare, Problema lui Cauchy Abstractă*, Proiectori Altomare, Editura Tehnică, Bucureşti, 1999. (in Romanian).