

Hamiltonian cycles of balanced hypercube with more faulty edges*

Ting Lan and Huazhong Lü[†]

School of Mathematical Sciences,
University of Electronic Science and Technology of China,
Chengdu, Sichuan, 610054, PR China
lvhz@uestc.edu.cn

Abstract

The balanced hypercube BH_n , a variant of the hypercube, is a novel interconnection network for massive parallel systems. It is known that the balanced hypercube remains Hamiltonian after deleting at most $4n - 5$ faulty edges if each vertex is incident with at least two edges in the resulting graph for all $n \geq 2$. In this paper, we show that there exists a fault-free Hamiltonian cycle in BH_n for $n \geq 2$ with $|F| < 5n - 7$ if the degree of every vertex in $BH_n - F$ is at least two and there exists no f_4 -cycles in $BH_n - F$, which improves some known results.

Key words: Interconnection network; Balanced hypercubes; Hamiltonian cycle; Fault-tolerance

1. Introduction

In parallel and distributed systems, interconnection network plays an important role in their performance and specific functions. The topological structure of an interconnection network is usually modeled by a simple undirected graph, where vertices represent processors and edges represent communication link between processors. The n -dimensional hypercube Q_n is one of the most efficient interconnection network models, which has numerous good properties such as high symmetry, large connectivity and recursive structure. The balanced hypercube, BH_n , is a variant of the hypercube, which not only retains most of the excellent properties of Q_n ,

*This research was partially supported by the National Natural Science Foundation of China (No. 11801061).

[†]Corresponding author.

but also possesses some outstanding features that Q_n does not have. For instance, the diameter of BH_n is less than the hypercube with the same number of vertices for odd n . Moreover, for each vertex of BH_n , there is a unique vertex (backup vertex) with the same neighborhood, implying that if a vertex fails then tasks running on it can be easily transferred to its backup vertex without modifying routing. With such good properties, BH_n has received much attention. Embedding properties, especially Hamiltonian path and cycle embedding of BH_n have been studied in [1, 2, 4, 8, 12, 21, 23, 26]. Reliability indices under different theoretical model of BH_n have been determined in [11, 22, 24]. Some other topological issues of BH_n have been investigated in [13, 19, 25].

A graph G is called *Hamiltonian* if there is a Hamiltonian cycle in G . A bipartite graph $G = (X \cup Y, E)$ is *Hamiltonian laceable* if there is a Hamiltonian path between any two vertices x and y in different partite sets. When designing and selecting the desired topology for an interconnected network, the Hamiltonian property is an important indicator for evaluating network performance [7]. Hamiltonian properties of the hypercube and its variants, as well as other famous networks have been widely studied [3, 5, 6, 15, 16].

In practical application, fault is inevitable in a large network, so it is necessary to consider fault-tolerance. A graph G is *k edge fault-tolerance Hamiltonian* if $G - F$ remains Hamiltonian for any $F \subset E(G)$ with $|F| \leq k$. And a bipartite graph G is *k edge fault-tolerance Hamiltonian laceable* if $G - F$ remains Hamiltonian laceable for any $F \subset E(G)$ with $|F| \leq k$. In [26], Zhou et al. proved that BH_n is $2n - 2$ edge fault-tolerant Hamiltonian laceable for $n \geq 2$. Recently, Li et al. [9] further showed that there exists a Hamiltonian cycle in $BH_n - F$ for $n \geq 2$ with $|F| \leq 4n - 5$ if the degree of every vertex in $BH_n - F$ is at least two. They also showed that the upper bound $4n - 5$ of the number of faulty edges can be tolerated is optimal. Actually, if a set F of $4n - 4$ edges is deleted from BH_n , then there exists no Hamiltonian cycles as a cycle of length four with a pair of nonadjacent vertices whose degrees are both two may occur in $BH_n - F$ (see Fig. 1). For convenience, we call such a cycle an *f_4 -cycle*. Thus, a natural question arises: what is the upper bound of the number of faulty edges can be tolerated if f_4 -cycle is forbidden in $BH_n - F$? Motivated by this, Liu and Wang [10] showed that the hypercube still has a Hamiltonian cycle in $Q_n - F$ if $\delta(Q_n - F) \geq 2$ ($|F| \leq 3n - 8$) and there exists no f_4 -cycles.

In this paper, we show that there exists a fault-free Hamiltonian cycle in BH_n for $n \geq 2$ with $|F| \leq 5n - 7$ if the degree of every vertex in $BH_n - F$ is at least two and there exists no f_4 -cycles in $BH_n - F$.

The rest of this paper is organized as follows. In Section 2, we give two equivalent definitions of balanced hypercubes and some useful properties of BH_n . In Section 3, we present the main result (Theorem 8) of this paper. Finally, we conclude this paper in Section 4.




Fig. 1. f_4 -cycle.

2. Some definitions and lemmas

Let $G = (V, E)$ be an undirected simple graph, where V is the vertex set of G and E is the edge set of G . We write $e = uv$ for an edge e with endpoints u and v . Let $N_G(u)$ be the set of vertices adjacent to a vertex u in G and $e(u)$ be the set of edges incident with u . The degree of u , denoted by $d(u)$, is the number of edges incident with u . Let $\delta(G)$ be the minimum degree of G . A graph G is bipartite if $V(G)$ is the union of two disjoint sets X, Y such that every edge has one endpoint in X and the other in Y . A path P from v_0 to v_n is a sequence of distinct vertices $v_0 v_1 \dots v_n$ that consecutive vertices are adjacent, denoted by $\langle v_0, P, v_n \rangle$. A cycle C is a closed path $\langle v_0, v_1, \dots, v_n, v_0 \rangle$. A *Hamiltonian cycle* (resp. path) in a graph is a cycle (resp. path) that passes through every vertex of the graph.

In the following, Wu and Huang [19] gave the definitions of BH_n .

Definition 1. An n -dimensional balanced hypercube BH_n consists of 2^{2n} vertices $(a_0, \dots, a_{i-1}, a_i, a_{i+1}, \dots, a_{n-1})$, where $a_i \in \{0, 1, 2, 3\} (0 \leq i \leq n-1)$. An arbitrary vertex $(a_0, \dots, a_{i-1}, a_i, a_{i+1}, \dots, a_{n-1})$ in BH_n has the following $2n$ neighbors:

- (1). $((a_0 + 1) \bmod 4, a_1, \dots, a_{i-1}, a_i, a_{i+1}, \dots, a_{n-1})$,
 $((a_0 - 1) \bmod 4, a_1, \dots, a_{i-1}, a_i, a_{i+1}, \dots, a_{n-1})$, and
- (2). $((a_0 + 1) \bmod 4, a_1, \dots, a_{i-1}, (a_i + (-1)^{a_0}) \bmod 4, a_{i+1}, \dots, a_{n-1})$,
 $((a_0 - 1) \bmod 4, a_1, \dots, a_{i-1}, (a_i + (-1)^{a_0}) \bmod 4, a_{i+1}, \dots, a_{n-1})$.

In BH_n , the first coordinate a_0 of $(a_0, a_1, \dots, a_{n-1})$ is called *inner index*, and other coordinate $a_i (1 \leq i \leq n-1)$ *i-dimensional index*. Let uv be an edge of BH_n . If u and v differ only the inner index, then uv is said to be a *0-dimensional edge*. If u and v differ not only the inner index, but also i -dimensional index, then uv is called an *i-dimensional edge*.

Definition 2. An n -dimensional balanced hypercube BH_n can be recursively constructed as follows:




Fig. 2. BH_1 .




Fig. 3. BH_2 .

- (1). BH_1 is a cycle of length four whose vertices are labeled clockwise as 0, 1, 2, 3, respectively.
- (2). BH_n is constructed from four disjoint BH_{n-1} s. These four BH_{n-1} s are labeled by BH_{n-1}^0 , BH_{n-1}^1 , BH_{n-1}^2 , BH_{n-1}^3 , where each vertex in BH_{n-1}^j ($0 \leq j \leq 3$) has j attached as the new n -dimensional index. Each vertex $(a_0, a_1, \dots, a_{n-2}, j)$ has two extra neighbors:
 - (a). $((a_0+1) \bmod 4, a_1, \dots, a_{n-2}, (j+1) \bmod 4)$ and $((a_0-1) \bmod 4, a_1, \dots, a_{n-2}, (j+1) \bmod 4)$, which are in BH_{n-1}^{j+1} if a_0 is even;
 - (b). $((a_0+1) \bmod 4, a_1, \dots, a_{n-2}, (j-1) \bmod 4)$ and $((a_0-1) \bmod 4, a_1, \dots, a_{n-2}, (j-1) \bmod 4)$, which are in BH_{n-1}^{j-1} if a_0 is odd.

We illustrate BH_1 and BH_2 in Figs. 2 and 3, respectively.

Let F be an edge subset and D_i be the set of i -dimensional edges of BH_n , respectively, and let $F_i = F \cap D_i$ for $i = 0, 1, 2, \dots, n-1$. Then, BH_n can be divided into four subcubes $BH_{n-1}^{j,i} \cong BH_{n-1}$ by deleting D_i , where $j = 0, 1, 2, 3$ and $i \in \{0, 1, 2, \dots, n-1\}$. We use BH_{n-1}^j to denote $BH_{n-1}^{j,n-1}$ for $j = 0, 1, 2, 3$ and let $F^j = F \cap E(BH_{n-1}^j)$. For simplicity, we assume that $|F^0| = \max \{|F^j| \mid j = 0, 1, 2, 3\}$.

In the following, we make an convention that the vertices with superscript j belong to $BH_{n-1}^{j,i}$, say $a^j, b^j, c^j, \dots \in V(BH_{n-1}^{j,i})$. An r -edge is an edge in $BH_{n-1}^{j,i}$ whose two endpoints are both incident with at least one i -dimensional nonfaulty edge, and if an edge in $BH_{n-1}^{j,i}$ has at least one endpoint incident with two i -dimensional faulty edges, we call it a *non- r -edge*. A *pivot vertex* is a vertex incident with exactly one non-faulty edge in $BH_{n-1}^{j,i}$.

In what follows, we shall present some basic properties of balanced hypercube, which will be used later.

Lemma 1 [18]. BH_n is bipartite.

Lemma 2 [25]. BH_n is vertex-transitive and edge-transitive.

Theorem 3 [12]. Let F be a set of edges with $|F| \leq 2n - 3$ and let $\{s_1, s_2\}$ and $\{t_1, t_2\}$ be two sets of vertices in different partite sets of BH_n for $n \geq 2$. Then $BH_n - F$ contains vertex-disjoint s_1, t_1 -path and s_2, t_2 -path that cover all vertices of it.

Theorem 4 [26]. BH_n is $2n - 2$ edge fault-tolerant Hamiltonian laceable for $n \geq 2$.

Theorem 5 [9]. Let F be an arbitrary edge set with $|F| \leq 4n - 5$ and $\delta(BH_n - F) \geq 2$. Then each edge in $BH_n - F$ lies on a fault-free Hamiltonian cycle for all $n \geq 2$.

Theorem 6 [14]. BH_n is hyper-Hamiltonian laceable for $n \geq 1$.

3. Main Results

Lemma 7. Let F be an edge subset of BH_3 with $|F| \leq 8$. $BH_3 - F$ is Hamiltonian if there exists no f_4 -cycles in $BH_3 - F$ and $\delta(BH_3 - F) \geq 2$.

Proof. Clearly, we need only to consider that $|F| = 8$. Since $\sum_{i=0}^2 |F_i| = |F| = 8$, there is an integer $i \in \{0, 1, 2\}$ such that $|F_i| \geq 3$. Without loss of generality, we assume $i = 2$, that is $|F_2| \geq 3$. In the following, we decompose BH_3 into four subcubes BH_2^j by deleting D_2 , where $j = 0, 1, 2, 3$. $|F^j| \leq \sum_{j=0}^3 |F^j| = |F| - |F_2| \leq 5$. We consider the following three cases in terms of the minimum degree of $BH_2^j - F^j$ and the cardinality of F^j , where $j = 0, 1, 2, 3$.

Case 1. $\delta(BH_2^j - F^j) \geq 2$ for all $j \in \{0, 1, 2, 3\}$.

Subcase 1.1. $|F^0| \leq 3$.

Since $\sum_{j=0}^3 |F^j| = |F| - |F_2| \leq 5$ and $|F^0| \leq 3$, $|F^j| \leq 2$ for $j = 1, 2, 3$. Obviously, there exists an r -edge $e_0 = a^0b^0$ in BH_2^0 . Let a^0b^1, a^3b^0 be the 2-dimensional nonfaulty edges. By Theorem 5, there is a fault-free Hamiltonian cycle C_0 in $BH_2^0 - F^0$ that contains e_0 . Let $H_0 = C_0 - e_0$ and let $a^1b^2, a^2b^3 \in D_2 - F_2$, by Theorem 4, there is a Hamiltonian path H_j in $BH_2^j - F^j$ that joins a^j and b^j for $j = 1, 2, 3$. Thus, $\langle a^0, b^1, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H_0, a^0 \rangle$ (see Fig. 4) is a fault-free Hamiltonian cycle in $BH_3 - F$.

Subcase 1.2. $|F^0| = 4$.

Clearly, $3 \leq |F_2| \leq 4$, and $|F^j| \leq 1$ for each $j = 1, 2, 3$. Since $|F_2| \leq 4$, there is an edge $f_0 = a^0b^0 \in F^0$ adjacent to at least one 2-dimensional nonfaulty edge, say a^0b^1 . By Theorem 5, there is a Hamiltonian cycle C_0 in $BH_2^0 - F^0 + f_0$ that contains f_0 . Let $H_0 = C_0 - f_0$.

If f_0 is an r -edge, assume that $a^3b^0 \in D_2 - F_2$. Let $a^1b^2, a^2b^3 \in D_2 - F_2$, by Theorem 4, there is a Hamiltonian path H_j in $BH_2^j - F^j$ that joins a^j and b^j for $j = 1, 2, 3$. Thus, $\langle a^0, b^1, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H_0, a^0 \rangle$ is a fault-free Hamiltonian cycle in $BH_3 - F$.




Fig. 4. Illustration for Subcase 1.1.

If f_0 is a non- r -edge, assume that b^0 is incident with two 2-dimensional faulty edges. Let $c^0 b^0 \notin E(C_0)$ with $c^0 \in V(C_0)$ and let d^0 and e^0 be the neighbors of c^0 satisfying $c^0 d^0, c^0 e^0 \in E(C_0)$. Let H_{00} be the path from a^0 to e^0 on C_0 and let H_{01} be the path from d^0 to b^0 on C_0 , then we denote C_0 by $\langle a^0, H_{00}, e^0, c^0, d^0, H_{01}, b^0, a^0 \rangle$. Since b^0 is incident with two 2-dimensional faulty edges and $|F_2| \leq 4$, d^0 or e^0 is incident with at least one 2-dimensional nonfaulty edge.

(1) If d^0 is incident with a 2-dimensional nonfaulty edge $a^3 d^0$, let $a^1 b^2, a^2 b^3 \in D_2 - F_2$. Since $|F^j| \leq 1$, by Theorem 4, there is a Hamiltonian path H_j in $BH_2^j - F^j$ that join a^j and b^j for $j = 1, 2, 3$. Then $\langle a^0, b^1, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, d^0, H_{01}, b^0, c^0, e^0, H_{00}, a^0 \rangle$ (see Fig. 5) is a fault-free Hamiltonian cycle in $BH_3 - F$.

(2) If d^0 is incident with two 2-dimensional faulty edges, then there exists an edge $c^3 e^0 \in D_2 - F_2$. We may assume that all edges in F^0 are non- r -edges, otherwise, we can replace f_0 by an r -edge in F^0 . So all edges in F^0 are incident with d^0 or b^0 . Since there exists no f_4 -cycles in $BH_3 - F$, the length of the cycle $C_{00} = \langle b^0, c^0, d^0, H_{01}, b^0 \rangle$ is at least 6. Hence, there exists an r -edge $g^0 f^0$ on C_{00} . Let H_{010} be the path from d^0 to f^0 on C_{00} and let H_{011} be the path from g^0 to b^0 on C_{00} . Then, we denote C_{00} by $\langle b^0, c^0, d^0, H_{010}, f^0, g^0, H_{011}, b^0 \rangle$. Let $g^0 d^1, a^3 f^0 \in D_2 - F_2$ satisfying $c^3 \neq a^3$ and $b^1 \neq d^1$ and let $a^j b^{j+1}, c^j d^{j+1} \in D_2 - F_2$ for $j = 1, 2$. By Theorem 3, there are two vertex-disjoint paths H_{j0} and H_{j1} in $BH_2^j - F^j$ such that H_{j0} joins a^j and b^j and H_{j1} joins c^j and d^j for $i = 1, 2, 3$. Then $\langle a^0, b^1, H_{10}, a^1, b^2, H_{20}, a^2, b^3, H_{30}, a^3, f^0, H_{010}, d^0, c^0, b^0, H_{011}, g^0, d^1, H_{11}, c^1, d^2, H_{21}, c^2, d^3, H_{31}, c^3, e^0, H_{00}, a^0 \rangle$ (see Fig. 6) is a fault-free Hamiltonian cycle in $BH_3 - F$.

Subcase 1.3. $|F^0| = 5$.

Then $|F^1| = |F^2| = |F^3| = 0$ and $|F_2| = 3$. Since $|F_2| = 3$ and $\delta(BH_2^0 - F^0) \geq 2$, there exist two r -edges $f_1 = a^0 b^0$ and $f_2 = c^0 d^0$ in F^0 . By Theorem 5, there is a Hamiltonian cycle C_0 in $BH_2^0 - F^0 + f_1 + f_2$ that contains f_1 . If $f_2 \notin E(C_0)$, the




Fig. 5. Illustration for Subcase 1.2.




Fig. 6. Illustration for Subcase 1.2.

situation is similar to the case that f_0 is r -edge in Subcase 1.2. So, we assume that $f_2 \in E(C_0)$.

Subcase 1.3.1. f_1 is adjacent to f_2 on C_0 .

Without loss of generality, we assume that $a^0 = c^0$. Assume that $a^3b^0, c^3d^0, a^0b^1 \in D_2 - F_2$ (possibly $a^3 = c^3$). Let $a^0p^0, a^0q^0 \notin E(C_0)$ with $p^0, q^0 \in V(C_0)$ and let x^0, y^0 be the neighbors of p^0 satisfying $x^0p^0, y^0p^0 \in E(C_0)$ and s^0, t^0 be the neighbors of q^0 satisfying $s^0q^0, t^0q^0 \in E(C_0)$ (possibly $y^0 = s^0$). Since $|F_2| = 3$, s^0, t^0, x^0 or y^0 is incident with a 2-dimensional nonfaulty edge that is not incident with b^1 . We may assume that $t^0d^1 \in D_2 - F_2$ with $d^1 \neq b^1$. Let H_{00} be the path from b^0 to t^0 on C_0 and H_{01} be the path from q^0 to d^0 on C_0 . We denote C_0 by $\langle a^0, b^0, H_{00}, t^0, q^0, H_{01}, d^0, a^0 \rangle$. Let $a^jb^{j+1}, c^jd^{j+1} \in D_2 - F_2$ for $j = 1, 2$.

If $c^3 \neq a^3$, by Theorem 3, there are two vertex-disjoint paths H_{j0} and H_{j1} in $BH_2^j - F^j$ such that H_{j0} joins a^j and b^j and H_{j1} joins c^j and d^j for $i = 1, 2, 3$. Then $\langle a^0, b^1, H_{10}, a^1, b^2, H_{20}, a^2, b^3, H_{30}, a^3, b^0, H_{00}, t^0, d^1, H_{11}, c^1, d^2, H_{21}, c^2, d^3, H_{31}, c^3, d^0, H_{01}, q^0, a^0 \rangle$ (see Fig. 7) is a fault-free Hamiltonian cycle in $BH_3 - F$.

If $c^3 = a^3$, by Theorem 6, there is a Hamiltonian path H_3 in $BH_2^3 - \{a^3\}$ that joins d^3 and b^3 . And by Theorem 3, there are two vertex-disjoint paths H_{j0} and H_{j1} in $BH_2^j - F^j$ such that H_{j0} joins a^j and b^j and H_{j1} joins c^j and d^j for $i = 1, 2$. Then, $\langle a^0, b^1, H_{10}, a^1, b^2, H_{20}, a^2, b^3, H_3, d^3, c^2, H_{21}, d^2, c^1, H_{11}, d^1, t^0, H_{00}, b^0, a^3, d^0, H_{01}, q^0, a^0 \rangle$ (see Fig. 8) is a fault-free Hamiltonian cycle in $BH_3 - F$.

Subcase 1.3.2. f_1 is nonadjacent to f_2 on C_0 .

We denote C_0 by $\langle a^0, b^0, H_{00}, c^0, d^0, H_{01}, a^0 \rangle$. Since $|F_2| = 3$ and a^0b^0, c^0d^0 are r -edges, a^0, b^0, c^0 and d^0 are incident with at least five 2-dimensional nonfaulty edges in total, and each is incident with at least one 2-dimensional nonfaulty edge. Then there must exist $c^3d^0, a^3b^0, a^0b^1, c^0d^1 \in D_2 - F_2$ satisfying $c^3 \neq a^3$ or $b^1 \neq d^1$. Without loss of generality, we assume that $b^1 \neq d^1$. Let $a^jb^{j+1}, c^jd^{j+1} \in D_2 - F_2$ for




Fig. 7. Illustration for Subcase 1.3.1.




Fig. 8. Illustration for Subcase 1.3.1.




Fig. 9. Illustration for Subcase 1.3.2.




Fig. 10. Illustration for Subcase 1.3.2.

$j = 1, 2$.

If $c^3 \neq a^3$, by Theorem 3, there are two vertex-disjoint paths H_{j0} and H_{j1} in $BH_2^j - F^j$ such that H_{j0} joins a^j and b^j and H_{j1} joins c^j and d^j for $i = 1, 2, 3$. Then $\langle a^0, b^1, H_{10}, a^1, b^2, H_{20}, a^2, b^3, H_{30}, a^3, b^0, H_{00}, c^0, d^1, H_{11}, c^1, d^2, H_{21}, c^2, d^3, H_{31}, c^3, d^0, H_{01}, a^0 \rangle$ (see Fig. 9) is a fault-free Hamiltonian cycle in $BH_3 - F$.

If $c^3 = a^3$, by Theorem 6, there is a Hamiltonian path H_3 in $BH_2^3 - \{a^3\}$ that joins d^3 and b^3 . And by Theorem 3, there are two vertex-disjoint paths H_{j0} and H_{j1} in $BH_2^j - F^j$ such that H_{j0} joins a^j and b^j and H_{j1} joins c^j and d^j for $i = 1, 2$. Then, $\langle a^0, b^1, H_{10}, a^1, b^2, H_{20}, a^2, b^3, H_3, d^3, c^2, H_{21}, d^2, c^1, H_{11}, d^1, c^0, H_{00}, b^0, a^3, d^0, H_{01}, a^0 \rangle$ (see Fig. 10) is a fault-free Hamiltonian cycle in $BH_3 - F$.

Case 2. $\delta(BH_2^j - F^j) = 1$ for some $j \in \{0, 1, 2, 3\}$.




Fig. 11. Illustration for Subcase 2.1.1.




Fig. 12. Illustration for Subcase 2.1.2.

Subcase 2.1. $|F^0| = 5$.

Then $|F^1| = |F^2| = |F^3| = 0$ and $|F_2| = 3$. Since $|F^0| = 5$, there are at most two pivot vertices in BH_2^0 .

Subcase 2.1.1. There is exactly one pivot vertex a^0 in BH_2^0 .

Let f_0 and f_1 be two r -edges in $e(a^0) \cap F^0$ and let $e_0 = a^0 b^0$ be the nonfaulty edge in $e(a^0) \cap E(BH_2^0)$. By Theorem 5, there is a Hamiltonian cycle C_0 in $BH_2^0 - F^0 + f_0 + f_1$ that contains e_0 . Notice also that one of f_0 and f_1 is on C_0 , say $f_0 = a^0 b^0 \in E(C_0)$. Let $H_0 = C_0 - f_0$. Then we can choose $b^0 a^3 \in D_2 - F_2$ and $a^j b^{j+1} \in D_2 - F_2$ for $j = 0, 1, 2$. By Theorem 4, there is a Hamiltonian path H_j in $BH_2^j - F^j$ that join a^j and b^j for $j = 1, 2, 3$. Then $\langle a^0, b^1, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H_0, a^0 \rangle$ (see Fig. 11) is a fault-free Hamiltonian cycle in $BH_3 - F$.

Subcase 2.1.2. There are exactly two pivot vertices a^0 and b^0 in BH_2^0 .

Then a^0 and b^0 are endpoints of a faulty edge $f_0 = a^0 b^0 \in F^0$. Let $f_1 \in e(a^0) \cap F^0 - f_0$ and $e_0 = a^0 d^0 \in E(BH_2^0) - F^0$. By Theorem 5, there exists a Hamiltonian cycle C_0 in $BH_2^0 - F^0 + f_0 + f_1$ that contains e_0 . Clearly, C_0 must contain f_0 , let $H_0 = C_0 - f_0$, so we denote C_0 by $\langle a^0, H_0, b^0, a^0 \rangle$. Let $b^0 a^3 \in D_2 - F_2$ and $a^j b^{j+1} \in D_2 - F_2$ for $j = 0, 1, 2$. Since $|F^j| = 0$, by Theorem 4, there is a Hamiltonian path H_j in $BH_2^j - F^j$ that join a^j and b^j for $j = 1, 2, 3$. Then $\langle a^0, b^1, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H_0, a^0 \rangle$ (see Fig. 12) is a fault-free Hamiltonian cycle in $BH_3 - F$.

Subcase 2.2. $|F^0| \leq 4$.

Since $\sum_{j=0}^3 |F^j| = |F| - |F_2| \leq 5$ and $|F^0| \leq 4$, there exists at most one pivot vertex a^0 in BH_2^0 . Then $3 \leq |F^0| \leq 4$ and $|F_2| \leq 5$. Assume that $f_0 = a^0 b^0$ is an r -edge in $e(a^0) \cap F^0$. By Theorem 5, there is a Hamiltonian cycle C_0 in $BH_2^0 - F^0 + f_0$. And C_0 must contain f_0 . Let $H_0 = C_0 - f_0$ and let $b^0 a^3 \in D_2 - F_2$, $a^j b^{j+1} \in D_2 - F_2$ for $j = 0, 1, 2$. Since $|F^j| \leq 8 - 3 - 3 = 2$, by Theorem 4,

there is a Hamiltonian path H_j in $BH_2^j - F^j$ that join a^j and b^j for $j = 1, 2, 3$. Then $\langle a^0, b^1, H_{10}, a^1, b^2, H_{20}, a^2, b^3, H_{30}, a^3, b^0, H_0, a^0 \rangle$ is a fault-free Hamiltonian cycle in $BH_3 - F$.




Fig. 13. Illustration for Case 3.

Case 3. $\delta(BH_2^j - F^j) = 0$ for some $j \in \{0, 1, 2, 3\}$.

Since $\sum_{j=0}^3 |F^j| \leq 5$, there is at most one isolated vertex in BH_2^j for $j = 0, 1, 2, 3$. By assumption the isolated vertex $a^0 \in V(BH_2^0)$. Let $a^0 b^1, a^0 d^1 \in D_2 - F_2$. Since $|F_2| \leq 4$, there exist two r -edges $f_1 = a^0 b^0$ and $f_2 = a^0 d^0$ in $e(a^0) \cap F^0$ such that $c^3 b^0, d^0 a^3 \in D_2 - F_2$ satisfying $a^3 \neq c^3$. Since $\delta(BH_2^0 - F^0 + f_1 + f_2) = 2$ and $|F^0 - f_1 - f_2| \leq 3$, by Theorem 5, there is a Hamiltonian cycle C_0 of $BH_2^0 - F^0 + f_1 + f_2$ containing f_1 and f_2 . Let $H_0 = C_0 - f_1 - f_2$. Then we can choose $a^j b^{j+1}, c^j d^{j+1} \in D_2 - F_2$ for $j = 1, 2$. Since $|F^j| \leq 8 - 3 - 4 = 1$, by Theorem 3, there are two vertex-disjoint paths H_{j0} and H_{j1} in $BH_2^j - F^j$ such that H_{j0} joins a^j and b^j and H_{j1} joins c^j and d^j for $i = 1, 2, 3$. Hence, $\langle a^0, b^1, H_{10}, a^1, b^2, H_{20}, a^2, b^3, H_{30}, a^3, d^0, H_0, b^0, c^3, H_{31}, d^3, c^2, H_{21}, d^2, c^1, H_{11}, d^1, a^0 \rangle$ (see Fig. 13) is a fault-free Hamiltonian cycle in $BH_3 - F$. $\square$

Theorem 8. Let F be an edge subset of BH_n with $|F| \leq 5n - 7$ for $n \geq 2$. $BH_n - F$ is Hamiltonian if there exists no f_4 -cycles in $BH_n - F$ and $\delta(BH_n - F) \geq 2$.

Proof. Clearly, it suffices to consider the condition that $|F| = 5n - 7$. We prove the theorem by induction on n . By Theorem 5, the assertion holds when $n = 2$. And it follows from Lemma 7 that the theorem holds when $n = 3$. Thus, we assume that the statement holds for $n - 1$ wherever $n \geq 4$. Next we consider BH_n . Since $\sum_{i=0}^{n-1} |F_i| = |F| = 5n - 7$, there is an integer $i \in \{0, 1, 2, \dots, n-1\}$ such that $|F_i| \geq 4$. Without loss of generality, we assume that $|F_{n-1}| \geq 4$. In the following, we decompose BH_n into four subcubes BH_{n-1}^j by deleting D_{n-1} , where $j = 0, 1, 2, 3$. We distinguish the following three cases.

Case 1. $\delta(BH_{n-1}^j - F^j) \geq 2$ for all $j \in \{0, 1, 2, 3\}$.

Subcase 1.1. There exists no f_4 -cycles in $BH_{n-1}^j - F^j$ for all $j \in \{0, 1, 2, 3\}$.

Subcase 1.1.1. $|F^0| = 5n - 11$.

Clearly, $|F_{n-1}| = 4$ and $|F^1| = |F^2| = |F^3| = 0$. Since $|F_{n-1}| = 4$ and $\delta(BH_{n-1}^0 - F^0) \geq 2$, there exist at least $5n - 11 - 2(2n - 4) = n - 3$ r -edges in F^0 for $n \geq 4$. Let $f_0 = a^0b^0$ be an r -edge in F^0 . By induction hypothesis, there is a Hamiltonian cycle C_0 in $BH_{n-1}^0 - F^0 + f_0$. If $f_0 \notin E(C_0)$, then there must exist an r -edge f_1 on C_0 . We may assume that $f_0 \in E(C_0)$. Let $H_0 = C_0 - f_0$. Then let $a^0b^1, a^3b^0, a^1b^2, a^2b^3$ be $(n - 1)$ -dimensional nonfaulty edges. By Theorem 4, there is a Hamiltonian path H_j in $BH_{n-1}^j - F^j$ that join a^j and b^j for $j = 1, 2, 3$. Then $\langle a^0, b^1, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H_0, a^0 \rangle$ is a fault-free Hamiltonian cycle in $BH_n - F$.

Subcase 1.1.2. $|F^0| \leq 5n - 12$.

By induction hypothesis, there is a Hamiltonian cycle C_0 in $BH_{n-1}^0 - F^0$. Since $|F_{n-1}| \leq 5n - 7$, there exist at most $\lfloor \frac{5n-7}{2} \rfloor$ vertices in $V(C_0)$ incident with two $(n - 1)$ -dimensional faulty edges. So there exists an r -edge $e_0 = a^0b^0$ in $E(C_0)$. Let $a^0b^1, a^3b^0 \in D_{n-1} - F_{n-1}$. Since $\sum_{j=0}^3 |F^j| = |F| - |F_{n-1}| \leq 5n - 11$, $|F^j| \leq 4n - 9$ for all $j = 1, 2, 3$ and there exists an integer $j \in \{1, 2, 3\}$ such that $|F^j| \leq 2n - 4$.

(1) $|F^2| \leq 2n - 4$. If there exists an r -edge $e_1 = a^1b^1 \in e(b^1) \cap E(BH_{n-1}^1)$, let $a^1b^2 \in D_{n-1} - F_{n-1}$. And if there exists an r -edge $e_3 = a^3b^3 \in e(a^3) \cap E(BH_{n-1}^3)$, let $a^2b^3 \in D_{n-1} - F_{n-1}$. Since $|F^j| \leq 4n - 9$ and $\delta(BH_{n-1}^j - F^j) \geq 2$, by Theorem 5, there is a fault-free Hamiltonian cycle C_j in $BH_{n-1}^j - F^j$ that contains e_j for $j = 1, 3$. Let $H_j = C_j - e_j$ for $j = 1, 3$. Since $|F^2| \leq 2n - 4$, by Theorem 4, there is a fault-free Hamiltonian path H_2 in $BH_{n-1}^2 - F^2$ that joins a^2 and b^2 . Thus, $\langle a^0, b^1, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H_0, a^0 \rangle$ is a fault-free Hamiltonian cycle in $BH_n - F$.

(2) $|F^3| \leq 2n - 4$ (the case $|F^1| \leq 2n - 4$ is similar). If there exists an r -edge $e_1 = a^1b^1 \in e(b^1) \cap E(BH_{n-1}^1)$, let $a^1b^2 \in D_{n-1} - F_{n-1}$. And if there exists an r -edge $e_2 = a^2b^2 \in e(b^2) \cap E(BH_{n-1}^2)$, let $a^2b^3 \in D_{n-1} - F_{n-1}$. By Theorem 5, there is a fault-free Hamiltonian cycle C_j in $BH_{n-1}^j - F^j$ that contains e_j for $j = 1, 2$. Let $H_j = C_j - e_j$ for $j = 1, 2$. Since $|F^3| \leq 2n - 4$, by Theorem 4, there is a fault-free Hamiltonian path H_3 in $BH_{n-1}^3 - F^3$ that joins a^3 and b^3 . Thus, $\langle a^0, b^1, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H_0, a^0 \rangle$ is a fault-free Hamiltonian cycle in $BH_n - F$.

If one of the aforementioned edges e_1, e_2 and e_3 does not exist, say e_1 , it means that all edges in $e(b^1) \cap E(BH_{n-1}^1)$ are non- r -edges, then there exist $2(2n - 2) = 4n - 4$ $(n - 1)$ -dimensional faulty edges adjacent to the edges in $e(b^1) \cap E(BH_{n-1}^1)$. It is easy to choose four edges $a^0b^1, a^1b^2, a^2b^3, a^3b^0 \in D_{n-1} - F_{n-1}$. Since $|F^j| \leq 5n - 7 - (4n - 4) = n - 3 \leq 2n - 4$, by Theorem 4, there is a fault-free Hamiltonian path H_j in $BH_{n-1}^j - F^j$ that joins a^j and b^j for $j = 0, 1, 2, 3$. Hence, $\langle a^0, b^1, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H_0, a^0 \rangle$ is a fault-free Hamiltonian cycle




Fig. 14. Illustration for Subcase 1.2.




Fig. 15. Illustration for Subcase 2.1.

in $BH_n - F$.

Subcase 1.2. There exists an f_4 -cycle in $BH_{n-1}^j - F^j$ for some $j \in \{0, 1, 2, 3\}$.

An f_4 -cycle is incident with $2(2n - 4) = 4n - 8$ faulty edges, so there exists at most one f_4 -cycle in $BH_{n-1}^j - F^j$. By our assumption, it follows that the f_4 -cycle lies in $BH_{n-1}^0 - F^0$. Let a^0, c^0 be the nonadjacent vertices whose degrees are both two in the f_4 -cycle. Since there exists no f_4 -cycle in $BH_n - F$, a^0 or c^0 , say a^0 , is incident with at least one $(n - 1)$ -dimensional nonfaulty edge $a^0b^1 \in D_{n-1} - F_{n-1}$. Since $|F_{n-1}| \leq 5n - 7 - (4n - 8) = n + 1 < 2(2n - 4)$ whenever $n \geq 4$, there exists a faulty r -edge $f_0 = a^0b^0$ in BH_{n-1}^0 , then let $a^3b^0 \in D_{n-1} - F_{n-1}$. Clearly, there exists no f_4 -cycles in $BH_{n-1}^0 - F^0 + f_0$, and $|F^0 - f_0| \leq 5n - 12$, by induction hypothesis, there is a Hamiltonian cycle C_0 in $BH_{n-1}^0 - F^0 + f_0$. Notice that the cycle C_0 must contain f_0 . Let $H_0 = C_0 - f_0$ and let $a^1b^2, a^2b^3 \in D_{n-1} - F_{n-1}$. Since $|F^j| \leq 5n - 7 - 4 - (4n - 8) = n - 3 \leq 2n - 4$, by Theorem 4, there is a fault-free Hamiltonian path H_j in $BH_{n-1}^j - F^j$ that joins a^j and b^j for $j = 1, 2, 3$. Thus, $\langle a^0, b^1, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H_0, a^0 \rangle$ (see Fig. 14) is a fault-free Hamiltonian cycle in $BH_n - F$.

Case 2. $\delta(BH_{n-1}^j - F^j) = 1$ for some $j \in \{0, 1, 2, 3\}$.

Since $|F| - |F_{n-1}| \leq 5n - 7 - 4 = 5n - 11$ and a pivot vertex in $BH_n - D_{n-1}$ is incident with $2n - 3$ faulty edges in $F - F_{n-1}$, then there exist at most two pivot vertices in $BH_n - D_{n-1}$.

Subcase 2.1. There is exactly one pivot vertex in $BH_{n-1}^j - F^j$ for some $j \in \{0, 1, 2, 3\}$.

Clearly, there exists no f_4 -cycle in each $BH_{n-1}^j - F^j$ since $2n - 3 + 4 + 2(2n - 4) = 6n - 7 \geq 5n - 7$. We may assume that the pivot vertex $u \in V(BH_{n-1}^0)$. Since $\delta(BH_n - F) \geq 2$, u is incident with at least one $(n - 1)$ -dimensional nonfaulty edge, say $ub^1 \in D_{n-1} - F_{n-1}$. Since $2n - 3 + 2(2n - 3) = 6n - 9 \geq 5n - 7$ whenever

$n \geq 4$, we can choose an r -edge $f_0 = ub^0 \in e(u) \cap F^0$. If there exists an f_4 -cycle in $BH_{n-1}^0 - F^0 + f_0$, then u must be a vertex in the f_4 -cycle. Since $|F^0| \leq 5n - 11$ and u is a pivot vertex, there exists at most one vertex with degree 2 in $BH_{n-1}^0 - F^0$, which is the vertex on the f_4 -cycle. Since $|F_{n-1}| \leq 5n - 7 - (2n - 3) - (2n - 4) = n < 2(2n - 3 - 1)$, there exists an r -edge $f_1 \in e(u) \cap F^0 - f_0$ and there exists no f_4 -cycles in $BH_{n-1}^0 - F^0 + f_1$. So, we may assume that there exists no f_4 -cycles in $BH_{n-1}^0 - F^0 + f_0$. Let $a^3b^0 \in D_{n-1} - F_{n-1}$. By induction hypothesis, there is a Hamiltonian cycle C_0 in $BH_{n-1}^0 - F^0 + f_0$. Clearly, the cycle C_0 must contain f_0 , let $H_0 = C_0 - f_0$. Notice that $|F^j| \leq \sum_{j=1}^3 |F^j| \leq 5n - 7 - (2n - 3) - 4 = 3n - 8 \leq 4n - 9$ for $j = 1, 2, 3$, and there exists an integer $j \in \{1, 2, 3\}$ such that $|F^j| \leq 2n - 4$.

(1) $|F^2| \leq 2n - 4$. Since $|F_{n-1}| \leq 5n - 7 - (2n - 3) = 3n - 4 < 2(2n - 2)$, there exist an r -edge $e_1 = a^1b^1 \in e(b^1) \cap E(BH_{n-1}^1)$ and an r -edge $e_3 = a^3b^3 \in e(a^3) \cap E(BH_{n-1}^3)$. Assume that $a^1b^2, a^2b^3 \in D_{n-1} - F_{n-1}$. By Theorem 5, there is a fault-free Hamiltonian cycle C_j in $BH_{n-1}^j - F^j$ that contains e_j for $j = 1, 3$. Let $H_j = C_j - e_j$ for $j = 1, 3$. By theorem 4, there is a fault-free Hamiltonian path H_2 in $BH_{n-1}^2 - F^2$ that joins a^2 and b^2 . Thus, $\langle u, b^1, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H_0, u \rangle$ (see Fig. 15) is a fault-free Hamiltonian cycle in $BH_n - F$.

(2) $|F^3| \leq 2n - 4$ (the case $|F^1| \leq 2n - 4$ is similar). Since $|F_{n-1}| \leq 5n - 7 - (2n - 3) = 3n - 4 < 2(2n - 2)$, there exists an r -edge $e_1 = a^1b^1 \in e(b^1) \cap E(BH_{n-1}^1)$. Assume that $a^1b^2 \in D_{n-1} - F_{n-1}$. And there exists an r -edge $e_2 = a^2b^2 \in e(b^2) \cap E(BH_{n-1}^2)$. In addition, assume that $a^2b^3 \in D_{n-1} - F_{n-1}$. By Theorem 5, there is a fault-free Hamiltonian cycle C_j in $BH_{n-1}^j - F^j$ that contains e_j for $j = 1, 2$. Let $H_j = C_j - e_j$ for $j = 1, 2$. Since $|F^3| \leq 2n - 4$, by Theorem 4, there is a fault-free Hamiltonian path H_3 in $BH_{n-1}^3 - F^3$ that joins a^3 and b^3 . Thus, $\langle u, b^1, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, b^0, H_0, u \rangle$ is a fault-free Hamiltonian cycle in $BH_n - F$.

Subcase 2.2. There are two pivot vertices u and v in $BH_{n-1}^j - F^j$ for some $j \in \{0, 1, 2, 3\}$.

If one of the following three conditions (1), (2) and (3) holds, then there exists an integer $i \in \{0, 1, \dots, n - 2\}$ such that u and v are both incident with two i -dimensional faulty edges and $|F_i| \geq 4$.

- (1) u and v are not adjacent.
- (2) uv is a nonfaulty edge.
- (3) uv is an $(n - 1)$ -dimensional faulty edge.

We may assume that $i = n - 2$, then we can decompose BH_n into four subcubes $BH_{n-1}^{j, n-2}$ by deleting D_{n-2} , where $j = 0, 1, 2, 3$. In $BH_{n-1}^{j, n-2}$ for $j = 0, 1, 2, 3$, u and v are two vertices whose degrees are 2 or 3, notice that there exists no isolated vertex and at most one pivot vertex in $BH_n - D_{n-2}$. Thus, the case is degenerated to the Case 1 or Subcase 2.1.

If uv is a faulty edge and uv is not an $(n - 1)$ -dimensional edge, we may assume that $u, v \in V(BH_{n-1}^0)$. Notice that uv is an r -edge, assume that ub^1, a^3v are $(n - 1)$ -dimensional nonfaulty edges. By induction hypothesis, there is a Hamiltonian cycle




Fig. 16. Illustration for Subcase 2.2.




Fig. 17. Illustration for Subcase 3.1.

C_0 in $BH_{n-1}^0 - F^0 + f_0$. Clearly, the cycle C_0 must contain f_0 , let $H_0 = C_0 - f_0$. Let $a^1b^2, a^2b^3 \in D_{n-1} - F_{n-1}$. Since $|F^j| \leq 5n - 7 - 4 - 2(2n - 3) + 1 = n - 4 < 2n - 4$, by Theorem 4, there is a fault-free Hamiltonian path H_j in $BH_{n-1}^j - F^j$ that joins a^j and b^j for $j = 1, 2, 3$. Therefore, $\langle u, b^1, H_1, a^1, b^2, H_2, a^2, b^3, H_3, a^3, v, H_0, u \rangle$ (see Fig. 16) is a fault-free Hamiltonian cycle in $BH_n - F$.

Case 3. $\delta(BH_{n-1}^j - F^j) = 0$ for some $j \in \{0, 1, 2, 3\}$.

Since $|F| - |F_{n-1}| \leq 5n - 7 - 4 = 5n - 11$, there exist at most two isolated vertices or at most one pivot vertex and one isolated vertex in $BH_{n-1}^j - F^j$ for some $j \in \{0, 1, 2, 3\}$.

Subcase 3.1. There is exactly one isolated vertex in $BH_{n-1}^j - F^j$ for some $j \in \{0, 1, 2, 3\}$.

We may assume that the isolated vertex $u \in BH_{n-1}^0$, then u is incident with $2n - 2$ edges of F^0 and two $(n - 1)$ -dimensional nonfaulty edges ub^1 and ud^1 .

If $|F_i| \leq 3$ for each $i \in \{0, 1, \dots, n - 2\}$, then $|F^j| \leq 3(n - 1) - (2n - 2) = n - 1 \leq 2n - 5$ for $j = 1, 2, 3$. Since $|F_{n-1}| \leq 5n - 7 - (2n - 2) = 3n - 5 < 2(2n - 2) - 3 = 4n - 7$, there exist two r -edges $f_1 = ub^0$ and $f_2 = ud^0$ in $e(u) \cap F^0$ satisfying $a^3b^0, c^3d^0 \in D_{n-1} - F_{n-1}$ and $a^3 \neq c^3$. Since $|F^0| \leq 3(n - 1)$ and u is an isolated vertex, there exists no vertices with degree at most two in $BH_{n-1}^0 - F^0$. So, there exists no f_4 -cycle in $BH_{n-1}^0 - F^0 + f_1 + f_2$. Since $|F^0 - f_1 - f_2| \leq 3(n - 1) - 2 = 3n - 5 \leq 5n - 12$, by induction hypothesis, there is a Hamiltonian cycle C_0 in $BH_{n-1}^0 - F^0 + f_1 + f_2$. Clearly, the cycle C_0 must contain f_1 and f_2 , let $H_0 = C_0 - f_1 - f_2$. And let $a^jb^{j+1}, c^jd^{j+1} \in D_{n-1} - F_{n-1}$ for $j = 1, 2$. By Theorem 3, there are two vertex-disjoint paths H_{j0} and H_{j1} in $BH_{n-1}^j - F^j$ such that H_{j0} joins a^j and b^j and H_{j1} joins c^j and d^j for $j = 1, 2, 3$. Then $\langle u, b^1, H_{10}, a^1, b^2, H_{20}, a^2, b^3, H_{30}, a^3, d^0, H_0, b^0, c^3, H_{31}, d^3, c^2, H_{21}, d^2, c^1, H_{11}, d^1, u \rangle$ (see Fig. 17) is a fault-free Hamiltonian cycle in $BH_n - F$.

If there exists an integer $i \in \{0, 1, \dots, n-2\}$ such that $|F_i| \geq 4$, say $|F_{n-2}| \geq 4$, then we decompose BH_n into four subcubes $BH_{n-1}^{j, n-2}$ by deleting D_{n-2} , where $j = 0, 1, 2, 3$. Notice that u is a vertex whose degree is 2 in $BH_{n-1}^{j, n-2}$ for some $j \in \{0, 1, 2, 3\}$. If there exists no isolated vertex in $BH_{n-1}^{j, n-2}$ for each $j \in \{0, 1, 2, 3\}$, then this case is degenerated to Case 1 or Case 2. Otherwise, if there still exists an isolated vertex v in $BH_{n-1}^{j, n-2}$ for some $j \in \{0, 1, 2, 3\}$, then there exists an integer $i \in \{0, 1, \dots, n-3\}$ such that u and v are both incident with two i -dimensional faulty edges and $|F_i| \geq 4$, we may assume that $i = n-3$. Then we decompose BH_n into four subcubes $BH_{n-1}^{j, n-3}$ by deleting D_{n-3} , where $j = 0, 1, 2, 3$. Notice that there exists no isolated vertex in $BH_{n-1}^{j, n-3}$ for $j \in \{0, 1, 2, 3\}$ as $5n-7-(2n-2)-(2n-2) = n-3$, the case is degenerated to Case 1 or Case 2.

Subcase 3.2. There are two isolated vertices u and v or a pivot vertex u and an isolated vertex v in $BH_{n-1}^j - F^j$ for some $j \in \{0, 1, 2, 3\}$.

It can be deduced that there exists an integer $i \in \{0, 1, \dots, n-2\}$ such that u and v are both incident with two i -dimensional faulty edges and $|F_i| \geq 4$, we may assume that $i = n-2$. Then we decompose BH_n into four subcubes $BH_{n-1}^{j, n-2}$ by deleting D_{n-2} , where $j = 0, 1, 2, 3$. Since $5n-7-(2n-2)-(2n-3) \geq n-2$, there exists no isolated vertex in $BH_{n-1}^{j, n-2}$ for $j \in \{0, 1, 2, 3\}$, the case is degenerated to Case 1 or Case 2. □

4. Conclusion

In this paper, we show that there exists a fault-free Hamiltonian cycle in $BH_n - F$ for $n \geq 2$ with $|F| \leq 5n-7$ if the degree of every vertex in $BH_n - F$ is at least two and there exists no f_4 -cycles in $BH_n - F$, which improves the upper bound of the number of faulty edges can be tolerated in BH_n . It is interesting to further consider Hamiltonicity of the faulty BH_n that contains no f_6 -cycle.

References

- [1] D. Cheng, R. Hao, Y. Feng, Two node-disjoint paths in balanced hypercubes, Appl. Math. Comput. 242 (2014) 127–142.
- [2] D. Cheng, R. Hao, Y. Feng, Vertex-fault-tolerant cycles embedding in balanced hypercubes, Inform. Sci. 288 (2014) 449–461.
- [3] J. Fan, X. Lin, X. Jia, Optimal path embeddings of paths with various lengths in twisted cubes, IEEE Trans. Parallel Distrib. Syst. 18(4) (2007) 511–521.
- [4] R. Hao, R. Zhang, Y. Feng, J. Zhou, Hamiltonian cycle embedding for fault tolerance in balanced hypercubes, Appl. Math. Comput. 244 (2014) 447–456.

- [5] S.-Y. Hsieh, G.H. Chen, C.W. Ho, Hamiltonian-laceability of star graphs, *Networks* 36 (2000) 225–232.
- [6] S.-Y. Hsieh, Y.-R. Cian, Conditional edge-fault Hamiltonicity of augment cubes, *Inform. Sci.* 180 (13) (2010) 2596–2617.
- [7] F.T. Leighton, *Introduction to Parallel Algorithms and Architectures*, Morgan Kaufmann Publishers, Calif, San Mateo, 1992.
- [8] P. Li, M. Xu, Edge-fault-tolerant edge-bipancyclicity of balanced hypercubes, *Appl. Math. Comput.* 307 (2017) 180–192.
- [9] P. Li, M. Xu, Fault-free Hamiltonian cycles in balanced hypercubes with conditional edge faults, *Int. J. Found. Comput. Sci.* 30(5) (2019) 693–717.
- [10] J. Liu, Y. Wang, Hamiltonian cycles in hypercubes with faulty edges, *Inform. Sci.* 256 (2014) 225–233.
- [11] H. Lü, On extra connectivity and extra edge-connectivity of balanced hypercubes, *Inter. J. Comput. Math.* 94 (2017) 813–820.
- [12] H. Lü, Paired many-to-many two-disjoint path cover of balanced hypercubes with faulty edges, *J. Supercomput.* 75 (2019) 400–424.
- [13] H. Lü, X. Li, H. Zhang, Matching preclusion for balanced hypercubes, *Theor. Comput. Sci.* 465 (2012) 10–20.
- [14] H. Lü, H. Zhang, Hyper-Hamiltonian laceability of balanced hypercubes, *J. Supercomput.* 68 (2014) 302–314.
- [15] J.-H. Park, Unpaired many-to-many disjoint path covers in restricted hypercube-like graphs, *Theor. Comput. Sci.* 617 (2016) 45–64.
- [16] C.H. Tsai, J. J. M. Tan, T. Liang, L. H. Hsu, Fault-tolerant Hamiltonian laceability of hypercubes, *Inform. Process. Lett.* 83 (2002) 301–306.
- [17] D.B. West, *Introduction to Graph Theory*, 2ed., Prentice Hall, 2001.
- [18] J. Wu, K. Huang, The balanced hypercube: a cube-based system for fault-tolerant applications, *IEEE Trans. Comput.* 46(4) (1997) 484–490.
- [19] J. Wu, K. Huang, Area efficient layout of balanced hypercubes, *Int. J. High Speed Electron. Syst.* 6(4) (1995) 631–646.
- [20] J.-M. Xu, *Topological structure and analysis of interconnection networks*, London, 2001.

- [21] M. Xu, H. Hu, J. Xu, Edge-pancyclicity and Hamiltonian laceability of the balanced hypercubes, *Appl. Math. Comput.* 189 (2007) 1393–1401.
- [22] D.-W. Yang, Y.-Q. Feng, J. Lee, J.-X. Zhou, On extra connectivity and extra edge-connectivity of balanced hypercubes, *Appl. Math. Comput.* 320 (2018) 464–473.
- [23] M. Yang, Bipanconnectivity of balanced hypercubes, *Comput. Math. Appl.* 60 (2010) 1859–1867.
- [24] M. Yang, Super connectivity of balanced hypercubes, *Appl. Math. Comput.* 219 (2012) 970–975.
- [25] J. Zhou, Z. Wu, S. Yang, K. Yuan, Symmetric property and reliability of balanced hypercube, *IEEE Trans. Comput.* 64(3) (2015) 876–881.
- [26] Q. Zhou, C. Chen, H. Lü, Fault-tolerant Hamiltonian laceability of balanced hypercubes, *Inform. Sci.* 300 (2015) 20–27.