

Efficient Concentration with Gaussian Approximation

Morgane Austern

Department of Statistics, Harvard University

MAUSTERN@FAS.HARVARD.EDU

Lester Mackey

Microsoft Research New England

LMACKEY@MICROSOFT.COM

Abstract

Concentration inequalities for the sample mean, like those due to Bernstein and Hoeffding, are valid for any sample size but overly conservative, yielding confidence intervals that are unnecessarily wide. The central limit theorem (CLT) provides asymptotic confidence intervals with optimal width, but these are invalid for all sample sizes. To resolve this tension, we develop new computable concentration inequalities with asymptotically optimal size, finite-sample validity, and sub-Gaussian decay. These bounds enable the construction of efficient confidence intervals with correct coverage for any sample size. We derive our inequalities by tightly bounding the Hellinger distance, Stein discrepancy, non-uniform Kolmogorov distance, and Wasserstein distance to a Gaussian, and, as a byproduct, we obtain the first explicit bounds for the Hellinger CLT.

Keywords: Efficient concentration inequality, Gaussian approximation, tail bound, quantile bound, confidence interval, Hellinger distance, Wasserstein distance, zero-bias coupling, Stein discrepancy, Stein kernel, central limit theorem

1. Introduction

Concentration inequalities for the sample mean are ubiquitous in probability theory, statistics, and machine learning. Given n independent and identically distributed (i.i.d.) observations W_1, \dots, W_n they allow us to give finite sample and high probability guarantees that the sample mean $\bar{W}_n \triangleq \frac{1}{n} \sum_{i=1}^n W_i$ is not too far away from the population mean $\mathbb{E}(W_1)$. Specifically, they provide upper bounds for the probability $P(\bar{W}_n - \mathbb{E}(W_1) \geq t/\sqrt{n})$ for each $t \geq 0$. Such inequalities lie at the heart of decision-making in reinforcement learning (Mnih et al., 2008; Audibert et al., 2009), generalization guarantees in high-dimensional statistics, machine learning, and deep learning (Wainwright, 2019; Bartlett and Mendelson, 2002; Zhou et al., 2018), and the design of efficient learning procedures (Maurer and Pontil, 2009).

However, standard concentration inequalities are overly conservative yielding confidence intervals that are unnecessarily wide and generalization guarantees that are weaker than needed. This is notably the case for the commonly used concentration inequalities of Hoeffding (1963) and Bernstein (1924). For random variables with bounded deviations $|W_1 - \mathbb{E}(W_1)| \leq R$ and variance $\sigma^2 \triangleq \text{Var}(W_1)$, the Hoeffding and Bernstein inequalities respectively state that

$$P(\bar{W}_n - \mathbb{E}(W_1) \geq t/\sqrt{n}) \leq e^{-\frac{t^2}{2R^2}} \quad \text{and} \quad (1)$$

$$P(\bar{W}_n - \mathbb{E}(W_1) \geq t/\sqrt{n}) \leq e^{-\frac{t^2}{2(\sigma^2 + Rt/3)}}. \quad (2)$$

Meanwhile, the central limit theorem (CLT) identifies the exact limit for these tail probabilities:

$$P(\bar{W}_n - \mathbb{E}(W_1) \geq t/\sqrt{n}) \xrightarrow{n \rightarrow \infty} \Phi^c\left(\frac{t}{\sigma}\right), \quad (3)$$

where $\Phi(\cdot)$ is the cumulative distribution function (CDF) of a standard normal distribution and $\Phi^c(\cdot) = 1 - \Phi(\cdot)$. As a result, standard confidence intervals based on the CLT are asymptotically exact and often much narrower than those obtained using concentration inequalities. However, these intervals are typically only asymptotically valid and provide incorrect coverage for every sample size n .

The choice between concentration inequalities that are finite-sample valid but loose and CLT-based approximations that are asymptotically tight but invalid is very unsatisfying. In this paper we derive new bounds that offer the best of both worlds: our new concentration inequalities are both finite-sample valid and *efficient*—that is, asymptotically of minimal width when scaled by \sqrt{n} . We achieve this by deriving explicit and practical bounds for the distance between the sample mean and a Gaussian with matching variance. In Secs. 3, 4, 5, and 6, we measure the distance in four different ways: via the Hellinger distance, the Stein discrepancy (as defined by [Ledoux et al. \(2015\)](#)), a non-uniform Kolmogorov distance, and the p -Wasserstein distance, respectively. In each case, our Gaussian approximation bounds yield efficient concentration inequalities of the form

$$\mathbb{P}(\bar{W}_n - \mathbb{E}(W_1) \geq t/\sqrt{n}) \leq \Phi^c(t/\sigma) + r_n(t/\sigma) \quad (4)$$

with explicit remainder terms $r_n(t/\sigma)$. For example, in the non-uniform Kolmogorov case, $r_n(t/\sigma)$ decreases at an $O(n^{-1/2})$ rate in n and at a sub-Gaussian $e^{-\Omega(t^2)}$ rate in t .

It is informative to compare these results with those obtained using classical CLT corrections. For random variables with bounded deviations, the Berry-Esseen bound ([Esseen, 1942](#)) guarantees that

$$|\mathbb{P}(\bar{W}_n - \mathbb{E}(W_1) \geq t/\sqrt{n}) - \Phi^c(t/\sigma)| \leq \frac{CR}{\sigma\sqrt{n}} \quad \text{for all } t \geq 0, \quad (5)$$

and some universal constant C . This yields an efficient concentration inequality, but the bound is overly conservative as the correction is independent of t . Non-uniform Berry-Esseen bounds ([Nagaev, 1965](#); [Bikyalis, 1966](#)) ameliorate this behavior by identifying a constant \tilde{C} satisfying

$$|\mathbb{P}(\bar{W}_n - \mathbb{E}(W_1) \geq t/\sqrt{n}) - \Phi^c(t/\sigma)| \leq \frac{\tilde{C}R}{\sigma\sqrt{n}(1+t/\sigma)^3} \quad \text{for all } t \geq 0.$$

Appealingly, this non-uniformity yields tighter bounds for larger t . However, the correction has only cubic, that is, $O(t^{-3})$, decay in t as the underlying argument only exploits the existence of a third moment of \bar{W}_n . Quantile coupling inequalities (see, e.g., [Mason and Zhou, 2012](#)) like the groundbreaking Komlós-Major-Tusnády approximations ([Komlós et al., 1975, 1976](#); [Bretagnolle and Massart, 1989](#), Thm. 1) and the strong embedding bounds of [Chatterjee \(2012\)](#); [Bhattacharjee and Goldstein \(2016\)](#) improve this t dependence for W_1 with finite exponential moments but provide at best exponential decay in t and $O(\log n/\sqrt{n})$ decay in n . By exploiting the sub-Gaussianity of W_1 , our theorems provide a correction term with *squared-exponential* $e^{-\Omega(t^2)}$ decay, leading to much sharper bounds for larger t . In addition, our non-uniform Kolmogorov bounds eliminate the extraneous $\log n$ factor from the decay rate.

As the Hellinger, Stein, non-uniform Kolmogorov, and Wasserstein distances each measure separate properties of the empirical average, the inequalities we obtain are complementary and hold under distinct conditions. For example, when controlling the Hellinger distance, the remainder term in Eq. (4) depends on the score function of the random variable W_1 . When controlling the Stein discrepancy, the remainder term instead depends on the moments of a quantity called the *Stein*

kernel, defined in Sec. 4. As the Stein kernel and the score function can be controlled in different settings those bounds complement each other. Finally, the bounds obtained using non-uniform Kolmogorov and p -Wasserstein distances apply to *all* bounded-deviation variables with no additional distributional assumptions. We note in general that when the random variables are continuous and the score function can be controlled (see Thm. 4 and Thm. 8) we can improve the tightness of our results by avoiding a lossy union bound. This is complemented by tighter efficient bounds for smoothed observations, which will hold under minimum conditions. We also propose new quantile bounds that are also valid for finite samples and efficient. Finally, with analogy to the empirical Bernstein bound (Mnih et al., 2008; Audibert et al., 2009; Maurer and Pontil, 2009), we develop new *empirical Berry-Esseen bounds* that are efficient and finite-sample valid even when the variance parameter σ^2 is unknown. We confirm numerically that these new bounds improve upon the most commonly used confidence intervals and the efficient $I_{n,3}$ confidence interval of Romano and Wolf (2000).

As a by-product of our inquiry we obtain the first computable bounds on the Hellinger distance to Gaussianity. Our bounds depend on the moments of the score function of W_1 . This is to contrast with the currently known rates of convergence of the entropic central limit theorems which are either not in closed form (Bobkov et al., 2013), require conditions on the spectral gap (Artstein et al., 2004), or depend on the Stein kernel (Ledoux et al., 2015). In addition, we propose a novel bound that, beyond empirical averages, upper-bounds the Hellinger distance between general random variables and a normal.

1.1. Overview

In Sec. 2 we define the notion of efficient concentration and quantiles, present some of the assumptions we will work with, and introduce notations. In Sec. 3 we propose efficient concentration inequalities using the Hellinger distance, under the condition that the random variables are continuous. Moreover, we propose the first-ever computable rate of convergence of the CLT in terms of the Hellinger distance. In Sec. 4 we propose three different efficient tail bounds under the condition that the random variables Y_1 admits a Stein kernel. Each one of those tail bounds will hold under different conditions and we contrast those to each other and previously known techniques. In Sec. 5 we use zero-bias couplings to bound non-uniform Kolmogorov distances and thereby obtain efficient tail bounds using only the bounded deviation assumption. In Sec. 6 we use the p -Wasserstein distances to obtain efficient tail bounds and quantile bounds under minimal conditions. In Sec. 7 we obtain even tighter efficient concentration inequalities for smoothed observations. Finally, in Sec. 8 we propose empirical efficient quantile bounds that can be computed without requiring any prior knowledge about the variance and present some numerical evaluations of these bounds. Throughout, we present sketches of the proofs of the derived results and defer full proofs to the appendices.

1.2. Notation

Throughout, we will let $\varphi(x) \triangleq \frac{1}{\sqrt{2\pi}}e^{-x^2/2}$ designate the probability density function (PDF) of a standard normal random variable and write $a_n \lesssim b_n$ to indicate that two sequences (a_n) and (b_n) satisfy $a_n \leq b_n[1 + o_n(1)]$. We define $a^+ = \max(a, 0)$ and make use of the following notation:

$$\begin{aligned} Y_i &\triangleq W_i - \mathbb{E}(W_1), & S_n &\triangleq \sqrt{n}(\bar{W}_n - \mathbb{E}(W_1)), & \tilde{S}_n &\triangleq \sigma^{-1}S_n, & \tilde{R} &\triangleq \sigma^{-1}R, \\ A_p &\triangleq \frac{\sqrt{e}\sqrt{p+2}(2e)^{1/p}}{\sqrt{2}}, & A_{n,p}^* &\triangleq \frac{(p+2)n^{1/p}}{2\sqrt{n}}, & \tilde{A}_{n,p} &\triangleq A_{n,p}^*\tilde{R}^{-2/p}. \end{aligned}$$

2. Defining Efficient Concentration

To match the setting of the classical Bernstein inequality Eq. (2), we will focus on random variables satisfying the following bounded deviations assumption.

Assumption (R, σ) *The scaled deviations $(S_n)_{n \geq 1}$ satisfy $S_n = \sqrt{n}(\bar{W}_n - \mathbb{E}(W_1))$ for a shared sequence of i.i.d. variables $(W_i)_{i \geq 1}$ with $\bar{W}_n \triangleq \frac{1}{n} \sum_{i=1}^n W_i$. We write the centered random variables $Y_i \triangleq W_i - \mathbb{E}(W_1)$, and we suppose that $\text{Var}(Y_1) = \sigma^2$ and $|Y_1| \leq R$ almost surely.*

Our first inferential goal is to tightly upper bound the tail probability $\mathbb{P}(S_n \geq t)$ for a given threshold $t \geq 0$. The CLT provides an asymptotic lower bound for this problem as $\mathbb{P}(S_n \geq t)$ is known to converge precisely to $\Phi^c(\frac{t}{\sigma})$ as n increases.

Proposition 1 (Asymptotic lower bound for valid tail bounds) *Fix any $t \geq 0$, any sequence of scaled deviations $(S_n)_{n \geq 1}$ satisfying Assump. (R, σ) , and any sequence of candidate tail bounds $(\delta_n(t))_{n \geq 1}$. If $\mathbb{P}(S_n \geq t) \leq \delta_n(t)$ for all n , then $\Phi^c(\frac{t}{\sigma}) \leq \liminf_{n \rightarrow \infty} \delta_n(t)$.*

Proof Suppose that a sequence $\delta_n(t)$ satisfies $\liminf_{n \rightarrow \infty} \delta_n(t) < \Phi^c(\frac{t}{\sigma})$. Then there exists an $\epsilon > 0$ such that $\delta_n(t) \leq \Phi^c(\frac{t}{\sigma}) - \epsilon$ for infinitely many n . However, by the CLT (see, e.g., [Durrett, 2019](#), Thm. 3.4.1), there exists an n_ϵ such that, for all $n > n_\epsilon$, $\Phi^c(\frac{t}{\sigma}) - \epsilon < \mathbb{P}(S_n \geq t)$. Therefore, $\delta_n(t) < \mathbb{P}(S_n \geq t)$ for infinitely many n , confirming the claim via its contrapositive. ■

Unfortunately, the CLT limit does not provide a suitable tail bound for any finite n , as $\mathbb{P}(S_n \geq t) > \Phi^c(\frac{t}{\sigma})$ for many S_n satisfying Assump. (R, σ) . However, by tightly bounding the distance between the distribution of S_n and the distribution of a Gaussian we can correct the asymptotic bound to obtain one that is both valid in finite samples and asymptotically exact. We will call such bounds *efficient concentration inequalities*.

Our second inferential goal is to tightly bound the quantiles of S_n . That is, given a tail probability $\delta \in (0, 1)$ we wish to find $q_n(R, \delta, \sigma)$ (a measurable function of $(W_i)_{i=1}^n, R, \sigma$, and δ) satisfying $\mathbb{P}(S_n \geq q_n(R, \delta, \sigma)) \leq \delta$. Such quantile bounds immediately deliver both one- and two-sided confidence intervals for the population mean $\mathbb{E}(W_1)$ as

$$\mathbb{P}(\bar{W}_n - \frac{q_n(R, \delta, \sigma)}{\sqrt{n}} \leq \mathbb{E}(W_1)) \wedge \mathbb{P}(|\mathbb{E}(W_1) - \bar{W}_n| \leq \frac{q_n(R, \delta/2, \sigma)}{\sqrt{n}}) \geq 1 - \delta.$$

The interval efficiency theory of [Romano and Wolf \(2000\)](#) implies that the CLT once again provides an asymptotic lower bound for any valid sequence of quantile bounds.

Proposition 2 (Asymptotic lower bound for valid quantile bounds) *Fix any $R > 0$, any $\delta \in (0, \frac{1}{2})$, and any nonnegative candidate quantile bounds $(q_n(R, \delta, \sigma))_{n \geq 1, \sigma > 0}$. Suppose that, for each $\sigma \in (0, R]$ and all $(S_n)_{n \geq 1}$ satisfying Assump. (R, σ) , $\mathbb{P}(S_n \geq q_n(R, \delta, \sigma)) \leq \delta$ for all n . Then $(q_n(R, \delta, \sigma))_{n \geq 1}$ is not asymptotically concentrated¹ on $[0, a]$ for any $a < -\sigma\Phi^{-1}(\delta)$.*

Proof The result follows by applying Thm. 2.1 of [Romano and Wolf \(2000\)](#) to the conservative confidence intervals $I_n(\delta, \sigma) = \bar{W}_n + \frac{1}{\sqrt{n}}[-q_n(R, \delta, \sigma), q_n(R, \delta, \sigma)]$ for the unknown mean $\mathbb{E}(W_1)$. ■

To construct *efficient quantile bounds*, we will once again tightly bound the distance between S_n and its Gaussian limit.

1. A sequence of nonnegative random variables $(X_n)_{n \geq 1}$ is asymptotically concentrated on $[0, a]$ if $(X_n - a)^+ \xrightarrow{P} 0$.

3. Efficient Concentration with Hellinger Approximation

For two continuous real-valued random variables X and Y be with respective probability density functions (PDFs) f_X and f_Y we define the Hellinger distance as

$$H(X, Y)^2 \triangleq \frac{1}{2} \int_{\mathbb{R}} (\sqrt{f_X(x)} - \sqrt{f_Y(x)})^2 dx.$$

In this section we show how the Hellinger distance can be used to obtain efficient concentration inequalities, and derive a bound for the Hellinger distance between S_n and its normal limit. Note that by doing so we establish, for the first time, an explicit rate of convergence for the central limit theorem in terms of the Hellinger distance, which is of independent interest.

Throughout this section, we suppose that the random variables (Y_i) are real valued continuous random variables.

Assumption (\tilde{f}) *The random variable Y_1 is continuous with a probability density function \tilde{f} that is continuously differentiable almost everywhere on its support.*

3.1. Concentration inequalities and the Hellinger distance

Under Assump. (\tilde{f}) the Hellinger distance between S_n and its normal limit is properly defined and upper bounded by 2. We show that this can be used to obtain efficient concentration inequalities for S_n if the assumption Assump. (R, σ) also holds. In this goal, we take $\hat{Q}_n(u)$ to be any known upper bound for $P(S_n \geq u\sigma)$ and $\hat{Q}_n^d(u)$ to be any known upper bound for $P(|S_n| \geq u\sigma)$. Possible choices are the classical Hoeffding or Bernstein bounds Eqs. (1) and (2) or the new Wasserstein-approximation bounds we derive in Sec. 6 (which hold whenever Assump. (R, σ) holds).

Theorem 3 (Efficient Hellinger tail bound) *Suppose that the random variables (Y_i) satisfy Assump. (R, σ) and Assump. (\tilde{f}) . Let $\hat{Q}_n(u)$ be an any arbitrary upper bound to $\mathbb{P}(S_n \geq \sigma u) \leq \hat{Q}_n(u)$. We denote by $H(\cdot, \cdot)$ the Hellinger distance and write $Z \sim N(0, 1)$ as a standard normal random variable. Then the following holds*

$$\mathbb{P}(S_n \geq \sigma u) \leq \Phi^c(u) + \sqrt{2}H(\sigma^{-1}S_n, Z) \left[\sqrt{\hat{Q}_n(u)} + \sqrt{\Phi^c(u)} \right].$$

Moreover let $\hat{Q}_n^d(u)$ be an any arbitrary upper bound to $\mathbb{P}(|S_n| \geq \sigma u) \leq \hat{Q}_n^d(u)$

$$P(|S_n| \geq \sigma u) \leq 2\Phi^c(u) + \sqrt{2}H(\sigma^{-1}S_n, Z) \left[\sqrt{2\Phi^c(u)} + \sqrt{\hat{Q}_n^d(u)} \right].$$

In Thm. 4 we prove that there is a constant K , that we derive, such that $\sqrt{2}H(\sigma^{-1}S_n, Z) \leq \frac{K \log(n)}{\sqrt{n}}$. This directly implies that:

$$\begin{aligned} \mathbb{P}(S_n \geq \sigma u) &\leq \Phi^c(u) + \frac{K \log(n)}{\sqrt{n}} \left[\sqrt{\hat{Q}_n(u)} + \sqrt{\Phi^c(u)} \right] \\ &\leq \Phi^c(u) + \frac{K \log(n)}{\sqrt{n}} \left[\sqrt{e^{-\frac{u^2 \sigma^2}{2R^2}}} + \sqrt{\Phi^c(u)} \right]. \end{aligned}$$

We note that in practice we will want to use the upper bound derived in Thm. 13 for $\hat{Q}_n(u)$ rather than the Hoeffding bound which will provide us with significantly tighter upper bounds. We also remark that the correction term $\frac{K \log(n)}{\sqrt{n}} \left[\sqrt{e^{-\frac{u^2 \sigma^2}{2R^2}}} + \sqrt{\Phi^c(u)} \right]$ is sub-Gaussian in u and decreases as $\frac{\log(n)}{\sqrt{n}}$.

3.2. CLT rate of convergence in terms of the Hellinger distance

In this section we derive the rate of convergence of S_n to its normal limit in terms of the Hellinger distance which is of independent interest as no explicit bound was previously known. To do so we need to impose some moment conditions on the score function of Y_1 which we denote $S_{Y_1}(\cdot)$ and is defined as $S_{Y_1}(\cdot) = -\frac{\tilde{f}'(\cdot)}{\tilde{f}(\cdot)}$.

Assumption (S_{Y_1}, p') Let $p' \geq 1$, we assume that the norm $\|S_{Y_1}(\sigma Y_1)\|_{\frac{2p'}{p'-1}} < \infty$ is finite. Write $\Omega \subset \mathbb{R}$ the support of Y_1 , we suppose that $\tilde{f}|_{\delta\Omega} = 0$.

Let $Z \sim N(0, 1)$ be a standard normal random variable. For all $p \in [1, p']$, we introduce the following notation for $A_{n, \frac{2p}{p-1}}$ and $A_{n, \frac{2p}{p-1}}^*$ defined in Sec. 1.2:

$$C_p \triangleq \min \left\{ \sqrt{\frac{2p}{p-1} - 1} \|S_{Y_1}(\sigma Y_1)\|_{\frac{2p}{p-1}}, A_{n, \frac{2p}{p-1}}^* \|S_{Y_1}(\sigma Y_1)\|_{\frac{2p}{p-1}} + A_{n, \frac{2p}{p-1}} \|S_{Y_1}(\sigma Y_1)\|_2 \right\}.$$

Theorem 4 (Hellinger CLT bound) Let $p' \geq 1$ be a real. Suppose that the random variables (Y_i) satisfy Assump. (R, σ) , Assump. (\tilde{f}) and Assump. (S_{Y_1}, p') . Write $\tilde{R} \triangleq \frac{R}{\sigma}$. Then we obtain that

$$\begin{aligned} \sqrt{2}H(\tilde{S}_n, Z) &\lesssim \inf_{\substack{D > 0 \\ \kappa \geq \frac{\tilde{R}^2}{\sqrt{n}D}}} \left\{ \frac{\sqrt{e}\tilde{R}^2(e^{2\kappa}-1)}{\sqrt{3}D\sqrt{n}\kappa} + \frac{\log(n/D^2)}{4\sqrt{2}\sqrt{n}}(\tilde{R}-1) - \frac{1}{2}\log(1 - \frac{D}{\sqrt{n}}) \right. \\ &\quad \left. + \inf_{p' \geq p \geq 1} C_p \left\{ \frac{\sqrt{2p-1}\tilde{R}}{\sqrt{n}\kappa} + \frac{2\sqrt{e}\sqrt{2p-1}\tilde{R}}{\sqrt{3n}\kappa} \left[2 - \frac{2}{3\tilde{R}} \right] \right. \right. \\ &\quad \left. \left. \times \left[e^{2(2p-1)\kappa} - 1 - 2(2p-1)\kappa \right] + \frac{\tilde{R}^2}{2n\kappa M_{n,\kappa}} \left[U_{n,2p} + \tilde{U}_{n,2p} \right] \right\} \right\}. \end{aligned}$$

Moreover they are constants K_1 , K_2 and K_3 given in Eq. (24) such that

$$\begin{aligned} \sqrt{2}H(\sigma^{-1}S_n, Z) &\lesssim \frac{K_1}{D\sqrt{n}} + \frac{D}{2\sqrt{n}} + \frac{\log(n/D^2)}{4\sqrt{2}\sqrt{n}}(\tilde{R}-1) + \|S_{Y_1}(\sigma Y_1)\|_2 \left\{ \frac{K_2\tilde{R}}{\sqrt{n}} + \frac{K_3}{\sqrt{n}} \right\} \\ &= O\left(\frac{\log(n)}{\sqrt{n}}\right). \end{aligned}$$

Note that an exact and a tighter computable bound can be found in Thm. 22. The latter is less interpretable but is useful for deriving as tight as possible quantiles bounds using Thm. 3. The proof can be found in App. C.

3.3. General bound for the Hellinger distance

We remark that Thm. 4 establishes a rate of convergence for the Hellinger distance under moment conditions for the score function of Y_1 . This result is novel. In this section we present the general bound we established and used to derive Thm. 4. Our proof technique relies on the Ornstein-Ulbeck interpolation, the heat equation, and the Stein exchangeable pair method. The latter takes inspiration from the ideas Bonis (2015) adapted to obtain a rate in the Wassertein distance. This provides us with an easily computable bound that only depends on the moments of the score function and the moments of the random variable itself. This is to contrast with the known bounds for the

KL divergence (which upper-bounds the square of the Hellinger distance), which are either not in closed form (Bobkov et al., 2013), require conditions on the spectral gap (Artstein et al., 2004) or depend on the moment of the Stein kernel (Ledoux et al., 2015) which is often significantly harder to upper-bound and does not always exist.

Let S be a real valued random variable with PDF $f_S \in \mathbb{C}^1(\mathbb{R})$ with support $\Omega_S \subset \mathbb{R}$. We write its score function $S_S(\cdot)$. Let $h_k(x) \triangleq e^{x^2/2} \frac{\partial^k}{\partial x} e^{-x^2/2}$ designate the Hermite polynomials, and denote $H_k \triangleq h_k(Z)$ where $Z \sim \mathcal{N}(0, 1)$ is a standard normal. We write the interpolated process as

$$X_t^S \triangleq e^{-t} S + \sqrt{1 - e^{-2t}} Z.$$

We denote by f_t^S the PDF of X_t^S and define $S_{X_t^S}$ to be the score function of X_t^S . We remark that for all $t > 0$ the function $f_t^S \in \mathbb{C}^\infty(\mathbb{R})$ is smooth and we write $I_t^S(\cdot) = \frac{f_t^{S''}(\cdot)}{f_t^S(\cdot)}$. For every $t \geq 0$ choose S^t to be a random variable such that (S, S^t) is an exchangeable pair. For all $t > 0$ we define

$$\tau_t^S \triangleq \sum_{k=1}^{\infty} \frac{e^{-kt}}{k! \sqrt{1 - e^{-2t}k}} \mathbb{E}((S^t - S)^k | S) H_{k-1}(Z); \quad (6)$$

$$\tau_t'^S \triangleq \sum_{k=1}^{\infty} \frac{e^{-kt}}{k! \sqrt{1 - e^{-2t}k}} \mathbb{E}((S^t - S)^k | S) H_k(Z). \quad (7)$$

Theorem 5 (Bounding Hellinger distance to a Gaussian) *Assume that S is standardized meaning that $\mathbb{E}(S) = 0$ and that $\text{var}(S) = 1$. Suppose that $f_S|_{\delta\Omega_S} = 0$. Then the following holds*

$$\begin{aligned} \sqrt{2}H(S, Z) &\leq \int_0^\infty \|I_t^S(X_t^S) - (X_t^S)^2 + 1 - X_t^S(S_{X_t^S} - X_t^S)\|_2 dt \\ &\leq \inf_{D>0} \left\{ \|S_S(S)\|_{\frac{2p}{p-1}} \int_0^{-\frac{1}{2} \log(1 - \frac{D}{\sqrt{n}})} \left\| \frac{Ze^{-2t}}{\sqrt{1 - e^{-2t}}} - e^{-t}S + \tau_t^S \right\|_{2p} dt \right. \\ &\quad \left. + \int_{-\frac{1}{2} \log(1 - \frac{D}{\sqrt{n}})}^\infty \left\| \frac{e^{-2t}}{1 - e^{-2t}} (Z^2 - 1) - \frac{e^{-t}}{\sqrt{1 - e^{-2t}}} SZ + \tau_t'^S \right\|_2 dt \right. \\ &\quad \left. - \frac{1}{2} \log(1 - \frac{D}{\sqrt{n}}) \right\}. \end{aligned}$$

This new upper-bound for the Hellinger distance is the key to the proof of Thm. 4. Outside of giving us a bound for $H(\tilde{S}_n, Z)$, this result can be used to obtain bounds on $H(S, Z)$ for any other random variable S . We remark that this upper bound will depend on the moments of S and of its score function. To do so one can pick any specific $D > 0$ and $p > 1$ and compute the corresponding bound. Note that we use this idea to obtain Thm. 4 and that any arbitrary choice of D and p give us that $\sqrt{2}H(\tilde{S}_n, Z)$ decreases at a rate of $O\left(\frac{\log(n)}{\sqrt{n}}\right)$. However depending on the choices of those coefficients the constants in the bound of $H(\tilde{S}_n, Z)$ will change. We note that this result plays a similar role to bounds that had been developed for the KL-divergence or the Wassertein distance see (Barron, 1986; Otto and Villani, 2000; Ledoux et al., 2015). The proof is presented in App. D.

In Bonis (2015) similar ideas could be exploited to obtain a bound for the p -Wassertein distance $\mathcal{W}_p(S, Z)$. Notably, it was established that

$$\mathcal{W}_p(S, Z) \leq \int_0^\infty \left\| \frac{e^{-2t}}{\sqrt{1 - e^{-2t}}} Z - e^{-t}S - \tau_t^S \right\|_p dt.$$

The Wassertein distance is a distance on the underlying geometry of the random variables. This is to contrast with the Hellinger distance which is an information-theoretic distance that intimately depends on the PDF and its smoothness. This difference explains why the latter can be upper-bounded independently from the score function.

3.4. Proof sketch for Thm. 3

In each of the proof sketches to follow, we assume, without loss of generality, that $\sigma = 1$.

The proof of Thm. 3 is a consequence of the Cauchy-Swartz inequality. Indeed let $\varphi(\cdot)$ be the PDF of a standard normal random variable, and \tilde{f}_n the PDF of S_n . If we write $h(t) = \frac{\tilde{f}_n(t)}{\varphi(t)}$ then

$$\begin{aligned} \mathbb{P}(S_n \geq u) - \Phi^c(u) &= \int_u^\infty (h(t) - 1)\varphi(t)dt \\ &\leq \int_u^\infty (\sqrt{h(t)} - 1)(\sqrt{h(t)} + 1)\varphi(t)dt \\ &\stackrel{(a)}{\leq} \sqrt{\int_u^\infty (\sqrt{h(t)} - 1)^2 \varphi(t)dt} \left[\sqrt{\Phi^c(u)} + \sqrt{\int_u^\infty h(t)\varphi(t)dt} \right], \end{aligned}$$

where (a) is consequence of Cauchy-Swartz inequality. Using the definition of $H(\cdot, \cdot)$ and $\hat{Q}_n(\cdot)$ we obtain the desired result.

3.5. Proof sketch for Thm. 5

The proof of Thm. 5 relies on the Ornstein-Ulbeck interpolation, the heat equation and the Stein exchangeable pair. To make this precise we define the interpolated process as $X_t^S \triangleq e^{-t}S + \sqrt{1 - e^{-2t}}Z$ where Z is an independent standard normal random variable. As t varies from 0 to ∞ X_t^S evolves from S to Z . Indeed we remark that $X_0^S = S$ and $\lim_{t \rightarrow \infty} X_t^S \rightarrow Z$. For all $t > 0$, as X_t^S is the sum of S with an independent normal it is a continuous random variable that admits a PDF f_t^S that is smooth. We write $h_t^S(x) \triangleq \frac{f_t^S(x)}{\varphi(x)}$.

Fix an arbitrary $T < \infty$. By definition we have

$$H(S, X_T^S)^2 = 1 - \int_{\mathbb{R}} \sqrt{f_S(x)} \sqrt{f_T^S(x)} dx.$$

We remark that for all $x \in \mathbb{R}$ the function $t \rightarrow f_t^S(x)$ is differentiable. Therefore we have $\int_0^T \partial_t \sqrt{f_t^S(x)} dt = \sqrt{f_T^S(x)} - \sqrt{f_S(x)}$ which implies that

$$H(S, X_T^S)^2 = -\frac{1}{2} \int_0^T \int_{\mathbb{R}} \sqrt{f_S(x)} \partial_t \sqrt{f_t^S(x)} dx dt. \quad (8)$$

We use this characterization of the Hellinger distance to bound it in terms of information-theoretic quantities. Indeed, by the chain rule and the heat equation we obtain that

$$\begin{aligned} \partial_t \sqrt{f_t^S(x)} &= \frac{1}{2\sqrt{f_t^S(x)}} \partial_t f_t^S(x) = \frac{\varphi(x)}{2\sqrt{f_t^S(x)}} \partial_t \frac{f_t^S(x)}{\varphi(x)} \\ &= \frac{\varphi(x)}{2\sqrt{f_t^S(x)}} \partial_t h_t^S(x) = \frac{\varphi(x)}{2\sqrt{f_t^S(x)}} [h_t^{S''}(x) - x h_t^{S'}(x)] \\ &= \frac{f_t^S(x)}{2\sqrt{f_t^S(x)}} \left[\frac{h_t^{S''}(x)}{h_t^S(x)} - x \frac{h_t^{S'}(x)}{h_t^S(x)} \right]. \end{aligned}$$

Therefore by combining this with Eq. (8) we obtain that

$$H(S, X_T^S)^2 = -\frac{1}{2} \int_0^T \mathbb{E} \left(\frac{\sqrt{f_S(X_t^S)}}{2\sqrt{f_t^S(X_t^S)}} \left[\frac{h_t^{S''}(X_t^S)}{h_t^S(X_t^S)} - X_t^S \frac{h_t^{S'}(X_t^S)}{h_t^S(X_t^S)} \right] \right) dt.$$

Using the fact that $\mathbb{E}\left(\frac{h_t^{S''}(X_t^S)}{h_t^S(X_t^S)} - X_t^S \frac{h_t^{S'}(X_t^S)}{h_t^S(X_t^S)}\right) = 0$, the Cauchy-Swartz inequality and the definition of the Hellinger distance we then establish that

$$H(S, X_T^S)^2 \leq \frac{1}{\sqrt{2}} \sup_{t \leq T} H(S, X_t^S) \int_0^T \left\| \frac{h_t^{S''}(X_t^S)}{h_t^S(X_t^S)} - X_t^S \frac{h_t^{S'}(X_t^S)}{h_t^S(X_t^S)} \right\|_2 dt.$$

As this holds for any arbitrary $T < \infty$ this implies that

$$H(S, Z) \leq \frac{1}{\sqrt{2}} \int_0^\infty \left\| \frac{h_t^{S''}(X_t^S)}{h_t^S(X_t^S)} - X_t^S \frac{h_t^{S'}(X_t^S)}{h_t^S(X_t^S)} \right\|_2 dt.$$

This bound can be seen as the equivalent of the De Bruijn identity (see, e.g., [Barron, 1986](#)) for the KL-divergence or similar expressions obtained for p -Wassertein distance in the celebrated work of [Otto and Villani \(2000\)](#) and [Ledoux et al. \(2015\)](#). As we do not know h_t^S , the right hand side of this equation is not easy to directly control. To go around this issue we re-express $\frac{h_t^{S''}(x)}{h_t^S(x)} - x \frac{h_t^{S'}(x)}{h_t^S(x)}$ in terms of the score function $S_{X_t^S}^S$ and I_t^S , which we can do as a simple consequence of the chain-rule. The advantage of this is that using the Stein identity we can re-express the score function and I_t^S . Indeed we remark that $S_t^S(X_t^S)$ is the unique X_t^S -measurable random variable such that for all $\phi \in \mathbb{C}^\infty(\mathbb{R})$ we have

$$\mathbb{E}(S_t^S(X_t^S)\phi(X_t^S)) = \mathbb{E}(\phi'(X_t^S)).$$

Similarly we remark that $I_t^S(X_t^S)$ is the unique X_t^S -measurable random variable such that for all $\phi \in \mathbb{C}^\infty(\mathbb{R})$ we have

$$\mathbb{E}(I_t^S(X_t^S)\phi(X_t^S)) = \mathbb{E}(\phi''(X_t^S)).$$

Therefore using the Stein identity we can prove that

$$\begin{aligned} S_{X_t^S}^S(X_t^S) &= \begin{cases} e^t \mathbb{E}(S_S(S)|X_t^S) \\ \frac{1}{\sqrt{1-e^{-2t}}} \mathbb{E}(Z|X_t^S) \end{cases} ; \\ I_t^S(X_t^S) &= \begin{cases} \frac{e^t}{\sqrt{1-e^{-2t}}} \mathbb{E}(S_S(S)Z|X_t^S) \\ \frac{1}{1-e^{-2t}} \mathbb{E}(Z^2 - 1|X_t^S). \end{cases} \end{aligned}$$

The last step of the proof is to notice that $\tau_t^{'S}$ and τ_t^S are centered with respect to the conditional expectation: $\mathbb{E}(\tau_t^S|X_t^S) = \mathbb{E}(\tau_t^{'S}|X_t^S) = 0$. Indeed we establish that for all smooth function $\Phi \in \mathbb{C}^\infty(\mathbb{R})$ we have

$$\begin{aligned} \mathbb{E}(\tau_t^{'S}\Phi(X_t^S)) &= \mathbb{E}(\Phi(X_t^S)) - \mathbb{E}(\Phi(e^{-t}S^t + \sqrt{1-e^{-2t}}Z)) \\ &\stackrel{(a)}{=} 0. \end{aligned}$$

where to obtain (a) we use the fact that (S, S^t) is an exchangeable pair. As this holds for arbitrary Φ this directly implies that $\mathbb{E}(\tau_t^{'S}|X_t^S) = 0$. Similarly we prove that $\mathbb{E}(\tau_t^S|X_t^S) = 0$. This establishes [Thm. 5](#).

3.6. Proof sketch for Thm. 4

To prove Thm. 4 we build an exchangeable pair (S_n, S_n^t) in the following way: We first start by drawing an independent copy (Y'_i) of (Y_i) . We set a $\kappa > 0$ and write $c(t) = \sqrt{\kappa}\sqrt{1 - e^{-2t}}$. Then for every $t > 0$ we choose an index $I \sim \text{unif}\{1, \dots, n\}$ at random and change the random variable Y_i by its independent copy Y'_i if and only if both $|Y_i/\sqrt{n}| \leq c(t)$ and $|Y'_i/\sqrt{n}| \leq c(t)$:

$$S_n^t \triangleq S_n + \frac{1}{\sqrt{n}} \mathbb{I}(|Y_i/\sqrt{n}|, |Y'_i/\sqrt{n}| \leq c(t)) (Y'_i - Y_i).$$

As the observations (Y_i) are i.i.d we remark that (S_n, S_n^t) is indeed an exchangeable pair. The advantage of this choice of exchangeable pair is that we have

$$\mathbb{E}(S_n^t | S_n) \approx S_n, \quad \frac{1}{2} \mathbb{E}((S_n^t - S_n)^2 | S_n) \approx 1.$$

This will allow us to obtain small upper bounds for $\| \frac{e^{-2t}}{\sqrt{1-e^{-2t}}} Z - e^{-t} \tilde{S}_n + \tau_t \|_{2p''}$ and $\| \frac{e^{-2t}}{1-e^{-2t}} (Z^2 - 1) - \frac{e^{-t}}{\sqrt{1-e^{-2t}}} \tilde{S}_n Z + \tau'_t \|_2$ and therefore control $H(S_n, Z)$.

4. Efficient Concentration with Stein Kernels

In the previous section, we showed how we can use the Hellinger distance to obtain an efficient concentration inequality when Assump. (\tilde{f}) holds. In this section, we will see how one can also obtain efficient concentration inequalities under the alternative condition than the random variables (Y_i) admit a Stein kernel. This can be useful notably for a certain number of discrete random variables that have a Stein kernel but not a score function. However, we note that not all random variables that satisfy Assump. (\tilde{f}) will also admit a Stein kernel or have it nicely controlled. Therefore the two approaches are complementary. Finally, we will see in Thm. 8 how when both exist we can exploit this to obtain an even tighter concentration inequality.

The notion of a Stein kernel was first used in [Cacoullos and Papathanasiou \(1992\)](#), although the term ‘‘Stein kernel’’ was introduced by [Ley et al. \(2017\)](#). A function τ is called the Stein kernel of a standardized random variable X if for all functions $\phi \in \mathbb{C}^1(\mathbb{R})$ we have:

$$\mathbb{E}(\phi(X)X) = \mathbb{E}(\tau(X)\phi'(X)).$$

If X is a continuous random variable with PDF f_X , and if f_X has as support a (potentially infinite) connected interval (a, b) then, under some regular stability conditions on f_X , the Stein kernel is given by $\tau(x) = \frac{1}{f_X(x)} \int_x^\infty t f_X(t) dt$ ([Ley et al., 2017](#)). Notably, if X is a standard normal random variable then its Stein kernel is given by $\tau(X) = 1$. When the random variable X does not possess such a density, the Stein kernel sometimes still exists, this is for example the case if the random variables satisfy a Poincare inequality ([Fathi, 2019](#)). In other cases, the existence has been established on a case-by-case basis.

This is notably true if X is a symmetric random variable with support $\{-1, +1\}$ ([Chatterjee, 2012](#)). See [Ley et al. \(2017\)](#) for other examples of Stein kernels and discussion about its existence. However, it is important to note that, in general, a Stein kernel does not always exist (even when X is continuous), and general conditions for its existence are not well determined.

The Stein kernel can measure the distance between X and a standard normal random variable by defining the *Stein discrepancy*:

$$\mathcal{S}(X||Z) \triangleq \|\tau(X) - 1\|_2.$$

Similarly for $p \geq 2$, we can define the following distance $\mathcal{S}_p(X||Z) \triangleq \|\tau(X) - 1\|_p$. While the Stein kernel can be hard to control, it has proven to be a useful quantity. Notably, [Ledoux et al. \(2015\)](#) has explored how it can be used to bound the p -Wassertein distance and the KL divergence between a random variable and a standard normal random variable. It has also been used in [Ledoux et al. \(2015\)](#) to upper-bound the moments of Lipchitz functions of S_n . In this section, we established how it can also be used to obtain efficient concentration inequalities.

Assumption $(\tau_{\tilde{Y}_1}, p')$ Assume that $\sigma^{-1}Y_1$ admit a Stein kernel $\tau_{\tilde{Y}_1}$ and that $\|\tau_{\tilde{Y}_1}(\sigma^{-1}Y_1)\|_{p'} < \infty$.

In the following we are going to take $\hat{Q}_n(x)$ to be such that for all $x > 0$

$$\sup_{t \geq 0} \mathbb{P}(e^{-t}\tilde{S}_n + \sqrt{1 - e^{-2t}}Z \geq x) \leq \hat{Q}_n(x).$$

We prove in Lem. 28 that $\hat{Q}_n(x)$ can be chosen to be the Hoeffding bound for R^2 sub-Gaussian random variables or the upper bound obtained in Thm. 13. For all $p > 1$ we write

$$D_p(\tau_{\tilde{Y}_1}) \triangleq \min \left\{ A_p \|\tau_{\tilde{Y}_1}(\sigma^{-1}Y_1) - 1\|_2 + A_{n,p}^* \|\tau_{\tilde{Y}_1}(\sigma^{-1}Y_1) - 1\|_p, \sqrt{p-1} \|\tau_{\tilde{Y}_1}(\sigma^{-1}Y_1) - 1\|_p \right\}.$$

Theorem 6 (Efficient Stein kernel tail bound) Let $p' \geq 2$. Suppose that the random variables (Y_i) satisfy Assump. [\(R, \$\sigma\$ \)](#) and Assump. $(\tau_{\tilde{Y}_1}, p')$. Choose $\alpha > 0$ and let $p \geq 1$ and $q \geq 2$. Define $u_\alpha(x) \triangleq \frac{1}{\alpha^2} \int_x^{x+\alpha} \int_t^{t+\alpha} \mathbb{I}(z \geq u) dz dt$ and let $Z \sim N(0, 1)$ then we have

$$\begin{aligned} & \mathbb{P}(S_n \geq \sigma u) \\ & \leq \mathbb{E}(u_\alpha(Z)) + \frac{D_p(\tau_{\tilde{Y}_1})}{\sqrt{n}} \min \left\{ \|Z\|_p \left[\|u'_\alpha(Z)\|_{\frac{p}{p-1}} + \frac{\pi \mathcal{W}_{\frac{pq}{(p-1)}}(G, \tilde{S}_n) \hat{Q}_n(u-2\alpha)^{\frac{(p-1)(q-1)}{pq}}}{4\alpha^2} \right] \right. \\ & \quad \left. \frac{1}{2\alpha^2} \hat{Q}_n(u-2\alpha)^{\frac{(p-1)}{p}} \right\} \\ & < \mathbb{P}(Z \geq u-2\alpha) + \frac{D_p(\tau_{\tilde{Y}_1})}{\sqrt{n}} \min \left\{ \|Z\|_p \left[\frac{1}{\alpha} \mathbb{P}(Z \in [u-2\alpha, u])^{\frac{p-1}{p}} + \frac{\pi}{4\alpha^2} \mathcal{W}_{\frac{pq}{(p-1)}}(G, \tilde{S}_n) \right. \right. \\ & \quad \left. \left. \hat{Q}_n(u-2\alpha)^{\frac{(p-1)(q-1)}{pq}} \right] \right. \\ & \quad \left. \frac{1}{2\alpha^2} \hat{Q}_n(u-2\alpha)^{\frac{(p-1)}{p}} \right\}. \end{aligned}$$

Remark 1 We note that a tighter but less interpretable bound is presented in Lem. 27. Moreover we remark that as p increases the terms $\|u'_\alpha(Z)\|_{\frac{p}{p-1}}$ and $\hat{Q}_n(u-2\alpha)^{\frac{(p-1)(q-1)}{pq}}$ decrease, and that both are asymptotically bounded by respectively $\mathbb{P}(Z \in [u-2\alpha, u]) \leq 2\alpha\varphi(u-2\alpha)$ and $e^{-\frac{(u-2\alpha)^2}{2R^2}}$.

We remark that similarly to Thm. 5 we can obtain a similar efficient two-sided tail bound. This is presented in Lem. 25. The proof can be found in App. E. See Sec. 4.1 for a proof sketch.

Let us compare this result to what has been previously obtained in the literature. Under the condition that the Stein kernel $\tau_{\tilde{Y}_1}(\cdot) - 1$ is almost surely bounded by $\mathcal{S}_\infty(\sigma^{-1}Y_1||Z)$ then Chatterjee (2012) proved that there is $G \sim N(0, 1)$ normally distributed such that for all $\theta > 0$

$$\mathbb{E}(e^{\theta|\tilde{S}_n - G|}) \leq 2e^{2\theta^2 \mathcal{S}_\infty(\sigma^{-1}Y_1||Z)^2}.$$

G is called a strong embedding of \tilde{S}_n . If the Stein kernel is not bounded or does not exist Bhattacharjee and Goldstein (2016) but $\mathbb{E}(Y_1^3) = 0$, then Bhattacharjee and Goldstein (2016) proposed to study a slightly smoothed empirical average and obtained a similar inequality. Using a union bound and a Markov inequality, this can be used to obtain efficient concentration inequalities for S_n :

Theorem 7 (Alternative Stein kernel tail bound) Suppose that the random variables satisfy Assump. (R, σ) and Assump. $(\tau_{\tilde{Y}_1}, p')$ with $p' = \infty$.

Then for all $u > 0$ and $0 < \alpha < u/2$ the following holds

$$\mathbb{P}(S_n \geq \sigma u) \leq \Phi^c(u - 2\alpha) + 2e^{-\frac{n\alpha^2}{2\mathcal{S}_\infty(\sigma^{-1}Y_1||Z)^2}}. \quad (9)$$

We remark that Thm. 7 requires stronger moment conditions on the Stein kernel than Thm. 6 does. Moreover for illustration if we choose $p = q = 2$ (which is not necessarily the optimal choice) then Thm. 6 tells us that

$$\begin{aligned} \mathbb{P}(S_n \geq \sigma u) \\ \leq \mathbb{E}(u_\alpha(Z)) + \frac{1}{\sqrt{n}} \|\tau_{\tilde{Y}_1}(\sigma^{-1}Y_1) - 1\|_2 \left[\|u'_\alpha(Z)\|_2 + \frac{\pi \mathcal{W}_4(G, \tilde{S}_n)}{4\alpha^2} \hat{Q}_n(u - 2\alpha)^{\frac{1}{4}} \right]. \end{aligned} \quad (10)$$

To contrast this with the bound obtained in Thm. 7, we note that $\mathbb{E}(u_\alpha(Z)) \leq \Phi^c(u - 2\alpha)$. Moreover we also remark that

$$\|u'_\alpha(Z)\|_2 \leq \alpha^{-1} \sqrt{\mathbb{P}(Z \in [u - 2\alpha, u])}.$$

Therefore we remark that the right-hand side of Eq. (10) is Sub-Gaussian in u . In contrast we notice that the second term in Eq. (9), $e^{-\frac{n\alpha^2}{2\mathcal{S}_\infty(\sigma^{-1}Y_1||Z)^2}}$, does not depend on u . Therefore for large u and moderate size n the bound in Thm. 6 will tend to be tighter. Finally using Lem. 34 we know that there is a constant K such that $\mathcal{W}_4(G, \tilde{S}_n) \leq \frac{\tilde{R}K}{\sqrt{n}}[1 + o_n(1)]$.

We observe that if Assump. (\tilde{f}) also holds then by using Ledoux et al. (2015) we can exploit Thm. 3 to obtain an efficient concentration inequality that is exactly sub-Gaussian in u .

Theorem 8 (Efficient Hellinger-Stein kernel tail bound) Set $p' = 2$. Suppose that the random variables (Y_i) satisfy Assump. (R, σ) , Assump. (\tilde{f}) , Assump. (S_{Y_1}, p') and Assump. $(\tau_{\tilde{Y}_1}, p')$. Let $\hat{Q}_n(u)$ and $\hat{Q}_n^d(u)$ be an any arbitrary upper bound to respectively $\mathbb{P}(S_n \geq \sigma u) \leq \hat{Q}_n(u)$ and $\mathbb{P}(|S_n| \geq \sigma u) \leq \hat{Q}_n^d(u)$. Suppose that $\|S_{Y_1}(\sigma Y_1)\|_2 < \infty$ and that $\mathcal{S}(\sigma^{-1}Y_1||Z) < \infty$. Then for all $u \geq 0$ the following bounds hold

$$\begin{aligned} \mathbb{P}(S_n \geq \sigma u) \leq \Phi^c(u) + \frac{\sqrt{2}\mathcal{S}_2(\sigma^{-1}Y_1||Z)}{\sqrt{n}} \sqrt{\log \left(1 + \frac{n\|S_{Y_1}(\sigma Y_1) - \sigma^{-1}Y_1\|_2^2}{\mathcal{S}_2(\sigma^{-1}Y_1||Z)^2} \right)} \\ \times \left[\sqrt{\hat{Q}_n(u)} + \sqrt{\Phi^c(u)} \right]; \end{aligned}$$

and

$$\begin{aligned} \mathbb{P}(|S_n| \geq \sigma u) &\leq 2\Phi^c(u) + \frac{\sqrt{2}S_2(\sigma^{-1}Y_1||Z)}{\sqrt{n}} \sqrt{\log\left(1 + \frac{n\|S_{Y_1}(\sigma Y_1) - \sigma^{-1}Y_1\|_2^2}{S_2(\sigma^{-1}Y_1||Z)^2}\right)} \\ &\quad \times \left[\sqrt{\hat{Q}_n^d(u)} + \sqrt{2\Phi^c(u)} \right]. \end{aligned}$$

See Sec. 4.2 for a proof sketch and App. E.3 for the proof. We note that the rate obtained in Thm. 8 is sub-Gaussian in u and that the reminder terms decreases at a rate of $\sqrt{\log(n)/n}$. Therefore if in addition to the Stein Kernel we can also control the score function then we can obtain a tighter rate than in Thm. 7 and Thm. 6.

4.1. Proof sketch for Thm. 6

We note that $u_\alpha(x) \leq \mathbb{I}(x \leq u)$. By the monotony of the expectation this directly implies that $\mathbb{E}(u_\alpha(\tilde{S}_n)) \leq \mathbb{P}(\tilde{S}_n \geq u)$. Therefore if we obtain a bound on $|\mathbb{E}(u_\alpha(\tilde{S}_n)) - \mathbb{E}(u_\alpha(Z))|$, we can use it to obtain a bound on $\mathbb{P}(\tilde{S}_n \geq u)$. The rest of the proof of Thm. 6 consists of using the Ornstein-Uhlenbeck interpolation and the heat equation.

To see this, we denote by $\tau_{\tilde{S}_n}(\cdot)$ the Stein kernel of \tilde{S}_n and take $Z \sim N(0, 1)$ to be an independent standard normal random variable. We define $P_t u_\alpha(x) \triangleq \mathbb{E}_Z(u_\alpha(e^{-t}x + \sqrt{1 - e^{-2t}}Z))$. We remark that $t \rightarrow P_t u_\alpha$ interpolates between $P_0 u_\alpha(x) = u_\alpha(x)$ and $P_\infty u_\alpha(x) = \mathbb{E}(u_\alpha(Z))$. Therefore we have:

$$\begin{aligned} |\mathbb{E}(u_\alpha(\tilde{S}_n)) - \mathbb{E}(u_\alpha(Z))| &= \left| \int_0^\infty \partial_t \mathbb{E}(P_t u_\alpha(\tilde{S}_n)) dt \right| \\ &\stackrel{(a)}{\leq} \left| \int_0^\infty \mathbb{E}(\Delta P_t u_\alpha(\tilde{S}_n) - \tilde{S}_n \nabla P_t u_\alpha(\tilde{S}_n)) dt \right| \\ &\stackrel{(b)}{\leq} \left| \int_0^\infty \mathbb{E}(\Delta P_t u_\alpha(\tilde{S}_n)(1 - T_{\tilde{S}_n}(\tilde{S}_n))) dt \right| \end{aligned}$$

where to obtain (a) we used the heat equation satisfied by the Ornstein-Uhlenbeck process and where (b) is a consequence of the definition of Stein kernels.

It is well known (see e.g. (Ledoux et al., 2015)) that $\tau_{\tilde{S}_n}(\tilde{S}_n) = \mathbb{E}(\frac{1}{n} \sum_{i=1}^n \tau_{\tilde{Y}_1}(Y_i) | \tilde{S}_n)$. Therefore as $\mathbb{E}(\tau_{\tilde{Y}_1}(Y_1)) = 1$ for all $p \geq 1$ there is a constant K_p such that

$$\|\tau_{\tilde{S}_n}(\tilde{S}_n) - 1\|_p \leq \left\| \frac{1}{n} \sum_{i=1}^n \tau_{\tilde{Y}_1}(Y_i) - 1 \right\|_p \leq \frac{K_p}{\sqrt{n}} \|\tau_{\tilde{Y}_1}(Y_1) - 1\|_p.$$

Therefore if we can control the size of $\partial_t^2 P_t u_\alpha(\tilde{S}_n)$ then we will obtain that $|\mathbb{E}(u_\alpha(\tilde{S}_n)) - \mathbb{E}(u_\alpha(Z))| = O(\frac{1}{\sqrt{n}})$. The remainder of the proof consists of controlling $\Delta P_t u_\alpha(\tilde{S}_n)$. This is what we do thanks to the Stein identity and the Wassertein distance.

4.2. Proof sketch of Thm. 8

The proof of Thm. 8 is based on Thm. 3 which guarantees that

$$\mathbb{P}(S_n \geq \sigma u) \leq \Phi^c(u) + \sqrt{2}H(\sigma^{-1}S_n, Z) \left[\sqrt{\hat{Q}_n(u)} + \sqrt{\Phi^c(u)} \right];$$

and Theorem 2.2 of Ledoux et al. (2015) which upper bounds the KL-divergence $KL(\tilde{S}_n || Z)$ in terms of the Stein kernel.

5. Efficient Concentration with Zero-Bias Couplings

In the prior two sections, we showed how to obtain efficient concentration given auxiliary knowledge of a score function or Stein kernel. In the remainder, we will establish efficient concentration without any additional knowledge beyond Assump. (R, σ) . We achieve this in the present section by tightly bounding a non-uniform Kolmogorov distance using *zero-bias couplings*. As in [Chen et al. \(2011, Prop. 2.1\)](#), we say that S^* has the *zero-bias distribution* for a mean-zero random variable S with $\text{Var}(S) < \infty$ if the distribution of S^* is absolutely continuous with Lebesgue density $p^*(x) = \mathbb{E}[S \mathbb{I}[S > x]] / \text{Var}(S)$ or, equivalently, if $\sigma^2 \mathbb{E}[f'(S^*)] = \mathbb{E}[Sf(S)]$ for all absolutely continuous f with $\mathbb{E}[|Sf(S)|] < \infty$. Our main result, proved in App. F, uses a close coupling of S_n and its zero-biased counterpart to establish efficient concentration.

Theorem 9 (Efficient zero-bias tail bound) *Suppose $S_n = \sum_{i=1}^n X_i$ for independent $(X_i)_{i \geq 1}$ satisfying $\mathbb{E}[X_i] = 0$, $\text{Var}(X_i) = \sigma_i^2$, $\sum_{i=1}^n \sigma_i^2 = \sigma^2$, and $|X_i| \leq \frac{R}{\sqrt{n}}$ almost surely. Define the auxiliary variable*

$$S'_n \triangleq S_n + X'_I - X_I \quad \text{for} \quad X'_i \triangleq X_i^* + U(X_i - X_i^*) \quad \text{and} \quad \mathbb{P}(I = i) = \frac{\sigma_i^2}{\sigma^2} \mathbb{I}[1 \leq i \leq n],$$

where each X_i^* has the zero-bias distribution of X_i , and $U \sim \text{Unif}([0, 1])$, I , and $(X_i^*)_{i \geq 1}$ are mutually independent and independent of $(X_i)_{i \geq 1}$. Then, for all $u \geq 0$ and $\lambda \in [0, 1]$,

$$\begin{aligned} \mathbb{P}(S_n > \sigma u + \frac{2R}{\sqrt{n}}) &\leq \frac{\sigma \sqrt{n}}{\sigma \sqrt{n+2R} \delta_u} \Phi^c(u) + \frac{2R}{\sigma \sqrt{n+2R} \delta_u} (h_u(u) - h_u(\lambda u)) \mathbb{P}(S'_n > \lambda \sigma u) \\ &\quad + \frac{2R}{\sigma \sqrt{n+2R} \delta_u} h_u(\lambda u) \end{aligned}$$

where, for all $w \leq u$,

$$h_u(w) \triangleq (w + (1 + w^2) \sqrt{2\pi} \exp(\frac{w^2}{2}) \Phi(w)) \Phi^c(u) \quad \text{and} \quad (11)$$

$$\delta_u \triangleq h_u(u) - \left(\frac{2}{u + \sqrt{u^2 + 8/\pi}} - \frac{8u}{\pi(u + \sqrt{u^2 + 8/\pi})^2} \right) \Phi(u). \quad (12)$$

If, in addition, $(X_i)_{i=1}^{n+1}$ are identically distributed, then

$$\begin{aligned} &\mathbb{P}(S_n > \sigma u \frac{\sqrt{n+1}}{\sqrt{n}} + \frac{R}{\sqrt{n}}) \\ &\leq \frac{\sigma \sqrt{n+1}}{\sigma \sqrt{n+1+2R} \delta_u} \Phi^c(u) + \frac{2R}{\sigma \sqrt{n+1+2R} \delta_u} (h_u(u) - h_u(\lambda u)) \mathbb{P}(S_n + X'_{n+1} > \lambda \sigma u) \\ &\quad + \frac{2R}{\sigma \sqrt{n+1+2R} \delta_u} h_u(\lambda u). \end{aligned}$$

Our next result, proved in App. G, shows how to convert concentration inequalities for bounded variables into valid tail bounds for S'_n and $S_n + X'_{n+1}$.

Proposition 10 (Properties of S'_n) *Under the notation and assumptions of Thm. 9, the following claims hold for each $i \geq 1$ and $t \in \mathbb{R}$.*

1. $|\mathbb{E}[X'_i]| \leq \frac{R}{4\sqrt{n}}$, $\text{Var}(X'_i) \in \frac{\sigma_i^2}{3} + [\frac{R^2}{36n}, \frac{R^2}{9n}]$, $|X'_i| \leq \frac{R}{\sqrt{n}}$ almost surely.
2. $\mathbb{P}(S'_n > t + \frac{R}{4\sqrt{n}}) \leq \sum_{i=1}^n \frac{\sigma_i^2}{\sigma^2} \mathbb{P}(X'_i - \mathbb{E}[X'_i] + \sum_{j \neq i} X_j > t)$.

3. If, in addition, $(X_i)_{i=1}^{n+1}$ are identically distributed, then $S_n + X'_{n+1} \stackrel{d}{=} \frac{\sqrt{n+1}}{\sqrt{n}} S'_{n+1}$.

Remark 2 A convenient default upper bound is given by the minimum of the Hoeffding, Bernstein (Boucheron et al., 2013, Cor. 2.11), and Berry-Esseen (Shevtsova, 2010) tail bounds for unidentically distributed summands and $\sigma_{\text{low},i}^2 \triangleq \sigma^2 + \frac{R^2}{36n} - \frac{2\sigma_i^2}{3}$.

$$\begin{aligned} & \mathbb{P}(S'_n > t + \frac{R}{4\sqrt{n}}) \\ & \leq \sum_{i=1}^n \frac{\sigma_i^2}{\sigma^2} \min\left(e^{-\frac{t^2}{2R^2 + \frac{9}{8n}R^2}}, e^{-\frac{t^2}{2(\tilde{s}_i^2 + \frac{Rt}{3\sqrt{n}})}}, \Phi^c\left(\frac{t}{\sigma_{\text{low},i}}\right) + \frac{0.56R}{\sqrt{n}} \frac{\sigma^2 + (5R^2 - 21n\sigma_i^2)/(36n)}{\sigma_{\text{low},i}^3}\right). \end{aligned}$$

where $\tilde{s}_i^2 := \sigma^2 + (25R^2 - 96n\sigma_i^2)/(144n)$.

Taken together, Thm. 9, Prop. 10, , and Rem. 2 imply that, for universal constants $c_1, c_2 > 0$ independent of (u, n, R, σ) ,

$$\mathbb{P}(S_n \geq \sigma u + \frac{2R}{\sqrt{n}}) \leq \Phi^c(u) + \frac{c_1 R}{\sigma \sqrt{n}} \exp(-c_2 \frac{u^2}{R^2})$$

whenever Assump. (R, σ) holds.

6. Efficient Concentration with Wasserstein Approximation

In the two previous sections, we showed how to obtain efficient concentration given auxiliary knowledge of a score function or Stein kernel. In this section, we again establish efficient concentration without any knowledge beyond Assump. (R, σ) . To do so, we measure how far S_n is from a normal by the p -Wasserstein distance $\mathcal{W}_p(\cdot, \cdot)$, defined in the following way:

$$\mathcal{W}_p(\nu, \mu) = \inf_{\gamma \in \varphi(\nu, \mu)} \mathbb{E}_{(X, Y) \sim \gamma} (|X - Y|^p)^{1/p},$$

where $\gamma(\nu, \mu)$ is the set of all possible couplings of the probability distributions ν and μ on \mathbb{R} .

6.1. Efficient tail bounds

When Assump. (R, σ) holds we obtain in Thm. 11 below that, for each $u > 0$,

$$\mathbb{P}(S_n \geq \sigma u) \leq \Phi^c\left(u - \frac{5.5R \log(n)}{\sigma \sqrt{n}}\right)[1 + o_n(1)] + \frac{5.5R}{\sqrt{n}\sigma} \varphi\left(u - \frac{5.5R \log(n)}{\sigma \sqrt{n}}\right)[1 + o_n(1)].$$

Notably, this bound is asymptotically exact, converging to the correct asymptotic tail bound $\Phi^c(u)$ as $n \rightarrow \infty$, with an error term that is asymptotically sub-Gaussian in u . To state our result more precisely, we define the growth constant

$$K \triangleq \frac{1}{\sqrt{0.4}} \left[R + \frac{\sqrt{0.4e}\sigma}{2\sqrt{2}} \right] + \frac{2\sqrt{e}R}{\sqrt{1.2}} \left[e^{0.8} - 1.8 \right] \left[2 - \frac{2\sigma}{3R} \right] \approx 3.71R,$$

the interpolation parameter

$$\begin{aligned} \rho_{n,u}^* & \triangleq \frac{\sigma}{eKu} \left(\sqrt{n + \frac{2eK}{\sigma^2} (\sqrt{n}u\sigma - eK(2 + \log(\frac{\sqrt{2\pi n}\sigma^2}{Ke}))^+} - \sqrt{n} \right) \\ & \approx 1 - \frac{eK\sigma}{\sqrt{nu}} \log\left(\frac{\sqrt{n}\sigma\sqrt{2\pi}}{eK}\right) \approx 1 - \frac{5.05R \log(0.25n\sigma^2/R^2)\sigma}{\sqrt{nu}}, \end{aligned}$$

and the correction power

$$L_{n,u} \triangleq \frac{(1-\rho_{n,u}^*)\sqrt{nu}\sigma}{Ke} - 2 \approx 5.05R \log(0.25n\sigma^2/R^2).$$

To first order, Thm. 11 establishes that $\mathbb{P}(S_n \geq \sigma u) \leq \Phi^c(\rho_{n,u}^* u)[1 + o_n(1)] + \frac{eK}{\sqrt{n}\sigma} \varphi(\rho_{n,u}^* u)[1 + o_n(1)]$. To make this bound exact we incorporate a small correction term $r_{n,u}$ of order $\frac{R^2 \log(n)^{5/2}}{\sigma^2 n} \varphi(\rho_{n,u}^* u)$ defined in App. H.1. Under this notation, we have the following efficient tail bound for S_n , proved in App. H.

Theorem 11 (Efficient tail bound) *If S_n satisfies Assump. (R, σ) and $2 \leq L_{n,u} \leq 1 + \frac{0.4\sigma^2 n}{R^2}$, then*

$$\begin{aligned} \mathbb{P}(S_n \geq \sigma u) &\leq \Phi^c(\rho_{n,u}^* u) + \frac{eK}{\sqrt{n}\sigma} \varphi(\rho_{n,u}^* u) [1 + r_{n,u}]^{L_{n,u}} \\ &\stackrel{\star}{\approx} \Phi^c\left(u - \frac{R \log(n)}{\sigma \sqrt{n}}\right) + \frac{R}{\sqrt{n}\sigma} \varphi\left(u - \frac{R \log(n)}{\sigma \sqrt{n}}\right) + O\left(\frac{R^2 \log(n)^{5/2}}{\sigma^2 n} \varphi(\rho_{n,u}^* u)\right), \end{aligned}$$

where $\star := 5.5R$.

Remark 3 *The constant 5.5 can be significantly reduced at the price of a more complicated expression. We also present a tighter but less interpretable bound that holds for all $u \geq 0$ in Sec. 6.3.*

6.2. Efficient quantile bounds

Similarly to how the Wassertein distance can be used to obtain efficient tail bounds it can also be used to obtain efficient quantile bounds. To do so we once again tightly bound the distance between S_n and its Gaussian limit. The quantile bounds we obtain in Thm. 12 are of the order

$$\begin{aligned} q_n(R, \delta, \sigma) &\underset{n \rightarrow \infty}{\sim} -\sigma \Phi^{-1}\left(\delta - \frac{eK}{\sigma \sqrt{n}} \varphi(\Phi^{-1}(\delta))\right) + \frac{eK}{\sqrt{n}} \log\left(\frac{\sqrt{n}\sigma}{eK} \varphi(\Phi^{-1}(\delta))^{-1}\right) \\ &\approx -\sigma \Phi^{-1}\left(\delta - \frac{10.1R}{\sigma \sqrt{n}} \varphi(\Phi^{-1}(\delta))\right) + \frac{10.1R}{\sqrt{n}} \log\left(\frac{\sqrt{n}\sigma}{10.1R} \varphi(\Phi^{-1}(\delta))^{-1}\right). \end{aligned}$$

Notably, as the sample size increases these bounds converge to the true asymptotic quantiles $-\sigma \Phi^{-1}(\delta)$, guaranteeing efficiency. To state our result in full, we introduce the shorthand

$$A_\delta^n \triangleq \log\left(\frac{\sqrt{n}\sigma}{eK} \varphi(\Phi^{-1}(\delta))^{-1}\right),$$

and an explicit correction term $s_{\delta,n}$ of order $\frac{R^2 \log(n)^{5/2}}{n\sigma}$ defined in App. H.3. With this notation in place, we have the following efficient quantile bound for S_n , proved in App. H.

Theorem 12 (Efficient quantile bound) *If S_n satisfies Assump. (R, σ) , $A_\delta^n \geq 2$, and $0 \leq \frac{R^2(A_\delta^n - 1)}{0.4n\sigma^2} \leq 1$ then $P(S_n \geq q_n(R, \delta, \sigma)) \leq \delta$ for*

$$\begin{aligned} q_n(R, \delta, \sigma) &\triangleq \sigma \Phi^{-1}\left(\delta - \frac{eK}{\sigma \sqrt{n}} \varphi(\Phi^{-1}(\delta))\right) + \frac{e[A_\delta^n + 2]}{\sqrt{n}} [K + s_{\delta,n}] \\ &\underset{n \rightarrow \infty}{\approx} -\sigma \Phi^{-1}\left(\delta - \frac{10.1R}{\sigma \sqrt{n}} \varphi(\Phi^{-1}(\delta))\right) + \frac{10.1R}{\sqrt{n}} \log\left(\frac{\sqrt{n}\sigma}{10.1R} \varphi(\Phi^{-1}(\delta))^{-1}\right). \end{aligned}$$

Remark 4 *This bound can be used to build efficient confidence intervals with nominal coverage. Moreover, the constant 10.1 can be significantly reduced at the price of a more complicated expression. In Sec. 6.3, we present a tighter but less interpretable quantile bound (Thm. 14) that holds for all $\delta \in (0, 1)$.*

6.3. Tighter computable bounds

In this section we propose tighter concentration inequalities and quantile approximations for S_n than the ones presented in Sec. 6. While these bounds are less interpretable, they are straightforwardly computed and hence can be used to produce tighter tail bounds and confidence intervals in practice.

First, we obtain a tighter bound for $\mathbb{P}(S_n \geq \sigma u)$, proved in App. I.

Theorem 13 (Tighter efficient tail bound) *If S_n satisfies Assump. (R, σ) , then, for any auxiliary upper bounds $\hat{Q}_n(u)$ and $\hat{Q}_n^d(u)$ on $P(S_n \geq \sigma u)$ and $P(|S_n| \geq \sigma u)$ respectively, we have*

$$P(S_n \geq \sigma u) \leq \min \left(\inf_{p \in [2, \infty) \cap \{1\}, \rho \in (0, 1)} \Phi^c(\rho u) + \frac{\omega_p^R(\sigma)^p}{(1-\rho)^p (u\sigma)^p}, \hat{Q}_n(u) \right); \quad \text{and}$$

$$P(|S_n| \geq \sigma u) \leq \min \left(\inf_{p \in [2, \infty) \cap \{1\}, \rho \in (0, 1)} 2\Phi^c(\rho u) + \frac{\omega_p^R(\sigma)^p}{(1-\rho)^p (u\sigma)^p}, \hat{Q}_n^d(u) \right).$$

for $\omega_p^R(\sigma)$ a computable bound on $\frac{1}{\sigma} \mathcal{W}_p(S_n, \mathcal{N}(0, \sigma^2))$ defined in App. I.1 that is non-decreasing in σ .

Remark 5 (Auxiliary tail bounds) *We include the auxiliary bounds $\hat{Q}_n(u)$ and $\hat{Q}_n^d(u)$ to emphasize that our efficient bounds can be paired with any valid tail bounds to simultaneously reap the large-sample benefits of the former and the small-sample benefits of the latter. The Hoeffding-Bernstein bounds $\hat{Q}_n(u) = \min(e^{-\frac{u^2 \sigma^2}{2R^2}}, e^{-\frac{u^2}{2(1+Ru/3\sigma)}})$ and $\hat{Q}_n^d(u) = 2\hat{Q}_n(u)$ provide a convenient default choice.*

By choosing $p = L_{n,u}$ and $\rho = \rho_{n,u}^*$ in Thm. 13 we obtain a tightened version of the inequality in Thm. 11. Further improvements are obtained in practice by explicitly optimizing over p and ρ .

Next we obtain tighter bounds for the quantiles of S_n , proved in App. I.

Theorem 14 (Tighter efficient quantile bound) *If S_n satisfies Assump. (R, σ) , and some auxiliary deterministic bounds \tilde{q}_n and \tilde{q}_n^d satisfy $P(S_n \geq \tilde{q}_n(R, \delta, \sigma)) \leq \delta$ and $P(|S_n| \geq \tilde{q}_n^d(R, \delta, \sigma)) \leq \delta$, then*

$$\mathbb{P}(S_n \geq q_n(R, \delta, \sigma)) \leq \delta, \quad \mathbb{P}(|S_n| \geq q_n^d(R, \delta, \sigma)) \leq \delta,$$

for $\omega_p^R(\sigma)$ a computable bound on $\frac{1}{\sigma} \mathcal{W}_p(S_n, \mathcal{N}(0, \sigma^2))$ defined in App. I.1 that is non-decreasing in σ and

$$q_n(R, \delta, \sigma) \triangleq \min \left(\inf_{p \geq 2, \rho \in (0, 1)} \frac{\sigma \omega_p^R(\sigma)}{(\delta(1-\rho))^{1/p}} - \sigma \Phi^{-1}(\delta \rho), \tilde{q}_n(R, \delta, \sigma) \right),$$

$$q_n^d(R, \delta, \sigma) \triangleq \min \left(\inf_{p \geq 2, \rho \in (0, 1)} \frac{\sigma \omega_p^R(\sigma)}{(\delta(1-\rho))^{1/p}} - \sigma \Phi^{-1}\left(\frac{\delta \rho}{2}\right), \tilde{q}_n^d(R, \delta, \sigma) \right).$$

Remark 6 (Auxiliary quantile bounds) *We again highlight that our efficient bounds can be paired with any valid quantile bounds $\tilde{q}_n(R, \delta, \sigma)$ and $\tilde{q}_n^d(R, \delta, \sigma)$ to inherit the benefits of each. Convenient defaults are provided by the minimum of the Bernstein quantiles (Boucheron et al., 2013, Thm. 2.10) and the Berry-Esseen quantiles (Shevtsova, 2011),*

$$\tilde{q}_n(R, \delta, \sigma) = \min \left(\frac{R}{3\sqrt{n}} \log(1/\delta) + \sigma \sqrt{2 \log(1/\delta)}, -\sigma \Phi^{-1} \left(\delta - \frac{C_{R,\sigma}}{\sqrt{n}} \right) \right),$$

$$\tilde{q}_n^d(R, \delta, \sigma) = \min \left(\frac{R}{3\sqrt{n}} \log(2/\delta) + \sigma \sqrt{2 \log(2/\delta)}, -\sigma \Phi^{-1} \left(\frac{\delta}{2} - \frac{C_{R,\sigma}}{\sqrt{n}} \right) \right).$$

where we set $C_{R,\sigma} = \min(.3328(\tilde{R} + .429), .33554(\tilde{R} + .415))$.

6.4. Proof sketch for Thms. 11 and 13

Similarly to the proof of Thm. 6 and Thm. 3, the main idea of the proofs is to use the fact that $S_n = \frac{1}{\sqrt{n}} \sum_{i \leq n} Y_i$ is close to being normally distributed if the sample size $n \in \mathbb{N}$ is large. For ease of notation, the rest of this section we will suppose that $\sigma = 1$. Note that there is no loss of generality by doing so. Fix $t \geq 0$, $\rho \in [0, 1)$, and $p \geq 1$. For all $\epsilon > 0$ we remark that there is a coupling (S_n, G) with $G \sim N(0, 1)$ such that $\|S_n - G\|_p \leq \mathcal{W}_p(S_n, G) + \epsilon$. Therefore we have

$$\begin{aligned} \mathbb{P}(S_n \geq t) &= \mathbb{P}(G + S_n - G \geq t) \\ &\leq \mathbb{P}(G \geq \rho t) + \mathbb{P}(S_n - G \geq (1 - \rho)t) \\ &\leq \Phi^c(\rho t) + \frac{\mathbb{E}(|S_n - G|^p)}{(1 - \rho)^p t^p} \\ &\leq \Phi^c(\rho t) + \frac{(\mathcal{W}_p(S_n, G) + \epsilon)^p}{(1 - \rho)^p t^p}. \end{aligned}$$

As this holds for any arbitrary $\epsilon > 0$ we get that

$$\begin{aligned} \mathbb{P}(S_n \geq t) &= \mathbb{P}(G + S_n - G \geq t) \\ &\leq \Phi^c(\rho t) + \frac{\mathcal{W}_p(S_n, G)^p}{(1 - \rho)^p t^p}. \end{aligned}$$

Imagine that $\mathcal{W}_p(S_n, G) \leq \frac{\tilde{K} p}{\sqrt{n}}$ for a constant \tilde{K} and that we set $p = \frac{\sqrt{n} t (1 - \rho)}{e \tilde{K}}$ then we obtain

$$\mathbb{P}(S_n \geq t) \leq \Phi^c(\rho t) + e^{-\frac{\sqrt{n} t (1 - \rho)}{e \tilde{K}}}.$$

If in addition we choose ,

$$\rho \triangleq \frac{-C_{n,t} + \sqrt{C_{n,t}^2 + 2t^2(C_{n,t} + 1 - \log(C_{n,t} \sqrt{2\pi}/t))}}{t^2}$$

with $C_{n,t} \triangleq \frac{\sqrt{n} t}{e \tilde{K}}$ then we have

$$\mathbb{P}(S_n \geq t) \lesssim \Phi^c(\rho t) + \frac{e \tilde{K}}{\sqrt{n}} \varphi(\rho t);$$

This gives us the desired bound.

6.5. Proof sketch for Thms. 12 and 14

Quantiles are obtained by following a similar reasoning. Let $\delta \in (0, \frac{1}{2})$ be a confidence level and $\alpha \in (0, \delta)$. Choose $\epsilon > 0$ and (S_n, G) to be a coupling of S_n with a standard normal random variable that satisfies $\|S_n - G\|_p \leq \mathcal{W}_p(S_n, \mathcal{N}(0, 1)) + \epsilon$. Then for all $p \geq 2$, if we write $u \triangleq \frac{\mathcal{W}_p(S_n, G)}{\alpha^{1/p}}$ then we obtain

$$\begin{aligned} \mathbb{P}(S_n \geq -\Phi^{-1}(\delta - \alpha) + u) &\leq \mathbb{P}(G \geq -\Phi^{-1}(\delta - \alpha)) + \mathbb{P}(S_n - G \geq u) \\ &\leq \delta - \alpha + \frac{(\|S_n - G\|_p + \epsilon)^p}{u^p}. \end{aligned}$$

As the choice of $\epsilon > 0$ can be arbitrarily small we obtain

$$\mathbb{P}(S_n \geq -\Phi^{-1}(\delta - \alpha) + u) \leq \delta - \alpha + \frac{\mathcal{W}_p(S_n, G)^p}{u^p} \leq \delta.$$

Therefore to obtain a good approximation of the quantiles of S_n we need to choose a candidate for α and p . If we choose $p = -\log(\alpha)$ then we obtain then we can upper bound the quantile of S_n by

$$-\Phi^{-1}(\delta - \alpha) + e\mathcal{W}_{-\log(\alpha)}(S_n, G) \triangleq O(-\Phi^{-1}(\delta - \alpha) + \frac{-\log(\alpha)}{\sqrt{n}}).$$

We choose $\alpha = O(\frac{\varphi(\Phi^{-1}(\delta))}{\sqrt{n}})$.

7. Efficient Concentration Inequalities for Smoothed Observations

We notice that when we had some additional information on the distribution of the random variables (Y_i) , e.g about their score function (see Thm. 4) and their Stein kernel (see Thm. 8), then we were able to avoid potentially lossy union bound and obtained tighter concentration inequalities. In this section we show that if the observations (Y_i) are smoothed then we can obtain (potentially) tighter concentration inequalities without requiring any additional properties of the distribution of the random variables (Y_i) . In this goal let $D > 0$ be a real and let Z be an independent standard normal random variable. We define

$$\tilde{S}_n^{\text{noisy}} \triangleq \sqrt{1 - \frac{D}{\sqrt{n}}} \tilde{S}_n + \frac{\sqrt{D}}{n^{1/4}} Z, \quad S_n^{\text{noisy}} \triangleq \sigma \tilde{S}_n^{\text{noisy}}.$$

We show how we can obtain efficient tail bounds for $\tilde{S}_n^{\text{noisy}}$. In this goal, we write $\omega_{1,p,q} = \mathcal{W}_{\frac{2pq}{q-1}}(\tilde{S}_n, Z)$ and $\omega_{2,p,q} = \|Z\|_{\frac{pq}{q-1}}$.

Theorem 15 (Efficient smoothed tail bound) *Assume that the random variables (Y_i) satisfy Assump. (R, σ) . Let $\hat{Q}_n(u)$ be an arbitrary upper bound for $\mathbb{P}(\tilde{S}_n \geq u) \leq \hat{Q}_n(u)$. Then the following holds for all $\kappa \geq \max(\frac{\tilde{R}^2}{n}, \frac{\tilde{R}^2}{\sqrt{n}D})$*

$$\begin{aligned} & \mathbb{P}(S_n^{\text{noisy}} \geq \sigma u) \\ & \lesssim \Phi^c(u) + \inf_{p \geq 2} \frac{\hat{Q}_n(u)^{\frac{p-1}{p}}}{\sqrt{n}} \left\{ \frac{\sqrt{e}(p-1)\tilde{R}^2(e^{2\kappa(p-1)}-1)}{\sqrt{3}D\kappa} + \frac{\log\left(\frac{n}{D^2}\right)\sqrt{p-1}A_p(\tilde{R}-1)}{4\sqrt{2}} \right\}. \end{aligned}$$

Remark 7 *The exact bound is presented in Thm. 36. We also note that this bound could not be obtained by directly applying Thm. 4 or Thm. 8. Indeed while the score function $\tilde{S}_n^{\text{noisy}}$ can be seen as an empirical average of continuous random variables the amount of Gaussian noise injected is not sufficient to control its score function or its Stein kernel.*

Note that one can straightforwardly bound the Wassertein distance between S_n^{noisy} and S_n and use this to obtain an efficient concentration inequality for the latter. Similarly if S_n admits a Stein kernel then one could use strong embeddings (Chatterjee, 2012, Thm. 1.2) to control the tail bound of $|S_n - S_n^{\text{noisy}}|$ and use Thm. 15 to obtain an efficient concentration inequality for S_n .

7.1. Proof sketch for Thm. 15

The proof relies on the Ornstein-Ulbeck interpolation and the heat equation. To make this precise we write the interpolated process $X_t \triangleq e^{-t}\tilde{S}_n + \sqrt{1 - e^{-2t}}Z$. As t varies from 0 to ∞ , then X_t

transforms from \tilde{S}_n to Z . We define $T_n \triangleq -\frac{1}{2} \log(1 - \frac{D}{\sqrt{n}})$. We note that X_{T_n} has the same distribution than $\tilde{S}_n^{\text{noisy}}$. The key to the proof is then to use the heat equation to establish the following equality:

$$\begin{aligned} & |\mathbb{P}(\tilde{S}_n^{\text{noisy}} \geq u) - \Phi^c(u)| \\ &= |\int_{T_n}^{\infty} \mathbb{E} \left[\mathbb{I}(X_t \geq u) \mathbb{E} \left[\frac{e^{-2t}}{1-e^{-2t}} (Z^2 - 1) - \frac{e^{-t}}{\sqrt{1-e^{-2t}}} \tilde{S}_n Z - \tau'_t | X_t \right] \right] dt|. \end{aligned}$$

where (τ'_t) is the quantity defined in Eq. (14). The proof is finished by using Hölder's inequality and bounding the norms $\| \frac{e^{-2t}}{1-e^{-2t}} (Z^2 - 1) - \frac{e^{-t}}{\sqrt{1-e^{-2t}}} \tilde{S}_n Z - \tau'_t \|_p$ (see Lem. 19 and Lem. 21).

8. Application: Efficient *Empirical Berry-Esseen* Bounds

The preceding sections assumed that the variance parameter σ^2 was a known quantity, but, in many applications, the variance is unknown and can only be estimated from data. In this section, we leverage efficient known-variance bounds to develop efficient quantile bounds that are valid even when σ is unknown. We refer to these constructions as *empirical Berry-Esseen* (EBE) bounds as they combine Gaussian approximation in the spirit of Berry-Esseen Eq. (5) with empirical variance estimation.

We begin by showing how to convert generic known-variance quantile bounds into valid empirical-variance quantile bounds. Our construction in Thm. 16 depends on the following readily-computed quantities:

- the empirical variance: $\hat{\sigma}^2 \triangleq \frac{1}{n} \sum_{i \leq n} (W_i - \bar{W}_n)^2$,
- the empirical centered third moment: $\hat{m}_3 \triangleq \frac{1}{n} \sum_{i \leq n} (W_i - \bar{W}_n)^3$,
- the empirical variance of $(W_i - \bar{W}_n)^2$: $\hat{m}_4 \triangleq \frac{1}{n} \sum_{i \leq n} ((W_i - \bar{W}_n)^2 - \hat{\sigma}^2)^2$,
- the $1 - \frac{a}{4}$ quantile of the two-sided Kolmogorov-Smirnov distribution: $q_n^{\text{KS}}(a/4)$,
- a lower estimate for σ^2 : $\sigma_{l,a}^2 \triangleq \hat{\sigma}^2 - R^2 q_n^{\text{KS}}(\frac{a}{4})$, and
- a lower estimate for $s_4^2 = \text{Var}((W_1 - \mathbb{E}(W_1))^2)$:

$$s_{l,a}^2 \triangleq \hat{m}_4 - R^4 q_n^{\text{KS}}(\frac{a}{4}) + \min_{|u| \leq 2Rq_n^{\text{KS}}(\frac{a}{4})} 4\hat{\sigma}^2 u^2 - 4\hat{m}_3 u,$$

and the proof can be found in App. K.

Theorem 16 (Empirical quantile bounds) *Consider any quantile bounds \tilde{q}_n^d and \tilde{q}_n with*

$$\begin{aligned} P(|S_n| \geq \tilde{q}_n^d(R, \delta, \sigma)) &\leq \delta \quad \text{and} \quad c_1^\delta(\sigma) \triangleq \frac{\tilde{q}_n^d(R, \delta, \sigma)^2}{n\sigma^2} \quad \text{non-increasing} \quad \text{and} \\ P(S_n > \tilde{q}_n(R, \delta, \sigma)) &\leq \delta \quad \text{and} \quad c_2^\delta(\sigma) \triangleq \frac{\tilde{q}_n(R, \delta, \sigma)^2}{n\sigma^2} \quad \text{non-increasing} \end{aligned}$$

for each $\delta \in (0, 1)$ and S_n satisfying Assump. (R, σ). If S_n satisfies Assump. (R, σ), then, for each confidence level $\delta \in (0, 1)$ and $a_n \in (0, \delta)$, we have

$$P(S_n \notin \sqrt{n} A_{\delta, a_n}^n) \leq \delta \quad \text{and hence} \quad P(|S_n| \geq \tilde{q}_n^d(R, \delta, a_n)) \leq \delta$$

for the empirical quantile bound $\hat{q}_n^d(R, \delta, a_n) \triangleq \sqrt{n} \max_{x \in A_{\delta, a_n}^n} |x|$ and the confidence set

$$A_{\delta, a}^n \triangleq \left\{ x \in \mathbb{R} \mid \left(x^2 - \hat{\sigma}^2 \frac{c_1^{\delta-a}(\sigma_{1,a})}{1 - c_1^{\delta-a}(\sigma_{1,a})} \right)^2 \leq \left(\frac{c_1^{\delta-a}(\sigma_{1,a})}{1 - c_1^{\delta-a}(\sigma_{1,a})} \right)^2 \frac{c_2^a(s_{1,a})}{1 - c_2^a(s_{1,a})} \left[\hat{m}_4 + 4\hat{\sigma}^2 x^2 - 4\hat{m}_3 x + R^4 q_n^{\text{KS}}\left(\frac{a}{4}\right) \right] \right\}.$$

Remark 8 The maximum and minimum elements of A_{δ, a_n}^n are roots of a depressed quartic equation that can be solved in closed form or with standard software.

Next we prove that efficient quantile bounds give rise to efficient empirical quantile bounds.

Theorem 17 (Efficient empirical Berry-Esseen bounds) Under the assumptions of Thm. 16, suppose $\tilde{q}_n^d(R, \sigma_n, \delta_n) \rightarrow -\sigma\Phi^{-1}(\frac{\delta}{2})$ whenever $(\sigma_n, \delta_n) \rightarrow (\sigma, \delta)$. If $a_n \rightarrow 0$, $\frac{\log(a_n)}{n} \rightarrow 0$, and $c_2^{a_n}(s_{1,a_n}) \xrightarrow{a.s.} 0$, then

$$\hat{q}_n^d(R, \delta, a_n) \xrightarrow{a.s.} -\sigma\Phi^{-1}\left(\frac{\delta}{2}\right).$$

Remark 9 As convenient default choices, we recommend setting $a_n = \frac{\log n}{n}$ and taking \tilde{q}_n^d and \tilde{q}_n to be the efficient quantile bounds q_n^d, q_n of Thm. 14 with the auxiliary bounds of Rem. 6.

Proof For brevity, we will adopt the shorthand $\hat{q}_n^d \triangleq \hat{q}_n^d(R, \delta, a_n)$ and $\tilde{q}_n^d \triangleq \tilde{q}_n^d(R, \delta - a_n, \sigma_{1,a_n})$. We will also write $T_{n,\delta}^d := \frac{\tilde{q}_n^d / \sigma_{1,a_n}^2}{1 - c_1^{\delta-a_n}(\sigma_{1,a_n})}$. Since $\sqrt{n}A_{\delta, a_n}^n$ is the sublevel set of the norm-coercive monic polynomial

$$y \mapsto \left(y^2 - \hat{\sigma}^2 T_{n,\delta}^d \right)^2 - (T_{n,\delta}^d)^2 \frac{c_2^{a_n}(s_{1,a_n}) \left(\hat{m}_4 + \frac{4\hat{\sigma}^2 y^2}{n} - \frac{4\hat{m}_3}{\sqrt{n}} + R^4 q_n^{\text{KS}}\left(\frac{a_n}{4}\right) \right)}{1 - c_2^{a_n}(s_{1,a_n})}, \quad (13)$$

\hat{q}_n^d is necessarily a root of this polynomial.

Moreover, the coefficients of this polynomial converge as follows. First, the Dvoretzky-Kiefer-Wolfowitz inequality (Dvoretzky et al., 1956) implies that

$$q_n^{\text{KS}}\left(\frac{a_n}{4}\right) \leq \sqrt{\frac{\log(C/a_n)}{n}}$$

for some constant C . Our assumption $\frac{\log(a_n)}{n} \rightarrow 0$ therefore yields $q_n^{\text{KS}}\left(\frac{a_n}{4}\right) \rightarrow 0$. Moreover, by the strong law of large numbers (Durrett, 2019, Thm. 2.4.1), $\hat{\sigma}^2 \xrightarrow{a.s.} \sigma^2$, $\hat{m}_4 \xrightarrow{a.s.} s_4^2$, and $\hat{m}_3 \xrightarrow{a.s.} \mathbb{E}[(W_i - \mathbb{E}W_1)^3]$ and hence $\sigma_{1,a}^2 \xrightarrow{a.s.} \sigma^2$ and $s_{1,a}^2 \xrightarrow{a.s.} s_4^2$ as well. Since $a_n \rightarrow 0$, we therefore have $\tilde{q}_n^d \xrightarrow{a.s.} -\sigma\Phi^{-1}(\frac{\delta}{2})$ and $c_1^{\delta-a_n}(\sigma_{1,a_n}) \xrightarrow{a.s.} 0$. Finally, as $c_2^{a_n}(s_{1,a_n}) \rightarrow 0$, the polynomial Eq. (13) converges almost surely to the monic polynomial $y \mapsto (y^2 - (-\sigma\Phi^{-1}(\frac{\delta}{2})))^2$. Hence, by Artin (2011, Prop. 5.2.1), \hat{q}_n^d converges almost surely to the unique positive root of this polynomial, $-\sigma\Phi^{-1}(\frac{\delta}{2})$. \blacksquare

8.1. Numerical evaluation

We next turn to a numerical evaluation of our efficient EBE bound. Python code implementing our bound and reproducing all plots can be found at https://github.com/lmackey/gauss_conc/. Fig. 1 compares the efficient EBE bound of Thm. 17 and Rem. 9 with the Hoeffding quantile bound, $\mathbb{P}(|S_n| \geq R\sqrt{2\log(2/\delta)}) \leq \delta$; the empirical Bernstein quantile bound (Maurer and Pontil, 2009, Thm. 4) commonly deployed in reinforcement learning (Mnih et al., 2008; Audibert et al., 2009), $\mathbb{P}(|S_n| \geq \hat{\sigma}\sqrt{2\log(2/\delta)\frac{n}{n-1}} + \frac{14}{3}R\log(2/\delta)\frac{\sqrt{n}}{n-1}) \leq \delta$; and the efficient $I_{n,3}$ quantile bound of Romano and Wolf (2000),² with its free parameter β_n set to match our default value, $a_n = \frac{\log n}{n}$.

As advertised, the efficient bounds converge to the optimal asymptotic size as n increases, while the Bernstein and Hoeffding inequalities remain bounded away from the optimum for all n . Moreover, the efficient EBE bound of Thm. 16 provides a tighter estimate than the optimized Romano-Wolf bound in all settings. Here the efficient empirical bound is constructed with $\hat{m}_3 = R\hat{\sigma}^2$ and $\hat{m}_4 = R^2\hat{\sigma}^2 - \hat{\sigma}^4$ and, as we show in App. M, changes only minimally as \hat{m}_3 and \hat{m}_4 vary.

9. Discussion and Related Work

In this work, we have derived new computable tail and quantile bounds for the scaled deviations $S_n = \sqrt{n}(\bar{W}_n - \mathbb{E}(W_1))$ with asymptotically optimal size, finite-sample validity, and sub-Gaussian decay. These bounds enable the construction of efficient confidence intervals with correct coverage for any sample size. Our concentration inequalities arise from new computable bounds on the Hellinger distance, Stein discrepancy, non-uniform Kolmogorov distance, and Wasserstein distance to a Gaussian, and, as a byproduct, we obtain the first explicit bounds for the Hellinger CLT. We also develop a general method for bounding the Hellinger distance between a Gaussian and any other random variable.

The notion of efficient confidence intervals for the mean was introduced by Romano and Wolf (2000). In their Thm. 2.1, Romano and Wolf showed that efficient—that is, asymptotically minimal-length when scaled by \sqrt{n} —confidence intervals must converge to the width of the asymptotic Gaussian intervals implied by the CLT Eq. (3). Moreover, Sec. 3 of Romano and Wolf surveys a number of procedures for constructing confidence intervals that are either finite-sample invalid (including the bootstrap (Efron, 1992) and methods based on Edgeworth expansions (Hall, 2013)) or inefficient (including the methods of Anderson (1969) and Gasko (1991)). Romano and Wolf conclude by developing an efficient valid confidence interval for the mean of variables supported on $[0, 1]$ but report that it is “unfortunately, much too wide for a reasonable sample.” Our new efficient bounds are developed in an entirely different manner, and, as we demonstrate in Sec. 8.1, improve upon the Romano-Wolf interval and the most commonly used empirical concentration inequalities.

Our results also suggest simple strategies for making any concentration inequality or confidence region efficient. For efficient concentration, one can simply take the minimum of any existing tail bound and Thm. 9 or Thm. 13 to simultaneously reap the small-sample benefits of the former and the large-sample benefits of the latter. For efficient confidence, one can divide the total confidence budget between an existing region and the efficient region of Thm. 17 and then intersect the two regions. The result will remain efficient if the budget allocated to the auxiliary region vanishes as n grows. These strategies are particularly relevant given the recent renewed interest in deriving

2. The Romano-Wolf bound can be infinite, but we constrain it to be no larger than the always-valid Hoeffding bound.




Figure 1: Empirical quantile bounds for $|S_n|$ with fixed boundedness parameter $R = 1$ and varying sample size n and empirical variance $\hat{\sigma}^2$. Unlike the empirical Bernstein and Hoeffding bounds, the efficient empirical Berry-Esseen bound of Thm. 16 converges to the ideal asymptotic Gaussian bound as n increases and provides a tighter estimate than the efficient $I_{n,3}$ interval of Romano and Wolf (2000) in all settings.

tighter concentration inequalities for bounded random variables (see, e.g., Jun and Orabona, 2019; Waudby-Smith and Ramdas, 2020; Orabona and Jun, 2021).

Our Hellinger bound is, to the best of our knowledge, the first known explicit rate of convergence for the central limit theorem in terms of the Hellinger distance. Indeed while rates were known for the KL divergence, none were explicitly derived for the Hellinger distance. Moreover, the KL rates are neither explicit nor computable and require stronger conditions. However, our proof technique

builds on previously developed ideas. Notably, we use the Ornstein-Uhlenbeck semigroup to interpolate between \tilde{S}_n and a normal. This allows us to re-express the Hellinger distance as an integral over the norm of information-theoretic quantities of the interpolated process. This can be seen as the equivalent of the De Bruijn identity (see e.g. (Barron, 1986)) for the KL-divergence or similar expressions obtained for p -Wassertein distance in the celebrated work of Otto and Villani (2000) and in Ledoux et al. (2015). Then we take inspiration from Bonis (2015) which proposed to use an exchangeable pair argument to give a closed form upper bound to the p -Wassertein distance and develop similar techniques. This provides us with an easily computable bound that only depends on the moments of the score function and the moments of the random variable itself. This is to contrast with the known bounds for the KL divergence, which are either not in closed form (Bobkov et al., 2013), require conditions on the spectral gap (Artstein et al., 2004) or depend on the moment of the Stein kernel (Ledoux et al., 2015).

The notion of a Stein kernel (sometimes known under the name of *Stein factor*, ω -*function*, or *covariance kernel* (Ley et al., 2017, Rem. 4.6)) is born from the literature of Stein’s approximation and is found implicitly in many of the earlier works (see for e.g. the original monograph (Stein, 1986)). It was however first formally introduced in Cacoullos and Papathanasiou (1992) and has been the subject of a lot of recent development including to establish multivariate central limit theorems (Nourdin et al., 2010), to compare univariate distributions (Ley et al., 2017) for its connection with the total variation distance (Cacoullos et al., 1994), the p -Wassertein distance (Ledoux et al., 2015; Bonis, 2015), the entropic central limit theorem (Nourdin et al., 2014; Ledoux et al., 2015), covariance bounds (Ernst et al., 2020) and concentration inequalities (Ledoux et al., 2015; Nourdin and Viens, 2009; Saumard, 2019). Our theorem Thm. 6 takes inspiration from the concentration inequalities obtained in Ledoux et al. (2015), but the inequality that we obtain is efficient and is on the tail of the S_n rather than on the moments of Lipschitz functions of S_n . The Stein kernel is however sadly not always guaranteed to exist. This led to a large body of work investigating under what conditions such a quantity exists and how to compute it (see e.g. (Courtade et al., 2017; Fathi, 2019)).

Our zero-bias coupling arguments generalize the uniform, u -independent bounds of Chen et al. (2011); Ross (2011) to derive tighter non-uniform bounds with sub-Gaussian decay. Our Wasserstein bounding arguments in Sec. 6 build on the pioneering work of Bonis (2015) who derived Wasserstein convergence rates for the CLT with inexplicit constants. Our new arguments lead to tighter estimates of the distance to Gaussianity and explicit, practical constants.

An important open question is if a similar approach could be used for other statistics, beyond sample means, that are known to be asymptotically normal. Another natural direction for future work is to relax the assumption that the random variables are bounded to conditions on the moments of $(W_i)_{i \geq 1}$.

Appendix

A	Appendix Overview and Notation	25
B	Proof of Thm. 3: Efficient Hellinger tail bound	26
C	Proof of Thm. 4: Hellinger CLT bound	26
C.1	Bounding the different terms	28

C.2	Combining the results together	43
D	Proof of Thm. 5: Bounding Hellinger distance to a Gaussian	44
E	Proof of Thms. 6, 7, and 8: Efficient Concentration with Stein Kernels	47
E.1	Proof of Thm. 6: Efficient Stein kernel tail bound	48
E.2	Proof of Thm. 7: Alternative Stein kernel tail bound	53
E.3	Proof of Thm. 8: Efficient Hellinger-Stein kernel tail bound	54
F	Proof of Thm. 9: Efficient zero-bias tail bound	54
F.1	Proof of Thm. 29: Zero-bias tail bounds	55
F.2	Proof of Lem. 30: Growth of h_u	56
F.3	Proof of Lem. 31: Growth of Φ^c	58
G	Proof of Prop. 10: Properties of S'_n	58
H	Proof of Thms. 11 and 12: Efficient Concentration with Wasserstein Approximation	58
H.1	Definition of $r_{n,t}$ in Thm. 11	59
H.2	Proof of Thm. 11: Efficient tail bound	60
H.3	Definition of $s_{n,\delta}$ in Thm. 12	62
H.4	Proof of Thm. 12: Efficient quantile bound	62
I	Proof of Thms. 13 and 14: Tighter computable bounds	64
I.1	Definitions of the Quantities in Sec. 6.3	64
I.2	Proof of Thms. 13 and 14	64
J	Proof of Thm. 15: Efficient smoothed tail bound	80
K	Proof of Thm. 16: Empirical quantile bounds	83
L	Additional Lemmas	86
M	Additional Plots	88

Appendix A. Appendix Overview and Notation

The proofs of Thms. 11 and 12 to 14 are each broken into two parts. In App. I.1 we derive upper bounds for the Wasserstein distances between S_n and its asymptotic distribution $\mathcal{N}(0, \sigma^2)$. In Apps. H.2, H.4, and I we exploit those upper bounds to establish the individual theorem claims.

Throughout the appendix, we will designate by $(X_i)_{i \geq 1}$ the rescaled random variables $X_i = \frac{W_i - \mathbb{E}(W_i)}{\sqrt{n}} = \frac{Y_i}{\sqrt{n}}$. We remark that $\mathbb{E}(X_1) = 0$, that $\text{Var}(X_1) = \frac{\text{Var}(Y_1)}{n} = \frac{\sigma^2}{n}$ and that $\|X_1\|_\infty \leq R/\sqrt{n}$. We also note that S_n can be re-expressed as $S_n = \sum_{i \leq n} X_i$. Moreover we use the following notation:

$$Y_i \triangleq W_i - \mathbb{E}(W_1), \quad S_n \triangleq \sqrt{n}(\bar{W}_n - \mathbb{E}(W_1)), \quad \tilde{S}_n \triangleq \sigma^{-1} S_n, \quad \tilde{R} \triangleq \sigma^{-1} R,$$

$$A_p \triangleq \frac{\sqrt{e}\sqrt{p+2}(2e)^{1/p}}{\sqrt{2}}, \quad A_{n,p}^* \triangleq \frac{(p+2)n^{1/p}}{2\sqrt{n}}, \quad \tilde{A}_{n,p} \triangleq A_{n,p}^* \tilde{R}^{-2/p}.$$

$$U_{n,p} \triangleq A_p + \tilde{R} \tilde{A}_{n,p}, \quad U_{n,p}^* \triangleq A_p + \tilde{R}^{2/p} \tilde{A}_{n,p}.$$

Finally, throughout the appendix we write for all $t > 0$

$$X_t \triangleq e^{-t} \tilde{S}_n + \sqrt{1 - e^{-2t}} Z, \quad \tilde{Y}_i = Y_i / \sigma.$$

Appendix B. Proof of Thm. 3: Efficient Hellinger tail bound

Write $f_n(\cdot)$ the PDF of $\sigma^{-1} S_n$ and let $Z \sim N(0, 1)$ be an independent standard normal random variable. Write $\varphi(\cdot)$ the PDF of Z and denote $h(\cdot) \triangleq \frac{f_n(\cdot)}{\varphi(\cdot)}$. Then we have

$$\begin{aligned} & |P(\sigma^{-1} S_n \geq u) - \mathbb{P}(Z \geq u)| \\ &= |\mathbb{E}((h(Z) - 1)\mathbb{I}(Z \geq u))| \\ &= |\mathbb{E}((\sqrt{h(Z)} - 1)(\sqrt{h(Z)} + 1)\mathbb{I}(Z \geq u))| \\ &\leq |\mathbb{E}((\sqrt{h(Z)} - 1)\mathbb{I}(Z \geq u))| \\ &\quad + |\mathbb{E}((\sqrt{h(Z)} - 1)\sqrt{h(Z)}\mathbb{I}(Z \geq u))| \\ &\stackrel{(a)}{\leq} \sqrt{\mathbb{E}((\sqrt{h(Z)} - 1)^2)} \sqrt{\mathbb{E}(\mathbb{I}(Z \geq u))} \\ &\quad + \sqrt{\mathbb{E}((\sqrt{h(Z)} - 1)^2)} \sqrt{\mathbb{E}(h(Z)\mathbb{I}(Z \geq u))} \\ &\leq \sqrt{\mathbb{E}((\sqrt{h(Z)} - 1)^2)} \left[\sqrt{\Phi^c(u)} + \sqrt{\mathbb{E}(h(Z)\mathbb{I}(Z \geq u))} \right] \\ &\leq \sqrt{\int_{-\infty}^{\infty} (\sqrt{f_n(x)} - \sqrt{\varphi(x)})^2 dx} \left[\sqrt{\Phi^c(u)} + \sqrt{\mathbb{E}(h(Z)\mathbb{I}(Z \geq u))} \right] \\ &\leq \sqrt{2} H(\sigma^{-1} S_n, Z) \left[\sqrt{\Phi^c(u)} + \sqrt{\mathbb{E}(h(Z)\mathbb{I}(Z \geq u))} \right] \\ &\leq \sqrt{2} H(\sigma^{-1} S_n, Z) \left[\sqrt{\Phi^c(u)} + \sqrt{\hat{Q}_n(u)} \right], \end{aligned}$$

where we obtain (a) as a consequence of Cauchy-Swartz inequality. Similarly we remark that:

$$\begin{aligned} |P(|\sigma^{-1} S_n| \geq u) - \mathbb{P}(|Z| \geq u)| &= |\mathbb{E}((h(Z) - 1)\mathbb{I}(|Z| \geq u))| \\ &\leq \sqrt{2} H(\sigma^{-1} S_n, Z) \left[\sqrt{2\Phi^c(u)} + \sqrt{\hat{Q}_n^d(u)} \right]. \end{aligned}$$

Appendix C. Proof of Thm. 4: Hellinger CLT bound

Throughout this section we fix a real $D > 0$ and take $\kappa \geq \max\left(\frac{\tilde{R}^2}{n}, \frac{\tilde{R}^2}{\sqrt{n}D}\right)$. Finally we set

$$S'_{n,t} \triangleq \tilde{S}_n + (X'_I - X_I)\mathbb{I}(|X'_I|, |X_I| \leq \sqrt{\kappa(1 - e^{-2t})}),$$

where $I \sim \text{unif}\{1, \dots, n\}$ and shorthand

$$Y_t = \tilde{S}_n - S'_{n,t}.$$

We first specialize Thm. 5 to \tilde{S}_n . In this goal we write $S_{\tilde{S}_n}(\cdot)$ the score function of \tilde{S}_n .

Lemma 18 Suppose that the random variables (Y_i) satisfy the condition of Thm. 4. Define $\tilde{S}_n \triangleq \sigma^{-1} S_n$ and for all $t > 0$ choose \tilde{S}_n^t to be such that $(\tilde{S}_n^t, \tilde{S}_n)$ forms an exchangeable pair. Let Z be an independent standard normal random variable. We define

$$\begin{aligned}\tau_t &\triangleq \sum_{k=1}^{\infty} \frac{e^{-kt}}{k! \sqrt{1-e^{-2t}}^{k-1}} \mathbb{E}((\tilde{S}_n^t - \tilde{S}_n)^k | S_n) H_{k-1}(Z); \\ \tau'_t &\triangleq \sum_{k=1}^{\infty} \frac{e^{-kt}}{k! \sqrt{1-e^{-2t}}^k} \mathbb{E}((\tilde{S}_n^t - \tilde{S}_n)^k | \tilde{S}_n) H_k(Z).\end{aligned}\tag{14}$$

Then the following holds for all $p \geq 1$ and all $D > 0$

$$\begin{aligned}\sqrt{2}H(\tilde{S}_n, Z) &\leq \int_{-\frac{1}{2} \log(1-\frac{D}{\sqrt{n}})}^{\infty} \left\| \frac{e^{-2t}}{1-e^{-2t}} (Z^2 - 1) - \frac{e^{-t}}{\sqrt{1-e^{-2t}}} \tilde{S}_n Z + \tau'_t \right\|_2 dt \\ &\quad + \int_0^{-\frac{1}{2} \log(1-\frac{D}{\sqrt{n}})} \inf_{p \geq 1} C_p \left\| \frac{Ze^{-2t}}{\sqrt{1-e^{-2t}}} - e^{-t} \tilde{S}_n + \tau_t \right\|_{2p} dt - \frac{1}{2} \log(1 - \frac{D}{\sqrt{n}}).\end{aligned}$$

Proof As a direct consequence of Thm. 5 applied to S_n , for all $D > 0, p \geq 1$ we have

$$\begin{aligned}\sqrt{2}H(\tilde{S}_n, Z) &\leq \int_{-\frac{1}{2} \log(1-\frac{D}{\sqrt{n}})}^{\infty} \left\| \frac{e^{-2t}}{1-e^{-2t}} (Z^2 - 1) - \frac{e^{-t}}{\sqrt{1-e^{-2t}}} \tilde{S}_n Z + \tau'_t \right\|_2 dt \\ &\quad + \int_0^{-\frac{1}{2} \log(1-\frac{D}{\sqrt{n}})} \|S_{\tilde{S}_n}(\tilde{S}_n)\|_{2\frac{p}{p-1}} \left\| \frac{Ze^{-2t}}{\sqrt{1-e^{-2t}}} - e^{-t} \tilde{S}_n + \tau_t \right\|_{2p} dt - \frac{1}{2} \log(1 - \frac{D}{\sqrt{n}}).\end{aligned}$$

Moreover, we know that the score function $S_{\tilde{S}_n}(\tilde{S}_n)$ is the only \tilde{S}_n -measurable random variable such that for all $\Phi \in \mathbb{C}^\infty(\mathbb{R})$ we have

$$\mathbb{E}(\Phi(\tilde{S}_n) S_{\tilde{S}_n}(\tilde{S}_n)) = \mathbb{E}(\Phi'(\tilde{S}_n)).$$

Therefore we note that

$$S_{\tilde{S}_n}(\tilde{S}_n) = \frac{1}{\sqrt{n}} \mathbb{E}(\sum_{i=1} S_{Y_1}(Y_i) | \tilde{S}_n).$$

Combined with the tower property we therefore have:

$$\|S_{\tilde{S}_n}(\tilde{S}_n)\|_{\frac{2p}{p-1}} \leq \left\| \frac{1}{\sqrt{n}} \sum_{i=1} S_{Y_1}(Y_i) \right\|_{\frac{2p}{p-1}}.$$

Therefore as $\mathbb{E}(S_{Y_1}(Y_1)) = 0$, then using Lem. 39 and Lem. 40 we observe that

$$\|S_{\tilde{S}_n}(\tilde{S}_n)\|_{\frac{2p}{p-1}} \leq C_p.$$

This directly implies that

$$\begin{aligned}\sqrt{2}H(\tilde{S}_n, Z) &\leq \int_{-\frac{1}{2} \log(1-\frac{D}{\sqrt{n}})}^{\infty} \left\| \frac{e^{-2t}}{1-e^{-2t}} (Z^2 - 1) - \frac{e^{-t}}{\sqrt{1-e^{-2t}}} \tilde{S}_n Z + \tau'_t \right\|_2 dt \\ &\quad + \int_0^{-\frac{1}{2} \log(1-\frac{D}{\sqrt{n}})} \inf_{p \geq 1} C_p \left\| \frac{Ze^{-2t}}{\sqrt{1-e^{-2t}}} - e^{-t} \tilde{S}_n + \tau_t \right\|_{2p} dt - \frac{1}{2} \log(1 - \frac{D}{\sqrt{n}}).\end{aligned}$$

■

C.1. Bounding the different terms

Throughout this subsection, we write $T_n \triangleq -\frac{1}{2} \log(1 - \frac{D}{\sqrt{n}})$

Lemma 19 *If Assump. (R, σ), Assump. (\tilde{f}) and Assump. (S_{Y_1}, p') hold then for all $\kappa \geq \max(\frac{\tilde{R}^2}{n}, \frac{\tilde{R}^2}{\sqrt{n}D})$ then for all $p \geq 1$ the following upper bound holds*

$$\begin{aligned} & \int_{-\frac{1}{2} \log(1 - \frac{D}{\sqrt{n}})}^{\infty} \left\| \frac{e^{-2t}}{1 - e^{-2t}} (Z^2 - 1) - \frac{e^{-t}}{\sqrt{1 - e^{-2t}}} \tilde{S}_n Z - \tau'_t \right\|_{2p} dt \\ & \leq \frac{\sqrt{3e} \tilde{R}^2 (2p-1) (\sqrt{n} + A_{2p} \tilde{R} + \tilde{R}^2 \tilde{A}_{n,2p})}{\kappa \sqrt{2} \sqrt{n}^3} \int_0^{1 - \frac{D}{\sqrt{n}}} \frac{(e^{2(2p-1)x\kappa} - 1)}{(1-x)^2} dx \\ & \quad + \frac{2\sqrt{e(2p-1)} \tilde{R}^2 (A_p + \tilde{R} \tilde{A}_{n,2p})}{\sqrt{3n\kappa}} \int_0^{\sqrt{1 - \frac{D}{\sqrt{n}}}} \frac{(e^{2x^2(2p-1)\kappa} - 1)}{\sqrt{1-x^2}^3} dx \\ & \quad + \frac{\log(n/D^2)}{4\sqrt{2}\sqrt{n}} \min \left\{ \sqrt{2p-1} \left(\frac{\tilde{R}^2}{\tilde{R}^{1/p}} - 1 \right) \right. \\ & \quad \left. (\tilde{R}^2 - \tilde{R}^{1/p}) \tilde{A}_{n,2p} + (\tilde{R} - 1) A_{2p} \right\}. \end{aligned}$$

Notably if $p = 1$ and if we write $M_D^n \triangleq \sqrt{1 - \frac{D}{\sqrt{n}}}$ then for all this can be further upper bounded as:

$$\begin{aligned} & \int_{T_n}^{\infty} \left\| \frac{e^{-2t}}{1 - e^{-2t}} (Z^2 - 1) - \frac{e^{-t}}{\sqrt{1 - e^{-2t}}} \tilde{S}_n Z - \tau'_t \right\|_{2p} dt \\ & \leq \frac{\sqrt{3e} \tilde{R}^2 (\sqrt{n} + \tilde{R}) (e^{2\kappa(M_D^n)^2} - 1)}{\kappa C \sqrt{2n}} + \frac{\sqrt{e} \tilde{R}^2 M_D^n (e^{2\kappa(M_D^n)^2} - 1)}{\sqrt{3D} \sqrt{n\kappa}} + \frac{\log(n/D^2)}{4\sqrt{2}\sqrt{n}} (\tilde{R} - 1) \\ & \sim \frac{\sqrt{e} \tilde{R}^2 (e^{2\kappa} - 1)}{\sqrt{3D} \sqrt{n\kappa}} + \frac{\log(n/D^2)}{4\sqrt{2}\sqrt{n}} (\tilde{R} - 1). \end{aligned}$$

Proof

Using the definition of τ'_t and triangle inequality we obtain that

$$\begin{aligned} & \int_{-\frac{1}{2} \log(1 - \frac{D}{\sqrt{n}})}^{\infty} \left\| \frac{e^{-2t}}{1 - e^{-2t}} (Z^2 - 1) - \frac{e^{-t}}{\sqrt{1 - e^{-2t}}} \tilde{S}_n Z - \tau'_t \right\|_{2p} dt \\ & \leq \int_{-\frac{1}{2} \log(1 - \frac{D}{\sqrt{n}})}^{\infty} \frac{e^{-t} \|H_1\|_2}{\sqrt{1 - e^{-2t}}} \|n\mathbb{E}(Y_t | \tilde{S}_n) - \tilde{S}_n\|_{2p} dt \\ & \quad + \int_{-\frac{1}{2} \log(1 - \frac{D}{\sqrt{n}})}^{\infty} \frac{e^{-2t} \|H_2\|_2}{1 - e^{-2t}} \left\| \frac{n}{2} \mathbb{E}((Y_t)^2 | \tilde{S}_n) - 1 \right\|_{2p} dt \\ & \quad + \int_{-\frac{1}{2} \log(1 - \frac{D}{\sqrt{n}})}^{\infty} \sum_{k \geq 3} \frac{e^{-tk} \|H_k\|_{2p}}{k! (\sqrt{1 - e^{-2t}})^k} n \|\mathbb{E}((Y_t)^k | \tilde{S}_n)\|_{2p} dt \\ & \triangleq (a_1) + (a_2) + (a_3). \end{aligned}$$

We bound each term successively. For ease of notation we write $c(t) \triangleq \sqrt{\kappa(1 - e^{-2t})}$.

Firstly we note that, if $t \geq -\frac{1}{2} \log(1 - \frac{\tilde{R}^2}{n\kappa}) \geq 0$ then $c(t) \geq \tilde{R}/\sqrt{n}$ therefore $Y_t = X_I - X'_I$. As I is a randomly drawn index in $\{1, \dots, n\}$, this implies that

$$\mathbb{E}(Y_t | \tilde{S}_n) = \frac{1}{n} \sum_{i \leq n} \mathbb{E}(X_i - X'_i | \tilde{S}_n) = \frac{1}{n} \sum_{i \leq n} X_i - \mathbb{E}(X_i) = \frac{1}{n} \tilde{S}_n; \quad (15)$$

which implies that $\|n\mathbb{E}(Y_t | \tilde{S}_n) - \tilde{S}_n\|_{2p} = 0$. As $-\frac{1}{2} \log(1 - \frac{\tilde{R}^2}{n\kappa}) \leq -\frac{1}{2} \log(1 - \frac{D}{\sqrt{n}})$ we have that $(a_1) = 0$.

We now work on upper bounding (a_2) . In this goal, we first remark that if $t \geq -\frac{1}{2} \log(1 - \frac{\tilde{R}^2}{n\kappa})$ we have $Y_t = X'_I - X_I$ which implies that:

$$\|\frac{n}{2}\mathbb{E}((Y_t)^2|\tilde{S}_n) - 1\|_{2p} \leq \|\frac{1}{2}\sum_{i \leq n} \mathbb{E}(X_i^2) + X_i^2 - 1\|_{2p} \leq \frac{1}{2}\|\sum_{i \leq n} X_i^2 - 1\|_{2p}.$$

We remark using Eq. (33) that $\|X_i^2 - 1/n\|_{2p} \leq \frac{1}{n}(\tilde{R}^{2-1/p} - 1)$ and $\|X_i^2 - 1/n\|_2 \leq \frac{1}{n}(\tilde{R} - 1)$. Therefore using Lem. 39 and Lem. 40 we obtain:

$$\|\frac{n}{2}\mathbb{E}((Y_t)^2|\tilde{S}_n) - 1\|_{2p} \leq \min \left\{ \frac{\sqrt{2p-1}}{2\sqrt{n}} \left(\frac{\tilde{R}^2}{\tilde{R}^{1/p}} - 1 \right) \right. \\ \left. \frac{1}{2\sqrt{n}} \left((\tilde{R}^2 - \tilde{R}^{1/p})\tilde{A}_{n,2p} + (\tilde{R} - 1)A_{2p} \right) \right\} \triangleq B_{n,2p}.$$

This implies that

$$\int_{-\frac{1}{2} \log(1 - \frac{D}{\sqrt{n}})}^{\infty} \|\frac{n}{2}\mathbb{E}((Y_t)^2|\tilde{S}_n) - 1\|_{2p} \frac{e^{-2t}}{1-e^{-2t}} dt \\ \leq B_{n,2p} \int_{-\frac{1}{2} \log(1 - \frac{D}{\sqrt{n}})}^{\infty} \frac{e^{-2t}}{1-e^{-2t}} dt = \frac{B_{n,2p}}{2} \int_0^{1 - \frac{D}{\sqrt{n}}} \frac{1}{1-t} dt \\ = \frac{B_{n,2p}}{4} \log(n/D^2) = \frac{\log(n/D^2)}{8\sqrt{n}} \min \left\{ \sqrt{2p-1} \left(\frac{\tilde{R}^2}{\tilde{R}^{1/p}} - 1 \right) \right. \\ \left. (\tilde{R}^2 - \tilde{R}^{1/p})\tilde{A}_{n,2p} + (\tilde{R} - 1)A_{2p} \right\}.$$

Therefore we obtain that

$$(a_2) \leq \frac{\log(n/D^2)\sqrt{2p-1}}{4\sqrt{2}\sqrt{n}} \min \left\{ \sqrt{2p-1} \left(\frac{\tilde{R}^2}{\tilde{R}^{1/p}} - 1 \right) \right. \\ \left. (\tilde{R}^2 - \tilde{R}^{1/p})\tilde{A}_{n,2p} + (\tilde{R} - 1)A_{2p} \right\}.$$

Finally we work on bounding (a_3) . We observe that if we write $Z_i^{t,k} \triangleq \mathbb{E}((X_i - X'_i)^k \mathbb{I}(|X'_i| \leq c(t)) | X_{1:n}) \mathbb{I}(|X_i| \leq c(t))$ then

$$n\mathbb{E}(Y_t^k | X_{1:n}) = \sum_{i \leq n} Z_i^{t,k}.$$

Let $k \geq 3$ be an *odd* integer. We observe that the random variables $(Z_i^{t,k})$ are symmetric and therefore have a mean of zero. This implies that $n\mathbb{E}(Y_t^k | \tilde{S}_n) = O_p(\frac{1}{\sqrt{n}})$.

To make this clear $\tilde{m}_{k,2p}(t) \triangleq c(t)^{k-2} \min(\sqrt{n}c(t), nc(t)^2, \tilde{R})$ then we also have

$$\|Z_i^{t,k}\|_{2p} \leq \frac{2^k \tilde{m}_{k,2p}(t)}{n}.$$

Therefore we obtain that for all $t > 0$:

$$n\|\mathbb{E}(Y_t^k | \tilde{S}_n)\|_2 \leq \frac{2^k}{\sqrt{n}} \tilde{m}_{k,2p}(t).$$

This implies that if $k \geq 3$ is odd we have

$$\int_{-\frac{1}{2} \log(1 - \frac{D}{\sqrt{n}})}^{\infty} \frac{e^{-kt}}{\sqrt{1-e^{-2t}}} n \|\mathbb{E}((Y_t)^k | \tilde{S}_n)\|_{2p} dt \\ \leq \frac{2^k \tilde{R}^2 (\tilde{R}\tilde{A}_{n,2p} + A_{2p})}{n} \int_{-\frac{1}{2} \log(1 - \frac{D}{\sqrt{n}})}^{\infty} \frac{e^{-tk} c(t)^{k-3}}{\sqrt{1-e^{-2t}}} dt \\ \leq \frac{2^k \tilde{R}^2 (\tilde{R}\tilde{A}_{n,2p} + A_{2p}) \sqrt{\kappa}^{k-3}}{n} \int_{-\frac{1}{2} \log(1 - \frac{D}{\sqrt{n}})}^{\infty} \frac{e^{-tk}}{\sqrt{1-e^{-2t}}} dt \\ \leq \frac{2^k \tilde{R}^2 (\tilde{R}\tilde{A}_{n,2p} + A_{2p}) \sqrt{\kappa}^{k-3}}{n} \int_0^{\sqrt{1 - \frac{D}{\sqrt{n}}}} \frac{x^{k-1}}{\sqrt{1-x^2}} dx.$$

We remark that

$$\begin{aligned} \sum_{\substack{k \geq 3 \\ k \text{ odd}}} \frac{2^k x^{k-1} \sqrt{2p-1}^k \sqrt{\kappa}^k}{\sqrt{k!}} &\leq \sqrt{2e\kappa} \sqrt{2p-1} \sum_{k \geq 1} \frac{2^k x^{2k} (2p-1)^k \kappa^k \sqrt{k+1}}{k! \sqrt{2k+1}} \\ &\leq \frac{2\sqrt{e\kappa} \sqrt{2p-1}}{\sqrt{3}} (e^{2x^2(2p-1)\kappa} - 1). \end{aligned}$$

Therefore we have:

$$\begin{aligned} \sum_{k \geq 3} \frac{\|H_k\|_{2p}}{k!} \int_0^\infty \frac{e^{-tk}}{\sqrt{1-e^{-2t}}} n \|\mathbb{E}((Y_t)^k | \tilde{S}_n)\|_{2p} dt \\ \leq \frac{2\sqrt{e(2p-1)} \tilde{R}^2 (\tilde{R} \tilde{A}_{n,2p} + A_{2p})}{\sqrt{3n\kappa}} \int_0^{\sqrt{1-\frac{D}{\sqrt{n}}}} \frac{(e^{2x^2(2p-1)\kappa} - 1)}{\sqrt{1-x^2}^3} dx. \end{aligned}$$

If $k \geq 4$ and is even then we have:

$$\begin{aligned} n \|\mathbb{E}(Y_t^k | \tilde{S}_n)\|_{2p} &\leq \|\sum_{i \leq n} \mathbb{E}[\mathbb{I}(|X_i|, |X'_i| \leq c(t)) (X_i - X'_i)^k | \tilde{S}_n]\|_{2p} \\ &\leq \|\sum_{i \leq n} \mathbb{E}[\mathbb{I}(|X_i|, |X'_i| \leq c(t)) (X_i - X'_i)^k | \tilde{S}_n] \\ &\quad - \mathbb{E}[\mathbb{I}(|X_i|, |X'_i| \leq c(t)) (X_i - X'_i)^k]\|_{2p} \\ &\quad + n \|\mathbb{E}[\mathbb{I}(|X_1|, |X'_1| \leq c(t)) (X_1 - X'_1)^k]\| \\ &\leq \frac{2^k c(t)^{k-4}}{\sqrt{n}^3} \tilde{R}^2 (\sqrt{n} + \tilde{R} U_{n,2p}). \end{aligned}$$

This directly implies that we have:

$$\begin{aligned} \int_{-\frac{1}{2} \log(1-\frac{D}{\sqrt{n}})}^\infty \frac{e^{-tk}}{\sqrt{1-e^{-2t}}} n \|\mathbb{E}((Y_t)^k | \tilde{S}_n)\|_{2p} dt \\ \leq \frac{2^k \tilde{R}^2 \sqrt{\kappa}^{k-4} (\sqrt{n} + A_{2p} \tilde{R} + \tilde{R}^2 \tilde{A}_{n,2p})}{\sqrt{n}^3} \int_{-\frac{1}{2} \log(1-\frac{D}{\sqrt{n}})}^\infty \frac{e^{-tk}}{\sqrt{1-e^{-2t}}^4} dt \\ \leq \frac{2^k \tilde{R}^2 \sqrt{\kappa}^{k-4} (\sqrt{n} + A_{2p} \tilde{R} + \tilde{R}^2 \tilde{A}_{n,2p})}{\sqrt{n}^3} \int_0^{\sqrt{1-\frac{D}{\sqrt{n}}}} \frac{x^{k-1}}{\sqrt{1-x^2}^4} dx. \end{aligned}$$

We remark that

$$\begin{aligned} \sum_{\substack{k \geq 4 \\ k \text{ even}}} \frac{2^k x^{k-1} \sqrt{2p-1}^k \sqrt{\kappa}^k}{\sqrt{k!}} &\leq \frac{\sqrt{e}}{\sqrt{2}} \sum_{k \geq 2} \frac{2^k x^{2k-1} (2p-1)^k \kappa^k \sqrt{k+1}}{k!} \\ &\leq \frac{\sqrt{3e} x \kappa (2p-1)}{\sqrt{2}} [e^{2x^2(2p-1)\kappa} - 1]. \end{aligned}$$

Therefore we have:

$$\begin{aligned} \sum_{\substack{k \geq 4 \\ k \text{ is even}}} \frac{\|H_k\|_{2p}}{k!} \int_{-\frac{1}{2} \log(1-\frac{D}{\sqrt{n}})}^\infty \frac{e^{-tk}}{\sqrt{1-e^{-2t}}} n \|\mathbb{E}((Y_t)^k | \tilde{S}_n)\|_{2p} dx \\ \leq \frac{\sqrt{3e} \tilde{R}^2 (\sqrt{n} + A_{2p} \tilde{R} + \tilde{R}^2 \tilde{A}_{n,2p}) (2p-1)}{2\sqrt{2}\kappa \sqrt{n}^3} \int_0^{\sqrt{1-\frac{D}{\sqrt{n}}}} \frac{x(e^{2x^2(2p-1)\kappa} - 1)}{\sqrt{1-x^2}^4} dx \\ \leq \frac{\sqrt{3e} \tilde{R}^2 (\sqrt{n} + A_{2p} \tilde{R} + \tilde{R}^2 \tilde{A}_{n,2p}) (2p-1)}{\kappa \sqrt{2} \sqrt{n}^3} \int_0^{1-\frac{D}{\sqrt{n}}} \frac{(e^{2x(2p-1)\kappa} - 1)}{(1-x)^2} dx. \end{aligned}$$

Therefore by combining this together we obtain that

$$\begin{aligned} (a_3) &\leq \frac{\sqrt{3e} \tilde{R}^2 (\sqrt{n} + A_{2p} \tilde{R} + \tilde{R}^2 \tilde{A}_{n,2p}) (2p-1)}{\kappa \sqrt{2} \sqrt{n}^3} \int_0^{1-\frac{D}{\sqrt{n}}} \frac{(e^{2(2p-1)x\kappa} - 1)}{(1-x)^2} dx \\ &\quad + \frac{2\sqrt{e(2p-1)} \tilde{R}^2 (A_p + \tilde{R} \tilde{A}_{n,2p})}{\sqrt{3n\kappa}} \int_0^{\sqrt{1-\frac{D}{\sqrt{n}}}} \frac{(e^{2x^2(2p-1)\kappa} - 1)}{\sqrt{1-x^2}^3} dx. \end{aligned}$$

Therefore we obtain that

$$\begin{aligned}
& \int_{-\frac{1}{2}}^{\infty} \log(1 - \frac{D}{\sqrt{n}}) \left\| \frac{e^{-2t}}{1-e^{-2t}} (Z^2 - 1) - \frac{e^{-t}}{\sqrt{1-e^{-2t}}} \tilde{S}_n Z - \tau'_t \right\|_{2p} \\
& \leq \frac{\sqrt{3e}\tilde{R}^2(2p-1)(\sqrt{n}+A_{2p}\tilde{R}+\tilde{R}^2\tilde{A}_{n,2p})}{\kappa\sqrt{2}\sqrt{n}^3} \int_0^{1-\frac{D}{\sqrt{n}}} \frac{(e^{2(2p-1)x\kappa}-1)}{(1-x)^2} dx \\
& \quad + \frac{2\sqrt{e(2p-1)}\tilde{R}^2(A_p+\tilde{R}\tilde{A}_{n,2p})}{\sqrt{3n\kappa}} \int_0^{\sqrt{1-\frac{D}{\sqrt{n}}}} \frac{(e^{2x^2(2p-1)\kappa}-1)}{\sqrt{1-x^2}^3} dx \\
& \quad + \frac{\log(n/D^2)}{4\sqrt{2}\sqrt{n}} \min \left\{ \sqrt{2p-1} \left(\frac{\tilde{R}^2}{\tilde{R}^{1/p}} - 1 \right) \right. \\
& \quad \left. (\tilde{R}^2 - \tilde{R}^{1/p})\tilde{A}_{n,2p} + (\tilde{R} - 1)A_{2p} \right\}.
\end{aligned}$$

■

Lemma 20 *If the conditions Assump. (R, σ), Assump. (f̃) and Assump. (S_{Y₁}, p') hold then we have*

$$\begin{aligned}
\int_0^{T_n} \left\| \frac{Ze^{-2t}}{\sqrt{1-e^{-2t}}} - e^{-t}\tilde{S}_n + \tau_t \right\|_{2p} dt & \leq \int_0^{T_n} e^{-t} \|n\mathbb{E}(Y_t|\tilde{S}_n) - \tilde{S}_n\|_{2p} dt \\
& \quad + \int_0^{T_n} \frac{e^{-2t}\|H_1\|_{2p}}{\sqrt{1-e^{-2t}}} \left\| \frac{n}{2} \mathbb{E}((Y_t)^2|\tilde{S}_n) - 1 \right\|_{2p} dt \\
& \quad + \sum_{k \geq 3} \int_0^{T_n} \frac{e^{-kt}\|H_{k-1}\|_{2p}}{k!(\sqrt{1-e^{-2t}})^{k-1}} n \|\mathbb{E}((Y_t)^k|\tilde{S}_n)\|_{2p} dt \\
& \triangleq (a'_{1,p}) + (a'_{2,p}) + (a'_{3,p}).
\end{aligned}$$

Proof By definition of τ_t we have

$$\begin{aligned}
\int_0^{T_n} \left\| \frac{Ze^{-2t}}{\sqrt{1-e^{-2t}}} - e^{-t}\tilde{S}_n + \tau_t \right\|_{2p} dt & \leq \int_0^{T_n} e^{-t} \|n\mathbb{E}(Y_t|\tilde{S}_n) - \tilde{S}_n\|_{2p} dt \\
& \quad + \int_0^{T_n} \frac{e^{-2t}\|H_1\|_{2p}}{\sqrt{1-e^{-2t}}} \left\| \frac{n}{2} \mathbb{E}((Y_t)^2|\tilde{S}_n) - 1 \right\|_{2p} dt \\
& \quad + \sum_{k \geq 3} \int_0^{T_n} \frac{e^{-kt}\|H_{k-1}\|_{2p}}{k!(\sqrt{1-e^{-2t}})^{k-1}} n \|\mathbb{E}((Y_t)^k|\tilde{S}_n)\|_{2p} dt \\
& \triangleq (a'_{1,p}) + (a'_{2,p}) + (a'_{3,p}).
\end{aligned}$$

■

Lemma 21 *The upper bound found for $(a'_{1,p})$, $(a'_{2,p})$ and $(a'_{3,p})$ found in Eq. (20), Eq. (22) and Eq. (23) are valid. Moreover the following also hold:*

$$\begin{aligned}
(a'_{1,p}) & \lesssim \frac{\tilde{R}^2}{2n\kappa M_{n,\kappa}} \left[U_{n,2p} + \tilde{U}_{n,2p} \right], \quad (a'_{2,p}) \lesssim \frac{\sqrt{2p-1}\tilde{R}}{\sqrt{n\kappa}}, \quad \text{and} \\
(a'_{3,p}) & \lesssim \frac{2\sqrt{e}\sqrt{2p-1}\tilde{R}}{\sqrt{3n\kappa}} \left[e^{2(2p-1)\kappa} - 1 - 2(2p-1)\kappa \right] \left[2 - \frac{2}{3\tilde{R}} \right].
\end{aligned}$$

Proof For ease of notation we write $c(t) \triangleq \sqrt{\kappa(1-e^{-2t})}$. Define (Z_i^t) as

$$Z_i^t \triangleq \begin{cases} -X_i & \text{if } |X_i| \geq c(t) \\ \mathbb{E}(X_i \mathbb{I}(|X_i| > c(t))) - X_i \mathbb{P}(|X_i| > c(t)) & \text{otherwise} \end{cases}.$$

Then we have $n\mathbb{E}(Y_t|\tilde{S}_n) - \tilde{S}_n = \sum_{i \leq n} Z_i^t$. Suppose that $t \leq -\frac{1}{2} \log(1 - \frac{\tilde{R}^2}{n\kappa})$ then $c(t) \leq \tilde{R}/\sqrt{n}$. Therefore using the triangle inequality we note that the norm of the variables (Z_i^t) are bounded by

$$\begin{aligned} \|Z_i^t\|_{2p} &\leq \|X_i\|_p + \|X_i \mathbb{I}(|X_i| \leq c(t))\|_{2p} \stackrel{(a)}{\leq} \frac{1}{\sqrt{n}} \frac{\tilde{R}}{\tilde{R}^{1/p}} + \min(n^{-1/2p} c(t)^{1-\frac{1}{p}}, c(t)); \\ \|Z_i^t\|_2 &\leq \|X_i\|_2 + \|X_i \mathbb{I}(|X_i| \leq c(t))\|_2 \stackrel{(a)}{\leq} \frac{1}{\sqrt{n}} + \mu_{n,\kappa,1}; \end{aligned}$$

where (a) is obtained using Eq. (33). Moreover according to Lem. 40 we have

$$\begin{aligned} \|n\mathbb{E}(Y_t|\tilde{S}_n) - \tilde{S}_n\|_{2p} & \\ &= A_{2p}(1 + \min(1, \sqrt{n}c(t))) + \tilde{A}_{n,2p}(\tilde{R} + \tilde{R}^{1/p} \sqrt{n} \min(n^{-\frac{1}{2p}} c(t)^{1-\frac{1}{p}}, c(t))). \end{aligned} \quad (16)$$

Moreover according to Lem. 39 we have

$$\|n\mathbb{E}(Y_t|\tilde{S}_n) - \tilde{S}_n\|_{2p} \leq \sqrt{2p-1} \left[\frac{\tilde{R}}{\tilde{R}^{1/p}} + \sqrt{n} \min(n^{-\frac{1}{2p}} c(t)^{1-\frac{1}{p}}, c(t)) \right]. \quad (17)$$

By combining Eq. (34) and Eq. (35) we obtain

$$\|n\mathbb{E}(Y_t|\tilde{S}_n) - \tilde{S}_n\|_{2p} \leq \min \left\{ \begin{aligned} &\sqrt{2p-1} \left[\frac{\tilde{R}}{\tilde{R}^{1/p}} + \sqrt{n} \min(n^{-\frac{1}{2p}} c(t)^{1-\frac{1}{p}}, c(t)) \right] \\ &\tilde{A}_{n,2p}(\tilde{R} + \tilde{R}^{1/p} \sqrt{n} \min(n^{-1/2p} c(t)^{1-\frac{1}{p}}, c(t))) \\ &+ A_{2p}(1 + \sqrt{n} \mu_{n,\kappa,1}). \end{aligned} \right. \quad (18)$$

Moreover, if $t \geq -\frac{1}{2} \log(1 - \frac{\tilde{R}^2}{n\kappa}) \geq 0$ then $c(t) \geq \tilde{R}/\sqrt{n}$ therefore $Y_t = X_I - X'_I$. As I is a randomly drawn index in $\{1, \dots, n\}$, this implies that

$$\mathbb{E}(Y_t|\tilde{S}_n) = \frac{1}{n} \sum_{i \leq n} \mathbb{E}(X_i - X'_i|\tilde{S}_n) = \frac{1}{n} \sum_{i \leq n} X_i - \mathbb{E}(X_i) = \frac{1}{n} \tilde{S}_n; \quad (19)$$

which implies that $\|n\mathbb{E}(Y_t|\tilde{S}_n) - \tilde{S}_n\|_{2p} = 0$.

Combining Eq. (36) and Eq. (37) we obtain that we can upper $(a'_{1,p})$ as:

$$\begin{aligned} (a'_{1,p}) &\leq \int_0^{T_n} e^{-t} \|n\mathbb{E}(Y_t|\tilde{S}_n) - \tilde{S}_n\|_{2p} dt \\ &\stackrel{(a)}{\leq} (\tilde{R} \tilde{A}_{n,2p} + A_{2p}) \int_0^{-\frac{1}{2} \log(1 - \frac{\tilde{R}^2}{n\kappa})} e^{-t} dt \\ &\quad + \tilde{R}^{1/p} \sqrt{n\kappa}^{1-\frac{1}{p}} \tilde{A}_{n,2p} \int_{-\frac{1}{2} \log(1 - \frac{\tilde{R}^2}{n\kappa})}^{-\frac{1}{2} \log(1 - \frac{1}{n\kappa})} e^{-t} \sqrt{1 - e^{-2t}}^{1-\frac{1}{p}} dt \\ &\quad + \sqrt{n\kappa} (\tilde{R}^{1/p} \tilde{A}_{n,2p} + A_{2p}) \int_0^{-\frac{1}{2} \log(1 - \frac{1}{n\kappa})} e^{-t} \sqrt{1 - e^{-2t}} dt \\ &\quad + A_{2p} \int_{-\frac{1}{2} \log(1 - \frac{\tilde{R}^2}{n\kappa})}^{-\frac{1}{2} \log(1 - \frac{1}{n\kappa})} e^{-t} dt \\ &= \tilde{R}^{1/p} \sqrt{n\kappa}^{1-\frac{1}{p}} \tilde{A}_{n,2p} \left(\tilde{M}_{n,\kappa} {}_2F_1\left(\frac{1}{2}, \frac{1}{2p} - \frac{1}{2}, \frac{3}{2}, \tilde{M}_{n,\kappa}^2\right) \right. \\ &\quad \left. - M_{n,\kappa} {}_2F_1\left(\frac{1}{2}, \frac{1}{2p} - \frac{1}{2}, \frac{3}{2}, M_{n,\kappa}^2\right) \right) + [A_{2p} + \tilde{R} \tilde{A}_{n,2p}] (1 - M_{n,\kappa}) \\ &\quad + \frac{\sqrt{n\kappa} (\tilde{R}^{1/p} \tilde{A}_{n,2p} + A_{2p})}{2} \left(\cos^{-1}(\tilde{M}_{n,\kappa}) - \frac{\tilde{M}_{n,\kappa}}{\sqrt{n\kappa}} \right) \\ &\quad + A_{2p} (\tilde{M}_{n,\kappa} - M_{n,\kappa}). \end{aligned}$$

where to obtain (a) we used the fact that $c(t) = \sqrt{\kappa}\sqrt{1 - e^{-2t}}$.

This can be further upper-bounded in the following way:

$$\begin{aligned}
(a'_{1,p}) &\leq \int_0^\infty e^{-t} \|n\mathbb{E}(Y_t|S_n) - S_n\|_{2p} dt \\
&\stackrel{(a)}{\leq} (\tilde{R}\tilde{A}_{n,2p} + 2A_{2p}) \int_0^{-\frac{1}{2}\log(1-\frac{\tilde{R}^2}{n\kappa})} e^{-t} dt \\
&\quad + \tilde{R}^{1/p} \sqrt{n\kappa} \tilde{A}_{n,2p} \int_0^{-\frac{1}{2}\log(1-\frac{\tilde{R}^2}{n\kappa})} e^{-t} \sqrt{1 - e^{-2t}} dt \\
&= \left[2A_{2p} + \tilde{R}\tilde{A}_{n,2p} \right] (1 - \sqrt{1 - \frac{\tilde{R}^2}{n\kappa}}) \\
&\quad + \frac{\tilde{R}^{1/p} \sqrt{n\kappa} \tilde{A}_{n,2p}}{2} K_{n,\kappa,p,3} \\
&= \frac{\tilde{R}^{1/p} \sqrt{n\kappa} \tilde{A}_{n,2p}}{2} K_{n,\kappa,p,3} \\
&\quad + \left[2A_{2p} + \tilde{R}\tilde{A}_{n,2p} \right] (1 - M_{n,\kappa}) \\
&\leq \frac{\tilde{R}^2}{2n\kappa M_{n,\kappa}} \left[U_{n,2p} + \tilde{U}_{n,2p} \right].
\end{aligned}$$

We remark that $(a'_{1,p})$ can be alternatively upper bounded as

$$\begin{aligned}
(a'_{1,p}) &\leq \int_0^\infty e^{-t} \|n\mathbb{E}(Y_t|S_n) - S_n\|_{2p} dt \\
&\leq \frac{\tilde{R}\sqrt{2p-1}}{\tilde{R}^{1/p}} \int_0^{-\frac{1}{2}\log(1-\frac{\tilde{R}^2}{n\kappa})} e^{-t} dt \\
&\quad + \sqrt{2p-1} \sqrt{\kappa n}^{1-\frac{1}{p}} \int_{-\frac{1}{2}\log(1-\frac{1}{n\kappa})}^{-\frac{1}{2}\log(1-\frac{\tilde{R}^2}{n\kappa})} e^{-t} \sqrt{1 - e^{-2t}}^{1-\frac{1}{p}} dt \\
&\quad + \sqrt{2p-1} \sqrt{\kappa n} \int_0^{-\frac{1}{2}\log(1-\frac{1}{n\kappa})} e^{-t} \sqrt{1 - e^{-2t}} dt \\
&= \sqrt{2p-1} \left\{ \sqrt{\kappa n}^{1-\frac{1}{p}} \left(\tilde{M}_{n,\kappa} {}_2F_1\left(\frac{1}{2}, \frac{1}{2p} - \frac{1}{2}, \frac{3}{2}, \tilde{M}_{n,\kappa}^2\right) \right. \right. \\
&\quad \left. \left. - M_{n,\kappa} {}_2F_1\left(\frac{1}{2}, \frac{1}{2p} - \frac{1}{2}, \frac{3}{2}, M_{n,\kappa}^2\right) \right) + \frac{\tilde{R}}{\tilde{R}^{1/p}} (1 - M_{n,\kappa}) \right. \\
&\quad \left. + \frac{\sqrt{\kappa n}}{2} \left[\cos^{-1}(\tilde{M}_{n,\kappa}) - \frac{\tilde{M}_{n,\kappa}}{\sqrt{n\kappa}} \right] \right\}.
\end{aligned}$$

Therefore we have

$$(a'_{1,p}) \leq \min \left\{ \begin{aligned} &\tilde{R}^{1/p} \sqrt{n\kappa}^{1-\frac{1}{p}} \tilde{A}_{n,2p} \left(\tilde{M}_{n,\kappa} {}_2F_1\left(\frac{1}{2}, \frac{1}{2p} - \frac{1}{2}, \frac{3}{2}, \tilde{M}_{n,\kappa}^2\right) \right. \\ &\quad \left. - M_{n,\kappa} {}_2F_1\left(\frac{1}{2}, \frac{1}{2p} - \frac{1}{2}, \frac{3}{2}, M_{n,\kappa}^2\right) \right) + \left[A_{2p} + \tilde{R}\tilde{A}_{n,2p} \right] (1 - M_{n,\kappa}) \\ &\quad + \frac{\sqrt{n\kappa} (\tilde{R}^{1/p} \tilde{A}_{n,2p} + A_{2p})}{2} \left(\cos^{-1}(\tilde{M}_{n,\kappa}) - \frac{\tilde{M}_{n,\kappa}}{\sqrt{n\kappa}} \right) + A_{2p} (\tilde{M}_{n,\kappa} - M_{n,\kappa}) \\ &\sqrt{2p-1} \left\{ \sqrt{\kappa n}^{1-\frac{1}{p}} \left(\tilde{M}_{n,\kappa} {}_2F_1\left(\frac{1}{2}, \frac{1}{2p} - \frac{1}{2}, \frac{3}{2}, \tilde{M}_{n,\kappa}^2\right) \right. \right. \\ &\quad \left. \left. - M_{n,\kappa} {}_2F_1\left(\frac{1}{2}, \frac{1}{2p} - \frac{1}{2}, \frac{3}{2}, M_{n,\kappa}^2\right) \right) + \frac{\tilde{R}}{\tilde{R}^{1/p}} (1 - M_{n,\kappa}) \right. \\ &\quad \left. + \frac{\sqrt{\kappa n}}{2} \left[\cos^{-1}(\tilde{M}_{n,\kappa}) - \frac{\tilde{M}_{n,\kappa}}{\sqrt{n\kappa}} \right] \right\}. \end{aligned} \right. \quad (20)$$

We now work on upper bounding $(a'_{2,p})$. In this goal, we first remark that if $t \geq -\frac{1}{2}\log(1 - \frac{\tilde{R}^2}{n\kappa})$ we have $Y_t = X'_I - X_I$ which implies that:

$$\|\frac{n}{2}\mathbb{E}((Y_t)^2|\tilde{S}_n) - 1\|_{2p} \leq \|\frac{1}{2}\sum_{i \leq n} \mathbb{E}(X_i^2) + X_i^2 - 1\|_{2p} \leq \frac{1}{2}\|\sum_{i \leq n} X_i^2 - 1\|_{2p}.$$

We remark using Eq. (33) that $\|X_i^2 - 1/n\|_{2p} \leq \frac{1}{n}(\tilde{R}^{2-1/p} - 1)$ and $\|X_i^2 - 1/n\|_2 \leq \frac{1}{n}(\tilde{R} - 1)$. Therefore using Lem. 39 and Lem. 40 we obtain:

$$\|\frac{n}{2}\mathbb{E}((Y_t)^2|\tilde{S}_n) - 1\|_{2p} \leq \min \left\{ \frac{\sqrt{2p-1}}{2\sqrt{n}} \left(\frac{\tilde{R}^2}{\tilde{R}^{1/p}} - 1 \right), \frac{1}{2\sqrt{n}} \left((\tilde{R}^2 - \tilde{R}^{1/p})\tilde{A}_{n,2p} + (\tilde{R} - 1)A_{2p} \right) \right\} \triangleq B_{n,2p}.$$

This implies that

$$\begin{aligned} & \int_{-\frac{1}{2}\log(1-\frac{D}{\sqrt{n}})}^{-\frac{1}{2}\log(1-\frac{R^2}{n\kappa})} \|\frac{n}{2}\mathbb{E}((Y_t)^2|\tilde{S}_n) - 1\|_{2p} \frac{e^{-2t}}{\sqrt{1-e^{-2t}}} dt \\ & \leq B_{n,2p} \int_{-\frac{1}{2}\log(1-\frac{R^2}{n\kappa})}^{T_n} \frac{e^{-2t}}{\sqrt{1-e^{-2t}}} dt = B_{n,2p} \left[\frac{\sqrt{D}}{n^{1/4}} - \frac{\tilde{R}}{\sqrt{n\kappa}} \right] \\ & = \frac{1}{2\sqrt{n}} \left[\frac{\sqrt{D}}{n^{1/4}} - \frac{\tilde{R}}{\sqrt{n\kappa}} \right] \min \left\{ \sqrt{2p-1} \left(\frac{\tilde{R}^2}{\tilde{R}^{1/p}} - 1 \right), (\tilde{R}^2 - \tilde{R}^{1/p})\tilde{A}_{n,2p} + (\tilde{R} - 1)A_{2p} \right\}. \end{aligned}$$

Similarly if $t \leq -\frac{1}{2}\log(1 - \frac{\tilde{R}^2}{n\kappa})$ then we have

$$\begin{aligned} \|\frac{n}{2}\mathbb{E}((Y_t)^2|\tilde{S}_n) - 1\|_{2p} & \leq 1 - \frac{n}{2}\mathbb{E}((Y_t)^2) + \|\frac{n}{2}\mathbb{E}((Y_t)^2|\tilde{S}_n) - \frac{n}{2}\mathbb{E}((Y_t)^2)\|_{2p} \\ & \leq 1 + \|\frac{n}{2}\mathbb{E}((Y_t)^2|\tilde{S}_n) - \frac{n}{2}\mathbb{E}((Y_t)^2)\|_{2p}. \end{aligned}$$

To control the size of this we remark that

$$\begin{aligned} \frac{n}{2}\mathbb{E}((Y_t)^2|\tilde{S}_n) & = \frac{1}{2}\mathbb{E}(X_1^2\mathbb{I}(|X_i| \leq c(t)) \sum_i \mathbb{I}(|X_i| \leq c(t)) \\ & \quad + \frac{1}{2}\mathbb{P}(|X_1| \leq c(t)) \sum_i X_i^2\mathbb{I}(|X_i| \leq c(t)) \\ & \quad - \mathbb{E}(X_1\mathbb{I}(|X_1| \leq c(t)) \sum_i X_i\mathbb{I}(|X_i| \leq c(t))). \end{aligned}$$

Therefore according to Lem. 40 we have

$$\begin{aligned} & \|\frac{n}{2}\mathbb{E}((Y_t)^2|\tilde{S}_n) - \frac{n}{2}\mathbb{E}((Y_t)^2)\|_{2p} \\ & \leq A_{2p} \left[\frac{1}{2\sqrt{n}} + \frac{\sqrt{n}}{2}\mu_{n,\kappa,1} + \sqrt{n}\mu_{n,\kappa,1}^2 \right] + \tilde{R}^{1/p}\tilde{A}_{n,2p} \left[\frac{1}{2\sqrt{n}} \right. \\ & \quad \left. + \min \left(\frac{c(t)^{2-\frac{1}{p}}\sqrt{n}^{1-\frac{1}{p}}}{2}, \frac{c(t)^2\sqrt{n}}{2} \right) + \min \left(c(t)^{1-\frac{1}{p}}n^{-1/2p}, \sqrt{n}c(t)^2 \right) \right]. \end{aligned}$$

Using also Lem. 39 this implies that

$$\begin{aligned} & \|\frac{n}{2}\mathbb{E}((Y_t)^2|\tilde{S}_n) - 1\|_{2p} \\ & \leq 1 + \left\{ A_{2p} \left[\frac{1}{2\sqrt{n}} + \frac{\sqrt{n}}{2}\mu_{n,\kappa,1} + \sqrt{n}\mu_{n,\kappa,1}^2 \right] + \tilde{R}^{1/p}\tilde{A}_{n,2p} \left[\frac{1}{2\sqrt{n}} \right. \right. \\ & \quad \left. \left. + \min \left(\frac{c(t)^{2-\frac{1}{p}}\sqrt{n}^{1-\frac{1}{p}}}{2}, \frac{c(t)^2\sqrt{n}}{2} \right) + \min \left(c(t)^{1-\frac{1}{p}}n^{-1/2p}, \sqrt{n}c(t)^2 \right) \right] \right. \\ & \quad \left. \sqrt{2p-1} \left[\frac{1}{2\sqrt{n}} + c(t)^2 \min \left(\frac{\sqrt{n}}{2c(t)^{\frac{1}{p}}n^{1/2p}}, \sqrt{n}/2 \right) + \min \left(c(t)^{1-\frac{1}{p}}n^{-1/2p}, \sqrt{n}c(t)^2 \right) \right] \right\}. \end{aligned}$$

Therefore to upper-bound $\int_0^{-\frac{1}{2} \log(1-\frac{\tilde{R}^2}{n\kappa})} \frac{e^{-2t}}{\sqrt{1-e^{-2t}}} \|\frac{n}{2} \mathbb{E}((Y_t)^2|S_n) - 1\|_{2p} dt$, we need remark that the following integrals are upper-bounded as:

$$\int_0^{-\frac{1}{2} \log(1-\frac{\tilde{R}^2}{n\kappa})} \frac{e^{-2t}}{\sqrt{1-e^{-2t}}} dt \leq \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^1 \frac{x}{\sqrt{1-x^2}} dx = \frac{\tilde{R}}{\sqrt{n\kappa}}.$$

Moreover we have

$$\int_{-\frac{1}{2} \log(1-\frac{\tilde{R}^2}{n\kappa})}^{-\frac{1}{2} \log(1-\frac{1}{n\kappa})} \frac{e^{-2t}}{\sqrt{1-e^{-2t}}} dt \leq \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} \frac{x}{\sqrt{1-x^2}} dx = \frac{(\tilde{R}-1)}{\sqrt{n\kappa}}.$$

Furthermore we observe that

$$\int_0^{-\frac{1}{2} \log(1-\frac{1}{n\kappa})} \sqrt{1-e^{-2t}} e^{-2t} dt \leq \frac{1}{2} \int_{1-\frac{1}{n\kappa}}^1 \sqrt{1-x} dx \leq \frac{1}{3(n\kappa)^{\frac{3}{2}}}.$$

In addition we have:

$$\begin{aligned} \int_{-\frac{1}{2} \log(1-\frac{\tilde{R}^2}{n\kappa})}^{-\frac{1}{2} \log(1-\frac{1}{n\kappa})} \sqrt{1-e^{-2t}}^{-1/p} e^{-2t} dt &\leq \frac{1}{2} \int_{1-\frac{1}{n\kappa}}^{1-\frac{\tilde{R}^2}{n\kappa}} \sqrt{1-x}^{-1/p} dx \\ &\leq \frac{2p}{2(2p-1)(n\kappa)^{-1/2p+1}} \left((\tilde{R}^2)^{1-1/2p} - 1 \right). \end{aligned}$$

And finally we also have

$$\begin{aligned} \int_{-\frac{1}{2} \log(1-\frac{\tilde{R}^2}{n\kappa})}^{-\frac{1}{2} \log(1-\frac{1}{n\kappa})} \sqrt{1-e^{-2t}}^{1-1/p} e^{-2t} dt &\leq \frac{1}{2} \int_{1-\frac{1}{n\kappa}}^{1-\frac{\tilde{R}^2}{n\kappa}} \sqrt{1-x}^{1-1/p} dx \\ &\leq \frac{2p}{(6p-2)(n\kappa)^{3/2-1/2p}} \left((\tilde{R}^2)^{3/2-1/2p} - 1 \right). \end{aligned}$$

and

$$\int_{-\frac{1}{2} \log(1-\frac{\tilde{R}^2}{n\kappa})}^{-\frac{1}{2} \log(1-\frac{1}{n\kappa})} e^{-2t} dt \leq \frac{1}{2n\kappa} [\tilde{R}^2 - 1].$$

Therefore we obtain that

$$\begin{aligned} &\int_0^{-\frac{1}{2} \log(1-\frac{\tilde{R}^2}{n\kappa})} \frac{e^{-2t}}{\sqrt{1-e^{-2t}}} \|\frac{n}{2} \mathbb{E}((Y_t)^2|S_n) - 1\|_{2p} dt \\ &\leq \min \left\{ \begin{aligned} &\frac{\tilde{R}}{\sqrt{n\kappa}} \left(1 + \frac{A_{2p} + \tilde{R}^{1/p} \tilde{A}_{n,2p}}{2\sqrt{n}} \right) + \frac{1}{n\sqrt{\kappa}} \left\{ \frac{(A_{2p} + \tilde{R}^{1/p} \tilde{A}_{n,2p})}{2} + \frac{p\tilde{R}^{1/p} \tilde{A}_{n,2p} ((\tilde{R}^2)^{1-\frac{1}{2p}} - 1)}{2(2p-1)} \right. \\ &\quad \left. + \frac{\tilde{R}^{1/p} \tilde{A}_{n,2p}}{2} \frac{2p}{(6p-2)} \left((\tilde{R}^2)^{3/2-1/2p} - 1 \right) + A_{2p}(\tilde{R} - 1) + \frac{A_{2p}}{4} [\tilde{R}^2 - 1] \right\} \\ &\frac{\tilde{R}}{\sqrt{n\kappa}} \left(1 + \frac{\sqrt{2p-1}}{2\sqrt{n}} \right) + \frac{1}{n\sqrt{\kappa}} \left\{ \frac{\sqrt{2p-1}}{2} \frac{1}{n\sqrt{\kappa}} + \sqrt{2p-1} \frac{2p}{2(2p-1)} \left((\tilde{R}^2)^{1-1/2p} - 1 \right) \right. \\ &\quad \left. + \frac{\sqrt{2p-1}}{2} \frac{2p}{(6p-2)} \left((\tilde{R}^2)^{3/2-1/2p} - 1 \right) \right\}. \end{aligned} \right. \end{aligned} \tag{21}$$

This directly implies

$$(a'_{2,p}) \leq \sqrt{2p-1} \min \left\{ \begin{aligned} & \frac{\tilde{R}}{\sqrt{n\kappa}} \left(1 + \frac{A_{2p} + \tilde{R}^{1/p} \tilde{A}_{n,2p}}{2\sqrt{n}} \right) + \frac{1}{n\sqrt{\kappa}} \left\{ \frac{(A_{2p} + \tilde{R}^{1/p} \tilde{A}_{n,2p})}{2} + A_{2p}(\tilde{R} - 1) \right. \\ & \quad \left. + \frac{p\tilde{R}^{1/p} \tilde{A}_{n,2p}}{2(2p-1)} \left((\tilde{R}^2)^{1-1/2p} - 1 \right) + \frac{\tilde{R}^{1/p} \tilde{A}_{n,2p}}{2} \frac{2p}{(6p-2)} \left((\tilde{R}^2)^{\frac{3}{2}-\frac{1}{2p}} - 1 \right) \right. \\ & \quad \left. \left. + \frac{A_{2p}}{4} [\tilde{R}^2 - 1] \right\} \right. \\ & \left. \frac{\tilde{R}}{\sqrt{n\kappa}} \left(1 + \frac{\sqrt{2p-1}}{2\sqrt{n}} \right) + \frac{1}{n\sqrt{\kappa}} \left\{ \frac{\sqrt{2p-1}}{2} \frac{1}{n\sqrt{\kappa}} + \sqrt{2p-1} \frac{2p \left((\tilde{R}^2)^{1-\frac{1}{2p}} - 1 \right)}{2(2p-1)} \right. \right. \\ & \quad \left. \left. + \frac{\sqrt{2p-1}}{2} \frac{2p}{(6p-2)} \left((\tilde{R}^2)^{\frac{3}{2}-\frac{1}{2p}} - 1 \right) \right\} \right. \\ & \left. + \frac{\sqrt{2p-1}}{2\sqrt{n}} \left[\frac{\sqrt{D}}{n^{1/4}} - \frac{\tilde{R}}{\sqrt{n\kappa}} \right] \min \left\{ \begin{aligned} & \sqrt{2p-1} \left(\frac{\tilde{R}^2}{\tilde{R}^{1/p}} - 1 \right) \\ & (\tilde{R}^2 - \tilde{R}^{1/p}) \tilde{A}_{n,2p} + (\tilde{R} - 1) A_{2p} \end{aligned} \right. \right. \end{aligned} \right. \quad (22)$$

Moreover to further upper bound $(a'_{2,p})$ we remark that :

$$\int_0^{-\frac{1}{2} \log(1-\frac{\tilde{R}^2}{n\kappa})} \sqrt{1-e^{-2t}} e^{-2t} dt \leq \frac{\tilde{R}^3}{3(n\kappa)^{\frac{3}{2}}} \quad \text{and} \quad \int_0^{-\frac{1}{2} \log(1-\frac{\tilde{R}^2}{n\kappa})} e^{-2t} dt \leq \frac{\tilde{R}^2}{2n\kappa}.$$

Therefore we obtain that $\int_0^{-\frac{1}{2} \log(1-\frac{\tilde{R}^2}{n\kappa})} \frac{e^{-2t}}{\sqrt{1-e^{-2t}}} \left\| \frac{n}{2} \mathbb{E}((Y_t)^2 | S_n) - 1 \right\|_{2p} dt$ can be further upper-bounded as

$$\begin{aligned} & \int_0^{-\frac{1}{2} \log(1-\frac{\tilde{R}^2}{n\kappa})} \frac{e^{-2t}}{\sqrt{1-e^{-2t}}} \left\| \frac{n}{2} \mathbb{E}((Y_t)^2 | S_n) - 1 \right\|_{2p} dt \\ & \leq \frac{\tilde{R}}{\sqrt{n\kappa}} \left(1 + \frac{\tilde{U}_{n,2p}}{2\sqrt{n}} \right) + \frac{\tilde{R}^2}{2n\sqrt{\kappa}} \left(\frac{3A_{2p}}{2} + \tilde{R}^{1+1/p} \tilde{A}_{n,2p} \right). \end{aligned}$$

Therefore this implies that $(a'_{2,p})$ can be further upper bounded as

$$\begin{aligned} (a'_{2,p}) & \leq \frac{\tilde{R}\sqrt{2p-1}}{\sqrt{n\kappa}} \left(1 + \frac{\tilde{U}_{n,2p}}{2\sqrt{n}} \right) + \frac{\tilde{R}^2}{2n\sqrt{\kappa}} \left(\frac{3A_{2p}}{2} + \tilde{R}^{1+1/p} \tilde{A}_{n,2p} \right) \\ & \quad + \frac{\sqrt{2p-1}}{2\sqrt{n}} \left(\frac{D}{n^{1/4}} - \frac{\tilde{R}}{\sqrt{n\kappa}} \right) \min \left\{ \begin{aligned} & \sqrt{2p-1} \left(\frac{\tilde{R}^2}{\tilde{R}^{1/p}} - 1 \right) \\ & (\tilde{R}^2 - \tilde{R}^{1/p}) \tilde{A}_{n,2p} + (\tilde{R} - 1) A_{2p} \end{aligned} \right. \\ & \sim \frac{\sqrt{2p-1} \tilde{R}}{\sqrt{n\kappa}}. \end{aligned}$$

Finally we work on bounding $(a'_{3,p})$. We observe that if we write $Z_i^{t,k} \triangleq \mathbb{E}((X_i - X'_i)^k \mathbb{I}(|X'_i| \leq c(t)) | X_{1:n}) \mathbb{I}(|X_i| \leq c(t))$ then

$$n\mathbb{E}(Y_t^k | X_{1:n}) = \sum_{i \leq n} Z_i^{t,k}.$$

Let $k \geq 3$ be an *odd* integer. We observe that the random variables $(Z_i^{t,k})$ are symmetric and therefore have a mean of zero. This implies that $n\mathbb{E}(Y_t^k | \tilde{S}_n) = O_p(\frac{1}{\sqrt{n}})$.

To make this clear, we shorthand $m_{k,2p}^*(t) \triangleq \min \left(n^{1-\frac{1}{p}} c(t)^{k-\frac{1}{p}}, nc(t)^k, \frac{1}{\sqrt{n}} c(t)^{k-3} \frac{\tilde{R}^3}{\tilde{R}^{1/p}} \right)$. We note that as $|X_1|^k \mathbb{I}(|X_1| \leq c(t)) \leq c(t)^{k-3} \min \left(c(t)^{3-\frac{1}{p}} |X_1|^{1/p}, |X_1|^3, c(t)^3 \right)$ then we have

$$\|Z_i^{t,k}\|_{2p} \leq 2^k \|X_1^k \mathbb{I}(|X_1| \leq c(t))\|_{2p} \leq \frac{2^k}{n} m_{k,2p}^*(t).$$

Similarly if we write $\tilde{m}_{k,2p}(t) \triangleq c(t)^{k-2} \min(\sqrt{n}c(t), nc(t)^2, \tilde{R})$ then we also have

$$\|Z_i^{t,k}\|_2 \leq \frac{2^k \tilde{m}_{k,2}(t)}{n}.$$

Therefore by using Lem. 39 and Lem. 40 we obtain that for all $t > 0$:

$$n\|\mathbb{E}(Y_t^k | \tilde{S}_n)\|_{2p} \leq \frac{2^k}{\sqrt{n}} \min \left\{ \sqrt{2p-1} m_{k,2p}^*(t), m_{k,2p}^*(t) \tilde{A}_{n,2p} \tilde{R}^{1/p} + \tilde{m}_{k,2}(t) A_{2p} \right\}.$$

This implies that if $k \geq 3$ is odd we have

$$\begin{aligned} & \int_0^{T_n} \frac{e^{-tk}}{\sqrt{1-e^{-2t^{k-1}}}} n\|\mathbb{E}((Y_t)^k | \tilde{S}_n)\|_{2p} dt \\ & \leq \sqrt{n} 2^k \tilde{R}^{1/p} \tilde{A}_{n,2p} n^{-1/2p} \int_{-\frac{1}{2} \log(1-\frac{\tilde{R}^2}{n\kappa})}^{-\frac{1}{2} \log(1-\frac{1}{n\kappa})} \frac{e^{-tk} c(t)^{k-\frac{1}{p}}}{\sqrt{1-e^{-2t^{k-1}}}} dt \\ & + \sqrt{n} 2^k (\tilde{R}^{1/p} \tilde{A}_{n,2p} + A_{2p}) \int_0^{-\frac{1}{2} \log(1-\frac{1}{n\kappa})} \frac{e^{-tk} c(t)^k}{\sqrt{1-e^{-2t^{k-1}}}} dt \\ & + \frac{2^k \tilde{R}^2 (\tilde{R} \tilde{A}_{n,2p} + A_{2p})}{n} \int_{-\frac{1}{2} \log(1-\frac{\tilde{R}^2}{n\kappa})}^{T_n} \frac{e^{-tk} c(t)^{k-3}}{\sqrt{1-e^{-2t^{k-1}}}} dt \\ & + 2^k A_{2p} \int_{-\frac{1}{2} \log(1-\frac{1}{n\kappa})}^{-\frac{1}{2} \log(1-\frac{\tilde{R}^2}{n\kappa})} \frac{e^{-tk} c(t)^{k-1}}{\sqrt{1-e^{-2t^{k-1}}}} dt. \end{aligned}$$

Using Eq. (45) we remark that for all $x \geq 1$ we have

$$\begin{aligned} & \sum_{\substack{k \geq 3 \\ k \text{ odd}}} \frac{\sqrt{2p-1}^{k-1} 2^k x^{k-1} \sqrt{\kappa}^k}{k \sqrt{(k-1)!}} \\ & \leq \sqrt{2e\kappa} \sum_{k \geq 1} \frac{(2p-1)^k 2^k x^{2k} \kappa^k \sqrt{k+1}}{(2k+1)} \frac{1}{k!} \\ & \leq \frac{2\sqrt{e\kappa}}{3} [e^{2(2p-1)x^2\kappa} - 1]. \end{aligned}$$

Therefore we have:

$$\begin{aligned} & \sum_{k \geq 3} \frac{\sqrt{2p-1}^{k-1}}{k \sqrt{(k-1)!}} \int_0^{T_n} \frac{e^{-tk}}{\sqrt{1-e^{-2t^{k-1}}}} n\|\mathbb{E}((Y_t)^k | S_n)\|_{2p} dt \\ & \leq \frac{2\sqrt{e}}{3} \sqrt{\kappa}^{1-\frac{1}{p}} \sqrt{n} \tilde{R}^{1/p} \tilde{A}_{n,2p} n^{-1/2p} \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} \sqrt{1-x^2}^{1-\frac{1}{p}} [e^{2(p-1)x^2\kappa} - 1] dx \\ & + \frac{2\sqrt{e}\sqrt{\kappa}}{3} \sqrt{n} (\tilde{R}^{1/p} \tilde{A}_{n,2p} + A_{2p}) \int_{\sqrt{1-\frac{1}{n\kappa}}}^1 \sqrt{1-x^2} [e^{2(2p-1)x^2\kappa} - 1] dx \\ & + \frac{2\sqrt{e}\tilde{R}^2 (\tilde{R} \tilde{A}_{n,2p} + A_{2p})}{3\kappa n} \int_{\sqrt{1-\frac{D}{\sqrt{n}}}}^{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}} \frac{[e^{2(2p-1)x^2\kappa} - 1]}{1-x^2} dx \\ & + \frac{2\sqrt{e}A_{2p}}{3} \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} [e^{2(2p-1)x^2\kappa} - 1] dx. \end{aligned}$$

Alternatively, we remark that

$$\begin{aligned}
& \sum_{k \geq 3} \frac{\sqrt{2p-1}^{k-1}}{k\sqrt{k-1}!} \int_0^{T_n} \frac{e^{-tk}}{\sqrt{1-e^{-2t}k-1}} n \|\mathbb{E}((Y_t)^k | S_n)\|_{2p} dt \\
& \leq \frac{2\sqrt{e}\sqrt{2p-1}\sqrt{n}^{1-\frac{1}{p}}}{3} \sqrt{\kappa}^{1-\frac{1}{p}} \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} \sqrt{1-x^2}^{1-\frac{1}{p}} [e^{2(2p-1)x^2\kappa} - 1] dx \\
& \quad + \frac{2\sqrt{e}\sqrt{2p-1}\sqrt{\kappa}}{3} \sqrt{n} \int_{\sqrt{1-\frac{1}{n\kappa}}}^1 \sqrt{1-x^2} [e^{2(2p-1)x^2\kappa} - 1] dx \\
& \quad + \frac{2\sqrt{e}\tilde{R}^3\sqrt{2p-1}}{3\kappa n \tilde{R}^{1/p}} \int_{\sqrt{1-\frac{D}{\sqrt{n}}}}^{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}} \frac{[e^{2(2p-1)x^2\kappa} - 1]}{1-x^2} dx.
\end{aligned}$$

We remark that this can be further upper bounded as

$$\begin{aligned}
& \sum_{k \geq 3} \frac{\sqrt{2p-1}^{k-1}}{k\sqrt{(k-1)!}} \int_0^{T_n} \frac{e^{-tk}}{\sqrt{1-e^{-2t}k-1}} n \|\mathbb{E}((Y_t)^k | S_n)\|_{2p} dt \\
& \leq \frac{2\sqrt{e}\sqrt{\kappa}}{3} \sqrt{n} (\tilde{R}^{1/p} \tilde{A}_{n,2p} + A_{2p}) \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^1 \sqrt{1-x^2} [e^{2(2p-1)x^2\kappa} - 1] dx \\
& \quad + \frac{2\sqrt{e}\tilde{R}^2(\tilde{R}\tilde{A}_{n,2p} + A_{2p})}{3\kappa n} \int_{\sqrt{1-\frac{D}{\sqrt{n}}}}^{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}} \frac{[e^{2(2p-1)x^2\kappa} - 1]}{1-x^2} dx \\
& \leq \frac{\sqrt{e}\tilde{R}^2 U_{n,2p} [e^{2(2p-1)M_{n,\kappa}^2} - 1]}{3\kappa n} \log \left(\frac{(M_{n,\kappa}+1)(1-\sqrt{1-\frac{D}{\sqrt{n}}})}{(1-M_{n,\kappa})(1+\sqrt{1-\frac{D}{\sqrt{n}}})} \right) \\
& \quad + \frac{2\sqrt{e}\tilde{R}^3 \tilde{U}_{n,2p}}{9n\kappa M_{n,\kappa}} [e^{2(2p-1)\kappa} - 1] \\
& \leq \frac{\sqrt{e}\tilde{R}^2}{3n\kappa} [e^{2(2p-1)\kappa} - 1] \left[\frac{2\tilde{R}\tilde{U}_{n,2p}}{3M_{n,\kappa}} + U_{n,2p} \log \left(\frac{(M_{n,\kappa}+1)(1-\sqrt{1-\frac{D}{\sqrt{n}}})}{(1-M_{n,\kappa})(1+\sqrt{1-\frac{D}{\sqrt{n}}})} \right) \right].
\end{aligned}$$

If $k \geq 4$ and is even then for all $t \geq 0$ such that $c(t) \leq \tilde{R}/\sqrt{n}$ then we have

$$\begin{aligned}
& n \|\mathbb{E}(Y_t^k | \tilde{S}_n)\|_{2p} \\
& \leq \|\sum_{i \leq n} \mathbb{E}[\mathbb{I}(|X_i|, |X'_i| \leq c(t))(X_i - X'_i)^k | \tilde{S}_n]\|_{2p} \\
& \leq \|\sum_{i \leq n} \mathbb{E}[\mathbb{I}(|X_i|, |X'_i| \leq c(t))(X_i - X'_i)^k | \tilde{S}_n] - \mathbb{E}[\mathbb{I}(|X_i|, |X'_i| \leq c(t))(X_i - X'_i)^k]\|_{2p} \\
& \quad + n \|\mathbb{E}[\mathbb{I}(|X_1|, |X'_1| \leq c(t))(X_1 - X'_1)^k]\| \\
& \stackrel{(a)}{\leq} 2^k \sqrt{2p-1} \min(\sqrt{n}^{1-\frac{1}{p}} c(t)^{k-\frac{1}{p}}, \sqrt{n} c(t)^k) + 2^k \min(c(t)^{k-2}, n c(t)^2).
\end{aligned}$$

where to get (a) we used Lem. 39 and the fact that as $k \geq 4$ is even we have

$$\begin{aligned}
& \|\mathbb{E}[\mathbb{I}(|X_i|, |X'_i| \leq c(t))(X_i - X'_i)^k] - \mathbb{E}[\mathbb{I}(|X_i|, |X'_i| \leq c(t))(X_i - X'_i)^k]\|_{2p} \\
& \leq \|\mathbb{E}[\mathbb{I}(|X_i|, |X'_i| \leq c(t))(X_i - X'_i)^k]\|_{2p} \leq 2^k \|\mathbb{E}[\mathbb{I}(|X_i| \leq c(t))X_i^k]\|_{2p}.
\end{aligned}$$

Moreover for all $t \geq -\frac{1}{2} \log(1 - \frac{\tilde{R}^2}{n\kappa})$ according to Lem. 39 we also have:

$$\begin{aligned}
n \|\mathbb{E}(Y_t^k | \tilde{S}_n)\|_{2p} &\leq \|\sum_{i \leq n} \mathbb{E}[\mathbb{I}(|X_i|, |X'_i| \leq c(t))(X_i - X'_i)^k | \tilde{S}_n]\|_{2p} \\
&\leq \|\sum_{i \leq n} \mathbb{E}[\mathbb{I}(|X_i|, |X'_i| \leq c(t))(X_i - X'_i)^k | \tilde{S}_n] \\
&\quad - \mathbb{E}[\mathbb{I}(|X_i|, |X'_i| \leq c(t))(X_i - X'_i)^k]\|_{2p} \\
&\quad + n |\mathbb{E}[\mathbb{I}(|X_1|, |X'_1| \leq c(t))(X_1 - X'_1)^k]| \\
&\leq \frac{\sqrt{2p-1}2^k}{\sqrt{n}^3} c(t)^{k-4} \tilde{R}^{4-1/p} + 2^k \frac{\tilde{R}^2}{n} c(t)^{k-4}.
\end{aligned}$$

This directly implies that we obtain

$$\begin{aligned}
&\int_0^{T_n} \frac{e^{-tk}}{\sqrt{1-e^{-2t}k-1}} n \|\mathbb{E}((Y_t)^k | \tilde{S}_n)\|_{2p} dt \\
&\leq \frac{2^k \sqrt{\kappa}^{k-4} \tilde{R}^2 (\sqrt{2p-1} \tilde{R}^{2-1/p} + \sqrt{n})}{\sqrt{n}^3} \int_{-\frac{1}{2} \log(1 - \frac{\tilde{R}^2}{n\kappa})}^{T_n} \frac{e^{-tk}}{\sqrt{1-e^{-2t}3}} dt \\
&\quad + 2^k \sqrt{2p-1} \sqrt{\kappa}^{k-\frac{1}{p}} \sqrt{n}^{1-1/p} \int_{-\frac{1}{2} \log(1 - \frac{\tilde{R}^2}{n\kappa})}^{-\frac{1}{2} \log(1 - \frac{1}{n\kappa})} e^{-tk} \sqrt{1-e^{-2t}1-\frac{1}{p}} dt \\
&\quad + 2^k \sqrt{\kappa}^k \sqrt{n} (\sqrt{n} + \sqrt{2p-1}) \int_0^{-\frac{1}{2} \log(1 - \frac{1}{n\kappa})} e^{-tk} \sqrt{1-e^{-2t}} dt \\
&\quad + 2^k \sqrt{\kappa}^{k-2} \int_{-\frac{1}{2} \log(1 - \frac{\tilde{R}^2}{n\kappa})}^{-\frac{1}{2} \log(1 - \frac{1}{n\kappa})} \frac{e^{-tk}}{\sqrt{1-e^{-2t}}} dt.
\end{aligned}$$

We remark that

$$\begin{aligned}
&\sum_{\substack{k \geq 4 \\ k \text{ even}}} \frac{\sqrt{2p-1}^{k-1} 2^k x^{k-1} \sqrt{\kappa}^k}{k \sqrt{(k-1)!}} \\
&\leq \sqrt{2e\kappa} \sqrt{2p-1} x \sum_{k \geq 1} \frac{(2p-1)^k 2^k x^{2k} \kappa^k}{\sqrt{k+1} \sqrt{2k+1}} \frac{1}{k!} \\
&\leq \frac{2\sqrt{e\kappa} \sqrt{2p-1} x}{\sqrt{3}} \left[e^{2(2p-1)x^2\kappa} - 1 - 2(2p-1)x^2\kappa \right].
\end{aligned}$$

Therefore we obtain that:

$$\begin{aligned}
& \sum_{\substack{k \geq 4 \\ \text{even}}} \frac{\sqrt{2p-1}^{k-1}}{k\sqrt{(k-1)!}} \int_0^{T_n} \frac{e^{-tk}}{\sqrt{1-e^{-2t}k-1}} n \|\mathbb{E}((Y_t)^k | \tilde{S}_n)\|_{2p} dt \\
& \leq \frac{2\sqrt{e}\sqrt{2p-1}}{\sqrt{3}} \left\{ \frac{\tilde{R}^2(\sqrt{2p-1}\tilde{R}^{2-1/p} + \sqrt{n})}{\sqrt{n^3}\kappa} \right. \\
& \quad \times \int_{\sqrt{1-\frac{D}{\sqrt{n}}}}^{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}} x \left[\frac{e^{2(2p-1)x^2\kappa} - 1 - 2(2p-1)x^2\kappa}{\sqrt{1-x^2}^3} \right] dx \\
& \quad + \sqrt{2p-1}\sqrt{\kappa}^{2-\frac{1}{p}}\sqrt{n}^{1-1/p} \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} \sqrt{1-x^2}^{1-\frac{1}{p}} x \left[e^{2(2p-1)x^2\kappa} - 1 - 2(2p-1)x^2\kappa \right] dx \\
& \quad + \sqrt{n}\kappa(\sqrt{n} + \sqrt{2p-1}) \int_{\sqrt{1-\frac{1}{n\kappa}}}^1 x \left[e^{2(2p-1)x^2\kappa} - 1 - 2(2p-1)x^2\kappa \right] \sqrt{1-x^2} dx \\
& \quad + \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} x \left[\frac{e^{2(2p-1)x^2\kappa} - 1 - 2(2p-1)x^2\kappa}{\sqrt{1-x^2}} \right] dx \\
& \leq \frac{\sqrt{e}\sqrt{2p-1}}{\sqrt{3}\kappa} \frac{\tilde{R}^2(\sqrt{2p-1}\tilde{R}^{2-1/p} + \sqrt{n})}{\sqrt{n^3}} \int_{\sqrt{1-\frac{D}{\sqrt{n}}}}^{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}} \left[\frac{e^{2(2p-1)x\kappa} - 1 - 2(2p-1)x\kappa}{\sqrt{1-x}^3} \right] dx \\
& \quad + \frac{\sqrt{e}}{\sqrt{3}} (2p-1) \sqrt{\kappa}^{2-\frac{1}{p}} \sqrt{n}^{1-1/p} \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} \sqrt{1-x}^{1-\frac{1}{p}} \left[e^{2(2p-1)x\kappa} - 1 - 2(2p-1)x\kappa \right] dx \\
& \quad + \frac{\sqrt{e}\kappa\sqrt{2p-1}}{\sqrt{3}} \sqrt{n}(\sqrt{n} + \sqrt{2p-1}) \int_{\sqrt{1-\frac{1}{n\kappa}}}^1 \left[e^{2(2p-1)x\kappa} - 1 - 2(2p-1)x\kappa \right] \sqrt{1-x} dx \\
& \quad + \frac{\sqrt{e}\sqrt{2p-1}}{\sqrt{3}} \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} \left[\frac{e^{2(2p-1)x\kappa} - 1 - 2(2p-1)x\kappa}{\sqrt{1-x}} \right] dx.
\end{aligned}$$

Similarly using Lem. 40 we have

$$\begin{aligned}
& n \|\mathbb{E}(Y_t^k | \tilde{S}_n)\|_{2p} \\
& \leq \begin{cases} \tilde{A}_{n,2p} \tilde{R}^{1/p} 2^k \min(\sqrt{n}^{1-\frac{1}{p}} c(t)^{k-\frac{1}{p}}, \sqrt{n} c(t)^k) + 2^k \min(c(t)^{k-1}, \sqrt{n} c(t)^k) A_{2p} \\ \quad + 2^k \min(c(t)^{k-2}, n c(t)^2). \\ \tilde{A}_{n,2p} \frac{2^k}{\sqrt{n}^3} c(t)^{k-4} \tilde{R}^4 + 2^k \frac{A_{2p}}{\sqrt{n}^3} c(t)^{k-4} \tilde{R}^3 + 2^k c(t)^{k-4} \tilde{R}^2 / n. \end{cases}
\end{aligned}$$

This directly implies that we have:

$$\begin{aligned}
& \sum_{\substack{k \geq 4 \\ \text{even}}} \frac{\sqrt{2p-1}^{k-1}}{k\sqrt{(k-1)!}} \int_0^{T_n} \frac{e^{-tk}}{\sqrt{1-e^{-2t}k-1}} n \|\mathbb{E}((Y_t)^k | \tilde{S}_n)\|_{2p} dt \\
& \leq \frac{\sqrt{e\kappa}\sqrt{2p-1}}{\sqrt{3}} \sqrt{n}(\tilde{U}_{n,2p} + \sqrt{n}) \int_{\sqrt{1-\frac{1}{n\kappa}}}^1 \left[e^{2(2p-1)x\kappa} - 1 - 2(2p-1)x\kappa \right] \sqrt{1-x} dx \\
& + \frac{\sqrt{e\kappa}\sqrt{2p-1}}{\sqrt{3}} A_{2p} \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} \left[e^{2(2p-1)x\kappa} - 1 - 2(2p-1)x\kappa \right] dx \\
& + \frac{\sqrt{e\kappa}\sqrt{2p-1}}{\sqrt{3}} \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} \frac{\left[e^{2(2p-1)x\kappa} - 1 - 2(2p-1)x\kappa \right]}{\sqrt{1-x}} dx \\
& + \frac{\sqrt{e(2p-1)\kappa}\tilde{A}_{n,2p}\tilde{R}^{1/p}}{\sqrt{3}} \sqrt{\kappa n}^{\frac{p-1}{p}} \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} \left[e^{2(2p-1)x\kappa} - 1 - 2(2p-1)x\kappa \right] \sqrt{1-x}^{1-\frac{1}{p}} dx \\
& + \frac{\sqrt{e\kappa}\sqrt{2p-1}}{\sqrt{3\kappa}} \frac{\tilde{R}^2(\sqrt{n} + \tilde{R}U_{n,2p})}{\sqrt{n}^3} \int_{\sqrt{1-\frac{D}{\sqrt{n}}}}^{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}} \frac{\left[e^{2(2p-1)x\kappa} - 1 - 2(2p-1)x\kappa \right]}{\sqrt{1-x}^3} dx.
\end{aligned}$$

This can be further simplified as

$$\begin{aligned}
& \sum_{\substack{k \geq 4 \\ \text{even}}} \frac{\sqrt{p-1}^{k-1}}{k\sqrt{(k-1)!}} \int_0^{T_n} \frac{e^{-tk}}{\sqrt{1-e^{-2t}k-1}} n \|\mathbb{E}((Y_t)^k | S_n)\|_{2p} dt \\
& \leq \frac{\sqrt{e\kappa}\sqrt{2p-1}}{\sqrt{3}} \sqrt{n}(A_{2p} + \tilde{A}_{n,2p}\tilde{R}^{1/p}) \int_{1-\frac{\tilde{R}^2}{n\kappa}}^1 \left[e^{2(2p-1)x\kappa} - 1 - 2(2p-1)x\kappa \right] \sqrt{1-x} dx \\
& + \frac{\sqrt{e\kappa}\sqrt{2p-1}}{\sqrt{3}} n \int_{1-\frac{1}{n\kappa}}^1 \left[e^{2(2p-1)x\kappa} - 1 - 2(2p-1)x\kappa \right] \sqrt{1-x} dx \\
& + \frac{\sqrt{e\kappa}\sqrt{2p-1}}{\sqrt{3\kappa}} \frac{\tilde{R}^2(\sqrt{n} + \tilde{R}A_{2p} + \tilde{R}^2\tilde{A}_{n,2p})}{\sqrt{n}^3} \int_{1-\frac{D}{\sqrt{n}}}^{1-\frac{\tilde{R}^2}{n\kappa}} \frac{\left[e^{2(2p-1)x\kappa} - 1 - 2(2p-1)x\kappa \right]}{\sqrt{1-x}^3} dx \\
& + \frac{\sqrt{e\kappa}\sqrt{2p-1}}{\sqrt{3}} \int_{1-\frac{\tilde{R}^2}{n\kappa}}^{1-\frac{1}{n\kappa}} \frac{\left[e^{2(2p-1)x\kappa} - 1 - 2(2p-1)x\kappa \right]}{\sqrt{1-x}} dx \\
& \leq \frac{2\sqrt{e\kappa}\sqrt{2p-1}}{3\sqrt{n\kappa}\sqrt{3}} \left[e^{2(2p-1)\kappa} - 1 - 2(2p-1)\kappa \right] \left[\frac{\tilde{R}^3}{\sqrt{n}} \tilde{U}_{n,2p} + 1 \right] \\
& + \frac{2\sqrt{e\kappa}\sqrt{2p-1}}{\sqrt{3\kappa}} \frac{\tilde{R}^2(\sqrt{n} + \tilde{R}U_{n,2p})}{\sqrt{n}} \left[\frac{\sqrt{n\kappa}}{\tilde{R}} - \frac{n^{1/4}}{\sqrt{D}} \right] \left[e^{2(2p-1)M_{n,\kappa}^2\kappa} - 1 - 2(2p-1)M_{n,\kappa}^2\kappa \right] \\
& + \frac{2\sqrt{e\kappa}\sqrt{2p-1}}{\sqrt{3n\kappa}} \left[e^{2(2p-1)\tilde{M}_{n,\kappa}^2\kappa} - 1 - 2(2p-1)\tilde{M}_{n,\kappa}^2\kappa \right] \left[\tilde{R} - 1 \right] \\
& \sim \frac{2\sqrt{e\kappa}\sqrt{2p-1}\tilde{R}}{\sqrt{3n\kappa}} \left[e^{2(2p-1)\kappa} - 1 - 2(2p-1)\kappa \right] \left[2 - \frac{2}{3\tilde{R}} \right].
\end{aligned}$$

This directly implies that

$$(a'_{3,p}) \triangleq (a'_{3,1,p}) + (a'_{3,2,p}). \quad (23)$$

where we set

$$(a'_{3,1,p}) \triangleq \min \left\{ \begin{aligned} & \frac{2\sqrt{e}}{3} \sqrt{\kappa}^{1-\frac{1}{p}} \sqrt{n} \tilde{R}^{1/p} \tilde{A}_{n,2p} n^{-1/2p} \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} \sqrt{1-x^2}^{1-\frac{1}{p}} [e^{2(p-1)x^2\kappa} - 1] dx \\ & + \frac{2\sqrt{e}\sqrt{\kappa}}{3} \sqrt{n} (\tilde{R}^{1/p} \tilde{A}_{n,2p} + A_{2p}) \int_{\sqrt{1-\frac{1}{n\kappa}}}^1 \sqrt{1-x^2} [e^{2(2p-1)x^2\kappa} - 1] dx \\ & + \frac{2\sqrt{e}\tilde{R}^2 (\tilde{R}\tilde{A}_{n,2p} + A_{2p})}{3\kappa n} \int_{\sqrt{1-\frac{D}{\sqrt{n}}}}^{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}} \frac{[e^{2(2p-1)x^2\kappa} - 1]}{1-x^2} dx \\ & + \frac{2\sqrt{e}A_{2p}}{3} \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} [e^{2(2p-1)x^2\kappa} - 1] dx \\ & \frac{2\sqrt{e}\sqrt{2p-1}\sqrt{n}^{1-\frac{1}{p}} \sqrt{\kappa}^{1-\frac{1}{p}}}{3} \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} \sqrt{1-x^2}^{1-\frac{1}{p}} [e^{2(2p-1)x^2\kappa} - 1] dx \\ & + \frac{2\sqrt{e}\sqrt{2p-1}\sqrt{\kappa}}{3} \sqrt{n} \int_{\sqrt{1-\frac{1}{n\kappa}}}^1 \sqrt{1-x^2} [e^{2(2p-1)x^2\kappa} - 1] dx \\ & + \frac{2\sqrt{e}\tilde{R}^3 \sqrt{2p-1}}{3\kappa n \tilde{R}^{1/p}} \int_{\sqrt{1-\frac{D}{\sqrt{n}}}}^{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}} \frac{[e^{2(2p-1)x^2\kappa} - 1]}{1-x^2} dx \end{aligned} \right.$$

and

$$(a'_{3,2,p}) \triangleq \min \left\{ \begin{aligned} & \frac{\sqrt{e\kappa}\sqrt{2p-1}}{\sqrt{3}} \sqrt{n} (\tilde{U}_{n,2p} + \sqrt{n}) \int_{\sqrt{1-\frac{1}{n\kappa}}}^1 [e^{2(2p-1)x\kappa} - 1 - 2(2p-1)x\kappa] \sqrt{1-x} dx \\ & + \frac{\sqrt{e}\sqrt{\kappa}\sqrt{2p-1}}{\sqrt{3}} A_{2p} \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} [e^{2(2p-1)x\kappa} - 1 - 2(2p-1)x\kappa] dx \\ & + \frac{\sqrt{e}\sqrt{2p-1}}{\sqrt{3}} \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} \frac{[e^{2(2p-1)x\kappa} - 1 - 2(2p-1)x\kappa]}{\sqrt{1-x}} dx \\ & + \frac{\sqrt{e}\sqrt{2p-1} \tilde{A}_{n,2p} \tilde{R}^{1/p} \sqrt{\kappa}^{2-\frac{1}{p}}}{\sqrt{3}} \sqrt{n}^{\frac{p-1}{p}} \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} [e^{2(2p-1)x\kappa} - 1 - 2(2p-1)x\kappa] \\ & \quad \times \sqrt{1-x}^{1-\frac{1}{p}} dx \\ & + \frac{\sqrt{e}\sqrt{2p-1}}{\sqrt{3\kappa}} \frac{\tilde{R}^2 (\sqrt{n} + \tilde{R}U_{n,2p})}{\sqrt{n}^3} \int_{\sqrt{1-\frac{D}{\sqrt{n}}}}^{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}} \frac{[e^{2(2p-1)x\kappa} - 1 - 2(2p-1)x\kappa]}{\sqrt{1-x^3}} dx \\ & \frac{\sqrt{e}\sqrt{2p-1}}{\sqrt{3\kappa}} \frac{\tilde{R}^2 (\sqrt{2p-1}\tilde{R}^{2-1/p} + \sqrt{n})}{\sqrt{n}^3} \int_{\sqrt{1-\frac{D}{\sqrt{n}}}}^{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}} \frac{[e^{2(2p-1)x\kappa} - 1 - 2(2p-1)x\kappa]}{\sqrt{1-x^3}} dx \\ & + \frac{\sqrt{e\kappa}}{(2p-1)} \sqrt{3} \sqrt{n\kappa}^{1-1/p} \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} \sqrt{1-x^2}^{1-\frac{1}{p}} [e^{2(2p-1)x\kappa} - 1 - 2(2p-1)x\kappa] dx \\ & + \frac{\sqrt{e\kappa}\sqrt{2p-1}}{\sqrt{3}} (\sqrt{n} + \sqrt{2p-1}) \int_{\sqrt{1-\frac{1}{n\kappa}}}^1 [e^{2(2p-1)x\kappa} - 1 - 2(2p-1)x\kappa] \sqrt{1-x} dx \\ & + \frac{\sqrt{e}\sqrt{2p-1}}{\sqrt{3}} \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} \frac{[e^{2(2p-1)x\kappa} - 1 - 2(2p-1)x\kappa]}{\sqrt{1-x}} dx \end{aligned} \right.$$

We remark that this can be further upper bounded as

$$\begin{aligned}
(a'_{3,p}) &\leq \frac{\sqrt{e}\tilde{R}^2}{3n\kappa} [e^{2(2p-1)\kappa} - 1] \left[\frac{2\tilde{R}\tilde{U}_{n,2p}}{3M_{n,\kappa}} + U_{n,2p} \log \left(\frac{(M_{n,\kappa}+1)(1-\sqrt{1-\frac{D}{\sqrt{n}}})}{(1-M_{n,\kappa})(1+\sqrt{1-\frac{D}{\sqrt{n}}})} \right) \right] \\
&\quad + \frac{2\sqrt{e}\sqrt{2p-1}\tilde{R}}{\sqrt{3n\kappa}} \left[e^{2(2p-1)\kappa} - 1 - 2(2p-1)\kappa \right] \left[\frac{\tilde{R}^2}{3\sqrt{n}}\tilde{U}_{n,2p} + 2 + \frac{\tilde{R}U_{n,2p}}{\sqrt{n}} - \frac{2}{3\tilde{R}} \right] \\
&\sim \frac{2\sqrt{e}\sqrt{2p-1}\tilde{R}}{\sqrt{3n\kappa}} \left[e^{2(2p-1)\kappa} - 1 - 2(2p-1)\kappa \right] \left[2 - \frac{2}{3\tilde{R}} \right].
\end{aligned}$$

■

C.2. Combining the results together

Theorem 22 Suppose that the conditions of Thm. 4. The upper bound found for $(a'_{1,p})$, $(a'_{2,p})$ and $(a'_{3,p})$ found in Eq. (20), Eq. (22) and Eq. (23) are valid. Moreover, we have

$$\begin{aligned}
\sqrt{2}H(\tilde{S}_n, Z) &\leq \inf_{\substack{D>0 \\ \kappa \geq \frac{\tilde{R}^2}{\sqrt{n}D}}} \left\{ \frac{\sqrt{3e}\tilde{R}^2(\sqrt{n}+\tilde{R})(e^{2\kappa(M_D^n)^2}-1)}{\kappa C\sqrt{2n}} + \frac{\sqrt{e}\tilde{R}^2 M_D^n (e^{2\kappa(M_D^n)^2}-1)}{\sqrt{3}D\sqrt{n\kappa}} \right. \\
&\quad + \frac{\log(n/D^2)}{4\sqrt{2}\sqrt{n}}(\tilde{R}-1) - \frac{1}{2}\log(1-\frac{D}{\sqrt{n}}) \\
&\quad \left. + \inf_{p \geq 1} C_p \left((a'_{1,p}) + (a'_{2,p}) + (a'_{3,p}) \right) \right\}.
\end{aligned}$$

Note that a looser but more concise upper bound for $H(\tilde{S}_n, Z)$ can be established.

Theorem 23 Suppose that the conditions of Thm. 4, hold then the upper bound in Thm. 22 is asymptotically equivalent to the following upper bound

$$\begin{aligned}
\sqrt{2}H(\tilde{S}_n, Z) &\lesssim \inf_{\substack{D>0 \\ \kappa \geq \frac{\tilde{R}^2}{\sqrt{n}D}}} \left\{ \frac{\sqrt{e}\tilde{R}^2(e^{2\kappa}-1)}{\sqrt{3}D\sqrt{n\kappa}} + \frac{\log(n/D^2)}{4\sqrt{2}\sqrt{n}}(\tilde{R}-1) - \frac{1}{2}\log(1-\frac{D}{\sqrt{n}}) \right. \\
&\quad + \inf_{p \geq 1} C_p \left\{ \frac{\sqrt{2p-1}\tilde{R}}{\sqrt{n\kappa}} + \frac{2\sqrt{e}\sqrt{2p-1}\tilde{R}}{\sqrt{3n\kappa}} \left[e^{2(2p-1)\kappa} - 1 - 2(2p-1)\kappa \right] \left[2 - \frac{2}{3\tilde{R}} \right] \right. \\
&\quad \left. \left. + \frac{\tilde{R}^2}{2n\kappa M_{n,\kappa}} \left[U_{n,2p} + \tilde{U}_{n,2p} \right] \right\} \right\}.
\end{aligned}$$

Moreover they are constants K_1 , K_2 and K_3 given in Eq. (24) such that

$$\begin{aligned}
\sqrt{2}H(\sigma^{-1}S_n, Z) &\lesssim \frac{K_1}{D\sqrt{n}} + \frac{\log(n/D^2)}{4\sqrt{2}\sqrt{n}}(\tilde{R}-1) + \frac{D}{2\sqrt{n}} \\
&\quad + \|S_{Y_1}(\sigma Y_1)\|_2 \left\{ \frac{K_2\tilde{R}}{\sqrt{n}} + \frac{K_3\tilde{R}}{\sqrt{n}} \left[2 - \frac{2}{3\tilde{R}} \right] \right\} = O\left(\frac{\log(n)}{\sqrt{n}}\right).
\end{aligned}$$

Proof Using Thm. 22 we obtain that for all $\kappa \geq \frac{\tilde{R}^2}{\sqrt{n}D}$

$$\begin{aligned}
& \sqrt{2}H(\tilde{S}_n, Z) \\
& \lesssim \frac{\sqrt{e}\tilde{R}^2(e^{2\kappa}-1)}{\sqrt{3}D\sqrt{n}\kappa} + \frac{\log(n/D^2)}{4\sqrt{2}\sqrt{n}}(\tilde{R}-1) - \frac{1}{2}\log(1 - \frac{D}{\sqrt{n}}) \\
& + \inf_{p \geq 1} C_p \left\{ \frac{\sqrt{2p-1}\tilde{R}}{\sqrt{n}\kappa} + \frac{2\sqrt{e}\sqrt{2p-1}\tilde{R}}{\sqrt{3n}\kappa} \left[e^{2(2p-1)\kappa} - 1 - 2(2p-1)\kappa \right] \left[2 - \frac{2}{3\tilde{R}} \right] \right. \\
& \left. + \frac{\tilde{R}^2}{2n\kappa M_{n,\kappa}} \left[U_{n,2p} + \tilde{U}_{n,2p} \right] \right\} \\
& \sim \frac{\sqrt{e}\tilde{R}^2(e^{2\kappa}-1)}{\sqrt{3}D\sqrt{n}\kappa} + \frac{\log(n/D^2)}{4\sqrt{2}\sqrt{n}}(\tilde{R}-1) \\
& + \inf_{p \geq 1} C_p \left\{ \frac{\sqrt{2p-1}\tilde{R}}{\sqrt{n}\kappa} + \frac{2\sqrt{e}\sqrt{2p-1}\tilde{R}}{\sqrt{3n}\kappa} \left[e^{2(2p-1)\kappa} - 1 - 2(2p-1)\kappa \right] \left[2 - \frac{2}{3\tilde{R}} \right] \right\}.
\end{aligned}$$

Write $C \triangleq \frac{3^{1/4}}{2\sqrt{2}\sqrt{3}e^{1/4}} \in [0.16, 0.17]$ and choose $\kappa = \max\left(\frac{3^{1/4}}{2(2p-1)\sqrt{2}\sqrt{3}e^{1/4}}, \frac{\tilde{R}^2}{\sqrt{n}D}\right)$ then we have

$$\begin{aligned}
\sqrt{2}H(\tilde{S}_n, Z) & \sim \frac{\sqrt{e}(2p-1)\tilde{R}^2(e^{\frac{C}{2p-1}}-1)}{\sqrt{3}D\sqrt{n}C} + \frac{\log(n/D^2)}{4\sqrt{2}\sqrt{n}}(\tilde{R}-1) + \frac{D}{2\sqrt{n}} \\
& + C_p \left\{ \frac{(2p-1)\tilde{R}}{\sqrt{n}C} + \frac{2\sqrt{e}(2p-1)\tilde{R}}{\sqrt{3n}C} \left[e^{2C} - 1 - 2C \right] \left[2 - \frac{2}{3\tilde{R}} \right] \right\}.
\end{aligned}$$

If we choose $p = 2$ then if we write

$$K_1 \triangleq 1.105, \quad K_2 \triangleq 35.01, \quad K_3 \triangleq 1.36, \quad (24)$$

we obtain that

$$\begin{aligned}
\sqrt{2}H(\sigma^{-1}S_n, Z) & \lesssim \frac{K_1}{D\sqrt{n}} + \frac{\log(n/D^2)}{4\sqrt{2}\sqrt{n}}(\tilde{R}-1) + \frac{D}{2\sqrt{n}} + \|S_{Y_1}(\sigma Y_1)\|_2 \left\{ \frac{K_2\tilde{R}}{\sqrt{n}} + \frac{K_3}{\sqrt{n}} \right\} \\
& = O\left(\frac{\log(n)}{\sqrt{n}}\right).
\end{aligned}$$

■

Appendix D. Proof of Thm. 5: Bounding Hellinger distance to a Gaussian

Lemma 24 *Under the conditions of Thm. 5, the following holds*

$$H(S, Z) \leq \frac{1}{\sqrt{2}} \int_0^\infty \|I_t^S(X_t) - (X_t^S)^2 + 1 - X_t^S(S_{X_t}^S - X_t^S)\|_2 dt.$$

Proof Let $\varphi(\cdot)$ be the PDF of a standard normal random variable. We write $h_t^S(x) \triangleq \frac{f_t^S(x)}{\varphi(x)}$. Fix an arbitrary $T < \infty$. By definition we have

$$H(S, X_T^S)^2 = 1 - \int_{\mathbb{R}} \sqrt{f_S(x)} \sqrt{f_T^S(x)} dx.$$

We remark that for all $x \in \mathbb{R}$ the function $t \rightarrow f_t^S(x)$ is differentiable. Therefore we have $\int_0^T \partial_t \sqrt{f_t^S(x)} dt = \sqrt{f_T^S(x)} - \sqrt{f_S(x)}$. Therefore this directly implies that

$$H(S, X_T^S)^2 = - \int_0^T \int_{\mathbb{R}} \sqrt{f_S(x)} \partial_t \sqrt{f_t^S(x)} dx dt.$$

We will re-express this in terms of information theoretic quantities. Indeed, by the chain rule and the heat equation we obtain that

$$\begin{aligned}\partial_t \sqrt{f_t^S(x)} &= \frac{1}{2\sqrt{f_t^S(x)}} \partial_t f_t^S(x) = \frac{\varphi(x)}{2\sqrt{f_t^S(x)}} \partial_t \frac{f_t^S(x)}{\varphi(x)} \\ &= \frac{\varphi(x)}{2\sqrt{f_t^S(x)}} \partial_t h_t^S(x) = \frac{\varphi(x)}{2\sqrt{f_t^S(x)}} [h_t^{S''}(S)(x) - x h_t^{S'}(x)] = \frac{f_t^S(x)}{2\sqrt{f_t^S(x)}} \left[\frac{h_t^{S''}(x)}{h_t^S(x)} - x \frac{h_t^{S'}(x)}{h_t^S(x)} \right].\end{aligned}\quad (25)$$

We therefore obtain that

$$\begin{aligned}H(S, X_T^S)^2 &= - \int_0^T \int_{\mathbb{R}} \sqrt{f_S(x)} \partial_t \sqrt{f_t^S(x)} dx dt \\ &\stackrel{(a)}{=} - \int_0^T \mathbb{E} \left(\frac{\sqrt{f_S(X_t^S)}}{2\sqrt{f_t^S(X_t^S)}} \left[\frac{h_t^{S''}(X_t^S)}{h_t^S(X_t^S)} - X_t^S \frac{h_t^{S'}(X_t^S)}{h_t^S(X_t^S)} \right] \right) dt \\ &\stackrel{(b)}{=} - \frac{1}{2} \int_0^T \mathbb{E} \left(\left[\frac{\sqrt{f_S(X_t^S)}}{\sqrt{f_t^S(X_t^S)}} - 1 \right] \left[\frac{h_t^{S''}(X_t^S)}{h_t^S(X_t^S)} - X_t^S \frac{h_t^{S'}(X_t^S)}{h_t^S(X_t^S)} \right] \right) dt \\ &\stackrel{(c)}{\leq} \frac{1}{2} \sup_{t \leq T} \|1 - \frac{\sqrt{f_S(X_t^S)}}{\sqrt{f_t^S(X_t^S)}}\|_2 \int_0^T \|I_t^S(X_t^S) - (X_t^S)^2 + 1 - X_t^S(S_{X_t^S}^S - X_t^S)\|_2 dt \\ &\leq \sup_{t < \infty} \frac{1}{\sqrt{2}} H(S, X_t^S) \int_0^T \|I_t^S(X_t^S) - (X_t^S)^2 + 1 - X_t^S(S_{X_t^S}^S - X_t^S)\|_2 dt.\end{aligned}$$

where (a) is a consequence of Eq. (25) and (b) is a consequence of the fact that

$$\mathbb{E} \left(\frac{h_t^{S''}(X_t^S)}{h_t^S(X_t^S)} - X_t^S \frac{h_t^{S'}(X_t^S)}{h_t^S(X_t^S)} \right) = 0,$$

and finally were to obtain (c) we used the fact that by the chain rule we know that:

$$\frac{h_t^{S'}(X_t^S)}{h_t^S(X_t^S)} = \frac{f_t^{S'}(X_t^S)}{f_t^S(X_t^S)} + X_t^S = -S_{X_t^S}(X_t^S) + X_t^S,$$

and

$$\begin{aligned}\frac{h_t^{S''}(X_t^S)}{h_t^S(X_t^S)} &= \frac{f_t^{S''}(X_t^S)}{f_t^S(X_t^S)} - (X_t^S)^2 + 1 + 2X_t^S \frac{h_t^{S'}(X_t^S)}{h_t^S(X_t^S)} \\ &= I_t(X_t^S) - (X_t^S)^2 + 1 + 2X_t^S(-S_{X_t^S}(X_t^S) + X_t^S) \\ &= I_t(X_t^S) + (X_t^S)^2 + 1 - 2X_t^S S_{X_t^S}(X_t^S).\end{aligned}$$

Therefore as this holds for all $T < \infty$ we obtain that

$$\sup_{t < \infty} H(S, X_t^S) \leq \frac{1}{\sqrt{2}} \int_0^\infty \|I_t^S(X_t^S) - (X_t^S)^2 + 1 - X_t^S(S_{X_t^S}^S - X_t^S)\|_2 dt.$$

This notably implies that

$$H(S, Z) \leq \frac{1}{\sqrt{2}} \int_0^\infty \|I_t^S(X_t^S) - (X_t^S)^2 + 1 - X_t^S(S_{X_t^S}^S - X_t^S)\|_2 dt.$$

■

Proof [Proof of Thm. 5] For all measurable function h , we write $P_t(h) : x \rightarrow \mathbb{E}(h(e^{-t}x + \sqrt{1 - e^{-2t}}Z))$.

By the linearity of the integration we know that

$$\begin{aligned}
& \int_0^\infty \|I_t^S(X_t^S) - (X_t^S)^2 + 1 - X_t^S(S_{X_t^S}^S - X_t^S)\|_2 dt \\
& \leq \int_0^{-\frac{1}{2} \log(1 + \frac{D}{\sqrt{n}})} \|I_t^S(X_t^S) - (X_t^S)^2 + 1 - X_t^S(S_{X_t^S}^S - X_t^S)\|_2 dt \\
& + \int_{-\frac{1}{2} \log(1 + \frac{D}{\sqrt{n}})}^\infty \|I_t^S(X_t^S) - (X_t^S)^2 + 1 - X_t^S(S_{X_t^S}^S - X_t^S)\|_2 dt \\
& \triangleq (b_1) + (b_2).
\end{aligned}$$

We bound each term successively. Firstly we remark that thanks to Lem. 37 we have $\mathbb{E}(\tau_t'^S | X_t^S) = \mathbb{E}(\tau_t^S | X_t^S) = 0$. Moreover, we remark that $S_t^S(X_t^S)$ is the unique X_t^S -measurable random variable such that for all $\phi \in \mathbb{C}^\infty(\mathbb{R})$ we have

$$\mathbb{E}(S_t^S(X_t^S)\phi(X_t^S)) = \mathbb{E}(\phi'(X_t^S)).$$

Moreover as Z is independent from S we remark that for all $\phi \in \mathbb{C}^\infty(\mathbb{R})$ we have

$$\mathbb{E}(\mathbb{E}(S_S(S) | X_t^S)\phi(X_t^S)) = \mathbb{E}(S_S(S)\phi(e^{-t}S + \sqrt{1 - e^{-2t}}Z)) = e^{-t}\mathbb{E}(\phi'(X_t^S)).$$

Therefore this implies that

$$S_t^S(X_t^S) = e^t \mathbb{E}(S_S(S) | X_t^S).$$

Moreover we remark that $I_t^S(X_t^S)$ is the unique X_t^S -measurable random variable such that for all $\phi \in \mathbb{C}^\infty(\mathbb{R})$ we have

$$\mathbb{E}(I_t^S(X_t^S)\phi(X_t^S)) = \mathbb{E}(\phi''(X_t^S)).$$

Therefore by the Stein identity we obtain that:

$$I_t^S(X_t^S) = \frac{e^t}{\sqrt{1 - e^{-2t}}} \mathbb{E}(S_S(S)Z | X_t^S).$$

This directly implies that

$$\begin{aligned}
I_t^S(X_t^S) - X_t^S S_t(X_t^S) &= e^t \mathbb{E}\left(S_S(S) \left[\frac{Z}{\sqrt{1 - e^{-2t}}} - e^{-t}S - \sqrt{1 - e^{-2t}}Z \right] | X_t^S\right) \\
&= e^t \mathbb{E}\left(S_S(S) \left[\frac{Ze^{-2t}}{\sqrt{1 - e^{-2t}}} - e^{-t}S \right] | X_t^S\right) \\
&= \mathbb{E}\left(S_S(S) \left[\frac{Ze^{-2t}}{\sqrt{1 - e^{-2t}}} - e^{-t}S \right] | X_t^S\right).
\end{aligned}$$

Therefore as $\mathbb{E}(\tau_t^S | X_t^S) = 0$, then by Hölder's inequality, we obtain that for all $p, p' \geq 1$

$$\begin{aligned}
& \|I_t^S(X_t^S) - (X_t^S)^2 + 1 - X_t^S(S_{X_t^S}^S - X_t^S)\|_2 \\
& \leq \|S_S(S)\|_{2\frac{p}{p-1}} \left\| \frac{Ze^{-2t}}{\sqrt{1 - e^{-2t}}} - e^{-t}S + \tau_t^S \right\|_{2p} + 1.
\end{aligned}$$

Therefore for all $p \geq 1$ we have

$$\begin{aligned}
(b_1) &\leq \int_0^{-\frac{1}{2} \log(1 - \frac{D}{\sqrt{n}})} \|I_t^S(X_t^S) - (X_t^S)^2 + 1 - X_t^S(S_{X_t^S}^S - X_t^S)\|_2 dt \\
&\leq \int_0^{-\frac{1}{2} \log(1 - \frac{D}{\sqrt{n}})} \|S_S(S)\|_{2\frac{p}{p-1}} \left\| \frac{Ze^{-2t}}{\sqrt{1 - e^{-2t}}} - e^{-t}S + \tau_t^S \right\|_{2p} dt - \frac{1}{2} \log(1 - \frac{D}{\sqrt{n}}).
\end{aligned}$$

Now that we have bounded (b_1) , we bound (b_2) . By the Stein identity we remark that

$$I_t^S(X_t^S) = \mathbb{E}\left(\frac{1}{1-e^{-2t}}(Z^2 - 1)|X_t^S\right).$$

This directly implies that

$$\begin{aligned} & I_t^S(X_t^S) - (X_t^S)^2 + 1 \\ &= \mathbb{E}\left(\frac{Z^2}{1-e^{-2t}} - (X_t^S)^2 | X_t^S\right) - \frac{e^{-2t}}{1-e^{-2t}} \\ &= \mathbb{E}\left(\frac{Z}{\sqrt{1-e^{-2t}}}\left(\frac{Z}{\sqrt{1-e^{-2t}}} - X_t^S\right) + X_t^S\left(\frac{Z}{\sqrt{1-e^{-2t}}} - X_t^S\right) | X_t^S\right) - \frac{e^{-2t}}{1-e^{-2t}} \\ &= \mathbb{E}\left(\frac{Z}{\sqrt{1-e^{-2t}}}\left(\frac{e^{-2t}}{\sqrt{1-e^{-2t}}}Z - e^{-t}S\right) | X_t^S\right) + X_t^S \mathbb{E}\left[\left(\frac{e^{-2t}}{\sqrt{1-e^{-2t}}}Z - e^{-t}S\right) | X_t^S\right] - \frac{e^{-2t}}{1-e^{-2t}}. \end{aligned}$$

Moreover we note that

$$\begin{aligned} & \mathbb{E}\left(\frac{Z}{\sqrt{1-e^{-2t}}}\left(\frac{e^{-2t}}{\sqrt{1-e^{-2t}}}Z - e^{-t}S\right) | X_t^S\right) - \frac{e^{-2t}}{1-e^{-2t}} \\ &= \mathbb{E}\left(\frac{e^{-2t}}{1-e^{-2t}}(Z^2 - 1) - \frac{e^{-t}}{\sqrt{1-e^{-2t}}}SZ | X_t^S\right). \end{aligned}$$

Moreover we also know by the Stein identity that

$$S_t^S(X_t^S) - X_t^S = \mathbb{E}\left[\left(\frac{e^{-2t}}{\sqrt{1-e^{-2t}}}Z - e^{-t}S\right) | X_t^S\right].$$

This directly implies that:

$$\begin{aligned} & I_t^S(X_t^S) - (X_t^S)^2 + 1 \\ &= \mathbb{E}\left(\frac{e^{-2t}}{1-e^{-2t}}(Z^2 - 1) - \frac{e^{-t}}{\sqrt{1-e^{-2t}}}SZ | X_t^S\right) + X_t^S(S_t^S(X_t^S) - X_t^S). \end{aligned}$$

Therefore we have

$$\begin{aligned} & I_t^S(X_t^S) - (X_t^S)^2 + 1 - X_t^S(S_t^S(X_t^S) - X_t^S) \\ &= \mathbb{E}\left(\frac{e^{-2t}}{1-e^{-2t}}(Z^2 - 1) - \frac{e^{-t}}{\sqrt{1-e^{-2t}}}SZ | X_t^S\right). \end{aligned}$$

As $\mathbb{E}(\tau_t'^X | X_t^S) = 0$, then by Jensen inequality this directly implies that

$$\begin{aligned} (b_2) &\leq \int_{-\frac{1}{2}\log(1-\frac{D}{\sqrt{n}})}^{\infty} \|I_t^S(X_t^S) - (X_t^S)^2 + 1 - X_t^S(S_t^S(X_t^S) - X_t^S)\|_2 dt \\ &\leq \int_{-\frac{1}{2}\log(1-\frac{D}{\sqrt{n}})}^{\infty} \|\mathbb{E}\left(\frac{e^{-2t}}{1-e^{-2t}}(Z^2 - 1) - \frac{e^{-t}}{\sqrt{1-e^{-2t}}}SZ | X_t^S\right)\|_2 dt \\ &\leq \int_{-\frac{1}{2}\log(1-\frac{D}{\sqrt{n}})}^{\infty} \|\frac{e^{-2t}}{1-e^{-2t}}(Z^2 - 1) - \frac{e^{-t}}{\sqrt{1-e^{-2t}}}SZ + \tau_t'^S\|_2 dt. \end{aligned}$$

■

Appendix E. Proof of Thms. 6, 7, and 8: Efficient Concentration with Stein Kernels

In this section we write $T_{\tilde{S}_n}(\cdot)$ the Stein kernel of $\tilde{S}_n \triangleq \sigma^{-1}S_n$.

E.1. Proof of Thm. 6: Efficient Stein kernel tail bound

In the following we are going to take $\hat{Q}_n^d(x)$ to be such that for all $x > 0$

$$\sup_{t \geq 0} \mathbb{P}(|e^{-t} \tilde{S}_n + \sqrt{1 - e^{-2t}} Z| \geq x) \leq \hat{Q}_n^d(x).$$

Lemma 25 *Let $p' \geq 2$. Suppose that the random variables (Y_i) satisfy Assump. (R, σ) and Assump. $(\tau_{\tilde{Y}_1}, p')$. Choose $\alpha > 0$ and let $p \geq 1$ and $q \geq 2$. Define $u_{d,\alpha}(x) \triangleq \frac{1}{\alpha^2} \int_x^{x+\alpha} \int_t^{t+\alpha} \mathbb{I}(|z| \geq u) dz dt$ and let $Z \sim N(0, 1)$ then we have*

$$\begin{aligned} \mathbb{P}(|S_n| \geq \sigma u) &\leq \mathbb{E}(u_{d,\alpha}(Z)) + \frac{D_p(\tau_{\tilde{Y}_1})}{\sqrt{n}} \\ &\times \min \left\{ \|Z\|_p \left[\|u'_{d,\alpha}(Z)\|_{\frac{p}{p-1}} + \frac{\pi \mathcal{W}_{\frac{pq}{(p-1)}}(G, \tilde{S}_n) \hat{Q}_n^d(u-2\alpha)^{\frac{(p-1)(q-1)}{pq}}}{4\alpha^2} \right] \right. \\ &\quad \left. \frac{1}{2\alpha^2} \hat{Q}_n^d(u-2\alpha)^{\frac{(p-1)}{p}} \right\}. \end{aligned}$$

As the proof is very similar to the proof of Lem. 26 we present only the proof of Lem. 26 in detail and highlight the differences with the proof of Lem. 25. A similar one-sided inequality can also be presented

Lemma 26 *If the conditions of Thm. 6 hold then the following upper bound is valid:*

$$\mathbb{P}(S_n \geq \sigma u) \leq \mathbb{E}(u_\alpha(Z)) + \frac{D_p(\tau_{\tilde{Y}_1})}{\sqrt{n}} \min \left\{ \|Z\|_p \left[\frac{\pi \mathcal{W}_{\frac{pq}{(p-1)}}(G, \tilde{S}_n) \hat{Q}_n(u-2\alpha)^{\frac{(p-1)(q-1)}{pq}}}{4\alpha^2} + \|u'_\alpha(Z)\|_{\frac{p}{p-1}} \right] \right. \\ \left. \frac{1}{2\alpha^2} \hat{Q}_n(u-2\alpha)^{\frac{(p-1)}{p}} \right\}.$$

Proof As the proof of Lem. 25 is very similar to the proof of Lem. 26 we present only the proof of Lem. 26 in detail and highlight the differences with the proof of Lem. 25.

Fix $\alpha > 0$ to be a constant. It will be useful to note that $u_\alpha \in C^2(\mathbb{R})$. Moreover we remark that $u_\alpha(x) \neq \mathbb{I}(x \geq u)$ and $u'_\alpha(x), u''_\alpha(x) \neq 0$ if and only if $x \in [u - 2\alpha, u]$. For all x we also have $|u'_\alpha(x)| \leq \frac{1}{\alpha}$ and $|u''_\alpha(x)| \leq \frac{1}{\alpha^2}$.

Moreover we denote

$$T_{\tilde{S}_n}(\tilde{S}_n) \triangleq \frac{1}{n} \mathbb{E}(\sum_{i=1}^n \tau_{\tilde{Y}_1}(\sigma^{-1} Y_i) | \tilde{S}_n).$$

We remark that $T_{\tilde{S}_n}$ is a Stein Kernel for \tilde{S}_n .

Let $Z \sim N(0, 1)$ be an independent standard normal random variable. Define $P_t u_\alpha(x) \triangleq \mathbb{E}_Z(u_\alpha(e^{-t}x + \sqrt{1 - e^{-2t}}Z))$. We remark that $P_0 u_\alpha(x) = u_\alpha(x)$ and $P_\infty u_\alpha(x) = \mathbb{E}(u_\alpha(Z))$. Therefore we have:

$$\begin{aligned} |\mathbb{E}(u_\alpha(\tilde{S}_n)) - \mathbb{E}(u_\alpha(Z))| &= |\int_0^\infty \partial_t \mathbb{E}(P_t u_\alpha(\tilde{S}_n)) dt| \\ &\stackrel{(a)}{\leq} |\int_0^\infty \mathbb{E}(\Delta P_t u_\alpha(\tilde{S}_n) - \tilde{S}_n \nabla P_t u_\alpha(\tilde{S}_n)) dt| \\ &\stackrel{(b)}{\leq} |\int_0^\infty \mathbb{E}(\Delta P_t u_\alpha(\tilde{S}_n)(1 - T_{\tilde{S}_n}(\tilde{S}_n))) dt|, \end{aligned}$$

where to obtain (a) we used the heat equation satisfied by the Ornstein-Uhlenbeck process and to obtain (b) we used the definition of Stein Kernels. Moreover, by the Stein identity, we note that we note that

$$\begin{aligned}\Delta P_t u_\alpha(x) &= e^{-2t} \mathbb{E}_Z(u''_\alpha(e^{-t}x + \sqrt{1-e^{-2t}}Z)) \\ &= \frac{e^{-2t}}{\sqrt{1-e^{-2t}}} \mathbb{E}_Z(Z u'_\alpha(e^{-t}x + \sqrt{1-e^{-2t}}Z)).\end{aligned}$$

Therefore for any arbitrary $p \geq 1$ we have

$$\begin{aligned}& |\mathbb{E}(u_\alpha(\tilde{S}_n)) - \mathbb{E}(u_\alpha(Z))| \\ & \leq \left| \int_0^\infty \frac{e^{-2t}}{\sqrt{1-e^{-2t}}} \mathbb{E}\left(Z u'_\alpha(e^{-t}\tilde{S}_n + \sqrt{1-e^{-2t}}Z)(1 - T_{\tilde{S}_n}(\tilde{S}_n))\right) dt \right| \\ & \leq \int_0^\infty \frac{e^{-2t}}{\sqrt{1-e^{-2t}}} \mathbb{E}\left(|Z(1 - T_{\tilde{S}_n}(\tilde{S}_n))| |u'_\alpha(e^{-t}\tilde{S}_n + \sqrt{1-e^{-2t}}Z)|\right) dt \\ & \stackrel{(a)}{\leq} \|Z\|_p \|1 - T_{\tilde{S}_n}(\tilde{S}_n)\|_p \int_0^\infty \frac{e^{-2t}}{\sqrt{1-e^{-2t}}} \|u'_\alpha(e^{-t}\tilde{S}_n + \sqrt{1-e^{-2t}}Z)\|_{\frac{p}{p-1}} dt.\end{aligned}$$

where (a) is a consequence of Holder inequality.

We can prove very similarly that

$$\begin{aligned}& |\mathbb{E}(u_{d,\alpha}(\tilde{S}_n)) - \mathbb{E}(u_{d,\alpha}(Z))| \\ & \leq \|Z\|_p \|1 - T_{\tilde{S}_n}(\tilde{S}_n)\|_p \int_0^\infty \frac{e^{-2t}}{\sqrt{1-e^{-2t}}} \|u'_{d,\alpha}(e^{-t}\tilde{S}_n + \sqrt{1-e^{-2t}}Z)\|_{\frac{p}{p-1}} dt.\end{aligned}$$

Now let $q \geq 2$ and let $G \sim N(0, 1)$ be an independent copy of Z that is such that $\|\tilde{S}_n - G\|_q = W_q(\tilde{S}_n, N(0, 1))$ (which exists) then we have

$$\begin{aligned}& \|u'_\alpha(e^{-t}\tilde{S}_n + \sqrt{1-e^{-2t}}Z)\|_{\frac{p}{p-1}} \\ & \stackrel{(a)}{\leq} \left\| \left[u'_\alpha(e^{-t}\tilde{S}_n + \sqrt{1-e^{-2t}}Z) - u'_\alpha(e^{-t}G + \sqrt{1-e^{-2t}}Z) \right] \mathbb{I}(X_t \in [u - 2\alpha, u]) \right\|_{\frac{p}{p-1}} \\ & + \|u'_\alpha(Z)\|_{\frac{p}{p-1}} \\ & \stackrel{(b)}{\leq} \|u'_\alpha(Z)\|_{\frac{p}{p-1}} + \frac{\mathcal{W}_{\frac{pq}{(p-1)}}(G, \tilde{S}_n)}{\alpha^2} e^{-t} \mathbb{P}(X_t \in [u - 2\alpha, u])^{\frac{(p-1)(q-1)}{pq}}.\end{aligned}$$

where (a) comes from the fact that $u'_\alpha(x) = 0$ for all $x \in [u - 2\alpha, u]$ and where (b) is a consequence of Hölder inequality and of the Lipchitzity of u'_α . Moreover we note that:

$$\mathbb{P}(X_t \in [u - 2\alpha, u]) \leq \mathbb{P}(X_t \geq u - 2\alpha) \leq \hat{Q}_n(u - 2\alpha).$$

This directly implies that

$$\begin{aligned}& \int_0^\infty \frac{e^{-2t}}{\sqrt{1-e^{-2t}}} \|u'_\alpha(e^{-t}\tilde{S}_n + \sqrt{1-e^{-2t}}Z)\|_{\frac{p}{p-1}} dt \\ & \leq \|u'_\alpha(Z)\|_{\frac{p}{p-1}} + \frac{\pi}{4} \frac{\mathcal{W}_{\frac{pq}{(p-1)}}(G, \tilde{S}_n) \hat{Q}_n(u - 2\alpha)^{\frac{(p-1)(q-1)}{pq}}}{\alpha^2}.\end{aligned}$$

Very similarly we can prove that

$$\begin{aligned}
& \|u'_{d,\alpha}(e^{-t}\tilde{S}_n + \sqrt{1 - e^{-2t}}Z)\|_{\frac{p}{p-1}} \\
& \leq \|u'_{d,\alpha}(Z)\|_{\frac{p}{p-1}} + \frac{\mathcal{W}_{\frac{pq}{(p-1)}}(G, \tilde{S}_n)}{\alpha^2} e^{-t} \mathbb{P}(|X_t| \in [u - 2\alpha, u])^{\frac{(p-1)(q-1)}{pq}} \\
& \leq \|u'_{d,\alpha}(Z)\|_{\frac{p}{p-1}} + \frac{\mathcal{W}_{\frac{pq}{(p-1)}}(G, \tilde{S}_n)}{\alpha^2} e^{-t} \hat{Q}_n^d(u - 2\alpha)^{\frac{(p-1)(q-1)}{pq}}.
\end{aligned}$$

Now we clearly notice that $\mathbb{P}(\tilde{S}_n \geq u) \leq \mathbb{E}(u_\alpha(\tilde{S}_n))$. Therefore we have

$$\begin{aligned}
& \mathbb{P}(S_n \geq \sigma u) \\
& \leq \|Z\|_p \|1 - T_{\tilde{S}_n}(\tilde{S}_n)\|_p \left[\|u'_\alpha(Z)\|_{\frac{p}{p-1}} + \frac{\pi \mathcal{W}_{\frac{pq}{(p-1)}}(G, \tilde{S}_n) \hat{Q}_n(u - 2\alpha)^{\frac{(p-1)(q-1)}{pq}}}{4\alpha^2} \right] \\
& \quad + \mathbb{E}(u_\alpha(Z)).
\end{aligned}$$

Moreover as we have

$$T_{\tilde{S}_n}(\tilde{S}_n) = \frac{1}{n} \mathbb{E}\left(\sum_{i=1}^n \tau_{\tilde{Y}_1}(\sigma^{-1}Y_i) | \tilde{S}_n\right).$$

Therefore by Lem. 39 and Lem. 40 this directly implies that

$$\begin{aligned}
\|1 - T_{\tilde{S}_n}(\tilde{S}_n)\|_p & \leq \left\| \frac{1}{n} \sum_{i \leq n} \mathbb{E}(\tau_{\tilde{Y}_1}(\sigma^{-1}Y_i) | \tilde{S}_n) - 1 \right\|_p \\
& \leq \frac{1}{\sqrt{n}} \min \left\{ A_p \|\tau_{\tilde{Y}_1}(\sigma^{-1}Y_1) - 1\|_2 + A_{n,p}^* \|\tau_{\tilde{Y}_1}(\sigma^{-1}Y_1) - 1\|_p \right. \\
& \quad \left. \sqrt{p-1} \|\tau_{\tilde{Y}_1}(\sigma^{-1}Y_1) - 1\|_p \right\}.
\end{aligned}$$

This directly implies

$$\begin{aligned}
& \mathbb{P}(S_n \geq \sigma u) \\
& \leq \mathbb{E}(u_\alpha(Z)) + \frac{\|Z\|_p}{\sqrt{n}} \min \left\{ A_p \|\tau_{\tilde{Y}_1}(\sigma^{-1}Y_1) - 1\|_2 + A_{n,p}^* \|\tau_{\tilde{Y}_1}(\sigma^{-1}Y_1) - 1\|_p \right. \\
& \quad \left. \sqrt{p-1} \|\tau_{\tilde{Y}_1}(\sigma^{-1}Y_1) - 1\|_p \right. \\
& \quad \left. \times \left[\|u'_\alpha(Z)\|_{\frac{p}{p-1}} + \frac{\pi \mathcal{W}_{\frac{pq}{(p-1)}}(G, \tilde{S}_n) \hat{Q}_n(u - 2\alpha)^{\frac{(p-1)(q-1)}{pq}}}{4\alpha^2} \right] \right\}.
\end{aligned}$$

This proves the first part of the minimum

To prove the second part, we note that $\Delta P_t u_\alpha(\cdot)$ can alternatively be bounded as

$$\begin{aligned}
|\Delta P_t u_\alpha(x)| & = e^{-2t} |\mathbb{E}_Z(u''_\alpha(e^{-t}x + \sqrt{1 - e^{-2t}}Z))| \\
& = e^{-2t} |\mathbb{E}_Z(u''_\alpha(e^{-t}x + \sqrt{1 - e^{-2t}}Z))| \\
& \leq \frac{e^{-2t}}{\alpha^2} \mathbb{E}_Z(I(e^{-t}x + \sqrt{1 - e^{-2t}}Z \in [u - 2\alpha, u])) \\
& \leq \frac{e^{-2t}}{\alpha^2} \mathbb{P}(e^{-t}x + \sqrt{1 - e^{-2t}}Z \in [u - 2\alpha, u]).
\end{aligned}$$

Therefore we obtain that

$$\begin{aligned}
& |\mathbb{E}(u_\alpha(\tilde{S}_n)) - \mathbb{E}(u_\alpha(Z))| \\
& \leq \left| \int_0^\infty e^{-2t} \mathbb{E} \left(u_\alpha''(e^{-t}\tilde{S}_n + \sqrt{1-e^{-2t}}Z)(1 - T_{\tilde{S}_n}(\tilde{S}_n)) \right) dt \right| \\
& \leq \|1 - T_{\tilde{S}_n}(\tilde{S}_n)\|_p \int_0^\infty e^{-2t} \|u_\alpha''(e^{-t}\tilde{S}_n + \sqrt{1-e^{-2t}}Z)\|_{\frac{p}{p-1}} dt \\
& \leq \frac{1}{\alpha^2} \|1 - T_{\tilde{S}_n}(\tilde{S}_n)\|_p \int_0^\infty e^{-2t} \|\mathbb{P}(e^{-t}\tilde{S}_n + \sqrt{1-e^{-2t}}Z \in [u-2\alpha, u]) | \tilde{S}_n)\|_{\frac{p}{p-1}} dt \\
& \leq \frac{\hat{Q}_n(u-2\alpha)^{\frac{p-1}{p}}}{\alpha^2} \|1 - T_{\tilde{S}_n}(\tilde{S}_n)\|_p \int_0^\infty e^{-2t} dt \\
& \leq \frac{D_p(\tau_{\tilde{Y}_1})}{2\sqrt{n}\alpha^2} \hat{Q}_n(u-2\alpha)^{\frac{p-1}{p}}.
\end{aligned}$$

As $\mathbb{E}(u_\alpha(S_n)) \geq \mathbb{P}(\tilde{S}_n \geq u)$ we obtain the desired result.

Following the same roadmap we obtain that

$$\begin{aligned}
|\mathbb{E}(u_{d,\alpha}(\tilde{S}_n)) - \mathbb{E}(u_{d,\alpha}(Z))| & \leq \frac{\hat{Q}_n^d(u-2\alpha)^{\frac{p-1}{p}}}{\alpha^2} \|1 - T_{\tilde{S}_n}(\tilde{S}_n)\|_p \int_0^\infty e^{-2t} dt \\
& \leq \frac{D_p(\tau_{\tilde{Y}_1})}{2\sqrt{n}\alpha^2} \hat{Q}_n^d(u-2\alpha)^{\frac{p-1}{p}}.
\end{aligned}$$

■

Lemma 27 *If the conditions of Thm. 6 hold then the following bound also holds:*

$$\begin{aligned}
\mathbb{P}(S_n \geq \sigma u) & \leq \mathbb{E}(u_\alpha(Z)) + \frac{\|Z\|_p D_p(\tau_{\tilde{Y}_1})}{\sqrt{n}} \\
& \left[\|u'_\alpha(Z)\|_{\frac{p}{p-1}} + \frac{\pi}{4\alpha^2} \mathcal{W}_{\frac{pq}{p-1}}(\tilde{S}_n, G) \mathbb{P}(Z \in [u-2\alpha-\rho, u+\rho])^{\frac{(q-1)(p-1)}{pq}} \right. \\
& \left. + \frac{\sqrt{\pi} \Gamma(\frac{q'(p-1)+2p}{2p})}{2\alpha^2 \Gamma(\frac{q'(p-1)+3p}{2p})} \frac{\mathcal{W}_{\frac{pq'}{p-1}}(\tilde{S}_n, Z)^{q'}}{\rho^{\frac{q'(p-1)}{p}}} \right].
\end{aligned}$$

Proof Fix $\alpha > 0$ to be a constant. It will be useful to note that $u_\alpha \in C^2(\mathbb{R})$. Moreover we remark that $u_\alpha(x) \neq \mathbb{I}(x \geq u)$ and $u'_\alpha(x), u''_\alpha(x) \neq 0$ if and only if $x \in [u-2\alpha, u]$. For all x we also have $|u'_\alpha(x)| \leq \frac{1}{\alpha}$ and $|u''_\alpha(x)| \leq \frac{1}{\alpha^2}$.

Let $Z \sim N(0, 1)$ be an independent standard normal random variable. Define $P_t u_\alpha(x) \triangleq \mathbb{E}_Z(u_\alpha(e^{-t}x + \sqrt{1-e^{-2t}}Z))$. We remark that $P_0 u_\alpha(x) = u_\alpha(x)$ and $P_\infty u_\alpha(x) = \mathbb{E}(u_\alpha(Z))$. Therefore we have:

$$\begin{aligned}
|\mathbb{E}(u_\alpha(\tilde{S}_n)) - \mathbb{E}(u_\alpha(Z))| & = \left| \int_0^\infty \partial_t \mathbb{E}(P_t u_\alpha(\tilde{S}_n)) dt \right| \\
& \stackrel{(a)}{\leq} \left| \int_0^\infty \mathbb{E}(\Delta P_t u_\alpha(\tilde{S}_n) - \tilde{S}_n \nabla P_t u_\alpha(\tilde{S}_n)) dt \right| \\
& \leq \left| \int_0^\infty \mathbb{E}(\Delta u_\alpha(\tilde{S}_n)(1 - T_{\tilde{S}_n}(\tilde{S}_n))) dt \right|;
\end{aligned}$$

where to obtain (a) we used the heat equation satisfied by the Ornstein-Uhlenbeck process. Moreover, by the Stein identity, we note that we note that

$$\begin{aligned}
\Delta P_t u_\alpha(x) & = e^{-2t} \mathbb{E}_Z(u_\alpha''(e^{-t}x + \sqrt{1-e^{-2t}}Z)) \\
& = \frac{e^{-2t}}{\sqrt{1-e^{-2t}}} \mathbb{E}_Z(Z u'_\alpha(e^{-t}x + \sqrt{1-e^{-2t}}Z)).
\end{aligned}$$

Therefore for any arbitrary $p \geq 1$ we have

$$\begin{aligned}
& |\mathbb{E}(u_\alpha(\tilde{S}_n)) - \mathbb{E}(u_\alpha(Z))| \\
& \leq \left| \int_0^\infty \frac{e^{-2t}}{\sqrt{1-e^{-2t}}} \mathbb{E} \left(Z u'_\alpha(e^{-t}\tilde{S}_n + \sqrt{1-e^{-2t}}Z)(1 - T_{\tilde{S}_n}(\tilde{S}_n)) \right) dt \right| \\
& \leq \int_0^\infty \frac{e^{-2t}}{\sqrt{1-e^{-2t}}} \mathbb{E} \left(|Z(1 - T_{\tilde{S}_n}(\tilde{S}_n))| |u'_\alpha(e^{-t}\tilde{S}_n + \sqrt{1-e^{-2t}}Z)| \right) dt \\
& \leq \|Z\|_p \|1 - T_{\tilde{S}_n}(\tilde{S}_n)\|_p \int_0^\infty \frac{e^{-2t}}{\sqrt{1-e^{-2t}}} \|u'_\alpha(e^{-t}\tilde{S}_n + \sqrt{1-e^{-2t}}Z)\|_{\frac{p}{p-1}} dt.
\end{aligned}$$

Now let $q \geq 2$ and $\rho \in (0, u - 2\alpha)$. Let G be independent from Z being such that $\|\tilde{S}_n - G\|_q = \mathcal{W}_q(\tilde{S}_n, G)$ then we have

$$\begin{aligned}
& \|u'_\alpha(e^{-t}\tilde{S}_n + \sqrt{1-e^{-2t}}Z)\|_{\frac{p}{p-1}} \\
& \leq \left\| \left[u'_\alpha(e^{-t}\tilde{S}_n + \sqrt{1-e^{-2t}}Z) - u'_\alpha(e^{-t}G + \sqrt{1-e^{-2t}}Z) \right] \mathbb{I}(X_t \in [u - 2\alpha, u]) \right\|_{\frac{p}{p-1}} \\
& \quad + \|u'_\alpha(Z)\|_{\frac{p}{p-1}} \\
& \leq \|u'_\alpha(Z)\|_{\frac{p}{p-1}} + \frac{1}{\alpha^2} e^{-t} \|(G - \tilde{S}_n) \mathbb{I}(X_t \in [u - 2\alpha, u])\|_{\frac{p}{p-1}} \\
& \leq \|u'_\alpha(Z)\|_{\frac{p}{p-1}} + \frac{1}{\alpha^2} e^{-t} \|(G - \tilde{S}_n) \mathbb{I}(X_t \in [u - 2\alpha, u])\|_{\frac{p}{p-1}} \\
& \leq \|u'_\alpha(Z)\|_{\frac{p}{p-1}} + \frac{1}{\alpha^2} e^{-t} \left[\|(G - \tilde{S}_n) \mathbb{I}(e^{-t}G + \sqrt{1-e^{-2t}}Z \in [u - 2\alpha - \rho, u + \rho])\|_{\frac{p}{p-1}} \right. \\
& \quad \left. + \|(G - \tilde{S}_n) \mathbb{I}(|G - \tilde{S}_n| \geq \rho e^t)\|_{\frac{p}{p-1}} \right] \\
& \stackrel{(a)}{\leq} \|u'_\alpha(Z)\|_{\frac{p}{p-1}} + \frac{1}{\alpha^2} e^{-t} \mathcal{W}_{\frac{pq}{p-1}}(\tilde{S}_n, G) \mathbb{P}(Z \in [u - 2\alpha - \rho, u + \rho])^{\frac{(q-1)(p-1)}{pq}} \\
& \quad + \frac{1}{\alpha^2} e^{-t(1+\frac{q'(p-1)}{p})} \frac{\mathcal{W}_{\frac{pq'}{p-1}}(\tilde{S}_n, Z)^{q'}}{\rho^{\frac{q'(p-1)}{p}}}.
\end{aligned}$$

where to obtain (a) we used Chebyshev inequality.

Therefore we obtain that

$$\begin{aligned}
& |\mathbb{E}(u_\alpha(S_n)) - \mathbb{E}(u_\alpha(Z))| \\
& \leq \|Z\|_p \|1 - T_{S_n}(S_n)\|_p \int_0^\infty \frac{e^{-2t}}{\sqrt{1-e^{-2t}}} \left[\|u'_\alpha(Z)\|_{\frac{p}{p-1}} + \frac{1}{\alpha^2} e^{-t(1+\frac{q'(p-1)}{p})} \right. \\
& \quad \left. \times \frac{\mathcal{W}_{\frac{pq'}{p-1}}(\tilde{S}_n, Z)^{q'}}{\rho^{\frac{q'(p-1)}{p}}} + \frac{1}{\alpha^2} e^{-t} \mathcal{W}_{\frac{pq}{p-1}}(\tilde{S}_n, G) \mathbb{P}(Z \in [u - 2\alpha - \rho, u + \rho])^{\frac{(q-1)(p-1)}{pq}} \right] dt.
\end{aligned}$$

Now we clearly notice that $\mathbb{P}(\tilde{S}_n \geq u) \leq \mathbb{E}(u_\alpha(\tilde{S}_n))$. Therefore we have

$$\begin{aligned}
& \mathbb{P}(S_n \geq \sigma u) \\
& \leq \mathbb{E}(u_\alpha(Z)) + \|Z\|_p \|1 - T_{\tilde{S}_n}(\tilde{S}_n)\|_p \left[\|u'_\alpha(Z)\|_{\frac{p}{p-1}} + \frac{\sqrt{\pi} \Gamma(\frac{q'(p-1)+2p}{2p})}{2\alpha^2 \Gamma(\frac{q'(p-1)+3p}{2p})} \frac{\mathcal{W}_{\frac{pq'}{p-1}}(\tilde{S}_n, Z)^{q'}}{\rho^{\frac{q'(p-1)}{p}}} \right. \\
& \quad \left. + \frac{\pi}{4\alpha^2} \mathcal{W}_{\frac{pq}{p-1}}(\tilde{S}_n, G) \mathbb{P}(Z \in [u - 2\alpha - \rho, u + \rho])^{\frac{(q-1)(p-1)}{pq}} \right].
\end{aligned}$$

■

Lemma 28 Suppose that the conditions of Thm. 6 hold. Write $B_R(\cdot)$ be the Hoeffding bound for a R^2 -subgaussian random variable. Then for all $t, u > 0$ we have

$$\mathbb{P}(X_t \geq u) \leq \min \left\{ B_R(u), \inf_{p \geq 2, \rho \in (0,1)} \Phi^c(\rho u) + \frac{\omega_p^R(1)}{(1-\rho)^p u^p} \right\};$$

where $\omega_p^R(\cdot)$ is defined in Eq. (30) of App. I.1.

Proof Let $p \geq 2$ be a real and let $G \sim N(0, 1)$ be a standard normal random variable independent from Z such that $\mathcal{W}_p(\tilde{S}_n, G) = \|G - \tilde{S}_n\|_p$ (exists). Let $u \geq 0$ then for all $\rho \in (0, 1)$ the following holds

$$\begin{aligned} \mathbb{P}(X_t \geq u) &\leq \mathbb{P}(X_t \geq u) \\ &\leq \left\{ \mathbb{P}(e^t G + \sqrt{1 - e^{-2t}} Z \geq u\rho) + e^{-tp} \frac{\|\tilde{S}_n - G\|_p^p}{(1-\rho)^p u^p} \right\} \\ &\leq \left\{ \Phi^c(u\rho) + e^{-tp} \frac{\mathcal{W}_p(S_n, Z)}{(1-\rho)^p u^p} \right\} \\ &\leq \left\{ \Phi^c(u\rho) + \frac{\mathcal{W}_p(\tilde{S}_n, Z)}{(1-\rho)^p u^p} \right\}. \end{aligned}$$

Therefore we obtain that

$$\mathbb{P}(X_t \geq u) \leq \inf_{p \geq 2, \rho \in (0,1)} \Phi^c(\rho u) + \frac{\omega_p^R(1)}{(1-\rho)^p u^p}.$$

Similarly we know that X_t is $\sqrt{1 - e^{-2t}} + e^{-t} R^2$ -subgaussian, it is therefore also R^2 subgaussian and this concluded the proof. ■

E.2. Proof of Thm. 7: Alternative Stein kernel tail bound

Thanks to Chatterjee (2012) we know that there is $G \sim N(0, 1)$ distributed such that for $\theta > 0$ we have

$$\mathbb{E}(e^{\theta|\tilde{S}_n - G|}) \leq 2e^{2\theta^2 \mathcal{S}_\infty(\sigma^{-1} Y_1 \| Z)^2}.$$

By Chebyshev inequality we therefore have for all $t > 0$ that

$$\mathbb{P}(|\tilde{S}_n - G| \geq t) \leq 2e^{-\theta t} e^{2\theta^2 \mathcal{S}_\infty(\sigma^{-1} Y_1 \| Z)^2}.$$

If we choose $\theta = \frac{t}{4\mathcal{S}_\infty(\sigma^{-1} Y_1 \| Z)^2}$ then we have

$$\mathbb{P}(|\tilde{S}_n - G| \geq t) \leq 2e^{-\frac{t^2}{8\mathcal{S}_\infty(\sigma^{-1} Y_1 \| Z)^2}}.$$

Therefore by the union bound we obtain that for all $u > 0$ and all $\alpha < u/2$ we have

$$\begin{aligned} P(\tilde{S}_n \geq u) &\leq P(\tilde{S}_n - G + G \geq u) \\ &\leq P(G \geq u - 2\alpha) + P(\tilde{S}_n - G \geq 2\alpha) \\ &\leq \Phi^c(u - 2\alpha) + 2e^{-\frac{\alpha^2}{2\mathcal{S}_\infty(\sigma^{-1} Y_1 \| Z)^2}}. \end{aligned}$$

E.3. Proof of Thm. 8: Efficient Hellinger-Stein kernel tail bound

We write $\tau_{\tilde{S}_n}$ the Stein kernel of \tilde{S}_n and its score function $S_{\tilde{S}_n}(\cdot)$. According to Thm. 3 we know that

$$\mathbb{P}(S_n \geq \sigma u) \leq \Phi^c(u) + \sqrt{2}H(\sigma^{-1}S_n, Z) \left[\sqrt{\hat{Q}_n(u)} + \sqrt{\Phi^c(u)} \right].$$

Moreover we know that $H(\sigma^{-1}S_n, Z) \leq \sqrt{KL(\tilde{S}_n||Z)}$. Furthermore according [Ledoux et al. \(2015\)](#) we know that

$$KL(\tilde{S}_n||Z) \leq \mathcal{S}_2^2(\tilde{S}_n||Z) \log \left(1 + \frac{\|S_{\tilde{S}_n}(\tilde{S}_n) - \tilde{S}_n\|_2^2}{\mathcal{S}_2^2(\tilde{S}_n||Z)} \right).$$

Firstly we note that the following is a Stein Kernel for \tilde{S}_n

$$\tau_{\tilde{S}_n}(\tilde{S}_n) \triangleq \frac{1}{n} \mathbb{E} \left(\sum_{i \leq n} \tau_{Y_1}(\sigma^{-1}Y_1) | \tilde{S}_n \right).$$

Therefore as a consequence of Jensen inequality we have: $\mathcal{S}_2(\tilde{S}_n||Z) \leq \frac{1}{\sqrt{n}} \mathcal{S}_2(\sigma^{-1}Y_1||S_n)$. As for all $c > 0$ we know that $x \rightarrow x \log(1 + \frac{c}{x})$ is a non-decreasing function then

$$KL(\tilde{S}_n||Z) \leq \frac{1}{n} \mathcal{S}_2^2(\sigma^{-1}Y_1||Z) \log \left(1 + \frac{n \|S_{\tilde{S}_n}(\tilde{S}_n) - \tilde{S}_n\|_2^2}{\mathcal{S}_2^2(\sigma^{-1}Y_1||Z)} \right).$$

Moreover we know that $S_{\tilde{S}_n}(\tilde{S}_n)$ is the only \tilde{S}_n -measurable random variable such that for all $\phi \in \mathbb{C}^\infty(\mathbb{R})$ we have

$$\mathbb{E}(\phi(\tilde{S}_n) S_{\tilde{S}_n}(\tilde{S}_n)) = \mathbb{E}(\phi'(\tilde{S}_n)).$$

Therefore we know that $S_{\tilde{S}_n}(\tilde{S}_n) = \mathbb{E}(\frac{1}{\sqrt{n}} \sum_{i \leq n} S_{Y_1}(\sigma Y_i) | \tilde{S}_n)$. Therefore by Jensen inequality, we obtain that:

$$\|S_{\tilde{S}_n}(\tilde{S}_n) - \tilde{S}_n\|_2 \leq \|S_{Y_1}(\sigma Y_1) - \sigma^{-1}Y_1\|_2.$$

This implies the desired result.

Appendix F. Proof of Thm. 9: Efficient zero-bias tail bound

The crux of our argument, inspired by [Ross \(2011, Thm. 3.27\)](#) and proved in App. F.1, shows that a zero-biased version S^* of a random variable S has tails close to those of a Gaussian whenever S^* and S have a close coupling.

Theorem 29 (Zero-bias tail bounds) *Suppose that S^* has the zero-bias distribution for a random variable S with $\mathbb{E}[S] = 0$ and $\text{Var}(S) = 1$ and that $S^* - S \leq \delta$ almost surely. For $U \sim \text{Unif}[0, 1]$ independent of (S, S^*) , define the intermediate variable $S' \triangleq S^* + U(S - S^*)$. Then, for all $u \geq 0$ and $\lambda \in [0, 1]$,*

$$\mathbb{P}(S^* > u) - \Phi^c(u) \leq \delta [h_u(\lambda u) + (h_u(u) - h_u(\lambda u))\mathbb{P}(S' > \lambda u) - \delta_u \mathbb{P}(S' > u)],$$

for h_u and δ_u defined in Eq. (11) and Eq. (12) respectively. If, in addition, $S - S^* \leq \delta$ almost surely, then

$$\mathbb{P}(S > u + \delta) \leq \frac{1}{1 + \delta \delta_u} \Phi^c(u) + \frac{\delta}{1 + \delta \delta_u} (h_u(\lambda u) + (h_u(u) - h_u(\lambda u))\mathbb{P}(S' > \lambda u)).$$

Hence, to establish the first claim, it only remains to construct a suitable zero-bias coupling for S_n/σ . By [Chen et al. \(2011, Lem. 2.8\)](#), $S_n^* = S_n + X_I^* - X_I$ has the zero-bias distribution for S_n . Therefore, by the zero-bias definition, S_n^*/σ has the zero-bias distribution for S_n/σ . Finally, $|S_n^*/\sigma - S_n/\sigma| = |X_I^* - X_I|/\sigma \leq 2R/(\sigma\sqrt{n})$ almost surely. The advertised result thus follows from the second claim of [Thm. 29](#) applied with $S = S_n/\sigma$, $S^* = S_n^*/\sigma$, and $\delta = 2R/(\sigma\sqrt{n})$.

Now suppose the random variables $(X_i)_{i=1}^{n+1}$ are identically distributed, define $S_{n+1} = \frac{\sqrt{n}}{\sqrt{n+1}}(S_n + X_{n+1})$, and consider an independently generated index $J \sim \text{Unif}(\{1, \dots, n+1\})$. By [Chen et al. \(2011, Lem. 2.8\)](#), $S_{n+1}^* = S_{n+1} + X_J^* - X_J$ has the zero-bias distribution for S_{n+1} . Define the intermediate variable $S'_{n+1} \triangleq (S_{n+1}^* + U(S_{n+1} - S_{n+1}^*))$. By symmetry, $S_n + X_{n+1}^* \stackrel{d}{=} \frac{\sqrt{n+1}}{\sqrt{n}} S_{n+1}^*$ and hence $S_n + X'_{n+1} \stackrel{d}{=} \frac{\sqrt{n+1}}{\sqrt{n}} S'_{n+1}$. In addition, since $X_{n+1}^* \wedge X'_{n+1} \geq -\frac{R}{\sqrt{n}}$,

$$\begin{aligned} \mathbb{P}(S > \sigma u \frac{\sqrt{n+1}}{\sqrt{n}} + \frac{R}{\sqrt{n}}) &\leq \mathbb{P}(S + X_{n+1}^* \wedge X'_{n+1} > \sigma u \frac{\sqrt{n+1}}{\sqrt{n}}) \\ &\leq \mathbb{P}(S_{n+1}^* > \sigma u) \wedge \mathbb{P}(S'_{n+1} > \sigma u). \end{aligned}$$

The result now follows from the first result of [Thm. 29](#) with $S = S_{n+1}/\sigma$, $S^* = S_{n+1}^*/\sigma$, and $\delta = 2R/(\sigma\sqrt{n+1})$.

F.1. Proof of [Thm. 29](#): Zero-bias tail bounds

Fix any $u \in \mathbb{R}$. Our proof structure, based on Stein's method, mimics that of [Ross \(2011, Thm. 3.27\)](#) but employs u -dependent bounds in place of the u -independent bounds invoked by [Ross](#).

Solving the Stein equation Next we define the *Stein equation* ([Stein, 1986](#))

$$f'_u(w) - wf_u(w) = \mathbb{I}[w \leq u] - \Phi(u).$$

By [Chen et al. \(2011, Lem. 2.2\)](#), the absolutely continuous function

$$f_u(w) = \sqrt{2\pi} \exp(\frac{w^2}{2}) \Phi(w \wedge u) \Phi^c(w \vee u) \quad (26)$$

solves the Stein equation, and we can therefore write

$$\mathbb{P}(S^* \leq u) - \Phi(u) = \mathbb{E}[f'_u(S^*) - S^* f_u(S^*)] =,$$

where the final equality uses the definition of the zero-biased distribution. Negating both sides, we obtain

$$\mathbb{P}(S^* > u) - \Phi^c(u) = \mathbb{E}[S^* f_u(S^*) - S f_u(S)].$$

Bounding the Stein solution Now let $g_u(w) = wf_u(w)$. By [Chen et al. \(2011, Eq. \(2.81\)\)](#), the function

$$\begin{aligned} h_u(w) &\triangleq \begin{cases} \sqrt{2\pi} \Phi^c(u) ((1 + w^2) \exp(w^2/2) \Phi(w) + w/\sqrt{2\pi}) & \text{if } w \leq u \\ \sqrt{2\pi} \Phi(u) ((1 + w^2) \exp(w^2/2) \Phi^c(w) - w/\sqrt{2\pi}) & \text{if } w > u \end{cases} \\ &= w(\Phi^c(u) - \mathbb{I}[w > u]) + (1 + w^2)f_u(w) \\ &= w(\mathbb{I}[w \leq u] \Phi^c(u) - \mathbb{I}[w > u] \Phi(u)) + (1 + w^2)f_u(w) \end{aligned} \quad (27)$$

matches the derivative of $g_u(w)$ whenever $w \neq u$. Since the absolute continuity of f_u implies that g_u is absolutely continuous on compact intervals, the fundamental theorem of calculus yields

$$\mathbb{E}[g_u(S^*) - g_u(S)] = \mathbb{E}[\int_0^1 h_u(S^* + x(S - S^*))(S^* - S)dx] = \mathbb{E}[h_u(S')(S^* - S)].$$

Our next result, proved in App. F.2 provides a suitable u -dependent bound on h_u .

Lemma 30 (Growth of h_u) For any $u \geq 0, \lambda \in [0, 1]$, and $w \in \mathbb{R}$,

$$0 \leq h_u(w) \leq h_u(\lambda u) + (h_u(u) - h_u(\lambda u))\mathbb{I}[w > \lambda u] - \delta_u \mathbb{I}[w > u].$$

for h_u and δ_u defined in Eq. (27) and Eq. (12) respectively.

S^* tail bound Fix any $\lambda \in [0, 1]$. Lem. 30, our almost sure assumption $S^* - S \leq \delta$, and the definition of S' together imply that

$$\begin{aligned} \mathbb{P}(S^* > u) - \Phi^c(u) &\leq \delta \mathbb{E}[h_u(S')] \\ &\leq \delta [h_u(\lambda u) + (h_u(u) - h_u(\lambda u))\mathbb{P}(S' > \lambda u) - \delta_u \mathbb{P}(S' > u)]. \end{aligned} \quad (28)$$

From S tails to S^* tails When $S - S^* \leq \delta$ almost surely, we additionally have

$$\begin{aligned} S - S' &\leq (S - S^*)(1 - U) \leq \delta \quad \text{and therefore} \\ \mathbb{P}(S > u + \delta) &= \mathbb{P}(S^* \wedge S' > u + \delta + S^* \wedge S' - S) \\ &\leq \mathbb{P}(S^* \wedge S' > u) \leq \mathbb{P}(S^* > u) \wedge \mathbb{P}(S' > u). \end{aligned}$$

Combining this inequality with our S^* tail bound Eq. (28) yields

$$\begin{aligned} \mathbb{P}(S > u + \delta) \\ \leq \Phi^c(u) + \delta [h_u(\lambda u) + (h_u(u) - h_u(\lambda u))\mathbb{P}(S' > \lambda u) - \delta_u \mathbb{P}(S > u + \delta)]. \end{aligned}$$

Rearranging the terms of this expression yields the final advertised result.

F.2. Proof of Lem. 30: Growth of h_u

Our proof makes use of the following bound on the growth of Φ^c , established in App. F.3.

Lemma 31 (Growth of Φ^c) If $w \geq 0$ then

$$\sqrt{2\pi}(1 + w^2) \exp(w^2/2) \Phi^c(w) - w \leq b(w) \triangleq \frac{2}{w + \sqrt{w^2 + 8/\pi}} - \frac{8w}{\pi(w + \sqrt{w^2 + 8/\pi})^2}.$$

Fix any $u \geq 0, \lambda \in [0, 1]$, and $w \in \mathbb{R}$. We will divide our proof into cases based on the value of w .

Lower bound, $w \neq u$: Since g_u is differentiable for $w \neq u$ and increasing by Chen et al. (2011, Lem. 2.3), $h_u(w) = g'_u(w) \geq 0$.

Lower bound, $w = u$: We have $h_u(u) = \lim_{v \uparrow u} h_u(v) \geq 0$.

Upper bound, $w > u$: From the definition of f_u Eq. (26) and the Φ^c growth bound (Lem. 31), we have

$$\begin{aligned}(1 + w^2)f_u(w)\mathbb{I}[w > u] &= \sqrt{2\pi}(1 + w^2)\exp(w^2/2)\Phi^c(w)\Phi(u)\mathbb{I}[w > u] \\ &\leq (b(u) + w)\Phi(u)\mathbb{I}[w > u].\end{aligned}$$

Hence,

$$h_u(w)\mathbb{I}[w > u] = (1 + w^2)f_u(w)\mathbb{I}[w > u] - w\Phi(u)\mathbb{I}[w > u] \leq b(u)\Phi(u)\mathbb{I}[w > u].$$

Upper bound, $\lambda u \geq w \geq 0$: Since g_u is increasing (Chen et al., 2011, Lem. 2.3), f_u is increasing for $w < u$, and $\lambda u f_u(\lambda u) + \Phi^c(u)$ is nonnegative, we have

$$\begin{aligned}h_u(w)\mathbb{I}[\lambda u \geq w \geq 0] &= (f_u(w) + w(wf_u(w) + \Phi^c(u)))\mathbb{I}[\lambda u \geq w \geq 0] \\ &\leq [f_u(\lambda u) + w(\lambda u f_u(\lambda u) + \Phi^c(u))]\mathbb{I}[\lambda u \geq w \geq 0] \\ &\leq [f_u(\lambda u) + \lambda u(\lambda u f_u(\lambda u) + \Phi^c(u))]\mathbb{I}[\lambda u \geq w \geq 0] \\ &= h_u(\lambda u)\mathbb{I}[\lambda u \geq w \geq 0] = h_u(\lambda u)(\mathbb{I}[w \geq 0] - \mathbb{I}[w > \lambda u]).\end{aligned}$$

Upper bound, $w < 0$: Since $w < 0 \leq u$, we use the definition of f_u Eq. (26), the Φ^c growth bound (Lem. 31), the nonnegativity of Φ^c , and the fact that b is decreasing to derive

$$\begin{aligned}(1 + w^2)f_u(w)\mathbb{I}[w < 0] &= \sqrt{2\pi}\exp(w^2/2)\Phi^c(|w|)\Phi^c(u)\mathbb{I}[w < 0] \\ &\leq (b(|w|) + |w|)\Phi^c(u)\mathbb{I}[w < 0] \\ &\leq (b(0) + |w|)\Phi^c(u)\mathbb{I}[w < 0].\end{aligned}$$

Since $\lambda u \geq 0$ by assumption, our prior derivation implies $h_u(\lambda u) \geq h_u(0)$ and hence,

$$\begin{aligned}h_u(w)\mathbb{I}[w < 0] &= ((b(0) + |w|) - |w|)\Phi^c(u)\mathbb{I}[w < 0] \\ &= b(0)\Phi^c(u)\mathbb{I}[w < 0] = \sqrt{\frac{\pi}{2}}\Phi^c(u)\mathbb{I}[w < 0] \\ &= h_u(0)\mathbb{I}[w < 0] \leq h_u(\lambda u)\mathbb{I}[w < 0].\end{aligned}$$

Upper bound, $u \geq w > \lambda u$: Since $u, w, f_u(u)$, and $\Phi^c(u)$ are nonnegative, g_u is increasing (Chen et al., 2011, Lem. 2.3), and $u \geq w$, we have

$$\begin{aligned}h_u(w)\mathbb{I}[u \geq w > \lambda u] &= (f_u(w) + w(wf_u(w) + \Phi^c(u)))\mathbb{I}[u \geq w > \lambda u] \\ &\leq (f_u(u) + w(uf_u(u) + \Phi^c(u)))\mathbb{I}[u \geq w > \lambda u] \\ &\leq (f_u(u) + u(uf_u(u) + \Phi^c(u)))\mathbb{I}[u \geq w > \lambda u] \\ &= h_u(u)\mathbb{I}[u \geq w > \lambda u] = h_u(u)(\mathbb{I}[w > \lambda u] - \mathbb{I}[w > u]).\end{aligned}$$

Complete upper bound Taken together, our upper bounds yield

$$\begin{aligned}h_u(w) &\leq b(u)\Phi(u)\mathbb{I}[w > u] + h_u(u)(\mathbb{I}[w > \lambda u] - \mathbb{I}[w > u]) \\ &\quad + h_u(\lambda u)(\mathbb{I}[w \geq 0] - \mathbb{I}[w > \lambda u]) + h_u(\lambda u)\mathbb{I}[w < 0] \\ &\leq (b(u)\Phi(u) - h_u(u))\mathbb{I}[w > u] + (h_u(u) - h_u(\lambda u))\mathbb{I}[w > \lambda u] + h_u(\lambda u).\end{aligned}$$

F.3. Proof of Lem. 31: Growth of Φ^c

By Abramowitz and Stegun (1964, 7.1.13)

$$\exp(w^2/2)\Phi^c(w) \leq \sqrt{2/\pi}/(w + \sqrt{w^2 + 8/\pi}).$$

Therefore,

$$\begin{aligned} & \sqrt{2\pi}(1 + w^2) \exp(w^2/2)\Phi^c(w) - w \\ & \leq 2(1 + w^2)/(w + \sqrt{w^2 + 8/\pi}) - w \\ & = (2(1 + w^2) - w^2 - w\sqrt{w^2 + 8/\pi})/(w + \sqrt{w^2 + 8/\pi}) \\ & = (2 + w(w - \sqrt{w^2 + 8/\pi}))/ (w + \sqrt{w^2 + 8/\pi}) \\ & = \frac{2}{w + \sqrt{w^2 + 8/\pi}} - \frac{8w}{\pi(w + \sqrt{w^2 + 8/\pi})^2}. \end{aligned}$$

Appendix G. Proof of Prop. 10: Properties of S'_n

Instantiate the notation and assumptions of Thm. 9, and fix any $i \in \{1, \dots, n\}$ and $t \in \mathbb{R}$. We invoke the definition of X'_i , Lem. 2.1(iv) of Goldstein and Reinert (1997), and our X_i boundedness assumption in turn to find that

$$|\mathbb{E}[X'_i]| = \frac{1}{2}|\mathbb{E}[X_i^*]| = \frac{1}{4\sigma_i^2}|\mathbb{E}[X_i^3]| \leq \frac{R}{4\sqrt{n}}.$$

The same invocations, coupled with the independence of X_i , U , and X_i^* , imply

$$\begin{aligned} \text{Var}(X'_i) &= \mathbb{E}[(UX_i + (1 - U)(X_i^* - \mathbb{E}[X_i^*]))^2] \\ &= \mathbb{E}[U^2]\text{Var}(X_i) + \mathbb{E}[(1 - U)^2]\text{Var}(X_i^*) \\ &= \frac{1}{3}\sigma_i^2 + \frac{1}{3}\text{Var}(X_i^*) = \frac{1}{3}\sigma_i^2 + \frac{1}{3}(\mathbb{E}[X_i^{*2}] - \mathbb{E}[X_i^*]^2) \\ &= \frac{1}{3}\sigma_i^2 + \frac{1}{9\sigma_i^2}\mathbb{E}[X_i^4] - \frac{1}{12\sigma_i^4}\mathbb{E}[X_i^3]^2 \leq \frac{1}{3}\sigma_i^2 + \frac{R^2}{9n}. \end{aligned}$$

Moreover, since $\mathbb{E}[X_i^3]^2 \leq \mathbb{E}[X_i^2]\mathbb{E}[X_i^4]$ by Cauchy Schwartz, we have

$$\text{Var}(X'_i) \geq \frac{1}{3}\sigma_i^2 + \frac{1}{9\sigma_i^2}\mathbb{E}[X_i^4] - \frac{1}{12\sigma_i^2}\mathbb{E}[X_i^4] = \frac{1}{3}\sigma_i^2 + \frac{1}{36\sigma_i^2}\mathbb{E}[X_i^4] \geq \frac{1}{3}\sigma_i^2 + \frac{R^2}{36n}.$$

Next, by Goldstein and Reinert (1997, Lem. 2.1(ii)), the support of X_i^* is the closed convex hull of the support of X_i . Therefore, by the triangle inequality,

$$|X'_i| \leq U|X_i| + (1 - U)|X_i^*| \leq \frac{R}{\sqrt{n}} \quad \text{almost surely.}$$

The final result is justified exactly as in App. F.

Appendix H. Proof of Thms. 11 and 12: Efficient Concentration with Wasserstein Approximation

The proofs of Thm. 11 and Thm. 12 exploit the following upper bound for $\mathcal{W}_p(S_n, \mathcal{N}(0, \sigma^2))$.

Lemma 32 *If the conditions of Thm. 11 hold and we write $D_{n,p} \triangleq \sqrt{1 - \frac{\tilde{R}^2(p-1)}{Cn}}$ where $C = 0.4$, then for all $p \geq 2$ the following upper-bound holds:*

$$\begin{aligned} \mathcal{W}_p(\tilde{S}_n, Z) &\leq \frac{(p-1)}{\sqrt{0.4n}} \left[\tilde{R} \left(1 + \frac{\tilde{U}_{n,p}}{2\sqrt{n}} \right) + \frac{\tilde{R}}{2\sqrt{n}} \left(\frac{3A_p}{2} + \tilde{R}^{1+2/p} \tilde{A}_{n,p} \right) \right. \\ &\quad \left. + \frac{\sqrt{0.4}}{2\sqrt{p-1}} \left(1 - \frac{\tilde{R}\sqrt{p-1}}{\sqrt{0.4n}} \right) \left[U_{n,p} - \tilde{R}^{-1} \tilde{U}_{n,p} \right] \right] \\ &\quad + \frac{\sqrt{e}(p-1)\tilde{R}^2}{1.2n} [e^{0.8} - 1] \left[\frac{2\tilde{R}\tilde{U}_{n,p}}{3D_{n,p}} + U_{n,p} \log \left(\frac{D_{n,p}+1}{1-D_{n,p}} \right) \right] \\ &\quad + \frac{2\sqrt{e}(p-1)\tilde{R}}{\sqrt{1.2n}} [e^{0.8} - 1.8] \left[\frac{\tilde{R}^2}{3\sqrt{n}} \tilde{U}_{n,p} + 2 + \frac{\tilde{R}U_{n,p}}{\sqrt{n}} - \frac{2}{3\tilde{R}} \right] \\ &\quad + \frac{\tilde{R}^2(p-1)}{0.8nD_{n,p}} [U_{n,p} + \tilde{U}_{n,p}]. \end{aligned}$$

Proof Using Lem. 34 we know that

$$\begin{aligned} \mathcal{W}_p(\tilde{S}_n, Z) &\leq \frac{\sqrt{p-1}}{\sqrt{n\kappa}} \left[\tilde{R} \left(1 + \frac{\tilde{U}_{n,p}}{2\sqrt{n}} \right) + \frac{\tilde{R}}{2\sqrt{n}} \left(\frac{3A_p}{2} + \tilde{R}^{1+2/p} \tilde{A}_{n,p} \right) \right. \\ &\quad \left. + \frac{\sqrt{\kappa}}{2} \left(1 - \frac{\tilde{R}}{\sqrt{n\kappa}} \right) \left[U_{n,p} - \tilde{R}^{-1} \tilde{U}_{n,p} \right] \right] \\ &\quad + \frac{\sqrt{e}\tilde{R}^2}{3n\kappa} [e^{2(p-1)\kappa} - 1] \left[\frac{2\tilde{R}\tilde{U}_{n,p}}{3M_{n,\kappa}} + U_{n,p} \log \left(\frac{M_{n,\kappa}+1}{1-M_{n,\kappa}} \right) \right] \\ &\quad + \frac{2\sqrt{e}\sqrt{p-1}\tilde{R}}{\sqrt{3n\kappa}} [e^{2(p-1)\kappa} - 1 - 2(p-1)\kappa] \left[\frac{\tilde{R}^2}{3\sqrt{n}} \tilde{U}_{n,p} + 2 + \frac{\tilde{R}U_{n,p}}{\sqrt{n}} - \frac{2}{3\tilde{R}} \right] \\ &\quad + \frac{\tilde{R}^2}{2n\kappa M_{n,\kappa}} [U_{n,p} + \tilde{U}_{n,p}]. \end{aligned}$$

We obtain the advertised bound by choosing $\kappa = C/(p-1)$ and $C = 0.4$. ■

H.1. Definition of $r_{n,t}$ in Thm. 11

The bound in Thm. 11 is stated in function of a small reminder term $r_{n,t}$ which we define in this section. We write

$$\begin{aligned} \frac{1}{e\sigma} r_{n,t} &\triangleq \frac{(p+2)}{\sqrt{0.4n}} \left[\tilde{R} \frac{\tilde{U}_n}{2\sqrt{n}} + \frac{\tilde{R}}{2\sqrt{n}} \left(\frac{A_{L_{n,t}}}{2} + \tilde{U}_n \right) + \frac{\sqrt{0.4}}{2\sqrt{L_{n,t}-1}} M_n \left[\frac{\sqrt{L_{n,t}+2}\sqrt{e}((2e)^{1/L_{n,t}}-1)}{\sqrt{2}} \right. \right. \\ &\quad \left. \left. - \tilde{R}^{-1} \tilde{U}_n \right] \right] + \frac{2\sqrt{e}(L_{n,t}-1)\tilde{R}}{\sqrt{1.2n}} [e^{0.8} - 1.8] \left[\frac{\tilde{R}^2}{3} \tilde{U}_n + \tilde{R}U_n \right] \\ &\quad + \frac{\tilde{R}^2(L_{n,t}-1)}{0.8nM_n} [U_n + \tilde{U}_n] + \frac{\sqrt{e}(L_{n,t}-1)\tilde{R}^2}{1.2n} [e^{0.8} - 1] \left[\frac{2\tilde{R}\tilde{U}_n}{3M_n} + U_n \log \left(\frac{M_n+1}{1-M_n} \right) \right], \end{aligned}$$

where we set

$$M_n \triangleq \sqrt{1 - \frac{\tilde{R}^2(L_{n,t}-1)}{0.4*n\sigma^2}}, \quad U_n \triangleq \frac{\sqrt{e}(2e)^{1/L_{n,t}}\sqrt{L_{n,t}+2}}{\sqrt{2}} + \frac{(L_{n,t}+2)}{2} \left(\frac{\tilde{R}}{\sqrt{n}} \right)^{1-2/L_{n,t}},$$

$$\text{and } \tilde{U}_n \triangleq \frac{\sqrt{e}(2e)^{1/L_{n,t}}\sqrt{L_{n,t}+2}}{\sqrt{2}} + \frac{(L_{n,t}+2)\tilde{R}^{2/L_{n,t}}}{2} \left(\frac{\tilde{R}}{\sqrt{n}} \right)^{1-2/L_{n,t}}.$$

H.2. Proof of Thm. 11: Efficient tail bound

As the proof of the two results is very similar we show only the details of the proof of the first result and highlight the differences with the proof of the second result.

The first step is to notice that by definition of the metric $\mathcal{W}_p(\cdot, \cdot)$ for all $\epsilon > 0$ we can find $G \sim \mathcal{N}(0, \sigma^2)$ a normally distributed random variable that respects $\|S_n - G\|_p \leq \epsilon + \mathcal{W}_p(S_n, \mathcal{N}(0, \sigma^2))$. This implies that for all $\rho \in (0, 1)$ we have

$$\begin{aligned} \mathbb{P}(S_n \geq t\sigma) &= \mathbb{P}(G + S_n - G \geq t\sigma) \\ &\leq \mathbb{P}(G \geq \rho t\sigma) + \mathbb{P}(S_n - G \geq (1 - \rho)t\sigma) \\ &\leq \Phi^c(\rho t) + \frac{\mathbb{E}(|S_n - G|^p)}{(1 - \rho)^p (t\sigma)^p} \\ &\leq \Phi^c(\rho t) + \frac{(\epsilon + \mathcal{W}_p(S_n, \mathcal{N}(0, \sigma^2)))^p}{(1 - \rho)^p (t\sigma)^p}. \end{aligned}$$

As this holds for arbitrary $\epsilon \in \mathbb{R}^+$, $p \geq 2$, and $\rho \in (0, 1)$ we obtain that

$$\mathbb{P}(S_n \geq t\sigma) \leq \inf_{p \geq 2, \rho \in (0, 1)} \Phi^c(\rho t) + \frac{\mathcal{W}_p^p(S_n, \mathcal{N}(0, \sigma^2))}{(1 - \rho)^p (t\sigma)^p}.$$

Our objective is to upper bound the right-hand side of this equation. In this goal and for ease of notation we shorthand $\omega_p \triangleq \mathcal{W}_p(S_n, \mathcal{N}(0, \sigma^2))$. This directly implies that:

$$\mathbb{P}(S_n \geq \sigma t) \leq \inf_{\substack{\rho \in [0, 1] \\ p \geq 2}} \Phi^c(\rho t) + \frac{\omega_p^p}{(1 - \rho)^p (t\sigma)^p}. \quad (29)$$

We proceed in two successive steps. First, for all $\rho \in (0, 1)$ we choose an integer $p_\rho \in \mathbb{N}$ that makes the right hand small and select a specific value for ρ only in a second stage. In this goal we first note that:

$$\begin{aligned} &\frac{\omega_p^p}{(1 - \rho)^p (t\sigma)^p} \\ &\leq \frac{1}{(1 - \rho)^p (t\sigma)^p} \left[\frac{(p+2)\sigma}{\sqrt{0.4n}} \left[\tilde{R} + \frac{\sqrt{0.4}\sqrt{e}}{2\sqrt{2}} \right] + \frac{2\sqrt{e}(p+2)\sigma\tilde{R}}{\sqrt{1.2n}} \left[e^{0.8} - 1.8 \right] \left[2 - \frac{2}{3\tilde{R}} \right] + \epsilon_{n,p} \right]^p \end{aligned}$$

where we have set

$$\begin{aligned} \frac{1}{\sigma} \epsilon_{n,p} &\triangleq \frac{(p+2)}{\sqrt{0.4n}} \left[\tilde{R} \frac{\tilde{U}_{n,p}}{2\sqrt{n}} + \frac{\tilde{R}}{2\sqrt{n}} \left(\frac{3A_p}{2} + \tilde{R}^{1+2/p} \tilde{A}_{n,p} \right) + \frac{\sqrt{0.4}}{2\sqrt{p-1}} \left(1 - \frac{\tilde{R}\sqrt{p-1}}{\sqrt{0.4n}} \right) \right. \\ &\quad \times \left. \left[\frac{\sqrt{p+2}\sqrt{e}((2e)^{1/p}-1)}{\sqrt{2}} - \tilde{R}^{-1} \tilde{U}_{n,p} \right] \right. \\ &\quad + \frac{\sqrt{e}(p-1)\tilde{R}^2}{1.2n} [e^{0.8} - 1] \left[\frac{2\tilde{R}\tilde{U}_{n,p}}{3D_{n,p}} + U_{n,p} \log \left(\frac{D_{n,p}+1}{1-D_{n,p}} \right) \right] \\ &\quad \left. + \frac{2\sqrt{e}(p-1)\tilde{R}}{\sqrt{1.2n}} [e^{0.8} - 1.8] \left[\frac{\tilde{R}^2}{3} \tilde{U}_{n,p} + \tilde{R}U_{n,p} \right] + \frac{\tilde{R}^2(p-1)}{0.8nD_{n,p}} [U_{n,p} + \tilde{U}_{n,p}] \right]. \end{aligned}$$

Minimizing $\frac{\omega_p^p}{(1 - \rho)^p t^p}$ in terms of p is complex. In the goal of obtaining a simple expression we instead choose to minimize the following alternative quantity

$$\frac{1}{(1 - \rho)^p (t\sigma)^p} \left[\frac{(p+2)\sigma}{\sqrt{0.4n}} \left[\tilde{R} + \frac{\sqrt{0.4}\sqrt{e}}{2\sqrt{2}} \right] + \frac{2\sqrt{e}(p+2)\sigma\tilde{R}}{\sqrt{1.2n}} [e^{0.8} - 1.8] \left[2 - \frac{2}{3\tilde{R}} \right] \right]^p.$$

In this goal we write

$$K \triangleq \frac{\sigma}{\sqrt{0.4}} \left[\tilde{R} + \frac{\sqrt{0.4e}}{2\sqrt{2}} \right] + \frac{2\sqrt{e}\sigma\tilde{R}}{\sqrt{1.2}} \left[e^{0.8} - 1.8 \right] \left[2 - \frac{2}{3\tilde{R}} \right].$$

and we define $h_\rho(p) \rightarrow \left[\frac{(p+2)K}{\sqrt{nt\sigma(1-\rho)}} \right]^p$. We remark that h_ρ is smooth and minimized for $p \geq 2$ satisfying $p + 2 = e^{-1} \frac{\sqrt{nt\sigma(1-\rho)}}{K} e^{-\frac{1}{p-1}}$. If $n \in \mathbb{N}$ is large then this equation is approximately solved by p solving the following alternative equation $p + 2 = e^{-1} \frac{\sqrt{nt\sigma(1-\rho)}}{K}$. Therefore we choose $p_\rho^* = e^{-1} \frac{\sqrt{nt\sigma(1-\rho)}}{K} - 2 \geq 2$. We remark that this implies that $\frac{K(p_\rho^*+2)}{\sqrt{nt\sigma(1-\rho)}} = e^{-1}$. Therefore we obtain

$$\begin{aligned} \left[\frac{\omega_{p_\rho^*}}{(1-\rho)t\sigma} \right]^{p_\rho^*} &\leq \left[e^{-1} + \epsilon_{n,p_\rho^*} \right]^{p_\rho^*} \\ &\leq e^{-e^{-1} \frac{\sqrt{nt\sigma(1-\rho)}}{K} + 2} [1 + e \times \epsilon_{n,p_\rho^*}]^{p_\rho^*}. \end{aligned}$$

Therefore according to Eq. (29) we obtain that

$$\mathbb{P}(S_n \geq \sigma t) \leq \min_{\rho \in (0,1)} \Phi^c(\rho t) + e^{-e^{-1} \frac{\sqrt{nt\sigma(1-\rho)}}{K} + 2} [1 + e \times \epsilon_{n,p_\rho^*}]^{p_\rho^*}.$$

To obtain a closed-form formula upper bound to Eq. (29) we want to minimize

$$g(\rho) \triangleq \Phi^c(\rho t) + e^{-e^{-1} \frac{\sqrt{nt\sigma(1-\rho)}}{K} + 2}.$$

In this goal denote $C_{n,t} \triangleq \frac{e^{-1}\sqrt{nt\sigma}}{K}$ and note that g is smooth and that its derivative is:

$$g'(\rho) = -t\varphi(\rho t) + C_{n,t}e^2e^{-C_{n,t}(1-\rho)}.$$

Therefore if $\frac{C_{n,t}}{t}e^{-C_{n,t}-1} \in \left[\frac{1}{\sqrt{2\pi}}, \varphi(t)e^{-C_{n,t}} \right]$ we have that $g(\cdot)$ is minimized by $\rho_{n,t}^* \in (0, 1)$ satisfying:

$$\varphi(\rho_{n,t}^* t) e^{-\rho_{n,t}^* C_{n,t}} = \frac{C_{n,t}}{t} e^{-C_{n,t}+2}.$$

This equation is solved by

$$\begin{aligned} \rho_{n,t}^* &\triangleq \frac{-C_{n,t} + \sqrt{C_{n,t}^2 + 2t^2(C_{n,t} - 2 - \log(C_{n,t}\sqrt{2\pi}/t))}}{t^2} \\ &\stackrel{(a)}{\sim} 1 - \frac{\log(C_{n,t}\sqrt{2\pi}/t)}{C_{n,t}}, \end{aligned}$$

where to get (a) we used the fact that $C_{n,t} = o_n(C_{n,t}^2)$ therefore we have

$$\begin{aligned} &\sqrt{C_{n,t}^2 + 2t^2(C_{n,t} - 2 - \log(C_{n,t}\sqrt{2\pi}/t))} \\ &\sim \sqrt{C_{n,t}^2} + \frac{1}{2\sqrt{C_{n,t}^2}} 2t^2(C_{n,t} - 2 - \log(C_{n,t}\sqrt{2\pi}/t)) \\ &\sim C_{n,t} + t^2 C_{n,t}^{-1} (C_{n,t} - 2 - \log(C_{n,t}\sqrt{2\pi}/t)). \end{aligned}$$

The advertised result now follows directly.

H.3. Definition of $s_{n,\delta}$ in Thm. 12

The bound in Thm. 12 has a small reminder term $s_{n,\delta}$ which we define in this section. We write

$$\begin{aligned} \frac{1}{\sigma} s_{n,\delta} &\triangleq \frac{1}{\sqrt{0.4}} \left[\tilde{R} \frac{\tilde{U}_n^\delta}{2\sqrt{n}} + \frac{\tilde{R}}{2\sqrt{n}} \left(\frac{A_\delta^n}{2} + U_n^\delta \right) + \frac{\sqrt{0.4}}{2\sqrt{A_\delta^n-1}} \right. \\ &\quad \times \left(1 - \frac{\tilde{R}\sqrt{A_\delta^n-1}}{\sqrt{0.4n}} \right) \left[\frac{\sqrt{A_\delta^n+2}\sqrt{e}((2e)^{1/A_\delta^n-1})}{\sqrt{2}} - \tilde{R}^{-1}\tilde{U}_n^\delta \right] \\ &\quad + \frac{\sqrt{e}(A_\delta^n-1)\tilde{R}^2}{1.2\sqrt{n}\sqrt{A_\delta^n+2}} [e^{0.8} - 1] \left[\frac{2\tilde{R}\tilde{U}_n^\delta}{3M_n^\delta} + U_n^\delta \log \left(\frac{M_n^\delta+1}{1-M_n^\delta} \right) \right] \\ &\quad + \frac{2\sqrt{e}(A_\delta^n-1)\tilde{R}}{\sqrt{1.2n}\sqrt{A_\delta^n+2}} [e^{0.8} - 1.8] \left[\frac{\tilde{R}^2}{3}\tilde{U}_n^\delta + \tilde{R}U_n^\delta \right] \\ &\quad + \frac{\tilde{R}^2(A_\delta^n-1)}{0.8\sqrt{n}(A_\delta^n+2)M_n^\delta} \left[U_{n,A_\delta^n} + \tilde{U}_{n,A_\delta^n} \right], \end{aligned}$$

where we set

$$M_n^\delta \triangleq \sqrt{1 - \frac{R^2 A_\delta^n}{0.4n\sigma^2}}, \quad U_n^\delta \triangleq \frac{\sqrt{e}(2e)^{1/A_\delta^n}\sqrt{A_\delta^n+2}}{\sqrt{2}} + \frac{(A_\delta^n+2)}{2} \left(\frac{\tilde{R}}{\sqrt{n}} \right)^{1-2/A_\delta^n},$$

$$\tilde{U}_n^\delta \triangleq \frac{\sqrt{e}(2e)^{1/A_\delta^n}\sqrt{A_\delta^n+2}}{\sqrt{2}} + \frac{(A_\delta^n+2)\tilde{R}^{2/A_\delta^n}}{2} \left(\frac{\tilde{R}}{\sqrt{n}} \right)^{1-2/A_\delta^n}.$$

H.4. Proof of Thm. 12: Efficient quantile bound

Let $G \sim \mathcal{N}(0, \sigma^2)$ be a normal random variable such that $\|S_n - G\|_p \leq \epsilon + \mathcal{W}_p(S_n, \mathcal{N}(0, \sigma^2))$.

Let $u > 0$ be such that $\frac{\mathcal{W}_p(S_n, \mathcal{N}(0, \sigma^2))^p}{u^p} \leq \alpha$ then for any $t < \frac{1}{2}$ we have:

$$\mathbb{P}(S_n \geq u - \sigma\Phi^{-1}(t)) \leq t + \left(\alpha^{1/p} + \frac{\epsilon}{u} \right)^p.$$

Therefore as $\epsilon > 0$ is arbitrary we have

$$\mathbb{P}(S_n \geq u - \sigma\Phi^{-1}(t)) \leq t + \alpha.$$

Therefore to obtain the desired result we work in two step: i) For every confidence $\alpha > 0$, we choose $u_\alpha^* > 0$ to be as small as possible while satisfying $\frac{\mathcal{W}_p(S_n, \mathcal{N}(0,1))^p}{u^p} \leq \alpha$ and (ii) choose α such that $u_\alpha^* - \sigma\Phi^{-1}(\delta - \alpha)$ is as small as possible.

Given a confidence level $\alpha > 0$, our first goal is to find the smallest u^* such that $\frac{\mathcal{W}_p(S_n, \mathcal{N}(0,1))^p}{u^p} \leq \alpha$. With that goal we define

$$g(p) = \frac{\mathcal{W}_p(S_n, \mathcal{N}(0, \sigma^2))}{\alpha^{1/p}}.$$

We remark that $u^* = \inf_p g(p)$. Using Lem. 32 we have:

$$g(p) = \frac{\mathcal{W}_p(S_n, \mathcal{N}(0, \sigma^2))}{\alpha^{1/p}} \leq \frac{1}{\alpha^{1/p}} \left[\frac{K(p+2)}{\sqrt{n}} + \epsilon_{n,p} \right],$$

where we set $\epsilon_{n,p}$ to be

$$\begin{aligned} \frac{1}{\sigma} \epsilon_{n,p} &\triangleq \frac{(p+2)}{\sqrt{0.4n}} \left[\tilde{R} \frac{\tilde{U}_{n,p}}{2\sqrt{n}} + \frac{\tilde{R}}{2\sqrt{n}} \left(\frac{3A_p}{2} + \tilde{R}^{1+2/p} \tilde{A}_{n,p} \right) \right. \\ &\quad \left. + \frac{\sqrt{0.4}}{2\sqrt{p-1}} \left(1 - \frac{\tilde{R}\sqrt{p-1}}{\sqrt{0.4n}} \right) \times \left[\frac{\sqrt{p+2}\sqrt{e}((2e)^{1/p}-1)}{\sqrt{2}} - \tilde{R}^{-1} \tilde{U}_{n,p} \right] \right. \\ &\quad \left. + \frac{\sqrt{e}(p-1)\tilde{R}^2}{1.2n} [e^{0.8} - 1] \left[\frac{2\tilde{R}\tilde{U}_{n,p}}{3D_{n,p}} + U_{n,p} \log \left(\frac{D_{n,p}+1}{1-D_{n,p}} \right) \right] \right. \\ &\quad \left. + \frac{2\sqrt{e}(p-1)\tilde{R}}{\sqrt{1.2n}} [e^{0.8} - 1.8] \left[\frac{\tilde{R}^2}{3} \tilde{U}_{n,p} + \tilde{R}U_{n,p} \right] + \frac{\tilde{R}^2(p-1)}{0.8nD_{n,p}} [U_{n,p} + \tilde{U}_{n,p}] \right]. \end{aligned}$$

As minimizing $\frac{1}{\alpha^{1/p}} \left[\frac{K(p+2)}{\sqrt{n}} + \epsilon_{n,p} \right]$ in terms of $p \geq 2$ is tricky we simplify things and instead choose to minimize $\frac{pK}{\sqrt{n}\alpha^{1/p}}$. This is minimized for $p_\alpha^* = -\log(\alpha)$. Therefore we have

$$g(p_\alpha^*) \leq e \left[\frac{K(-\log(\alpha)+2)}{\sqrt{n}} + r_{n,\alpha} \right];$$

where we have set

$$\begin{aligned} \frac{1}{\sigma} r_{n,\alpha} &\triangleq \frac{(-\log(\alpha)+2)}{\sqrt{0.4n}} \left[\tilde{R} \frac{\tilde{U}_{n,-\log(\alpha)}}{2\sqrt{n}} + \frac{\tilde{R}}{2\sqrt{n}} \left(\frac{3A_{-\log(\alpha)}}{2} + \tilde{R}^{1-2/\log(\alpha)} \tilde{A}_{n,p} \right) \right. \\ &\quad \left. + \frac{\sqrt{0.4}}{2\sqrt{-\log(\alpha)-1}} \times \left(1 - \frac{\tilde{R}\sqrt{-\log(\alpha)-1}}{\sqrt{0.4n}} \right) \left[-\tilde{R}^{-1} \tilde{U}_{n,-\log(\alpha)} \right. \right. \\ &\quad \left. \left. + \frac{\sqrt{-\log(\alpha)+2}\sqrt{e}((2e)^{-1/\log(\alpha)}-1)}{\sqrt{2}} \right] \right. \\ &\quad \left. + \frac{\sqrt{e}(-\log(\alpha)-1)\tilde{R}^2}{1.2n} [e^{0.8} - 1] \left[\frac{2\tilde{R}\tilde{U}_{n,-\log(\alpha)}}{3D_{n,-\log(\alpha)}} + U_{n,-\log(\alpha)} \right. \right. \\ &\quad \left. \left. \times \log \left(\frac{D_{n,-\log(\alpha)}+1}{1-D_{n,-\log(\alpha)}} \right) \right] \right. \\ &\quad \left. + \frac{2\sqrt{e}(-\log(\alpha)-1)\tilde{R}}{\sqrt{1.2n}} [e^{0.8} - 1.8] \left[\frac{\tilde{R}^2}{3} \tilde{U}_{n,-\log(\alpha)} + \tilde{R}U_{n,-\log(\alpha)} \right] \right. \\ &\quad \left. + \frac{\tilde{R}^2(-\log(\alpha)-1)}{0.8nD_{n,-\log(\alpha)}} [U_{n,-\log(\alpha)} + \tilde{U}_{n,-\log(\alpha)}] \right]. \end{aligned}$$

Therefore for any choice of $\alpha \in (0, \delta)$ we have:

$$\mathbb{P}(S_n \geq -\sigma\Phi^{-1}(\delta - \alpha) + g(p_\alpha^*)) \leq \delta.$$

Let $h(\alpha) \triangleq -\sigma z_{\delta-\alpha} + e \frac{K(-\log(\alpha)+2)}{\sqrt{n}}$. We want to minimize $h(\cdot)$. To do so we note that h is smooth and we have:

$$h'(\alpha) = \sqrt{2\pi\sigma^2} e^{\frac{\Phi^{-1}(\delta-\alpha)^2}{2}} - \frac{eK}{\sqrt{n}\alpha}.$$

We propose the following approximate solution to $h'(\alpha) = 0$ and choose $\alpha^* \triangleq \frac{eK}{\sqrt{n}\sigma} \varphi(\Phi^{-1}(\delta))$. We set $A_\delta^n \triangleq \log \left(\frac{\sqrt{n}\sigma}{eK} \varphi(\Phi^{-1}(\delta))^{-1} \right)$; and write

$$q_n(R, \delta, \sigma) \triangleq \sigma z_{\delta - \frac{eK}{\sigma\sqrt{n}} \varphi(\Phi^{-1}(\delta))} + \frac{e[A_\delta^n+2]}{\sqrt{n}} [K + s_{\delta,n}]$$

where we have set

$$\begin{aligned}
\frac{1}{\sigma} s_{n,\delta} &\triangleq \frac{1}{\sqrt{0.4}} \left[\tilde{R} \frac{\tilde{U}_n^\delta}{2\sqrt{n}} + \frac{\tilde{R}}{2\sqrt{n}} \left(\frac{A_\delta^n}{2} + U_n^\delta \right) + \frac{\sqrt{0.4}}{2\sqrt{A_\delta^n-1}} \right. \\
&\quad \times \left(1 - \frac{\tilde{R}\sqrt{A_\delta^n-1}}{\sqrt{0.4n}} \right) \left[\frac{\sqrt{A_\delta^n+2}\sqrt{e}((2e)^{1/A_\delta^n-1})}{\sqrt{2}} - \tilde{R}^{-1}\tilde{U}_n^\delta \right] \\
&\quad + \frac{\sqrt{e}(A_\delta^n-1)\tilde{R}^2}{1.2\sqrt{n}\sqrt{A_\delta^n+2}} [e^{0.8} - 1] \left[\frac{2\tilde{R}\tilde{U}_n^\delta}{3M_n^\delta} + U_n^\delta \log \left(\frac{M_n^\delta+1}{1-M_n^\delta} \right) \right] \\
&\quad + \frac{2\sqrt{e}(A_\delta^n-1)\tilde{R}}{\sqrt{1.2n}\sqrt{A_\delta^n+2}} [e^{0.8} - 1.8] \left[\frac{\tilde{R}^2}{3}\tilde{U}_n^\delta + \tilde{R}U_n^\delta \right] \\
&\quad + \frac{\tilde{R}^2(A_\delta^n-1)}{0.8\sqrt{n(A_\delta^n+2)}M_n^\delta} \left[U_{n,A_\delta^n} + \tilde{U}_{n,A_\delta^n} \right].
\end{aligned}$$

Appendix I. Proof of Thms. 13 and 14: Tighter computable bounds

I.1. Definitions of the Quantities in Sec. 6.3

In section Sec. 6.3, we propose some tighter concentration inequalities. Those bounds are dependent on a quantity $\omega_p^{R,\kappa}$ that we define rigorously in this section. For all choices of $p \geq 2$, $\kappa > 0$ and $\tilde{\sigma} > 0$ we write

$$\tilde{R}_{\tilde{\sigma}} \triangleq R/\tilde{\sigma}.$$

When $\tilde{\sigma} = \sigma$ is equal to the true variance we shorthand $M_{n,\kappa} \triangleq M_{n,\kappa}(\sigma)$.

We define the following candidate for an upper bound to $\mathcal{W}_p(S_n, \mathcal{N}(0, \sigma^2))$:

$$\omega_p^{R,\kappa}(\tilde{\sigma}) \triangleq \tilde{\sigma} \left\{ (a_{1,1}^{\kappa,p,\tilde{R}_{\tilde{\sigma}}}) + (a_{2,1}^{\kappa,p,\tilde{R}_{\tilde{\sigma}}}) + (a_{3,1}^{\kappa,p,\tilde{R}_{\tilde{\sigma}}}) \right\},$$

where $(a_{1,1}^{\kappa,p,\tilde{R}_{\tilde{\sigma}}})$, $(a_{2,1}^{\kappa,p,\tilde{R}_{\tilde{\sigma}}})$ and $(a_{3,1}^{\kappa,p,\tilde{R}_{\tilde{\sigma}}})$ are defined in respectively Eq. (39), Eq. (43) and Eq. (47). We remark that if $\tilde{\sigma} = \sigma$ is the true variance of Y_1 then $\omega_p^{R,\kappa}(\sigma)$ is an upper bound on the Wassertein distance $\mathcal{W}_p(S_n, Z)$ (see Lem. 34 for a proof). Finally, we define:

$$\omega_p^R(\tilde{\sigma}) \triangleq \begin{cases} \inf_{\kappa \in (\frac{R^2}{\tilde{\sigma}^2 n}, \infty)} \omega_p^{R,\kappa}(\tilde{\sigma}) & \text{if } p \geq 2 \\ \frac{\tilde{R}_{\tilde{\sigma}}}{\sqrt{n}} & \text{if } p = 1. \end{cases} \quad (30)$$

I.2. Proof of Thms. 13 and 14

To prove Thm. 13 and Thm. 14 we first establish the following result that any upper bound on the Wassertein distance $\mathcal{W}_p(S_n, \mathcal{N}(0, \sigma^2))$ also yields a tail bound and a quantile bound for S_n .

Lemma 33 *Let $(X_i)_{i \geq 1}$ be a sequence of random variable satisfying the conditions of Thm. 13. For all $p \geq 2$ let ω_p be an upper bound to $\mathcal{W}_p(S_n, \mathcal{N}(0, \sigma^2)) \leq \omega_p$ then the following holds*

$$\begin{aligned}
P(S_n \geq t) &\leq \inf_{p \geq 2, \rho \in (0,1)} \left\{ \Phi^c\left(\frac{t\rho}{\sigma}\right) + \frac{\omega_p^p}{(1-\rho)^{p t^p}} \right\}, \\
P(|S_n| \geq t) &\leq \inf_{p \geq 2, \rho \in (0,1)} \left\{ 2\Phi^c\left(\frac{t\rho}{\sigma}\right) + \frac{\omega_p^p}{(1-\rho)^{p t^p}} \right\}.
\end{aligned}$$

Moreover for all $\delta \in (0, 1)$ for all $\rho \in [0, 1]$ if we define $u_\delta^\rho \triangleq \frac{\omega_p}{(\delta(1-\rho))^{1/p}} - \sigma\Phi^{-1}(\delta\rho)$ then we have

$$\mathbb{P}(S_n \geq u_\delta^\rho) \leq \delta;$$

finally if we define $u_\delta^{d,\rho} \triangleq \frac{\omega_p}{(\delta(1-\rho))^{1/p}} - \sigma\Phi^{-1}(\frac{1}{2}\delta\rho)$ then we have

$$\mathbb{P}(|S_n| \geq u_\delta^{d,\rho}) \leq \delta.$$

Proof Choose reals $\rho \in (0, 1)$ and $p \geq 2$. For all $\epsilon > 0$ there is a Gaussian random variable $G \sim N(0, \sigma^2)$ such that $\|S_n - G\|_p \leq \mathcal{W}_p(S_n, N(0, \sigma^2)) + \epsilon$. By the triangle inequality and the union bound we have

$$\begin{aligned} P(S_n \geq t) &= P(S_n - G + G \geq t) \\ &\leq \mathbb{P}(G \geq \rho t) + \mathbb{P}(S_n - G \geq (1 - \rho)t) \\ &\leq \mathbb{P}(G \geq \rho t) + \frac{\|S_n - G\|_p^p}{(1 - \rho)^p t^p} \\ &\leq \mathbb{P}(G \geq \rho t) + \frac{(\mathcal{W}_p(S_n, \mathcal{N}(0, \sigma^2)) + \epsilon)^p}{(1 - \rho)^p t^p} \\ &\leq \Phi^c\left(\frac{\rho t}{\sigma}\right) + \frac{(\mathcal{W}_p(S_n, \mathcal{N}(0, \sigma^2)) + \epsilon)^p}{(1 - \rho)^p t^p}. \end{aligned}$$

Similarly by the triangle inequality we also observe that:

$$\begin{aligned} P(|S_n| \geq t) &= P(|S_n - G + G| \geq t) \\ &\leq \mathbb{P}(|G| \geq \rho t) + \mathbb{P}(|S_n - G| \geq (1 - \rho)t) \\ &\leq \mathbb{P}(|G| \geq \rho t) + \frac{\|S_n - G\|_p^p}{(1 - \rho)^p t^p} \\ &\leq \mathbb{P}(|G| \geq \rho t) + \frac{(\mathcal{W}_p(S_n, \mathcal{N}(0, \sigma^2)) + \epsilon)^p}{(1 - \rho)^p t^p} \\ &\leq 2\Phi^c\left(\frac{\rho t}{\sigma}\right) + \frac{(\mathcal{W}_p(S_n, \mathcal{N}(0, \sigma^2)) + \epsilon)^p}{(1 - \rho)^p t^p}. \end{aligned}$$

As this holds for any arbitrary choice of $\epsilon > 0$ we deduce that

$$P(S_n \geq t) \leq \Phi^c\left(\frac{\rho t}{\sigma}\right) + \frac{\mathcal{W}_p(S_n, \mathcal{N}(0, \sigma^2))^p}{(1 - \rho)^p t^p}.$$

and

$$P(|S_n| \geq t) \leq 2\Phi^c\left(\frac{\rho t}{\sigma}\right) + \frac{\mathcal{W}_p(S_n, \mathcal{N}(0, \sigma^2))^p}{(1 - \rho)^p t^p}.$$

For all $\delta \in (0, 0.5)$ and all $\rho \in (0, 1)$ if we write $u_\delta \triangleq \frac{\mathcal{W}_p(S_n, \mathcal{N}(0, \sigma^2))}{(\delta(1-\rho))^{1/p}} - \sigma\Phi^{-1}(\delta\rho)$. We remark that

$$\begin{aligned} P(S_n \geq u_\delta) &= P(\tilde{S}_n - G + G \geq u_\delta) \\ &\leq P(G \geq -\sigma\Phi^{-1}(\delta\rho)) + \mathbb{P}(S_n - G \geq \frac{\omega_p}{(\delta(1-\rho))^{1/p}}) \\ &\leq \delta\rho + \frac{\delta(1-\rho)(\mathcal{W}_p(S_n, \mathcal{N}(0, \sigma^2)) + \epsilon)^p}{\omega_p^p}. \end{aligned}$$

As this holds for any arbitrary choice of $\epsilon > 0$ we deduce that

$$\begin{aligned} P(S_n \geq u_\delta) &= P(\tilde{S}_n - G + G \geq u_\delta) \\ &\leq P(G \geq -\sigma\Phi^{-1}(\delta\rho)) + \mathbb{P}(S_n - G \geq \frac{\omega_p}{(\delta(1-\rho))^{1/p}}) \\ &\leq \delta\rho + \delta(1 - \rho) = \delta. \end{aligned}$$

In the same way we can prove that if we write $u_\delta^d \triangleq \frac{\mathcal{W}_p(S_n, \mathcal{N}(0, \sigma^2))}{(\delta(1-\rho))^{1/p}} - \sigma \Phi^{-1}(\frac{\delta\rho}{2})$. We remark that

$$\begin{aligned} P(|S_n| \geq u_\delta^d) &= P(\tilde{S}_n - G + G \geq u_\delta^d) \\ &\leq P(|G| \geq -\sigma \Phi^{-1}(\frac{\delta\rho}{2})) + \mathbb{P}(|S_n - G| \geq \frac{\omega_p}{(\delta(1-\rho))^{1/p}}) \\ &\leq \delta\rho + \frac{\delta(1-\rho)(\mathcal{W}_p(S_n, \mathcal{N}(0, \sigma^2)) + \epsilon)^p}{\omega_p^p}. \end{aligned}$$

As this holds for any arbitrary choice of $\epsilon > 0$ we deduce that

$$P(|S_n| \geq u_\delta) \leq \delta\rho + \delta(1-\rho) = \delta.$$

■

We then propose an upper bound for $\mathcal{W}_p(S_n, \mathcal{N}(0, \sigma^2))$.

Lemma 34 *Let $Z \sim \mathcal{N}(0, 1)$ be a standard normal random variable. Let $(X_i)_{i \geq 1}$ be random variables satisfying the conditions of Thm. 11. If the conditions of Thm. 13 holds then we have:*

$$\sigma^{-1} \mathcal{W}_p(S_n, \sigma Z) \leq \inf_{\kappa \geq \frac{\tilde{R}^2}{n}} \left\{ (a_{1,1}^{\kappa,p,\tilde{R}}) + (a_{2,1}^{\kappa,p,\tilde{R}}) + (a_{3,1}^{\kappa,p,\tilde{R}}) \right\}, \quad (31)$$

where $(a_{1,1}^{\kappa,p,\tilde{R}})$, $(a_{2,1}^{\kappa,p,\tilde{R}})$ and $(a_{3,1}^{\kappa,p,\tilde{R}})$ are defined in respectively Eq. (38), Eq. (42) and Eq. (46). Moreover this can be further upper-bounded as

$$\sigma^{-1} \mathcal{W}_p(S_n, \sigma Z) \leq \inf_{\kappa \geq \frac{\tilde{R}^2}{n}} \left\{ (a_{1,2}^{\kappa,p}) + (a_{2,2}^{\kappa,p}) + (a_{3,2}^{\kappa,p}) \right\}. \quad (32)$$

where $(a_{1,2}^{\kappa,p})$, $(a_{2,2}^{\kappa,p})$ and $(a_{3,2}^{\kappa,p})$ are defined in respectively Eq. (39), Eq. (43) and Eq. (47).

Proof As the proof both for Eq. (31) and Eq. (32) are very similar we only present in details the proof for Eq. (31) and when needed we highlight the differences with the proof of Eq. (32) in proof-asides.

We first notice that if $\sigma > 0$ then we have $\mathcal{W}_p(S_n, \sigma Z) \leq \sigma \mathcal{W}_p(\frac{S_n}{\sigma}, Z)$. Therefore we can assume without loss of generality that $\text{Var}(X_1) = \frac{1}{n}$ and that the random variables $(\sqrt{n}X_i)$ are bounded by $\tilde{R} \triangleq \frac{R}{\sigma}$. This notably implies that for all $k \geq 0$ the moments of X_1^k can be upper bounded as

$$\|X_1^k\|_p = (\mathbb{E}(|X_1|^{kp}))^{1/p} \leq (\|X_1\|_\infty^{kp-2} \mathbb{E}(|X_1|^2))^{1/p} \leq \frac{\tilde{R}^k}{\tilde{R}^{2/p} \sqrt{n}^k} \quad (33)$$

$$\|X_1^k \mathbb{I}(|X_1| \leq c(t))\|_p \leq (\|X_1 \mathbb{I}(|X_1| \leq c(t))\|_\infty^{kp-2} \mathbb{E}(|X_1|^2))^{1/p} \leq n^{-1/p} c(t)^{1-2/p}.$$

where $I \sim \text{unif}\{1, \dots, n\}$ Let $\kappa \in \mathbb{R}^+$ be a positive real that satisfies $\kappa \geq \frac{\tilde{R}^2}{n}$. We set

$$S'_{n,t} \triangleq S_n + (X'_I - X_I) \mathbb{I}(|X'_I|, |X_I| \leq \sqrt{\kappa(1 - e^{-2t})})$$

where $I \sim \text{unif}\{1, \dots, n\}$ and shorthand

$$Y_t = S_n - S'_{n,t}.$$

Let $h_k(x) \triangleq e^{x^2/2} \frac{\partial^k}{\partial x} e^{-x^2/2}$ designate the Hermite polynomials, and denote the $H_k \triangleq h_k(Z)$ where $Z \sim \mathcal{N}(0, 1)$ is a standard normal. By slightly modifying Theorem 5 of [Bonis \(2015\)](#) we have:

$$\begin{aligned} \mathcal{W}_p(S_n, Z) &\leq \int_0^\infty e^{-t} \|n\mathbb{E}(Y_t|S_n) - S_n\|_p dt \\ &\quad + \int_0^\infty \frac{e^{-2t} \|H_1\|_p}{\sqrt{1-e^{-2t}}} \left\| \frac{n}{2} \mathbb{E}((Y_t)^2|S_n) - 1 \right\|_p dt \\ &\quad + \sum_{k \geq 3} \int_0^\infty \frac{e^{-kt} \|H_{k-1}\|_p}{k!(\sqrt{1-e^{-2t}})^{k-1}} n \|\mathbb{E}((Y_t)^k|S_n)\|_p dt \\ &\leq (a_1) + (a_2) + (a_3). \end{aligned}$$

We bound each term, (a_1) (a_2) and (a_3) , successively. For ease of notation we write $c(t) \triangleq \sqrt{\kappa(1 - e^{-2t})}$. Moreover for all $p \geq 2$, and $\kappa > \frac{\tilde{R}^2}{n}$ we write

$$M_{n,\kappa} \triangleq \sqrt{1 - \frac{R^2}{n\sigma^2\kappa}}, \quad \tilde{M}_{n,\kappa} \triangleq \sqrt{1 - \frac{1}{n\kappa}}.$$

Moreover we set

$$\begin{aligned} K_{n,\kappa,p,1} &\triangleq \tilde{M}_{n,\kappa} {}_2F_1\left(\frac{1}{2}, \frac{1}{p} - \frac{1}{2}, \frac{3}{2}, \tilde{M}_{n,\kappa}^2\right) - M_{n,\kappa} {}_2F_1\left(\frac{1}{2}, \frac{1}{p} - \frac{1}{2}, \frac{3}{2}, M_{n,\kappa}^2\right) \\ K_{n,\kappa,p,2} &\triangleq \cos^{-1}(\tilde{M}_{n,\kappa}) - \frac{\tilde{M}_{n,\kappa}}{\sqrt{n\kappa}}, \quad K_{n,\kappa,p,2} \triangleq \cos^{-1}(M_{n,\kappa}) - \frac{M_{n,\kappa}}{\sqrt{n\kappa}} \\ \mu_{n,\kappa,1} &:= \min(c(t), \frac{1}{\sqrt{n}}), \quad \mu_{n,p,\kappa,2} \triangleq \min(n^{-\frac{1}{p}} c(t)^{1-\frac{2}{p}}, c(t)) \\ \mu_{n,p,\kappa,3} &\triangleq \min(c(t)^{1-\frac{2}{p}} n^{-1/p}, \sqrt{n} c(t)^2) \end{aligned}$$

Define (Z_i^t) as

$$Z_i^t \triangleq \begin{cases} -X_i & \text{if } |X_i| \geq c(t) \\ \mathbb{E}(X_i \mathbb{I}(|X_i| > c(t))) - X_i \mathbb{P}(|X_i| > c(t)) & \text{otherwise} \end{cases}.$$

Then we have $n\mathbb{E}(Y_t|S_n) - S_n = \sum_{i \leq n} Z_i^t$. Suppose that $t \leq -\frac{1}{2} \log(1 - \frac{\tilde{R}^2}{n\kappa})$ then $c(t) \leq \tilde{R}/\sqrt{n}$. Therefore using the triangle inequality we note that the norm of the variables (Z_i^t) are bounded by

$$\begin{aligned} \|Z_i^t\|_p &\leq \|X_i\|_p + \|X_i \mathbb{I}(|X_i| \leq c(t))\|_p \stackrel{(a)}{\leq} \frac{1}{\sqrt{n}} \frac{\tilde{R}}{\tilde{R}^{2/p}} + \mu_{n,p,\kappa,2}; \\ \|Z_i^t\|_2 &\leq \|X_i\|_2 + \|X_i \mathbb{I}(|X_i| \leq c(t))\|_2 \stackrel{(a)}{\leq} \frac{1}{\sqrt{n}} + \mu_{n,\kappa,1}, \end{aligned}$$

where (a) is obtained using Eq. (33). Moreover according to Lem. 40 we have

$$\begin{aligned} &\|n\mathbb{E}(Y_t|S_n) - S_n\|_p \\ &= A_p(1 + \min(1, \sqrt{n}c(t))) + \tilde{A}_{n,p}(\tilde{R} + \tilde{R}^{2/p} \sqrt{n} \mu_{n,p,\kappa,2}). \end{aligned} \tag{34}$$

Moreover according to Lem. 39 we have

$$\|n\mathbb{E}(Y_t|S_n) - S_n\|_p \leq \sqrt{p-1} \left[\frac{\tilde{R}}{\tilde{R}^{2/p}} + \sqrt{n} \mu_{n,p,\kappa,2} \right]. \tag{35}$$

By combining Eq. (34) and Eq. (35) we obtain

$$\|n\mathbb{E}(Y_t|S_n) - S_n\|_p \leq \min \left\{ \sqrt{p-1} \left[\frac{\tilde{R}}{\tilde{R}^{2/p}} + \sqrt{n}\mu_{n,p,\kappa,2} \right] \right. \\ \left. A_p(1 + \sqrt{n}\mu_{n,\kappa,1}) + \tilde{A}_{n,p}(\tilde{R} + \tilde{R}^{2/p}\sqrt{n}\mu_{n,p,\kappa,2}). \right. \quad (36)$$

Moreover, if $t \geq -\frac{1}{2} \log(1 - \frac{\tilde{R}^2}{n\kappa}) \geq 0$ then $c(t) \geq \tilde{R}/\sqrt{n}$ therefore $Y_t = X_I - X'_I$. As I is a randomly drawn index in $\{1, \dots, n\}$, this implies that

$$\mathbb{E}(Y_t|S_n) = \frac{1}{n} \sum_{i \leq n} \mathbb{E}(X_i - X'_i|S_n) = \frac{1}{n} \sum_{i \leq n} X_i - \mathbb{E}(X_i) = \frac{1}{n} S_n; \quad (37)$$

which implies that $\|n\mathbb{E}(Y_t|S_n) - S_n\|_p = 0$.

Combining Eq. (36) and Eq. (37) we obtain that we can upper (a_1) as:

$$\begin{aligned} (a_1) &\leq \int_0^\infty e^{-t} \|n\mathbb{E}(Y_t|S_n) - S_n\|_p dt \\ &\stackrel{(a)}{\leq} U_{n,p} \int_0^{-\frac{1}{2} \log(1 - \frac{\tilde{R}^2}{n\kappa})} e^{-t} dt \\ &\quad + \tilde{R}^{2/p} \sqrt{n\kappa}^{1-\frac{2}{p}} \tilde{A}_{n,p} \int_{-\frac{1}{2} \log(1 - \frac{\tilde{R}^2}{n\kappa})}^{-\frac{1}{2} \log(1 - \frac{1}{n\kappa})} e^{-t} \sqrt{1 - e^{-2t}}^{1-\frac{2}{p}} dt \\ &\quad + \sqrt{n\kappa} U_{n,p}^* \int_0^{-\frac{1}{2} \log(1 - \frac{1}{n\kappa})} e^{-t} \sqrt{1 - e^{-2t}} dt + A_p \int_{-\frac{1}{2} \log(1 - \frac{\tilde{R}^2}{n\kappa})}^{-\frac{1}{2} \log(1 - \frac{1}{n\kappa})} e^{-t} dt \\ &= \tilde{R}^{2/p} \sqrt{n\kappa}^{1-\frac{2}{p}} \tilde{A}_{n,p} K_{n,\kappa,p,1} + U_{n,p}(1 - M_{n,\kappa}) \\ &\quad + \frac{\sqrt{n\kappa} U_{n,p}^*}{2} K_{n,\kappa,p,2} + A_p(\tilde{M}_{n,\kappa} - M_{n,\kappa}), \end{aligned}$$

where to obtain (a) we used the fact that $c(t) = \sqrt{\kappa} \sqrt{1 - e^{-2t}}$.

We remark that (a_1) can be alternatively upper bounded as

$$\begin{aligned} (a_1) &\leq \int_0^\infty e^{-t} \|n\mathbb{E}(Y_t|S_n) - S_n\|_p dt \\ &\leq \frac{\tilde{R} \sqrt{p-1}}{\tilde{R}^{2/p}} \int_0^{-\frac{1}{2} \log(1 - \frac{\tilde{R}^2}{n\kappa})} e^{-t} dt \\ &\quad + \sqrt{p-1} \sqrt{\kappa n}^{1-\frac{2}{p}} \int_{-\frac{1}{2} \log(1 - \frac{\tilde{R}^2}{n\kappa})}^{-\frac{1}{2} \log(1 - \frac{1}{n\kappa})} e^{-t} \sqrt{1 - e^{-2t}}^{1-\frac{2}{p}} dt \\ &\quad + \sqrt{p-1} \sqrt{\kappa n} \int_0^{-\frac{1}{2} \log(1 - \frac{1}{n\kappa})} e^{-t} \sqrt{1 - e^{-2t}} dt \\ &= \sqrt{p-1} \left\{ \sqrt{\kappa n}^{1-\frac{2}{p}} K_{n,\kappa,p,1} + \frac{\tilde{R}}{\tilde{R}^{2/p}} (1 - M_{n,\kappa}) + \frac{\sqrt{\kappa n}}{2} K_{n,\kappa,p,2} \right\}. \end{aligned}$$

Therefore we have

$$\begin{aligned} (a_1) &\leq (a_{1,1}^{\kappa,p,\tilde{R}}) \triangleq \min \left\{ \begin{aligned} &\tilde{R}^{2/p} \sqrt{n\kappa}^{1-\frac{2}{p}} \tilde{A}_{n,p} K_{n,\kappa,p,1} + U_{n,p}(1 - M_{n,\kappa}) \\ &+ \frac{\sqrt{n\kappa} U_{n,p}^*}{2} K_{n,\kappa,p,2} + A_p(\tilde{M}_{n,\kappa} - M_{n,\kappa}) \end{aligned} \right. \\ &\quad \left. \sqrt{p-1} \left\{ \sqrt{\kappa n}^{1-\frac{2}{p}} K_{n,\kappa,p,1} + \frac{\tilde{R}}{\tilde{R}^{2/p}} (1 - M_{n,\kappa}) + \frac{\sqrt{\kappa n}}{2} K_{n,\kappa,p,2} \right\} \right\}. \end{aligned} \quad (38)$$

We remark that $(a_{1,1}^{\kappa,p,\tilde{R}})$ can be further upper bounded by a more compact expression. In this goal we can remark that

$$\begin{aligned}
(a_1) &\leq \int_0^\infty e^{-t} \|n\mathbb{E}(Y_t|S_n) - S_n\|_p dt \\
&\stackrel{(a)}{\leq} (\tilde{R}\tilde{A}_{n,p} + 2A_p) \int_0^{-\frac{1}{2}\log(1-\frac{\tilde{R}^2}{n\kappa})} e^{-t} dt \\
&\quad + \tilde{R}^{2/p} \sqrt{n\kappa} \tilde{A}_{n,p} \int_0^{-\frac{1}{2}\log(1-\frac{\tilde{R}^2}{n\kappa})} e^{-t} \sqrt{1-e^{-2t}} dt \\
&= \left[A_p + U_{n,p} \right] (1 - M_{n,\kappa}) + \frac{\tilde{R}^{2/p} \sqrt{n\kappa} \tilde{A}_{n,p}}{2} K_{n,\kappa,p,3} \\
&\leq \frac{\tilde{R}^2}{2n\kappa M_{n,\kappa}} \left[U_{n,p} + \tilde{U}_{n,p} \right].
\end{aligned}$$

Therefore (a_1) can be further upper-bounded by the more compact bound:

$$(a_1) \leq (a_{1,2}^{\kappa,p}) \triangleq \frac{\tilde{R}^2}{2n\kappa M_{n,\kappa}} \left[U_{n,p} + \tilde{U}_{n,p} \right]. \quad (39)$$

We now work on upper bounding (a_2) . In this goal, we first remark that if $t \geq -\frac{1}{2}\log(1 - \frac{\tilde{R}^2}{n\kappa})$ we have $Y_t = X'_I - X_I$ which implies that:

$$\|\frac{n}{2}\mathbb{E}((Y_t)^2|S_n) - 1\|_p \leq \|\frac{1}{2}\sum_{i \leq n} \mathbb{E}(X_i^2) + X_i^2 - 1\|_p \leq \frac{1}{2}\|\sum_{i \leq n} X_i^2 - 1\|_p.$$

We remark using Eq. (33) that $\|X_i^2 - 1/n\|_p \leq \frac{1}{n}(\tilde{R}^{2-2/p} - 1)$ and $\|X_i^2 - 1/n\|_2 \leq \frac{1}{n}(\tilde{R} - 1)$. Therefore using Lem. 39 and Lem. 40 we obtain:

$$\|\frac{n}{2}\mathbb{E}((Y_t)^2|S_n) - 1\|_p \leq \min \left\{ \frac{\sqrt{p-1}}{2\sqrt{n}} \left(\frac{\tilde{R}^2}{\tilde{R}^{2/p}} - 1 \right), \frac{1}{2\sqrt{n}} \left((\tilde{R}^2 - \tilde{R}^{2/p}) \tilde{A}_{n,p} + (\tilde{R} - 1) A_p \right) \right\} \triangleq B_{n,p}.$$

This implies that

$$\begin{aligned}
&\int_{-\frac{1}{2}\log(1-\frac{\tilde{R}^2}{n\kappa})}^\infty \|\frac{n}{2}\mathbb{E}((Y_t)^2|S_n) - 1\|_p \frac{e^{-2t}}{\sqrt{1-e^{-2t}}} dt \\
&\leq B_{n,p} \int_{-\frac{1}{2}\log(1-\frac{\tilde{R}^2}{n\kappa})}^\infty \frac{e^{-2t}}{\sqrt{1-e^{-2t}}} dt = B_{n,p} \int_0^{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}} \frac{t}{\sqrt{1-t^2}} dt \\
&= \frac{1}{2\sqrt{n}} \left(1 - \frac{\tilde{R}}{\sqrt{n\kappa}} \right) \min \left\{ \sqrt{p-1} \left(\frac{\tilde{R}^2}{\tilde{R}^{2/p}} - 1 \right), (\tilde{R}^2 - \tilde{R}^{2/p}) \tilde{A}_{n,p} + (\tilde{R} - 1) A_p \right\}.
\end{aligned} \quad (40)$$

We note that this can be further upper-bounded as

$$\begin{aligned}
&\int_{-\frac{1}{2}\log(1-\frac{\tilde{R}^2}{n\kappa})}^\infty \|\frac{n}{2}\mathbb{E}((Y_t)^2|S_n) - 1\|_p \frac{e^{-2t}}{\sqrt{1-e^{-2t}}} dt \\
&\leq \frac{1}{2\sqrt{n}} \left(1 - \frac{\tilde{R}}{\sqrt{n\kappa}} \right) \left[\tilde{R} U_{n,p} - \tilde{U}_{n,p} \right].
\end{aligned}$$

Similarly if $t \leq -\frac{1}{2}\log(1 - \frac{\tilde{R}^2}{n\kappa})$ then we have

$$\begin{aligned}\|\frac{n}{2}\mathbb{E}((Y_t)^2|S_n) - 1\|_p &\leq 1 - \frac{n}{2}\mathbb{E}((Y_t)^2) + \|\frac{n}{2}\mathbb{E}((Y_t)^2|S_n) - \frac{n}{2}\mathbb{E}((Y_t)^2)\|_p \\ &\leq 1 + \|\frac{n}{2}\mathbb{E}((Y_t)^2|S_n) - \frac{n}{2}\mathbb{E}((Y_t)^2)\|_p.\end{aligned}$$

To control the size of this we remark that

$$\begin{aligned}\frac{n}{2}\mathbb{E}((Y_t)^2|S_n) &= \frac{1}{2}\mathbb{E}(X_1^2\mathbb{I}(|X_1| \leq c(t)) \sum_i \mathbb{I}(|X_i| \leq c(t)) + \frac{1}{2}\mathbb{P}(|X_1| \leq c(t)) \sum_i X_i^2\mathbb{I}(|X_i| \leq c(t)) \\ &\quad - \mathbb{E}(X_1\mathbb{I}(|X_1| \leq c(t)) \sum_i X_i\mathbb{I}(|X_i| \leq c(t))).\end{aligned}$$

Therefore according to Lem. 40 we have

$$\begin{aligned}\|\frac{n}{2}\mathbb{E}((Y_t)^2|S_n) - \frac{n}{2}\mathbb{E}((Y_t)^2)\|_p &\leq A_p \left[\frac{1}{2\sqrt{n}} + \frac{\sqrt{n}}{2}\mu_{n,\kappa,1} + \sqrt{n}\mu_{n,\kappa,1}^2 \right] + \tilde{R}^{2/p}\tilde{A}_{n,p} \left[\frac{1}{2\sqrt{n}} \right. \\ &\quad \left. + \min\left(\frac{c(t)^{2-\frac{2}{p}}\sqrt{n}^{1-\frac{2}{p}}}{2}, \frac{c(t)^2\sqrt{n}}{2}\right) + \mu_{n,p,\kappa,3} \right].\end{aligned}$$

Using also Lem. 39 this implies that

$$\begin{aligned}\|\frac{n}{2}\mathbb{E}((Y_t)^2|S_n) - 1\|_p &\leq 1 + \begin{cases} \tilde{R}^{2/p}\tilde{A}_{n,p} \left[\frac{1}{2\sqrt{n}} + \min\left(\frac{c(t)^{2-\frac{2}{p}}\sqrt{n}^{1-\frac{2}{p}}}{2}, \frac{c(t)^2\sqrt{n}}{2}\right) + \mu_{n,p,\kappa,3} \right] \\ + A_p \left[\frac{1}{2\sqrt{n}} + \frac{\sqrt{n}}{2}\mu_{n,\kappa,1} + \sqrt{n}\mu_{n,\kappa,1}^2 \right] \\ \sqrt{p-1} \left[\frac{1}{2\sqrt{n}} + \min\left(\frac{c(t)^{2-\frac{2}{p}}\sqrt{n}}{2n^{1/p}}, \sqrt{n}c(t)^2/2\right) + \mu_{n,p,\kappa,3} \right]. \end{cases}\end{aligned}$$

We note that this can be further upper bound as

$$\|\frac{n}{2}\mathbb{E}((Y_t)^2|S_n) - 1\|_p \leq 1 + \frac{1}{2\sqrt{n}}\tilde{U}_{n,p} + \frac{3}{2}[A_p c(t) + \sqrt{n}c(t)^2 \tilde{R}^{2/p}\tilde{A}_{n,p}].$$

Therefore to upper-bound $\int_0^{-\frac{1}{2}\log(1-\frac{\tilde{R}^2}{n\kappa})} \frac{e^{-2t}}{\sqrt{1-e^{-2t}}} \|\frac{n}{2}\mathbb{E}((Y_t)^2|S_n) - 1\|_p dt$, we remark that the following integrals are upper-bounded as:

$$\int_0^{-\frac{1}{2}\log(1-\frac{\tilde{R}^2}{n\kappa})} \frac{e^{-2t}}{\sqrt{1-e^{-2t}}} dt \leq \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^1 \frac{x}{\sqrt{1-x^2}} dx = \frac{\tilde{R}}{\sqrt{n\kappa}}.$$

Moreover we have

$$\int_{-\frac{1}{2}\log(1-\frac{1}{n\kappa})}^{-\frac{1}{2}\log(1-\frac{\tilde{R}^2}{n\kappa})} \frac{e^{-2t}}{\sqrt{1-e^{-2t}}} dt \leq \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} \frac{x}{\sqrt{1-x^2}} dx = \frac{(\tilde{R}-1)}{\sqrt{n\kappa}}.$$

Furthermore we observe that

$$\int_0^{-\frac{1}{2}\log(1-\frac{1}{n\kappa})} \sqrt{1-e^{-2t}} e^{-2t} dt \leq \frac{1}{2} \int_{1-\frac{1}{n\kappa}}^1 \sqrt{1-x} dx \leq \frac{1}{3(n\kappa)^{\frac{3}{2}}}$$

In addition we have:

$$\begin{aligned} \int_{-\frac{1}{2}\log(1-\frac{\tilde{R}^2}{n\kappa})}^{-\frac{1}{2}\log(1-\frac{1}{n\kappa})} \sqrt{1-e^{-2t}}^{-2/p} e^{-2t} dt &\leq \frac{1}{2} \int_{1-\frac{1}{n\kappa}}^{1-\frac{\tilde{R}^2}{n\kappa}} \sqrt{1-x}^{-2/p} dx \\ &\leq \frac{p}{2(p-1)(n\kappa)^{-1/p+1}} \left((\tilde{R}^2)^{1-1/p} - 1 \right). \end{aligned}$$

And finally we also have

$$\begin{aligned} \int_{-\frac{1}{2}\log(1-\frac{\tilde{R}^2}{n\kappa})}^{-\frac{1}{2}\log(1-\frac{1}{n\kappa})} \sqrt{1-e^{-2t}}^{1-2/p} e^{-2t} dt &\leq \frac{1}{2} \int_{1-\frac{1}{n\kappa}}^{1-\frac{\tilde{R}^2}{n\kappa}} \sqrt{1-x}^{1-2/p} dx \\ &\leq \frac{p}{(3p-2)(n\kappa)^{3/2-1/p}} \left((\tilde{R}^2)^{3/2-1/p} - 1 \right); \end{aligned}$$

and

$$\int_{-\frac{1}{2}\log(1-\frac{\tilde{R}^2}{n\kappa})}^{-\frac{1}{2}\log(1-\frac{1}{n\kappa})} e^{-2t} dt \leq \frac{1}{2n\kappa} [\tilde{R}^2 - 1].$$

Therefore we obtain that

$$\begin{aligned} &\int_0^{-\frac{1}{2}\log(1-\frac{\tilde{R}^2}{n\kappa})} \frac{e^{-2t}}{\sqrt{1-e^{-2t}}} \left\| \frac{n}{2} \mathbb{E}((Y_t)^2 | S_n) - 1 \right\|_p dt \\ &\leq \min \left\{ \begin{aligned} &\frac{\tilde{R}}{\sqrt{n\kappa}} \left(1 + \frac{U_{n,p}^*}{2\sqrt{n}} \right) + \frac{1}{n\sqrt{\kappa}} \left\{ \frac{U_{n,p}^*}{2} + \frac{p\tilde{R}^{2/p}\tilde{A}_{n,p}}{2(p-1)} \left((\tilde{R}^2)^{1-1/p} - 1 \right) \right. \\ &\quad \left. + \frac{\tilde{R}^{2/p}\tilde{A}_{n,p}}{2} \frac{p}{(3p-2)} \left((\tilde{R}^2)^{3/2-1/p} - 1 \right) + A_p(\tilde{R} - 1) + \frac{A_p}{4} [\tilde{R}^2 - 1] \right\} \\ &\frac{\tilde{R}}{\sqrt{n\kappa}} \left(1 + \frac{\sqrt{p-1}}{2\sqrt{n}} \right) + \frac{1}{n\sqrt{\kappa}} \left\{ \frac{\sqrt{p-1}}{2} \frac{1}{n\sqrt{\kappa}} + \sqrt{p-1} \frac{p}{2(p-1)} \left((\tilde{R}^2)^{1-1/p} - 1 \right) \right. \\ &\quad \left. + \frac{\sqrt{p-1}}{2} \frac{p}{(3p-2)} \left((\tilde{R}^2)^{3/2-1/p} - 1 \right) \right\}. \end{aligned} \right\} \end{aligned} \quad (41)$$

We can note that similarly we have:

$$\int_0^{-\frac{1}{2}\log(1-\frac{\tilde{R}^2}{n\kappa})} \sqrt{1-e^{-2t}} e^{-2t} dt \leq \frac{\tilde{R}^3}{3(n\kappa)^{\frac{3}{2}}};$$

and

$$\int_0^{-\frac{1}{2}\log(1-\frac{\tilde{R}^2}{n\kappa})} e^{-2t} dt \leq \frac{\tilde{R}^2}{2n\kappa}.$$

Therefore we obtain that $\int_0^{-\frac{1}{2}\log(1-\frac{\tilde{R}^2}{n\kappa})} \frac{e^{-2t}}{\sqrt{1-e^{-2t}}} \left\| \frac{n}{2} \mathbb{E}((Y_t)^2 | S_n) - 1 \right\|_p dt$ can be further upper-bounded as

$$\begin{aligned} &\int_0^{-\frac{1}{2}\log(1-\frac{\tilde{R}^2}{n\kappa})} \frac{e^{-2t}}{\sqrt{1-e^{-2t}}} \left\| \frac{n}{2} \mathbb{E}((Y_t)^2 | S_n) - 1 \right\|_p dt \\ &\leq \frac{\tilde{R}}{\sqrt{n\kappa}} \left(1 + \frac{\tilde{U}_{n,p}}{2\sqrt{n}} \right) + \frac{\tilde{R}^2}{2n\sqrt{\kappa}} \left(\frac{3A_p}{2} + \tilde{R}^{1+2/p} \tilde{A}_{n,p} \right). \end{aligned}$$

Therefore by combining Eq. (41) and Eq. (40) we obtain that

$$(a_2) \leq (a_{2,1}^{\kappa,p,\tilde{R}})$$

where we set

$$(a_{2,1}^{\kappa,p,\tilde{R}}) \triangleq \min \left\{ \begin{aligned} & \sqrt{p-1} \left\{ \frac{\tilde{R}}{\sqrt{n\kappa}} \left(1 + \frac{U_{n,p}^*}{2\sqrt{n}} \right) + \frac{1}{n\sqrt{\kappa}} \left\{ \frac{U_{n,p}^*}{2} + \frac{p\tilde{R}^{2/p}\tilde{A}_{n,p}}{2(p-1)} \left((\tilde{R}^2)^{1-1/p} - 1 \right) \right. \right. \\ & \quad \left. \left. + \frac{\tilde{R}^{2/p}\tilde{A}_{n,p} \left((\tilde{R}^2)^{3/2-1/p} - 1 \right)}{2} - \frac{p}{(3p-2)} + A_p(\tilde{R} - 1) + \frac{A_p}{4} [\tilde{R}^2 - 1] \right\} \right\} \\ & \sqrt{p-1} \left\{ \frac{\tilde{R}}{\sqrt{n\kappa}} \left(1 + \frac{\sqrt{p-1}}{2\sqrt{n}} \right) + \frac{1}{n\sqrt{\kappa}} \left\{ \frac{\sqrt{p-1}}{2} \frac{1}{n\sqrt{\kappa}} + \frac{p \left((\tilde{R}^2)^{1-1/p} - 1 \right)}{2\sqrt{p-1}} \right. \right. \\ & \quad \left. \left. + \frac{\sqrt{p-1}}{2} \frac{p}{(3p-2)} \left((\tilde{R}^2)^{3/2-1/p} - 1 \right) \right\} \right\} \\ & + \frac{\sqrt{p-1}}{2\sqrt{n}} \left(1 - \frac{\tilde{R}}{\sqrt{n\kappa}} \right) \min \left\{ \begin{aligned} & \sqrt{p-1} \left(\frac{\tilde{R}^2}{\tilde{R}^{2/p}} - 1 \right) \\ & (\tilde{R}^2 - \tilde{R}^{2/p}) \tilde{A}_{n,p} + (\tilde{R} - 1) A_p \end{aligned} \right\} \end{aligned} \right. \quad (42)$$

We note that (a_2) can be further upper-bounded as

$$(a_2) \leq (a_{2,2}^{\kappa,p}) \triangleq \frac{\sqrt{p-1}}{\sqrt{n\kappa}} \left[\tilde{R} \left(1 + \frac{\tilde{U}_{n,p}}{2\sqrt{n}} \right) + \frac{\tilde{R}}{2\sqrt{n}} \left(\frac{3A_p}{2} + \tilde{R}^{1+2/p} \tilde{A}_{n,p} \right) \right. \\ \left. + \frac{\sqrt{\kappa}}{2} \left(1 - \frac{\tilde{R}}{\sqrt{n\kappa}} \right) \left[U_{n,p} - \tilde{R}^{-1} \tilde{U}_{n,p} \right] \right] \quad (43)$$

Finally we work on bounding (a_3) . We observe that if we write $Z_i^{t,k} \triangleq \mathbb{E}((X_i - X'_i)^k \mathbb{I}(|X'_i| \leq c(t)) | X_{1:n}) \mathbb{I}(|X_i| \leq c(t))$ then

$$n\mathbb{E}(Y_t^k | X_{1:n}) = \sum_{i \leq n} Z_i^{t,k}.$$

Let $k \geq 3$ be an *odd* integer. We observe that the random variables $(Z_i^{t,k})$ are symmetric and therefore have a mean of zero. This implies that $n\mathbb{E}(Y_t^k | S_n) = O_p(\frac{1}{\sqrt{n}})$.

To make this clear, we shorthand $m_{k,p}^*(t) \triangleq \min(n^{1-1/p} c(t)^{k-\frac{2}{p}}, nc(t)^k, \frac{1}{\sqrt{n}} c(t)^{k-3} \frac{\tilde{R}^3}{\tilde{R}^{2/p}})$. We note that as $|X_1|^k \mathbb{I}(|X_1| \leq c(t)) \leq c(t)^{k-3} \min(c(t)^{3-\frac{2}{p}} |X_1|^{2/p}, |X_1|^3, c(t)^3)$ then we have

$$\|Z_i^{t,k}\|_p \leq 2^k \|X_1^k \mathbb{I}(|X_1| \leq c(t))\|_p \leq \frac{2^k}{n} m_{k,p}^*(t).$$

Similarly if we write $\tilde{m}_{k,p}(t) \triangleq c(t)^{k-2} \min(\sqrt{n} c(t), nc(t)^2, \tilde{R})$ then we also have

$$\|Z_i^{t,k}\|_2 \leq \frac{2^k \tilde{m}_{k,p}(t)}{n}.$$

Therefore by using Lem. 39 and Lem. 40 we obtain that for all $t > 0$:

$$n\|\mathbb{E}(Y_t^k | S_n)\|_p \leq \frac{2^k}{\sqrt{n}} \min \left\{ \begin{aligned} & \sqrt{p-1} m_{k,p}^*(t) \\ & m_{k,p}^*(t) \tilde{A}_{n,p} \tilde{R}^{2/p} + \tilde{m}_{k,p}(t) A_p \end{aligned} \right\}.$$

This implies that if $k \geq 3$ is odd we have

$$\begin{aligned}
& \int_0^\infty \frac{e^{-tk}}{\sqrt{1-e^{-2t^{k-1}}}} n \|\mathbb{E}((Y_t)^k | S_n)\|_p dt \\
& \leq \sqrt{n} 2^k \tilde{R}^{2/p} \tilde{A}_{n,p} n^{-1/p} \int_{-\frac{1}{2} \log(1-\frac{\tilde{R}^2}{n\kappa})}^{-\frac{1}{2} \log(1-\frac{1}{n\kappa})} \frac{e^{-tk} c(t)^{k-\frac{2}{p}}}{\sqrt{1-e^{-2t^{k-1}}}} dt \\
& + \sqrt{n} 2^k (\tilde{R}^{2/p} \tilde{A}_{n,p} + A_p) \int_0^{-\frac{1}{2} \log(1-\frac{1}{n\kappa})} \frac{e^{-tk} c(t)^k}{\sqrt{1-e^{-2t^{k-1}}}} dt \\
& + \frac{2^k \tilde{R}^2 (\tilde{R} \tilde{A}_{n,p} + A_p)}{n} \int_{-\frac{1}{2} \log(1-\frac{\tilde{R}^2}{n\kappa})}^\infty \frac{e^{-tk} c(t)^{k-3}}{\sqrt{1-e^{-2t^{k-1}}}} dt \\
& + 2^k A_p \int_{-\frac{1}{2} \log(1-\frac{\tilde{R}^2}{n\kappa})}^{-\frac{1}{2} \log(1-\frac{1}{n\kappa})} \frac{e^{-tk} c(t)^{k-1}}{\sqrt{1-e^{-2t^{k-1}}}} dt.
\end{aligned}$$

We note that this alternatively can be upper-bounded as

$$\begin{aligned}
& \int_0^\infty \frac{e^{-tk}}{\sqrt{1-e^{-2t^{k-1}}}} n \|\mathbb{E}((Y_t)^k | S_n)\|_p dt \\
& \leq \sqrt{n} 2^k (\tilde{R}^{2/p} \tilde{A}_{n,p} + A_p) \int_0^{-\frac{1}{2} \log(1-\frac{\tilde{R}^2}{n\kappa})} \frac{e^{-tk} c(t)^k}{\sqrt{1-e^{-2t^{k-1}}}} dt \\
& + \frac{2^k \tilde{R}^2 (\tilde{R} \tilde{A}_{n,p} + A_p)}{n} \int_{-\frac{1}{2} \log(1-\frac{\tilde{R}^2}{n\kappa})}^\infty \frac{e^{-tk} c(t)^{k-3}}{\sqrt{1-e^{-2t^{k-1}}}} dt.
\end{aligned}$$

To bound this we want to control $\sqrt{(k-1)!}$. In this goal note that

$$\begin{aligned}
\sqrt{(2m+1)!} &= \sqrt{m! \prod_{i=m+1}^{2m+1} i} \geq \sqrt{(2m)!} \sqrt{2m+1} \geq m! 2^m \sqrt{\frac{2(2m+1)}{(m+1)(1+\frac{1}{m})^m}} \\
&\geq m! 2^m \sqrt{\frac{2(2m+1)}{e(m+1)}}.
\end{aligned} \tag{44}$$

We therefore note that

$$\begin{aligned}
& \sum_{\substack{k \geq 3 \\ k \text{ odd}}} \frac{\sqrt{p-1}^{k-1} 2^k x^{k-1} \sqrt{\kappa}^k}{k \sqrt{(k-1)!}} \\
& \leq \sqrt{2e\kappa} \sum_{k \geq 1} \frac{(p-1)^k 2^k x^{2k} \kappa^k \sqrt{k+1}}{(2k+1)} \frac{1}{k!} \\
& \leq \frac{2\sqrt{e\kappa}}{3} \left[e^{2(p-1)x^2\kappa} - 1 \right].
\end{aligned}$$

Therefore we have:

$$\begin{aligned}
& \sum_{k \geq 3} \frac{\sqrt{p-1}^{k-1}}{k \sqrt{(k-1)!}} \int_0^\infty \frac{e^{-tk}}{\sqrt{1-e^{-2t^{k-1}}}} n \|\mathbb{E}((Y_t)^k | S_n)\|_p dt \\
& \leq \frac{2\sqrt{e}}{3} \sqrt{\kappa}^{1-\frac{2}{p}} \sqrt{n} \tilde{R}^{2/p} \tilde{A}_{n,p} n^{-1/p} \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} \sqrt{1-x^2}^{1-\frac{2}{p}} [e^{2(p-1)x^2\kappa} - 1] dx \\
& + \frac{2\sqrt{e}\sqrt{\kappa}}{3} \sqrt{n} U_{n,p}^* \int_{\sqrt{1-\frac{1}{n\kappa}}}^1 \sqrt{1-x^2} [e^{2(p-1)x^2\kappa} - 1] dx \\
& + \frac{2\sqrt{e}\tilde{R}^2 U_{n,p}}{3\kappa n} \int_0^{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}} \frac{[e^{2(p-1)x^2\kappa} - 1]}{1-x^2} dx + \frac{2\sqrt{e}A_p}{3} \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} [e^{2(p-1)x^2\kappa} - 1] dx.
\end{aligned}$$

Alternatively, we remark that

$$\begin{aligned}
& \sum_{k \geq 3} \frac{\sqrt{p-1}^{k-1}}{k\sqrt{k-1}!} \int_0^\infty \frac{e^{-tk}}{\sqrt{1-e^{-2t}k-1}} n \|\mathbb{E}((Y_t)^k | S_n)\|_p dt \\
& \leq \frac{2\sqrt{e}\sqrt{p-1}\sqrt{n}^{1-\frac{2}{p}}}{3} \sqrt{\kappa}^{1-\frac{2}{p}} \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} \sqrt{1-x^2}^{1-\frac{2}{p}} [e^{2(p-1)x^2\kappa} - 1] dx \\
& + \frac{2\sqrt{e}\sqrt{p-1}\sqrt{\kappa}}{3} \sqrt{n} \int_{\sqrt{1-\frac{1}{n\kappa}}}^1 \sqrt{1-x^2} [e^{2(p-1)x^2\kappa} - 1] dx \\
& + \frac{2\sqrt{e}\tilde{R}^3\sqrt{p-1}}{3\kappa n \tilde{R}^{2/p}} \int_0^{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}} \frac{[e^{2(p-1)x^2\kappa} - 1]}{1-x^2} dx.
\end{aligned}$$

We remark that this can be further upper bounded as

$$\begin{aligned}
& \sum_{k \geq 3} \frac{\sqrt{p-1}^{k-1}}{k\sqrt{(k-1)!}} \int_0^\infty \frac{e^{-tk}}{\sqrt{1-e^{-2t}k-1}} n \|\mathbb{E}((Y_t)^k | S_n)\|_p dt \\
& \leq \frac{2\sqrt{e}\sqrt{\kappa}}{3} \sqrt{n} U_{n,p}^* \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^1 \sqrt{1-x^2} [e^{2(p-1)x^2\kappa} - 1] dx \\
& + \frac{2\sqrt{e}\tilde{R}^2 U_{n,p}}{3\kappa n} \int_0^{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}} \frac{[e^{2(p-1)x^2\kappa} - 1]}{1-x^2} dx \\
& \leq \frac{2\sqrt{e}\tilde{R}^3 \tilde{U}_{n,p}}{9n\kappa M_{n,\kappa}} [e^{2(p-1)\kappa} - 1] + \frac{\sqrt{e}\tilde{R}^2 U_{n,p} [e^{2(p-1)M_{n,\kappa}^2\kappa} - 1]}{3\kappa n} \log \left(\frac{M_{n,\kappa}+1}{1-M_{n,\kappa}} \right) \\
& \leq \frac{\sqrt{e}\tilde{R}^2}{3n\kappa} [e^{2(p-1)\kappa} - 1] \left[\frac{2\tilde{R}\tilde{U}_{n,p}}{3M_{n,\kappa}} + U_{n,p} \log \left(\frac{M_{n,\kappa}+1}{1-M_{n,\kappa}} \right) \right].
\end{aligned}$$

If $k \geq 4$ and is even then for all $t \geq 0$ such that $c(t) \leq \tilde{R}/\sqrt{n}$ then we have

$$\begin{aligned}
n \|\mathbb{E}(Y_t^k | S_n)\|_p & \leq \left\| \sum_{i \leq n} \mathbb{E}[\mathbb{I}(|X_i|, |X'_i| \leq c(t)) (X_i - X'_i)^k | S_n] \right\|_p \\
& \leq \left\| \sum_{i \leq n} \mathbb{E}[\mathbb{I}(|X_i|, |X'_i| \leq c(t)) (X_i - X'_i)^k | S_n] \right. \\
& \quad \left. - \mathbb{E}[\mathbb{I}(|X_i|, |X'_i| \leq c(t)) (X_i - X'_i)^k] \right\|_p \\
& \quad + n \|\mathbb{E}[\mathbb{I}(|X_1|, |X'_1| \leq c(t)) (X_1 - X'_1)^k]\| \\
& \stackrel{(a)}{\leq} 2^k \sqrt{p-1} \min(\sqrt{n}^{1-\frac{2}{p}} c(t)^{k-\frac{2}{p}}, \sqrt{n} c(t)^k) + 2^k \min(c(t)^{k-2}, n c(t)^2),
\end{aligned}$$

where to get (a) we used Lem. 39 and the fact that as $k \geq 4$ is even we have

$$\begin{aligned}
& \|\mathbb{E}[\mathbb{I}(|X_i|, |X'_i| \leq c(t)) (X_i - X'_i)^k] - \mathbb{E}[\mathbb{I}(|X_i|, |X'_i| \leq c(t)) (X_i - X'_i)^k]\|_p \\
& \leq \|\mathbb{E}[\mathbb{I}(|X_i|, |X'_i| \leq c(t)) (X_i - X'_i)^k]\|_p \leq 2^k \|\mathbb{E}[\mathbb{I}(|X_i| \leq c(t)) X_i^k]\|_p.
\end{aligned}$$

Moreover for all $t \geq -\frac{1}{2} \log(1 - \frac{\tilde{R}^2}{n\kappa})$ according to Lem. 39 we also have:

$$\begin{aligned}
n\|\mathbb{E}(Y_t^k|S_n)\|_p &\leq \|\sum_{i \leq n} \mathbb{E}[\mathbb{I}(|X_i|, |X'_i| \leq c(t))(X_i - X'_i)^k | S_n]\|_p \\
&\leq \|\sum_{i \leq n} \mathbb{E}[\mathbb{I}(|X_i|, |X'_i| \leq c(t))(X_i - X'_i)^k | S_n] \\
&\quad - \mathbb{E}[\mathbb{I}(|X_i|, |X'_i| \leq c(t))(X_i - X'_i)^k]\|_p \\
&\quad + n\|\mathbb{E}[\mathbb{I}(|X_1|, |X'_1| \leq c(t))(X_1 - X'_1)^k]\|_p \\
&\leq \frac{\sqrt{p-1}2^k}{\sqrt{n^3}} c(t)^{k-4} \tilde{R}^{4-2/p} + 2^k \frac{\tilde{R}^2}{n} c(t)^{k-4}.
\end{aligned}$$

This directly implies that we obtain

$$\begin{aligned}
&\int_0^\infty \frac{e^{-tk}}{\sqrt{1-e^{-2t}k-1}} n\|\mathbb{E}((Y_t)^k|S_n)\|_p dt \\
&\leq \frac{2^k \sqrt{\kappa}^{k-4} \tilde{R}^2 (\sqrt{p-1} \tilde{R}^{2-2/p} + \sqrt{n})}{\sqrt{n^3}} \int_{-\frac{1}{2} \log(1 - \frac{\tilde{R}^2}{n\kappa})}^\infty \frac{e^{-tk}}{\sqrt{1-e^{-2t}k-1}} dt \\
&\quad + 2^k \sqrt{p-1} \sqrt{\kappa}^{k-\frac{2}{p}} \sqrt{n}^{1-2/p} \int_{-\frac{1}{2} \log(1 - \frac{\tilde{R}^2}{n\kappa})}^\infty e^{-tk} \sqrt{1-e^{-2t}k-1}^{1-\frac{2}{p}} dt \\
&\quad + 2^k \sqrt{\kappa}^k \sqrt{n} (\sqrt{n} + \sqrt{p-1}) \int_0^{-\frac{1}{2} \log(1 - \frac{1}{n\kappa})} e^{-tk} \sqrt{1-e^{-2t}k-1} dt \\
&\quad + 2^k \sqrt{\kappa}^{k-2} \int_{-\frac{1}{2} \log(1 - \frac{1}{n\kappa})}^{-\frac{1}{2} \log(1 - \frac{\tilde{R}^2}{n\kappa})} \frac{e^{-tk}}{\sqrt{1-e^{-2t}k-1}} dt.
\end{aligned}$$

To propose a good upper bound to this we first re-express $\sqrt{(k-1)!}$.

We note that for all $m \in \mathbb{N} \setminus \{0\}$ we have

$$\sqrt{2m!} \geq \sqrt{m! \Pi_{i=m+1}^{2m}} i. \quad (45)$$

Hence $\frac{m!}{\sqrt{2m!}} \leq \sqrt{\Pi_{i=m+1}^{2m} \frac{i-m}{i}}$. By exploiting the monotony of $x \rightarrow \frac{m}{x}$ we obtain;

$$\begin{aligned}
\log\left(\frac{\Pi_{i=m+1}^{2m}}{m!}\right) &= \sum_{i=1}^m \log\left(1 + \frac{m}{i}\right) \\
&\geq \sum_{i=1}^{m-1} \int_i^{i+1} \log\left(1 + \frac{m}{x}\right) dx + \log(2) \\
&= \int_1^m \log\left(1 + \frac{m}{x}\right) dx + \log(2) \\
&\geq (2m+1) \log(2) - \log(m+1) - m \log\left(1 + \frac{1}{m}\right).
\end{aligned}$$

This directly implies that $\sqrt{2m!} \geq m! 2^m \sqrt{\frac{2}{(m+1)(1+\frac{1}{m})^m}} \geq m! 2^m \sqrt{\frac{2}{e(m+1)}}$. We remark that

$$\begin{aligned}
\sum_{\substack{k \geq 4 \\ k \text{ even}}} \frac{\sqrt{p-1}^{k-1} 2^k x^{k-1} \sqrt{\kappa}^k}{k \sqrt{(k-1)!}} &\leq \sqrt{2e\kappa} \sqrt{p-1} x \sum_{k \geq 1} \frac{(p-1)^k 2^k x^{2k} \kappa^k}{\sqrt{k+1} \sqrt{2k+1}} \frac{1}{k!} \\
&\leq \frac{2\sqrt{e\kappa} \sqrt{p-1} x}{\sqrt{3}} \left[e^{2(p-1)x^2 \kappa} - 1 - 2(p-1)x^2 \kappa \right].
\end{aligned}$$

Therefore we obtain that:

$$\begin{aligned}
& \sum_{\substack{k \geq 4 \\ \text{even}}} \frac{\sqrt{p-1}^{k-1}}{k\sqrt{(k-1)!}} \int_0^\infty \frac{e^{-tk}}{\sqrt{1-e^{-2t}k-1}} n \|\mathbb{E}((Y_t)^k | S_n)\|_p dt \\
& \leq \frac{2\sqrt{e}\sqrt{p-1}}{\sqrt{3}\kappa} \frac{\tilde{R}^2(\sqrt{p-1}\tilde{R}^{2-2/p} + \sqrt{n})}{\sqrt{n}^3} \int_0^{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}} \frac{x \left[e^{2(p-1)x^2\kappa} - 1 - 2(p-1)x^2\kappa \right]}{\sqrt{1-x^2}^3} dx \\
& + \frac{2\sqrt{e}(p-1)}{\sqrt{3}} \sqrt{\kappa}^{2-\frac{2}{p}} \sqrt{n}^{1-2/p} \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} \sqrt{1-x^2}^{1-\frac{2}{p}} x \left[e^{2(p-1)x^2\kappa} - 1 - 2(p-1)x^2\kappa \right] dx \\
& + \frac{2\sqrt{e}\kappa\sqrt{p-1}}{\sqrt{3}} \sqrt{n}(\sqrt{n} + \sqrt{p-1}) \int_{\sqrt{1-\frac{1}{n\kappa}}}^1 x \left[e^{2(p-1)x^2\kappa} - 1 - 2(p-1)x^2\kappa \right] \sqrt{1-x^2} dx \\
& + \frac{2\sqrt{e}\sqrt{p-1}}{\sqrt{3}} \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} \frac{x \left[e^{2(p-1)x^2\kappa} - 1 - 2(p-1)x^2\kappa \right]}{\sqrt{1-x^2}} dx \\
& \leq \frac{\sqrt{e}\sqrt{p-1}}{\sqrt{3}\kappa} \frac{\tilde{R}^2(\sqrt{p-1}\tilde{R}^{2-2/p} + \sqrt{n})}{\sqrt{n}^3} \int_0^{1-\frac{\tilde{R}^2}{n\kappa}} \frac{\left[e^{2(p-1)x\kappa} - 1 - 2(p-1)x\kappa \right]}{\sqrt{1-x}^3} dx \\
& + \frac{\sqrt{e}}{\sqrt{3}} (p-1) \sqrt{\kappa}^{2-\frac{2}{p}} \sqrt{n}^{1-2/p} \int_{1-\frac{\tilde{R}^2}{n\kappa}}^{1-\frac{1}{n\kappa}} \sqrt{1-x}^{1-\frac{2}{p}} \left[e^{2(p-1)x\kappa} - 1 - 2(p-1)x\kappa \right] dx \\
& + \frac{\sqrt{e}\kappa\sqrt{p-1}}{\sqrt{3}} \sqrt{n}(\sqrt{n} + \sqrt{p-1}) \int_{1-\frac{1}{n\kappa}}^1 \left[e^{2(p-1)x\kappa} - 1 - 2(p-1)x\kappa \right] \sqrt{1-x} dx \\
& + \frac{\sqrt{e}\sqrt{p-1}}{\sqrt{3}} \int_{1-\frac{\tilde{R}^2}{n\kappa}}^{1-\frac{1}{n\kappa}} \frac{\left[e^{2(p-1)x\kappa} - 1 - 2(p-1)x\kappa \right]}{\sqrt{1-x}} dx.
\end{aligned}$$

Similarly using Lem. 40 we have

$$\begin{aligned}
& n \|\mathbb{E}(Y_t^k | S_n)\|_p \\
& \leq \begin{cases} \tilde{A}_{n,p} \tilde{R}^{2/p} 2^k \min(\sqrt{n}^{1-\frac{2}{p}} c(t)^{k-\frac{2}{p}}, \sqrt{n} c(t)^k) + (2c(t))^k \min(c(t)^{-1}, \sqrt{n}) A_p \\ + 2^k \min(c(t)^{k-2}, n c(t)^2). \\ \frac{\tilde{A}_{n,p} 2^k}{\sqrt{n}^3} c(t)^{k-4} \tilde{R}^4 + 2^k \frac{A_p}{\sqrt{n}^3} c(t)^{k-4} \tilde{R}^3 + 2^k c(t)^{k-4} \tilde{R}^2 / n. \end{cases}
\end{aligned}$$

This directly implies that we have:

$$\begin{aligned}
& \sum_{\substack{k \geq 4 \\ \text{even}}} \frac{\sqrt{p-1}^{k-1}}{k\sqrt{(k-1)!}} \int_0^\infty \frac{e^{-tk}}{\sqrt{1-e^{-2t}k-1}} n \|\mathbb{E}((Y_t)^k | S_n)\|_p dt \\
& \leq \frac{2\sqrt{e}\kappa\sqrt{p-1}}{\sqrt{3}} \sqrt{n} (A_p + \tilde{A}_{n,p} \tilde{R}^{2/p} + \sqrt{n}) \\
& \quad \times \int_{\sqrt{1-\frac{1}{n\kappa}}}^1 x \left[e^{2(p-1)x^2\kappa} - 1 - 2(p-1)x^2\kappa \right] \sqrt{1-x^2} dx \\
& + \frac{2\sqrt{e}\sqrt{\kappa}\sqrt{p-1}}{\sqrt{3}} A_p \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} x \left[e^{2(p-1)x^2\kappa} - 1 - 2(p-1)x^2\kappa \right] dx \\
& + \frac{2\sqrt{e}\sqrt{p-1}}{\sqrt{3}} \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} x \frac{\left[e^{2(p-1)x^2\kappa} - 1 - 2(p-1)x^2\kappa \right]}{\sqrt{1-x^2}} dx \\
& + \frac{2\sqrt{e}\sqrt{p-1}}{\sqrt{3}} \tilde{A}_{n,p} \tilde{R}^{2/p} n^{-1/p} \sqrt{\kappa}^{2-\frac{2}{p}} \sqrt{n} \\
& \quad \times \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} x \left[e^{2(p-1)x^2\kappa} - 1 - 2(p-1)x^2\kappa \right] \sqrt{1-x^2}^{1-\frac{2}{p}} dx \\
& + \frac{2\sqrt{e}\sqrt{p-1}}{\sqrt{3}\kappa} \frac{\tilde{R}^2(\sqrt{n} + \tilde{R}A_p + \tilde{R}^2\tilde{A}_{n,p})}{\sqrt{n}^3} \int_0^{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}} x \frac{\left[e^{2(p-1)x^2\kappa} - 1 - 2(p-1)x^2\kappa \right]}{\sqrt{1-x^2}^3} dx \\
& \leq \frac{\sqrt{e}\kappa\sqrt{p-1}}{\sqrt{3}} \sqrt{n} (A_p + \tilde{A}_{n,p} \tilde{R}^{2/p} + \sqrt{n}) \int_{1-\frac{1}{n\kappa}}^1 \left[e^{2(p-1)x\kappa} - 1 - 2(p-1)x\kappa \right] \sqrt{1-x} dx \\
& + \frac{\sqrt{e}\sqrt{\kappa}\sqrt{p-1}}{\sqrt{3}} A_p \int_{1-\frac{\tilde{R}^2}{n\kappa}}^{1-\frac{1}{n\kappa}} \left[e^{2(p-1)x\kappa} - 1 - 2(p-1)x\kappa \right] dx \\
& + \frac{\sqrt{e}\sqrt{p-1}}{\sqrt{3}} \int_{1-\frac{\tilde{R}^2}{n\kappa}}^{1-\frac{1}{n\kappa}} \frac{\left[e^{2(p-1)x\kappa} - 1 - 2(p-1)x\kappa \right]}{\sqrt{1-x}} dx \\
& + \frac{\tilde{A}_{n,p} \tilde{R}^{2/p} \sqrt{e(p-1)\kappa}}{\sqrt{3}} \sqrt{n\kappa}^{1-\frac{1}{2p}} \int_{1-\frac{\tilde{R}^2}{n\kappa}}^{1-\frac{1}{n\kappa}} \left[e^{2(p-1)x\kappa} - 1 - 2(p-1)x\kappa \right] \sqrt{1-x}^{1-\frac{2}{p}} dx \\
& + \frac{\sqrt{e}\sqrt{p-1}}{\sqrt{3}\kappa} \frac{\tilde{R}^2(\sqrt{n} + \tilde{R}A_p + \tilde{R}^2\tilde{A}_{n,p})}{\sqrt{n}^3} \int_0^{1-\frac{\tilde{R}^2}{n\kappa}} \frac{\left[e^{2(p-1)x\kappa} - 1 - 2(p-1)x\kappa \right]}{\sqrt{1-x}^3} dx.
\end{aligned}$$

Note that this can be further upper bounded as

$$\begin{aligned}
& \sum_{\substack{k \geq 4 \\ e \vee n}} \frac{\sqrt{p-1}^{k-1}}{k\sqrt{(k-1)!}} \int_0^\infty \frac{e^{-tk}}{\sqrt{1-e^{-2t}k-1}} n \|\mathbb{E}((Y_t)^k | S_n)\|_p dt \\
& \leq \frac{\sqrt{e\kappa}\sqrt{p-1}}{\sqrt{3}} \sqrt{n} (A_p + \tilde{A}_{n,p} \tilde{R}^{2/p}) \int_{1-\frac{\tilde{R}^2}{n\kappa}}^1 \left[e^{2(p-1)x\kappa} - 1 - 2(p-1)x\kappa \right] \sqrt{1-x} dx \\
& + \frac{\sqrt{e\kappa}\sqrt{p-1}}{\sqrt{3}} n \int_{1-\frac{1}{n\kappa}}^1 \left[e^{2(p-1)x\kappa} - 1 - 2(p-1)x\kappa \right] \sqrt{1-x} dx \\
& + \frac{\sqrt{e}\sqrt{p-1}}{\sqrt{3\kappa}} \frac{\tilde{R}^2(\sqrt{n} + \tilde{R}A_p + \tilde{R}^2\tilde{A}_{n,p})}{\sqrt{n^3}} \int_0^{1-\frac{\tilde{R}^2}{n\kappa}} \frac{\left[e^{2(p-1)x\kappa} - 1 - 2(p-1)x\kappa \right]}{\sqrt{1-x^3}} dx \\
& + \frac{\sqrt{e}\sqrt{p-1}}{\sqrt{3}} \int_{1-\frac{\tilde{R}^2}{n\kappa}}^{1-\frac{1}{n\kappa}} \frac{\left[e^{2(p-1)x\kappa} - 1 - 2(p-1)x\kappa \right]}{\sqrt{1-x}} dx \\
& \leq \frac{2\sqrt{e}\sqrt{p-1}}{3\sqrt{n\kappa}\sqrt{3}} \left[e^{2(p-1)\kappa} - 1 - 2(p-1)\kappa \right] \left[\frac{\tilde{R}^3}{\sqrt{n}} \tilde{U}_{n,p} + 1 \right] \\
& + \frac{2\sqrt{e}\sqrt{p-1}}{\sqrt{3}\sqrt{\kappa}} \frac{\tilde{R}(\sqrt{n} + \tilde{R}U_{n,p})}{n} \left[e^{2(p-1)M_{n,\kappa}^2} - 1 - 2(p-1)M_{n,\kappa}^2 \right] \\
& + \frac{2\sqrt{e}\sqrt{p-1}}{\sqrt{3n\kappa}} \left[e^{2(p-1)\tilde{M}_{n,\kappa}^2} - 1 - 2(p-1)\tilde{M}_{n,\kappa}^2 \right] [\tilde{R} - 1] \\
& \leq \frac{2\sqrt{e}\sqrt{p-1}\tilde{R}}{\sqrt{3n\kappa}} \left[e^{2(p-1)\kappa} - 1 - 2(p-1)\kappa \right] \left[\frac{\tilde{R}^2}{3\sqrt{n}} \tilde{U}_{n,p} + 2 + \frac{\tilde{R}U_{n,p}}{\sqrt{n}} - \frac{2}{3\tilde{R}} \right].
\end{aligned}$$

Therefore to conclude we obtain that

$$\begin{aligned}
 (a_3) & \leq (a_{3,1}^{\kappa,p,\tilde{R}}) \triangleq \left\{ \begin{aligned} & \frac{2\sqrt{e}}{3} \sqrt{\kappa}^{1-\frac{2}{p}} \sqrt{n} \tilde{R}^{2/p} \tilde{A}_{n,p} n^{-1/p} \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} \sqrt{1-x^2}^{1-\frac{2}{p}} [e^{2(p-1)x^2\kappa} - 1] dx \\ & + \frac{2\sqrt{e}\sqrt{\kappa}}{3} \sqrt{n} U_{n,p}^* \int_{\sqrt{1-\frac{1}{n\kappa}}}^1 \sqrt{1-x^2} [e^{2(p-1)x^2\kappa} - 1] dx \\ & + \frac{2\sqrt{e}\tilde{R}^2 U_{n,p}}{3\kappa n} \int_0^{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}} \frac{[e^{2(p-1)x^2\kappa} - 1]}{1-x^2} dx \\ & + \frac{2\sqrt{e}A_p}{3} \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} [e^{2(p-1)x^2\kappa} - 1] dx \\ & + \frac{2\sqrt{e}\sqrt{p-1}\sqrt{n}^{1-\frac{2}{p}}}{3} \sqrt{\kappa}^{1-\frac{2}{p}} \int_{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}}^{\sqrt{1-\frac{1}{n\kappa}}} \sqrt{1-x^2}^{1-\frac{2}{p}} [e^{2(p-1)x^2\kappa} - 1] dx \\ & + \frac{2\sqrt{e}\sqrt{p-1}\sqrt{\kappa}}{3} \sqrt{n} \int_{\sqrt{1-\frac{1}{n\kappa}}}^1 \sqrt{1-x^2} [e^{2(p-1)x^2\kappa} - 1] dx \\ & + \frac{2\sqrt{e}\tilde{R}^3\sqrt{p-1}}{3\kappa n \tilde{R}^{2/p}} \int_0^{\sqrt{1-\frac{\tilde{R}^2}{n\kappa}}} \frac{[e^{2(p-1)x^2\kappa} - 1]}{1-x^2} dx \end{aligned} \right. \\
 & + \left\{ \begin{aligned} & \frac{\sqrt{e}\sqrt{p-1}}{\sqrt{3}\kappa} \frac{\tilde{R}^2(\sqrt{p-1}\tilde{R}^{2-2/p} + \sqrt{n})}{\sqrt{n}^3} \int_0^{1-\frac{\tilde{R}^2}{n\kappa}} \frac{[e^{2(p-1)x\kappa} - 1 - 2(p-1)x\kappa]}{\sqrt{1-x^3}} dx \\ & + \frac{\sqrt{e}}{\sqrt{3}} (p-1) \sqrt{\kappa}^{2-\frac{2}{p}} \sqrt{n}^{1-2/p} \int_{1-\frac{\tilde{R}^2}{n\kappa}}^{1-\frac{1}{n\kappa}} \sqrt{1-x}^{1-\frac{2}{p}} [e^{2(p-1)x\kappa} - 1 - 2(p-1)x\kappa] dx \\ & + \frac{\sqrt{e}\kappa\sqrt{p-1}}{\sqrt{3}} \sqrt{n}(\sqrt{n} + \sqrt{p-1}) \int_{1-\frac{1}{n\kappa}}^{1-\frac{\tilde{R}^2}{n\kappa}} [e^{2(p-1)x\kappa} - 1 - 2(p-1)x\kappa] \sqrt{1-x} dx \\ & + \frac{\sqrt{e}\sqrt{p-1}}{\sqrt{3}} \int_{1-\frac{\tilde{R}^2}{n\kappa}}^{1-\frac{1}{n\kappa}} \frac{[e^{2(p-1)x\kappa} - 1 - 2(p-1)x\kappa]}{\sqrt{1-x}} dx \\ & + \frac{\sqrt{e}\kappa\sqrt{p-1}}{\sqrt{3}} \sqrt{n}(A_p + \tilde{A}_{n,p}\tilde{R}^{2/p} + \sqrt{n}) \int_{1-\frac{1}{n\kappa}}^{1-\frac{\tilde{R}^2}{n\kappa}} [e^{2(p-1)x\kappa} - 1 - 2(p-1)x\kappa] \sqrt{1-x} dx \\ & + \frac{\sqrt{e}\sqrt{\kappa}\sqrt{p-1}}{\sqrt{3}} A_p \int_{1-\frac{\tilde{R}^2}{n\kappa}}^{1-\frac{1}{n\kappa}} [e^{2(p-1)x\kappa} - 1 - 2(p-1)x\kappa] dx \\ & + \frac{\sqrt{e}\sqrt{p-1}}{\sqrt{3}} \int_{1-\frac{\tilde{R}^2}{n\kappa}}^{1-\frac{1}{n\kappa}} \frac{[e^{2(p-1)x\kappa} - 1 - 2(p-1)x\kappa]}{\sqrt{1-x}} dx + \frac{\sqrt{e}\sqrt{p-1}}{\sqrt{3}} \tilde{A}_{n,p} \tilde{R}^{2/p} n^{-1/p} \sqrt{\kappa}^{2-\frac{2}{p}} \\ & \sqrt{n} \int_{1-\frac{\tilde{R}^2}{n\kappa}}^{1-\frac{1}{n\kappa}} [e^{2(p-1)x\kappa} - 1 - 2(p-1)x\kappa] \sqrt{1-x}^{1-\frac{2}{p}} dx \\ & + \frac{\sqrt{e}\sqrt{p-1}}{\sqrt{3}\kappa} \frac{\tilde{R}^2(\sqrt{n} + \tilde{R}A_p + \tilde{R}^2\tilde{A}_{n,p})}{\sqrt{n}^3} \int_0^{1-\frac{\tilde{R}^2}{n\kappa}} \frac{[e^{2(p-1)x\kappa} - 1 - 2(p-1)x\kappa]}{\sqrt{1-x^3}} dx. \end{aligned} \right.
 \end{aligned} \tag{46}$$

We remark that (a_3) can be further upper-bounded by

$$\begin{aligned}
 (a_3) & \leq (a_{3,2}^{\kappa,p}) \triangleq \frac{\sqrt{e}\tilde{R}^2}{3n\kappa} [e^{2(p-1)\kappa} - 1] \left[\frac{2\tilde{R}\tilde{U}_{n,p}}{3M_{n,\kappa}} + U_{n,p} \log \left(\frac{M_{n,\kappa}+1}{1-M_{n,\kappa}} \right) \right] \\
 & + \frac{2\sqrt{e}\sqrt{p-1}\tilde{R}}{\sqrt{3n\kappa}} [e^{2(p-1)\kappa} - 1 - 2(p-1)\kappa] \left[\frac{\tilde{R}^2}{3\sqrt{n}} \tilde{U}_{n,p} + 2 + \frac{\tilde{R}U_{n,p}}{\sqrt{n}} - \frac{2}{3\tilde{R}} \right].
 \end{aligned} \tag{47}$$

■

According to Lem. 34 we know that if $p \geq 2$ then $\mathcal{W}_p(S_n, \mathcal{N}(0, \sigma^2)) \leq \omega_p^{R, \kappa}(\sigma)$. Finally, by Chen et al. (2011, Cor. 4.2), we have $\mathcal{W}_1(S_n, \mathcal{N}(0, \sigma^2)) \leq \frac{R}{\sqrt{n}} = \omega_1^R(\sigma)$. Thm. 13 and Thm. 14 directly follow using Lem. 33.

Appendix J. Proof of Thm. 15: Efficient smoothed tail bound

For ease of notations we write $\omega_{1,p,q} = \mathcal{W}_{\frac{2pq}{q-1}}(\tilde{S}_n, Z)$ and $\omega_{2,p,q} = \|Z\|_{\frac{pq}{q-1}}$.

Theorem 35 Assume that the random variables (Y_i) satisfy the conditions of Thm. 15. Let $\hat{Q}_n(u)$ be an arbitrary upper bound for $\mathbb{P}(\tilde{S}_n \geq u) \leq \hat{Q}_n(u)$. Let $D > 0$ be a constant and write $T_n \triangleq -\frac{1}{2} \log(1 - \frac{D}{\sqrt{n}})$. Then the following holds

$$\mathbb{P}(X_{T_n} \geq u) \leq \Phi^c(u) + \int_{T_n}^{\infty} \inf_{p,q \geq 1} \left\{ \hat{Q}_n(u)^{\frac{p-1}{p}} \left\| \frac{e^{-2t}}{1-e^{-2t}}(Z^2 - 1) - \frac{e^{-t}}{\sqrt{1-e^{-2t}}} \tilde{S}_n Z - \tau'_t \right\|_p \right\} dt.$$

where τ'_t is defined in Eq. (14).

Proof Write $\varphi(\cdot)$ the distribution of G and denote P_t the following functional

$$P_t f(x) \triangleq \int_{\mathbb{R}} f(e^{-t}x + \sqrt{1-e^{-2t}}z) d\varphi(z).$$

We write h the density of the distribution of \tilde{S}_n with respect to φ and h_t the density of the distribution of X_t with respect to φ . We know that $h_t(x) = P_t h(x)$. Therefore we have

$$\mathbb{P}(X_t \geq u) = \int_u^{\infty} P_t h(x) d\varphi(x).$$

Moreover we note that by the heat equation we know that:

$$\partial_t P_t h(x) = h_t''(x) - x h_t'(x).$$

This directly implies that

$$\begin{aligned} \mathbb{P}(G \geq u) - \mathbb{P}(X_{T_n} \geq u) &= \int_{T_n}^{\infty} \partial_t \mathbb{P}(X_t \geq u) dt \\ &= \int_{T_n}^{\infty} \int_u^{\infty} h_t''(x) - x h_t'(x) d\varphi(x) dt \\ &= \int_{T_n}^{\infty} \int_u^{\infty} \left[\frac{h_t''(x)}{h_t(x)} - x \frac{h_t'(x)}{h_t(x)} \right] h_t(x) d\varphi(x) dt \\ &= \int_{T_n}^{\infty} \mathbb{E} \left(\mathbb{I}(X_t \geq u) \left[\frac{h_t''(X_t)}{h_t(X_t)} - X_t \frac{h_t'(X_t)}{h_t(X_t)} \right] \right) dt. \end{aligned}$$

We remark that the PDF of X_t is $f_t(x) = h_t(x)\varphi(x)$. Therefore we have

$$S_t(x) = -\frac{h_t'(x)}{h_t(x)} + x.$$

Moreover define $I_t(x) \triangleq \frac{f_t''(x)}{f_t(x)}$. We have

$$\frac{f_t''(x)}{f_t(x)} = \frac{h_t''(x)}{h_t(x)} - 2x \frac{h_t'(x)}{h_t(x)} + x^2 - 1.$$

Therefore we have

$$\begin{aligned}
\partial_t \mathbb{P}(X_t \geq u) &= \mathbb{E} \left(\mathbb{I}(X_t \geq u) \left[I_t(X_t) - 2X_t S_{X_t} - X_t^2 + 1 - X_t(X_t - S_{X_t}(X_t)) \right] \right) \\
&= \mathbb{E} \left(\mathbb{I}(X_t \geq u) \left[I_t(X_t) - X_t S_{X_t} + 1 \right] \right) \\
&= \mathbb{E} \left(\mathbb{I}(X_t \geq u) \left[I_t(X_t) - X_t^2 + 1 - X_t(S_{X_t} - X_t) \right] \right).
\end{aligned}$$

Moreover using the Stein identity we know that

$$S_t(X_t) - X_t = \mathbb{E} \left(\frac{e^{-2t}}{\sqrt{1-e^{-2t}}} Z - e^{-t} \tilde{S}_n | X_t \right).$$

Moreover we note that by the Stein identity we also obtain that

$$I_t(X_t) = \mathbb{E} \left(\frac{1}{1-e^{-2t}} (Z^2 - 1) | X_t \right).$$

This directly implies that

$$\begin{aligned}
&I_t(X_t) - X_t^2 + 1 \\
&= \mathbb{E} \left(\frac{Z^2}{1-e^{-2t}} - X_t^2 | X_t \right) - \frac{e^{-2t}}{1-e^{-2t}} \\
&= \mathbb{E} \left(\frac{Z}{\sqrt{1-e^{-2t}}} \left(\frac{Z}{\sqrt{1-e^{-2t}}} - X_t \right) + X_t \left(\frac{Z}{\sqrt{1-e^{-2t}}} - X_t \right) | X_t \right) - \frac{e^{-2t}}{1-e^{-2t}} \\
&= \mathbb{E} \left(\frac{Z}{\sqrt{1-e^{-2t}}} \left(\frac{e^{-2t}}{\sqrt{1-e^{-2t}}} Z - e^{-t} \tilde{S}_n \right) | X_t \right) + X_t \mathbb{E} \left[\left(\frac{e^{-2t}}{\sqrt{1-e^{-2t}}} Z - e^{-t} \tilde{S}_n \right) | X_t \right] \\
&\quad - \frac{e^{-2t}}{1-e^{-2t}} \\
&= \mathbb{E} \left(\frac{Z}{\sqrt{1-e^{-2t}}} \left(\frac{e^{-2t}}{\sqrt{1-e^{-2t}}} Z - e^{-t} \tilde{S}_n \right) | X_t \right) + X_t (S_{X_t} - X_t) - \frac{e^{-2t}}{1-e^{-2t}}.
\end{aligned}$$

Moreover, we note that

$$\begin{aligned}
&\mathbb{E} \left(\frac{Z}{\sqrt{1-e^{-2t}}} \left(\frac{e^{-2t}}{\sqrt{1-e^{-2t}}} Z - e^{-t} \tilde{S}_n \right) | X_t \right) - \frac{e^{-2t}}{1-e^{-2t}} \\
&= \mathbb{E} \left(\frac{e^{-2t}}{1-e^{-2t}} (Z^2 - 1) - \frac{e^{-t}}{\sqrt{1-e^{-2t}}} \tilde{S}_n Z | X_t \right).
\end{aligned}$$

This directly implies that

$$\begin{aligned}
&I_t(X_t) - X_t^2 + 1 - X_t(S_t(X_t) - X_t) \\
&= \mathbb{E} \left(\frac{Z}{\sqrt{1-e^{-2t}}} \left(\frac{e^{-2t}}{\sqrt{1-e^{-2t}}} Z - e^{-t} \tilde{S}_n \right) | X_t \right) - \frac{e^{-2t}}{1-e^{-2t}}.
\end{aligned}$$

This directly implies that

$$\begin{aligned}
&\mathbb{P}(X_{T_n} \geq u) \\
&\leq \Phi^c(u) + \int_{-\frac{1}{2}}^{\infty} \log(1 - \frac{D}{\sqrt{n}}) |\mathbb{E} \left(\mathbb{I}(X_t \geq u) \left[\mathbb{E} \left(\frac{e^{-2t}}{1-e^{-2t}} (Z^2 - 1) - \frac{e^{-t}}{\sqrt{1-e^{-2t}}} \tilde{S}_n Z | X_t \right) \right] \right) | dt.
\end{aligned}$$

Now we have proven in Lem. 37 that $\mathbb{E}(\tau'_t | X_t) = \mathbb{E}(\tau_t | X_t) = 0$. Therefore

This directly implies that

$$\begin{aligned}
& \mathbb{P}(X_{T_n} \geq u) \\
& \leq \Phi^c(u) + \int_{-\frac{1}{2}}^{\infty} \log(1 - \frac{D}{\sqrt{n}}) |\mathbb{E}(\mathbb{I}(X_t \geq u) \left[\mathbb{E}\left(\frac{e^{-2t}}{1-e^{-2t}}(Z^2 - 1) - \frac{e^{-t}}{\sqrt{1-e^{-2t}}} \tilde{S}_n Z - \tau'_t | X_t \right) \right] | \\
& \stackrel{(a)}{\leq} \Phi^c(u) + \int_{T_n}^{\infty} \inf_{p \geq 2} \mathbb{P}(X_t \geq u) \frac{p-1}{p} \left\{ \left\| \frac{e^{-2t}}{1-e^{-2t}}(Z^2 - 1) - \frac{e^{-t}}{\sqrt{1-e^{-2t}}} \tilde{S}_n Z - \tau'_t \right\|_p \right\} \\
& \leq \Phi^c(u) + \int_{T_n}^{\infty} \inf_{p \geq 2} \hat{Q}_n(u) \frac{p-1}{p} \left\{ \left\| \frac{e^{-2t}}{1-e^{-2t}}(Z^2 - 1) - \frac{e^{-t}}{\sqrt{1-e^{-2t}}} \tilde{S}_n Z - \tau'_t \right\|_p \right\}.
\end{aligned}$$

■

Theorem 36 Assume that the random variables (Y_i) satisfy the conditions of Thm. 15 hold. Let $D > 0$ be the constant set in Thm. 15. Let $\hat{Q}_n(u)$ be an arbitrary upper bound for $\mathbb{P}(\tilde{S}_n \geq u) \leq \hat{Q}_n(u)$. Then the following holds

$$\begin{aligned}
& \mathbb{P}(S_n^{\text{noisy}} \geq \sigma u) \\
& \leq \Phi^c(u) + \inf_{p \geq 2} \hat{Q}_n(u) \frac{p-1}{p} \left\{ \frac{\sqrt{3}e(p-1)\tilde{R}^2(\sqrt{n} + A_p\tilde{R} + \tilde{R}^2\tilde{A}_{n,p})}{\kappa\sqrt{2}\sqrt{n}^3} \right. \\
& \quad \times \int_0^{1-\frac{D}{\sqrt{n}}} \frac{(e^{2(p-1)x(p-1)\kappa} - 1)}{(1-x)^2} dx + \frac{\log(n/D^2)\sqrt{p-1}}{4\sqrt{2}\sqrt{n}} \min \left\{ \begin{aligned} & \sqrt{p-1} \left(\frac{\tilde{R}^2}{\tilde{R}^{2/p}} - 1 \right) \\ & (\tilde{R}^2 - \tilde{R}^{2/p})\tilde{A}_{n,p} \\ & + (\tilde{R} - 1)A_p \end{aligned} \right. \\
& \quad \left. + \frac{2\sqrt{e(p-1)}\tilde{R}^2(A_p + \tilde{R}\tilde{A}_{n,p})}{\sqrt{3}n\kappa} \int_0^{\sqrt{1-\frac{D}{\sqrt{n}}}} \frac{(e^{2x^2(p-1)\kappa} - 1)}{\sqrt{1-x^2}^3} dx \right\} \\
& \lesssim \Phi^c(u) + \inf_{p \geq 2} \frac{\hat{Q}_n(u) \frac{p-1}{p}}{\sqrt{n}} \left\{ \frac{\sqrt{e}\tilde{R}^2(e^{2\kappa} - 1)}{\sqrt{3}D\sqrt{n}\kappa} + \frac{\log(n/D^2)\sqrt{p-1}}{4\sqrt{2}\sqrt{n}} (\tilde{R} - 1)A_p \right\}.
\end{aligned}$$

Proof The first step is to remark that $X_{T_n} \stackrel{d}{=} \tilde{S}_n^{\text{noisy}}$. Therefore without a loss of generality, we obtain a tail bound for X_{T_n} . To do we exploit Thm. 35 which guarantees that

$$\begin{aligned}
\mathbb{P}(X_{T_n} \geq u) & \leq \int_{T_n}^{\infty} \inf_{p,q \geq 1} \left\{ \hat{Q}_n(u) \frac{p-1}{p} \left\| \frac{e^{-2t}}{1-e^{-2t}}(Z^2 - 1) - \frac{e^{-t}}{\sqrt{1-e^{-2t}}} \tilde{S}_n Z - \tau'_t \right\|_p \right\} dt. \\
& \quad + \Phi^c(u).
\end{aligned}$$

Moreover, according to Lem. 19 we have that:

$$\begin{aligned}
& \int_{T_n}^{\infty} \left\| \frac{e^{-2t}}{1-e^{-2t}} (Z^2 - 1) - \frac{e^{-t}}{\sqrt{1-e^{-2t}}} \tilde{S}_n Z - \tau'_t \right\|_p \\
& \leq \frac{\sqrt{3}e(p-1)\tilde{R}^2(\sqrt{n}+A_p\tilde{R}+\tilde{R}^2\tilde{A}_{n,p})}{\kappa\sqrt{2}\sqrt{n}^3} \int_0^{1-\frac{D}{\sqrt{n}}} \frac{(e^{2(p-1)x(p-1)\kappa}-1)}{(1-x)^2} dx \\
& \quad + \frac{2\sqrt{e(p-1)}\tilde{R}^2(A_p+\tilde{R}\tilde{A}_{n,p})}{\sqrt{3}n\kappa} \int_0^{\sqrt{1-\frac{D}{\sqrt{n}}}} \frac{(e^{2x^2(p-1)\kappa}-1)}{\sqrt{1-x^2}^3} dx \\
& \quad + \frac{\log(n/D^2)\sqrt{p-1}}{4\sqrt{2}\sqrt{n}} \min \left\{ \sqrt{p-1} \left(\frac{\tilde{R}^2}{\tilde{R}^{2/p}} - 1 \right) \right. \\
& \quad \left. \left(\tilde{R}^2 - \tilde{R}^{2/p} \right) \tilde{A}_{n,p} + (\tilde{R} - 1)A_p \right\} \\
& \lesssim \frac{\sqrt{e}\tilde{R}^2(e^{2\kappa}-1)}{\sqrt{3}D\sqrt{n}\kappa} + \frac{\log(n/D^2)\sqrt{p-1}}{4\sqrt{2}\sqrt{n}} (\tilde{R} - 1)A_p.
\end{aligned}$$

■

Appendix K. Proof of Thm. 16: Empirical quantile bounds

To simplify our presentation, we introduce the following notation

$$\hat{\mu} \triangleq \frac{1}{n} \sum_{i \leq n} W_i, \quad \hat{m}_2(m) = \frac{1}{n} \sum_{i \leq n} (W_i - m)^2, \quad \hat{m}_3(m) = \frac{1}{n} \sum_{i \leq n} (W_i - m)^3,$$

$$v_4(m, s) \triangleq \frac{1}{n} \sum_{i \leq n} ((W_i - m)^2 - s)^2, \quad s_4^2 = \text{var}(Y_1^2 - \sigma^2),$$

$$\mu \triangleq \mathbb{E}(W_1), \quad \Delta\hat{\mu} = \hat{\mu} - \mu.$$

and note that $\hat{\sigma}^2 = \hat{m}_2(\hat{\mu})$, $\hat{m}_3 = \hat{m}_3(\hat{\mu})$, and $\hat{m}_4 = v_4(\hat{\mu}, \hat{\sigma}^2)$.

Fix any $\delta > 0$ and $a \in (0, \delta)$, and define the event $E_1 \triangleq \{|\frac{1}{n} \sum_{i \leq n} Y_i|^2 \leq c_1^{\delta-a_n}(\sigma)\sigma^2\}$. Since

$$\hat{\sigma}^2 = \hat{m}_2(\hat{\mu}) = \frac{1}{n} \sum_{i \leq n} (W_i - \hat{\mu})^2 = \hat{m}_2(\mu) + \Delta\hat{\mu}^2.$$

we have

$$\sigma^2 = \sigma^2 - \hat{\sigma}^2 + \hat{\sigma}^2 = \hat{\sigma}^2 + \Delta\hat{\mu}^2 + \sigma^2 - \hat{m}_2(\mu). \quad (48)$$

Therefore if E_1 holds we have

$$\Delta\hat{\mu}^2 \leq c_1^{\delta-a_n}(\sigma)[\hat{\sigma}^2 + \Delta\hat{\mu}^2 + \sigma^2 - \hat{m}_2(\mu)].$$

This directly implies that

$$\Delta\hat{\mu}^2 \leq \frac{c_1^{\delta-a_n}(\sigma)}{1-c_1^{\delta-a_n}(\sigma)}[\hat{\sigma}^2 + \sigma^2 - \hat{m}_2(\mu)].$$

We note that $\sigma \rightarrow c_1^{\delta-a_n}(\sigma)$ is a non-increasing function. Therefore for any nonnegative value $c < \sigma$ we have

$$\Delta\hat{\mu}^2 \leq \frac{c_1^{\delta-a_n}(c)}{1-c_1^{\delta-a_n}(c)}[\hat{\sigma}^2 + \sigma^2 - \hat{m}_2(\mu)].$$

We, therefore, want to lower-bound σ with high probability. According to Lem. 41 we have:

$$P(|\hat{m}_2(\mu) - \sigma^2| \geq R^2 q_n^{\text{KS}}(\frac{a_n}{4})) \leq \frac{a_n}{4}.$$

We define the event $E_2 \triangleq \{(|\hat{m}_2(\mu) - \sigma^2| \leq R^2 q_n^{\text{KS}}(\frac{a_n}{4}))\}$ and remark that $\sigma^2 \geq \sigma_{1,a_n}^2$ when E_2 holds by Eq. (48). Therefore we have

$$\Delta \hat{\mu}^2 \leq \frac{c_1^{\delta-a_n}(\sigma_{1,a_n})}{1-c_1^{\delta-a_n}(\sigma_{1,a_n})} [\hat{\sigma}^2 + \sigma^2 - \hat{m}_2(\mu)]. \quad (49)$$

To further upper bound this we want again to upper bound $\sigma^2 - \hat{m}_2(\mu)$. To do so we propose to use Thm. 14. Indeed remark that

$$\hat{m}_2(\mu) - \sigma^2 = \frac{1}{n} \sum_{i \leq n} Y_i^2 - \sigma^2.$$

We also notice that $\|Y_1^2 - \sigma^2\|_\infty \leq R^2$ and that $\text{Var}(Y_1^2 - \sigma^2) = s_4$. Therefore according to Thm. 14 we have:

$$\mathbb{P}(\sigma^2 \leq \hat{m}_2(\mu) + \tilde{q}_n(R^2, \frac{a_n}{4}, s_4)) \geq 1 - \frac{a_n}{4}.$$

We write the event $E_3 \triangleq \{\sigma^2 \leq \hat{m}_2(\mu) + \tilde{q}_n(R^2, \frac{a_n}{4}, s_4)\}$. As s_4 is unknown to be able to use this result we want to bound s_4 . Now we notice that

$$\begin{aligned} v_4(\mu, \hat{m}_2(\mu)) &= \frac{1}{n} \sum_{i \leq n} [(W_i - \mu)^2 - (\frac{1}{n} \sum_{i \leq n} (W_i - \mu)^2)^2] \\ &= v_4(\mu, \sigma^2) - (\sigma^2 - \hat{m}_2(\mu))^2 \\ &= s_4^2 + v_4(\mu, \sigma^2) - s_4^2 - (\sigma^2 - \hat{m}_2(\mu))^2. \end{aligned}$$

This directly implies that

$$s_4^2 = v_4(\mu, \hat{m}_2(\mu)) + (\sigma^2 - \hat{m}_2(\mu))^2 + s_4^2 - v_4(\mu, \sigma^2). \quad (50)$$

Now we will re-express $v_4(\mu, \hat{m}_2(\mu))$ in terms of $v_4(\hat{\mu}, \hat{m}_2(\hat{\mu}))$. In this goal for all $i, j \leq n$ we remember that $Y_i = W_i - \mu$ and write $\Delta \hat{\mu} \triangleq \hat{\mu} - \mu$. We remark that

$$Y_i^2 - Y_j^2 = (Y_i - \Delta \hat{\mu})^2 - (Y_j - \Delta \hat{\mu})^2 + 2\Delta \hat{\mu}(Y_i - Y_j).$$

Therefore we have:

$$\begin{aligned} & \left[(Y_i^2 - Y_j^2)^2 - [(Y_i - \Delta \hat{\mu})^2 - (Y_j - \Delta \hat{\mu})^2]^2 \right] \\ &= \left[(Y_i^2 - Y_j^2) - [(Y_i - \Delta \hat{\mu})^2 - (Y_j - \Delta \hat{\mu})^2] \right] \\ & \times \left[(Y_i^2 - Y_j^2) + [(Y_i - \Delta \hat{\mu})^2 - (Y_j - \Delta \hat{\mu})^2] \right] \\ &= 4\Delta \hat{\mu}^2 (Y_i - Y_j)^2 + 4\Delta \hat{\mu} (Y_i - Y_j) [(Y_i - \Delta \hat{\mu})^2 - (Y_j - \Delta \hat{\mu})^2]. \end{aligned}$$

By traditional arguments, we remark that

$$\begin{aligned} & \frac{1}{2n^2} \sum_{i,j \leq n} (Y_i - Y_j)^2 \\ &= \frac{1}{2n^2} \sum_{i,j \leq n} (Y_i - \Delta \hat{\mu})^2 + (Y_j - \Delta \hat{\mu})^2 - 2(Y_i - \Delta \hat{\mu})(Y_j - \Delta \hat{\mu}) = \hat{m}_2(\hat{\mu}). \end{aligned}$$

where to obtain the last equality we used the fact that

$$\frac{1}{2n^2} \sum_{i,j \leq n} (Y_i - \Delta\hat{\mu})(Y_j - \Delta\hat{\mu}) = 0.$$

Similarly we have

$$\begin{aligned} & \frac{1}{2n^2} \sum_{i,j \leq n} ((Y_i - \Delta\hat{\mu})^2 - (Y_j - \Delta\hat{\mu})^2)(Y_i - \Delta\hat{\mu} - (Y_j - \Delta\hat{\mu})) \\ &= \frac{1}{n} \sum_{i \leq n} (Y_i - \Delta\hat{\mu})^3 = \hat{m}_3(\hat{\mu}). \end{aligned}$$

This directly implies that

$$v_4(\mu, \hat{m}_2(\mu)) - v_4(\hat{\mu}, \hat{m}_2(\hat{\mu})) = 4\hat{m}_2(\hat{\mu})\Delta\hat{\mu}^2 - 4\Delta\hat{\mu}\hat{m}_3(\hat{\mu}). \quad (51)$$

We define $c_3 = 4\hat{m}_2(\hat{\mu})$ and $c_4 = -4\hat{m}_3(\hat{\mu})$ and remark that by combining Eq. (50) and Eq. (51) we have

$$s_4^2 = v_4(\hat{\mu}, \hat{m}_2(\hat{\mu})) + c_3\Delta\hat{\mu}^2 + \Delta\hat{\mu}c_4 + (\sigma^2 - \hat{m}_2(\mu))^2 + s_4^2 - v_4(\mu, \sigma^2).$$

Therefore when the event E_3 holds we obtain that

$$\begin{aligned} & [(\sigma^2 - \hat{m}_2(\mu))\mathbb{I}(E_2)]^2 \\ & \leq c_2^{a_n}(s_4) \left[v_4(\hat{\mu}, \hat{m}_2(\hat{\mu})) + c_3\Delta\hat{\mu}^2 + \Delta\hat{\mu}c_4 + (\sigma^2 - \hat{m}_2(\mu))^2 + s_4^2 - v_4(\mu, \sigma^2) \right]. \end{aligned}$$

This directly implies that

$$\begin{aligned} & [(\sigma^2 - \hat{m}_2(\mu))\mathbb{I}(E_2)]^2 \\ & \leq \frac{c_2^{a_n}(s_4)}{1 - c_2^{a_n}(s_4)} \left[v_4(\hat{\mu}, \hat{m}_2(\hat{\mu})) + c_3\Delta\hat{\mu}^2 + \Delta\hat{\mu}c_4 + s_4^2 - v_4(\mu, \sigma^2) \right] \end{aligned}$$

According to Lem. 41, we have:

$$P(|s_4^2 - v_4(\mu, \sigma^2)| \geq R^4 q_n^{\text{KS}}(\frac{a_n}{4})) \leq \frac{a_n}{4}.$$

We write the event $E_4 \triangleq \{|s_4^2 - v_4(\sigma^2)| \leq R^4 q_n^{\text{KS}}(\frac{a_n}{4})\}$. Then when the event E_3 and E_4 both hold then

$$s_4^2 \geq v_4(\hat{m}_2(\hat{\mu})) + c_3\hat{\mu}^2 + \hat{\mu}c_4 + (\sigma^2 - \hat{m}_2(\mu))^2 - R^4 q_n^{\text{KS}}(\frac{a_n}{4}).$$

When the event $E_5 \triangleq \{|\Delta\hat{\mu}| \leq R q_n^{\text{KS}}(a_n/4)\}$ holds we have the further lower bound

$$s_4^2 \geq s_{1,a_n}^2.$$

We remark that $s \rightarrow c_2^{a_n}(s)$ is a non increasing function therefore we have that under E_3 , E_4 and E_5 we have

$$[(\sigma^2 - \hat{m}_2(\mu))\mathbb{I}(E_2)]^2 \leq \frac{c_2^{a_n}(s_{1,a_n})}{1 - c_2^{a_n}(s_{1,a_n})} \left[v_4(\hat{\mu}, \hat{m}_2(\hat{\mu})) + c_3\Delta\hat{\mu}^2 + \Delta\hat{\mu}c_4 + R^2 q_n^{\text{KS}}(\frac{a_n}{4}) \right].$$

Therefore by combining this with Eq. (49) we obtain that

$$\begin{aligned} & \left[\Delta \hat{\mu}^2 - \frac{c_1^{\delta-an}(\sigma_{1,a_n})}{1-c_1^{\delta-an}(\sigma_{1,a_n})} \hat{\sigma}^2 \right]^2 \\ & \leq \left[\frac{c_1^{\delta-an}(\sigma_{1,a_n})}{1-c_1^{\delta-an}(\sigma_{1,a_n})} \right]^2 \left[\frac{c_2^{a_n}(s_{1,a_n})}{1-c_2^{a_n}(s_{1,a_n})} \left[v_4(\hat{\mu}, \hat{m}_2(\hat{\mu})) + c_3 \Delta \hat{\mu}^2 + \Delta \hat{\mu} c_4 + R^2 q_n^{\text{KS}}\left(\frac{a_n}{4}\right) \right] \right]. \end{aligned} \quad (52)$$

Note that Eq. (52) holds only if

$$\begin{aligned} & \Delta \hat{\mu}^4 + \left[\frac{c_1^{\delta-an}(\sigma_{1,a_n})}{1-c_1^{\delta-an}(\sigma_{1,a_n})} \right]^2 \hat{\sigma}^4 - 2 \Delta \hat{\mu}^2 \hat{\sigma}^2 \frac{c_1^{\delta-an}(\sigma_{1,a_n})}{1-c_1^{\delta-an}(\sigma_{1,a_n})} \\ & \leq \left[\frac{c_1^{\delta-an}(\sigma_{1,a_n})}{1-c_1^{\delta-an}(\sigma_{1,a_n})} \right]^2 \frac{c_2^{a_n}(s_{1,a_n})}{1-c_2^{a_n}(s_{1,a_n})} \left[v_4(\hat{\mu}, \hat{m}_2(\hat{\mu})) + c_3 \Delta \hat{\mu}^2 + c_4 \Delta \hat{\mu} + R^4 q_n^{\text{KS}}\left(\frac{a_n}{4}\right) \right]. \end{aligned}$$

Therefore we have

$$P\left(\Delta \hat{\mu} \in A_{\delta,a}^n\right) \leq \mathbb{P}(E_1) + \mathbb{P}(E_2) + \mathbb{P}(E_3) + \mathbb{P}(E_4) + \mathbb{P}(E_5) \leq \delta.$$

Appendix L. Additional Lemmas

Lemma 37 *Let S be a standardized bounded random variable, and let Z be an independent standard normal random variable. For all $t > 0$, write $X_t^S \triangleq e^{-t}S + \sqrt{1-e^{-2t}}Z$. Let S^t be another real valued bounded random variables such that (S, S^t) are an exchangeable pair. Let $\tau_t^{'S}$ and τ_t^S be the quantities defined respectively in Eq. (6) and Eq. (7) then*

$$\mathbb{E}(\tau_t^{'S} | X_t^S) = \mathbb{E}(\tau_t^S | X_t^S) = 0.$$

Proof For all measurable function h , we write $P_t(h) : x \rightarrow \mathbb{E}(h(e^{-t}x + \sqrt{1-e^{-2t}}Z))$. Let $\Phi \in \mathbb{C}^\infty(\mathbb{R})$ be a smooth function then

$$\begin{aligned} \mathbb{E}(\mathbb{E}(\tau_t^{'S} | X_t^S) \Phi(X_t^S)) &= \sum_{k=1}^{\infty} \frac{e^{-kt}}{k! \sqrt{1-e^{-2t}}^k} \mathbb{E}\left(\mathbb{E}((S^t - S)^k | S) H_k(Z) \Phi(X_t^S)\right) \\ &\stackrel{(a)}{=} \sum_{k=1}^{\infty} \frac{e^{-kt}}{k!} \mathbb{E}\left(\mathbb{E}((S^t - S)^k | S) \Phi^{(k)}(X_t^S)\right) \\ &= \sum_{k=1}^{\infty} \frac{e^{-kt}}{k!} \mathbb{E}\left((S^t - S)^k \mathbb{E}(\Phi^{(k)}(X_t^S) | S)\right) \\ &= \sum_{k=1}^{\infty} \frac{e^{-kt}}{k!} \mathbb{E}\left((S^t - S)^k (P_t \Phi)^{(k)}(S)\right) \\ &= \mathbb{E}\left((P_t \Phi)(S^t) - (P_t \Phi)(S)\right) \\ &= 0, \end{aligned}$$

where (a) is a consequence of the Stein identity that implies for any measurable function $f \in \mathbb{C}^k(\mathbb{R})$ we have $\mathbb{E}(f(Z) H_k(Z)) = \mathbb{E}(f^{(k)}(Z))$ and where the last equality is obtained by exploiting the fact that $S \stackrel{d}{=} S^t$. Similarly we can prove that $\mathbb{E}(\tau_t^S | X_t^S) = 0$. \blacksquare

Lemma 38 *Let $h_k(x) \triangleq e^{x^2/2} \frac{\partial^k}{\partial x} e^{-x^2/2}$. Shorthand $H_K \triangleq h_k(Z)$. Then the following holds for all $k, p \in \mathbb{N}$:*

$$\|H_k\|_p \leq \sqrt{k!} \sqrt{p-1}^k.$$

Lemma 39 (Marcinkiewicz-Zygmund inequality (Rio, 2009)) *Let (\tilde{X}_i) be a sequence of centered i.i.d observations. Assume that they admit a p -th moment then the following holds for all $p \geq 2$*

$$\|\frac{1}{\sqrt{n}} \sum_{i \leq n} \tilde{X}_i\|_p \leq \sqrt{p-1} \|\tilde{X}_1\|_p.$$

Lemma 40 (Rosenthal inequality with explicit constants) *Let (\tilde{X}_i) be a sequence of centered i.i.d observations. Assume that they admit a p -th moment then the following holds for all $p \geq 2$*

$$\|\frac{1}{\sqrt{n}} \sum_{i \leq n} \tilde{X}_i\|_p \leq (\frac{p}{2} + 1) n^{1/p-1/2} \|\tilde{X}_1\|_p + 2^{1/p} \sqrt{p/2 + 1} e^{\frac{1}{2} + \frac{1}{p}} \|\tilde{X}_1\|_2.$$

Proof According to [Nagaev and Pinelis \(1978, Thm. 2\)](#) we have

$$\|\frac{1}{\sqrt{n}} \sum_{i \leq n} \tilde{X}_i\|_p^p \leq \inf_{c > \frac{p}{2}} c^p n^{1-\frac{p}{2}} \|\tilde{X}_1\|_p^p + p c^{p/2} e^c B(\frac{p}{2}, c - \frac{p}{2}) \|\tilde{X}_1\|_2^p,$$

where $B(\cdot, \cdot)$ is the Beta function. Choose $c = \frac{p}{2} + 1$ then we obtain

$$\begin{aligned} \|\frac{1}{\sqrt{n}} \sum_{i \leq n} \tilde{X}_i\|_p^p &\leq (\frac{p}{2} + 1)^p n^{1-\frac{p}{2}} \|\tilde{X}_1\|_p^p + p (\frac{p}{2} + 1)^{p/2} e^{\frac{p}{2}+1} B(\frac{p}{2}, 1) \|\tilde{X}_1\|_2^p \\ &= (\frac{p}{2} + 1)^p n^{1-\frac{p}{2}} \|\tilde{X}_1\|_p^p + 2 (\frac{p}{2} + 1)^{\frac{p}{2}} e^{\frac{p}{2}+1} \|\tilde{X}_1\|_2^p. \end{aligned}$$

Therefore this directly implies that

$$\|\frac{1}{\sqrt{n}} \sum_{i \leq n} \tilde{X}_i\|_p \leq \frac{(p+2)}{2} n^{\frac{1}{p}-\frac{1}{2}} \|\tilde{X}_1\|_p + (2e)^{1/p} \sqrt{\frac{p+2}{2}} \sqrt{e} \|\tilde{X}_1\|_2.$$

■

Lemma 41 ((Romano and Wolf, 2000)) *Let (X_i) be a sequence of i.i.d random variables taking value in $[0, 1]$. Let $\alpha > 0$ be a level of confidence. Let $q_n^{\text{KS}}(\alpha)$ be the $1 - \alpha$ th quantile of a two-sided Kolmogorov-Smirnov distribution with parameter n . Let $k \in \mathbb{N}$ be an integer, then the following holds:*

$$P\left(|\hat{m}_k - m_k| \geq q_n^{\text{KS}}(\alpha)\right) \leq \alpha,$$

where $\hat{m}_k \triangleq \frac{1}{n} \sum_{i \leq n} X_i^k$.

Proof According to Proposition 3.1 of [\(Romano and Wolf, 2000\)](#) if you denote by d_K the Kolmogorov-Smirnov distance then for any two distributions on $[0, 1]$ μ, ν and any $k \in \mathbb{N}$ the following holds

$$|\mathbb{E}_{X \sim \nu}(X^k) - \mathbb{E}_{X \sim \mu}(X^k)| \leq d_K(\mu, \nu).$$

Write $KS(n)$ the Kolmogorov-Smirnov distribution with parameter n and F_{X_1} the distribution of X_1 . Now suppose that the (X_i) are continuous random variables then we know that $d_K(\frac{1}{n} \sum_{i \leq n} \delta_{\cdot=X_i}, F_{X_1}) \sim KS(n)$. Therefore this implies that

$$P\left(|\hat{m}_k - m_k| \geq q_n^{\text{KS}}(\alpha)\right) \leq P\left(d_K(\frac{1}{n} \sum_{i \leq n} \delta_{\cdot=X_i}, F_{X_1}) \geq q_n^{\text{KS}}(\alpha)\right) \leq \alpha.$$

Now if (X_i) are not continuous, let (U_i) be an iid sequence of uniform random variables in $[0, 1]$. Define:

$$X_i^\epsilon \triangleq \frac{1}{1+\epsilon}(X_i + \epsilon U_i).$$

Then the (X_i^ϵ) are continuous random variables on $[0, 1]$ we know that:

$$P\left(\left|\frac{1}{n}\sum_{i \leq n}(X_i^\epsilon)^k - \mathbb{E}((X_1^\epsilon)^k)\right| \geq q_n^{\text{KS}}(\alpha)\right) \leq \alpha.$$

As this holds for any arbitrary choice of $\epsilon > 0$ we get the desired result. ■

Appendix M. Additional Plots

The supplementary plots in this section demonstrate that the efficient EBE bounds of Fig. 1 vary only minimally as the empirical parameters \hat{m}_3 and \hat{m}_4 vary over their feasible ranges $[-R\hat{\sigma}^2, R\hat{\sigma}^2]$ and $[0, R^2\hat{\sigma}^2 - \hat{\sigma}^4]$ respectively.




Figure 2: Empirical quantile bound comparison with efficient EBE inputs $\hat{m}_3 = -R\hat{\sigma}^2$, $\hat{m}_4 = 0$.

References

Milton Abramowitz and Irene A Stegun. *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, volume 55. US Government printing office, 1964.

Theodore Wilbur Anderson. Confidence limits for the expected value of an arbitrary bounded random variable with a continuous distribution function. Technical report, STANFORD UNIV CA DEPT OF STATISTICS, 1969.

M. Artin. *Algebra*. Pearson Education, 2011. ISBN 9780132413770.




Figure 3: Empirical quantile bound comparison with efficient EBE inputs $\hat{m}_3 = -R\hat{\sigma}^2$, $\hat{m}_4 = \frac{1}{2}(R^2\hat{\sigma}^2 - \hat{\sigma}^4)$.




Figure 4: Empirical quantile bound comparison with efficient EBE inputs $\hat{m}_3 = -R\hat{\sigma}^2$, $\hat{m}_4 = R^2\hat{\sigma}^2 - \hat{\sigma}^4$.




Figure 5: Empirical quantile bound comparison with efficient EBE inputs $\hat{m}_3 = R\hat{\sigma}^2$, $\hat{m}_4 = 0$.




Figure 6: Empirical quantile bound comparison with efficient EBE inputs $\hat{m}_3 = R\hat{\sigma}^2$, $\hat{m}_4 = \frac{1}{2}(R^2\hat{\sigma}^2 - \hat{\sigma}^4)$.




Figure 7: Empirical quantile bound comparison with efficient EBE inputs $\hat{m}_3 = 0$, $\hat{m}_4 = R^2 \hat{\sigma}^2 - \hat{\sigma}^4$.




Figure 8: Empirical quantile bound comparison with efficient EBE inputs $\hat{m}_3 = R\hat{\sigma}^2$, $\hat{m}_4 = 0$.




Figure 9: Empirical quantile bound comparison with efficient EBE inputs $\hat{m}_3 = R\hat{\sigma}^2$, $\hat{m}_4 = \frac{1}{2}(R^2\hat{\sigma}^2 - \hat{\sigma}^4)$.

-
- Shiri Artstein, Keith M Ball, Franck Barthe, and Assaf Naor. On the rate of convergence in the entropic central limit theorem. *Probability theory and related fields*, 129(3):381–390, 2004.
- Jean-Yves Audibert, Rémi Munos, and Csaba Szepesvári. Exploration–exploitation tradeoff using variance estimates in multi-armed bandits. *Theoretical Computer Science*, 410(19):1876–1902, 2009.
- Andrew R Barron. Entropy and the central limit theorem. *The Annals of probability*, pages 336–342, 1986.
- Peter L Bartlett and Shahar Mendelson. Rademacher and gaussian complexities: Risk bounds and structural results. *Journal of Machine Learning Research*, 3(Nov):463–482, 2002.
- Sergei Bernstein. On a modification of chebyshev’s inequality and of the error formula of laplace. *Ann. Sci. Inst. Sav. Ukraine, Sect. Math*, 1(4):38–49, 1924.
- Chinmoy Bhattacharjee and Larry Goldstein. On strong embeddings by stein’s method. *Electronic Journal of Probability*, 21:1–30, 2016.
- A Bikyalis. Estimates of the remainder term in the central limit theorem, litovsk. *Mat. Sb*, 6: 323–346, 1966.
- Sergey G Bobkov, Gennadiy P Chistyakov, and Friedrich Götze. Rate of convergence and edgeworth-type expansion in the entropic central limit theorem. *The Annals of Probability*, pages 2479–2512, 2013.
- Thomas Bonis. Rates in the central limit theorem and diffusion approximation via stein’s method. *arXiv preprint arXiv:1506.06966*, 2015.
- Stéphane Boucheron, Gábor Lugosi, and Pascal Massart. *Concentration inequalities: A nonasymptotic theory of independence*. Oxford university press, 2013.
- Jean Bretagnolle and Pascal Massart. Hungarian constructions from the nonasymptotic viewpoint. *The Annals of Probability*, pages 239–256, 1989.
- T Cacoullos and V Papathanasiou. Lower variance bounds and a new proof of the central limit theorem. *Journal of multivariate analysis*, 43(2):173–184, 1992.
- Theophilos Cacoullos, Vasilis Papathanasiou, and Sergey A Utev. Variational inequalities with examples and an application to the central limit theorem. *The Annals of Probability*, pages 1607–1618, 1994.
- Sourav Chatterjee. A new approach to strong embeddings. *Probability Theory and Related Fields*, 152(1):231–264, 2012.
- Louis HY Chen, Larry Goldstein, and Qi-Man Shao. *Normal approximation by Stein’s method*, volume 2. Springer, 2011.
- T Courtade, M Fathi, and A Pananjady. Existence of stein kernels via spectral gap, and discrepancy bounds. to appear in *ann. ihp: Probab. Stat*, 2017.

-
- Rick Durrett. *Probability: theory and examples*, volume 49. Cambridge university press, 2019.
- A. Dvoretzky, J. Kiefer, and J. Wolfowitz. Asymptotic Minimax Character of the Sample Distribution Function and of the Classical Multinomial Estimator. *The Annals of Mathematical Statistics*, 27(3):642 – 669, 1956. doi: 10.1214/aoms/1177728174. URL <https://doi.org/10.1214/aoms/1177728174>.
- Bradley Efron. Bootstrap methods: another look at the jackknife. In *Breakthroughs in statistics*, pages 569–593. Springer, 1992.
- Marie Ernst, Gesine Reinert, and Yvik Swan. First-order covariance inequalities via stein’s method. *Bernoulli*, 26(3):2051–2081, 2020.
- Carl-Gustav Esseen. On the liapunov limit error in the theory of probability. *Ark. Mat. Astr. Fys.*, 28:1–19, 1942.
- Max Fathi. Stein kernels and moment maps. *The Annals of Probability*, 47(4):2172–2185, 2019.
- M Gasko. A new distribution-free lower confidence bound for the mean of a positive random variable. Technical report, Working Paper QS 92-006, San Jose State Univ. College of Business, 1991.
- Larry Goldstein and Gesine Reinert. Stein’s method and the zero bias transformation with application to simple random sampling. *The Annals of Applied Probability*, 7(4):935–952, 1997.
- Peter Hall. *The bootstrap and Edgeworth expansion*. Springer Science & Business Media, 2013.
- Wassily Hoeffding. Probability inequalities for sums of bounded random variables. *Journal of the American Statistical Association*, 58(301):13–30, 1963.
- Kwang-Sung Jun and Francesco Orabona. Parameter-free online convex optimization with sub-exponential noise. In *Conference on Learning Theory*, pages 1802–1823. PMLR, 2019.
- János Komlós, Péter Major, and Gábor Tusnády. An approximation of partial sums of independent rv’-s, and the sample df. i. *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete*, 32(1):111–131, 1975.
- János Komlós, Péter Major, and Gábor Tusnády. An approximation of partial sums of independent rv’s, and the sample df. ii. *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete*, 34(1):33–58, 1976.
- Michel Ledoux, Ivan Nourdin, and Giovanni Peccati. Stein’s method, logarithmic sobolev and transport inequalities. *Geometric and Functional Analysis*, 25(1):256–306, 2015.
- Christophe Ley, Gesine Reinert, and Yvik Swan. Stein’s method for comparison of univariate distributions. *Probability Surveys*, 14:1–52, 2017.
- David M Mason and Harrison H Zhou. Quantile coupling inequalities and their applications. *Probability Surveys*, 9:439–479, 2012.

-
- Andreas Maurer and Massimiliano Pontil. Empirical bernstein bounds and sample-variance penalization. In *COLT*, 2009.
- Volodymyr Mnih, Csaba Szepesvári, and Jean-Yves Audibert. Empirical bernstein stopping. In *Proceedings of the 25th international conference on Machine learning*, pages 672–679, 2008.
- Sergey Victorovich Nagaev. Some limit theorems for large deviations. *Theory of Probability & Its Applications*, 10(2):214–235, 1965.
- SV Nagaev and IF Pinelis. Some inequalities for the distribution of sums of independent random variables. *Theory of Probability & Its Applications*, 22(2):248–256, 1978.
- Ivan Nourdin and Frederi Viens. Density formula and concentration inequalities with malliavin calculus. *Electronic Journal of Probability*, 14:2287–2309, 2009.
- Ivan Nourdin, Giovanni Peccati, and Anthony Réveillac. Multivariate normal approximation using stein’s method and malliavin calculus. In *Annales de l’IHP Probabilités et statistiques*, volume 46, pages 45–58, 2010.
- Ivan Nourdin, Giovanni Peccati, and Yvik Swan. Entropy and the fourth moment phenomenon. *Journal of Functional Analysis*, 266(5):3170–3207, 2014.
- Francesco Orabona and Kwang-Sung Jun. Tight concentrations and confidence sequences from the regret of universal portfolio. *arXiv preprint arXiv:2110.14099*, 2021.
- Felix Otto and Cédric Villani. Generalization of an inequality by talagrand and links with the logarithmic sobolev inequality. *Journal of Functional Analysis*, 173(2):361–400, 2000.
- Emmanuel Rio. Moment inequalities for sums of dependent random variables under projective conditions. *Journal of Theoretical Probability*, 22(1):146–163, 2009.
- Joseph P Romano and Michael Wolf. Finite sample nonparametric inference and large sample efficiency. *Annals of Statistics*, pages 756–778, 2000.
- Nathan Ross. Fundamentals of stein’s method. *Probability Surveys*, 8:210–293, 2011.
- Adrien Saumard. Weighted poincaré inequalities, concentration inequalities and tail bounds related to stein kernels in dimension one. *Bernoulli*, 25(4B):3978–4006, 2019.
- I. G. Shevtsova. An improvement of convergence rate estimates in the lyapunov theorem. *Doklady Mathematics*, 82(3):862–864, 2010. doi: 10.1134/S1064562410060062. URL <https://doi.org/10.1134/S1064562410060062>.
- Irina Shevtsova. On the absolute constants in the berry-esseen type inequalities for identically distributed summands. *arXiv preprint arXiv:1111.6554*, 2011.
- Charles Stein. Approximate computation of expectations. IMS, 1986.
- Martin J Wainwright. *High-dimensional statistics: A non-asymptotic viewpoint*, volume 48. Cambridge University Press, 2019.

Ian Waudby-Smith and Aaditya Ramdas. Estimating means of bounded random variables by betting. *arXiv preprint arXiv:2010.09686*, 2020.

Wenda Zhou, Victor Veitch, Morgane Austern, Ryan P Adams, and Peter Orbanz. Non-vacuous generalization bounds at the imagenet scale: a pac-bayesian compression approach. *arXiv preprint arXiv:1804.05862*, 2018.