

VARIATIONS OF CENTRAL LIMIT THEOREMS AND STIRLING NUMBERS OF THE FIRST KIND

BERNHARD HEIM AND MARKUS NEUHAUSER

ABSTRACT. We construct a new parametrization of double sequences $\{A_{n,k}(s)\}_{n,k}$ between $A_{n,k}(0) = \binom{n-1}{k-1}$ and $A_{n,k}(1) = \frac{1}{n!} [n]_k$, where $[n]_k$ are the unsigned Stirling numbers of the first kind. For each s we prove a central limit theorem and a local limit theorem. This extends the de Moivre–Laplace central limit theorem and Goncharov’s result, that unsigned Stirling numbers of the first kind are asymptotically normal. Here-with, we provide several applications.

1. INTRODUCTION

The central limit theorem is one of the most remarkable theorems in science [Fe45, Fe71, Fi11]. From the de Moivre–Laplace theorem in combinatorics and probability, involving binomial distributions, the central limit theorem culminated in a universal law of nature ([KM20], section 3). In this paper, we construct a parametrization of double sequences $\{A_{n,k}(s)\}_{n,k}$ between $A_{n,k}(0) = \binom{n-1}{k-1}$ and $A_{n,k}(1) = \frac{1}{n!} [n]_k$, where $[n]_k$ are the unsigned Stirling numbers of the first kind. Finally, we apply Harper’s method [Ha67] and prove a central limit theorem and a local limit theorem for each s .

Let $Z_n \in \{0, 1, \dots, n\}$ denote a random variable with binomial distribution

$$\mathbb{P}(Z_n = k) = \binom{n}{k} p^k q^{n-k},$$

where $0 < p < 1$ and $q = 1 - p$. The central limit theorem by de Moivre–Laplace states, that the normalized random variables Z_n , converge in distribution against the standard normal distribution $N(0, 1)$:

$$\frac{Z_n - np}{\sqrt{npq}} \xrightarrow{D} N(0, 1).$$

In asymptotic analysis [Be73, Ca15], one is interested in the asymptotic normality of sequences. Goncharov [Go44, Go62] proved in 1944, that the unsigned Stirling numbers of the first kind $[n]_k$ have this property. More than 20 years later, Harper [Ha67] discovered

2020 *Mathematics Subject Classification.* Primary 05A16, 60F05; Secondary 05A16, 11B39.

Key words and phrases. Central Limit Theorem, Local Limit Theorem, Probabilistic Number Theory, Singularity Analysis.

a more conceptional method, proving also that the Stirling numbers of the second kind are asymptotically normal. When describing Goncharov's proof, Harper writes "Goncharov . . . by brute force torturously manipulates the characteristic functions of the distributions until they approach $\exp(-x^2/c)$, c a positive constant."

Recently, Harper's method has been applied by Gawronski and Neuschel [GN13], investing in Euler–Frobenius numbers and by Kahle and Stump [KS19]. Note, that in several cases, also other non-gaussian distributions need to be considered (e. g. limiting Betti distributions of Hilbert schemes of n points, where the Gumbel distribution was the right limit distribution, Griffin et al. [GORT22]).

Moreover, related to the topic, since we deal with unimodal sequences, we suggest the analysis of properties of the modes. We utilize a result of Darroch [Da64, Be96] and study the modes of $\{A_{n,k}(s)\}$. This is connected to Erdős' proof [Er53] of the Hammersley conjecture [Ha51], related to the peaks of $\{\left[n\right]_k\}$. For other sequences, we refer to Bringmann et al. [BJSMR19].

2. MAIN RESULTS

Let $h_s(n) := n^s$ for $s \in \mathbb{R}$. Let $P_n^{h_s}(x) := \frac{x}{h_s(n)} \sum_{k=0}^{n-1} P_k^{h_s}(x)$ with initial value $P_0^{h_s}(x) := 1$. We are interested in the coefficients of these polynomials:

$$P_n^{h_s}(x) = \sum_{k=0}^n A_{n,k}(s) x^k.$$

Let $n \geq 1$. Then $A_{n,n}(s) = (n!)^{-s}$ and $A_{n,0}(s) = 0$. We deduce from [HN21], Example 1 and Example 3 for the values $s = 0$ and $s = 1$:

$$(2.1) \quad A_{n,k}(0) = \binom{n-1}{k-1} \text{ and } A_{n,k}(1) = \frac{1}{n!} \left[n\right]_k.$$

The unsigned Stirling numbers of the first kind $\left[n\right]_k$, denote the number of all permutations of a set of n elements with exactly k distinct cycles. We refer to Bóna [Bo12].

We mainly focus on $s \in [0, 1]$ due to (2.1). Then, we obtain central and local limit theorems for the double sequence $\{A_{n,k}(s)\}_{n,k}$. The case $s \in [-1, 0]$ can be reduced to the case $[0, 1]$, since

$$P_n^{h_s}(x) = h_s(n) \prod_{k=1}^n h_s(k) x^{n+1} P_n^{h_{-s}}(x^{-1}).$$

2.1. Central limit theorem. The classical central limit theorem of de Moivre (1738) and Laplace (1812), was developed from the results in probability theory [Fi11, KM20] to a general theorem, without direct reference to concepts as random variable, expected value, and variance. We refer to Feller [Fe45] and Canfield [Ca15] for excellent surveys.

The modern version of the central limit theorem can also be considered as a theorem on the asymptotic normality of a sequence of non-negative numbers in singularity analysis [Be73]. In this spirit, we state our first result in the most general form.

Theorem 2.1. *Suppose $s \in [0, 1]$. Then there exist real sequences $\{a_n(s)\}_n$ and $\{b_n(s)\}_n$ with $b_n(s)$ positive for almost all n , so that*

$$(2.2) \quad \lim_{n \rightarrow \infty} \sup_{x \in \mathbb{R}} \left| \frac{1}{P_n^{h_s}(1)} \sum_{k \leq a_n(s) + x b_n(s)} A_{n,k}(s) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-\frac{t^2}{2}} dt \right| = 0.$$

Theorem 2.1 is proven with Harper's method [Ha67]. The sequences $\{a_n(s)\}_n$ and $\{b_n(s)\}_n$ are provided by the expected values and variances of a suitable sequence of random variables. Let $h_s(0) := 0$. It is essential, that

$$(2.3) \quad P_n^{h_s}(x) = (n!)^{-s} \prod_{k=0}^{n-1} (x + h_s(k))$$

has real roots $-h_s(k) \leq 0$.

Further, we utilize the Berry–Esseen theorem [Ca15] to control the convergence rate. The corresponding expected values and variances are

$$(2.4) \quad \mu_n(s) := 1 + \sum_{k=1}^{n-1} \frac{1}{1 + k^s},$$

$$(2.5) \quad \sigma_n^2(s) := \sum_{k=1}^{n-1} \frac{k^s}{(1 + k^s)^2}.$$

Then $\mu_n(0) = \frac{n+1}{2}$ and $\mu_n(1) = H_n$, where H_n is the n th harmonic number. Moreover, let generally $H_n^{(r)} := \sum_{k=1}^n k^{-r}$. We have $\sigma_n^2(0) = \frac{n-1}{4}$ and $\sigma_n^2(1) = H_n - H_n^{(2)}$. Most importantly, let $s \in [0, 1]$. Then

$$\lim_{n \rightarrow \infty} \sigma_n(s) \longrightarrow \infty.$$

Therefore, we obtain:

Theorem 2.2. *Suppose $s \in [0, 1]$. There exists a positive constant C so that*

$$\lim_{n \rightarrow \infty} \sup_{x \in \mathbb{R}} \left| \frac{1}{P_n^{h_s}(1)} \sum_{k \leq \mu_n(s) + x \sigma_n(s)} A_{n,k}(s) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-\frac{t^2}{2}} dt \right| \leq C \frac{1}{\sigma_n(s)}.$$

The standard deviation $\sigma_n(s)$ approaches infinity.

Remarks.

a) Theorem 2.2 implies Theorem 2.1. In singularity analysis [Be73, Ca15], one is most

interested in the asymptotic behaviors of $\{a_n\}_n$ and $\{b_n\}_n$, rather than in the concrete realization, as given in Theorem 2.2.

b) The constant be chosen as $C = 0.7975$ (we refer to [vB72], and the survey article [Pi97]).

c) Let $s > 1$. Then $\lim_{n \rightarrow \infty} \sigma_n(s) \leq \lim_{n \rightarrow \infty} \sum_{k=1}^{n-1} k^{-s} = \zeta(s) < \infty$. Here we denote by $\zeta(s)$ the Riemann zeta function.

2.2. Local limit theorem. We refer to Section 4 for an introduction. We prove:

Theorem 2.3. *Let $s \in \mathbb{R}$. Then there exists a universal constant $K > 0$, so that*

$$\max_k \left| \frac{\sigma_n(s)}{P_n^{hs}(1)} A_{n,k}(s) - \frac{e^{-\frac{(x_n(s))^2}{2}}}{\sqrt{2\pi}} \right| < \frac{K}{\sigma_n(s)},$$

for $x_n(s) = (k - \mu_n(s))/\sigma_n(s)$. Uniformly for $k - \mu_n(s) = O(\sigma_n(s))$ we have

$$\frac{A_{n,k}(s)}{P_n^{hs}(1)} \sim \frac{e^{-(x_n(s))^2/2}}{\sigma_n(s) \sqrt{2\pi}}.$$

2.3. Peaks and plateaux. The polynomials $P_n^{hs}(x)$ are real-rooted. Therefore, a theorem by Newton implies, that the sequence $\{A_{n,k}(s)\}_k$ is unimodal and has two modes at most. Either we have one peak, or a plateau.

In the case $s = 0$, we have a peak for n odd at $k = \frac{n+1}{2}$ and a plateau for n even at $k = \frac{n}{2}$ and $\frac{n+2}{2}$. This is obvious, since the $A_{n,k}(0)$ are binomial coefficients. The case $s = 1$ is more delicate. Let $n \geq 3$. Hammersley [Ha51] conjectured in the context of Stirling numbers of the first kind that there is always a peak. This was proved by Erdős [Er53]. The proof depends on the fact, that $\left\{ \begin{bmatrix} n \\ k \end{bmatrix} \right\}_k$ are natural numbers. This allows Erdős to apply special results related to the prime number theorem and certain divisibility properties of the Stirling numbers of the first kind. Our goal is to contribute to the case $s \in (0, 1)$ and obtain information for $s = 1$. But this seems to be very difficult, since in general, the numbers $A_{n,k}(s)$ are not integers. Nevertheless, by utilizing a theorem by Darroch [Da64] we obtain:

Theorem 2.4. *Let $n \geq 6$. Let $k_0 \in \mathbb{N}$ be any integer associated with $s \in (0, 1)$ so that*

$$k_0 = 1 + \sum_{k=1}^{n-1} \frac{1}{1 + k^s}.$$

Then the sequence $\{A_{n,k}(s)\}$ has a peak at k_0 . The number of possible k_0 is given by the number of integers between H_n and $\frac{n+1}{2}$.

3. THE PROBABILISTIC VIEWPOINT ON ASYMPTOTIC NORMALITY

We begin with a useful tool from probability.

3.1. The Berry–Esseen theorem [Ca15], **theorem 3.2.4.** Let X be a random variable. We denote by $\mathbb{E}(X)$ and $\mathbb{V}(X)$ the expected value and variance of X .

Theorem 3.1. Let $X_{n,k}$ for $1 \leq k \leq n$ be independent random variables with values in $\{0, 1, \dots, n\}$. Let $\mu_{n,k}$ be the expected values, $\sigma_{n,k}^2$ the variances, and

$$\rho_{n,k} = \mathbb{E}(|X_{n,k} - \mu_{n,k}|^3) < \infty$$

the absolute third central moment. Let $\mu_n := \sum_{k=1}^n \mu_{n,k}$, $\sigma_n^2 := \sum_{k=1}^n \sigma_{n,k}^2$, and $Z_n := \sum_{k=1}^n X_{n,k}$. Let $Z_n^* = (Z_n - \mu_n)/\sigma_n$. Then

$$\left\| \mathbb{P}(Z_n^* < x) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-\frac{t^2}{2}} dt \right\|_{\mathbb{R}} \leq C \frac{\sum_{k=1}^n \rho_{n,k}}{\sigma_n^3},$$

where $\|f(x)\|_{\mathbb{R}}$ denotes the supremum norm of f on \mathbb{R} and $C > 0$ is a universal constant. This constant can be chosen as $C = 0.7975$ [vB72].

Let $P_n(x) = \sum_{k=0}^n a_{n,k} x^k$ be a monic polynomial of degree n with $a_{n,k} \geq 0$. Suppose, the roots of $P_n(x)$ are real and $P_n(x) = \prod_{k=1}^n (x + r_k)$. Harper [Ha67] introduced a triangular array of Bernoulli random variables $X_{n,j}$ with distribution

$$\mathbb{P}(X_{n,j} = 0) := \frac{r_j}{1 + r_j} \text{ and } \mathbb{P}(X_{n,j} = 1) := \frac{1}{1 + r_j}.$$

Let $Z_n := \sum_{j=1}^n X_{n,j}$. Then $\mathbb{P}(Z_n = k) = \frac{a_{n,k}}{P_n(1)}$. Let $X_{n,j}$ be given. Then

$$\mathbb{E}(X_{n,j}) = \frac{1}{1 + r_j}, \quad \mathbb{V}(X_{n,j}) = \frac{r_j}{(1 + r_j)^2}, \quad \mathbb{E}(|X_{n,j} - \mathbb{E}(X_{n,j})|^3) = \frac{r_j(1 + r_j^2)}{(1 + r_j)^4}.$$

This implies that

$$(3.1) \quad \mathbb{E}(|X_{n,j} - \mathbb{E}(X_{n,j})|^3) < \mathbb{V}(X_{n,j}).$$

We apply Harper's method setting $P_n(x) = (n!)^s P_n^{hs}(x)$. The expected value of $Z_n(s)$ is given by $\mu_n(s)$ and the variance by $\sigma_n(s)^2$, as recorded in (2.4) and (2.5).

Lemma 3.2. a) Let $s \in [0, 1)$. Then

$$\mu_n(s) \sim \frac{n^{1-s}}{1-s} \text{ and } \sigma_n(s) \asymp n^{1-s}.$$

b) We obtain

$$\mu_n(1) \sim \ln n \text{ and } \sigma_n^2(1) \asymp \ln n.$$

Recall that $f \asymp g$ is an abbreviation for $f = O(g)$ and $g = O(f)$.

Proof. a) Let $n \geq 2$. We have $\frac{1}{1+k^s} \leq \frac{1}{k^s}$. For $k \leq t \leq k+1$ we have $(k+1)^{-s} \leq t^{-s} \leq k^{-s}$. Therefore, $1 + \sum_{k=1}^{n-1} \frac{1}{1+k^s} \leq 2 + \int_1^{n-1} t^{-s} dt$. Now, for $0 \leq s < 1$, we obtain $\sum_{k=1}^{n-1} \frac{1}{1+k^s} \leq 2 + \frac{(n-1)^{1-s}-1}{1-s}$.

For $k \geq K$, we have $\frac{1}{1+k^s} \geq \frac{1}{K^{-s+1}} k^{-s}$. Therefore, $1 + \sum_{k=1}^{n-1} \frac{1}{1+k^s} \geq \frac{1}{K^{-s+1}} \int_K^n t^{-s} dt$. For $0 \leq s < 1$, this shows $1 + \sum_{k=1}^{n-1} \frac{1}{1+k^s} \geq \frac{1}{K^{-s+1}} \frac{n^{1-s} - K^{1-s}}{1-s}$.

This indicates that for $0 \leq s < 1$ and arbitrarily small $\varepsilon > 0$, there is an $N \in \mathbb{N}$ so that $(1 - \varepsilon) \frac{n^{1-s}}{1-s} \leq \mu_n(s) \leq (1 + \varepsilon) \frac{n^{1-s}}{1-s}$ for all $n \geq N$. Therefore, for $0 \leq s < 1$ and all $\varepsilon > 0$, there is an $N \in \mathbb{N}$, such that for all $n \geq N$ holds $1 - \varepsilon \leq \frac{\mu_n(s)}{\frac{n^{1-s}}{1-s}} \leq 1 + \varepsilon$.

We have $\sigma_n^2(s) \geq \sum_{k=1}^{n-1} \frac{1}{4k^s} \geq \frac{1}{4} \int_1^n t^{-s} dt$. So, for $0 \leq s < 1$, we obtain $\sigma_n^2(s) \geq \frac{n^{1-s} - 1}{4 - 4s}$. Additionally, we have $\sigma_n^2(s) \leq \sum_{k=1}^{n-1} k^{-s} = \mu_n(s)$, meaning that the same upper bounds also apply here.

b) Similarly, for $s = 1$, we obtain $\frac{\ln(n) - \ln(K)}{K^{-s+1}} \leq 1 + \sum_{k=1}^{n-1} \frac{1}{1+k^s} \leq 2 + \ln(n-1)$, $1 - \varepsilon \leq \frac{\mu_n(1)}{\ln(n)} \leq 1 + \varepsilon$, and $\sigma_n^2(1) \geq \frac{1}{4} \ln(n)$. \square

Corollary 3.3. *Let $s \in [0, 1]$. Then $\lim_{n \rightarrow \infty} \sigma_n(s) = \infty$.*

Remark. Let $0 \leq s_1 \leq s_2$. Then $\sigma_n(s_1) \geq \sigma_n(s_2)$, since $\frac{d}{ds} \sigma_n(s) < 0$.

3.2. Proof of Theorem 2.2. Let $s \in [0, 1]$. Let $X_{n,k}(s) \in \{0, 1\}$ for $1 \leq k \leq n$ be a random variable with values 0 and 1. Here

$$\mathbb{P}(X_{n,k}(s) = 1) = \frac{1}{1 + r_{n,k}(s)},$$

where $r_{n,k}(s) = h_s(k-1)$. We put $Z_n(s) := \sum_{k=1}^n X_{n,k}(s)$. Then

$$\mathbb{P}(Z_n(s) = k) = \frac{A_{n,k}(s)}{P_n^{h_s}(1)}.$$

Thus, we obtain

$$\mathbb{E}(Z_n(s)) = \mu_n(s) = 1 + \sum_{k=1}^{n-1} \frac{1}{1+k^s} \text{ and } \mathbb{V}(Z_n(s)) = \sigma_n(s)^2 = \sum_{k=1}^{n-1} \frac{k^s}{(1+k^s)^2}.$$

Corollary 3.3 states, that the variance approaches infinity. This proves Theorem 2.2.

3.3. Proof of Theorem 2.1. This follows from Theorem 2.2. The crucial part of successfully applying Harper's method and the Berry–Esseen theorem is that the variance approaches infinity.

4. LOCAL LIMIT THEOREM

A double indexed sequence $\{a_{n,k}\}_{n,k}$ satisfies a local limit theorem on a set S of real numbers provided

$$\sup_{x \in S} \left| \frac{\sigma_n a(n, \lfloor \mu_n + x \sigma_n \rfloor)}{\sum_k a(n, k)} - \frac{e^{-x^2/2}}{\sqrt{2\pi}} \right| \rightarrow 0$$

(cf. Canfield [Ca15], section 3.7). We recall the following result due to Bender [Be73].

Theorem 4.1 (Bender). *Suppose, that the $\{a(n, k)\}_k$ for $n \in \mathbb{N}$ are asymptotically normal, and $\sigma_n^2 \rightarrow \infty$. If for each n the sequence $\{a(n, k)\}_k$ is log-concave in k , then $\{a(n, k)\}_k$ satisfies a local limit theorem on $S = \mathbb{R}$.*

4.1. Pólya frequency sequences and limit theorems. We follow the excellent survey by Pitman [Pi97] and apply several results to the sequences $\{A_{n,k}(s)\}_k$.

Let (a_0, a_1, \dots, a_n) be a sequence of non-negative real numbers. Let $P_n(x) := \sum_{k=0}^n a_k x^k$ be real rooted and $P_n(1) > 0$. Then the sequence is called a (finite) Pólya frequency sequence. Let $Z_n \in \{0, 1, \dots, n\}$ be a random variable with $\mathbb{P}(Z_n = k) := \frac{a_k}{P_n(1)}$, mean μ , and variance σ^2 . Then

$$\max_k \left| \mathbb{P}(0 \leq Z_n \leq k) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\frac{k-\mu}{\sigma}} e^{-\frac{t^2}{2}} dt \right| < \frac{0.7975}{\sigma}$$

(see [Pi97], formula (24) or [vB72] for a reference).

Further, there exists a universal constant K so that

$$\max_k \left| \sigma \mathbb{P}(Z_n = k) - \frac{1}{\sqrt{2\pi}} e^{-(\frac{k-\mu}{\sigma})^2/2} \right| < \frac{K}{\sigma}.$$

The bound is due to Platanov [Pl80] (see also [Pi97], formula (25)).

Proof of Theorem 2.3. It follows from our previous considerations, that $\{A_{n,k}(s)\}_k$ is a Pólya frequency sequence for all $s \in \mathbb{R}$. This implies the theorem. \square

5. PEAKS OF $\{A_{n,k}(s)\}_k$

We recall Darroch's theorem [Da64]. Let (a_0, a_1, \dots, a_n) be a Pólya frequency sequence. Let $p(x) := \sum_{k=0}^n a_k x^k$ with $p(1) > 0$. Let $\mu_n := \frac{p'(1)}{p(1)}$. Newton's theorem implies, that the sequence $\{a_k\}_k$ is unimodal and has two modes at most. Darroch proved, that the modes have distance less than 1 from μ_n . Armed with the results of the previous sections we have:

Proof of Theorem 2.4.

We consider the polynomials $P_n^{h_s}(x)$. Then $\mu_n(s) = 1 + \sum_{k=1}^{n-1} \frac{1}{1+k^s}$. Suppose, that $\mu_n(s)$ is an integer. Then $\{A_{n,k}(s)\}_k$ has a peak. Let us restrict μ_n to $[0, 1]$. In this case

$$\mu_n : [0, 1] \longrightarrow [\mu_n(1), \mu_n(0)], \quad s \mapsto \mu_n(s).$$

It construes that μ_n is differentiable and strictly monotone for $n \geq 3$. We have

$$\frac{d}{ds} \mu_n(s) = - \sum_{k=2}^{n-1} \frac{k^s \ln k}{(1+k^s)^2} < 0.$$

Therefore, μ_n is bijective. Let $n \geq 6$. Then $\mu_n(0) - \mu_n(1) > 1$. This implies that integers $k_0 \in (\mu_n(1), \mu_n(0))$ exist and are realizable by suitable $s \in (0, 1)$:

$$k_0 = 1 + \sum_{k=1}^{n-1} \frac{1}{1 + k^s}.$$

Let such an s be given. Then $\{A_{n,k}(s)\}_k$ has a peak at k_0 . \square

We add the following approximation of $\mu_n(s)$.

Remark. Let $s \in [0, 1]$ and $n \geq 3$. There exists a $\xi \in [0, 1]$ so that

$$\mu_n(s) = \frac{n+1}{2} - \left(\frac{1}{4} \sum_{k=2}^{n-1} \ln k \right) s + \frac{\mu_n''(\xi)}{2} s^2.$$

Finally, we provide an illustration of Theorem 2.4 by Table 1. Let $m_n(1)$ be the unique mode of $\{A_{n,k}(1)\}$, as proven by Erdős.

n	1	2	3	4	5	6	7	8	9	10	100	1000
H_n	1	1.5	1.83	2.08	2.28	2.45	2.59	2.72	2.83	2.93	5.19	7.49
$m_n(1)$	—	—	2	2	2	2	2	3	3	3	5	7
$\frac{n+1}{2}$	1	1.5	2	2.5	3	3.5	4	4.5	5	5.5	50.5	500.5

TABLE 1. Modes for $s = 1$ and related values.

6. APPLICATIONS

6.1. Approximation of Stirling numbers of the first kind. Wilf [Wi93] contributed to the asymptotic behavior of Stirling numbers of the first kind. Several asymptotic formulas are provided. Wilf also compares his results with Jordan's formula in the case $\left[\begin{smallmatrix} 100 \\ 5 \end{smallmatrix} \right]$ and presents numerical data. Although, the focus of this paper is not to obtain best approximations, the numerical value we obtain is already solid. We refer to Theorem 2.3 and the approximation

$$(6.1) \quad \frac{1}{100!} \left[\begin{smallmatrix} 100 \\ 5 \end{smallmatrix} \right] \approx \frac{e^{-\left(\frac{5 - \mu_{100}(1)}{\sigma_{100}(1)} \right)^2 / 2}}{\sigma_{100}(1) \sqrt{2\pi}}.$$

We have $\mu_{100}(1) \approx 5.19$ and $\sigma_{100}(1) \approx 1.88477$. Wilf considers the value of $\left[\begin{smallmatrix} 100 \\ 5 \end{smallmatrix} \right] / 99!$ (Table 2).

	$\frac{[100]_5}{99!}$	Error
Exact value	21.1204415...	-
Jordan formula	18.740...	$\approx 11\%$
3 term of Eq. (9) (Wilf)	21.24986...	$\approx 0.613\%$
Theorem 1 above (Wilf)	20.960...	$\approx 0.76\%$
Approximation (6.1) (this paper)	21.062180...	$\approx 0.28\%$
Eq. (7) to order $1/n$ (Wilf)	21.12070...	$\approx 0.0012\%$
Eq. (7) to order $1/n^2$ (Wilf)	21.1204409...	$\approx 0.000003\%$

TABLE 2. Several approximations of the maximal value for $n = 100$ (see Wilf [Wi93], page 349 for details).

6.2. One mode property of the Stirling numbers of the first kind. Let $n \geq 3$. Erdős [Er53] proved that $\left\{ \begin{bmatrix} n \\ k \end{bmatrix} \right\}_k$ has one mode. We give a new proof for infinitely many n . A variant of Darroch's theorem [Da64, Pi97] states: Let $\{a_k\}_k$ be a Pólya frequency sequence. Let $\mu_n = \frac{p'_n(1)}{p_n(1)}$, then the sequence has exactly one mode k_0 , if

$$k_0 \leq \mu_n < k_0 + \frac{1}{k_0 + 2} \quad \text{or} \quad k_0 - \frac{1}{n - k_0 + 2} < \mu_n \leq k_0.$$

Let the sequence $A_{n,k}(1)$ be given. Then $\mu_n(1) = H_n$. Since H_n is unbounded and $H_{n+1} = H_n + \frac{1}{n+1}$, we can find infinitely many pairs (n, k_0) , such that $H_n \in [k_0, k_0 + \frac{1}{k_0+2}]$. This implies, that $\{A_{n,k}(1)\}_k$ has one mode and thus, $\left\{ \begin{bmatrix} n \\ k \end{bmatrix} \right\}_k$ has one mode.

Example. From Table 1 we see that for $n = 10$, we have $k_0 = 3$ and $3 - \frac{1}{9} < 2.9 < H_{10} = \mu_{10}(1) \leq k_0$. For $n = 30$, we have $\mu_{30} = H_{30} \approx 3.995$ and therefore, $k_0 = 4$ is the unique mode, as $4 - \frac{1}{28} < 3.97 < \mu_{30} \leq 4$. For $83 \leq n \leq 95$, we have $5.002 \approx \mu_{83} = H_{83} \leq \mu_n \leq \mu_{95} \approx 5.136$ and therefore, $k_0 = 5$ is the unique mode as $5 \leq \mu_n < 5.14 < 5 + \frac{1}{7}$.

One can vary the argument and show:

Theorem 6.1. *Let $n \geq 6$. We consider the sequence $\{A_{n,k}(s)\}_k$ for $s \in (0, 1)$. Let $k_0 = \mu_n(s_0) \in \mathbb{N}$ with $s_0 \in (0, 1)$. Then there exists an $\varepsilon > 0$, so that for all $s \in (s_0 - \varepsilon, s_0 + \varepsilon)$, the sequences $\{A_{n,k}(s)\}_k$ have exactly one mode.*

REFERENCES

- [vB72] P. van Beek: *An application of Fourier methods to the problem of sharpening the Berry–Esseen inequality*. Z. Wahrsch. Verw. Gebiete **23**, (1972), 187–196.
- [Be73] E. Bender: *Central and local limit theorems applied to asymptotic enumeration*. J. Comb. Theory (A) **15**, (1973), 91–111.
- [Be96] M. Benoumhani: *Sur une propriété des polynômes à racines réelles négatives*. J. Math. Pures Appl., (1996), 85–110.
- [Bo12] M. Bóna: *Combinatorics of Permutations*. Chapman and Hall/CRC (2012).

- [BJSMR19] K. Bringmann, C. Jennings-Shaffer, K. Mahlburg, R. Rhoades: *Peak positions of strongly unimodal sequences*. Trans. American Math. Society **372** (10), (2019), 7087–7109.
- [Ca15] E. R. Canfield: *Asymptotic normality in enumeration*. In: Miklós Bóna (ed.) Handbook of Enumeration, CRC Press. Discrete Mathematics and its Applications (2015), 255–280.
- [Da64] J. N. Darroch: *On the distribution of the number of successes in independent trials*. Ann. Math. Statist. **35** (3), (1964), 1317–1321.
- [Er53] P. Erdős: *On a conjecture of Hammersley*. Journal London Math. Soc. **28**, (1953), 232–236.
- [Fe45] W. Feller: *The fundamental limit theorems in probability*. Bull. AMS **51**, (1945), 800–832.
- [Fe71] W. Feller: *An Introduction to Probability Theory and Applications*. **II**. Wiley (1971).
- [Fi11] H. Fischer: *A History of the Central Limit Theorem*. From Classical to Modern Probability Theory. Springer, New York, NY (2011).
- [GN13] W. Gawronski, T. Neuschel: *Euler–Frobenius numbers*. Integral transforms and special functions **24** (10), (2013), 817–830.
- [Go44] V. Goncharov: *Du domaine de l’analyse combinatoire*. Bull. Acad. Sci. URSS Sér. Math. **8**, (1944), 3–48.
- [Go62] V. Goncharov: *On the field of combinatorial analysis*. Am. Math. Soc., Transl., II. Ser. **19**, (1962), 1–46.
- [GORT22] M. Griffin, K. Ono, L. Rolen, W.-L. Tsai: *Limiting Betti distribution of Hilbert schemes on n points*. Canadian Math. Bulletin, (2022), 1–16.
- [Ha51] J. M. Hammersley: *The sum of products of the natural numbers*. Proc. London Math. Soc. **1** (3), (1951), 435–452.
- [Ha67] L. Harper: *Stirling behaviour is asymptotically normal*. Ann. Math. Stat. **38**, (1967), 410–414.
- [HN21] B. Heim, M. Neuhauser: *Horizontal and vertical log-concavity*. Research in Number Theory **7** (18), (2021), 12pp.
- [KS19] T. Kahle, C. Stump: *Counting inversions and descents of random elements in finite Coxeter groups*. Mathematics of Computation **89** (321), (2020), 437–464.
- [KM20] A. Kar, M. Murty: *The central limit theorem in algebra and number theory*. In: B. Ramakrishnan, Bernhard Heim, Brundaban Sahu (eds.) Modular Forms and Related Topics in Number Theory. Kozhikode, India, December 10–14, 2018. Springer Proceed. Math. & Statistics **340**, (2020), 101–124.
- [Pi97] J. Pitman: *Probabilistic bounds on the coefficients of polynomials with only real zeros*. J. of Combinat. Theory, Series A **77**, (1997), 279–303.
- [Pl80] M. L. Platonov: *Combinatorial Numbers of a Class of Mappings and Their Applications*. Nauka, Moscow, 1980. [Russian]
- [Wi93] H. Wilf: *The asymptotic behavior of the Stirling numbers of the first kind*. J. of Comb. Theory, Series A **64**, (1993), 344–349.

LEHRSTUHL A FÜR MATHEMATIK, RWTH AACHEN UNIVERSITY, 52056 AACHEN, GERMANY
Email address: `bernhard.heim@rwth-aachen.de`

KUTAI SI INTERNATIONAL UNIVERSITY, 5/7, YOUTH AVENUE, KUTAI SI, 4600 GEORGIA

LEHRSTUHL A FÜR MATHEMATIK, RWTH AACHEN UNIVERSITY, 52056 AACHEN, GERMANY
Email address: `markus.neuhauser@kiu.edu.ge`