

Counting Cycles on Planar Graphs in Subexponential Time

Jin-Yi Cai¹ and Ashwin Maran¹

University of Wisconsin-Madison, USA

Abstract. We study the problem of counting all cycles or self-avoiding walks (SAWs) on triangulated planar graphs. We present a subexponential $2^{O(\sqrt{n})}$ time algorithm for this counting problem. Among the technical ingredients used in this algorithm are the planar separator theorem and a delicate analysis using pairs of Motzkin paths and Motzkin numbers. We can then adapt this algorithm to uniformly sample SAWs, in subexponential time. Our work is motivated by the problem of gerrymandered districting maps.

Keywords: counting cycles · planar graphs · planar separator · Motzkin paths

1 Introduction

We consider the problem of counting all (simple) cycles, or self-avoiding walks (SAWs), in a planar undirected graph. well known algorithms by Tiernan [17], Tarjan [16], and Johnson [10] enumerate cycles in a graph, such that it outputs one cycle per polynomial number of steps. However, many planar graphs (e.g., $n \times O(1)$ grid graphs) have $2^{\Omega(n)}$ cycles. Bousquet-Mélou et al. [4] solved the problem of counting SAWs for $n \times n$ grid graphs exactly¹. Dorn et al. [6] solved the decision version of the weighted Hamiltonian cycle problem on planar graphs in $2^{O(\sqrt{n})}$ time using planar branch decomposition, with $n = |V(G)|$, as well as the decision problem of cycles of length $\geq k$ in time $O(2^{13.6\sqrt{k}}n + n^3)$ on planar graphs (see also [3]). We give a $2^{O(\sqrt{n})}$ time algorithm for the problem of counting all cycles in a planar undirected 3-regular graph. Compared to the approaches in [6,3], our algorithm is more elementary in the use of structural graph properties. It is well known [9] that counting leads to uniform sampling. We also extend our algorithm to count the number of ways a planar graph can be partitioned into two simply connected components, in $2^{O(\sqrt{n})}$ time. It is known [14] that this problem cannot be solved in polynomial time unless $\text{NP} = \text{RP}$.

1.1 Overview

A basic ingredient to achieving this time complexity is the planar separator theorem [11,12,2,15,8]. We consider our plane graph H is embedded on a sphere, and let $G = H^*$ be its dual graph. The overall idea is to divide the sphere by a cycle separator of G and solve a more general problem in two smaller topological semi-spheres. We use (a version of) the planar separator theorem to find a planar separator which is a cycle on G . We then use the fact that cycles on G correspond to cuts on the primal H , to derive the number of cycles on the primal graph based on structural properties of certain quantities we recursively compute for the topological semi-spheres.

A Motzkin path has many equivalent definitions. We can think of a (cyclic) Motzkin path as encoding a (cyclic) string on the alphabet set $\{0, (,)\}$ of matching left and right parentheses. We make the crucial observation that each cycle on the original primal graph, being self-avoiding, intersects with the cycle separator in a way that corresponds to a Motzkin path on the cycle separator (with the symbol “0” indicating non-crossing at that location). Therefore the number of cycles is intimately related to the Motzkin number which counts Motzkin paths on the cut generated by the cycle separator. It is well known [1,13] that the

¹ The algorithm in [4] is applicable for grid graphs only, and it was explicitly calculated that the number of self avoiding walks connecting two diagonal corners in a 19×19 grid graph is $1523344971704879993080742810319229690899454255323294555776029866737355060592877569255844 > 10^{88}$. Our algorithm for planar graphs is based on a recursive, thus different, approach.

Motzkin number of a cycle of size k is $2^{O(k)}$, and that the Motzkin paths can be enumerated. Note that our separator has size $k = O(\sqrt{n})$. We use this fact to efficiently keep track of all possible ways that cycles on the primal graph can intersect with the cut-set generated by the cyclic planar separator.

We then need to somehow count the number of cycles on the primal graph that intersects with the cut-set in a way specified by a Motzkin path. We look at each of the two subgraphs that the planar separator divides the dual graph into. We would like to count the number of traces that are the intersections of cycles on each of these subgraphs that traverse the cut-set specified by a Motzkin path, and then combine them to count the number of cycles on the original primal graph. However, this problem turns out to be a little tricky. It turns out that just keeping track of the intersection of the cycles with the cut-set is not enough. Given a trace which is an intersection of a cycle with one of the two subgraphs, it is possible to have extensions into the subgraph on the other semi-sphere, that form cycles, but that are topologically distinct (see Fig. 3). This complication makes it harder to disentangle what happens in the two subgraphs, and in turn, this makes it harder to synthesize the counts from the two subgraphs and find the number of cycles on the original graph.

We solve this problem by keeping track of not just the intersections of the traces with the cut-set, but also the *order* in which they are crossed. At face value, this seems to introduce an additional factor $k!$ for k crossings. This would ruin our $2^{O(\sqrt{n})}$ time bound. However, we argue that by using a pair of Motzkin paths this difficulty can be overcome. Abstractly, suppose $K, K' \subset S^2$ are homeomorphic to S^1 and they intersect at k points transversely, then there are only $2^{O(k)}$ many topologically distinct ways this can happen (see Remark 1).

In Section 2, we introduce necessary definitions and notations, and we show in Section 3.1 that the problem of counting cycles on a 3-regular planar graph can be reduced to the problem of counting self-avoiding walks (SAWs) on subgraphs that pass through some fixed edges of the subgraph in a given (cyclic) order. In Section 3.2, we show that this problem can be further reduced to counting cycles on a smaller planar graph, that pass through some fixed edges in a given order. In Section 3.3, we inductively solve this problem, and show that the number of cycles on a planar graph can be counted in subexponential time. There is a 1-1 correspondence between cycles and partitions of a planar graph (on S^2) into two simply connected regions (viewed in the dual graph). In Section 4, we extend the algorithm to count the number of partitions of a planar graph into two simply connected regions. It is well known that counting leads to uniform sampling [9]. Our work is motivated by the problem of detecting and certifying gerrymandered districting maps. Sampling can provide a provably unbiased argument that a given gerrymandered map is an extreme outlier as measured by various fairness criteria; see [7].

2 Self-Avoiding Walks and Partial Self-Avoiding Walks

Let H be a connected 3-regular planar graph with an embedding on the sphere S^2 . Our technical problem is to count the number of cycles on the graph H . (Our graph H can have parallel edges but no loops.) We denote by $G = (V, E, F)$ the plane dual graph of H , and denote H by $H = (\tilde{F}, \tilde{E}, \tilde{V})$ (see Remark 2 in the Appendix A). We will use induction on this graph G to find the number of cycles on H . So, we will define the set of cycles on the graph H using its dual graph G as index.

Definition 1. *A simple path, or self-avoiding walk (SAW), in H is a vertex (and edge) disjoint path (also a sequence of distinct adjacent triangular faces in G). A cycle is a special case of SAW with the same beginning and ending vertex. Denote by \mathcal{W}_G the set of all cycles on the graph H .*

While our goal is to count cycles on H , it turns out that cycles on G are extremely helpful. Let $A = (V_A, E_A)$ be any cycle in G . This cycle divides S^2 into two parts D_1 and D_2 , each homeomorphic to a disc with A as its boundary. Thus, this cycle A gives rise to two subgraphs $G_{D_i} = (V_{D_i}, E_{D_i}, F_{D_i})$ ($i = 1, 2$) on these two discs.

Let D be one of D_1 or D_2 , and let D' be the other one. Consider the graph G_D , and its dual H_D . Note that H_D has a special vertex v_∞ (of degree $|V_A| = |E_A|$) which corresponds to the disc D' on the sphere. Suppose $\{u, v\} \in \tilde{E}$ is an edge from H , such that $u \in \tilde{F}_D$ but $v \in \tilde{F}_{D'}$. This is precisely when $\{u, v\} \in \tilde{E}_A$.

Now we modify H_D as follows: We remove v_∞ and its incident edges, and for each $\{u, v\} \in \tilde{E}_A$ with $v \in \tilde{F}_D$, we add a *dangling edge* $\{u, v^*\}$ to \tilde{E}_D with v^* being a new *dangling vertex* (see [Remark 3](#) in the [Appendix A](#)).

Since A is a cycle on the graph G , the edges \tilde{E}_A in the graph H act as a cut between the vertices in \tilde{F}_{D_1} and \tilde{F}_{D_2} (see [Fig. 1](#)). Therefore, if any $W \in \mathcal{W}_G$ has to go from \tilde{F}_{D_1} to \tilde{F}_{D_2} , it has to go across \tilde{E}_A (a consequence of the Jordan Curve Theorem). Also the fact that W is non-self-intersecting places strong constraints on how it will interact with \tilde{E}_A .




Fig. 1: The red cycle A dividing the graph G embedded on S^2 , and generating the two subgraphs G_{D_1} and G_{D_2} .

Let $W \in \mathcal{W}_G$, and let $P = W \cap \tilde{E}_D$, for $D \in \{D_1, D_2\}$. Then for this D we will use this set of edges P in H to define a function ℓ_P on $E_A \subseteq E$. Consider any edge $e \in E_A$. Let its corresponding edge in the graph H be $\tilde{e} \in \tilde{E}_A \subseteq \tilde{E}$. If $\tilde{e} \notin P$, then we define $\ell_P(e) = 0$. Suppose $\tilde{e} \in P$.

Lemma 1. *Let $\tilde{e} = \{u_1, u_2\} \in P \cap \tilde{E}_A$. Without loss of generality $u_1 \notin \tilde{F}_D$, and $u_2 \in \tilde{F}_D$. Then there exist unique vertices $u_3, \dots, u_{k-1} \in \tilde{F}_D$, and $u_k \notin \tilde{F}_D$ ($k \geq 3$), such that $\{u_i, u_{i+1}\} \in P$ for $1 \leq i < k$.*

The proof can be found in the [Appendix A](#).

Let $\tilde{e}' = \{u_{k-1}, u_k\}$ be the edge such that $u_{k-1} \in \tilde{F}_D$ and $u_k \notin \tilde{F}_D$ as guaranteed by [Lemma 1](#). We note that $\tilde{e}' \neq \tilde{e}$ (see [Remark 4](#) in the [Appendix A](#)). Then $\tilde{e}' \in P \cap \tilde{E}_A$, and let $e' \in E_A$ be the corresponding edge in E_A . Now we define $\ell_P(e) = e'$, which completes the definition of the function $\ell_P : E_A \rightarrow E_A \cup \{0\}$.

For any $P = W \cap \tilde{E}_D$ where $W \in \mathcal{W}_G$, the function ℓ_P represents a set of non-intersecting chords on the cycle A . Now, let us characterize the set of all functions that represent non-intersecting chords on the cycle A . Consider the following three properties for a function $\ell : E_A \rightarrow E_A \cup \{0\}$:

1. For any $e \in E_A$, $\ell(e) \neq e$.
2. For any $e, e' \in E_A$, $\ell(e) = e' \iff \ell(e') = e$.
3. For any $e, e' \in E_A$, if $\ell(e) = e'$, then edges e, e' partition the cycle E_A into two segments $\sigma_1 \sqcup \sigma_2 = E_A \setminus \{e, e'\}$ such that for no two edges $e_1 \in \sigma_1, e_2 \in \sigma_2$ is $\ell(e_1) = e_2$.

Let \mathcal{L}_A be the set of all functions $\ell : E_A \rightarrow E_A \cup \{0\}$ that satisfy the above properties. Every ℓ_P defined by $P = W \cap \tilde{E}_D$ for any $W \in \mathcal{W}_G$ belongs to \mathcal{L}_A . Thus,

$$\{\ell_{W \cap \tilde{E}_D} : W \in \mathcal{W}_G\} \subseteq \mathcal{L}_A, \quad \forall D \in \{D_1, D_2\}.$$

Each $\ell \in \mathcal{L}_A$ represents a Motzkin path on E_A (see [Lemma 10](#) in the [Appendix A](#)), and $|\mathcal{L}_A|$ is a Motzkin number [\[13, 1\]](#). The following is a well known fact about Motzkin numbers.

Fact 1. *The number of Motzkin paths of length n is $2^{O(n)}$. Thus, there exists a constant κ such that ² for any planar triangulated graph G and a cycle A on G , $|\mathcal{L}_A| \leq \kappa^{|E_A|}$.*

² It is known that asymptotically we have $|\mathcal{L}_A| \leq O(3^{|E_A|})$.

Now, let

$$\mathcal{C}(\ell_1, \ell_2) = \{W \in \mathcal{W}_G : \ell_{W \cap \tilde{E}_{D_1}} = \ell_1 \text{ and } \ell_{W \cap \tilde{E}_{D_2}} = \ell_2\}.$$

The following lemma shows that $|\mathcal{W}_G|$ is determined by $|\mathcal{C}(\ell_1, \ell_2)|$ for all $\ell_1, \ell_2 \in \mathcal{L}_A$.

Lemma 2.

$$|\mathcal{W}_G| = \sum_{\ell_1 \in \mathcal{L}_A} \sum_{\ell_2 \in \mathcal{L}_A} |\mathcal{C}(\ell_1, \ell_2)|.$$

The proof can be found in the [Appendix A](#).

Now, let us see how we can compute $|\mathcal{C}(\ell_1, \ell_2)|$. Essentially, for $\ell_1, \ell_2 \in \mathcal{L}_A$, we will count $|\{W \cap \tilde{E}_{D_1} : W \in \mathcal{W}_G, \ell_{W \cap \tilde{E}_{D_1}} = \ell_1\}|$ and $|\{W \cap \tilde{E}_{D_2} : W \in \mathcal{W}_G, \ell_{W \cap \tilde{E}_{D_2}} = \ell_2\}|$. With some care, we will be able to combine them to find the value of $|\mathcal{C}(\ell_1, \ell_2)|$.

In order to formalize this notion, we will now define *partial self-avoiding walks* (PSAW) on the disc D (see [Remark 5](#) in the [Appendix A](#)). Let $\ell \in \mathcal{L}_A$ be a labelling. We saw that ℓ represents a Motzkin path, which corresponds to a set of non-intersecting chords on E_A . Let D and D' be as before. We will view this labelling ℓ as being embedded in the disc D' . For any $e \in E_A$, we have a dangling edge \tilde{e} incident to a newly created dangling vertex v^* not in \tilde{F}_D , which we will denote by $\pi_D(e)$. If $\ell(e) = e'$ we also have a dangling vertex $\pi_D(e')$.

Now, let

$$\mu_D(\ell) = \{\{\pi_D(e), \pi_D(e')\} : e, e' \in E_A \text{ such that } \ell(e) = e' \text{ and } \ell(e') = e\}.$$

We can think of the dangling vertices as belonging to the other disc D' (see [Fig. 6](#) in [Appendix A](#) for an example of the dangling edges). Then, $\mu_D(\ell)$ can be viewed as a set of edges between these dangling vertices that are embedded on D' .

Definition 2. Let D be one of D_1 or D_2 . A set of edges $P \subseteq \tilde{E}_D$ is called a *partial self-avoiding walk* (PSAW) on the graph H_D , if there exists an $\ell \in \mathcal{L}_A$, such that $P \cup \mu_D(\ell)$ is a cycle. Denote by \mathcal{P}_D the set of all PSAW on H_D .

Note that P is a set of edges in \tilde{E}_D , some of which may be incident on the dangling vertices which are thought to be in D' , and $\mu_D(\ell)$ is a set of edges between these vertices as described above. (Despite the notation $\mu_D(\ell)$, intuitively one should think of $\ell \in \mathcal{L}_A$ as referring to pairings that take place in D' . See the proof of [Lemma 3](#) in the [Appendix A](#).) But we observe that the definition of P being a PSAW does not explicitly refer to the existence of some cycle $W \in \mathcal{W}_G$.

Lemma 3. For any planar triangulated graph G and any $D \in \{D_1, D_2\}$ determined by a cycle A on G ,

$$\{W \cap \tilde{E}_D : W \in \mathcal{W}_G\} \subseteq \mathcal{P}_D.$$

The proof can be found in the [Appendix A](#). An example of $W \cap \tilde{E}_D \in \mathcal{P}_D$ can also be seen in [Fig. 6](#) in [Appendix A](#).

More importantly, as a corollary of [Lemma 1](#), given $P \in \mathcal{P}_D$ for any $D \in \{D_1, D_2\}$, since there exists an $\ell \in \mathcal{L}_A$ such that $P \cup \mu_D(\ell)$ is a cycle with $(P \cup \mu_D(\ell)) \cap \tilde{E}_D = P$, we see that ℓ_P is well-defined and $\ell_P \in \mathcal{L}_A$. We can then define, for any $\ell \in \mathcal{L}_A$,

$$\mathcal{C}_D(\ell) = \{P \in \mathcal{P}_D : \ell_P = \ell\}.$$

Note that, if $\mathcal{C}(\ell_1, \ell_2) \neq \emptyset$, then ℓ_1 and ℓ_2 must assign the same set of edges $e \in E_A$ to nonzero, even though the pairings by ℓ_1 and ℓ_2 are typically different. Thus the nonzero places of ℓ_1 and ℓ_2 mutually determine each other. Our idea is to try to compute $|\mathcal{C}_{D_1}(\ell_1)|$ and $|\mathcal{C}_{D_2}(\ell_2)|$, and then to use it to compute $|\mathcal{C}(\ell_1, \ell_2)|$. Intuitively, we will use ℓ_2 to “stitch” up the disc D_2 in H_{D_1} to construct a new planar triangulated graph G_1 (see [Fig. 2](#) for an example, a more detailed discussion on the exact procedure to be followed can be found in [Section 3.2](#)). If we can somehow count all the cycles in the graph H_1 that passes through certain special edges, we will be able to compute $|\{P \in \mathcal{P}_{D_1} : P \cup \mu_{D_1}(\ell_2) \text{ is a cycle}\}|$.




Fig. 2: During the stitching process, edges e, e' such that $\ell_2(e) = e'$ and $\ell_2(e') = e$ are identified with each other. SAWs in the original graph that induce the labelling ℓ_2 on D_2 correspond to SAWs in the stitched graph that pass through the edges that were identified.

However, as the example in Fig. 3 shows, given $\ell_2 \in \mathcal{L}_A$, there can be multiple ℓ_1 such that $P \cup \mu_{D_1}(\ell_2)$ is a cycle for all $P \in \mathcal{C}_{D_1}(\ell_1)$. Therefore, the stitching alone unfortunately does not allow us to compute $|\mathcal{C}_{D_1}(\ell_1)|$ directly. However, the next key insight is that given $\ell_1, \ell_2 \in \mathcal{L}_A$ with $\mathcal{C}(\ell_1, \ell_2) \neq \emptyset$, any cycle $W \in \mathcal{C}(\ell_1, \ell_2)$ crosses the same set of edges from \tilde{E}_A , and more importantly, these edges from \tilde{E}_A are crossed in the same cyclic order (upto reversal) by all cycles from $\mathcal{C}(\ell_1, \ell_2)$ (see Remark 6 in the Appendix A).




Fig. 3: In this graph G , the blue cycles are $W_1, W_2 \in \mathcal{W}_G$. $\ell_{W_1 \cap \tilde{E}_{D_1}} \neq \ell_{W_2 \cap \tilde{E}_{D_1}}$, but $\ell_{W_1 \cap \tilde{E}_{D_2}} = \ell_{W_2 \cap \tilde{E}_{D_2}}$. Therefore, given a labelling $\ell_2 \in \mathcal{L}_A$, there can be multiple ℓ_1 such that $P \cup \mu_{D_1}(\ell_2)$ is a cycle for all $P \in \mathcal{C}_{D_1}(\ell_1)$.

For any two adjacent vertices u, v in H connected by an edge $\{u, v\}$ (or two triangular faces of G that share an edge), we use $(u \rightarrow v)$ to refer to an *arc* in H from u to v in that direction. Given an arc $\tilde{a} = (u \rightarrow v)$ in H , let $\varepsilon(\tilde{a}) \in \tilde{E}$ denote the underlying edge $\{u, v\}$. Let $S = (\tilde{a}_1, \dots, \tilde{a}_k)$ be an ordered set of arcs in the graph H . We denote by $\varepsilon(S) = \{\varepsilon(\tilde{a}_1), \dots, \varepsilon(\tilde{a}_k)\}$ the unordered set of edges corresponding to the arcs in S .

Definition 3. Let $S = ((u_1 \rightarrow v_1), \dots, (u_k \rightarrow v_k))$ be an ordered set of arcs in H . We say that a cycle W conforms to S if $\varepsilon(S) \subseteq W$, and if in one cyclic traversal of the cycle W we encounter the vertices $u_1, v_1, u_2, v_2, \dots, u_k, v_k$ in this order.

Note that Definition 3 is defined in terms of some cyclic traversal of W . If W conforms to S , then W also conforms to $((u_2 \rightarrow v_2), \dots, (u_k \rightarrow v_k), (u_1 \rightarrow v_1))$, as well as its reversal.

We can define a relation \sim on ordered sets of arcs as follows:

$$((u_1 \rightarrow v_1), (u_2 \rightarrow v_2), \dots, (u_k \rightarrow v_k)) \sim ((u_2 \rightarrow v_2), \dots, (u_k \rightarrow v_k), (u_1 \rightarrow v_1)).$$

$$((u_1 \rightarrow v_1), \dots, (u_k \rightarrow v_k)) \sim ((v_k \rightarrow u_k), \dots, (v_1 \rightarrow u_1)).$$

For brevity we will also denote by \sim its reflexive, symmetric, and transitive closure. Thus, for $S \sim T$, we have W conforms to S iff W conforms to T .

We will now define a few terms to refer to cycles that conform to a given ordered set of arcs.

Definition 4. Let S be an ordered set of arcs in H . We define

$$\mathcal{W}_{G,S} = \{W \in \mathcal{W}_G : W \text{ conforms to } S\}.$$

For any $\ell_1, \ell_2 \in \mathcal{L}_A$,

$$\mathcal{C}_S(\ell_1, \ell_2) = \{W \in \mathcal{W}_{G,S} : \ell_{W \cap \tilde{E}_{D_1}} = \ell_1 \text{ and } \ell_{W \cap \tilde{E}_{D_2}} = \ell_2\}.$$

Definition 5. Let $D \in \{D_1, D_2\}$. Let S be an ordered subset of arcs in H_D such that $\varepsilon(S) \subseteq \tilde{E}_D$. We define

$$\mathcal{P}_{D,S} = \{P \in \mathcal{P}_D : P \cup \mu_D(\ell) \text{ conforms to } S, \text{ for some } \ell \in \mathcal{L}_A\}.$$

Finally, for any $\ell \in \mathcal{L}_A$,

$$\mathcal{C}_{D,S}(\ell) = \{P \in \mathcal{P}_{D,S} : \ell_P = \ell\}.$$

Our main technical result in this paper is an algorithm to compute $|\mathcal{W}_{G,S}|$ for a general ordered set S of arcs in H . This will give us a subexponential time algorithm to compute $|\mathcal{W}_G|$.

3 The Algorithm

Continuing from the previous section, consider an ordered set of arcs S from the graph H . Our overall design of the algorithm to compute $|\mathcal{W}_{G,S}|$ is a divide-and-conquer strategy, and the following planar separator theorem of Lipton and Tarjan [11] plays a pivotal role. Here we use a version of the theorem due to Alon et al. using a simple cycle separator [12, 2, 15, 8]³.

Theorem 1. Given a planar triangulated graph $G = (V, E, F)$, there exists a cycle $A = (V_A, E_A)$ in G with vertices $V_A \subseteq V$ and edges $E_A \subseteq E$ of size $|V_A| = |E_A| \leq \sqrt{8|V|}$, which partitions the vertices $V \setminus V_A$ into two parts V_B, V_C such that $|V_B|, |V_C| \leq \frac{2|V|}{3}$. Moreover, this cycle A can be found in polynomial time, as a function of $|V|$.

We use Theorem 1 to find the cycle A on the graph G . We may assume that the vertices $V_A \cup V_B$ belong to the disc D_1 , and the vertices $V_A \cup V_C$ belong to the disc D_2 . The following is a generalization of Lemma 2; the proof is a simple adaptation. It implies that if we can compute $|\mathcal{C}_S(\ell_1, \ell_2)|$ for all $\ell_1, \ell_2 \in \mathcal{L}_A$, we will be able to compute $|\mathcal{W}_{G,S}|$.

Lemma 4.

$$|\mathcal{W}_{G,S}| = \sum_{\ell_1 \in \mathcal{L}_A} \sum_{\ell_2 \in \mathcal{L}_A} |\mathcal{C}_S(\ell_1, \ell_2)|.$$

3.1 Computing $|\mathcal{C}_S(\ell_1, \ell_2)|$

We now need to compute $|\mathcal{C}_S(\ell_1, \ell_2)|$. Given labellings $\ell_1, \ell_2 \in \mathcal{L}_A$, we will say they are *compatible* if they pass the test defined by the algorithm **CompatibilityCheck**.

We note that if the labellings ℓ_1, ℓ_2 are defined by some $W \in \mathcal{W}_{G,S}$ then they are compatible. The test is trying to reflect how $W \in \mathcal{C}_S(\ell_1, \ell_2)$ crosses \tilde{E}_A and forms a cycle (see Fig. 4). We use a pair of Motzkin paths to get around the difficulty described in Fig. 3. While **CompatibilityCheck** is not sufficient to conclude that $|\mathcal{C}_S(\ell_1, \ell_2)| > 0$, it is a necessary condition. By dealing only with pairs of Motzkin paths of length k , which have order $2^{O(k)}$, we avoid the complexity of order $k!$.

³ For simplicity in stating our results, we are not using optimal constants, see [2, 15, 5] for more details.




Fig. 4: Given $\ell_2 \in \mathcal{L}_A$, we see two examples $\ell_1, \ell_1' \in \mathcal{L}_A$ that are compatible with ℓ_2 , but one example ℓ_1'' is not.

Algorithm $\text{CompatibilityCheck}(G = (V, E, F), A = (V_A, E_A), \ell_1 \in \mathcal{L}_A, \ell_2 \in \mathcal{L}_A)$

```

1: if  $\{e \in E_A : \ell_1(e) = 0\} \neq \{e \in E_A : \ell_2(e) = 0\}$  then
2:   return false
3: if  $\ell_1(e) = 0 \forall e \in E_A$  then
4:   return true
5: Pick  $e \in E_A$  such that  $\ell_1(e) \neq 0$ 
6:  $M \leftarrow \{e\}$ 
7: while  $\ell_1(e) \notin M$  do
8:    $e' \leftarrow \ell_1(e)$ 
9:    $e \leftarrow \ell_2(e')$ 
10:   $M \leftarrow M \cup \{e', e\}$ 
11: if  $M \neq \{e \in E_A : \ell_1(e) \neq 0\}$  then
12:   return false
13: return true

```

Lemma 5. *If $\mathcal{C}_S(\ell_1, \ell_2) \neq \emptyset$, then ℓ_1 and ℓ_2 are compatible as defined by $\text{CompatibilityCheck}$.*

The proof can be found in the [Appendix B](#).

Combining [Lemma 5](#) and [Lemma 4](#), we have

$$|\mathcal{W}_{G,S}| = \sum_{\text{compatible } \ell_1, \ell_2 \in \mathcal{L}_A} |\mathcal{C}_S(\ell_1, \ell_2)|.$$

Now consider two compatible labellings ℓ_1 and ℓ_2 . We deal with the special case, where $\ell_1 = \ell_2 = 0$ in [Lemma 11](#) in [Appendix B](#). We will instead consider here, the more interesting case where $(\ell_1, \ell_2) \neq (0, 0)$. Before we see how to compute $|\mathcal{C}_S(\ell_1, \ell_2)|$, let us first tackle the simpler problem of computing $|\mathcal{C}_\emptyset(\ell_1, \ell_2)|$. We know that any cycle $W \in \mathcal{C}_\emptyset(\ell_1, \ell_2)$ satisfies the following condition:

$$W \cap \tilde{E}_A = \{\tilde{e} \in \tilde{E}_A : \ell_1(e) \neq 0\} = \{\tilde{e} \in \tilde{E}_A : \ell_2(e) \neq 0\}.$$

So, let us consider $\{\tilde{e} \in \tilde{E}_A : \ell_1(e) \neq 0\}$, and assume it is nonempty. We can now create an ordered set $T = T(\ell_1, \ell_2)$ of arcs on H , representing the order in which any $W \in \mathcal{C}_\emptyset(\ell_1, \ell_2)$ would go across these edges. We construct T as follows: First, we arbitrarily pick some $e_1 \in E_A$ such that $\ell_1(e_1) \neq 0$. Let $\tilde{e}_1 = \{u_1, v_1\}$, such that $u_1 \in \tilde{F}_{D_2}, v_1 \in \tilde{F}_{D_1}$. Then, we add the arc $(u_1 \rightarrow v_1)$ to T . Next, let $e_2 = \ell_1(e_1)$ and let $\tilde{e}_2 = \{u_2, v_2\}$, such that $u_2 \in \tilde{F}_{D_1}, v_2 \in \tilde{F}_{D_2}$. Then we add the arc $(u_2 \rightarrow v_2)$ as the next element to T . We then move on to $e_3 = \ell_2(e_2)$. If $e_3 = e_1$ then T is completed; otherwise, let $\tilde{e}_3 = \{u_3, v_3\}$, such that $u_3 \in \tilde{F}_{D_2}, v_3 \in \tilde{F}_{D_1}$. Then we add $(u_3 \rightarrow v_3)$ to T . We repeat this process just as in the algorithm $\text{CompatibilityCheck}$, until an edge is visited again, at which point we stop and T is completed.

As a cyclically ordered set of arcs under the equivalence of \sim , the definition of T does not depend on the choice of $e_1 \in E_A$. It is important to note that the ordered set T is a function of the labellings ℓ_1 and ℓ_2 ; a crucial point is that a compatible pair (ℓ_1, ℓ_2) together determines not only the subset of E_A where ℓ_1 and ℓ_2 are nonzero, but also the order they are to be traversed by any $W \in \mathcal{C}_\emptyset(\ell_1, \ell_2)$. This T represents the order and orientation in which these edges on \tilde{E}_A will be visited in a cyclic traversal by any $W \in \mathcal{C}_\emptyset(\ell_1, \ell_2)$.

When there is a need to indicate this dependence, we will use $T(\ell_1, \ell_2)$ to denote this ordered set of arcs (see [Remark 7](#) in the [Appendix B](#)).

Lemma 6. *Given a compatible $(\ell_1, \ell_2) \neq (0, 0)$,*

$$|\mathcal{C}_\emptyset(\ell_1, \ell_2)| = |\mathcal{C}_{D_1, T}(\ell_1)| \times |\mathcal{C}_{D_2, T}(\ell_2)|.$$

The proof can be found in [Appendix B](#).

Remark 1. Going back to the statement at the end of [Section 1](#) that two cycles homeomorphic to S^1 intersecting at k points transversely can have only $2^{O(k)}$ many topologically distinct ways, we have shown how to handle this problem using two compatible Motzkin paths. We showed that to compute $|\mathcal{C}_{D_1}(\ell_1)|$, we do not need to find the union of $\mathcal{C}_{D_1, T'}(\ell_1)$ over all ordered sets T' such that $\varepsilon(T') = \{\tilde{e} \in \tilde{E}_A : \ell_1(e) \neq 0\}$. If $|\varepsilon(T')| = k$ then there are up to $k!$ many ordered sets T' to consider. Instead, we can restrict our attention to just $\{T(\ell_1, \ell_2) : \ell_2 \in \mathcal{L}_A \text{ compatible with } \ell_1\}$, which has size only $2^{O(k)}$. On the other hand, to achieve this saving, the definition of these sets and the underlying concept become more delicate and we have had to use some more involved notations.

We have successfully reduced the problem of computing $|\mathcal{C}_\emptyset(\ell_1, \ell_2)|$ to the problem of computing $|\mathcal{C}_{D, T}(\ell)|$, as promised at the beginning of this section. Now, we need to extend this to the problem of computing $|\mathcal{C}_S(\ell_1, \ell_2)|$, for a general $S \neq \emptyset$.

Given an ordered set S of arcs in H , and the ordered set T described above, we now need to somehow combine these two ordered sets into one. Informally, we say that I is an interleaving of S and T , if I contains the arcs from S and T in the same cyclic order upto reversals. The main idea here is that if a cycle W conforms to both S and T , it will have to conform to some interleaving I of S and T as well. In order to formally define it, we will need the equivalence relation \sim from earlier. We say an ordered set I of arcs in H is an *interleaving* of S and T if the following conditions hold:

1. $\varepsilon(I) = \varepsilon(S) \cup \varepsilon(T)$.
2. $I \cap \varepsilon(S) \sim S$, and $I \cap \varepsilon(T) \sim T$.

Here, $I \cap \varepsilon(S)$ denotes the ordered set of arcs on the underlying (unordered) edge set $\varepsilon(S)$ with the induced orientation and order from I , and $I \cap \varepsilon(T)$ is similarly defined. Note that in item 2 the equivalence relation \sim allows taking reversals independently for S and T . We then use $\mathcal{I}_{S, T}$ to denote the set of all the equivalence classes of interleavings of S and T with each equivalence class represented by a single interleaving.

Now that we have defined interleavings, we can show that

$$|\mathcal{C}_S(\ell_1, \ell_2)| = \sum_{I \in \mathcal{I}_{S, T}} |\mathcal{C}_I(\ell_1, \ell_2)|.$$

The formal proof of this claim can be found as [Lemma 15](#) in [Appendix B](#).

We showed in [Lemma 5](#) that incompatible labelling pairs $\ell_1, \ell_2 \in \mathcal{L}_A$ can be omitted, since in that case $\mathcal{C}_S(\ell_1, \ell_2) = \emptyset$. Going one step further, even if $\ell_1, \ell_2 \in \mathcal{L}_A$ are compatible we may still have $\mathcal{C}_I(\ell_1, \ell_2) = \emptyset$ if I is not an appropriate interleaving of S and $T = T(\ell_1, \ell_2)$. Now, we will define what a *valid* interleaving is, and then show that $\mathcal{C}_I(\ell_1, \ell_2) = \emptyset$ for the invalid ones.

Definition 6. *Let $\ell_1, \ell_2 \in \mathcal{L}_A$ be compatible labellings, and $T = T(\ell_1, \ell_2)$. We say $I \in \mathcal{I}_{S, T}$ is a valid interleaving of S and T if the following properties hold:*

1. *There exists some ordered subset T' of T such that*

$$S \cap \tilde{E}_A \sim T'.$$

2. *If $((u_1^T \rightarrow v_1^T), (u_1^S \rightarrow v_1^S), \dots, (u_k^S \rightarrow v_k^S), (u_2^T \rightarrow v_2^T))$ are consecutive arcs in $I' \sim I$ such that $\{\{u_1^T, v_1^T\}, \{u_2^T, v_2^T\}\} \subseteq I' \cap \varepsilon(T)$, and $\{\{u_1^S, v_1^S\}, \dots, \{u_k^S, v_k^S\}\} \subseteq I' \cap (\varepsilon(S) \setminus \varepsilon(T))$, then*

$$u_1^T, v_2^T \in \tilde{F}_D, \text{ and } v_1^T, u_1^S, v_1^S, \dots, u_k^S, v_k^S, u_1^T \notin \tilde{F}_D \text{ for some } D \in \{D_1, D_2\}.$$

Informally, condition (1) says that the subset of S that are on \tilde{E}_A , must appear in T in the right order. Condition (2) constrains that walks in $\mathcal{C}_I(\ell_1, \ell_2)$ can only jump between discs D_1 and D_2 using the arcs from T . Formally, we have the following lemma, whose proof can be found in [Appendix B](#).

Lemma 7. *Given compatible labellings $\ell_1, \ell_2 \in \mathcal{L}_A$ such that $(\ell_1, \ell_2) \neq (0, 0)$,*

$$|\mathcal{C}_S(\ell_1, \ell_2)| = \sum_{\text{valid } I \in \mathcal{I}_{S,T}} |\mathcal{C}_I(\ell_1, \ell_2)|,$$

where $T = T(\ell_1, \ell_2)$.

Finally, we have the main lemma of this section, which proves that the interleavings allows us to circumvent the issue we observed in [Fig. 3](#).

Lemma 8. *Given compatible $\ell_1, \ell_2 \in \mathcal{L}_A$ such that $(\ell_1, \ell_2) \neq (0, 0)$, and $T = T(\ell_1, \ell_2)$,*

$$|\mathcal{C}_S(\ell_1, \ell_2)| = \sum_{\text{valid } I \in \mathcal{I}_{S,T}} \left| \mathcal{C}_{D_1, I \cap \tilde{E}_{D_1}}(\ell_1) \right| \times \left| \mathcal{C}_{D_2, I \cap \tilde{E}_{D_2}}(\ell_2) \right|.$$

The proof can be found in [Appendix B](#).

We have now succeeded in reducing the problem of computing $|\mathcal{W}_{G,S}|$ to the problem of computing $|\mathcal{C}_{D, I \cap \tilde{E}_D}(\ell)|$.

3.2 Computing $|\mathcal{C}_{D,S}(\ell)|$

In this section, given compatible (ℓ_1, ℓ_2) (including the case $(\ell_1, \ell_2) = (0, 0)$), and valid $I \in \mathcal{I}_{S,T}$, we will construct two graphs G_1, G_2 , and two ordered sets S_1, S_2 of arcs on the duals H_1, H_2 , and use $|\mathcal{W}_{G_1, S_1}|$ and $|\mathcal{W}_{G_2, S_2}|$ to compute $|\mathcal{C}_{D_1, I \cap \tilde{E}_{D_1}}(\ell_1)|$, and $|\mathcal{C}_{D_2, I \cap \tilde{E}_{D_2}}(\ell_2)|$ respectively.

The reason why we want this reduction is that G_1 and G_2 will have substantially fewer vertices than G , and S_1 and S_2 will not be much larger than S . Therefore, this reduction will allow us to reduce the problem of computing the number of cycles on a graph G to that of computing the number of cycles on smaller graphs.

We will briefly describe the construction of G_1 here. Intuitively, we will use the labelling ℓ_2 to *stitch up* the disc D_2 . The construction of G_2 will be analogous, using ℓ_1 to stitch up the disc D_1 .

We start with the graph G_{D_1} , which is just the graph G restricted to the disc D_1 . In other words, it is the plane graph containing the vertices V_{D_1} , the edges E_{D_1} , as well as the faces $F_{D_1} \cup \{f^*\}$, where f^* is the face that is bounded by the edges E_A , which can be thought of as the disc D_2 without internal edges in our spherical embedding of G . We will then repeatedly use the three contractions seen in [Fig. 5](#) to shrink the cycle E_A away, until the face f^* disappears (see [Fig. 2](#) for an example of this stitching).

We use single-edge contractions (see [Fig. 5a](#)) to shrink away edges of E_A for which $\ell_2(e) = 0$, and edge-pair contractions (see [Fig. 5b](#)) to merge together edges e, e' that are such that $\ell_2(e) = e'$, and $\ell_2(e') = e$. We also add the merged edge to the ordered set of arcs S_1 , so that we only count cycle that cross this edge in the correct orientation. Finally, parallel-edge contractions (see [Fig. 5c](#)) are used to ensure that the resulting graph G_1 has no loops. As $|E_A|$ eventually shrinks and disappears, the face f^* disappears, and we are left with the graph G_1 .

We will leave the details of the exact construction of G_1, S_1 and G_2, S_2 , including the exact definitions of the contractions, and the order in which they are to be applied, to [Appendix C](#). The construction of these graphs allows us to prove the following theorem. Its proof can also be found in [Appendix C](#).

Theorem 2. *Given a graph $G = (V, E, F)$ with $|V| = \Omega(1)$ with a cycle A dividing into two discs D, D' , a labelling $\ell \in \mathcal{L}_A$, and a set I of ordered arcs such that $\varepsilon(I) \subseteq \tilde{E}_D$, we can compute $|\mathcal{C}_{D, I \cap \tilde{E}_D}(\ell)|$ in polynomial time, or construct a graph $G_1 = (V_1, E_1, F_1)$, and an ordered set of arcs S_1 , such that $|V_1| \leq \frac{3|V|}{4}$, and $|\mathcal{C}_{D, I \cap \tilde{E}_D}(\ell)|$ can be computed from $|\mathcal{W}_{G_1, S_1}|$ in polynomial time.*




Fig. 5: Examples of the three contractions that will be used to stitch up the discs and construct G_1 and G_2 .

3.3 Computing $|\mathcal{W}_G|$

For the purpose of complexity claims on our algorithm to compute $|\mathcal{W}_G|$, we may assume the number of vertices m of G is greater than a fixed constant; smaller graphs can be handled by brute-force in constant time, which will serve as our base case in an induction. As $\mathcal{W}_G = \mathcal{W}_{G,\emptyset}$, we have

$$|\mathcal{W}_G| = \sum_{\text{compatible } (\ell_1, \ell_2) \neq (0,0)} \left(\sum_{\text{valid } I \in \mathcal{I}_{\emptyset, T(\ell_1, \ell_2)}} \left| \mathcal{C}_{D_1, I \cap \tilde{E}_{D_1}}(\ell_1) \right| \times \left| \mathcal{C}_{D_2, I \cap \tilde{E}_{D_2}}(\ell_2) \right| \right) + \left(\left| \mathcal{C}_{D_1, \emptyset}(0) \right| + \left| \mathcal{C}_{D_2, \emptyset}(0) \right| - 1 \right).$$

As seen in [Theorem 2](#), the problem of computing $\left| \mathcal{C}_{D_1, I \cap \tilde{E}_{D_1}}(\ell_1) \right|$ and $\left| \mathcal{C}_{D_2, I \cap \tilde{E}_{D_2}}(\ell_2) \right|$ can be reduced to the problems of the form of computing $|\mathcal{W}_{G_1, S_1}|$. Now note that if $G_1 = (V_1, E_1, F_1)$, then [Lemma 17](#) says that if $|V_1| \geq 3$, G_1 is a planar triangulated graph with no vertices of degree less than 2, and the dual H_1 of G_1 is a 3-regular planar multi-graph (with no loops). We have therefore reduced the problem of computing $|\mathcal{W}_G|$ to that of computing $|\mathcal{W}_{G_1, S_1}|$ for a number of graphs with at most $\frac{3|V|}{4}$ vertices, or we have reduced it to a problem of $O(1)$ size.

So, when G_1 is large, one can recursively further reduce these smaller problems. This allows us to construct a tree \mathcal{T} with the problem $|\mathcal{W}_G|$ at the root, with the children of each node being the sub-problems that need to be solved to solve the parent problem. The leaves of this tree are problems of $O(1)$ size.

Starting at the root level $|\mathcal{W}_{G_0, S_0}|$ at level 0 with $G_0 = G$ and $S_0 = \emptyset$, a problem $|\mathcal{W}_{G_k, S_k}|$ on level k of the tree \mathcal{T} has $|V_k| = |V(G_k)| \leq m_k = (3/4)^k |V|$. So the height of the tree is $L = O(\log m)$. Moreover, for a problem on level k of \mathcal{T} , the constructed ordered set of arcs for any compatible pair (ℓ_1, ℓ_2) has size

$$|T_k| \leq |E_{A_k}| \leq \sqrt{8m_k}.$$

This is the maximum number of edges that can be added to S_k . Thus,

$$|S_{k+1}| \leq |S_k| + |T_k| \leq |S_k| + t_k$$

where $t_k = \sqrt{8m_k}$. Since $|S_0| = 0$, this means that

$$|S_k| \leq \sum_{0 \leq i < k} t_i < \sum_{i=0}^{\infty} t_i = O(\sqrt{|V|}).$$

We now have the following lemma, whose proof can be found in [Appendix D](#).

Lemma 9. *There is a constant α , such that any problem $|\mathcal{W}_{G_l, S_l}|$ on level l of the tree \mathcal{T} can be computed in time*

$$\alpha^{\sqrt{m_l}} \frac{(t_0 + \dots + t_{L-1})!}{(t_0 + \dots + t_{l-1})!(t_l)!\dots(t_{L-1})!}.$$

Finally, we have our main theorem. Its proof can also be found in [Appendix D](#).

Theorem 3. *There exists a constant β such that given any connected planar 3-regular multi-graph (with no loops) H with n vertices, the number of cycles on H can be computed in $\beta^{\sqrt{n}}$ time.*

4 Extensions

We will now extend the algorithm presented in [Section 3](#) to count the number of ways a planar triangulated graph can be partitioned into two regions such that the faces are simply connected (in other words, the dual graph is simply connected).

Definition 7. Let $X, Y \subseteq \tilde{E}$ be two disjoint subsets. Then,

$$\mathcal{W}_G^{X,Y} = \{W \in \mathcal{W}_G : X \subseteq W, Y \cap W = \emptyset\}.$$

In other words, $\mathcal{W}_G^{X,Y}$ is the set of cycles in H that traverse every edge in X , and do not traverse any edge in Y . We define $\mathcal{P}_D^{X,Y}$, $\mathcal{W}_{G,S}^{X,Y}$, $\mathcal{P}_{D,S}^{X,Y}$, $\mathcal{C}_{D,S}^{X,Y}(\ell)$ and $\mathcal{C}_S^{X,Y}(\ell_1, \ell_2)$ similarly.

The stipulation that cycles in $\mathcal{W}_G^{X,Y}$ must pass through all edges in X (and avoid Y) may appear to be of the same kind as the requirement in $\mathcal{W}_{G,S}$ that they must pass through S . However, there is a crucial difference, namely cycles $W \in \mathcal{W}_{G,S}$ are required to traverse the edges $\varepsilon(S)$ in the directed order as arcs and in the order specified by S , whereas there is no such requirement for $W \in \mathcal{W}_G^{X,Y}$ with respect to X . Our $2^{O(\sqrt{n})}$ time algorithm is for $|\mathcal{W}_{G,\emptyset}|$. With respect to an arbitrary S , no subexponential run time claim was made for $|\mathcal{W}_{G,S}|$ in its dependence on S . However, this time bound $2^{O(\sqrt{n})}$ does hold for counting $|\mathcal{W}_G^{X,Y}|$, which has the same complexity as counting $|\mathcal{W}_{G,\emptyset}|$, and we can extend our algorithm to this setting without much effort. A more detailed discussion of [Theorem 4](#) can be found in [Appendix E](#).

Theorem 4. *There exists a constant β such that given any 3-regular planar multi-graph (with no loops) H with n vertices, and any $X, Y \subseteq \tilde{E}$, the number of cycles on H that traverse each edge in X and do not traverse any edge in Y can be computed in $\beta^{\sqrt{n}}$ time.*

We can apply [Theorem 4](#) to count the number of ways of partitioning a planar triangulated graph into two simply connected graphs (i.e. the faces are connected). Given a graph $H = (\tilde{F}, \tilde{E}, \tilde{V})$ with the border $B = (\tilde{F}_B, \tilde{E}_B)$, we consider an arbitrary pair of vertices $a, b \in \tilde{F}_B$, such that $a \neq b$. Since $a, b \in \tilde{F}_B$ and B is a cycle, by traversing this cycle in two different directions, there exist two SAWs within \tilde{E}_B between the vertices a and b . Let $B_{a,b} \subseteq \tilde{E}_B$ be one of these SAWs with the endpoints a, b .

We can now let $X = B_{a,b}$, and $Y = \{\tilde{e} \in \tilde{E} : \tilde{e} \text{ is incident on } \tilde{F}_B \setminus \{a, b\}\} \setminus X$. Then, we can use [Theorem 4](#) to compute $|\mathcal{W}_G^{X,Y}|$. If we repeat this for all possible pairs of $a \neq b$, we will be done.

Using standard techniques, this exact counting algorithm can also be used to draw samples from the uniform distribution over all partitions into simply connected regions (or cycles in planar graphs). Therefore, this algorithm yields a $2^{O(\sqrt{n})}$ time algorithm to uniformly sample from the space of all partitions of a 3-regular planar graph into two simply connected regions (i.e. such that the faces of the regions are connected, or alternately, the vertices of the two regions in the dual graph are connected).

However, a vast number of boundaries of such partitions are likely to be space filling curves. It is possible to extend the algorithm above so that it only counts partitions where the boundary is not space filling, so we can sample from a space of more natural partitions. A sketch of this extension is to be found in [Appendix E](#).

One restriction of the above algorithm is that it is currently limited to 3-regular graphs. However, the general problem can be reduced to a generalized version of the 3-regular graph problem with only a little effort. We can simply replace all vertices with degree > 3 with a path of vertices all with degree 3. The only issue this causes is that a cycle on this modified graph need not be a cycle on the original graph, as the path corresponding to a single vertex may be visited multiple times by a cycle. However, this can be easily worked around, by making use of the parameters X, Y on the paths that cut across the planar separator A . It can be shown that if a graph has vertices with degrees bounded by some integer d , counting the number of cycles on the graph introduces only an additional factor of $d^{O(\sqrt{n})}$.

References

1. Aigner, M.: Motzkin numbers. *European Journal of Combinatorics* **19**(6), 663–675 (1998)

2. Alon, N., Seymour, P., Thomas, R.: Planar separators. *SIAM Journal on Discrete Mathematics* **7**(2), 184–193 (1994)
3. Bodlaender, H.L., Cygan, M., Kratsch, S., Nederlof, J.: Deterministic single exponential time algorithms for connectivity problems parameterized by treewidth. *Information and Computation* **243**, 86–111 (2015)
4. Bousquet-Mélou, M., Guttmann, A.J., Jensen, I.: Self-avoiding walks crossing a square. *Journal of Physics A: Mathematical and General* **38**(42), 9159 (2005)
5. Djidjev, H.N., Venkatesan, S.M.: Reduced constants for simple cycle graph separation. *Acta informatica* **34**(3), 231–243 (1997)
6. Dorn, F., Penninkx, E., Bodlaender, H.L., Fomin, F.V.: Efficient exact algorithms on planar graphs: Exploiting sphere cut decompositions. *Algorithmica* **58**(3), 790–810 (2010)
7. Ellenberg, J.: Geometry, inference, complexity, and democracy. *Bulletin of the American Mathematical Society* **58**(1), 57–77 (2021)
8. Har-Peled, S.: A simple proof of the existence of a planar separator. *arXiv preprint arXiv:1105.0103* (2011)
9. Jerrum, M.R., Valiant, L.G., Vazirani, V.V.: Random generation of combinatorial structures from a uniform distribution. *Theoretical computer science* **43**, 169–188 (1986)
10. Johnson, D.B.: Finding all the elementary circuits of a directed graph. *SIAM Journal on Computing* **4**(1), 77–84 (1975)
11. Lipton, R.J., Tarjan, R.E.: A separator theorem for planar graphs. *SIAM Journal on Applied Mathematics* **36**(2), 177–189 (1979)
12. Miller, G.L.: Finding small simple cycle separators for 2-connected planar graphs. *Journal of Computer and system Sciences* **32**(3), 265–279 (1986)
13. Motzkin, T.: Relations between hypersurface cross ratios, and a combinatorial formula for partitions of a polygon, for permanent preponderance, and for non-associative products. *Bulletin of the American Mathematical Society* **54**(4), 352–360 (1948)
14. Najt, E., Deford, D., Solomon, J.: Complexity and geometry of sampling connected graph partitions. *arXiv preprint arXiv:1908.08881* (2019)
15. Spielman, D.A., Teng, S.H.: Disk packings and planar separators. In: *Proceedings of the twelfth annual symposium on Computational geometry*. pp. 349–358 (1996)
16. Tarjan, R.: Enumeration of the elementary circuits of a directed graph. *SIAM Journal on Computing* **2**(3), 211–216 (1973)
17. Tiernan, J.C.: An efficient search algorithm to find the elementary circuits of a graph. *Communications of the ACM* **13**(12), 722–726 (1970)

A Appendix A

Remark 2. By the abstract definition of the dual graph, $\tilde{F}, \tilde{E}, \tilde{V}$ are just F, E, V respectively (or in 1-1 correspondence), but we use the notation $\tilde{F}, \tilde{E}, \tilde{V}$ to indicate intuitively to which graph an object belongs. E.g., by $v \in \tilde{F}$ we wish to indicate that it is a vertex v in H (but it can also be considered as a triangular face in G), and by $\{u, v\} \in \tilde{E}$ we think of it as an edge between two adjacent vertices u and v in H (and also two adjacent triangular faces in G).

Remark 3. In effect, we split v_∞ into $|E_A|$ many dangling vertices, and think of them as lying in D' . When there is no confusion we will identify $\{u, v\} \in \tilde{E}_A$ with $\{u, v^*\}$ in (the modified) H_D , except if two adjacent edges of E_A in G are on the same triangular face in D' , in which case we will treat the dangling vertices as distinct and different, even though the two vertices are identified in the original graph.

Lemma 1. *Let $\tilde{e} = \{u_1, u_2\} \in P \cap \tilde{E}_A$. Without loss of generality $u_1 \notin \tilde{F}_D$, and $u_2 \in \tilde{F}_D$. Then there exist unique vertices $u_3, \dots, u_{k-1} \in \tilde{F}_D$, and $u_k \notin \tilde{F}_D$ ($k \geq 3$), such that $\{u_i, u_{i+1}\} \in P$ for $1 \leq i < k$.*

Proof. By definition, $P = W \cap \tilde{E}_D$ for some $W \in \mathcal{W}_G$. Clearly, $\tilde{e} \in W$. Starting from $u_2 \in \tilde{F}_D$ in the direction away from u_1 we can follow the cycle W , identifying the vertices $u_3, \dots, u_{k-1} \in \tilde{F}_D$, until we reach a vertex $u_k \notin \tilde{F}_D$. Such a vertex u_k exists, since W is a cycle and $u_1 \notin \tilde{F}_D$ lies on it.

Note that this choice of u_3, \dots, u_k does not depend on our particular choice of W , but only depends on P and the initial \tilde{e} . To see this, suppose $P = W_1 \cap \tilde{E}_D = W_2 \cap \tilde{E}_D$. For either $W = W_1$ or W_2 , every vertex on the intersection $P = W \cap \tilde{E}_D$ within D has degree 2, and $\{u_i, u_{i+1}\} \in P$ for all $1 \leq i < k$. So the same sequence u_3, \dots, u_k will be selected.

Remark 4. This is true even when $k = 3$ and $u_3 = u_1$ in which case \tilde{e}' and \tilde{e} are a pair of parallel edges between u_1 and u_2 .

Lemma 10. *Let ℓ_P be the labelling of E_A defined by P on D for some $D \in \{D_1, D_2\}$. For any $e \in E_A$, if $\ell_P(e) = e'$ for some $e' \in E_A$, then $\ell_P(e') = e$. If $P = W \cap \tilde{E}_D$, then the edges where ℓ_P is nonzero coincides with where W crosses A ; in particular, ℓ_P is identically 0 iff $P \cap \tilde{E}_A = \emptyset$.*

Proof. If following a walk from \tilde{e} leads to \tilde{e}' , then following the walk back from \tilde{e}' leads to \tilde{e} . Therefore, $\ell_P(e) = e'$ implies that $\ell_P(e') = e$. The last statement is obvious.

Lemma 2.

$$|\mathcal{W}_G| = \sum_{\ell_1 \in \mathcal{L}_A} \sum_{\ell_2 \in \mathcal{L}_A} |\mathcal{C}(\ell_1, \ell_2)|.$$

Proof. We show that \mathcal{W}_G can be expressed as a disjoint union

$$\mathcal{W}_G = \bigcup_{\ell_1 \in \mathcal{L}_A} \bigcup_{\ell_2 \in \mathcal{L}_A} \mathcal{C}(\ell_1, \ell_2).$$

By definition, $\mathcal{C}(\ell_1, \ell_2) \subseteq \mathcal{W}_G$, for every $\ell_1, \ell_2 \in \mathcal{L}_A$. Now, let $W \in \mathcal{W}_G$. Let $P_1 = W \cap \tilde{E}_{D_1}$ and $P_2 = W \cap \tilde{E}_{D_2}$. Then $\ell_{P_1}, \ell_{P_2} \in \mathcal{L}_A$, and $W \in \mathcal{C}(\ell_{P_1}, \ell_{P_2})$. This is a disjoint union, for if $W \in \mathcal{C}(\ell_1, \ell_2)$ and also $W \in \mathcal{C}(\ell'_1, \ell'_2)$, then $\ell_1 = \ell_{W \cap \tilde{E}_{D_1}} = \ell'_1$ and $\ell_2 = \ell_{W \cap \tilde{E}_{D_2}} = \ell'_2$.

Remark 5. Intuitively a PSAW is the intersection of a cycle on H with D (i.e. $W \cap \tilde{E}_D$ for $W \in \mathcal{W}_G$). However, in order to work effectively with this notion, we have to define it in a way that only refers to H_D , and does not refer to any unknown cycle in the graph H . See Fig. 6 for an example of a PSAW where this occurs. Even if two edges e, e' from W are incident on the same face of D_2 , when we look at the same two dangling edges in $W \cap \tilde{E}_{D_1}$, we see that they are now no longer incident on the same vertex.




Fig. 6: Here, we have a segment of a cycle $W \in \mathcal{W}_G$, and the two induced PSAWs $W \cap \tilde{E}_{D_2}$ and $W \cap \tilde{E}_{D_1}$. Notice the dangling edges in the PSAWs that are incident on the other disc.

Lemma 3. *For any planar triangulated graph G and any $D \in \{D_1, D_2\}$ determined by a cycle A on G ,*

$$\{W \cap \tilde{E}_D : W \in \mathcal{W}_G\} \subseteq \mathcal{P}_D.$$

Proof. Let $D' \in \{D_1, D_2\}$ be the disc other than D . Let $P = W \cap \tilde{E}_D$ for some $W \in \mathcal{W}_G$. If there exists an edge $\tilde{e} = \{u, v\} \in W$ such that $u \in \tilde{F}_D, v \in \tilde{F}_{D'}$, we may equivalently consider this edge is $\{u, \pi_D(e)\}$, which is incident on a dangling vertex instead of the vertex v . Now, let $Q = W \cap \tilde{E}_{D'}$, and consider ℓ_Q . It follows from the construction of this function that $P \cup \mu_D(\ell_Q)$ is a cycle. Therefore, $P \in \mathcal{P}_D$.

Remark 6. Even though the labelling ℓ_2 does not uniquely determine the labelling ℓ_1 , the labelling ℓ_2 together with a specific (cyclic) order to traverse those edges from \tilde{E}_A indicated by ℓ_2 does uniquely determine the labelling ℓ_1 .

B Appendix B

Lemma 5. *If $\mathcal{C}_S(\ell_1, \ell_2) \neq \emptyset$, then ℓ_1 and ℓ_2 are compatible as defined by `CompatibilityCheck`.*

Proof. Let $W \in \mathcal{C}_S(\ell_1, \ell_2)$. Then $\ell_{W \cap \tilde{E}_{D_1}} = \ell_1$ and $\ell_{W \cap \tilde{E}_{D_2}} = \ell_2$, and since W is a cycle on H , we have the same set where they are nonzero on \tilde{E}_A which is $W \cap \tilde{E}_A$.

Now suppose $\{e \in E_A : \ell_1(e) \neq 0\} = \{e \in E_A : \ell_2(e) \neq 0\} \neq \emptyset$. The set M defined in `CompatibilityCheck` remains a subset of E_A and strictly grows during the **while** loop. As E_A is finite, the **while** loop must terminate. Starting with some arbitrary $e_1 \in E_A$, let $M(e_1)$ be the set M that is obtained at the end of the **while** loop if we start with the edge e_1 . Let $e_2 = \ell_1(e_1)$. This means that any $W \in \mathcal{C}_S(\ell_1, \ell_2)$ when enters \tilde{F}_{D_1} through the edge \tilde{e}_1 must exit through the edge \tilde{e}_2 . As $\ell_1(e_2) \neq 0$, we also have $\ell_2(e_2) \neq 0$. Now, if we let $e_3 = \ell_2(e_2)$, we see that W has to enter \tilde{F}_{D_2} through \tilde{e}_2 , and exit through \tilde{e}_3 . We can repeat this process until we reach an edge e_{k+1} such that $\ell_1(e_{k+1}) \in M$. Since W is a cycle, this must be e_1 , and the cycle has been traversed. Therefore, we have found that $W \cap \tilde{E}_A = \{\tilde{e}_1, \dots, \tilde{e}_k\}$. It follows that $M(e_1)$ has collected all the edges where W crosses A , $M(e_1) = \{e_1, e_2, \dots, e_k\}$, thus $M(e_1) = \{e \in E_A : \ell_1(e) \neq 0\}$, and `CompatibilityCheck` returns **true**.

Remark 7. It should be noted that T can also be viewed as an ordered set of arcs in H_{D_1} and H_{D_2} , since $\varepsilon(T) \subseteq \tilde{E}_A$. If T is viewed as an ordered set of arcs in H_{D_1} , then we can replace all the vertices in \tilde{F}_{D_2} in the arcs with the corresponding dangling vertices in the graph, and the same applies for H_{D_2} . We will not distinguish between these cases any further, and use the same symbol T in all three settings, as the meaning will be clear from the context.

Lemma 11.

- If $\varepsilon(S) \cap \tilde{E}_{D_1} \neq \emptyset$ and $\varepsilon(S) \cap \tilde{E}_{D_2} \neq \emptyset$, then $|\mathcal{C}_S(0,0)| = 0$.
- If $\emptyset \neq \varepsilon(S) \subseteq \tilde{E}_D$ for $D \in \{D_1, D_2\}$, then $|\mathcal{C}_S(0,0)| = |\mathcal{C}_{D,S}(0)|$.
- If $\varepsilon(S) = \emptyset$, then $|\mathcal{C}_S(0,0)| = |\mathcal{C}_{D_1,S}(0)| + |\mathcal{C}_{D_2,S}(0)| - 1$.

Proof. If $W \in \mathcal{C}_S(0,0)$, then $W \cap \tilde{E}_A = \emptyset$, as otherwise there would be some edge $\tilde{e} \in W \cap \tilde{E}_A$, which would imply that $\ell_{W \cap \tilde{E}_{D_1}}(e) \neq 0$. Thus, W cannot have edges in both \tilde{E}_{D_1} and \tilde{E}_{D_2} . In particular, if $\varepsilon(S) \cap \tilde{E}_{D_1} \neq \emptyset$, and $\varepsilon(S) \cap \tilde{E}_{D_2} \neq \emptyset$, then $\mathcal{C}_S(0,0) = \emptyset$.

Now consider the case where $\emptyset \neq \varepsilon(S) \subseteq \tilde{E}_{D_1}$. For any $W \in \mathcal{W}_{G,S}$ since $\varepsilon(S) \subseteq W$, it follows that $W \cap \tilde{E}_{D_1} \neq \emptyset$. Suppose $W \in \mathcal{C}_S(0,0)$. Then, as we have already seen, $W \cap \tilde{E}_A = \emptyset$. Since W is a cycle with $W \cap \tilde{E}_{D_1} \neq \emptyset$ and $W \cap \tilde{E}_A = \emptyset$, we have $W \subseteq \tilde{E}_{D_1}$, and as a consequence $W = W \cap \tilde{E}_{D_1}$. Applying [Definition 2](#), we see that $W \in \mathcal{P}_{D_1}$ with the empty $\mu_{D_1}(0)$ extension. And since W conforms to S , we have $W \in \mathcal{P}_{D_1,S}$. Finally as $\ell_W = \ell_{W \cap \tilde{E}_{D_1}} = 0$, we get $W \in \mathcal{C}_{D_1,S}(0)$.

Conversely, let $P \in \mathcal{C}_{D_1,S}(0)$. Then by definition $P \in \mathcal{P}_{D_1,S}$, $P \subseteq \tilde{E}_{D_1}$ and $\ell_P = \ell_{P \cap \tilde{E}_{D_1}} = 0$, hence $P \cap \tilde{E}_A = \emptyset$. By the definition of $\mathcal{P}_{D_1,S}$, there is some $\ell \in \mathcal{L}_A$ such that $P \cup \mu_{D_1}(\ell)$ is a cycle. We claim such ℓ must be 0. Indeed, for any e if $\ell(e) \neq 0$ then this e must belong to $P \cap \tilde{E}_A$, a contradiction to $P \cap \tilde{E}_A = \emptyset$. Hence this $\ell = 0$, and $\mu_{D_1}(\ell) = \emptyset$. Therefore P itself is a cycle. We have $P \in \mathcal{W}_{G,S}$, and since $P \subseteq \tilde{E}_{D_1}$, we have $\ell_{P \cap \tilde{E}_{D_2}} = 0$ in addition to $\ell_{P \cap \tilde{E}_{D_1}} = 0$. It follows that $P \in \mathcal{C}_S(0,0)$. We have proved that, when $\emptyset \neq \varepsilon(S) \subseteq \tilde{E}_{D_1}$,

$$\mathcal{C}_S(0,0) = \mathcal{C}_{D_1,S}(0).$$

Finally, consider the case $\varepsilon(S) = \emptyset$. The empty cycle consisting of no vertex and no edge is in $\mathcal{C}_\emptyset(0,0)$ as well as in both $\mathcal{C}_{D_1,\emptyset}(0)$ and $\mathcal{C}_{D_2,\emptyset}(0)$.

Now consider nonempty $W \in \mathcal{C}_\emptyset(0,0)$. We know that $W \cap \tilde{E}_A = \emptyset$. Hence, $W \subseteq \tilde{E}_{D_1}$ or $W \subseteq \tilde{E}_{D_2}$, but not both. Say $W \subseteq \tilde{E}_{D_1}$. Then $W \in \mathcal{P}_{D_1} = \mathcal{P}_{D_1,\emptyset}$ with the empty extension $\mu_{D_1}(0) = \emptyset$. Hence, $W \in \mathcal{C}_{D_1,\emptyset}(0)$. Conversely, if $W \in \mathcal{C}_{D_1,\emptyset}(0)$, then $W \subseteq \tilde{E}_{D_1}$, and $\ell_W = \ell_{W \cap \tilde{E}_{D_1}} = 0$. Hence $W \cap \tilde{E}_A = \emptyset$. By the definition of \mathcal{P}_{D_1} , any labelling $\ell \in \mathcal{L}_A$ such that $W \cup \mu_{D_1}(\ell)$ is a cycle implies that $\ell = 0$, and so W is a cycle. It follows that $W \in \mathcal{C}_\emptyset(0,0)$.

We have shown that a nonempty W belongs to $\mathcal{C}_\emptyset(0,0)$ iff it belongs to exactly one of $\mathcal{C}_{D_1,\emptyset}(0)$ or $\mathcal{C}_{D_2,\emptyset}(0)$. Taking into account the empty cycle, we have $|\mathcal{C}_\emptyset(0,0)| = |\mathcal{C}_{D_1,\emptyset}(0)| + |\mathcal{C}_{D_2,\emptyset}(0)| - 1$.

Lemma 6. *Given a compatible $(\ell_1, \ell_2) \neq (0,0)$,*

$$|\mathcal{C}_\emptyset(\ell_1, \ell_2)| = |\mathcal{C}_{D_1,T}(\ell_1)| \times |\mathcal{C}_{D_2,T}(\ell_2)|.$$

Proof. [Lemma 12](#) implies that if $W \in \mathcal{C}_\emptyset(\ell_1, \ell_2)$, then $W \cap \tilde{E}_{D_1} \in \mathcal{C}_{D_1,T}(\ell_1)$, and $W \cap \tilde{E}_{D_2} \in \mathcal{C}_{D_2,T}(\ell_2)$.

[Lemma 13](#) implies that the converse is also true. Therefore, for compatible $(\ell_1, \ell_2) \neq (0,0)$, the function $\phi : \mathcal{C}_{D_1,T}(\ell_1) \times \mathcal{C}_{D_2,T}(\ell_2) \rightarrow \mathcal{C}_\emptyset(\ell_1, \ell_2)$ defined as

$$\phi(P_1, P_2) = P_1 \cup P_2$$

is a well-defined surjective function. It is easy to see that the function is also injective, since $P_1 \cup P_2 = P'_1 \cup P'_2$ implies that

$$P_1 = (P_1 \cup P_2) \cap \tilde{E}_{D_1} = (P'_1 \cup P'_2) \cap \tilde{E}_{D_1} = P'_1,$$

and similarly, $P_2 = P'_2$. Therefore, ϕ is a bijection. It follows that, for compatible $(\ell_1, \ell_2) \neq (0,0)$,

$$|\mathcal{C}_\emptyset(\ell_1, \ell_2)| = |\mathcal{C}_{D_1,T}(\ell_1)| \times |\mathcal{C}_{D_2,T}(\ell_2)|.$$

Lemma 12. *If $W \in \mathcal{C}_\emptyset(\ell_1, \ell_2)$ for $(\ell_1, \ell_2) \neq (0,0)$, then $W \cap \tilde{E}_{D_1} \in \mathcal{C}_{D_1,T}(\ell_1)$, and $W \cap \tilde{E}_{D_2} \in \mathcal{C}_{D_2,T}(\ell_2)$.*

Proof. Let $P_1 = W \cap \tilde{E}_{D_1}$. We know from [Lemma 3](#) that $P_1 \in \mathcal{P}_{D_1}$. It also follows that $P_1 \in \mathcal{C}_{D_1}(\ell_1)$, since $\ell_{P_1} = \ell_1$, by our choice of W . Similarly, $P_2 \in \mathcal{C}_{D_2}(\ell_2)$, where $P_2 = W \cap \tilde{E}_{D_2}$. Moreover, it follows from the

construction of P_1 , that $P_1 \cup \mu_{D_1}(\ell_2)$ is a cycle. Now, we show that $P_1 \cup \mu_{D_1}(\ell_2)$ (and similarly, $P_2 \cup \mu_{D_2}(\ell_1)$) conforms to T . Assume that $T = ((u_1 \rightarrow v_1), \dots, (u_k \rightarrow v_k))$.

For convenience, we will use $e_i \in E_A$ to refer to the edge for which $\tilde{e}_i = \{u_i, v_i\}$. From the construction of T , we know that $\ell_1(e_1) = e_2$. Therefore, from the definition of the function $\ell_{P_1} = \ell_1$, we can see that P_1 must enter \tilde{F}_{D_1} through the edge (u_1, v_1) , and exit through the edge (u_2, v_2) , and visit the vertices u_1, v_1, u_2, v_2 , in that order.

Once again, from the construction of T , we know that $\ell_2(e_2) = e_3$. Therefore, we know that $\{v_2, u_3\} \in \mu_{D_1}(\ell_2)$. So, it picks up from P_2 at the dangling vertex v_2 , and joins it again at u_3 . Combining with our earlier observation, we see that $P_1 \cup \mu_{D_1}(\ell_2)$ visits the vertices $u_1, v_1, u_2, v_2, u_3, v_3$ in that order. This pattern then repeats for e_3, \dots, e_k . So, we know that $\varepsilon(T) \subseteq W$, and since $P_1 \cup \mu_{D_1}(\ell_2)$ is a cycle, we can see that it visits the vertices $u_1, v_1, \dots, u_k, v_k$ in that order. Therefore, $P_1 \in \mathcal{C}_{D_1, T}(\ell_1)$. By an analogous argument, we also have $P_2 \in \mathcal{C}_{D_2, T}(\ell_2)$.

Lemma 13. *If $P_1 \in \mathcal{C}_{D_1, T}(\ell_1)$, and $P_2 \in \mathcal{C}_{D_2, T}(\ell_2)$ for compatible $(\ell_1, \ell_2) \neq (0, 0)$, then $P_1 \cup P_2 \in \mathcal{C}_\emptyset(\ell_1, \ell_2)$.*

Proof. Note that since $P_1 \subseteq \tilde{E}_{D_1}$ and $P_2 \subseteq \tilde{E}_{D_2}$, it follows that $P_1 \cup P_2 \subseteq \tilde{E}$. Here a dangling edge $\tilde{e} \in P_1$ (or P_2) is identified with the corresponding edge in \tilde{E} .

First, we show that $P_1 \cup P_2$ contains a cycle in $\mathcal{W}_{G, \emptyset}$. Note that $P_1 \cap P_2 \subseteq \tilde{E}_A$. The compatibility of $\ell_1, \ell_2 \in \mathcal{L}_A$ implies that

$$P_1 \cap \tilde{E}_A = \{e \in E_A : \ell_1(e) \neq 0\} = \{e \in E_A : \ell_2(e) \neq 0\} = P_2 \cap \tilde{E}_A,$$

and hence both are equal to $P_1 \cap P_2$.

Just like before, we may assume that $T = ((u_1 \rightarrow v_1), \dots, (u_k \rightarrow v_k))$, such that $\tilde{e}_i = \{u_i, v_i\}$ for $e_i \in E_A$. Since ℓ_1 and ℓ_2 are compatible and $T = T(\ell_1, \ell_2)$, we have $\ell_1(e_1) = e_2, \ell_2(e_2) = e_3, \ell_1(e_3) = e_4, \dots, \ell_2(e_k) = e_1$. Importantly, we also have that $\{\tilde{e}_1, \dots, \tilde{e}_k\} = P_1 \cap P_2$.

So, starting with e_1 , we know that since $\ell_1(e_1) = e_2$, P_1 enters \tilde{F}_{D_1} through \tilde{e}_1 , and exits through \tilde{e}_2 . Then, since $\ell_2(e_2) = e_3$, we know that P_2 enters \tilde{F}_{D_2} through \tilde{e}_2 , and exits through \tilde{e}_3 . Repeating this process until we reach e_k , we see that $P_1 \cup P_2$ contains a cycle W that passes through $\{\tilde{e}_1, \dots, \tilde{e}_k\}$.

Now we show that $P_1 \cup P_2$ is itself the cycle W we have identified, i.e., no edge of $P_1 \cup P_2$ is left out from the cycle W . Let $e \in P_1$ be an arbitrary edge of P_1 . Since $P_1 \in \mathcal{C}_{D_1, T}(\ell_1)$, there exists some $\ell \in \mathcal{L}_A$ such that $P_1 \cup \mu_{D_1}(\ell)$ is a cycle. This ℓ and ℓ_1 must be compatible, so they share all the nonzero locations on \tilde{E}_A . We are given $\ell_1 \neq 0$, and thus e is traversed on the cycle $P_1 \cup \mu_{D_1}(\ell)$ on a simple path segment beginning and ending at some edges on \tilde{E}_A . As $P_1 \cap \tilde{E}_A = P_2 \cap \tilde{E}_A$, and this is the same as the underlying edge set $\varepsilon(T)$, when we traverse the cycle $P_1 \cup \mu_{D_1}(\ell)$ exactly the same simple path segments of P_1 as indicated by ℓ_1 are traversed, as in the cyclic traversal of W indicated by T , possibly in some different order. In particular, $e \in P_1$ is traversed in the cycle W . Similarly all edges of P_2 are also contained in W .

Therefore, $W = P_1 \cup P_2$ is a cycle, and $W \in \mathcal{W}_{G, \emptyset}$. Finally, as $W \cap \tilde{E}_{D_1} = P_1$, by construction, we have $\ell_{P_1} = \ell_1$. Similarly, $\ell_{P_2} = \ell_2$. Therefore, $P_1 \cup P_2 \in \mathcal{C}_\emptyset(\ell_1, \ell_2)$.

Another immediate corollary of [Lemma 12](#) and [Lemma 13](#) is the following:

Corollary 1. *Given compatible $(\ell_1, \ell_2) \neq (0, 0)$,*

$$\mathcal{C}_{D_1}(\ell_1) = \bigcup_{\substack{\ell_2 \in \mathcal{L}_A: \\ (\ell_1, \ell_2) \text{ compatible}}} \mathcal{C}_{D_1, T(\ell_1, \ell_2)}(\ell_1).$$

Lemma 14. *Let $t \geq 1$. If $|S| \leq s$ and $|T| \leq t$, then*

$$|\mathcal{I}_{S, T}| \leq 2t \binom{s+t}{t}.$$

Proof. Without loss of generality $|S| = s$ and $|T| = t$, as the upper bound is monotonic in both s and t . If $s = 0$ then the upper bound is obvious. Let $s \geq 1$. Since $\mathcal{I}_{S,T}$ only contains one representative of each equivalence class, we may choose it so that if $I \in \mathcal{I}_{S,T}$, then $I \cap \varepsilon(S) = S$. Picking the first element of S as the starting position, after that in the forward direction of the cyclic order of S we can set $s - 1 + t$ positions, of which we may pick any t positions to be occupied by $\varepsilon(T)$. Then there are t choices for the starting element of T and two possible cyclic directions for T relative to the cyclic order of S . This gives us an upper bound

$$2t \binom{s+t-1}{t} \leq 2t \binom{s+t}{t}.$$

Lemma 15. *Let S be an ordered set of arcs in the graph H . Given compatible labellings $\ell_1, \ell_2 \in \mathcal{L}_A$, let $T = T(\ell_1, \ell_2)$ be the ordered set of arcs in H defined above. Then,*

$$|\mathcal{C}_S(\ell_1, \ell_2)| = \sum_{I \in \mathcal{I}_{S,T}} |\mathcal{C}_I(\ell_1, \ell_2)|.$$

Proof. We show that $\mathcal{C}_S(\ell_1, \ell_2)$ is a disjoint union,

$$\mathcal{C}_S(\ell_1, \ell_2) = \bigcup_{I \in \mathcal{I}_{S,T}} \mathcal{C}_I(\ell_1, \ell_2).$$

It is easy to see that if $W \in \mathcal{C}_I(\ell_1, \ell_2)$ for some $I \in \mathcal{I}_{S,T}$, then $W \in \mathcal{C}_S(\ell_1, \ell_2)$ since S is an ordered subset of I . Similarly, if $W \in \mathcal{C}_S(\ell_1, \ell_2)$, note that W conforms to T by construction of $T = T(\ell_1, \ell_2)$. Therefore, there is some $I \in \mathcal{I}_{S,T}$ such that $W \in \mathcal{C}_I(\ell_1, \ell_2)$.

To show that it is a disjoint union, let $W \in \mathcal{C}_{I_1}(\ell_1, \ell_2) \cap \mathcal{C}_{I_2}(\ell_1, \ell_2)$ for $I_1, I_2 \in \mathcal{I}_{S,T}$. If either S or $T = \emptyset$, then it is clear that $I_1 \sim I_2$. Suppose $S, T \neq \emptyset$. By a cyclic shift and reversal we may assume in both I_1 and I_2 , the ordered set of arcs S appears as $((u_1 \rightarrow v_1), \dots, (u_k \rightarrow v_k))$, with $(u_1 \rightarrow v_1)$ appearing first.

We pick a traversal of W to start at $(u_1 \rightarrow v_1)$ in S , in the order given by I_1 above. Then we follow the appearances in W of the arcs whose edges are from $\varepsilon(S)$ and $\varepsilon(T)$. The sequence of these appearances shows that $I_1 \sim I_2$. This shows that the union is disjoint, one term per each equivalence class in $\mathcal{I}_{S,T}$.

Lemma 7. *Given compatible labellings $\ell_1, \ell_2 \in \mathcal{L}_A$ such that $(\ell_1, \ell_2) \neq (0, 0)$,*

$$|\mathcal{C}_S(\ell_1, \ell_2)| = \sum_{\text{valid } I \in \mathcal{I}_{S,T}} |\mathcal{C}_I(\ell_1, \ell_2)|,$$

where $T = T(\ell_1, \ell_2)$.

Proof. Note that condition (1) is not dependent on the specific choice of I . By the construction of $T = T(\ell_1, \ell_2)$, we know that if $W \in \mathcal{C}_I(\ell_1, \ell_2) \subseteq \mathcal{C}_\emptyset(\ell_1, \ell_2)$, then $W \cap \tilde{E}_A = \varepsilon(T)$. So, it is not possible to find an edge \tilde{e}' in W such that $\tilde{e}' \in (\varepsilon(S) \cap \tilde{E}_A) \setminus \varepsilon(T)$. Moreover, if $W \in \mathcal{C}_I(\ell_1, \ell_2)$, it has to conform to both S and T . So, if S and T require that the vertices be visited in different orders, no walk can conform to both S and T . Therefore, if condition (1) is not satisfied, $\mathcal{C}_I(\ell_1, \ell_2) = \emptyset$.

Now, let us consider condition (2). Since \tilde{E}_A is a cut between \tilde{F}_{D_1} and \tilde{F}_{D_2} , if a walk W goes from a vertex in \tilde{F}_D to another vertex not in \tilde{F}_D , it has to go through some edge in \tilde{E}_A . Since $\varepsilon(T) = W \cap \tilde{E}_A$ for all $W \in \mathcal{C}_I(\ell_1, \ell_2)$, it can be seen immediately that if condition (2) is violated, then $\mathcal{C}_I(\ell_1, \ell_2) = \emptyset$.

Therefore, from Lemma 15 we have that

$$|\mathcal{C}_S(\ell_1, \ell_2)| = \sum_{\text{valid } I \in \mathcal{I}_{S,T}} |\mathcal{C}_I(\ell_1, \ell_2)|,$$

where $T = T(\ell_1, \ell_2)$

Lemma 8. Given compatible $\ell_1, \ell_2 \in \mathcal{L}_A$ such that $(\ell_1, \ell_2) \neq (0, 0)$, and $T = T(\ell_1, \ell_2)$,

$$|\mathcal{C}_S(\ell_1, \ell_2)| = \sum_{\text{valid } I \in \mathcal{I}_{S,T}} \left| \mathcal{C}_{D_1, I \cap \tilde{E}_{D_1}}(\ell_1) \right| \times \left| \mathcal{C}_{D_2, I \cap \tilde{E}_{D_2}}(\ell_2) \right|.$$

Proof. For a compatible $(\ell_1, \ell_2) \neq (0, 0)$, and a valid $I \in \mathcal{I}_{S,T}$ (where $T = T(\ell_1, \ell_2)$), consider the following function $\phi_I : \mathcal{C}_{D_1, I \cap \tilde{E}_{D_1}}(\ell_1) \times \mathcal{C}_{D_2, I \cap \tilde{E}_{D_2}}(\ell_2) \rightarrow \mathcal{C}_I(\ell_1, \ell_2)$ defined as:

$$\phi_I(P_1, P_2) = P_1 \cup P_2$$

Now, from Lemma 16, we see that

$$|\mathcal{C}_S(\ell_1, \ell_2)| = \sum_{\text{valid } I \in \mathcal{I}_{S,T}} \left| \mathcal{C}_{D_1, I \cap \tilde{E}_{D_1}}(\ell_1) \right| \times \left| \mathcal{C}_{D_2, I \cap \tilde{E}_{D_2}}(\ell_2) \right|.$$

Lemma 16. The function ϕ_I is bijective.

Proof. First, we need to verify that the function is well-defined. Note that Lemma 13 implies that since $\mathcal{C}_{D_1, I \cap \tilde{E}_{D_1}}(\ell_1) \subseteq \mathcal{C}_{D_1, T}(\ell_1)$ and $\mathcal{C}_{D_2, I \cap \tilde{E}_{D_2}}(\ell_2) \subseteq \mathcal{C}_{D_2, T}(\ell_2)$, it follows that $P_1 \cup P_2 \in \mathcal{C}_\emptyset(\ell_1, \ell_2)$.

Now, consider the cycle $W = P_1 \cup P_2$ in H . Since I is a valid interleaving, we may assume that $I = ((u^{T,1} \rightarrow v^{T,1}), (u_1^{S,1} \rightarrow v_1^{S,1}), \dots, (u_{t_1}^{S,1} \rightarrow v_{t_1}^{S,1}), (u^{T,2} \rightarrow v^{T,2}), (u_1^{S,2} \rightarrow v_1^{S,2}), \dots, (u_{t_2}^{S,2} \rightarrow v_{t_2}^{S,2}), \dots, (u^{T,k} \rightarrow v^{T,k}), (u_1^{S,k} \rightarrow v_1^{S,k}), \dots, (u_{t_k}^{S,k} \rightarrow v_{t_k}^{S,k}))$, where

$$\{\{u^{T,1}, v^{T,1}\}, \dots, \{u^{T,k}, v^{T,k}\}\} = \varepsilon(I) \cap \tilde{E}_A$$

and k is even, being the number of times a cycle crosses A . We may further assume without loss of generality that $u^{T,1}, v^{T,2}, u^{T,3}, v^{T,4}, \dots, v^{T,k} \in \tilde{F}_{D_2}$, $v^{T,1}, u^{T,2}, \dots, u^{T,k} \in \tilde{F}_{D_1}$, $u_i^{S,1}, v_i^{S,1}, u_i^{S,3}, v_i^{S,3}, \dots, u_i^{S,k-1}, v_i^{S,k-1} \in \tilde{F}_{D_1}$, and that $u_i^{S,2}, v_i^{S,2}, u_i^{S,4}, v_i^{S,4}, \dots, u_i^{S,k}, v_i^{S,k} \in \tilde{F}_{D_2}$ for all i .

In this case, as $W = P_1 \cup P_2$, since $P_1 \in \mathcal{C}_{D_1, I \cap \tilde{E}_{D_1}}(\ell_1)$ and $P_2 \in \mathcal{C}_{D_2, I \cap \tilde{E}_{D_2}}(\ell_2)$, it follows that $W \supseteq \varepsilon(I)$. Moreover, W visits the vertices $u^{T,1}, v^{T,1}, u_1^{S,1}, v_1^{S,1}, \dots, u_{t_1}^{S,1}, v_{t_1}^{S,1}, u^{T,2}, v^{T,2}$ in \tilde{E}_{D_1} in that order. From $u^{T,2}, v^{T,2}$, it then visits the vertices $u_1^{S,2}, \dots, v_{t_2}^{S,2}, u^{T,3}, v^{T,3}$ in \tilde{E}_{D_2} in that order, and so on. Therefore, when we put these segments of W together, we see that W conforms to I . Therefore, $W \in \mathcal{C}_I(\ell_1, \ell_2)$, and ϕ_I is well-defined.

Now, we show that this function is injective. If there exist (P_1, P_2) and (P'_1, P'_2) such that $P_1 \cup P_2 = P'_1 \cup P'_2$, we know that

$$P_1 = (P_1 \cup P_2) \cap \tilde{E}_{D_1} = (P'_1 \cup P'_2) \cap \tilde{E}_{D_1} = P'_1$$

Therefore, $P_1 = P'_1$. Similarly, $P_2 = P'_2$. Therefore, the function is injective.

Finally, we show that ϕ_I is surjective. Given $W \in \mathcal{C}_I(\ell_1, \ell_2) \subseteq \mathcal{C}_\emptyset(\ell_1, \ell_2)$, Lemma 12 tells us that $W \cap \tilde{E}_{D_1} \in \mathcal{C}_{D_1, T}(\ell_1)$, and $W \cap \tilde{E}_{D_2} \in \mathcal{C}_{D_2, T}(\ell_2)$. Now, let $P_1 = W \cap \tilde{E}_{D_1}$, and $P_2 = W \cap \tilde{E}_{D_2}$. Since W conforms to I , and therefore $I \cap \tilde{E}_{D_1}$, it is clear that $P_1 \cup \mu_{D_1}(\ell_2)$ conforms to $I \cap \tilde{E}_{D_1}$ as well. Therefore, $P_1 \in \mathcal{C}_{I \cap \tilde{E}_{D_1}}(\ell_1)$, and similarly, $P_2 \in \mathcal{C}_{I \cap \tilde{E}_{D_2}}(\ell_2)$. Since, $P_1 \cup P_2 = W$ by construction, it follows that ϕ_I is surjective.

C Appendix C

We will go through the construction of the graph G_1 , and the ordered set of arcs S_1 in more detail here. The construction of G_1 occurs inductively. Starting from $D^0 = G_{D_1}$, we iteratively construct $D^1, \dots, D^n = G_1$. Assume that $D^i = (V^i, E^i, F^i)$. We know that D^0 contains the cycle $A^0 = A$ which bounds the face f^* . During each iteration, we construct D^{i+1} by modifying D^i with boundary A^i , so that D^{i+1} is a graph on a disc with boundary A^{i+1} . We will abuse the notation a bit, and refer to the face bounded by the edges E_A^i of A^i in each D^i still as f^* .

First, we will formally describe the contractions we will use to transform D^i into D^{i+1} .

1. **Single-edge contraction:** We are given some edge $e = \{a, b\} \in E_A^i$. As e is an edge on the boundary A^i of D^i , there is a unique triangular face within D_1 containing this edge e . Let $e' = \{a, c\}, e'' = \{b, c\}$ both in E^i , such that $\{e, e', e''\} \in F^i$ is that face. We assume $\deg(c) > 2$, i.e., there is no parallel edge $e^* = \{a, b\} \in E^i$ such that $\{e^*, e', e''\} \in F^i$ (otherwise we can not perform single-edge contraction on e). To perform single-edge contraction on e , we identify the vertices a and b with each other, and the edges e' and e'' with each other. We also delete the edge e and the face $\{e, e', e''\}$. The degree of c is decreased by 1. In this case, $E_A^{i+1} = E_A^i \setminus \{e\}$; however, if $E_A^i = \{e, e', e''\}$, then $E_A^{i+1} = \emptyset$. See Fig. 5a.
2. **Edge-pair contraction:** We have a pair of consecutive edges $e, e' \in E_A^i$ on A^i , $e = \{a, b\}$ and $e' = \{b, c\}$. Moreover, there is no $e'' = \{a, c\} \in E^i$ such that $\{e, e', e''\} \in F^i$, in other words, $\deg(b) > 2$. To perform edge-pair contraction on $\{e, e'\}$, we identify the vertices a and c with each other, and the edges e and e' with each other. The degree of b is decreased by 1. In this case, V_A^{i+1} is $V_A^i \setminus \{b\}$ with a and c identified, and $E_A^{i+1} = E_A^i \setminus \{e, e'\}$. See Fig. 5b.
3. **Parallel-edge contraction:** We are given two parallel edges $e = \{a, b\} \in E_A^i, e' = \{a, b\} \in E^i$, and two adjacent faces $\{e, e'', e'''\}, \{e', e'', e'''\} \in F^i$, where $e'' = \{a, c\}, e''' = \{c, b\} \in E^i$. To perform parallel-edge contraction on $\{e, e'\}$, we delete the vertex c , and the edges e'', e''' , and identify the edges e and e' with each other. The two faces $\{e, e'', e'''\}, \{e', e'', e'''\} \in F^i$ are also removed. In this case, $E_A^{i+1} = E_A^i$ (with the identified e and e' taking the place of e), but $|V^{i+1}| < |V^i|$. However, if $E_A^i = \{e, e'\}$, then $E_A^{i+1} = \emptyset$. See Fig. 5c.

Now, let us see the precise order in which these contractions are applied. For the graph D^0 , we define the function $\ell_2^0 = \ell_2$ on $E_A^0 = E_A$. Inductively assume that $\ell_2^i \in \mathcal{L}_{A^i}$, we will define ℓ_2^{i+1} as we construct D^{i+1} from D^i .

To construct D^{i+1} , we first find an edge $e = \{a, b\} \in E_A^i$ such that $\ell_2^i(e) = 0$, if such an edge exists. We check if this edge e is eligible for a parallel-edge contraction (i.e., we check if there is a parallel edge $e' = \{a, b\} \in E^i$, and edges $e'' = \{a, c\}, e''' = \{c, b\} \in E^i$ such that $\{e, e'', e'''\}, \{e', e'', e'''\} \in F^i$.) See Fig. 5c. If this is the case, we apply a parallel-edge contraction to obtain D^{i+1} , and let $\ell_2^{i+1} = \ell_2^i$. Clearly, $\ell_2^{i+1} \in \mathcal{L}_{A^{i+1}}$ in this case. If a parallel-edge contraction is not possible for the edge e , then let the unique triangular face in F^i containing e be $\{e, e', e''\}$, and $e' = \{a, c\}, e'' = \{b, c\}$, we must have $\deg(c) > 2$, and single-edge contraction on e is applicable. See Fig. 5a. We apply a single-edge contraction on the edge e , and let $\ell_2^{i+1} = \ell_2^i|_{E_A^{i+1}}$. Since $\ell_2^i(e) = 0$, and $E_A^{i+1} = E_A^i \setminus \{e\}$, it is clear that $\ell_2^{i+1} \in \mathcal{L}_{A^{i+1}}$.

Now we assume that there are no edges $e \in E_A^i$ such that $\ell_2^i(e) = 0$, but $E_A^i \neq \emptyset$. Since $\ell_2^i \in \mathcal{L}_{A^i}$, the pairing of edges by ℓ_2^i corresponds to a Motzkin path which denotes a set of non-intersecting chords. We claim that there are two adjacent edges $e, e' \in E_A^i$ such that $\ell_2^i(e) = e'$ and $\ell_2^i(e') = e$. To see that, start with any paired edges $e_1, e_2 \in E_A^i$ with $\ell_2^i(e_1) = e_2$ and $\ell_2^i(e_2) = e_1$. If e_1, e_2 are not adjacent edges in E_A^i , then they separate A^i into two segments, and the pairings according to ℓ_2^i are confined within each segment, or else there would be intersecting chords. Restricted within each segment, ℓ_2^i still denotes a Motzkin path which assigns no 0 values, and by induction we can get two adjacent paired edges $e, e' \in E_A^i$ as claimed.

Assume $e = \{a, b\}, e' = \{b, c\}$. If there exists some edge $e'' = \{a, c\} \in E^i$, such that $\{e, e', e''\} \in F^i$, i.e., $\deg(b) = 2$, then we do not construct D^{i+1} at all. We can directly claim that $|\mathcal{C}_{D_1, I \cap \tilde{E}_{D_1}}(\ell_1)| = 1$, if $\varepsilon(I \cap \tilde{E}_{D_1}) = \{\tilde{e}, \tilde{e}'\}$; we set $|\mathcal{C}_{D_1, I \cap \tilde{E}_{D_1}}(\ell_1)| = 0$, otherwise (see Lemma 18). On the other hand, if there is no such face in F^i , then edge-pair contraction on e, e' is applicable; see Fig. 5b. We perform an edge-pair contraction on e, e' to construct D^{i+1} , and let $\ell_2^{i+1} = \ell_2^i|_{E_A^{i+1}}$. By our choice that e, e' are paired adjacent edges, it follows that $\ell_2^{i+1} \in \mathcal{L}_{A^{i+1}}$.

This describes the construction of D^{i+1} when $E_A^i \neq \emptyset$, with the one exception of that special case as specified, which immediately concludes the computation of $|\mathcal{C}_{D_1, I \cap \tilde{E}_{D_1}}(\ell_1)|$. Since during each contraction step the graph D^i does not increase but either $|E_A^i|$ or $|V^i|$ strictly decreases, this process will eventually terminate. Assume this process continues until $i = \eta$, at which point $E_A^\eta = \emptyset$, and we let $G_1 = D^\eta$. We note that our description of constructing D^η from D^0 depends on a particular order of applicable contraction steps and does not assert that the process produces a canonical or unique graph G_1 . However, we will not need this uniqueness.

Lemma 17. *The graph $G_1 = (V_1, E_1, F_1)$ is a planar triangulated graph. Moreover, if $|V_1| \geq 3$, then G_1 has no vertices of degree less than 2.*

Proof. Clearly starting from a plane graph the contraction steps maintain planarity. Therefore G_1 is planar.

Now we show that G_1 is triangulated and that when $|V_1| \geq 3$, G_1 has no vertex of degree 1. We will inductively show that for each i , the faces $F^i \setminus \{f^*\}$ are triangles. Since f^* does not exist in $D^\eta = G_1$, this would imply that G_1 is triangulated. Similarly, we will inductively show that the degrees of all vertices stay greater than 1, as long as $|V^i| \geq 3$. Since $|V^i|$ decreases as i increases, this completes the proof.

Consider the case where a parallel-edge contraction (Fig. 5c) is applied to D^i . Consider the edge e' . If $e' \in E_A^i$, then $E_A^i = \{e, e'\}$, as they form a cycle, and $E^i = \{e, e', e'', e'''\}$ with $V^i = \{a, b, c\}$. Therefore, after the contraction, $E_A^{i+1} = \emptyset$, and $V^{i+1} = V^\eta = V_1 = \{a, b\}$. In this case, G_1 is vacuously triangulated.

Now, assume $e' \notin E_A^i$. Then, e' is shared by two triangular faces. One triangular face is $\{e', e'', e'''\}$. Let the other one be $\{e', e_1, e_2\}$. The effect of this parallel-edge contraction step is simply to make $\{e, e_1, e_2\}$ a new triangular face replacing three triangular faces, and nothing outside of $\{e, e_1, e_2\}$ is changed (see Fig. 5c). Since all the other faces in $F^i \setminus \{f^*\}$ were triangular, every $f \in F^{i+1} \setminus \{f^*\}$ is also triangular.

Moreover, the only vertices in V^{i+1} coming from V^i whose degrees are changed by the parallel-edge contraction are the vertices a and b . But we know that when $e' \notin E_A^i$, the edges e, e_1 and the edges e, e_2 are incident to a, b respectively, in D^{i+1} . Therefore, all vertices in V^{i+1} have degree at least 2.

Next consider the case where a single-edge contraction (Fig. 5a) is applied to D^i . Let $e = \{a, b\} \in E_A^i, e' = \{a, c\}, e'' = \{b, c\} \in E^i$ be the edges involved. If $e', e'' \in E_A^i$, as well, then, $E_A^i = \{e, e', e''\}$, as these edges form a cycle. Then, after the contraction, $E_A^{i+1} = \emptyset$, and $V^{i+1} = V^\eta = V_1 = \{a, c\}$. So, G_1 is vacuously triangulated.

Now, assume without loss of generality that $e' \notin E_A^i$. So, e' is shared by two triangular faces in D^i . One of these faces is $\{e, e', e''\}$. Let the other one be $\{e', e_1, e_2\}$. Note that we perform this single-edge contraction step only after we have performed all parallel-edge contraction steps. So in particular, $e'' \notin \{e', e_1, e_2\}$. Moreover, since $e \in E_A^i$, it belongs to only one face in F^i . Therefore, $e \notin \{e', e_1, e_2\}$. So, after the contraction, edges in $\{e', e_1, e_2\}$ are not deleted or identified with each other, and the face remains a triangle in F^{i+1} . Similarly, if $e'' \notin E_A^i$ is also true, the triangle $\{e'', e_3, e_4\}$ (other than $\{e, e', e''\}$) that contains it remains a triangle in F^{i+1} . Since these are the only faces in $F^{i+1} \setminus \{f^*\}$ that are affected by the contraction (note that $\{e, e', e''\}$ is deleted), every face $f \in F^{i+1} \setminus \{f^*\}$ is triangular.

Moreover, the only vertices in V^{i+1} whose degrees are changed by the single-edge contraction are the vertices $a(=b)$, and c . But we know that when $e' \notin E_A^i$, there exists a face $\{e', e_1, e_2\} \in F^{i+1}$ that contains both vertices. So, their degree is at least 2. Therefore, all vertices in V^{i+1} have degree at least 2.

Finally consider the case where an edge-pair contraction (Fig. 5b) is applied on D^i . Let $e = \{a, b\}, e' = \{b, c\} \in E_A^i$ be the edges involved. Note that we perform the edge-pair contraction only if there is no edge $e'' = \{a, c\}$ such that $\{e, e', e''\} \in F^i$, i.e., $\deg(b) > 2$. Therefore, there exist $f_1 = \{e, e_1, e_2\}, f_2 = \{e', e_3, e_4\} \in F^i$ such that $f_1 \neq f_2$. Since $e' \notin f_1$, and no edges are deleted by the contraction, the triangle f_1 remains a triangle in F^{i+1} . The same applies for f_2 . Since these are the only faces in $F^{i+1} \setminus \{f^*\}$ which are affected by the contraction, every face $f \in F^{i+1} \setminus \{f^*\}$ is triangular.

Moreover, the only vertices in V^{i+1} whose degrees are changed by the edge-pair contraction are the vertices $a(=c)$ and b . For b since we had $\deg(b) > 2$ in F^i , the degree remains no less than 2. For the combined vertex formed by the merging of a and c the two triangular faces f_1 and f_2 remain two distinct faces incident to it, its degree also remains at least 2.

Before proceeding further, let us also justify why the construction of G_1 was unnecessary in the special case during the edge-pair contraction, where we found the face $\{e, e', e''\} \in F^i$.

Lemma 18. *While performing an edge-pair contraction on a graph D^i on (e, e') , where $e = \{a, b\} \in E_A^i$ and $e' = \{b, c\} \in E_A^i$, if there exists some $e'' = \{a, c\}$ such that $\{e, e', e''\} \in F^i$, then $|\mathcal{C}_{D_1, I \cap E_{D_1}}(\ell_1)| = 1$ iff $\varepsilon(I \cap \tilde{E}_{D_1}) = \{\tilde{e}, \tilde{e}'\}$, and $|\mathcal{C}_{D_1, I \cap E_{D_1}}(\ell_1)| = 0$ otherwise.*

Proof. Let $f = \{e, e', e''\} \in F^i$. Observe that we only perform edge-pair contraction when there are no 0 labelled edges on E_A^i , which implies that $|E_A^i|$ is even as nonzero labels are paired. In particular, it is not the

case that E_A^i consists of precisely these three edges $\{e, e', e''\}$. Having excluded this particular possibility, we observe that none of the three types of operations (see Fig. 5) creates a triangular face containing two consecutive edges of the boundary curve. It follows that f is already a face in F^0 .

Now consider $D^0 = G_{D_1}$. The condition that led us to perform this edge-pair contraction step on (e, e') is because $\ell_2(e) = e'$ and $\ell_2(e') = e$. This stipulates that a cycle enters and then immediately exits D_1 on the two edges $\{e, e'\}$ of the same face f , and the cycle must complete itself in the disc D_2 as stipulated by the labelling ℓ_2 matching e and e' . Thus the unique possibility for the cycle is to have a simple path in D_2 connecting \tilde{e} and \tilde{e}' and finish off within D_1 by the trivial incursion in D_1 with f . This happens if and only if $\varepsilon(I \cap \tilde{E}_{D_1}) = \{\tilde{e}, \tilde{e}'\}$ (and in fact as a consequence in this case, they are consecutive in I and the arc orientations of \tilde{e}, \tilde{e}' in I match).

Therefore, $|\mathcal{C}_{D_1, I \cap E_{D_1}}(\ell_1)| = 1$ if $\varepsilon(I \cap \tilde{E}_{D_1}) = \{\tilde{e}, \tilde{e}'\}$, and is 0 otherwise.

With the graph G_1 constructed as above, we now need to construct an ordered set S_1 using $I \cap \tilde{E}_{D_1}$. We will construct S_1 in conjunction with G_1 . We will start with $S^0 = I \cap \tilde{E}_{D_1}$, and as we construct D^i using a contraction, we will also iteratively construct S^i , until we reach $S^\eta = S_1$.

In addition to constructing S^i , we need to construct a function ℓ_1^i (the construction of ℓ_2^i was already described earlier for D^i) on the set E_A^i , such that ℓ_1^i and ℓ_2^i are compatible, and $T(\ell_1^i, \ell_2^i) = \varepsilon(S^i \cap \tilde{E}_A^i)$. We start with $\ell_1^0 = \ell_1$, which clearly satisfies all requirements. Then, if D^{i+1} is constructed by a parallel-edge contraction, we let $\ell_1^{i+1} = \ell_1^i$. If D^{i+1} is constructed by a single-edge contraction, we let $\ell_1^{i+1} = \ell_1^i|_{E_A^{i+1}}$. If D^{i+1} is constructed by an edge-pair contraction, we have two cases to deal with. Let $e, e' \in E_A^i$ be the two edges that are identified with each other. If $\ell_1^i(e) = e'$ (and vice versa), we simply let $\ell_1^{i+1} = \ell_1^i|_{E_A^{i+1}}$. Otherwise, if $\ell_1^i(e) = e''$ and $\ell_1^i(e') = e'''$ (and vice versa), we let $\ell_1^{i+1}(e'') = e'''$ (and vice versa), and let ℓ_1^{i+1} be the same as ℓ_1^i for all other edges. It is apparent from the construction, that $\ell_1^{i+1}, \ell_2^{i+1}$ are compatible, if ℓ_1^i, ℓ_2^i are compatible. Note that after this edge-pair contraction on (e, e') , these two edges are removed from E_A^i , and ℓ_2^{i+1} is the restriction of ℓ_2^i on the remaining edges of E_A^i , namely E_A^{i+1} .

Now, getting back to the construction of S^{i+1} , we will ensure that S^{i+1} is an ordered set of arcs on the dual H^{i+1} of the graph D^{i+1} . Moreover, we will ensure that during the construction of S^{i+1} , we will never remove or change the order in which arcs appear in S^i . At most, we will identify arcs with each other, if the underlying edges are identified in D^{i+1} , and we will remove duplicates of the same arc if the duplicates appear consecutively after identification. We will then establish a recursive relation between $|\mathcal{C}_{D^i, S^i}(\ell_1^i)|$ and $|\mathcal{C}_{D^{i+1}, S^{i+1}}(\ell_1^{i+1})|$.

However, before we show that, let us consider a few cases where the construction of S^{i+1} , and with it the construction of G_1 , can be easily terminated.

Lemma 19.

1. Let D^{i+1} be constructed by a single-edge contraction. Let e be the edge that is deleted, and let e', e'' be the edges that are identified with each other (see Fig. 5a). Let f be the face that is deleted by the identification of e' with e'' , and $\tilde{e}' = \{\tilde{f}_1, \tilde{f}\}$, and $\tilde{e}'' = \{\tilde{f}, \tilde{f}_2\}$.
 - (a) If $\tilde{e} \in \varepsilon(S^i)$, then $|\mathcal{C}_{D^i, S^i}(\ell_1^i)| = 0$.
 - (b) If $\{\tilde{e}', \tilde{e}''\} \subseteq \varepsilon(S^i)$, but $((\tilde{f}_1 \rightarrow \tilde{f}), (\tilde{f} \rightarrow \tilde{f}_2))$ does not appear as consecutive arcs in the cyclic order of S^i , then $|\mathcal{C}_{D^i, S^i}(\ell_1^i)| = 0$.
2. Let D^{i+1} be constructed by a parallel-edge contraction. Let e, e' be the parallel edges, and let e'', e''' be the edges that are deleted (see Fig. 5c).
 - (a) If $(\ell_1^i, \ell_2^i) \neq (0, 0)$, and $\{\tilde{e}'', \tilde{e}'''\} \cap \varepsilon(S^i) \neq \emptyset$, then $|\mathcal{C}_{D^i, S^i}(\ell_1^i)| = 0$.
 - (b) If $(\ell_1^i, \ell_2^i) = (0, 0)$, and $\{\tilde{e}'', \tilde{e}'''\} \cap \varepsilon(S^i) \neq \emptyset$, but $\varepsilon(S^i) \not\subseteq \{\tilde{e}'', \tilde{e}'''\}$, then $|\mathcal{C}_{D^i, S^i}(\ell_1^i)| = 0$.
 - (c) If $(\ell_1^i, \ell_2^i) = (0, 0)$, and $\emptyset \neq \varepsilon(S^i) \subseteq \{\tilde{e}'', \tilde{e}'''\}$, then $|\mathcal{C}_{D^i, S^i}(\ell_1^i)| = 1$.
 - (d) If $\{\tilde{e}, \tilde{e}'\} \cap \varepsilon(S^i) \neq \emptyset$, then $|\mathcal{C}_{D^i, S^i}(\ell_1^i)| = 0$.

Proof. Recall that we perform single-edge contraction or parallel-edge contraction only when $\ell_2^i(e) = 0$. In case (1a), this contradicts the requirement that it must pass \tilde{e} . Similarly, in case (1b), given that no passage

is allowed at \tilde{e} , (by $\ell_2^i(e) = 0$), any passage using one of $\{\tilde{e}', \tilde{e}''\}$ must also use the other, and they must be for consecutive steps in the cyclic order of the interleaving, and in the correct orientation of the arcs.

In case (2) part (a)(b)(c), consider a PSAW $P \in \mathcal{C}_{D^i, S^i}(\ell_1^i)$. It has either \tilde{e}'' or \tilde{e}''' . Since a parallel-edge contraction was applied, $\ell_1^i(e) = \ell_2^i(e) = 0$. Given any vertex (other than the vertex \tilde{f}^*) in \tilde{F}^i , P must have either 0 or 2 edges incident on it. Thus P has both \tilde{e}'' or \tilde{e}''' . But since \tilde{e}'' and \tilde{e}''' form a cycle in H^i , it must be the case that $P = \{\tilde{e}'', \tilde{e}'''\}$.

This immediately proves (2a), (2b), and (2c). For (2d), given $\{\tilde{e}, \tilde{e}'\} \cap \varepsilon(S^i) \neq \emptyset$, any passage of a cycle using one of $\{\tilde{e}, \tilde{e}'\}$ must use both. But this contradicts $\ell_2(e) = 0$, which is the reason we performed this parallel-edge contraction on e .

Remark. Item 2 part (a)(b)(c) of [Lemma 19](#) combined takes care of the case $\{\tilde{e}'', \tilde{e}'''\} \cap \varepsilon(S^i) \neq \emptyset$ in a parallel-edge contraction step. After all cases in item 2 part (a)(b)(c)(d) are taken care of, we may assume $\{\tilde{e}, \tilde{e}', \tilde{e}'', \tilde{e}'''\} \cap \varepsilon(S^i) = \emptyset$ in a parallel-edge contraction step.

We now see how to construct S^{i+1} , assuming none of these special cases occur. First, let us consider the case where D^{i+1} is constructed from D^i by performing an edge-pair contraction on e and $e' \in E_A$. This step identifies e and e' with each other, but all the other edges stay as before. With a little abuse of notation we may assume that $e, e' \in E_A^0 = E_A$ (as no contraction step creates any edge on the boundary), such that $\ell_2(e) = e'$ (and vice versa). This means that $\tilde{e}, \tilde{e}' \in \varepsilon(T)$ by construction of T . Let $\tilde{e} = \{u_1, v_1\}$, and $\tilde{e}' = \{u_2, v_2\}$ with $u_1, v_1 \in \tilde{F}_{D_1}$, and $v_1, u_2 \notin \tilde{F}_{D_1}$ are dangling vertices. Since $\ell_2(e) = e'$, by the construction of T , we can see that $((u_1 \rightarrow v_1), (u_2 \rightarrow v_2))$ are consecutive arcs in T . Moreover, since $I \in \mathcal{I}_{S, T(\ell_1, \ell_2)}$ is a valid interleaving, we know that $((u_1 \rightarrow v_1), (u_2 \rightarrow v_2))$ are consecutive arcs in $I \cap \tilde{E}_{D_1}$ as any arcs between these two in I must belong to $\tilde{E}_{D_2} \setminus \tilde{E}_A$.

By the induction hypothesis, we assumed that during the construction of S^1, \dots, S^i , these arcs were not removed, or had their order changed. Therefore, $((u_1 \rightarrow v_1), (u_2 \rightarrow v_2))$ must also be consecutive arcs in S^i . Now, if we assume that f_1 is the unique face in \tilde{F}^i that contains the edge e , and f_2 is the unique face containing e' , we know by our choice of u_1, v_2 , that $\tilde{f}_1 = u_1$ and $\tilde{f}_2 = v_2$. Moreover, we have already taken care of the special case where $f_1 = f_2$ in [Lemma 18](#), so here, we may assume that $f_1 \neq f_2$. Now, when we identify the edges e and e' with each other in D^{i+1} , we see that f_1 and f_2 are now adjacent to each other. In other words, $(u_1, v_2) \in \tilde{E}^{i+1}$. Therefore, in S^{i+1} , we can replace the consecutive arcs $((u_1 \rightarrow v_1), (u_2 \rightarrow v_2))$ with the arc $((u_1 \rightarrow v_2))$. Note that S^{i+1} now satisfies the induction hypothesis.

Now, let us consider the case where D^{i+1} is constructed by performing a parallel-edge construction on D^i . Note that items 2 (a)(b)(c) in [Lemma 19](#) dealt with the case when $\{\tilde{e}'', \tilde{e}'''\} \cap \varepsilon(S^i) \neq \emptyset$, and then after items 2 (d), we only need to consider case $\{\tilde{e}, \tilde{e}', \tilde{e}'', \tilde{e}'''\} \cap \varepsilon(S^i) = \emptyset$. But in this case no edge in S^i is affected by the construction of D^{i+1} . So, we may let $S^{i+1} = S^i$.

Finally, let us consider the case where D^{i+1} is constructed by performing a single-edge contraction on D^i . We identify the edges $e' = \{a, c\}, e'' = \{b, c\} \in E^i$ with each other, and the edge $e = \{a, b\} \in E^i$ is deleted. But, all other edges stay as before. Since the special cases from [Lemma 19](#) have already been dealt with, we may assume that $\tilde{e} \notin \varepsilon(S^i)$. As for \tilde{e}' and \tilde{e}'' , unlike in the case of the edge-pair contractions, these edges are not guaranteed to belong to $\varepsilon(S^i)$. So, there are various possibilities in which these edges can appear (or not) in $\varepsilon(S^i)$. If neither of the edges appear, then we have nothing more to do, and the induction is already complete. Let us consider the case, where exactly one of the edges (say \tilde{e}' appears in $\varepsilon(S^i)$). Let $\tilde{e}' = \{u_1, v_1\}$, and $\tilde{e}'' = \{u_2, v_2\}$. Since e' and e'' both belong to the face $f = \{e, e', e''\}$, we may further assume that $\tilde{f} = v_1 = u_2$, and that $\tilde{f}_1 = u_1, \tilde{f}_2 = v_2$. In this case, note that because of the contraction, the faces f_1 and f_2 are now adjacent to each other. So, $\{u_1, v_2\} \in \tilde{E}^{i+1}$. Therefore, in S^{i+1} , we can replace any instance of $(u_1 \rightarrow v_1)$ with $(u_1 \rightarrow v_2)$, and any instance of $(v_1 \rightarrow u_1)$ with $(v_2 \rightarrow u_1)$. As we can see, in this case too, S^{i+1} satisfies the induction hypothesis.

Finally, let us consider the case where \tilde{e} and \tilde{e}' both belong to $\varepsilon(S^i)$. Once again, since the special cases from [Lemma 19](#) are already taken care of, we know that $((u_1 \rightarrow v_1), (u_2 \rightarrow v_2))$ (here $v_1 = u_2$) are consecutive arcs in S^i . So, we can replace these arcs with $(u_1 \rightarrow v_2)$. We have already seen how this satisfies our induction hypothesis.

Now that we have constructed G_1 and S_1 , we need to prove that they work as we intended.

Lemma 20. *Let the graph D^i and the ordered set S^i be such that they do not fall into the special cases from Lemma 19. Let D^{i+1} , and S^{i+1} be constructed as above. If D^{i+1} is constructed by an edge-pair contraction, or single-edge contraction, then*

$$|\mathcal{C}_{D^i, S^i}(\ell_1^i)| = |\mathcal{C}_{D^{i+1}, S^{i+1}}(\ell_1^{i+1})|.$$

If D^{i+1} is constructed by a parallel-edge contraction, then

$$|\mathcal{C}_{D^i, S^i}(\ell_1^i)| = |\mathcal{C}_{D^{i+1}, S^{i+1}}(\ell_1^{i+1})| + \mathbb{1}_{\{(\ell_1^i, \ell_2^i) = (0, 0); \varepsilon(S^i) = \emptyset\}}.$$

Proof. Consider $P^i \in \mathcal{C}_{D^i, S^i}(\ell_1^i)$. Since ℓ_1^i is compatible with ℓ_2^i by construction, it follows that $P^i \cup \mu_{D^i}(\ell_2^i)$ is a cycle in $H^i = H_{D^i}$. We will now use P^i to construct a $P^{i+1} \in \mathcal{C}_{D^{i+1}, S^{i+1}}(\ell_1^{i+1})$.

First, let us consider the case where D^{i+1} was constructed using an edge-pair contraction. As we have seen during the construction of S^{i+1} , this means that the edges $\tilde{e} = \{u_1, v_1\} \in \tilde{E}_A^i$ and $\tilde{e}' = \{u_2, v_2\} \in \tilde{E}_A^i$ are replaced with the edge $\{u_1, v_2\}$. Now, since $\ell_2^i(e) \neq 0$ and $\ell_2^i(e') \neq 0$ (this is because we perform edge-pair contraction only for the paired edge by ℓ_2), we know that $\{\{u_1, v_1\}, \{u_2, v_2\}\} \subseteq P^i$. We construct P^{i+1} by just replacing these two edges in P^i with $\{u_1, v_2\}$.

Clearly, $P^{i+1} \subseteq \tilde{E}_{D^{i+1}}$. Consider $P^{i+1} \cup \mu_{D^{i+1}}(\ell_2^{i+1})$. We already know that $P^i \cup \mu_{D^i}(\ell_2^i)$ is a cycle in H^i . In this cycle, we just need to replace the segment $\{\{u_1, v_1\}, \{v_1, u_2\}, \{u_2, v_2\}\}$ (where $\{u_1, v_1\}$ and $\{u_2, v_2\}$ belong to P^i and $\{v_1, u_2\}$ belongs to $\mu_{D^i}(\ell_2^i)$) with the edge $\{u_1, v_2\}$ to get $P^{i+1} \cup \mu_{D^{i+1}}(\ell_2^{i+1})$. So, $P^{i+1} \in \mathcal{P}_{D^{i+1}}$.

Now, we need to show that $P^{i+1} \cup \mu_{D^{i+1}}(\ell_2^{i+1})$ conforms to S^{i+1} . By our choice of S^{i+1} , this is apparent, since we replaced the consecutive arcs $((u_1 \rightarrow v_1), (u_2 \rightarrow v_2))$ in S^i with $(u_1 \rightarrow v_2)$. So, if P^i conforms to S^i , P^{i+1} has to conform to S^{i+1} . Moreover, by definition of ℓ_1^{i+1} , it is easy to see that $\ell_{P^{i+1}} = \ell_1^{i+1}$. Therefore, $P^{i+1} \in \mathcal{C}_{D^{i+1}, S^{i+1}}(\ell_1^{i+1})$.

Clearly, this is an invertible transformation. So, when D^{i+1} is constructed using an edge-pair contraction,

$$|\mathcal{C}_{D^i, S^i}(\ell_1^i)| = |\mathcal{C}_{D^{i+1}, S^{i+1}}(\ell_1^{i+1})|.$$

Now, let us consider the case where D^{i+1} was constructed using a single-edge contraction. Once again, as we saw during the construction of S^{i+1} , the edges $\{u_1, v_1\}, \{v_1, v_2\} \in \tilde{E}^i$ are replaced by the edge $\{u_1, v_2\} \in \tilde{E}^{i+1}$. If P^i does not contain either of the deleted edges, we have nothing to do. It is immediate that $P^{i+1} \in \mathcal{C}_{D^i, S^i}(\ell_1^i)$. On the other hand, assume that P^i contains one of these edges (say $\{u_1, v_1\}$). We know that $P^i \cup \mu_{D^i}(\ell_2^i)$ is a cycle in H^i . So, we know that there has to be another edge in P^i that is incident on v_1 . But this edge cannot be \tilde{e} , since $\ell_2^i(e) = 0$ by our choice of e (as the edge where we perform a single-edge contraction). Therefore, P^i must contain the only other edge $\{v_1, v_2\}$ that is incident on v_1 . Therefore, in P^{i+1} , we may replace these two edges with $\{u_1, v_2\}$.

We need to show that $P^{i+1} \in \mathcal{P}_{D^{i+1}}$. To do that, consider $P^{i+1} \cup \mu_{D^{i+1}}(\ell_2^{i+1})$. Note that in the cycle $P^i \cup \mu_{D^i}(\ell_2^i)$, we are replacing the segment $\{\{u_1, v_1\}, \{v_1, v_2\}\}$ with the edge $\{u_1, v_2\}$ to obtain $P^{i+1} \cup \mu_{D^{i+1}}(\ell_2^{i+1})$. So, $P^{i+1} \in \mathcal{P}_{D^{i+1}}$. Moreover, since the special case from Lemma 19 has not occurred, we know that even if $\{\{u_1, v_1\}, \{v_1, v_2\}\} \subseteq \varepsilon(S^i)$, from the construction of S^{i+1} and ℓ_1^{i+1} , we see that $P^{i+1} \in \mathcal{C}_{D^{i+1}, S^{i+1}}(\ell_1^{i+1})$.

Clearly, this is also an invertible transformation. So, when D^{i+1} is constructed using a single-edge contraction,

$$|\mathcal{C}_{D^i, S^i}(\ell_1^i)| = |\mathcal{C}_{D^{i+1}, S^{i+1}}(\ell_1^{i+1})|.$$

Finally, we consider the case where D^{i+1} was constructed using a parallel-edge contraction. Since the special cases from Lemma 19 have not come to pass, we know that $\{\tilde{e}, \tilde{e}', \tilde{e}'', \tilde{e}'''\} \cap \varepsilon(S^i) = \emptyset$. In this case, as we saw in the proof of Lemma 19, if \tilde{e}'' or \tilde{e}''' belongs to P^i , then in fact, $P^i = \{\tilde{e}'', \tilde{e}'''\}$. So, if $(\ell_1^i, \ell_2^i) \neq (0, 0)$, or if $\varepsilon(S^i) \neq \emptyset$, it follows that $\{\tilde{e}'', \tilde{e}'''\} \cap P^i = \emptyset$. Since each vertex in H^i can have either 0 or 2 edges from P^i incident on it, we also find that $\tilde{e}, \tilde{e}' \notin P^i$. Therefore, we can let $P^{i+1} = P^i$.

Clearly, $P^{i+1} \cup \mu_{D^{i+1}}(\ell_2^{i+1})$ is a cycle, and it conforms to S^{i+1} , since $S^{i+1} = S^i$, and $\{\tilde{e}, \tilde{e}', \tilde{e}'', \tilde{e}'''\} \cap \varepsilon(S^i) = \emptyset$. Therefore, $P^{i+1} \in \mathcal{C}_{D^{i+1}, S^{i+1}}(\ell_1^{i+1})$.

Clearly, this is an invertible transformation. So, when D^{i+1} is constructed using a parallel-edge contraction, when $(\ell_1^i, \ell_2^i) \neq (0, 0)$ or $\varepsilon(S^i) \neq \emptyset$,

$$|\mathcal{C}_{D^i, S^i}(\ell_1^i)| = |\mathcal{C}_{D^{i+1}, S^{i+1}}(\ell_1^{i+1})|.$$

The last case to deal with is when $(\ell_1^i, \ell_2^i) = (0, 0)$, and $\varepsilon(S^i) = \emptyset$. Assume $P^i \neq \{\tilde{e}'', \tilde{e}'''\}$. Once again, we let $P^{i+1} = P^i$. Clearly, $P^{i+1} \in \mathcal{C}_{D^{i+1}, S^{i+1}}(\ell_1^{i+1})$, just as we saw in the previous case. So, this transformation from $\mathcal{C}_{D^i, S^i}(\ell_1^i) \setminus \{\{\tilde{e}'', \tilde{e}'''\}\}$ to $\mathcal{C}_{D^i, S^i}(\ell_1^i)$ is invertible. However, in this case, $P^i = \{\tilde{e}'', \tilde{e}'''\}$ is also possible. Therefore, when $(\ell_1^i, \ell_2^i) = (0, 0)$, and $\varepsilon(S^i) = \emptyset$,

$$|\mathcal{C}_{D^i, S^i}(\ell_1^i)| = |\mathcal{C}_{D^{i+1}, S^{i+1}}(\ell_1^{i+1})| + 1$$

Theorem 2. *Given a graph $G = (V, E, F)$ with $|V| = \Omega(1)$ with a cycle A dividing into two discs D, D' , a labelling $\ell \in \mathcal{L}_A$, and a set I of ordered arcs such that $\varepsilon(I) \subseteq \tilde{E}_D$, we can compute $|\mathcal{C}_{D, I \cap \tilde{E}_D}(\ell)|$ in polynomial time, or construct a graph $G_1 = (V_1, E_1, F_1)$, and an ordered set of arcs S_1 , such that $|V_1| \leq \frac{3|V|}{4}$, and $|\mathcal{C}_{D, I \cap \tilde{E}_D}(\ell)|$ can be computed from $|\mathcal{W}_{G_1, S_1}|$ in polynomial time.*

Proof. In the case where G_1 is constructed, this follows directly from Lemma 20, since the construction takes polynomial time, and we know that $|\mathcal{C}_{D^0, S^0}(\ell_1^0)| = |\mathcal{C}_{D_1, I \cap \tilde{E}_{D_1}}(\ell_1)|$, and that $|\mathcal{C}_{D^\eta, S^\eta}(\ell_1^\eta)| = |\mathcal{W}_{G_1, S_1}|$.

There are two cases in which G_1 would not be constructed. The first is the case when applying an edge-pair contraction on edges $e, e' \in \tilde{E}_A^i$, we found another edge e'' , such that $\{e, e', e''\} \in F^i$. The correctness in this case was established in Lemma 18.

The other case is when we came across the special cases from Lemma 19 during the construction of S^{i+1} . Since $\varepsilon(S^i) \neq \emptyset$ in this case anyway, from Lemma 20, we see that $|\mathcal{C}_{D_1, I \cap \tilde{E}_{D_1}}(\ell_1)| = |\mathcal{C}_{D^0, S^0}(\ell_1^0)| = |\mathcal{C}_{D^i, S^i}(\ell_1^i)|$, and the rest follows from Lemma 19.

D Appendix D

Lemma 9. *There is a constant α , such that any problem $|\mathcal{W}_{G_l, S_l}|$ on level l of the tree \mathcal{T} can be computed in time*

$$\alpha^{\sqrt{m_l}} \frac{(t_0 + \dots + t_{L-1})!}{(t_0 + \dots + t_{l-1})!(t_l)! \dots (t_{L-1})!}.$$

Proof. As leaf nodes have constant size, the base case is clearly true. Assume the lemma statement is true for all problems at level $> k$. Consider a problem $|\mathcal{W}_{G_k, S_k}|$ at level k of \mathcal{T} .

We know from Fact 1 that $|\mathcal{L}_{(A_k)}|$ is the number of Motzkin paths on E_{D_k} , which is at most κ^{t_k} for a fixed constant κ . Now, we know that $|S_k| \leq t_0 + \dots + t_{k-1}$, and for any compatible (ℓ_1, ℓ_2) , we have $|T(\ell_1, \ell_2)| \leq t_k$. Therefore, from Lemma 14, we get that for each compatible (ℓ_1, ℓ_2) ,

$$|\mathcal{I}_{S_k, T(\ell_1, \ell_2)}| \leq 2t_k \binom{t_0 + \dots + t_k}{t_k}.$$

Since each compatible (ℓ_1, ℓ_2) and a valid I result in at most two subproblems, the number of children of this problem at level k is at most

$$4(\kappa^{t_k})^2 t_k \binom{t_0 + \dots + t_k}{t_k}.$$

By inductive hypothesis we know that solving each of these subproblems takes time at most

$$\alpha^{\sqrt{m_{k+1}}} \frac{(t_0 + \dots + t_{L-1})!}{(t_0 + \dots + t_k)!(t_{k+1})! \dots (t_{L-1})!}.$$

Now, note that given the problem $|\mathcal{W}_{G_k, S_k}|$, checking if any given $\ell_1, \ell_2 \in \mathcal{L}_{(A_k)}$ are compatible, and checking if $I \in \mathcal{I}_{S_k, T(\ell_1, \ell_2)}$ is valid takes polynomial time (as a function of the number of vertices in the graph). So does the reduction from $|\mathcal{C}_{(D_k), S_k}(\ell)|$ to $|\mathcal{W}_{G_{k+1}, S_{k+1}}|$. In total, we can assume that determining and finally combining the solutions to each of the children of the problem $|\mathcal{W}_{G_k, S_k}|$ takes $p(m_k)$ time, where $p(\cdot)$ is a polynomial function.

Combining everything, the total time taken to solve the problem at level k is at most:

$$\begin{aligned} & 4t_k p(m_k) \left(\kappa^{2t_k} \frac{(t_0 + \dots + t_k)!}{(t_0 + \dots + t_{k-1})!(t_k)!} \right) \left(\alpha^{\sqrt{m_{k+1}}} \frac{(t_0 + \dots + t_{L-1})!}{(t_0 + \dots + t_k)!(t_{k+1})! \dots (t_{L-1})!} \right) \\ &= 4t_k p(m_k) (\kappa^{2t_k} \alpha^{\sqrt{m_{k+1}}}) \left(\frac{(t_0 + \dots + t_{L-1})!}{(t_0 + \dots + t_{k-1})!(t_k)! \dots (t_{L-1})!} \right) \end{aligned}$$

Note that $4t_k p(m_k) = 4(\sqrt{8m_k})p(m_k)$ can be bounded by a polynomial of m_k . So there is a constant γ , such that the time taken to solve the problem at level k is at most

$$\gamma^{\sqrt{m_k}} \alpha^{\sqrt{\frac{3}{4}m_k}} \frac{(t_0 + \dots + t_{L-1})!}{(t_0 + \dots + t_{k-1})!(t_k)! \dots (t_{L-1})!},$$

and $\gamma^{\sqrt{m_k}} \alpha^{\sqrt{\frac{3}{4}m_k}} < \alpha^{\sqrt{m_k}}$ for a large constant α . This completes the induction.

Theorem 3. *There exists a constant β such that given any connected planar 3-regular multi-graph (with no loops) H with n vertices, the number of cycles on H can be computed in $\beta^{\sqrt{n}}$ time.*

Proof. Let $G = (V, E, F)$ be the dual of the graph H with $|V| = m$. By the 3-regularity of H , and by Euler's formula, $m = n/2 + 2$. By Lemma 9, computing $|\mathcal{W}_G|$ takes time

$$\alpha^{\sqrt{m}} \frac{(t_0 + \dots + t_{L-1})!}{(t_0)! \dots (t_{L-1})!}.$$

Applying Lemma 21 with $r = \sqrt{3/4}$, and $a = \sqrt{8m}$ completes the proof.

Lemma 21. *For any $0 \leq r < 1$, there exists a constant δ such that for any $a \geq 0$ and $k \geq 0$,*

$$\frac{(a + ar + \dots + ar^k)!}{(a)!(ar)! \dots (ar^k)!} \leq \delta^{a+ar+\dots+ar^k}.$$

Proof. Take $\delta = 2^{\frac{1}{1-r}}$. We prove this by induction on k . When $k = 0$, the statement is clearly true as $\delta \geq 1$. Assume that the statement is true for $k - 1$. Then,

$$\begin{aligned} \frac{(a + ar + \dots + ar^k)!}{(a)!(ar)! \dots (ar^k)!} &= \frac{(a + ar + \dots + ar^k)!}{a!(ar + \dots + ar^k)!} \frac{(ar + \dots + ar^k)!}{(ar)! \dots (ar^k)!} \\ &\leq 2^{a+\dots+ar^k} \delta^{ar+\dots+ar^k} \\ &\leq \left(2^{\frac{1}{1-r}} \right)^a \delta^{ar+\dots+ar^k} \\ &= \delta^{a+ar+\dots+ar^k}. \end{aligned}$$

E Appendix E

In order to prove Theorem 4, we will need the following analogue of Lemma 4:

Lemma 22.

$$|\mathcal{W}_{G,S}^{X,Y}| = \sum_{\text{compatible } \ell_1, \ell_2 \in \mathcal{L}_A} |\mathcal{C}_S^{X,Y}(\ell_1, \ell_2)|.$$

Let $X_D = X \cap \tilde{E}_D, Y_D = Y \cap \tilde{E}_D$ for $D \in \{D_1, D_2\}$. If $(\ell_1, \ell_2) \neq (0, 0)$ is a compatible pair of labellings then

$$|\mathcal{C}_S^{X,Y}(\ell_1, \ell_2)| = \sum_{\text{valid } I \in \mathcal{I}(S, T(\ell_1, \ell_2))} |\mathcal{C}_{D_1, I \cap \tilde{E}_{D_1}}^{X_{D_1}, Y_{D_1}}(\ell_1)| \times |\mathcal{C}_{D_2, I \cap \tilde{E}_{D_2}}^{X_{D_2}, Y_{D_2}}(\ell_2)|$$

If $(\ell_1, \ell_2) = (0, 0)$, then

- If $\emptyset \neq \varepsilon(S) \subseteq \tilde{E}_D$ and $X \subseteq \tilde{E}_D$ for $D \in \{D_1, D_2\}$, then $\left| \mathcal{C}_S^{X,Y}(0,0) \right| = \left| \mathcal{C}_{D,S}^{X,Y_D}(0) \right|$.
- If $\varepsilon(S) = \emptyset$ and $\emptyset \neq X \subseteq \tilde{E}_D$ for $D \in \{D_1, D_2\}$, then $\left| \mathcal{C}_S^{X,Y}(0,0) \right| = \left| \mathcal{C}_{D,S}^{X,Y_D}(0) \right|$.
- If $\varepsilon(S) = \emptyset$ and $X = \emptyset$, then $\left| \mathcal{C}_S^{X,Y}(0,0) \right| = \left| \mathcal{C}_{D_1,S}^{X,Y_{D_1}}(0) \right| + \left| \mathcal{C}_{D_2,S}^{X,Y_{D_2}}(0) \right| - 1$.
- In all other cases, $\left| \mathcal{C}_S^{X,Y}(0,0) \right| = 0$.

The problem of computing $\left| \mathcal{C}_{D_1, I \cap \tilde{E}_{D_1}}^{X_{D_1}, Y_{D_1}}(\ell_1) \right|$ and $\left| \mathcal{C}_{D_2, I \cap \tilde{E}_{D_2}}^{X_{D_2}, Y_{D_2}}(\ell_2) \right|$ for any $I \in \mathcal{I}(S, T(\ell_1, \ell_2))$ is also similar to the previous setting. The stitching technique used in [Section 3.2](#) can be used to compute these quantities. There are a few things we need to be careful about. During the construction of G_1 (and similarly, G_2), while performing a single-edge contraction or parallel-edge contraction, if we delete an edge e from the graph, we need to verify that $\tilde{e} \notin X$. We also need to ensure that while performing any contraction, we never identify two edges e and e' with each other such that $\tilde{e} \in X$ and $\tilde{e}' \in Y$. If these two events occur, then $\left| \mathcal{C}_{D_1, I \cap \tilde{E}_{D_1}}^{X_{D_1}, Y_{D_1}}(\ell_1) \right| = 0$. Otherwise, the reduction works similar to [Lemma 20](#). Overall, if we account for X, Y while choosing the values for the base cases, the proof of [Theorem 4](#) is nearly identical to that of [Theorem 3](#).

We will now extend the algorithm to count cycles of a fixed length. We will need to define a few terms to do this.

Definition 8. Let $X, Y \subseteq \tilde{E}$, and let $l \geq 0$. Then,

$$\mathcal{W}_G^{X,Y,l} = \{W \in \mathcal{W}_G^{X,Y} : |W| = l\}.$$

Let $X, Y \subseteq \tilde{E}_D$ for some $D \in \{D_1, D_2\}$, and let $l \geq 0$. Then,

$$\mathcal{P}_D^{X,Y,l} = \{P \in \mathcal{P}_D^{X,Y} : |P| = l\}.$$

Similarly, we define $\mathcal{W}_{G,S}^{X,Y,l}$, $\mathcal{P}_{D,S}^{X,Y,l}$, $\mathcal{C}_{D,S}^{X,Y,l}(\ell)$, and $\mathcal{C}_S^{X,Y,l}(\ell_1, \ell_2)$.

Lemma 23.

$$\left| \mathcal{W}_{G,S}^{X,Y,l} \right| = \sum_{\text{compatible } \ell_1, \ell_2 \in \mathcal{L}_A} \left| \mathcal{C}_S^{X,Y,l}(\ell_1, \ell_2) \right|.$$

Let $X_D = X \cap \tilde{E}_D, Y_D = Y \cap \tilde{E}_D$ for $D \in \{D_1, D_2\}$. If $(\ell_1, \ell_2) \neq (0,0)$ is a compatible pair of labellings then

$$\left| \mathcal{C}_S^{X,Y,l}(\ell_1, \ell_2) \right| = \sum_{\text{valid } I \in \mathcal{I}(S, T(\ell_1, \ell_2))} \sum_{i=0}^{l-l'} \left| \mathcal{C}_{D_1, I \cap \tilde{E}_{D_1}}^{X_{D_1}, Y_{D_1}, l'+i}(\ell_1) \right| \times \left| \mathcal{C}_{D_2, I \cap \tilde{E}_{D_2}}^{X_{D_2}, Y_{D_2}, l'+(l-i)}(\ell_2) \right|$$

where $l' = |\{e \in E_A : \ell_1(e) \neq 0\}|$.

If $(\ell_1, \ell_2) = (0,0)$ then

- If $\emptyset \neq \varepsilon(S) \subseteq \tilde{E}_D$ and $X \subseteq \tilde{E}_D$ for $D \in \{D_1, D_2\}$, then $\left| \mathcal{C}_S^{X,Y,l}(0,0) \right| = \left| \mathcal{C}_{D,S}^{X,Y_D,l}(0) \right|$.
- If $\varepsilon(S) = \emptyset$ and $\emptyset \neq X \subseteq \tilde{E}_D$ for $D \in \{D_1, D_2\}$, then $\left| \mathcal{C}_S^{X,Y,l}(0,0) \right| = \left| \mathcal{C}_{D,S}^{X,Y_D,l}(0) \right|$.
- If $\varepsilon(S) = \emptyset$ and $X = \emptyset$, then $\left| \mathcal{C}_S^{X,Y,l}(0,0) \right| = \left| \mathcal{C}_{D_1,S}^{X,Y_{D_1},l}(0) \right| + \left| \mathcal{C}_{D_2,S}^{X,Y_{D_2},l}(0) \right| - \mathbb{1}_{\{l=0\}}$, where $\mathbb{1}_{\{l=0\}} = 1$ if $l = 0$, and 0 otherwise.
- In all other cases, $\left| \mathcal{C}_S^{X,Y,l}(0,0) \right| = 0$.

Now, consider the problem of computing $\left| \mathcal{C}_{D_1, I \cap \tilde{E}_{D_1}}^{X_{D_1}, Y_{D_1}, j}(\ell_1) \right|$, where $T = T(\ell_1, \ell_2)$, and $I \in \mathcal{I}_{S,T}$. We use the same stitching technique detailed in [Section 3.2](#) to construct the graph G_1 . However, (referring to notations

in Section 3.2) because of the way edges in D may get identified with each other, the relations between the lengths of the cycles in G_1 , and the lengths of PSAWs in D can get complicated. Let

$$\psi : \mathcal{C}_{D_1, I \cap \tilde{E}_{D_1}}^{X_{D_1}, Y_{D_1}}(\ell_1) \rightarrow \mathcal{W}_{G_1, S_1}^{X_{D_1}, Y_{D_1}}$$

be the invertible function, whose existence is guaranteed by Theorem 2.

Let R be the set of all edges in G_{D_1} that are either merged with another edge, or deleted, by a single-edge contraction or edge-pair contraction, during the construction of G_1 . Note that

$$R \subseteq \bigcup \left\{ \tilde{e}, \tilde{e}', \tilde{e}'' : \{e, e', e''\} \in F_{D_1}, e \in E_A \right\}$$

i.e., R is a subset of the set of all edges belonging to the faces in D_1 that contain edges from A . Therefore, $|R| \leq 3|E_A|$. We have a disjoint union

$$\mathcal{C}_{D_1, I \cap \tilde{E}_{D_1}}^{X_{D_1}, Y_{D_1}, l}(\ell_1) = \bigsqcup_{R_X \subseteq R} \mathcal{C}_{D_1, I \cap \tilde{E}_{D_1}}^{X_{D_1} \cup R_X, Y_{D_1} \cup (R \setminus R_X), l}(\ell_1)$$

Consider some $R_X \subseteq R$. For simplicity, let $R_Y = R \setminus R_X$. We now want to compute $\left| \mathcal{C}_{D_1, I \cap \tilde{E}_{D_1}}^{X_{D_1} \cup R_X, Y_{D_1} \cup R_Y, l}(\ell_1) \right|$. The reduction to G_1 is the same as before, but this time, we have a much better handle on the size of the cycles and PSAWs. Just like with S^{i+1} and X^{i+1}, Y^{i+1} , we also iteratively construct $R_X^{i+1} \subseteq R_X^i$ such that R_X^{i+1} is a set of all the distinct edges in R_X^i from E^{i+1} . We construct R_Y^{i+1} similarly. Following the same ideas as in Lemma 19 and Lemma 20, we can compute $\left| \mathcal{C}_{D_1, I \cap \tilde{E}_{D_1}}^{X_{D_1} \cup R_X, Y_{D_1} \cup R_Y, l}(\ell_1) \right|$.

The only thing that would have to be changed in Lemma 19 is in case (2c). When we apply a parallel-edge contraction, and delete the edges e'', e''' . Let d'', d''' be the number of distinct edges in E_{D_1} that are identified with e'', e''' respectively, in the graph D^i . We modify case (2c) to say $\left| \mathcal{C}_{D^i, S^i}^{X^i \cup R_X^i, Y^i \cup R_Y^i, l} \right| = 1$ if and only if $l = d'' + d'''$, and otherwise, it is 0. It goes without saying that we should also account for X^i, Y^i here, and claim that it is 1 only if $X^i \subseteq \{\tilde{e}'', \tilde{e}'''\}$.

Then, we use the following lemma to compute $\left| \mathcal{C}_{D_1, I \cap \tilde{E}_{D_1}}^{X_{D_1} \cup R_X, Y_{D_1} \cup (R \setminus R_X), l}(\ell_1) \right|$.

Lemma 24. *Let the graph D^{i+1} be such that it does not fall into the special cases from Lemma 19. Let $S^{i+1}, X^{i+1}, Y^{i+1}, R_X^{i+1}, R_Y^{i+1}$ be constructed as explained. Let $r = |R_X^i|$, and $r' = |R_X^{i+1}|$. If D^{i+1} is constructed by an edge-pair contraction, or single-edge contraction, then*

$$\left| \mathcal{C}_{D^i, S^i}^{X^i \cup R_X^i, Y^i \cup R_Y^i, l}(\ell_1^i) \right| = \left| \mathcal{C}_{D^{i+1}, S^{i+1}}^{X^{i+1} \cup R_X^{i+1}, Y^{i+1} \cup R_Y^{i+1}, l+r'-r}(\ell_1^{i+1}) \right|.$$

If D^{i+1} is constructed by a parallel-edge contraction, with edges e'', e''' deleted from D^i , let d'', d''' be the number of edges in E_{D_1} identified with e'', e''' in D^i , then

$$\left| \mathcal{C}_{D^i, S^i}^{X^i \cup R_X^i, Y^i \cup R_Y^i, l}(\ell_1^i) \right| = \left| \mathcal{C}_{D^{i+1}, S^{i+1}}^{X^{i+1} \cup R_X^{i+1}, Y^{i+1} \cup R_Y^{i+1}, l+r'-r}(\ell_1^{i+1}) \right| + \mathbb{1}_{\{(\ell_1^i, \ell_2^i) = (0, 0); \varepsilon(S^i) = \emptyset; l = d'' + d'''\}}.$$

Therefore, the algorithm in Section 3.3 can be easily adapted to compute $|\mathcal{W}_G^{X, Y, l}|$. During each level of the tree we constructed in Section 3.3, given a problem $|\mathcal{W}_{G_k, S_k}^{X_k, Y_k}|$, we compute $|\mathcal{W}_{G_k, S_k}^{X_k, Y_k, l}|$ for all $l \leq |\tilde{E}_k|$. The correctness of this algorithm can be easily verified. Let us now consider the time complexity of this algorithm.

In Section 3.3, we saw that a problem at level k of the tree has at most $4(\kappa^{t_k})^2 t_k \binom{t_0 + \dots + t_k}{t_k}$ subproblems as children. In this setting, the number of children can be bounded by

$$4(\kappa^{t_k})^2 t_k \binom{t_0 + \dots + t_k}{t_k} O(m_k 2^{3t_k})$$

The 2^{3t_k} factor is because that is the maximum possible number of choices for $R_X \subseteq R$, and the factor $O(m_k)$ is a bound on all possible lengths of cycles we have to consider. It is easy to see from the proof of Theorem 3 that this extra factor can be absorbed into the $\beta^{\sqrt{n}}$ term.

Theorem 5. *There exists a constant β such that given any connected 3-regular planar multi-graph (with no loops) H with n vertices, and any $X, Y \subseteq \tilde{E}$, the number of cycles on H of a fixed length l , that traverse each edge in X , and do not traverse any edge in Y , can be computed in $\beta^{\sqrt{n}}$ time.*

Just as we are able to count the number of cycles on H of a fixed length, we can count the number of cycles on H that divide H into two subgraphs with an p faces on one subgraph, and $|\tilde{V}| - p$ faces on the other. The extra factor this produces can be absorbed into the $\beta^{\sqrt{n}}$ term.