Digraph Networks and Groupoids

Nyumbu Chishwashwa The University of the Western Cape Cape Town, South Africa

Vance Faber and Noah Streib Center for Computing Sciences Bowie, Maryland

Abstract. We answer a question posed in [3]. We prove that every vertex transitive digraph has a spanning factorization; in fact, this is a necessary and sufficient condition. We show that 1-factorization of a regular digraph is closely related to the notion of a Cayley graph of a groupoid and as such, the theorem we prove on spanning factorizations can be translated to a 2006 theorem of Mwambene (4; Theorem 9) on groupoids. We also show that groupoids are a powerful tool for examining network routing on general regular digraphs. We show there is a 1-1 relation between regular connected digraphs of degree d and the Cayley graphs of groupoids (not necessarily associative but with left identity and right cancellation) with d generators. This enables us to provide compact algebraic definitions for some important graphs that are either given as explicit edge lists or as the Cayley coset graphs of groups larger than the graph. One such example is a single expression for the Hoffman-Singleton graph.

Introduction. In this paper, we define a regular digraph D to be a set of vertices V and directed edges E such that the in-degree and out-degree of every vertex is a constant d and there is a directed path between any two vertices. (For technical reasons, we allow multiple edges but no loops.) These digraphs are often employed as the model for network connections in parallel computing, and a critical problem in that regard is optimization of the network with respect to global communication tasks. See for example [3]. In this note, we associate an algebraic structure with every regular digraph which allows us to extend notions used in highly symmetric graphs to more general graphs. There are several definitions of groupoids. Here, we define a groupoid as a set Γ equipped with a binary (not necessarily associative) relation * called *product* (which we often suppress). A *word* w in the elements of Γ is a finite sequence of elements. The *value* of w is the element formed by taking the product in the order given. A subset S of Γ is (non-trivial) generating set for Γ (which we denote by $\Gamma = \langle S \rangle$) if Γ is the set of products of finite words in S. In addition, we impose some additional conditions on our finitely generated groupoids:

- 1) the empty word which we call e is a left identity,
- 2) $x \notin xS$ for all $x \in \Gamma$,
- 3) vs = us for $v, u \in \Gamma$ and $s \in S$ only if v = u (right cancellation).

Given a finitely generated groupoid $\Gamma = \langle S \rangle$, we denote each element by a minimum length word which generates it. The properties 1 and 2 can be simply stated as right multiplication by an element of *S* is a fixed point-free permutation on Γ . It should be noted that property 1 is independent of the other two properties by examining a simple example (see the particular assignment for the 6 vertex Kautz graph in Example 2 below.) In many cases, we want our groupoids to have an additional property we call *left cancellation on S*:

4} vs = vt for $v \in \Gamma$ and $s, t \in S$ only if s = t.

Factorizations and spanning factorizations. This section reviews the definitions from [3].

A *factorization* $F = \{F_1, F_2, ..., F_d\}$ of a regular digraph of degree *d* is disjoint decomposition of the edges such each vertex is the in-vertex for one edge from each F_i . By Petersen's Theorem, every regular digraph has a factorization.

Fact 1. Every directed graph G where the in-degree and out-degree of every vertex is d has an edge disjoint decomposition into d 1-factors.

Proof. Form an auxiliary graph *B* with two new vertices u' and u'' for each vertex u. The edges of *B* are the pairs (u',v'') where (u,v) is a directed edge in *G*. The undirected graph *B* is bipartite and regular with degree *d* and so by Hall's Marriage Theorem, it can be decomposed into *d* 1-factors. Each of these 1-factors corresponds to a directed 1-factor in *G*.

Definition. Let F_1 , F_2 , \cdots , F_d be the factors in a 1-factoring of Γ . We call a finite string of symbols from the set $\{F_i\}$ a word. If v is a vertex and ω is a word, then $v\omega$ denotes the directed path (and its endpoint) in Γ starting at v and proceeding along the unique edge corresponding to each consecutive factor represented in the word ω . We say that a 1-factoring is a *spanning factorization* of the digraph Γ with n vertices if there exists a set $W = \{\emptyset, \omega_1, \omega_2, \dots, \omega_{n-1}\}$ of n words such that for every v the vertices $v\omega_i$ are distinct. We say the factorization is *tree-like* if W is a tree.

Definition. A schedule associated with a factorization is an assignment of a time (a label) to each occurrence of each factor in the words of W such that no time is assigned more than once to a particular factor and times assigned to the factors in a single word are increasing. The *time of a schedule* is the largest time assigned to any of the factors. If T is the total time, the schedule can be thought of as a $d \times T$ array where each row corresponds to a factor and an entry in that row indicates which occurrence of that factor has been assigned the corresponding time. An entry in a row in the array can be empty indicating no occurrence of that factor has been assigned the given time.

The power of a spanning factorization lies in the fact that a schedule can be used to describe an algorithm for conflict free global exchange of information between the vertices of the graph. If the factorization is tree-like, the schedule also provides an algorithm for other communication primitives like global broadcast or summation. This is discussed at length in [5].

Fact 2. Suppose we have a schedule for a factorization of the graph *G*. Then the collection of directed label-increasing paths $v\omega_i$ for all v and non-empty ω_i have the property that no edge in the graph is assigned the same time twice. A schedule for a spanning factorization yields a time labeled directed path between every two vertices so that no edge is labeled with the same time twice.

Proof. Each edge in the graph is assigned to a single one factor. Assume there is an edge in the one factor F that has been assigned the same time twice. Since every occurrence of F in the words in W has been assigned a unique time, this can only mean that there are two different vertices u and v and an initial subword ω of a word in W such that the edges $(u\omega, u\omega F)$ and

 $(v\omega, v\omega F)$ are the same edge. Then $u\omega$ and $v\omega$ must be the same vertex. Let us assume that this is the shortest ω for which this happens. The word ω cannot be empty since u and v are different. But then the last factor in ω must also be the same edge, a contradiction. If we start with a spanning factorization, then all the non-empty paths from v are unique, there are n-1 of them and none of them can return to v so they must reach to every other vertex in the graph.

Vertex symmetric graphs. A digraph G is *vertex transitive* if for any two vertices u and v there is an automorphism of G which maps u to v. Suppose G is a vertex transitive. We want to know if it has a spanning factorization.

We will also need to use the Cayley coset representation of a vertex transitive digraph.

Definition (Cayley coset graph). Let Γ be a finite group, H a subgroup and S a subset. Suppose

- (i) $S \cap H = \emptyset$ and Γ is generated by $\Delta \cup H$,
- (ii) $HSH \subseteq SH$,
- (iii) S is a set of distinct coset representatives of H in Γ .

Then we can form the *Cayley coset digraph* $G = (\Gamma, S, H)$ with the cosets $\{gH : g \in \Gamma\}$ as vertices and the set of pairs (gH, gsH) with $s \in S$ as edges. When H is the identity subgroup, the graph is a *Cayley digraph*.

The classic proof of Sabidussi [13] shows that a digraph is vertex transitive if and only if it is a Cayley coset digraph. An important aspect of the proof shows that one can construct a Cayley coset digraph from a vertex transitive digraph by using the automorphism group as the group Γ required in the definition and the subgroup of automorphisms that fix a vertex as the required subgroup H. The generators *S* correspond to automorphisms that map a vertex to a neighbor.

Example of what can go wrong. We want to show that every vertex transitive digraph (Cayley coset digraph) has a spanning factorization. In fact, we want to create an algorithm that can find a spanning factorization for any vertex transitive graph. We know from Fact 1 above that a regular digraph has a 1-factorization. However, that 1-factorization may not be spanning. In [3], it took a great deal of effort to create a spanning factorization for some particular vertex transitive digraphs. Alternatively, we can try to use a set of coset representatives to label the vertices of the Cayley coset diagraph, but there is no obvious way to create a factorization which is necessary for creating conflict-free network routing. Our main result will resolve this problem but first we give an example that illustrates it.

We start with a directed version of the Petersen graph. We can use the Cayley coset definition. The group is a semidirect product group on the two generators α and θ with

 $\alpha^4 = 1 = \theta^5$

 $\alpha^{-1}\theta\alpha = \theta^2$.

The subgroup is $H = \{1, \alpha^2\}$. The edges in the graph are the cycle $(H, \theta H, \theta^2 H, \theta^3 H, \theta^4 H)$, the edges $(\theta^i H, \theta^i \alpha H)$ and $(\theta^i \alpha H, \theta^i H)$ and the cycle $(\alpha H, \alpha \theta H, \alpha \theta^2 H, \alpha \theta^3 H, \alpha \theta^4 H)$. (Note that it takes some calculation to determine the labels for this last set cosets.) If you form a 1-factor with these labels, you get something that can be used to obtain a spanning factorization. However, these are not the only choices for coset representatives; in each case, there are two choices. It turns out that if you choose the other coset representative for the 5 vertices in the interior of the pentagon you get trouble. In

clockwise order, these new labels are $\theta^i \alpha^2 \theta^{-1}$ for i = 0, 1, 2, 3, 4. The problem comes when you try to write down the 1-factor using these coset representatives and the generator θ . The outside cycle works as expected. But the two distinct vertices $u = \theta^{-1}H$ and $v = \alpha^2 \theta^{-1}H$ give $u\theta = v\theta$ so these edges can't both be a part of a 1-factor.

Our next definition is motivated by the fact that if the subgroup H is normal in Γ , then the spanning factorization is unique up to permutation of the generators. This factorization was the original routing algorithm used in [5].

Definition. A Cayley coset digraph $G = (\Gamma, S, H)$ is irreducible, if given *s* and *t* in *S*, there exists an $h \in H$ such that sH = htH. In other words, *H* acts transitively on *SH*.

Fact 3. A digraph is both vertex and edge transitive if and only if it is an irreducible Cayley coset digraph.

Proof. Let the vertex transitive graph be $G = (\Gamma, S, H)$. If this graph is edge transitive, then for each *s* and *t* in *S*, there exists an automorphism θ which maps (H, sH) to (H, tH). In particular, since θ fixes the vertex *H*, it must be an element of *H*. But since it also maps sH to tH this says that *H* acts transitively on *SH*. Conversely, suppose *H* acts transitively on *SH* and (uH, usH) and (vH, vtH) are two edges in *G*. Let sH = htH. Then

(uH, usH) = (uhH, uhtH)

so the automorphism $v(uh)^{-1}$ maps (uH, usH) to (vH, vtH).

Notation. We let g(H,sH) = (gH,gsH).

Theorem 1. Let $G = (\Gamma, S, H)$ is an irreducible Cayley coset digraph. Then given any 1-factor F and $r \in S$ there is a choice of coset representatives R and such that $F = F_r = \{g(H, rH) | g \in R\}$. Furthermore, given s = hr, $F_s = \{hg(H, rH) | g \in R\}$ is also a 1-factor. The set $\{F_s | s \in S\}$ formed in this way is a 1-factorization of G.

Proof. We want to create a set of coset representatives R which satisfy the theorem. We construct F_r first. Start with any 1-factor D in some 1-factorization and any initial set of coset representatives Q. Now we create R from Q by

replacing $f \in Q$ with a suitable g in the same coset. Suppose $f(H, sH) \in D$. There exists an $h \in H$ such that s = hr so

$$f(H,sH) = f(H,hrH) = fh(H,rH)$$

and if we replace f by g = fh, we have f(H, sH) = g(H, rH) so the edges $F_r = \{g(H, rH) | g \in R\}$ are identical to the given 1-factor D.

We want to show that if we have any 1-factor $F_1 = \{g(H, sH) | g \in R\}$ and $h \in H$ such that hs = t then $F_2 = \{hg(H, sH) | g \in R\}$ is also a 1-factor which is edge-disjoint from F_1 . It's clearly a 1-factor because h is an automorphism. If F_1 and F_2 are not disjoint, then there exists $f, g \in R$ such that g(H, sH) = hf(H, sH) for some $f, g \in R$. This is a pair of equations

gH = hfH

and

$$gsH = hfsH$$
).

Thus there exists $k \in H$ such that g = hfk which gives gs = hfks. But then

$$hfsH = gsH = gsH = hfksH$$

which yields

$$sH = ksH = tH$$

which is false.

Theorem 2. If (Γ, S, H) is an irreducible Cayley coset digraph and F is a 1-factorization constructed in Theorem 1, then given any set of words in F,

 $W = \{w_0 = \emptyset, w_1 = F_1, w_2 = F_2, \dots, w_d = F_d, w_{d+1}, \dots, w_{n-1}\},\$

comprising a unique path from a given root vertex to each other vertices in G forms a spanning factorization.

Proof. By theorem 1, the words can also be taken to be words in the group elements s_i . Given $gw_iH = gw_jH$ then clearly $w_iH = w_jH$. Thus W is a spanning factorization of G.

Theorem 3. Given a Cayley coset digraph $G = (\Gamma, S, H)$, there is a decomposition of *S* into disjoint subsets S_i such that each $G_i = (\Gamma, S_i, H)$ is irreducible.

Proof. We form an equivalence relation on S, $s \sim t$ if and only if there exists an $h \in H$ such that hsH = tH. This is clearly an equivalence relation and each equivalence class S_i has the property that given s and t in S_i there is an h such that hsH = tH. Now given any $h \in H$ and $s \in S_i$ then there is some $t \in S$ such that hsH = tH. This t must be an element of S_i because S_i is an equivalence class. In particular, $HS_iH = S_iH$.

First, we show the graph G_i is independent of coset representatives. Consider an edge g(H,tH) with $g \in \Gamma$ and $t \in S_i$. We must show that g(hH,hsH) = g(H,tH) for some $s \in S_i$. But this follows from $HS_iH = S_iH$.

Second, we show Γ acts transitively on G_i . Let $g \in \Gamma$. Then if (H, sH) with $s \in S_i$ is an edge at the vertex H in G_i , g maps it to (gH, gsH) which is still an edge in G_i .

This shows that each G_i is an irreducible Cayley coset digraph.

Corollary 4. Given any Cayley coset digraph $G = (\Gamma, S, H)$, it has a 1-factorization with factors labeled by elements of *S*.

Proof. Since G_i is an irreducible Cayley coset graph, any 1-factorization has a representation by group elements from Theorem 1. The union of these 1-factorizations is a 1-factorization of G.

Theorem 5. If (Γ, S, H) is a Cayley coset digraph, it has a spanning tree-like factorization.

Proof. Suppose *F* is a factorization given by Corollary 4. Take a tree *W* of paths from *H* to each *gH* consisting of words in in *F*. If *w* is a path, the product of the edge labels along the path correspond to an element $\overline{w}H$ in *G*. Let *fH* be any other vertex in *G* and suppose (fH)u = fHv for words *u* and *v* in *W*.

Then $fh\overline{u} = f\overline{v}$ for some $h \in H$ so $h\overline{u} = \overline{v}$. This means that $H\overline{u} = H\overline{v}$ in the group and so eu = ev in the graph so u = v. This proves F is a spanning factorization.

Theorem 6. If G is a digraph with a spanning factorization

 $W = \{w_0 = \emptyset, w_1 = F_1, w_2 = F_2, \dots, w_d = F_d, w_{d+1}, \dots, w_{n-1}\}$

then G is vertex transitive.

Proof. First, we show that for each vertex F_i there is an automorphism θ_i with $\theta_i(u) = F_i u$. If $\theta_i(u) = \theta_i(v)$ then $F_i u = F_i v$ so by the definition of spanning factorization, u = v; θ_i is 1-1. Now suppose (u, uF_j) is an edge. Then since $\theta_i(u) = F_i u$ and $\theta_i(uF_j) = F_i uF_j$, $(\theta_i(u), \theta_i(uF_j))$ is also an edge. This shows that θ_i is an automorphism. Second, using induction, we can construct an automorphism mapping vertices u to v by taking a word w in the F_i such that uw = v.

Theorem 7. A digraph is vertex transitive if and only if it has a spanning factorization. In that case, it also has a tree-like spanning factorization.

Proof. This is just a summary of the previous statements.

Partial groupoid tables and their extensions. In a finitely generated groupoid $\Gamma = \langle S \rangle$ we can assume that every element has been labeled by a word in a breadth first spanning tree on *S* starting with the empty word *e*. Then the *d* columns corresponding to the generators encode the *d* edges in the graph emanating from each vertex *u*. In other words, the edges are exactly (u, u * s). We call these *d* columns a partial groupoid and we have assumed that they are fixed point free permutations of the elements. We call any groupoid with the same columns an *extension* of the partial groupoid. One way of extending the table to the whole groupoid is by defining *uw* inductively by using associativity on the words in the set *W*. We call this the canonical extension from *W*. We have to show this is a well-formed definition, that is, it satisfies the property of right cancellation.

Theorem 8. The canonical extension is a groupoid.

Proof. We have to show that the extension has right cancellation. We can do this by induction on the tree W. Suppose we start with two distinct elements u and v and the shortest word w such that $uw \neq vw$. Clearly w is not a generator

because that would violate right cancellation on the partial groupoid. Thus there is a word z in W and generator s such that z is the parent of w, that is, w = zs. Now we have defined

uw = (uz)s

and

vw = (vz)s.

Again by right cancellation of the partial groupoid

uz = vz.

Since z is shorter than w, we have right cancellation and we have reached a contradiction on the assumption that

 $u \neq v$.

Associated digraph of a groupoid. Given a groupoid $\Gamma = \langle S \rangle$, we can create an associated digraph G(S) with directed edges (u, us) for each $u \in \Gamma$ and $s \in S$. By analogy from group theory, we call this the *Cayley graph* of the groupoid.

Definition. Given an ordering on $S = \{s_1, s_2, ..., s_d\}$ we can label the vertices and edges of G(S) recursively using a spanning tree of words in S. The vertex e is called level L_0 . Given vertices in level L_j , consider the equivalence relation on $L_j \times S$ given by $(u,s) \sim (v,t)$ if us = vt. Create the vertices in level L_{j+1} by taking one vertex ws_i from each equivalence class. We call this a *tree-like labeling* of the groupoid. We can create a 1-factorization of G(S) by letting $F_i = \{L_j s_i \mid j \ge 0\}$. We call this the *associated factorization* of the labeling.

Lemma 9. A tree-like labeling assigns a label to every element in Γ .

Proof. If not, let w be the shortest word not labeled. Then the initial word v of w has been labeled since it is shorter and there is an s such that vs = w. Thus the algorithm will label w with s.

Note that as far as the Cayley graph of the groupoid is concerned, it is completely determined by the partial groupoid. All that is needed is to fill the other columns with fixed point free permutations. There can be no inconsistency problem because there is no way to calculate vw from the table when w is not a

generator. As soon as w is the product of two or more generators, say st then we need to know if v(st) can be deduced from the partial table which would require associativity. Generally, we choose to use the canonical extension.

Characterization. We show there is a 1-1 relation between a regular connected digraph with a factorization F and a finitely generated groupoid with associated digraph G(S) and a tree-like labeling.

Theorem 10. The digraph G(S) is regular and the sets of edges $F_{s_i} = \{(u, us_i) | u \in \Gamma\}$ from a tree-like labeling form a factorization. Conversely, given a regular digraph D with factorization $F = \{F_i\}$ and root \emptyset , we can form a finitely generated groupoid Γ .

Proof. Clearly the out-degree of G(S) is d. So the average in-degree is also d. If not all vertices have in-degree d then there is some vertex u with in-degree greater than d. This in turn means there is some s and distinct vertices x and y such that xs = ys. But by right cancellation (property 3), this means that x = y, a contradiction. This shows G(S) is regular. The argument used to show that all the elements are labeled shows that the digraph is strongly connected. We have defined the F_{s_i} so that they are a factorization. Conversely, if D is a regular digraph with factorization $F = \{F_i\}$, we can label the vertices of D with the algorithm above so that each vertex is given by a unique word in F. If we want to form u * v where u and v are two vertices labeled this way, the result is the vertex found by starting at \emptyset following the path to u and then following the path given by the word v.

This theorem allows us to represent a regular digraph by an algebraic structure. We can sum up the result like this:

There is a one-to-one mapping between Cayley graphs of groupoids with d generators and regular digraphs with degree d.

Theorem 11. Given a groupoid Γ with generating set S, the digraph G(S) is vertex transitive if and only if there is a tree-like labeling of the vertices such that the canonical extension has left cancellation.

Proof. This is just a translation of Theorem 7.

Checking vertex transitivity of a Cayley digraph. It would be good to have an algorithm to check if a Cayley graph G(S) of a groupoid Γ is vertex transitive. If we can find the subgroup H of the automorphism group which fixes the identity,

we can use the methods in the proof of Theorem 7 to efficiently make this check. The first step is to break S into subsets invariant under the action of H. Then G(S) is vertex transitive if and only if the subgraphs determined by these invariant subsets are vertex transitive. So without loss of generality, we can assume that H acts transitively on S. Following the construction in the proof of Theorem 1, we start by assuming we have a fixed tree-like labeling of the vertices of G(S) and any arbitrary 1-factor D. Now we are going to replace the groupoid Γ with a new groupoid Υ which has the same vertices and the same Cayley graph. We fix an element r in S and suppose (u, us) is in the 1-factor D. We define the new operation * in Υ so that for this particular u, u * t = u(ht) where hr = s. Now D consists of edges (u, u * r). At this point, a 1-factorization of Υ is given by factors $F_s = \{(u, u * s) | u \in \Upsilon\}$. Furthermore, we showed that the Cayley graph is vertex transitive if and only if any tree-like labeling of Υ is a spanning factorization, in other words, Υ has left cancellation.

Examples. We give some examples that illustrate how groupoids can be used to describe digraphs.

Example 1. This first example is a groupoid on $\mathbb{Z}_2 \times \mathbb{Z}_3$. Multiplication is defined by the table

	00	01	02	10	11	12
00	00	01	02	10	11	12
01	01	02	10	12	00	01
02	02	10	11	01	02	10
10	10	11	12	00	01	02
11	11	12	00	02	10	11
12	12	00	01	11	12	00

Note that in fact the columns are fixed point free permutations but the rows are not. The generators are t = (1,0) and s = (0,1). The generator *s* produces a 6-cycle in the Cayley graph. The set $H = \{e,t\}$ is a subgroup and Hs and Hs^2 are disjoint cosets. However, in the Cayley graph, *H* is a 2-cycle and $Hs \cup Hs^2$ is a 4-cycle. This graph is the Kautz graph G(2,3).

Example 2. The following groupoid on $\mathbb{Z}_2 \times \mathbb{Z}_3$ satisfies the 2nd and 3rd axioms but not the 1st. It is finitely generated by t = (1,0) and s = (0,1) but the empty word can't be assigned a consistent meaning. The problem is that it functions on the right as an identity but not on the left. This shows that axiom 1 is independent of the other axioms.

	00	01	02	10	11	12
00	00	01	02	11	12	10
01	01	02	00	10	11	12
02	02	00	01	12	10	11
10	10	11	12	01	02	00
11	11	12	10	00	01	02
12	12	10	11	02	00	01

Even though this is not a groupoid in the sense we are using, it still has a Cayley graph. The graph is the directed graph as in Example 1.

Example 3. We define a groupoid on $\mathbb{Z}_2 \times \mathbb{Z}_p \times \mathbb{Z}_p$ by

$$(a,b,c)*(x,y,z) = (a+x,b-bx+y,c+(-1)^{x}by+2^{a}z)$$

for p a prime. In the case p = 5,

$$S = \{(0,0,1), (0,0,4), (1,0,0), (1,1,0), (1,2,0), (1,3,0), (1,4,0)\}$$

produces the Hoffman-Singleton graph [2] which is an undirected graph of degree 7 and diameter 2.

Example 4. The Alegre graph [1] is the largest known degree 2 diameter 4 digraph. It has 25 vertices. We can represent it as the Cayley graph G of a groupoid on $\mathbb{Z}_5 \times \mathbb{Z}_5$. Again the elements are $t^i s^j$. This time we use the lexicographic ordering to represent the elements as natural numbers from 0 to 24 on the Hamiltonian cycle determined by s. Then it is only necessary to describe the factor determined by t. It consists of a 5-cycle starting at 0, another 5-cycle starting at 3 and a 15-cycle. We denote this by

(05101520)(32318138)(117242112191671411296).

Cyclic difference sets. The degree 2 examples above all have the property that one factor is a Hamiltonian cycle of some length n = ab and the other has a disjoint cycle containing 0 and of length a. We say that the factor system (and equivalently the groupoid) is derived from a *cyclic difference set*. We show how to use cyclic difference sets in general to produce groupoids with 2 generators.

We call the two 1-factors Y and Z. If the n vertices V are labeled from the elements of \mathbb{Z}_n , then each of the 1-factors is a permutation in S_n . In order to simplify the problem, we will assume that the cycle structure of the 1-factors is

derived from a difference set. In particular, we will assume that G is Hamiltonian and Z is a spanning cycle.

First, we start with a subgroup U of V with b = |U| a divisor of n. We take the canonical set of coset representatives of U in V to be $0 \le i \le a - 1$. Let π be a permutation on $0 \le i \le a - 1$. We let $\{v_0, v_1, \dots, v_{a-1}\}$ be a sequence of elements of U where n = ab. Consider the graph

$$Y(\pi, (v_i)) = \{(i+u, \pi(i)+u+v_i) \mid 0 \le i \le a-1, u \in U\}.$$

Lemma 11. The map given by Y is a permutation on V.

Proof. By definition every element of *V* has a unique representation i + u so we only need to prove that *Y* is 1-1. Suppose $\pi(i) + u + v_i = \pi(j) + v + v_j$. But then $\pi(i) = \pi(j)$ and since π is a permutation, i = j. Thus $u + v_i = v + v_i$ which means that u = v.

Lemma 12. The cycle of the permutation *Y* containing the element i + u has length αc where *c* is the length of the cycle of π containing *i* and $\alpha > 0$ is smallest such that $\alpha v = 0$ in *U* where

$$v = v_i + v_{\pi(i)} + v_{\pi^2(i)} + \ldots + v_{\pi^{t-1}(i)}.$$

Proof. Start at i + u and apply Y repeatedly. We get a sequence of elements

$$\pi^{J}(i) + u + v_{i} + v_{\pi(i)} + v_{\pi^{2}(i)} + \dots + v_{\pi^{j-1}(i)}$$

and if this repeats i + u, we must have

$$i = \pi^{j}(i)$$

and

$$0 = v_i + v_{\pi(i)} + v_{\pi^2(i)} + \ldots + v_{\pi^{j-1}(i)}.$$

This can only happen when $j = \alpha c$ and so

$$\pi^{\alpha c}(i) + u + \alpha (v_i + v_{\pi(i)} + v_{\pi^2(i)} + \dots + v_{\pi^{t-1}(i)}) = i + u + \alpha v$$

and the theorem follows.

If we choose the following values for the parameters:

 $V = \mathbb{Z}_{25}$ n = 25 a = 5 b = 5 $\pi = (0, 2, 4)$ $U = \{0, 5, 10, 15, 20\}$ $(v_0, v_1, v_2, v_3, v_4) = (5, 20, 20, 5, 20)$

then 1-factor Y is

(0,7,4,20,2,24,15,22,19,10,17,14,5,12,9)(1,21,16,11,6)(3,8,13,18,23).

Lemma 13. Let w be any element of U. The map $\theta: v \to v + w$ is an automorphism of G.

Proof. Note that $\theta(i+u) = i + (u+w)$, $\theta(i+u+1) = i + (u+w) + 1$ and $\theta(\pi(i) + u + v_i) = \pi(i) + (u+w) + v_i$.

Lemma 14. The map $\sigma(i) = i + 1 \pmod{a}$ creates an isomorphism between the graph $Y(\pi, (v_i))$ and $Y(\pi', (w_i))$ where $\pi'(i) = \pi(i-1) + 1 \pmod{a}$ and $w_0 = v_{a-1} - a$, $w_{\pi^{-1}(a-1)+1} = v_{\pi^{-1}(a-1)} + a$ and $w_i = v_{i-1}$ otherwise.

Proof. This isomorphism just renames the vertex $j \in \mathbb{Z}_n$ by j+1 and adjusts the sequence (v_i) accordingly.

If we apply this isomorphism twice to the graph defined just above, we get a new set of parameters for an isomorphic graph:

 $V = \mathbb{Z}_{25}$ n = 25 a = 5 b = 5 $\pi = (4,1,2)$ $U = \{0,5,10,15,20\}$ $(v_0, v_1, v_2, v_3, v_4) = (20,15,5,5,0)$

and Y factor as given in Example 4 for the Alegre graph.

Search complexity. If we want to search the space of all $Y(\pi, (v_i))$, the number we have to search is nominally $a!b^a$. However, there is no point in considering the isomorphic graphs from Lemma 4 so this reduces the space to the size $(a-1)!b^{a-1}$. There may be other symmetries, for example, we can negate all the entries in \mathbb{Z}_n to get an isomorphic graph.

Calculating the diameter. To calculate the diameter, we only need to calculate the distance from each vertex *i* with $0 \le i \le a - 1$ because of Lemma 3.

References

- 1. Miguel A. Fiol, J. Luis Andres Yebra and Ignacio Alegre (1984), Line digraph iterations and the (d,k) digraph problem, *IEEE Transactions on Computers*, C-33(5):400-403.
- 2. Alan J. Hoffman, Robert R. Singleton (1960), Moore graphs with diameter 2 and 3, *IBM Journal of Research and Development*, 5(4):497-504.
- 3. Randall Dougherty and Vance Faber (2014), Network routing on regular directed graphs from spanning factorizations, *Bulletin of Mathematical Sciences and Applications*, 20:9-24.
- 4. Eric Mwambene (2006), Representing vertex-transitive graphs on groupoids, Quaestiones Mathemaicae 29(2006), 279-284.
- 5. Vance Faber, Global communication algorithms for hypercubes and other Cayley coset graphs, Technical Report, LA-UR 87-3136, Los Alamos National Laboratory (1987).