Recent results from the TOTEM collaboration and the discovery of the odderon

C. Royon[1],

1 Department of Physics and Astronomy, The University of Kansas, Lawrence, USA
* christophe.royon@cern.ch

August 24, 2022

21st International Symposium on Very High Energy Cosmic Ray Interactions (ISVHECRI 2022)
Online, 23-27 May 2022
doi:10.21468/SciPostPhysProc.?

arXiv:2208.10782v1 [hep-ex] 23 Aug 2022

Abstract

We describe the most recent results from the TOTEM collaboration on elastic, inelastic and total cross sections as well as the odderon discovery by the D0 and TOTEM collaborations.

Contents

1 Measurements of the elastic, inelastic and total cross sections by the TOTEM collaboration 1

2 The odderon discovery by the TOTEM and D0 collaborations 4

3 Conclusion 7

References 8

In this short report, we will first discuss the most recent measurements from the TOTEM collaboration of elastic, inelastic and total cross sections at different center-of-mass energies of the LHC, namely 2.76, 7, 8 and 13 TeV. In a second part of this report, we will discuss the odderon discovery obtained by the D0 and TOTEM collaborations by comparing pp and $p\bar{p}$ elastic interactions.

1 Measurements of the elastic, inelastic and total cross sections by the TOTEM collaboration

To study elastic collisions at the LHC between protons, one detects and measures the intact protons in dedicated detectors called roman pots. Each proton is scattered at some angles and can lose

or gain some momentum. The fact that the protons are intact is due to the fact that a colorless object (at least two gluons) is exchanged between the two protons. The advantage of the TOTEM detector is that there are two inelastic telescopes called $T1$ and $T2$ that were installed in the very forward region covering the domain in rapidity of $3.1 < |\eta| < 4.7$ and $5.3 < |\eta| < 6.5$ allowing to veto on any particle produced in the very forward region, or to select cleanly elastic events. In addition, it is possible to run the LHC for different center-of-mass energies (2.76, 7, 8 and 13 TeV), with different values of β^* allowing to increase the domain of measurement in $|t|$, the 4-momentum transferred square at the proton vertex.

The TOTEM collaboration developed different methods to measure the total cross section σ_{tot} based on the optical theorem

$$L\sigma_{tot}^2 = \frac{16\pi}{1+\rho^2}(dN_{el}/dt)_{t=0} \quad (1)$$

$$L\sigma_{tot} = N_{el} + N_{inel} \quad (2)$$

where L is the luminosity, $\rho = \frac{Re(A^N(0))}{Im(A^N(0))}$, the ratio of the real and the imaginary part of the nuclear amplitude at $t = 0$, and N_{el} and N_{inel} the number of elastic and inelastic events. Three different methods can thus be defined to measure the total cross section σ_{tot}, namely the luminosity independent measurement

$$\sigma_{tot} = \frac{16\pi}{(1+\rho^2)}\frac{(dN_{el}/dt)_{t=0}}{(N_{el}+N_{inel})} \quad (3)$$

the luminosity dependent measurement

$$\sigma_{tot}^2 = \frac{16\pi}{(1+\rho^2)}\frac{1}{L}(dN_{el}/dt)_{t=0} \quad (4)$$

and the ρ independent measurement

$$\sigma_{tot} = \sigma_{el} + \sigma_{inel}. \quad (5)$$

All methods lead to consistent measurement of σ_{tot} at different center-of-mass energies [1–5].

The measurements by the TOTEM collaboration of the elastic, inelastic and total cross sections are displayed in Fig. 1. There is a good agreement with the cosmic ray measurements at high energies (within uncertainties) and a discrepancy of about 1.9σ has been found with ATLAS [6] at 8TeV [1].

The TOTEM collaboration also measured the differential elastic pp $d\sigma/dt$ cross sections at center-of-mass energies of 2.76, 7, 8 and 13 TeV, that allows to measure the B slope of the elastic cross section as a function of $\sqrt{s}$ as shown in Fig. 2, left. The B slope is found to be larger at 13 TeV and a linear behavior (in lns) of the elastic $d\sigma/dt$ cross section is only observed for $\sqrt{s} < 3$ TeV, whereas it is no longer the case at higher energies [8–10]. The increase of σ_{el}/σ_{tot} with energy is confirmed at LHC energies, as shown in Fig. 2, right.

A simple exponential fit to $d\sigma/dt$ at low t fails to describe the elastic data at low $|t|$. The TOTEM collaboration performed an exponential fit of elastic data as $d\sigma/dt = A\exp(-B(t)|t|)$ where different polynomial fits of $B(t)$ can be performed

- $N_b = 1$ $B = b_1$

[1] A recent measurement of the total cross section from ATLAS at 13 TeV was recently released at the ICHEP 2022 conference after the ISVHECRI workshop and shows the same level of discrepancy between ATLAS and TOTEM [7].




Figure 1: Elastic, inelastic and total cross section measurements as a function of the center-of-mass energy $\sqrt{s}$.




Figure 2: B slope as a function of $\sqrt{s}$ and increase of the elastic cross section as a function of energy.

- $N_b = 2, B = b_1 + b_2 t$
- $N_b = 3, B = b_1 + b_2 t + b_3 t^2$

where N_b gives the number of parameters in the fit. A pure simple exponential form ($N_b = 1$) is excluded at 7.2 σ with 8 TeV data [9] and similar results were found using using 13 TeV data [10]. Data cannot distinguish between a linear or a quartic dependence of B on $|t|$.

The very precise TOTEM elastic data at 13 TeV and the fact that it was possible to use very high β^* data (about 2 km) allowed to perform a precise measurement of elastic pp $d\sigma/dt$ over a wide range in $|t|$ and to cover the Coulomb-Nuclear interference (CNI) region at $|t| \sim 10^{-4} - 10^{-3}$ GeV^2 where $d\sigma/dt$ is sensitive to the amplitude in the Coulomb and nuclear regions

$$\frac{d\sigma}{dt} \sim |A^C + A^N(1 - \alpha G(t))|^2. \tag{6}$$

In the CNI region, both the modulus and the phase of the nuclear amplitude can be used to determine the ρ parameter, where the modulus is constrained by the measurement in the hadronic

region and the phase by the t dependence. The total cross section and the ρ measurements performed by the TOTEM experiment are displayed in Fig. 3 as a function of $\sqrt{s}$ [11, 12]. The total cross section measurements are compared with different series of parametrizations from COMPETE [13], namely $\sigma_{tot} = a + bln s + cln^2 s$ in blue, $\sigma_{tot} = a + bln^2 s$ in purple, and $\sigma_{tot} = a + bln s$ in green. It is clear that σ_{tot} favors the blue series of parametrizations whereas ρ favors the green ones, especially at 13 TeV where $\rho = 0.09 \pm 0.01$. This creates a tension between these two measurements. It is worth noting that none of these parametrizations contains an odderon contribution, and adding the exchange of the Odderon in addition to the Pomeron at high energies removes the tension between these measurements. This result can thus be interpreted as a first evidence for the odderon. For the models included in COMPETE and the Durham models [14], the TOTEM ρ measurement at 13 TeV provides a 3.4 to 4.6σ significance.




Figure 3: Measurement of ρ and σ_{tot} from TOTEM as a function of $\sqrt{s}$.

2 The odderon discovery by the TOTEM and D0 collaborations

In this section, we will describe the odderon discovery by the TOTEM and D0 collaborations [15–17]. As we already mentioned in the previous section, elastic pp or $p\bar{p}$ scattering at high energies is due to the exchange of colorless objects, namely Pomeron and Odderon [2]. Pomeron and Odderon [20–27] correspond respectively to positive and negative C parity. From the QCD point of view, the pomeron is made of and even number of gluons (2, 4, 6, etc) which leads to a (+1) parity whereas the odderon is made of and odd number of gluons (3, 5, 7, etc) corresponding to a (−1) parity. The scattering amplitudes can be written as:

$$A_{pp} = Even + Odd \quad (7)$$
$$A_{p\bar{p}} = Even - Odd. \quad (8)$$

From the equations above, it is clear that observing a difference between pp and $p\bar{p}$ interactions would be a clear way to observe the odderon, and this is the strategy followed by the TOTEM and D0 collaborations.

The elastic pp $d\sigma/dt$ cross section measurements for different $\sqrt{s}$ by the TOTEM collaboration [8–10, 18, 19] are shown in Fig. 4 together with the measurement for $p\bar{p}$ at the Tevatron by the D0 collaboration. pp elastic cross sections always show the same features, namely a decreasing cross section at low $|t|$, the presence of a minimum, the “dip" and of a maximum, the “bump"

[2] At lower energies such as the ISR, the situation is more complicated since there can be in addition exchanges of mesons and reggeons.

at medium $|t|$ and finally a decreasing cross section at higher $|t|$. On the contrary, the $p\bar{p}$ cross section does not show any bump or dip. The first observable that shows a difference between pp and $p\bar{p}$ elastic $d\sigma/dt$ cross sections is the ratio of the cross sections at the bump and at the dip as a function of $\sqrt{s}$ [15, 28] as displayed in Fig. 5 for LHC and Tevatron data [29, 30] as well as from measurements from fixed targets or colliders at lower $\sqrt{s}$ [31–35]. The ratio in pp elastic collisions decreases as a function of $\sqrt{s}$ up to $\sim$ 100 GeV and is flat above. The D0 $p\bar{p}$ data show a ratio of 1.00±0.21 given the fact that no bump nor dip is observed in the data within uncertainties. It leads to a difference of more than 3σ between pp and $p\bar{p}$ elastic data (assuming a flat behavior above $\sqrt{s} = 100GeV$ for the pp ratio).




Figure 4: Left: Comparison of elastic pp $d\sigma/dt$ cross section measurements for different $\sqrt{s}$ by the TOTEM collaboration. Right: Elastic $p\bar{p}$ $d\sigma/dt$ cross section measurements from the D0 collaboration at 1.96 TeV.




Figure 5: Bump over dip ratio from the elastic $d\sigma/dt$ cross section measurements for pp and $p\bar{p}$ interactions.

In order to get a better comparison of pp and $p\bar{p}$ elastic data, it is needed to extrapolate the elastic pp $d\sigma/dt$ data to 1.96 TeV, the center-of-mass energy of the Tevatron [3]. In order to predict the pp elastic $d\sigma/dt$ cross section at 1.96 TeV, we define eight characteristic points of the behavior of the elastic cross section at 2.76, 7, 8 and 13 TeV and study the dependence of the $|t|$ and $d\sigma/dt$ values of these characteristic points as a function of $\sqrt{s}$. As we mentioned already,

[3] Unfortunately, running the LHC at 1.96 TeV, the Tevatron energy, is not easy and would lead to zero acceptance in the dip/bump region where we compare pp and $p\bar{p}$ elastic data because of the roman pot detector acceptance.

the $d\sigma/dt$ distributions for pp collisions show a dip and a bump as a function of $|t|$ (see Fig. 4). The chosen characteristic points are thus the dip. the $bump$, $dip-2$ and $bump-2$ (same $d\sigma/dt$ as a the dip and the bump but at lower and higher $t|$), $mid1$, $mid2$ (where the cross section is the average between the ones at the bump and at the dip), and $bump+5$ and $bump+10$ (where the difference between the cross sections is 5 or 10 times the differences between the ones at the bump and at the dip) as shown in Fig. 6, left. This leads to a distribution of t and $d\sigma/dt$ values as a function of $\sqrt{s}$ for all characteristic points as shown in Fig. 6, middle and right [15]. We notice that the same $\sqrt{s}$ dependence can be used for all characteristic points for the $|t|$ and $d\sigma/dt$ values, namely

$$|t| = a\log(\sqrt{s}[\text{TeV}]) + b \tag{9}$$

$$(d\sigma/dt) = c\sqrt{s}\,[\text{TeV}] + d. \tag{10}$$

From the fits, we determine the $|t|$ and $d\sigma/dt$ values of all characteristic points at the Tevatron $\sqrt{s} = 1.96$ TeV as illustrated from the stars in Fig. 6.

The last step is to predict the $d\sigma/dt$ values of elastic pp interactions in the same $|t|$ bins as the D0 measurement in order to make a direct comparison. We fit the reference points extrapolated to 1.96 TeV from TOTEM measurements using a double exponential fit ($\chi^2 = 0.63$ per dof)

$$h(t) = a_1 e^{-b_1|t|^2 - c_1|t|} + d_1 e^{-f_1|t|^3 - g_1|t|^2 - h_1|t|}. \tag{11}$$

This function is chosen for fitting purposes only. The two exponential terms cross around the dip, one rapidly falling and becoming negligible in the high t-range where the other term rises above the dip. In addition, differences in normalization taken into account by adjusting TOTEM and D0 data sets to have the same cross sections at the optical point $d\sigma/dt(t=0)$.

Fig. 7 provides the final result leading to the odderon discovery [15]. We display the comparison between the D0 pp $d\sigma/dt$ and the extrapolated pp $d\sigma/dt$ TOTEM measurements in the medium $|t|$ region of the dip and the bump. The χ^2 test with six degrees of freedom yields the p-value of 0.00061, corresponding to a significance of 3.4σ.

This result can be combined with the independent evidence of the odderon found by the TOTEM Collaboration using ρ and total cross section measurements at low $|t|$ at 13 TeV in a completely different kinematical domain described in the previous section. The combined significance ranges from 5.3 to 5.7σ depending on the model. This constitutes the first experimental observation of the odderon, a major discovery at CERN and Tevatron.

Let us discuss in addition the fact that the same $\sqrt{s}$ dependence can be used for all characteristic points. It indicates that there is a new scaling [36] in the elastic data measured by TOTEM as shown in Fig. 8. If we plot $\mathrm{d}\sigma/\mathrm{d}|t| \times s^{-0.305}$ as a function of $t^{**} = s^{0.065}(|t|)^{0.72}$, we see that all data for different $\sqrt{s}$ superimpose each other as shown on Fig. 8, left. In Fig. 8, right, we display λ as a function of b, the impact parameter, for various reference $\sqrt{s}$ pairs where λ is defined as

$$\lambda = \frac{1}{\ln(s_1/s_2)} \ln\left(\frac{\mathrm{Re}\Gamma(s_1,b)}{\mathrm{Re}\Gamma(s_2,b)}\right). \tag{12}$$

where s_1 and s_2 are two center-of-mass energies square, and the Γs are the profile functions in the b-parameter space. λ is equal to 0.06 at small values of b and goes up to 0.4 at large values of $b = 2$ fm. The power exponent λ has a weak dependence on the reference energies $\sqrt{s}$ used to extract it. In particular, at $b = 0$, the value of $\lambda = 0.06$ does not depend on the reference energies and is directly related to the scaling properties. The λ values at small b are compatible with expectations from a dense object, such as a black disc [36].




Figure 6: (a) Schematic definition of the characteristic points in the TOTEM differential cross section data. (b) and (c) Characteristic points in (b) $|t|$ and (c) $d\sigma/dt$ from TOTEM measurements at 2.76, 7, 8, and 13 TeV (circles) as a function of $\sqrt{s}$ extrapolated to Tevatron center-of-mass energy (stars).




Figure 7: Comparison between the D0 elastic $p\bar{p}$ $d\sigma/dt$ cross section measurement at 1.96 TeV and the extrapolated TOTEM measurement for pp interactions

3 Conclusion

In this short report, we first described the total, inelastic and elastic cross sections measured by the TOTEM collaboration at 2.76, 7, 8 and 13 TeV, that led to a non-exponential behavior of $d\sigma/dt$. A detailed comparison between $p\bar{p}$ (1.96 TeV from D0) and pp (2.76, 7, 8, 13 TeV from TOTEM) elastic $d\sigma/dt$ data led to the fact that pp and $p\bar{p}$ cross sections differ with a significance of 3.4σ in a model-independent way and thus provides evidence that the Colorless C-odd gluonic compound i.e. the odderon is needed to explain elastic scattering at high energies. When combined with the ρ and total cross section result at 13 TeV, the significance is in the range 5.3 to 5.7σ and thus constitutes the first experimental observation of the odderon, a major discovery at CERN and Tevatron.

Many additional recent physics results of the TOTEM and CMS collaborations include the sensitivity to quartic anomalous couplings, benefitting from the fact that the LHC can be considered as a $\gamma\gamma$ collider, quasi-real photons being emitted by the protons. This leads to unprecedented sensitivities to quartic anomalous couplings such as $\gamma\gamma\gamma\gamma$ [37–44] $\gamma\gamma WW$ [45,46], $\gamma\gamma ZZ$, $\gamma\gamma\gamma Z$ [47]




Figure 8: Left: $d\sigma/dt^*$ as a function of t^{**} showing the scaling of all TOTEM elastic scattering data at $\sqrt{s} = 2.76$, 7, 8 and 13 TeV. Right: Power growth exponent λ as a function of b for various reference $\sqrt{s}$ pairs.

and $\gamma\gamma t\bar{t}$ [48, 49] and to axion-like particles at high mass [50, 51] with reach usually better by two or three orders of magnitude with respect to standard methods at the LHC without tagging the intact protons in the final state.

References

[1] G. Antchev *et al.*, TOTEM Collaboration, First measurement of elastic, inelastic and total cross-section at $\sqrt{s}$ =13 TeV by TOTEM and overview of cross-section data at LHC energies, Eur. Phys. J. C **79** (2019) 2, 103.

[2] G. Antchev *et al.*, TOTEM Collaboration, Measurement of proton-proton elastic scattering and total cross-section at $\sqrt{s}$ =7 TeV, EPL **101** (2013) 2, 21002.

[3] G. Antchev *et al.*, TOTEM Collaboration, Luminosity-independent measurements of total, elastic and inelastic cross-sections at $\sqrt{s} = 7$ TeV, EPL **101** (2013) 2, 21004.

[4] G. Antchev *et al.*, TOTEM Collaboration, Luminosity-Independent Measurement of the Proton-Proton Total Cross Section at $\sqrt{s}$ =8 TeV, Phys. Rev. Lett. **111** (2013) 1, 012001.

[5] G. Antchev *et al.*, TOTEM Collaboration, First measurement of the total proton-proton cross section at the LHC energy of $\sqrt{s}$ =7 TeV, EPL **96** (2011) 2, 21002.

[6] M. Aaboud *et al.*, ATLAS Collaboration, Measurement of the total cross section from elastic scattering in pp collisions at $\sqrt{s}$ =8 TeV with the ATLAS detector, Phys. Lett. B **761** (2016) 158.

[7] ATLAS Collaboration, Measurement of the total cross section and ρ-parameter from elastic scattering in pp collisions at $\sqrt{s}$ =13 TeV with the ATLAS detector, preprint arXiv:2207.12246 .

[8] G. Antchev *et al.*, TOTEM Collaboration, Proton-proton elastic scattering at the LHC energy of $\sqrt{s} = 7$ TeV, EPL **95**, no. 41004 (2011).

[9] G. Antchev *et al.*,TOTEM Collaboration, Evidence for non-exponential elastic pp differential cross-section at low $|t|$ and $\sqrt{s} = 8$ TeV by TOTEM, Nucl. Phys. B **899**, 527 (2015).

[10] G. Antchev *et al.*, TOTEM Collaboration, Elastic differential cross-section measurement at $\sqrt{s}$ =13 TeV by TOTEM, Eur. Phys. J. C**79**, no.10, 861 (2019).

[11] G. Antchev *et al.*, TOTEM Collaboration, Measurement of elastic pp scattering at $\sqrt{s}$ =8 TeV in the Coulomb?nuclear interference region: determination of the ρ parameter and the total cross-section, Eur. Phys. J. C **76** (2016) 12, 661.

[12] G. Antchev *et al.*, TOTEM Collaboration, First determination of the ρ parameter at $\sqrt{s}$ =13 TeV – probing the existence of a colourless C-odd three-gluon compound state, Eur. Phys. J. C**79**, no.9, 785 (2019).

[13] J.R. Cudell *et al.*, COMPETE Coll., COMPETE Collaboration, Benchmarks for the forward observables at RHIC, the Tevatron-run II and the LHC, Phys. Rev. Lett. **89** (2002) 201801.

[14] V.A. Khoze, A.D. Martin, M.G. Ryskin, Elastic and diffractive scattering at the LH, Phys. Lett. B **784** (2018) 192.

[15] D0 and TOTEM Collaborations, V.M. Abazov *et al.*, Odderon Exchange from Elastic Scattering Differences between pp and $\bar{p}p$ Data at 1.96 TeV and from pp Forward Scattering Measurements, Phys. Rev. Lett. **127** (2021) 6, 062003.

[16] C. Royon, The odderon discovery by the D0 and TOTEM collaborations, Rev. Mex. Fis. Suppl. **3** (2022) 2, 020708.

[17] K. Osterberg, Odderon observation: explanations and answers to questions/objections regarding the PRL publication, preprint ArXiv 2202.03724.

[18] G. Antchev *et al.*, TOTEM Collaboration, Performance of the TOTEM detector at the LHC, Int. J. Mod. Phys. A **28**, 1330046 (2013).

[19] G. Antchev *et al.*, TOTEM Collaboration, Elastic differential cross-section $d\sigma/dt$ at $\sqrt{s}$ =2.76 TeV and implications on the existence of a colourless C-odd three-gluon compound state, Eur. Phys. J. C**80**, no.2, 91 (2020).

[20] L. Lukaszuk, B. Nicolescu, A Possible interpretation of pp rising total cross sections, Lett. Nuovo Cim. **8**, 405 (1973).

[21] E. Martynov, B. Nicolescu, Odderon effects in the differential cross-sections at Tevatron and LHC energies, Eur. Phys. J. C **79**, no.6, 461 (2019).

[22] F. Low, Model of the bare Pomeron, Phys. Rev. D**12**,163

[23] S. Nussinov, Colored-Quark Version of Some Hadronic Puzzles, Phys. Rev. Lett. **34**, 1286 (1975).

[24] P. Gauron, L. Lukaszuk, B. Nicolescu, Consistency of the maximal odderon approach with the QFT constraints, Phys. Lett. B **294**, 298 (1992).

[25] A. Bouquet, B. Diu, E. Leader, B. Nicolescu, A possible incompatibility between the NN and $\bar{N}N$ total cross sections and the Regge pole model, Nuovo Cim. **29A**, 30 (1975).

[26] D. Joynson, E. Leader, B. Nicolescu, C. Lopez, Non-regge and hyper-regge effects in pion-nucleon charge exchange scattering at high energies, Nuovo Cim. **A 30**, 345 (1975).

[27] A.D. Donnachie, P.V. Landshoff, Multi-gluon exchange in pp elastic scattering, Phys. Lett. B **123B**, no 5, 345 (1983).

[28] G. Antchev *et al.*, TOTEM Collaboration, Characterisation of the dip-bump structure observed in proton proton elastic scattering at $\sqrt{s}$ =8 TeV, Eur. Phys. J. C **82** (2022) 3, 263.

[29] V.M. Abazov *et al.*, D0 Collaboration, The upgraded D0 detector, Nucl. Instrum. Methods A **565** 463 (2006).

[30] V.M. Abazov *et al.*, D0 Collaboration, Measurement of the differential cross section $d\sigma/dt$ in elastic $p\bar{p}$ scattering at $\sqrt{s}$ =1.96 TeV, Phys. Rev. D **86**, 012009 (2012).

[31] A. Breakstone *et al.*, Measurement of $\bar{p}p$ and pp Elastic Scattering in the Dip Region at $\sqrt{s}$ =53 GeV, Phys. Rev. Lett. **54**, 2180 (1985).

[32] S. Erhan et al., Comparison of $\bar{p}p$ and pp elastic scattering with $0.6 < t < 2.1$ GeV^2 at the CERN-ISR, Phys. Lett. B **152**, 131 (1985).

[33] D. Bernard *et al.*, UA4 Collaboration, Large-t elastic scattering at the CERN SPS collider at $\sqrt{s}$ =630 GeV, Phys. Lett. B **171**, 142 (1986).

[34] M. Bozzo *et al.*, UA4 Coll., Elastic scattering at the CERN SPS collider up to a four-momentum transfer of 1.55 GeV^2, Phys. Lett. B **155**, 197 (1985).

[35] E. Nagy *et al.*, Measurement of Proton-Proton Elastic Scattering at Large Momentum Transfer at the CERN Intersecting Storage Rings, Nucl. Phys. B **150**, 221 (1979).

[36] C. Baldenegro, C. Royon, A. Sasto, Scaling properties of elastic proton-proton scattering at LHC, Phys. Lett. B **830** (2022) 137141.

[37] E. Chapon, C. Royon, O. Kepka, Anomalous quartic W W gamma gamma, Z Z gamma gamma, and trilinear WW gamma couplings in two-photon processes at high luminosity at the LHC, Phys. Rev. D **81** (2010) 074003.

[38] O. Kepka, C. Royon, Anomalous γWW coupling in photon-induced processes using forward detectors at the LHC, Phys. Rev. D **78** (2008) 073005.

[39] S. Fichet, G. von Gersdorff, C. Royon, Measuring the Diphoton Coupling of a 750 GeV Resonance, Phys. Rev. Lett. **116** (2016) 23, 231801.

[40] S. Fichet, G. von Gersdorff, C. Royon, Scattering light by light at 750 GeV at the LHC, Phys. Rev. D **93** (2016) 7, 075031.

[41] S. Fichet, G. von Gersdorff, B. Lenzi, C. Royon, M. Saimpert, Light-by-light scattering with intact protons at the LHC: from Standard Model to New Physics, JHEP **02** (2015) 165.

[42] S. Fichet , G. von Gersdorff , O. Kepka, B. Lenzi , C. Royon, Probing new physics in diphoton production with proton tagging at the Large Hadron Collider, Phys. Rev. D **89** (2014) 114004.

[43] CMS and TOTEM Collaborations, Search for high-mass exclusive diphoton production with tagged protons, CMS-PAS-EXO-21-007, http://cms-results.web.cern.ch/cms-results/public-results/preliminary-results/EXO-21-007/index.html.

[44] A. Tumasyan *et al.*, CMS and TOTEM Collaborations, First search for exclusive diphoton production at high mass with tagged protons in proton-proton collisions at $\sqrt{s}$ =13 TeV, Phys. Rev. Lett. **129** (2022) 011801.

[45] C. Baldenegro, G. Biagi, G. Legras, C. Royon, Central exclusive production of W boson pairs in pp collisions at the LHC in hadronic and semi-leptonic final states, JHEP **12** (2020) 165.

[46] CMS and TOTEM Collaborations, Search for exclusive $\gamma\gamma \to WW$ and $\gamma\gamma \to ZZ$ production in final states with jets and forward protons, CMS-PAS-SMP-21-014, http://cms-results.web.cern.ch/cms-results/public-results/preliminary-results/SMP-21-014/index.html.

[47] C. Baldenegro, S. Fichet, G. von Gersdorff, C. Royon, Probing the anomalous $\gamma\gamma\gamma Z$ coupling at the LHC with proton tagging, JHEP **06** (2017) 142.

[48] C. Baldenegro, A. Bellora, S. Fichet, G. von Gersdorff, M. Pitt, C. Royon, Searching for anomalous top quark interactions with proton tagging and timing detectors at the LHC, JHEP **08** (2022) 021.

[49] CMS and TOTEM Collaborations, Search for central exclusive production of top quark pairs in proton-proton collisions at $\sqrt{s}$ =g13 TeV with tagged protons, CMS-PAS-TOP-21-007, http://cms-results.web.cern.ch/cms-results/public-results/preliminary-results/TOP-21-007/index.html

[50] C. Baldenegro, S. Fichet, G. von Gersdorff, C. Royon, Searching for axion-like particles with proton tagging at the LHC, JHEP **06** (2018) 131.

[51] C. Baldenegro, S. Hassani, C. Royon, L. Schoeffel, Extending the constraint for axion-like particles as resonances at the LHC and laser beam experiments, Phys. Lett.B **795** (2019) 339.