

Proof of an open problem on the Sombor index

Hechao Liu

School of Mathematical Sciences, South China Normal University, Guangzhou, 510631, P. R. China, e-mail: hechaoliu@m.scnu.edu.cn

Received 18 August 2022

Abstract The Sombor index is one of the geometry-based descriptors, which was defined as

$$SO(G) = \sum_{uv \in E(G)} \sqrt{d^2(u) + d^2(v)},$$

where $d(u)$ (resp. $d(v)$) denotes the degree of vertex u (resp. v) in G .

In this note, we determine the maximum and minimum graphs with respect to the Sombor index among the set of graphs with vertex connectivity (resp. edge connectivity) at most k , which solves an open problem on the Sombor index proposed by Hayat and Rehman [On Sombor index of graphs with a given number of cut-vertices, *MATCH Commun. Math. Comput. Chem.* **89** (2023) 437–450]. For some of the conclusions of the above paper, we give some counterexamples. At last, we give the QSPR analysis with regression modeling and Sombor index.

Keywords Sombor index, vertex connectivity, edge connectivity.

Mathematics Subject Classification: 05C07, 05C09, 05C92

1 Introduction

Let G be a simple undirected connected graph with vertex set $V(G)$ and edge set $E(G)$. Let $N_G(u)$ (or $N(u)$ for short) be the neighbor of vertex u in G , then the degree of vertex u is $d(u) = |N(u)|$.

Let $uv \notin E(G)$ (resp. $uv \in E(G)$), then $G + uv$ (resp. $G - uv$) denotes the graphs obtained from G by adding (resp. deleting) the edge uv . Denote by C_n , K_n , S_n , P_n , the cycle, complete graph, star graph, path with order n , respectively. In this paper, all notations and terminologies used but not defined can refer to Bondy and Murty [1].

One of the geometry-based indices, Sombor index [5] was defined as

$$SO(G) = \sum_{uv \in E(G)} \sqrt{d^2(u) + d^2(v)},$$

where $d(u)$ (resp. $d(v)$) denotes the degree of vertex u (resp. v) in G . One can refer to [2–4, 6, 7, 9] for more details about Sombor index.

We call a graph G is k -connected if $G \cong K_{k+1}$, or $|V(G)| \geq k+2$ and G has no $(k-1)$ -vertex cut. We call a graph G is k -edge-connected ($k \geq 1$) if $|V(G)| \geq 2$ and G has no $(k-1)$ -edge cut. Denote by $\kappa(G) = \max\{k \mid G \text{ is } k\text{-connected}\}$ the connectivity of connected graph G . Denote by $\kappa'(G) = \max\{k \mid G \text{ is } k\text{-edge-connected}\}$ the edge connectivity of connected graph G . Then $\kappa(G) \leq \kappa'(G) \leq n-1$ and $\kappa(G) = n-1 \Leftrightarrow \kappa'(G) = n-1 \Leftrightarrow G \cong K_n$.

Let \mathbb{V}_n^k (resp. \mathbb{E}_n^k) be the set of graphs with n vertices and $\kappa(G) \leq k \leq n-1$ (resp. $\kappa'(G) \leq k \leq n-1$). Suppose G_1 and G_2 are two disjoint graphs. Let $G_1 \vee G_2$ be the graph obtained from $G_1 \cup G_2$ by adding edges between any vertex of G_1 and any vertex of G_2 . Let $K_n^k \triangleq K_1 \vee K_k \vee K_{n-k-1}$.

The remainder of this paper is organized as follow. In Section 2, we determine the maximum and minimum graphs with respect to the Sombor index among the set of graphs with vertex connectivity (resp. edge connectivity) at most k . In Section 3, we give some counterexamples about some results of [8]. In Section 3, we give the QSPR analysis with regression modeling and Sombor index.

2 Extremal graphs with connectivity at most k

In one recently published paper [8], Hayat and Rehman proposed the open problem that considering the maximum and minimum graphs with respect to the Sombor index among the set of graphs with vertex connectivity (resp. edge connectivity) at most k . In the section, we completely solve the problem.

Lemma 2.1 [5] *Let T be a tree with n vertices. Then*

$$SO(P_n) \leq SO(T) \leq SO(S_n)$$

with equality if and only if $T \cong P_n$ or $T \cong S_n$.

Lemma 2.2 [10] *Let $x > a \geq 1, y > 0$, $f(x, y) = \sqrt{x^2 + y^2} - \sqrt{(x-a)^2 + y^2}$. Then $f(x, y)$ is strictly increasing with x , strictly decreasing with y .*

Lemma 2.3 If $u, v \in V(G)$ and $uv \notin E(G)$, then $SO(G) < SO(G + uv)$.




Figure 1: The graphs G and G_1 of Lemma 2.4.

Lemma 2.4 Suppose G is a connected graph, $u, v \in V(G)$, and $u_1, u_2, \dots, u_t \in N_G(u) \setminus \{N_G(v) \cap N_G(u)\}$ for $1 \leq t \leq d_u$ and $u_1, u_2, \dots, u_t \notin P_{uv}$ where P_{uv} is a path from u to v . Let $G_1 = G - \{uu_1, uu_2, \dots, uu_t\} + \{vu_1, vu_2, \dots, vu_t\}$ (see Figure 1). If $d_u \leq d_v$ and $uv \notin E(G)$, then $SO(G_1) > SO(G)$.

Proof. For convenience, let $M = N_G(v) \cap N_G(u)$, $M_1 = N_G(u) \setminus \{N_G(v) \cap N_G(u)\}$, $M_2 = N_G(v) \setminus \{N_G(v) \cap N_G(u)\}$. By the definition of Sombor index and the structure of graphs G and G_1 , we have

$$\begin{aligned}
 SO(G_1) - SO(G) &= \sum_{v_i \in M_2} \sqrt{(d_v + t)^2 + d_{v_i}^2} - \sum_{v_i \in M_2} \sqrt{d_v^2 + d_{v_i}^2} \\
 &\quad + \sum_{v_i \in M_1} \sqrt{(d_v + t)^2 + d_{v_i}^2} - \sum_{v_i \in M_1} \sqrt{d_u^2 + d_{v_i}^2} \\
 &\quad + \left(\sum_{v_i \in M} \sqrt{(d_v + t)^2 + d_{v_i}^2} - \sum_{v_i \in M} \sqrt{d_v^2 + d_{v_i}^2} \right) \\
 &\quad - \left(\sum_{v_i \in M} \sqrt{d_u^2 + d_{v_i}^2} - \sum_{v_i \in M} \sqrt{(d_u - t)^2 + d_{v_i}^2} \right) \\
 &> 0. \quad (\text{by Lemma 2.2 and } d_u \leq d_v)
 \end{aligned}$$

This completes the proof. ■

Recalling that $G_1 \vee G_2$ is the graph obtained from $G_1 \cup G_2$ by adding edges between any vertex of G_1 and any vertex of G_2 .

Theorem 2.5 Suppose that $G(i, n - k - i) = K_i \vee H_k \vee K_{n-k-i}$ is the graph with n vertices and H_k is a graph with $k \geq 1$ vertices. Then $SO(G(i, n - k - i)) < SO(G(1, n - k - 1))$ for $2 \leq i \leq \frac{n-k}{2}$.

Proof. Suppose that $V(K_i) = \{u_1, u_2, \dots, u_i\}$, $V(K_{n-k-i}) = \{v_1, u_2, \dots, u_{n-k-i}\}$. Since $2 \leq i \leq \frac{n-k}{2}$, then $d_{u_1} \leq d_{v_1}$. By Lemma 2.4, we have $SO(G_1) > SO(G(i, n-k-i))$, where $G_1 = G - \{u_1u_2, u_1u_3, \dots, u_1u_i\} + \{v_1u_2, v_1u_3, \dots, v_1u_i\}$. We can also know that $G(1, n-k-1) = G_1 + \{v_2u_2, v_2u_3, \dots, v_2u_i\} + \{v_3u_2, v_3u_3, \dots, v_3u_i\} + \dots + \{v_{n-k-i}u_2, v_{n-k-i}u_3, \dots, v_{n-k-i}u_i\}$. Then by Lemma 2.3, we have $SO(G_1) < SO(G(1, n-k-1))$. Thus $SO(G(i, n-k-i)) < SO(G(1, n-k-1))$ for $2 \leq i \leq \frac{n-k}{2}$.

This completes the proof. ■

Theorem 2.6 *Let $G \in \mathbb{V}_n^k$. Then we have $SO(G) \leq k\sqrt{k^2 + (n-1)^2} + k(n-k-1)\sqrt{(n-1)^2 + (n-2)^2} + \frac{\sqrt{2}}{2}k(k-1)(n-1) + \frac{\sqrt{2}}{2}(n-k-1)(n-k-2)(n-2)$, with equality if and only if $G \cong K_n^k$.*

Proof. If $k = n-1$, then $K_n^{n-1} \cong K_n \in \mathbb{V}_n^{n-1}$, thus the $SO(G) \leq SO(K_n^k)$ for $k = n-1$. Next we only consider the case $1 \leq k \leq n-2$.

Suppose that $SO(G) \leq SO(G_1)$ for any $G \in \mathbb{V}_n^k$ ($1 \leq k \leq n-2$). Thus $G_1 \not\cong K_n$, then G_1 has a k -vertex cut that is $W = \{u_1, u_2, \dots, u_k\}$. Denote by $\omega(G)$ the number of connected components in graph G .

We first proof that $\omega(G_1 - W) = 2$. Otherwise, $\omega(G_1 - W) \geq 3$. We suppose that $G_1 - W = \{H_1, H_2, \dots, H_l\}$ ($l \geq 3$). Let $u \in H_i$ and $v \in H_j$ where $i, j \in \{1, 2, \dots, l\}$ and $i \neq j$. We find that W is still a k -vertex cut of $G_1 + uv$, i.e., $G_1 + uv \in \mathbb{V}_n^k$. Since $SO(G_1 + uv) > SO(G_1)$, which is a contradiction with that $SO(G) \leq SO(G_1)$ for any $G \in \mathbb{V}_n^k$. Thus we have $\omega(G_1 - W) = 2$.

We suppose that $G_1 - W = \{H_1, H_2\}$.

Next we proof that $G_1[V(H_1) \cup W]$ and $G_1[V(H_2) \cup W]$ are all cliques. Otherwise, $G_1[V(H_1) \cup W]$ is not a clique. Then there exist vertices $u, v \in V(H_1) \cup W$, such that $uv \notin E(G_1[V(H_1) \cup W])$. Since $SO(G_1 + uv) > SO(G_1)$ and $G_1 + uv \in \mathbb{V}_n^k$, which is a contradiction with that $SO(G) \leq SO(G_1)$ for any $G \in \mathbb{V}_n^k$. Thus $G_1[V(H_1) \cup W]$ is a clique. Similarly, we also have that $G_1[V(H_2) \cup W]$ is a clique.

Since $G_1[V(H_1) \cup W]$ and $G_1[V(H_2) \cup W]$ are all cliques, then H_1 and H_2 are all cliques. For convenience, we denote H_1 (resp. H_2) as K_{n_1} (resp. K_{n_2}).

At last we proof that $n_1 = 1$ or $n_2 = 1$. Otherwise, we have $n_1 \geq 2$ and $n_2 \geq 2$. Suppose that $G_2 = K_1 \vee G_1[W] \vee K_{n-k-1}$, it is obvious that $G_2 \in \mathbb{V}_n^k$. By the conclusion of Lemma 2.5, we know that $SO(G_1) < SO(G_2)$, which is a contradiction with that $SO(G) \leq SO(G_1)$ for any $G \in \mathbb{V}_n^k$. Thus $n_1 = 1$ or $n_2 = 1$.

By a simple calculation, we know that $SO(G) \leq k\sqrt{k^2 + (n-1)^2} + k(n-k-1)\sqrt{(n-1)^2 + (n-2)^2} + \frac{\sqrt{2}}{2}k(k-1)(n-1) + \frac{\sqrt{2}}{2}(n-k-1)(n-k-2)(n-2)$, with equality if and only if $G \cong K_n^k$.

This completes the proof. ■

Corollary 2.7 *Let $G \in \mathbb{E}_n^k$. Then we have $SO(G) \leq k\sqrt{k^2 + (n-1)^2} + k(n-k-1)\sqrt{(n-1)^2 + (n-2)^2} + \frac{\sqrt{2}}{2}k(k-1)(n-1) + \frac{\sqrt{2}}{2}(n-k-1)(n-k-2)(n-2)$, with equality if and only if $G \cong K_n^k$.*

Proof. Since $K_n^k \in \mathbb{E}_n^k \subseteq \mathbb{V}_n^k$, then by Theorem 2.6, the conclusion holds. ■

Theorem 2.8 *Let $G \in \mathbb{V}_n^k$ ($k \geq 1$) be a connected graph. Then we have $SO(G) \geq 2\sqrt{2}(n-3) + 2\sqrt{5}$, with equality if and only if $G \cong P_n$.*

Proof. Note that for $e = uv \in E(G)$ and $G \in \mathbb{V}_n^k$, then $G - uv \in \mathbb{V}_n^k$. Thus if $G \in \mathbb{V}_n^k$ be a connected graph with minimum Sombor index, then G must be a trees. Then by Lemma 2.1, we have P_n is the minimum connected graph in \mathbb{V}_n^k with respect to Sombor index. ■

Corollary 2.9 *Let $G \in \mathbb{E}_n^k$ ($k \geq 1$) be a connected graph. Then we have $SO(G) \geq 2\sqrt{2}(n-3) + 2\sqrt{5}$, with equality if and only if $G \cong P_n$.*

Proof. Since $P_n \in \mathbb{E}_n^k \subseteq \mathbb{V}_n^k$, then by Theorem 2.8, the conclusion holds. ■

3 Counterexamples

In one recently published paper [8], there are some flaws in their Lemmas. We give some counterexamples about their results.




Figure 2: A counterexamples of Lemma 3 of [8].

Lemma 3.1 [8] *Let Γ_α be the α -switched graph of a graph Γ with n vertices and k cut vertices (see Figure 1 of [8]). Then $SO(\Gamma_\alpha) < SO(\Gamma)$.*

The conclusion of Lemma 3.1 of [8] is not entirely correct. A counterexamples of Lemma 3 of [8] can see Figure 2. If $d_{x_1} = 8$ and $d_{y_1} = 8$, then $SO(\Gamma) \approx 135.716 < 136.169 \approx SO(\Gamma_\alpha)$, which is a contradiction with the conclusion of Lemma 3.1 of [8]. In fact, if $d_{x_1} \geq 8$ and $d_{y_1} \geq 8$, then the conclusion of Lemma 3.1 of [8] is wrong.

Let $x, y \geq 2$ and $f(x, y) = \sqrt{x^2 + y^2} - \sqrt{x^2 + 2^2} - \sqrt{y^2 + 2^2}$. Then $\frac{\partial f}{\partial x} = \frac{x}{\sqrt{x^2 + y^2}} - \frac{x}{\sqrt{x^2 + 2^2}} < 0$. Thus $f(x, y)$ is monotonically decreasing on x or y . However, in the proof of Lemma 3.1 of [8], the authors think that $f(x, y)$ is monotonically increasing on x or y .

The conclusions of Lemma 4 and Lemma 5 of [8] is also not entirely correct. We can also find counterexamples. The main problem is that the author misunderstood the monotonicity of the above function $f(x, y)$.

4 QSPR analysis with regression modeling and Sombor index

The experimental values of enthalpie of combustion (resp. enthalpie of formation of liquid, enthalpie of sublimation, enthalpie of vaporization) of 19 monocarboxylic acids of Table 1 are taken from [11].

The regression modeling between Sombor index and enthalpie of combustion (ΔH_c^o), enthalpie of formation of liquid (ΔH_f^o), enthalpie of sublimation (ΔH_{sub}^o), enthalpie of vaporization (ΔH_{vap}^o) of 19 monocarboxylic acids are, respectively,

$$\Delta H_c^o = 229.7 \times SO(G) - 1263, \quad \Delta H_f^o = 10.65 \times SO(G) + 369.2,$$

$$\Delta H_{sub}^o = 1.212 \times SO(G) + 36.41, \quad \Delta H_{vap}^o = 2.559 \times SO(G) + 21.83.$$

By comparison, we find that the Sombor index exerts a better predictive capability than other vertex degree-based indices.

Table 1: Experimental values of enthalpie of combustion ($\Delta H_c^\circ KJ/mol$), enthalpie of formation of liquid ($\Delta H_f^\circ KJ/mol$), enthalpie of sublimation ($\Delta H_{sub}^\circ KJ/mol$), enthalpie of vaporization ($\Delta H_{vap}^\circ KJ/mol$) and Sombor index of 19 monocarboxylic acids

Compounds	$\Delta H_c^\circ KJ/mol$	$\Delta H_f^\circ KJ/mol$	$\Delta H_{sub}^\circ KJ/mol$	$\Delta H_{vap}^\circ KJ/mol$	SO
<i>Aceticacid</i>	875.16	483.5	46.3	49.7	9.48683
<i>Propanoicacid</i>	1527.3	510.8	50.0	56.1	12.1662
<i>Butanoicacid</i>	2183.5	533.9	54.9	62.9	14.9946
<i>Pentanoicacid</i>	2837.8	558.9	58.2	69.0	17.8230
<i>Hexanoicacid</i>	3494.3	581.8	63.0	75.0	20.6515
<i>Heptanoicacid</i>	4146.9	608.5	64.8	81.7	23.4799
<i>Octanoicacid</i>	4799.9	634.8	69.4	86.9	26.3083
<i>Nonanoicacid</i>	5456.1	658.0	72.3	93.6	29.1367
<i>Decanoicacid</i>	6079.3	713.7	76.3	100.8	31.9652
<i>Undecanoicacid</i>	6736.5	736.2	78.9	106.7	34.7936
<i>Dodecanoicacid</i>	7333.0	775.1	82.2	115.9	37.6220
<i>Tridecanoicacid</i>	8024.2	807.2	84.9	121.2	40.4504
<i>Tetradecanoicacid</i>	8676.7	834.1	87.7	130.2	43.2789
<i>Pentadecanoicacid</i>	9327.7	862.4	91.4	136.5	46.1073
<i>Hexadecanoicacid</i>	9977.2	892.2	94.5	144.3	48.9357
<i>Heptadecanoicacid</i>	10624.4	924.4	100.7	159.6	51.7642
<i>Octadecanoicacid</i>	11280.1	947.2	102.8	164.7	54.5926
<i>Nonadecanoicacid</i>	11923.4	984.1	105.0	172.9	57.4210
<i>Eicosanoicacid</i>	12574.2	1012.6	109.9	179.2	60.2494

Table 2: The R^2 and $RMSE$ of regression modeling between Sombor index and enthalpie of combustion ($\Delta H_c^\circ KJ/mol$), enthalpie of formation of liquid ($\Delta H_f^\circ KJ/mol$), enthalpie of sublimation ($\Delta H_{sub}^\circ KJ/mol$), enthalpie of vaporization ($\Delta H_{vap}^\circ KJ/mol$) of 19 monocarboxylic acids.

Physico-chemical property	R^2	$RMSE$
enthalpie of combustion	0.99998	17.987
enthalpie of formation of liquid	0.99737	8.9567
enthalpie of vaporization	0.99745	1.0034
enthalpie of vaporization	0.99355	3.2771

References

- [1] J. A. Bondy, U. S. R. Murty, *Graph Theory*, Springer, New York, 2008.
- [2] R. Cruz, I. Gutman, J. Rada, Sombor index of chemical graphs, *Appl. Math. Comput.* **399** (2021) #126018.
- [3] H. Chen, W. Li, J. Wang, Extremal values on the Sombor index of trees, *MATCH Commun. Math. Comput. Chem.* **87** (2022) 23–49.
- [4] H. Deng, Z. Tang, R. Wu, Molecular trees with extremal values of Sombor indices, *Int. J. Quantum. Chem.* **121** (2021) #e26622.

- [5] I. Gutman, Geometric approach to degree-based topological indices: Sombor indices, *MATCH Commun. Math. Comput. Chem.* **86** (2021) 11–16.
- [6] I. Gutman, Sombor index-one year later, *Bull. Acad. Serb. Sci. Arts (Cl. Sci. Math. Natur.* **153** (2020) 43–55.
- [7] I. Gutman, Some basic properties of Sombor indices, *Open J. Discr. Appl. Math.* **4** (2021) 1–3.
- [8] S. Hayat, A. Rehman, On Sombor index of graphs with a given number of cut-vertices, *MATCH Commun. Math. Comput. Chem.* **89** (2023) 437–450
- [9] S. Li, Z. Wang, M. Zhang, On the extremal Sombor index of trees with a given diameter, *Appl. Math. Comput.* **416** (2022) #126731.
- [10] H. Liu, L. You, Y. Huang, Z. Tang, On extremal Sombor indices of chemical graphs, and beyond, *MATCH Commun. Math. Comput. Chem.* **89** (2023) 415–436.
- [11] F. Shafiei, Relationship between topological indices and thermodynamic properties and of the monocarboxylic acids applications in QSPR, *Iran. J. Math. Chem.* **6** (2015) 15–28.