

The rank of the walk matrix of the extended Dynkin graph \tilde{D}_n

Sunyo Moon* and Seungkook Park†

Abstract

In this paper, we provide an explicit formula for the rank of the walk matrix of the extended Dynkin graph \tilde{D}_n .

1 Introduction

Let G be a simple graph with n vertices and let A be an adjacency matrix of G . The *walk matrix* $W(G)$ of G is defined by

$$W(G) = \begin{bmatrix} e_n & Ae_n & \cdots & A^{n-1}e_n \end{bmatrix},$$

where e_n is the all-one vector of length n . The (i, j) -entry of the walk matrix $W(G)$ counts the number of walks in G of length $j - 1$ from the vertex i . A tree obtained from the path of order $n - 1$ by adding a pendant edge at the second vertex is called a *Dynkin graph* D_n (See Figure 1) and by adding a pendant edge to the second to last vertex of D_n , we get the *extended Dynkin graph* \tilde{D}_n (See Figure 2), where $n \geq 4$.




Figure 1: Dynkin graph D_n

Dynkin graphs and extended Dynkin graphs are widely used in the study of simple Lie algebras [3, 6, 7], representation theory [1, 5, 8, 11, 12, 14, 16], and spectral theory [2, 4, 13].

*Research Institute of Natural Sciences, Hanyang University, Seoul, Republic of Korea (symoon89@hanyang.ac.kr)

†Department of Mathematics and Research Institute of Natural Sciences, Sookmyung Women's University, Seoul, Republic of Korea (skpark@sookmyung.ac.kr)




Figure 2: Extended Dynkin graph \tilde{D}_n

In [15], Wang, Wang and Guo gave a formula for the rank of the walk matrix of the Dynkin graph D_n ($n \geq 4$). In this paper, we find a formula for the rank of the walk matrix of the extended Dynkin graph \tilde{D}_n ($n \geq 4$) which is $\text{rank } W(\tilde{D}_n) = \lfloor \frac{n}{2} \rfloor$.

For a graph G with n vertices, an eigenvalue of the adjacency matrix of G is said to be *main* if its corresponding eigenvector is not orthogonal to e_n . In [10], Hagos showed that the rank of the walk matrix of a graph G is equal to the number of its main eigenvalues. Thus our formula also provides the number of the main eigenvalues of the walk matrix of the extended Dynkin graph.

2 Rank of the extended Dynkin graph

Let \tilde{D}_n ($n \geq 4$) be the extended Dynkin graph with $n + 1$ vertices. We label the vertices of \tilde{D}_n as shown in Figure 2. Then the partition

$$\Pi = \{\{1, 2\}, \{3\}, \dots, \{n-1\}, \{n, n+1\}\}$$

of $V(\tilde{D}_n)$ is equitable. The $(n+1) \times (n-1)$ characteristic matrix P of Π and the $(n-1) \times (n-1)$ divisor matrix B of Π are

$$P = \begin{bmatrix} 1 & 0 & 0 & & \\ 1 & 0 & 0 & & \\ 0 & 1 & 0 & & \\ & \ddots & \ddots & \ddots & \\ & & 0 & 1 & 0 \\ & & 0 & 0 & 1 \\ & & 0 & 0 & 1 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 0 & 1 & & & \\ 2 & 0 & 1 & & \\ & 1 & 0 & 1 & \\ & & \ddots & \ddots & \ddots \\ & & & 1 & 0 & 1 \\ & & & & 1 & 0 & 2 \\ & & & & & 1 & 0 \end{bmatrix},$$

respectively.

For an $n \times n$ matrix M , the *Smith normal form* of M is the diagonal matrix $\text{diag}(d_1, d_2, \dots, d_r, 0, \dots, 0)$, where $r = \text{rank}(M)$ and each d_i divides d_{i+1} for $i = 1, \dots, r-1$. Two integral matrices M_1 and M_2 are *integrally equivalent* if there are unimodular matrices P and Q such that $M_2 = PM_1Q$, that is, M_2 can be obtained from M_1 by a finite sequence of elementary row

operations and column operations. Two matrices are integrally equivalent if and only if they have the same Smith normal form (See [9] for details).

Let $\hat{W}(\tilde{D}_n)$ be the $(n-1) \times (n-1)$ matrix obtained by removing the first and last rows and the last two columns from $W(\tilde{D}_n)$. Then we define an $(n+1) \times (n+1)$ matrix $W'(\tilde{D}_n)$ in which $\hat{W}(\tilde{D}_n)$ is a submatrix as follows:

$$W'(\tilde{D}_n) = \begin{bmatrix} 0 & 0 & 0 \\ \hat{W}(\tilde{D}_n) & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

For an $n \times n$ matrix M , the *walk matrix* $W(M)$ of M is defined by

$$W(M) = \begin{bmatrix} e_n & Me_n & \cdots & M^{n-1}e_n \end{bmatrix}.$$

The following lemma provides a method for computing the rank of a walk matrix.

Lemma 2.1. [15, Lemma 2] *Let M be a real $n \times n$ matrix which is diagonalizable over the real field \mathbb{R} . Let v_1, v_2, \dots, v_n be n linearly independent eigenvectors of M^T corresponding to eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$, respectively. Then we have*

$$\det W(M) = \frac{\prod_{1 \leq k < j \leq n} (\lambda_j - \lambda_k) \prod_{j=1}^n e_n^T v_j}{\det[v_1, v_2, \dots, v_n]}.$$

Moreover, if $\lambda_1, \lambda_2, \dots, \lambda_n$ are pairwise different, then

$$\text{rank } W(M) = |\{j : 1 \leq j \leq n \text{ and } e_n^T v_j \neq 0\}|.$$

We outline the steps for establishing our formula for the rank of the extended Dynkin graph \tilde{D}_n .

- (1) Using Lemma 2.1, we find a formula for the rank of the walk matrix of B .
- (2) We show that $\hat{W}(\tilde{D}_n) = W(B)$, which implies $\text{rank } \hat{W}(\tilde{D}_n) = \text{rank } W(B)$.
- (3) We prove that the Smith normal form of $W(\tilde{D}_n)$ and $W'(\tilde{D}_n)$ are identical.
- (4) We conclude that

$$\text{rank } W(\tilde{D}_n) = \text{rank } W'(\tilde{D}_n) = \text{rank } \hat{W}(\tilde{D}_n) = \text{rank } W(B).$$

We start by finding the eigenvalues of B^T and the corresponding eigenvectors.

Proposition 2.2. Let $\lambda_k = 2 \cos \frac{k\pi}{n-2}$ for $k = 0, \dots, n-3$ and $\lambda_{n-2} = -2$. Let $v_k = \begin{bmatrix} 1 & \cos \frac{k\pi}{n-2} & \cos \frac{2k\pi}{n-2} & \dots & \cos \frac{(n-3)k\pi}{n-2} & \cos k\pi \end{bmatrix}^T$ for $k = 0, \dots, n-3$ and $v_{n-2} = \begin{bmatrix} 1 & -1 & 1 & -1 & \dots \end{bmatrix}^T$. Then v_k is an eigenvector of B^T corresponding to the eigenvalue λ_k , that is, $B^T v_k = \lambda_k v_k$ for $k = 0, \dots, n-2$.

Proof. Since $v_0 = e_{n-1}$ and $\lambda_0 = 2$, $B^T v_0 = \lambda_0 v_0$. It is easily check that $B^T v_{n-2} = \lambda_{n-2} v_{n-2}$. Now, we show that $B^T v_k = \lambda_k v_k$ for $k = 1, \dots, n-3$. Let $\theta = k\pi/(n-2)$. Then

$$B^T v_k = \begin{bmatrix} 0 & 2 & & & & \\ 1 & 0 & 1 & & & \\ & 1 & 0 & 1 & & \\ & & \ddots & \ddots & \ddots & \\ & & & 1 & 0 & 1 \\ & & & & 2 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ \cos \theta \\ \cos 2\theta \\ \vdots \\ \cos(n-3)\theta \\ \cos k\pi \end{bmatrix} = \begin{bmatrix} 2 \cos \theta \\ 1 + \cos 2\theta \\ \cos \theta + \cos 3\theta \\ \vdots \\ \cos(n-4)\theta + \cos k\pi \\ 2 \cos(n-3)\theta \end{bmatrix}.$$

Since $\lambda_k = 2 \cos \theta$ and $\cos m\theta + \cos(m+2)\theta = 2 \cos \theta \cos(m+1)\theta$, we obtain

$$\cos m\theta + \cos(m+2)\theta = \lambda_k \cos(m+1)\theta$$

for $m = 0, \dots, n-4$. We note that

$$2 \cos(n-3)\theta = 2(\cos(n-2)\theta \cos \theta + \sin(n-2)\theta \sin \theta).$$

Since $\sin(n-2)\theta = \sin k\pi = 0$ for all $k = 1, \dots, n-3$, we have

$$2 \cos(n-3)\theta = 2 \cos \theta \cos(n-2)\theta = \lambda_k \cos k\pi.$$

Hence $B^T v_k = \lambda_k v_k$ for all $k = 0, \dots, n-2$. □

The following lemma will be used in the proof of Lemma 2.4.

Lemma 2.3. [15, Lemma 7]

$$\sum_{k=1}^n \cos(ak+b)x = \frac{1}{2 \sin \frac{1}{2}ax} \left(\sin \left(\left(n + \frac{1}{2} \right) a + b \right) x - \sin \left(\frac{1}{2} a + b \right) x \right)$$

for $\sin \frac{1}{2}ax \neq 0$.

Lemma 2.4. Let $v_k = \begin{bmatrix} 1 & \cos \frac{k\pi}{n-2} & \cos \frac{2k\pi}{n-2} & \dots & \cos \frac{(n-3)k\pi}{n-2} & \cos k\pi \end{bmatrix}^T$ for $k = 0, \dots, n-3$ and $v_{n-2} = \begin{bmatrix} 1 & -1 & 1 & -1 & \dots \end{bmatrix}^T$. Then

$$e_{n-1}^T v_0 = n-1 \quad \text{and} \quad e_{n-1}^T v_k = \begin{cases} 1, & \text{if } k \text{ is even} \\ 0, & \text{if } k \text{ is odd} \end{cases}$$

for $k = 1, \dots, n-2$.

Proof. It is straightforward to check that

$$e_{n-1}^T v_0 = n-1 \quad \text{and} \quad e_{n-1}^T v_{n-2} = \begin{cases} 1, & \text{if } n \text{ is even} \\ 0, & \text{if } n \text{ is odd.} \end{cases}$$

Now, we give the proof for $k = 1, \dots, n-3$. Let $\theta = k\pi/(n-2)$. Then

$$e_{n-1}^T v_k = 1 + \cos k\pi + \sum_{m=1}^{n-3} \cos m\theta.$$

By Lemma 2.3, we have

$$\begin{aligned} \sum_{m=1}^{n-3} \cos m\theta &= \frac{1}{2 \sin \frac{1}{2}\theta} \left(\sin \left(n - \frac{5}{2} \right) \theta - \sin \frac{1}{2} \theta \right) \\ &= \frac{1}{\sin \frac{1}{2}\theta} \cos \frac{k}{2} \pi \sin \frac{n-3}{2} \theta. \end{aligned}$$

If k is odd, then $\cos k\pi = -1$ and $\cos \frac{k}{2}\pi = 0$ and hence $e_{n-1}^T v_k = 0$. Suppose that k is even. Then $\cos k\pi = 1$ and $\cos \frac{k}{2}\pi = \pm 1$. Since

$$\begin{aligned} \sin \frac{n-3}{2} \theta &= \sin \left(\frac{n-2}{2} \theta - \frac{1}{2} \theta \right) \\ &= \sin \frac{n-2}{2} \theta \cos \frac{1}{2} \theta - \sin \frac{1}{2} \theta \cos \frac{n-2}{2} \theta \\ &= \sin \frac{k}{2} \pi \cos \frac{1}{2} \theta - \sin \frac{1}{2} \theta \cos \frac{k}{2} \pi \\ &= -\sin \frac{1}{2} \theta \cos \frac{k}{2} \pi, \end{aligned}$$

we obtain

$$\sum_{m=1}^{n-3} \cos m\theta = -\cos^2 \frac{k}{2} \pi = -1.$$

Hence $e_{n-1}^T v_k = 1$. □

We can now formulate the rank of the walk matrix of B .

Theorem 2.5. *The rank of the walk matrix of B is $\lfloor \frac{n}{2} \rfloor$.*

Proof. Let v_0, \dots, v_{n-2} be the eigenvectors of B^T in Proposition 2.2. Then there are $1 + \lfloor \frac{n-2}{2} \rfloor = \lfloor \frac{n}{2} \rfloor$ eigenvectors of B^T such that $e_{n-1} v_k \neq 0$ by Lemma 2.4. We note that all

eigenvalues of B^T are pairwise different, since the cosine function is monotonically decreasing on $[0, \pi]$. By Lemma 2.1, we have

$$\text{rank } W(B) = \left\lfloor \frac{n}{2} \right\rfloor.$$

□

Lemma 2.6. *Let \tilde{D}_n be the extended Dynkin graph with $n + 1$ vertices. Then $\hat{W}(\tilde{D}_n) = W(B)$.*

Proof. Since $AP = PB$, $A^k P = PB^k$ for all $k \geq 0$. Then we have $A^k e_{n+1} = PB^k e_{n-1}$ for all $k \geq 0$, because $Pe_{n-1} = e_{n+1}$. It follows that

$$\begin{bmatrix} e_{n+1} & Ae_{n+1} & \cdots & A^{n-2}e_{n+1} \end{bmatrix} = P \begin{bmatrix} e_{n-1} & Be_{n-1} & \cdots & B^{n-2}e_{n-1} \end{bmatrix}.$$

If we delete the first and last rows from $\begin{bmatrix} e_{n+1} & Ae_{n+1} & \cdots & A^{n-2}e_{n+1} \end{bmatrix}$ and P , we get $\hat{W}(\tilde{D}_n)$ and the identity matrix, respectively. Hence $\hat{W}(\tilde{D}_n) = W(B)$. □

Remark 2.7. By Theorem 2.5 and Lemma 2.6, we have

$$\text{rank } \hat{W}(\tilde{D}_n) = \text{rank } W(B) = \left\lfloor \frac{n}{2} \right\rfloor.$$

In the next proposition, we show that $W(\tilde{D}_n)$ and $W'(\tilde{D}_n)$ are integrally equivalent and hence proving that they have the same Smith normal form.

Proposition 2.8. *Let \tilde{D}_n be the extended Dynkin graph with $n + 1$ vertices. Then $W(\tilde{D}_n)$ and $W'(\tilde{D}_n)$ have the same Smith normal form.*

Proof. Let $r = \text{rank } W(\tilde{D}_n)$. Then the first r columns of $W(\tilde{D}_n)$ are linearly independent. Note that the first two rows of $W(\tilde{D}_n)$ are the same and the last two rows of $W(\tilde{D}_n)$ are the same. Since $r \leq n - 1$, the last two columns of $W(\tilde{D}_n)$ can be written as linear combinations of $e_{n+1}, Ae_{n+1}, \dots, A^{r-1}e_{n+1}$ with integer coefficients. Hence we obtain $W'(\tilde{D}_n)$ by using the elementary row and column operations on $W(\tilde{D}_n)$. Therefore, $W(\tilde{D}_n)$ and $W'(\tilde{D}_n)$ have the same Smith normal form. □

Corollary 2.9. *Let \tilde{D}_n be the extended Dynkin graph with $n + 1$ vertices. Then*

$$\text{rank } W(\tilde{D}_n) = \left\lfloor \frac{n}{2} \right\rfloor.$$

Proof. By Proposition 2.8 and Remark 2.7,

$$\text{rank } W(\tilde{D}_n) = \text{rank } W'(\tilde{D}_n) = \text{rank } \hat{W}(\tilde{D}_n) = \text{rank } W(B) = \left\lfloor \frac{n}{2} \right\rfloor.$$

□

We give an example of Proposition 2.8.

Example 2.10. Let \tilde{D}_8 be the extended Dynkin graph. Then

$$W(\tilde{D}_8) = \begin{bmatrix} 1 & 1 & 3 & 4 & 11 & 16 & 43 & 64 & 171 \\ 1 & 1 & 3 & 4 & 11 & 16 & 43 & 64 & 171 \\ 1 & 3 & 4 & 11 & 16 & 43 & 64 & 171 & 256 \\ 1 & 2 & 5 & 8 & 21 & 32 & 85 & 128 & 341 \\ 1 & 2 & 4 & 10 & 16 & 42 & 64 & 170 & 256 \\ 1 & 2 & 5 & 8 & 21 & 32 & 85 & 128 & 341 \\ 1 & 3 & 4 & 11 & 16 & 43 & 64 & 171 & 256 \\ 1 & 1 & 3 & 4 & 11 & 16 & 43 & 64 & 171 \\ 1 & 1 & 3 & 4 & 11 & 16 & 43 & 64 & 171 \end{bmatrix}$$

and

$$W'(\tilde{D}_8) = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 3 & 4 & 11 & 16 & 43 & 0 & 0 \\ 1 & 3 & 4 & 11 & 16 & 43 & 64 & 0 & 0 \\ 1 & 2 & 5 & 8 & 21 & 32 & 85 & 0 & 0 \\ 1 & 2 & 4 & 10 & 16 & 42 & 64 & 0 & 0 \\ 1 & 2 & 5 & 8 & 21 & 32 & 85 & 0 & 0 \\ 1 & 3 & 4 & 11 & 16 & 43 & 64 & 0 & 0 \\ 1 & 1 & 3 & 4 & 11 & 16 & 43 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

The Smith normal forms of $W(\tilde{D}_8)$ and $W'(\tilde{D}_8)$ are $\text{diag}(1, 1, 1, 7, 0, 0, 0, 0, 0)$ and the rank of $W(\tilde{D}_8)$ is 4.

Since the number of main eigenvalues of $W(\tilde{D}_n)$ is equal to the rank of $W(\tilde{D}_n)$, we have the following corollary.

Corollary 2.11. *Let \tilde{D}_n be the extended Dynkin graph with $n+1$ vertices. Then the number of main eigenvalues of \tilde{D}_n is $\lfloor \frac{n}{2} \rfloor$.*

3 Future works

In this paper, we show that the rank of the walk matrix of the extended Dynkin graph \tilde{D}_n is $\lfloor \frac{n}{2} \rfloor$. The next step would be to find the Smith normal form of the walk matrix of \tilde{D}_n . We give the following conjecture.

Conjecture 1. *Let \tilde{D}_n be the extended Dynkin graph with $n + 1$ vertices. Then the Smith normal form of $W(\tilde{D}_n)$ is*

$$\begin{cases} \text{diag}(\underbrace{1, \dots, 1}_{r-1}, n-1, 0, \dots, 0), & \text{if } n \text{ is even,} \\ \text{diag}(\underbrace{1, \dots, 1}_{r-1}, \frac{n-1}{2}, 0, \dots, 0), & \text{if } n \text{ is odd,} \end{cases}$$

where $r = \lfloor \frac{n}{2} \rfloor$.

References

- [1] Albeverio S, Ostrovskyi V, Samoilenko Y. On functions on graphs and representations of a certain class of \ast -algebras, J. Algebra 203 (2007) 567–582.
- [2] Cvetković DM, Gutman I., On spectral structure of graphs having the maximal eigenvalue not greater than two, Publ. Inst. Math. (Beograd) (N.S.) 18 (32) (1975) 39–45.
- [3] Draper Fontanals C. Notes on G_2 : the Lie algebra and the Lie group, Differential Geom. Appl. 57 (2018) 23–74.
- [4] Dokuchaev MA, Gubareni NM, Futorny VM, Khibina MA, Kirichenko VV. Dynkin diagrams and spectra of graphs, São Paulo J. Math. Sci. 7 (2013) 83–104.
- [5] Donovan P, Freislich MR. The representation theory of finite graphs and associated algebras, Carleton Mathematical Lecture Notes, No. 5, Carleton University, Ottawa, Ont., (1973).
- [6] Dynkin E. Classification of the simple Lie groups, Rec. Math. [Mat. Sbornik] N. S. 18 (60) (1946) 347–352.
- [7] Frappat I, Sciarrino A, Sorba P. Structure of basic Lie superalgebras and of their affine extensions, Comm. Math. Phys. 121 (1989) 457–500.
- [8] Gabriel P. Unzerlegbare Dartellungen I, Manuscr. Math, 6 (1972) 71–103.

- [9] Gantmacher FR. The theory of matrices Vols. 1, Chelsea Publishing Co. New York, 1959.
- [10] Hagos EM. Some results on graph spectra, *Linear Algebra Appl.* 356 (2002) 103–111.
- [11] Kruglyak SA, Nazarova LA, Roĭter AV., Orthoscalar Quiver Representations Corresponding to Extended Dynkin Graphs in the Category of Hilbert Spaces, *Funktsional. Anal. i Prilozhen.* 44 (2010) 57–73.
- [12] Nazarova LA. Representations of quivers of infinite type, *Izv. Akad. Nauk SSSR Ser. Mat.* 37 (1973) 752–791.
- [13] Ostrovs’kii VL, Samoilenko YS. On spectral theorems for families of linearly connected selfadjoint operators with prescribed spectra associated with extended Dynkin graphs, *Ukrain. Mat. Zh.* 58 (2006) 1556–1570.
- [14] Redchuk IK, Roĭter AV. Singular locally scalar representations of quivers in Hilbert spaces and separating functions, *Ukrain. Mat. Zh.* 56 (2004) 796–809.
- [15] Wang W., Wang C, Guo S. On the walk matrix of the Dynkin graph D_n , *Linear Algebra Appl.* 653 (2022) 193–206.
- [16] Yusenko K. On existence of $*$ -representations of certain algebras related to extended Dynkin graphs, *Methods Funct. Anal. Topology* 12 (2006) 197–204.