A Polynomial Decision for **3-sat**Revisited

M Angela Weiss

weiss@ime.usp.br, Universidade de São Paulo, São Paulo - Brasil

Abstract

We propose a polynomially bounded, in time and space, method to decide whether a given **3-sat** formula is satisfiable or not.

The tools we use here are, in fact, very simple. We first decide satisfiability for a particular **3-sat** formula, called *pivoted* **3-sat** and, after a plain transformation, still keeping the polynomial boundaries, it is shown that **3-sat** formulas can be written as pivoted formulas.

1 Introduction

After the preliminary definitions that establish a common language, we define a *Pivoted* **3-sat** formula, the main subject we deal with. After that, we show that any **3-sat** formula has a stronger version in the format of a Pivoted **3-sat** in the sense that a Pivoted **3-sat** formula is unsatisfiable if and only if the original **3-sat** formula is unsatisfiable and, if the original **3-sat** formula is a theorem (true under any valuation), its pivoted version is not a theorem

Secondly, we show that we can solve, that is, we can decide if the pivoted formula is **Satisfiable** or **Unsatisfiable**, in a polynomial use of space and time. Our resolution methods will require no more than basic knowledge of Boolean Logic and some basic skills.

An excellent way to start on the Complexity topic is by reading Scott Aaronson breathtaking survey,

https://www.scottaaronson.com/papers/pnp.pdf

The question is, "What is computing?". What kind of reasoning, operation, can be performed using an algorithm" and how fast the operation can be performed. See [2] for a general overview, or consult the roots, [12], [10], [11], [5], [1], [7], [8], [9].

A precise definition of *algorithm* was given by Alan Turing in 1937 (see [13]). The natural question that arises is: *What is the computational difficulty of performing such an algorithm?* See, in chronological order, [11], [10], [7], [5], [2]. Classification of complexity is found in [6], [4] and [2]. We aim to open a new trail to understand complexity questions on polynomially solving our problem that obviously can be solved in exponential time and space and draw new lines on the complexity classification.

In Section 2, we provide classical definitions of valuation over Boolean Logic, define SAT, pose some more standard definitions, and introduce Pivoted **3-sat** the special kind of **3-sat** we focus on. To solve the central problem, decide satisfiability or unsatisfiability, of Pivoted **3-sat** we write all the conjunctions of formulas as labeled edges of a digraph, the *cylindrical digraph*. We show how can we rewrite any **3-sat** formula as a pivoted **3-sat**.

In Section 3, we fully describe the algorithms we use in the process of deciding whether the closed digraph represents all 2^{m} combinations of valuation over the conjugated literals of a Pivoted **3-sat**. We will not write down all possible combinations to have a polynomial bound in our solution. Rather, we analyze the plain question of whether there is a satisfiable combination by generating a digraph.

Section 4 is the connecting link between a **3-sat** formula and a pivoted **3-sat** formula. We demonstrate that given a **3-sat** formula Ψ , then there exists a strong version of Ψ , Ψ_T , so that Ψ is unsatisfiable if and only if Ψ_T is unsatisfiable.

Section 5 is quite a technical section and, as usual, a profound and critical reading follows after unraveling the basic Section, 2 and the foundation part, Section 3.

Finally, a very long part of this paper is devoted to the examples, Section 6. This section can be regarded as a companion section.

This work presents a short version, revisited and with shorter proofs of a previuous published "A Polynomial Decision for 3-SAT", submitted to *arrxiv*, [14].

2 Basic Definitions

We work with a Boolean Language whose basic symbols are \lor, \land, \neg endowed with a finite set of atoms \mathcal{A} . The set of literals, \mathcal{L} , is the set $\mathcal{A} \cup \{\neg p | p \in \mathcal{A}\}$. A pair formed by a literal together with its negation is called a conjugated pair. Besides a finite set of atoms, we have the symbols \top and \bot .

Here we give the (classical) valuation definition over a Boolean formula. That is the definition we will use henceforth.

Definition 2.1. Given a finite set of atoms, \mathcal{A} , ψ belongs to the set of Boolean formulas F if

- 1. $\psi \in \mathcal{A}$;
- 2. $\psi = \chi \land \phi$ and both χ and ϕ belong to F;
- 3. $\psi = \chi \lor \phi$ and both χ and ϕ belong to F;
- 4. $\psi = \neg \chi$ and χ belongs to F;
- 5. ψ is either \top or \bot .

We use the symbols

1.
$$\psi \implies \chi$$
 to denote $\neg \psi \lor \chi$;

2. $\psi \equiv \chi$ to denote $(\psi \implies \chi) \land (\chi \implies \psi)$.

Definition 2.2. A valuation over a set of atoms \mathcal{A} is a mapping v from \mathcal{A} to the set {True, False} and, as usual, we extend the mapping v from a formula ψ to by reducing the complexity of writing a formula. Define the extension mapping v to a mapping v' from the formulas into the set {True, False} as above,

- 1. If $\psi \in \mathcal{A}$, $v'(\psi) = True$ if $v(\psi) = True$;
- 2. $\psi = \chi \land \phi$, $v'(\psi) = True$ if $v'(\chi) = True$ and $v'(\phi) = True$;
- 3. $\psi = \chi \lor \phi$, $v'(\psi) = True$ if $v'(\chi) = True$ or $v'(\phi) = True$;
- 4. $\psi = \neg \chi$, $v'(\psi) = False$ if and only if $v'(\chi) = True$;
- 5. $v'(\bot) = False and v'(\intercal) = True$.

If we do not have $v'(\psi) = True$, then $v'(\psi) = False$.

Definition 2.3 (SAT). A formula χ in Boolean Logic is said satisfiable if there is a valuation v from the set {True, False} onto the set of atoms of χ so that $v(\chi)$ is **True**. If no such valuation exists, that is $v(\chi)$ is **False** for any valuation, then we say that χ is unsatisfiable.

Any formula in the Boolean Logic is *decidable*, that is, given a formula χ , we can decide if χ is **satisfiable** (True for some valuation) or **unsatisfiable** (False for any valuation). The complexity problem consists of deciding between the values, **True** or **False** in the most optimal way, see [4].

We use the symbols: AND, OR, NOT and IMPLIES to denote external logical symbols. **Definition 2.4.** A 2-sat formula Ψ is a conjunction of a finite number of a disjunction of at most two literals. We write

$$\Psi \equiv (l_1^1 \lor l_1^2) \land \dots \land (l_s^1 \lor l_s^2) \equiv C_1 \land \dots \land C_s$$

A subformula $C_k \equiv l_k^1 \vee l_k^2, \ 1 \leq k \leq \mathbf{s}$ of Ψ is called a clause.

A 3-sat formula Ψ is a conjunction of a finite number of a disjunction of at most three literals. We write

$$\Psi \equiv (l_1^1 \lor l_1^2 \lor l_1^3) \land \dots \land (l_s^1 \lor l_s^2 \lor l_s^5) \equiv C_1 \land \dots \land C_s$$

 $A \ subformula \ C_k \equiv l_k^1 \lor l_k^2 \lor l_k^3, \ 1 \leq k \leq \mathbf{s} \ of \ \Psi \ is \ called \ a \ \text{clause}.$

Define pivoted **3-sat** formulas,

Definition 2.5. A Pivoted 3-sat formula is a 3-sat formula of the form,

$$\begin{array}{l} (a_1 \lor p_1^1 \lor q_1^1) \land \dots \land (a_1 \lor p_{k_{a_1}}^1 \lor q_{k_{a_1}}^1) \land \\ (\neg a_1 \lor r_1^1 \lor s_1^1) \land \dots \land (\neg a_1 \lor r_{k_{\neg a_1}}^1 \lor s_{k_{\neg a_1}}^1) \land \dots \land \\ (a_{\mathbf{m}} \lor p_1^{\mathbf{m}} \lor q_1^{\mathbf{m}}) \land \dots \land (a_{\mathbf{m}} \lor p_{k_{a_{\mathbf{m}}}}^{\mathbf{m}} \lor q_{k_{a_{\mathbf{m}}}}^{\mathbf{m}}) \land \\ (\neg a_{\mathbf{m}} \lor r_1^{\mathbf{m}} \lor s_1^{\mathbf{m}}) \land \dots \land (\neg a_{\mathbf{m}} \lor r_{k_{a\neg \mathbf{m}}}^{\mathbf{m}} \lor s_{k_{\neg \mathbf{m}}}^{\mathbf{m}}) \end{array}$$

or, in factorized form,

$$\begin{array}{l} (a_1 \lor \left((p_1^1 \lor q_1^1) \land \dots \land (p_{ka_1}^1 \lor q_{ka_1}^1) \right) \land \\ (\neg a_1 \lor \left((r_1^1 \lor s_1^1) \land \dots \land (r_{k_{\neg a_1}}^1 \lor s_{k_{\neg a_1}}^1) \right) \land \dots \land \\ (a_{\mathbf{m}} \lor \left((p_1^{\mathbf{m}} \lor q_1^{\mathbf{m}}) \land \dots \land (p_{ka_{\mathbf{m}}}^{\mathbf{m}} \lor q_{ka_{\mathbf{m}}}^{\mathbf{m}}) \right) \land \\ (\neg a_{\mathbf{m}} \lor \left((r_1^{\mathbf{m}} \lor s_1^{\mathbf{m}}) \land \dots \land (r_{ka_{\neg \mathbf{m}}}^{\mathbf{m}} \lor s_{k_{\neg a_{\mathbf{m}}}}^{\mathbf{m}}) \right) \end{array}$$

where the set of pivots, $\{a_1, \neg a_1, \ldots, a_m, \neg a_m\}$ has no intersection with the set of entries,

$$\{ p_1^1, q_1^1, \dots, p_{ka_1}^1, q_{ka_1}^1, r_1^1, s_1^1, \dots, r_{k_{\neg a_1}}^1, s_{k_{\neg a_1}}^1, \dots, p_1^{\mathbf{m}}, q_1^{\mathbf{m}}, p_{k_{\mathbf{m}}}^{\mathbf{m}}, q_{k_{\mathbf{m}}}^{\mathbf{m}}, r_1^{\mathbf{m}}, s_1^{\mathbf{m}}, \dots, r_{k_{\mathbf{m}}}^{\mathbf{m}}, s_{k_{\mathbf{m}}}^{\mathbf{m}} \}$$

Denote the pivot a_i by 1*i* and its conjugated pair $\neg a_i$ by 2*i* Call the set of entries of a_i the set of conjunctions,

$$(p_1^i \lor q_1^i) \land \dots \land (p_{ka_1}^i \lor q_{ka_1}^i)$$

and the set of entries of $\neg a_i$ the set of conjunctions,

$$(r_1^i \lor s_1^i) \land \dots \land (r_{k \neg a_1}^i \lor s_{k \neg a_1}^i)$$

We show, in Appendix 4 that any **3-sat** formula can be rewritten as a *logically equivalent* Pivoted **3-sat** formula.

Definition 2.6. Given a Pivoted formula Ψ , replace any clause of the form $a_i \lor p$, where a_i is a pivot, by the clauses $(a_i \lor p \lor q_p) \land (a_i \lor p \lor \neg q_p)$, where the pairs of conjugated, q_p and $\neg q_p$ are fresh new literals.

A pivoted **3-sat** formula is said complete if all of its clauses contain three literals.

Lemma 2.7. Any pivoted **2-sat** formula is logically equivalent to its complete version.

From now on, we work with complete **3-sat** pivoted formulas.

It is well known that **2-sat** formulas can be polynomially solved, a result originally shown in [3]. See [4]. We will store unsatisfiable groups of pivoted **3-sat** formulas in groups of **2-sat**.

Proposition 2.8. Given 2-sat formula Ψ ,

$$(p_1 \lor q_1) \land \cdots \land (p_k \lor q_k)$$

then, ϕ is unsatisfiable if and only if it there is a pair of conjugated literals, $\{a, \neg a\}$ and a set of conjugated literals

$$\{b_1, \neg b_1 \dots, b_r, \neg b_r, c_1, \neg c_1, \dots, c_s, \neg c_s\}$$

so that

 $(a \lor b_1) \land (\neg b_1 \lor b_2) \land \dots \land (\neg b_r \lor a)$

and

$$(\neg a \lor c_1) \land (\neg c_1 \lor c_2) \land \dots \land (\neg c_s \lor \neg a)$$

are subformulas of Ψ .

Explore the above result as a support tool to achieve our claim. We store the data contained in a pivoted **3-sat** formula in a cylindrical digraph.

Definition 2.9. Let Ψ be a pivoted (complete) **3-sat** formula. The cylindrical digraph associated to \mathcal{P} , $\mathbb{C}lndr = \langle V, E, label \rangle$, where V is a set of vertices, E is a set of edges and label is a mapping from E to the set of parts of the set of entries is,

- 1. The set of vertices is the set of literals together with their negations;
- 2. If $a \lor b$ belongs to the set of entries $\{i_1j_1, i_2j_2, \ldots, i_kj_k\}$, then,

(a) ¬a ⇒ b and ¬b ⇒ a belong to the set edges, E;
(b) i₁j₁, i₂j₂,..., i_kj_k is the label of both edges ¬a ⇒ b and ¬b ⇒ a.

Sequences of vertices connected by \Rightarrow are a path, as we define below,

Definition 2.10. Let a and b be two literals, there is a path from a to b if there is a subdigraph of \mathbb{C} Indr the form

$$a \stackrel{l_1}{\Rightarrow} c^1 \stackrel{l_2}{\Rightarrow} c^2 \stackrel{l_3}{\Rightarrow} \dots \stackrel{l_{k-1}}{\Rightarrow} c^{k-1} \stackrel{l_k}{\Rightarrow} b \tag{1}$$

Some subgraphs of the cylindrical digraph will be associated to unsatisfiable formulas and we keep track of these subgraphs and, thus, we can decide, in an optimal way whether they represent or not all possible combinations.

We search for sequences of the form,

$$\neg a \Rightarrow b^{1} \Rightarrow b^{2} \Rightarrow \dots \Rightarrow b^{t_{1}} \Rightarrow a$$

$$and$$

$$a \Rightarrow c^{1} \Rightarrow c^{2} \Rightarrow \dots \Rightarrow c^{t_{2}} \Rightarrow \neg a$$
(2)

We conclude that a Pivoted **3-sat** formula Ψ has an unsatisfiable combination, with the entries chosen among the set of labels over sequences depicted in Equation 2.

Notation 2.11. Let $2^{\mathbf{m}}$, the set of all mappings from \mathbf{m} to $2 = \{1, 2\}$. Write an element of $2^{\mathbf{m}}$ as $\{1j_1, 2j_2, \ldots, \mathbf{m}j_{\mathbf{m}}\}$, that is, the above set represents the graphic of a mapping $f : \mathbf{m} \mapsto 2$ and an entry, ij has the meaning that f(i) = j.

We identify an entry a_p with $\mathbf{p}1$ and an entry $\neg a_p$ with $\mathbf{p}2$, $1 \leq \mathbf{p} \leq \mathbf{m}$.

Definition 2.12. Given a Pivoted **3-sat** formula, C, our task is to verify, in the most efficient way, regarding the use of time and space, whether for all all combinations σ in $2^{\mathbf{m}}$, for all literals in σ regarded as False whether the resulting $2 - \mathbf{sat}$ formula is unsatisfiable. That is, we verify whether there is an element $\sigma = \{1j_1, 2j_2, \dots, \mathbf{mjm}\}$ in $2^{\mathbf{m}}$, in which the combination,

$$\bigwedge_{1 \le r \le \mathbf{m}} \{ \wedge (p^{rj_r} \lor q^{rj_r}) | (p^{rj_r} \lor q^{rj_r}) \in \text{ contents } rj_r \}$$
(3)

is satisfiable.

In short, we ask if the Pivoted **3-sat** formula is *unsatisfiable* or *sat-isfiable*, by first making valuations ranging over the set of pivots. Using Proposition 2.8, we search the sequences that lead to unsatisfiable formulas

and, thereafter, we decide whether we wrote all possible combinations or not.

We set the keystones by defining the cylindrical digraph associated with a given pivoted **sat** formula. Any disjunction contained in an entry plays the role of a vertex in a cylindrical digraph.

Definition 2.13. Given a Pivoted Formula Ψ , an interval [p,q] is the subdigraph that contains all sequences in between p and q and contains no loops, that is there is not a vertex r so that a sequence

 $r \Rightarrow \cdots \Rightarrow r$

belongs to [p,q] and the set of necessarily true literals, \mathcal{NEC} , is the set of all pair of conjugated literals so that the intervals $[\neg a, a]$ and $[a, \neg a]$ are non-empty.

Definition 2.14. A pair of necessarily true literals is a conjugated pair of literals of a given pivoted formula so that the sequences from $\neg a$ to a and from a to $\neg a$ are non empty.

The set of necessarily true literals is denoted by \mathcal{NEC} .

For each pair of conjugated literals, $\{a, \neg a\} \subseteq \mathcal{NEC}$, we write distinct copies of $[\neg a, a]$ and $[a, \neg a]$. As usual, distinct copies are fabricated using Cartesian Products to differentiate intervals.

Definition 2.15. Given a cylindrical digraph, suppose that the cardinality of \mathcal{NEC} is an even number $r \geq 2$. Enumerate the set of necessarily true literals as $\{p_1, \neg p_1, \ldots, p_{r/2}, \neg p_{r/2}\}$.

For each digraph, consider the product,

- 1. $[p_l, \neg p_l] \times \{2l\}, \ 1 \le l \le r/2;$
- 2. $[\neg p_l, p_l] \times \{2l 1\}, 1 \le l \le r/2.$

Define the edges and labels as the edges and labels inherited by the projection to the interval. Define a closed digraph as a digraph obtained with the union of intervals $[p_{2l-1}, \neg p_{2l-1}] \cup [\neg p_{2l}, p_{2l}], 1 \leq l \leq r/2$, endowed with the following identification $(p_{2l-1}, 2l-1) \sim (p_{2l}, 2l)$

The set of closed digraph, CSD is the union of all closed digraphs

Definition 2.16. A chain is a sequence contained in the set of closed digraphs,

$$(\neg a, 2l-1) \stackrel{l_0}{\Rightarrow} (c^1, 2l-1) \stackrel{l_1}{\Rightarrow} (c^2, 2l-1) \stackrel{l_2}{\Rightarrow} \dots \stackrel{l_{k-1}}{\Rightarrow} (c^k, 2l-1) \stackrel{l_k}{\Rightarrow} (a, 2l-1) \sim (a, 2l) \stackrel{l_{k+1}}{\Rightarrow} (c^{k+1}, 2l) \stackrel{l_{k+2}}{\Rightarrow} (c^{k+2}, 2l) \stackrel{l_{k+2}}{\Rightarrow} \dots \stackrel{l_{t-1}}{\Rightarrow} (c^{t+1}, 2l) \stackrel{l_t}{\Rightarrow} (\neg a, 2l)$$

The set of all chains is denoted by \mathcal{P} .

Consider the set of all chains in a closed digraph. Enumerate them by $\{1, \ldots, s\}$. To each chain Ch_t enumerated by t, consider the Cartesian Product $Ch_t \times \{t\}$, labels, edges inherited by the labels and edges in Ch.

Definition 2.17. The mapping

$$\begin{array}{ll} Ch: & \mathcal{P} \mapsto \mathcal{CSD} \\ & (v,i) \to v \end{array}$$

is the projection of vertices of \mathcal{P} onto \mathcal{CSD} .

The expansion to \mathcal{P} is the fixed (exponential) writing of all maximal chains contained in a closed digraph. The enumeration of the chains is arbitrary and, from now on, remains fixed.

Given the set of chains, if (v,t) belongs to a chain in \mathcal{P} , then v is a vertex in the cylindrical digraph and (u,v) is an edge of the cylindrical digraph if there is a chain Br_t and two arrows $(u,t) \Rightarrow (v,t)$ in Br_t , in Br_t .

Definition 2.18. Consider the edge $e = ((u,t) \Rightarrow (v,t))$ in \mathcal{P} . Define Ch(e) as

$$Ch(u,t) \Rightarrow Ch(v,t)$$

Definition 2.19. A compatible set of entries is a set $U = \{i_1 j_1, \ldots, i_n j_n\}$ so that for all pair i_r and i_s , if $i_r = i_s$ then $j_r = j_s$. If a set of entries is not compatible, it is said an incompatible set.

A set of edges $\mathbb{E} = \{e_1, e_2, \dots, e_k\}$ is said incompatible if the union of it set of labels, $U = \{i_1j_1, \dots, i_nj_n\}$ is an incompatible set of entries. If a set of edges is not incompatible, we say that \mathbb{V} is compatible.

We wrote digraphs that encode all unsatisfiable choices. We must decide whether we have, in fact, the set of all unsatisfiable choices. Again, we will not spell all the compatible, if any, choices. It is an expsize task.

Any chain Br and a compatible choice in Br is an unsatisfiable combination. After writing all unsatisfiable combinations we must decide whether we wrote all possible combinations (we have an unsatisfiable formula) or not (the pivoted formula is satisfiable).

Definition 2.20. Given a chain in \mathcal{P} ,

$$\mathbf{p} = c^0 \stackrel{l_1}{\Rightarrow} c^1 \stackrel{l_2}{\Rightarrow} c^2 \stackrel{l_3}{\Rightarrow} \dots \stackrel{l_{k-1}}{\Rightarrow} c^{k-1} \stackrel{l_k}{\Rightarrow} c^k$$

a compatible choice in **p** is a set of compatible entries $\mathcal{E} = \{j_1 i_1, \ldots, j_l i_l\}$, so that for each edge $e_i = c^i \Rightarrow c^{i+1}$, $0 \le i < k$, there is an entry $ij \in \mathcal{E}$ so that $ij \in label(e_i)$. If one casts a tableau where all combinations of pivots, $\sigma \in 2^{\mathbf{m}}$ labeled as *False* and there is a sequence

$$\begin{array}{c} \neg a \stackrel{q_0}{\Rightarrow} c^1 \stackrel{q_1}{\Rightarrow} c^2 \stackrel{q_2}{\Rightarrow} \dots \stackrel{q_{k-2}}{\Rightarrow} c^{k-1} \stackrel{q_{k-1}}{\Rightarrow} \\ a \stackrel{q_k}{\Rightarrow} c^{k+1} \stackrel{q_{k+1}}{\Rightarrow} c^{k+2} \stackrel{q_{k+2}}{\Rightarrow} \dots \stackrel{q_{t-2}}{\Rightarrow} c^{t-1} \stackrel{q_{t-1}}{\Rightarrow} \neg a = c^t \end{array}$$

where $\{q_0, q_1, q_2, \ldots, q_{k-1}, q_k, q_{k+1}, \ldots, q_{t-1}, q_t\} \subseteq \sigma$ and each q_j belongs to the label of the label l_j , $0 \leq j \leq t-1$, then the branch that falsifies the elements of σ closes. So, the next question is whether all possible sequences appear in \mathcal{P} and, therefore the pivoted formula is unsatisfiable or not, that is, the pivoted formula is satisfiable.

Given a Pivoted **3-sat** formula Ψ , the set of all compatible choices in \mathcal{P} is the set of all unsatisfiable combinations we can perform in Ψ . We now ask how one can decide whether the set of all compatible choices entails all possible combinations, thus, the Pivoted **3-sat** is unsatisfiable.

Define the set of vertices *orthogonal* to the set of chains.

Definition 2.21. A set of edges $\tilde{seq} = \{e_0, \ldots, e_n\}$ is called an antichain if for all $seq \in \mathcal{P}$, there is an element $e \in seq$ so that e is an edge of seq.

We do not focus on the set of vertices any more. Instead, focus on the set of *edges*, or, more specifically, on all the compatible choices of labels in \mathcal{P} . If all possible combinations were written, the formula would be unsatisfiable. Otherwise, the formula is satisfiable. Do all possible choices encompass all the possible combinations?

Definition 2.22 (ENTAILS). Let $T(2^{\mathbf{m}})$ be a set of partial mappings of \mathbf{m} into $\{1,2\}$. We say that $T(2^{\mathbf{m}})$ entails $2^{\mathbf{m}}$ if for all $\gamma \in 2^{\mathbf{m}}$, there is an $\eta \in T(2^{\mathbf{m}})$ so that η is a restriction of γ , that is, η is a subset of some $\gamma = \{i_11, i_22, \ldots, i_{\mathbf{m}}\mathbf{m}\}.$

We search for chains in CSD. Writing all the paths is a expsize long task. Recall that in Definition 2.15, \mathcal{P} denotes the set of all chains in CSD.

Definition 2.23. Let τ be the set of all entries $\eta = \{1j_1, \ldots, mj_m\}$ contained in 2^m so that

 $\exists seq \in \mathcal{P} \quad \forall e \in seq \quad \exists 1 \leq l \leq \mathbf{m}(i_l j_l \in label(e))$

Let $\tilde{\tau}$ be the complementary of τ , that is, the set of all entries $\tilde{\eta} = \{1j_1, \ldots, \mathbf{m}j_{\mathbf{m}}\}$ so that

$$\forall seq \in \mathcal{P} \; \exists e \in seq \; \forall 1 \leq l \leq \mathbf{m}(i_l j_l \notin label(e))$$

Proposition 2.24 below marks a decisive step to build our theoremhood. After building the closed digraphs, and applying Proposition 2.24, our search is focused on looking for compatible antichains. Again, we emphasize that our answer is a plain output, **YES** or **NO**, without naming the compatible antichains if any. In other words, we have a plain program and we expect an answer **YES** or **NO** regarding the possibility of the existence of some kind of input.

Proposition 2.24. The following assertions are equivalent,

- 1. There is no compatible set of entries in $\tilde{\tau}$;
- 2. τ entails 2^{m} ;
- 3. There is no compatible antichain in \mathcal{P} .

Proof. Clearly 1 and 2 are equivalent. If seq is a compatible set of entries in $\tilde{\tau}$, then, seq $\notin \tau$ for the sets are complementary.

1 and 2 imply 3. Suppose that there is a compatible antichain in \mathcal{P} , $Ach = \{e_1, \ldots, e_k\}$. Let lb(Ach) be the union of the set of labels of each vertex of Ach. As Ach is compatible, lb(Ach) has the form

$$\{t_1r_1, t_2r_2, \ldots, t_lr_l\}$$

where $1 \le t_1 < t_2 < ... < t_l \le \mathbf{m}$ and r_i is either 1 or 2. That is, for all $1 \le n \le l$ we have only one choice, either $r_n = 1$ or $r_n = 2$. So, for all $1 \le s \le \mathbf{m}$, we can choice a $\tilde{t}_s s$ so that $\tilde{t}_s \notin ach$.

Therefore, there is a non-empty set

$$Comp = \{\tilde{t_1}1, \tilde{t_2}2, \dots, \tilde{t_m}\mathbf{m}\}$$

that is compatible and $Comp \cap Ach = \emptyset$. Conclude that Comp belongs to $\tilde{\tau}$ and, thus, there is a compatible array in $\tilde{\tau}$.

Suppose that 1 is false and $s = \{t_1r_1, t_2r_2, \ldots, t_{\mathbf{m}}r_{\mathbf{m}}\}$ is a compatible set of entries in $\tilde{\tau}$ and, then, for all sequence $s_r \in \mathcal{P}$, there is an edge e_r so that $label(e_r) \cap s = \emptyset$. Let $A = \{e_r | e_r \in s_r AND \ (label(e_r) \cap s = \emptyset)\}$. As s is compatible, A is compatible, so, A is compatible antichain in \mathcal{P} . \Box

Due to Theorem 2.24, we can solve our game by deciding whether all antichains are incompatible or if there is a compatible antichain. Bear in mind that the word "entails $2^{\mathbf{m}}$ " encodes the fact that all the combinations in the pivoted **3-sat** formula are unsatisfiable.

3 The Search For Compatible Antichains

In this section, we develop the tools we use for deciding whether a closed digraph entails 2^m. Using Proposition 2.24, we search for compatible antichains in the closed digraph.

There is no need to search for unsatisfiable combinations. Using Proposition 2.24, we search compatible antichains, that is, we deal with pivots. Sequences of vertices were used to build the closed digraph. By using Proposition 2.24, our attention goes to the question of whether there is a compatible antichain in the closed digraphs and, thus, the set of closed digraphs do not entail 2^{m} . Again, we stress that we do not develop tools to write out all the antichains and, therefore, spell out all the possible compatible combinations which is an expsize problem. The polysize problem consists of the search for a plain output **YES**, there are antichains or **NO**, there is no antichain.

The set of all chains stores all possible unsatisfiable combinations one can perform in a card. We already wrote the closed digraph and, now, our attention is devoted to the search for compatible antichains, a search we perform in the set of labels. Each edge of a digraph has a label that will be used to point out the relevant parts that weigh in our search for antichains.

Given a closed digraph, we first modify the shape of the digraph to avoid exponential branching, the subject of the first Subsection 3.1. We must show that the reshaping, the *Linearized Digraph* does not change the set of compatible antichains.

In the second step, Subsection 3.2, after modifying the closed digraph we write a digraph that marks the compatible antichains orthogonal to the chains of the Linearized Digraph. A result of an empty digraph signalizes no compatible antichains in the Linearized Digraphs and, as we show in this Section, no compatible antichain in \mathcal{P} . Proposition 2.24 below marks a decisive step to build our theoremhood. After building the closed digraphs, and applying Proposition 2.24, our search is focused on looking for compatible antichains. Again, we emphasize that our answer is a plain output, **YES** or **NO**, without naming the compatible antichains, if any. In other words, we have a plain program and we expect an answer **YES** or **NO** regarding the possibility of the existence of some kind of input.

3.1 A Simpler Digraph

Here, we work to give a new shape to the set of closed digraphs. To avoid an exponential search, and keeping in mind that the search for compatible antichains has to be unchanged, we prove that there is a digraph, the *Linearized Digraph* less complex than the closed digraph. Moreover, we show there are compatible antichains in the Linearized Digraph if and only if there are compatible antichains in the closed digraph. Thereafter, we show that the search for compatible antichains over the Linearized Digraph is polynomial in time and space.

In this section, we develop the tools we use for deciding whether a closed digraph entails $2^{\mathbf{m}}$. Using Proposition 2.24, we can search for compatible antichains in the closed digraph but, instead of searching for unsatisfiable combinations, we search for compatible antichains, that is, we deal with pivots, subject of Subsection 3.2. The set of vertices, sequences of vertices, played the main role in building closed digraphs. Guided by Proposition 2.24, our attention goes to the question of whether the labels contain a compatible antichain and, thus, the set of closed digraphs do not entail $2^{\mathbf{m}}$. Again, we stress that we do not develop tools to write out all the antichains and, therefore, spell out all the possible compatible combinations which is, clearly, an expsize problem. The polysize problem consists of the search for a plain output **YES**, there are antichains or **NO**, there is no antichain.

The set of all chains stores all possible unsatisfiable combinations one can perform in a card and we already wrote the closed digraph.

Given a closed digraph, we first modify the shape of the digraph to avoid exponential branching, the subject of the first Subsection 3.1. We must show that the reshaping, the *Linearized Digraph* does not change the set of compatible antichains.

Our proof strategy, schematized to avoid exponential multiplication on writing the set of chains, is the transformation of closed digraphs into linear digraphs, that is, digraphs with no source or drain beside the top and root. Using Lifting, Multiplication of Branches and the Addition of Labels, we define next, we modify the branching to focus on the question: "Are there compatible antichains?" instead of asking: "Can we name all of them?".

Definition 3.1. Given a closed digraph, define,

- 1. R, the set of roots is $\{a \in V | \nexists b \in V((a \Rightarrow b) \in E)\};$
- 2. T, the set of tops is $\{a \in V | \nexists b \in V((b \Rightarrow a) \in E\}$;
- 3. If there are at least two edges

$$\begin{array}{l} a \Rightarrow b_1 \\ a \Rightarrow b_2 \end{array}$$

$$b_1 \Rightarrow a \\ b_2 \Rightarrow a$$

where $b_1 \neq b_2$, then a is a branching.

Definition 3.2. A sequence of labeled edges in a closed digraph is called a maximal branch if seq is of the form

$$c^{0} \stackrel{l_{1}}{\Rightarrow} c^{1} \stackrel{l_{2}}{\Rightarrow} c^{2} \stackrel{l_{3}}{\Rightarrow} c^{3} \dots c^{k-1} \stackrel{l_{k}}{\Rightarrow} c^{k} \tag{4}$$

where c^k is either a root or a branching and c^0 is either a top or a branching and no other branching belongs to the sequence.

If an interval [a,b] contains no branching besides a and b, then it is called a linear interval.

A linear interval [a, b] where b is a root, is called a root interval.

Definition 3.3. We use the term branch to design a linear sequence

$$c^0 \Rightarrow c^1 \Rightarrow \dots \Rightarrow c^{k-1} \Rightarrow c^k$$

where except by, perhaps, c^0 and c^k , no other vertex is a branching.

We will define the operations over the closed digraphs, Lifting, Multiplication of Branches and the Addition of Labels, the operations, 3.4, 3.5 and 3.7 and, without any loss in our search for compatible antichains, the given closed digraph is reshaped into a simpler digraph, the *Linearized Digraph*. Of course, we have to show that we do not lose or gain information about compatible antichains. Examples of the operations and more explanations can be found in the institutional page,

In addition, Example 6.1 illustrates the use of all operations defined next.

Definition 3.4 (Lifting). Given a closed digraph Gr,

• If B and R are two consecutive linear intervals, that is, they share a common branching, v,

$$br_{11} = v_{11} \Rightarrow \dots \Rightarrow v_{1t_{11}} \Rightarrow v$$
$$\dots$$
$$br_{1k} = v_{1k} \Rightarrow \dots \Rightarrow v_{kt_{1k}} \Rightarrow v$$

or

and

$$br_{21} = v \Rightarrow v_{21} \Rightarrow \dots \Rightarrow v_{2t_{21}}$$
$$\dots$$
$$br_{2k} = v \Rightarrow v_{2k} \Rightarrow \dots \Rightarrow v_{2t_{2k}}$$

the Lifting of Gr by B and R is $Gr(B \star R) = \langle V_{B\star R}, E_{B\star R}, label \rangle$,

$$V_{B\star R} = (V_G \smallsetminus \{v\}) \cup (\{v\} \times \{1, \dots, k\})$$

$$E_{B\star R} = (E_{Gr} \smallsetminus \{v_{1t_{11}} \Rightarrow v, \dots, v_{kt_{ik}} \Rightarrow v, \dots, v \Rightarrow v_{21}, \dots, v \Rightarrow v_{2k}\})$$

$$\cup \{v_{1t_{11}} \Rightarrow (v, 1), \dots, v_{kt_{1k}} \Rightarrow (v, k),$$

$$\dots, (v, 1) \Rightarrow v_{21}, \dots, (v, k) \Rightarrow v_{2k}\}$$

The labels of each $w \Rightarrow (v, j)$ and $(v, j) \Rightarrow w$ are the same labels associated with $w \Rightarrow v$ and $v \Rightarrow w$, respectively.

• If R is of the form

$$br_1 = v_1 \Rightarrow \dots \Rightarrow v_{t_1} \Rightarrow r$$
$$\dots$$
$$br_n = v_n \Rightarrow \dots \Rightarrow v_{t_n} \Rightarrow r$$

where r is a root, the Lifting of Gr by R is $Gr(R) = \langle V_R, E_R, label \rangle$ given by,

$$V_R = (V \setminus \{r\}) \cup (\{r\} \times \{1, \dots, n\})$$

$$E_R = (E \setminus \{v_{t_1} \Rightarrow r, \dots, v_{t_n} \Rightarrow r\}) \cup$$

$$\{v_{t_1} \Rightarrow (r, 1), \dots, v_{t_n} \Rightarrow (r, n)\}$$

The labels of each $v_{t_j} \Rightarrow (r, j)$ are the same labels of $v_{t_j} \Rightarrow r$.

• If B is of the form

$$br_{11} = a \Rightarrow v_1 \Rightarrow \dots \Rightarrow v_{t_1}$$
$$\dots$$
$$br_{1k} = a \Rightarrow v_m \Rightarrow \dots \Rightarrow v_{t_m}$$

where a is a top, the Lifting of Gr by B is $Gr(B) = \langle V_B, E_R, label \rangle$ given by,

$$V_B = (V \setminus \{a\}) \cup (\{a\} \times \{1, \dots, m\})$$
$$E_B = (E \setminus \{a \Rightarrow v_1, \dots, a \Rightarrow v_m\}) \cup$$
$$\{(a, 1) \Rightarrow v_1, \dots, (a, m) \Rightarrow v_m\}$$

The labels of each $(a, j) \Rightarrow (v_j)$ are the labels of $a \Rightarrow v_j$

Definition 3.5 (Multiplication of a branch). Let

 $br = v \Rightarrow v_1 \Rightarrow \dots \Rightarrow v_j \Rightarrow u$

be a maximal branch contained in a linear interval $[v, u] = \langle V, E, label \rangle$, m = $\{1, \ldots, m\}$ a finite set and $M = \{v_1, \ldots, v_j\} \times \{1, \ldots, m\}$.

The Multiplication of the branch br is the the interval [v, u]', where the vertices are given by the set $(V \setminus \{v_1, \ldots, v_j\}) \cup M$ and the edges of $V \setminus \{v_1, \ldots, v_j\}$ plus the edges in the new sequences,

$$v \Rightarrow (v_1, i) \Rightarrow \dots \Rightarrow (v_j, i) \Rightarrow u$$

 $1 \leq i \leq m$.

As one cannot form edges among vertices in distinct linear sequences, the definition is consistent.

Definition 3.6. Given a closed digraph and a vertex a, define Up_a and $Down_a$, respectively as sets of vertices,

$$\{ v \in V | \exists n \in \mathbb{N} \exists \{ v_1, v_2, \dots, v_n \} \subseteq V | (v \Rightarrow v_1 \Rightarrow v_2 \Rightarrow \dots \Rightarrow v_n \Rightarrow a \}$$
$$\{ w \in V | \exists n \in \mathbb{N} \exists \{ w_1, w_2, \dots, w_n \} \subseteq V | (a \Rightarrow w_n \Rightarrow \dots \Rightarrow w_2 \Rightarrow w_1 \Rightarrow w \}$$

An edge $v_1 \Rightarrow v_2$ belongs to Up_a if v_1 belongs to Up_a . An analogous definition is given to $Down_a$, that is, v_2 belongs to $Down_a$.

Next, we present a tool to avoid the creation of new compatible antichains in the linearized digraph. Suppose there is no addition of new labels. In that case, the Linearized Digraph we build next as a simpler digraph will present new compatible antichains with no corresponding, under a suitable mapping, antichain in \mathcal{P} .

Definition 3.7 (Adding a Label). Let the set of the branching in a closed digraph be $B = \{a_1, \ldots, a_t\}$. Consider a set of conjugated pairs of literals, $B = \{p_{a_1}, p_{\neg a_1}, \ldots, p_{a_t}, p_{\neg a_t}\}$ that are all distinct from the literals we have been using. The addition of the new pair of literals to the set of branching is the new set of labels, label' defined by,

$$label'(e) = label(e) \cup \{p_{a_i} | e \in Down(a_i)\} \cup \{\neg p_{a_i} | e \in Up(a_i)\}$$

We do not overload our manuscript, so we drop the prime symbol, *label'* and write only *label*,

In Example 6.1, we emphasize the role of adding a pair of conjugated literals that work to prevent the creation of a new antichain that never existed in the originally set \mathcal{P} .

To perform Lifting, Multiplication of Branches and Addition of a Label to obtain a simpler digraph, we must show that there will be no prejudice in counting antichains. We cannot create or erase antichains.

Reshape a closed digraph into a set of linear intervals on Procedure 3.8. The mainstay to sustain our construction is that, after writing Procedure 3.8, we obtain a linear digraph that is equivalent to the originally closed digraph. We show that our remodeling will not impair the preexisting antichains or create new antichains.

Lifting, Multiplying Branches and Adding labels in a closed digraph are operations used to generate linear digraphs with no branching and, mainly, no loss of information about compatible antichains stored in \mathcal{P} with no exponential multiplication of branches. The operations are illustrated in Example 6.1.

Procedure 3.8 (Changing a Closed Digraph into a Linear Digraph). Given a closed digraph CSD, and a set of conjugated literals distinct of any literals from the set of labels of CSD, $S = \{p_1, \neg p_1, \ldots, p_r, \neg p_r\}$, where V is the cardinality of literals (vertices). First, we transform the closed digraph, by Multiplying branches, into a digraph whose intervals, from branching to a root, are maximal linear sequences. After we obtain a digraph whose intervals, from branching to a root, are maximal linear sequences, we multiply branches, add suitable labels, and perform lifting.

1. Let the roots of the closed digraph be $\{e_1, \ldots, e_n\}$ and let the set of all root intervals be

$$\begin{split} r_{11} &\Rightarrow \cdots \Rightarrow e_1 \\ \vdots \\ r_{l_11} &\Rightarrow \cdots \Rightarrow e_1 \\ \vdots \\ r_{1s} &\Rightarrow \cdots \Rightarrow e_n \\ \vdots \\ r_{l_s} &\Rightarrow \cdots \Rightarrow e_n \end{split}$$

lift each maximal branch. Obtain the digraph Modf,

```
r_{11} \Rightarrow \dots \Rightarrow (e_1, 1)
\vdots
r_{l_11} \Rightarrow \dots \Rightarrow (e_1, l_1)
\vdots
r_{1s} \Rightarrow \dots \Rightarrow (e_r, 1)
\vdots
r_{l_ss} \Rightarrow \dots \Rightarrow (e_r, l_r)
```

2. Suppose we obtain sets of root intervals of the form,

```
\begin{array}{l} u \Rightarrow \cdots \Rightarrow v_1 \\ \vdots \\ u \Rightarrow \cdots \Rightarrow v_n \end{array}
```

where $\{v_1, \ldots, v_n\}$ are roots in the digraph.

Let all the maximal branches ending on u be,

$$w_1 \Rightarrow \dots \Rightarrow u$$

:
$$w_t \Rightarrow \dots \Rightarrow u$$

Multiply the branches by $m = max\{n,t\}$ and lift. Add to the set of labels a pair of conjugated literals p_{u_i} and $\neg p_{u_i}$ never used in our process, to, respectively the edges in $Down_{u_i}$ and Up_{u_i} . Rewrite Modf as,

$$\begin{split} w_1 &\Rightarrow \cdots \Rightarrow (u,1) \Rightarrow \cdots \Rightarrow v_1 \\ \vdots \\ w_m &\Rightarrow \cdots \Rightarrow (u,m) \Rightarrow \cdots \Rightarrow v_m \end{split}$$

Proceed until all intervals are lifted.

3. Finally, we have sets of maximal branches of the form,

$$\begin{array}{l} v \Rightarrow \cdots \Rightarrow u \Rightarrow \cdots \Rightarrow t_1 \\ \vdots \\ v \Rightarrow \cdots \Rightarrow u \Rightarrow \cdots \Rightarrow t_o \end{array}$$

where v is a top and $\{t_1, \ldots, t_o\}$ are roots. Lift all the branches and obtain

$$(v,1) \Rightarrow \dots \Rightarrow u \Rightarrow \dots \Rightarrow t_1 \\ \vdots \\ (v,o) \Rightarrow \dots \Rightarrow u \Rightarrow \dots \Rightarrow t_o$$

Call the Linearized Closed Digraph to the linear digraphs obtained after the successive applications of Lifting, Multiplication and Addition of a label Denote the Linearized Closed Digraph by Linclsd.

Suppose that each linear branch has the cardinality k and write each linear branch in Lincloid as Br_i as a short for $Br_i \times \{i\}$. Define the edges whose arrow is inherited by $v \Rightarrow u$.

We show that a Linearized Digraph, despite being simpler than the set of chains, preserves compatible antichains in the sense that we do not create nor erase information about compatible antichains, Lemma 3.17. Recall that the vertices of either \mathcal{P} and *Linclsd* are pairs, a vertex in the closed digraph and a number.

Definition 3.9. The mapping

$$\begin{array}{ll} Proj: & Linclsd \to \mathcal{CSD} \\ & (v,j) \mapsto v \end{array}$$

is the projection of vertices of Linclsd onto CSD.

Definition 3.10. Consider the edge $e' = (u' \Rightarrow v')$ in Linclsd. Define Proj(e') as the edge,

$$Proj(u') \Rightarrow Proj(v')$$

Definitions 3.10 and 2.18 are well posed for if $u' \Rightarrow v'$ and $u'' \Rightarrow v''$ are two edges, respectively in *Linclsd* and in \mathcal{P} , then $Proj(u') \Rightarrow Proj(v')$ and $Ch(u'') \Rightarrow Ch(v'')$ are, equally, edges in \mathcal{CSD} .

Lemma 3.11. Let ach be an antichain in the Linearized Digraph. Then, for all branching a, either $Proj(Ach) \cap Up(a)$ or $Proj(Ach) \cap Down(a)$ is empty.

Proof. Due to the addition of labels, there is no branching $a \in CSD$ so that two edgers $e_1 \in Up(a)$ and $e_2 \in Down(a)$ belong to a compatible antichain *ach* in the Linearized Digraph because both are incompatible.

Definition 3.12. The set of all antichains in \mathcal{P} is denoted by Σ and the set of all antichains in Linclsd is denoted by Θ .

Lemma 3.13. The mapping $Proj : \Theta \mapsto \mathcal{P}(Linclsd)$, where $\mathcal{P}(Linclsd)$ is the set of parts of Linclsd, given by

$$Proj(\sigma) = \{v \in Linclsd | \exists w \in \sigma Proj(w) = v\}$$

is an injective mapping.

Have we created or erased antichains? As the *relevant antichains* are kept, we search for compatible antichains in *Linclsd*. We show that there are compatible antichains in *Linclsd* if and only if there are compatible antichains in \mathcal{P} .

Given a branching a, recall the definition of a branch in Up(a) is a linear sequence, in \mathcal{P} of the form,

$$v_1 \Rightarrow v_2 \Rightarrow \cdots \Rightarrow a$$

Definition 3.14. For all set of edges Z in \mathcal{P} and branching a, we say that Z represents Up(a) if for all branching Br in Up(a), there is a $z \in Ch(Z)$ so that z belongs to Br. A symmetric definition applies to Down(a), that is, for all branching Br in Down(a), there is a $z \in Ch(Z)$ so that z belongs to Br.

For all set of edges Z in \mathcal{P} , Z involves a set of branching μ if for all branching $a \in \mu$, either Z represents Up(a) or Z represents Down(a).

Proposition 3.15. Given an antichain σ in \mathcal{P} , for all branching a, σ represents Up(a) or Down(a).

Proof. Let σ be an antichain in \mathcal{P} . The antichain Up(a) or Down(a) where "or" is not exclusive.

Suppose otherwise. Let the branches of Up(a) and Down(a) as, respectively,

Combine the above chains and obtain the chains in \mathcal{P} ,

and

If the antichain σ has no edges in columns, say r and s in, respectively Up(a) and Down(a), then the combination of the two columns $r \times s$, does not contain any edge of the antichain.

In conclusion, for any antichain σ , for any branching a, either all edges of $Ch(\sigma)$ are a member of one edge in Up(a) or all edges of $Ch(\sigma)$ belong to Down(a).

Proposition 3.16. Given an antichain σ in \mathcal{P} , σ involves the set of branching.

Proof. Let $a_1 < \cdots < a_s$ be an arbitrary ordering of all branches in the closed digraph.

Use Proposition 3.15 to a_1 . If σ represents $Up(a_1)$, obtain a new antichain σ_1 by defining,

1. Let $Br_{11}, \ldots, Br_{1s_1}$ be the set of branches in $Up(a_1)$, still in the closed digraph. For all $1 \le i \le s_1$, choose a $x_i \to y_i \in Br_{1i} \cap Ch(\sigma)$. Erase any edge whose projection via Ch in Br_{1i} is different from $x_i \to y_i$ and add to σ_1 all $(x_i, t) \to (y_i, t)$ whose projection is $x_i \to y_i$.

Otherwise, σ represents $Down(a_1)$ and proceed similarly.

Obtain a new antichain σ_1 . Notice that $Ach(\sigma_1) \subseteq Ach(\sigma)$. Suppose that we obtained σ_i , $1 \le i \le t < r$ and that $Ach(\sigma_t) \subseteq \cdots \subseteq Ach(\sigma_1) \subseteq Ach(\sigma)$. Suppose that σ_i represents $Ur(\sigma_i)$

Suppose that σ_t represents $Up(a_{t+1})$.

1. Let $Br_{t+11}, \ldots, Br_{t+1s_{t+1}}$ be the branches in $Up(a_{t+1})$. For all $1 \leq i \leq s_{t+1}$, choose a $x_i \to y_i \in Br_{t+1i} \cap Ch(\sigma_t)$. Erase any edge whose projection via Ch in Br_{1t+i} is different from $x_i \to y_i$ and add to σ_{t+1} all $(x_i, t) \to (y_i, t)$ whose projection is $x_i \to y_i$.

Otherwise, σ_t represents $Down(a_{t+1})$ and a mirror reasoning applies.

As a result, obtain a series of nested antichain that we build (can build in several random ways), $Ach(\sigma) \supseteq \cdots \supseteq Ach(\sigma_t)$, where the cardinality of t does not exceed the cardinality of the set of branching. Given the already chosen ordering, for all branching a, we choose preferentially Up(a), if possible, and, if we do not have this choice, we have the choice Down(a).

We have that $Ach(\sigma_t)$ cannot be reduced and, using Proposition 3.15 represents any branching in the set of branches, thus, σ , as well as its reduction, involves the set of branching.

Lemma 3.17. There is an onto mapping Eq from Σ to Θ .

Proof. The operations of Lifting, Multiplying and Adding Labels are fixed, so, after reducing an antichain σ to its minimal version σ_t , we obtain, in the Linearized Digraph, an antichain $\overline{\sigma_t}$, the result of the fixed operation over σ_t in the Linearized Digraph. The onto mapping is given by $Eq(\sigma)$ is the expansion of $Ch(\sigma_t)$.

In short, our quest is whether there is a compatible antichain in a closed digraph is equivalent to the quest whether there is a compatible antichain in a linearized digraph.

3.2 Digraph of Compatible Antichains

Now, our attention is devoted to the search for compatible antichains, which we perform in the set of labels. Each edge of a digraph has a label that will be used to point out the relevant parts that weigh in our search for antichains.

In outline, once we have linear digraphs, k linear digraphs, each linear digraph in the column r endowed l_r vertices. To decide the existence of compatible antichains in the Linearized Digraph, we start with all the vertices in the first column in interaction with each vertex from the second column to the k^{th} column, We obtain, after performing over all vertices in the first label, a nested compatible digraph of depth 1, We proceed column by column. In the h^{th} , each vertex v_i^h , $1 \le i \le l_h$ visits all vertex v_j^n , n > h, $1 \le j \le l_n$, and built a nested digraph with dept h whose root is v_j^n that encodes compatible antichains that passes through each v_i^h and v_j^n .

We introduce *Nested Digraphs*, the tool to find out compatible antichains in the linearized digraphs. In the world of Nested Digraphs, the set of labels plays the role of vertices. To avoid a heavy load, we discard unnecessary elements. Labels are the only fundamental elements and, as we advance our search, labels play the role of vertices and the connection if given by compatibility.

Up to now, we no longer need the set of literals, that played the role of vertices in the Linearized Digraph and whose sequences encode unsatisfiable combinations. We pursued for new strategy for solving a card, once we cleaned new paths. The maximal chains in the Closed Digraph were used to mark the unsatisfiable combinations and our search is centered on the search for a compatible antichain. So, we are dealing only with the set of labels.

It is time to discharge vertices and edges in the Linearized Digraph because these elements were used to mark unsatisfiable choices and we no longer use them. We use lighter gadgets ruling out the set of vertices, the set of literals.

A sequence of linear digraphs of the form,

$$a \stackrel{l_1}{\Rightarrow} b_1 \stackrel{l_2}{\Rightarrow} \dots \stackrel{l_t}{\Rightarrow} b_t \stackrel{l_{t+1}}{\Rightarrow} c$$

corresponds to the sequence of edges

$$l_1 \rightarrow l_2 \rightarrow \cdots \rightarrow l_t \rightarrow l_{t+1}$$

After modifying the closed digraph we write a digraph that marks the compatible antichains orthogonal to the chains of the Linearized Digraph. A result of an empty digraph signalizes no compatible antichains in the Linearized Digraphs and, as we show in this Section, no compatible antichain in \mathcal{P} .

Let (r, s) be a enumeration of the pair *column* and *location* in the array of k columns each with m_k elements. Write the pair edge together with the enumeration. An edge $(p, i, j) \Rightarrow (q, i, j+1) \Rightarrow (r, i, j+2)$ belongs to column i and the vertices belong to, respectively, the j, j+1 and j+2 places. Define,

Definition 3.18. Given a Linearized Digraph, define its Companion Digraph as the digraph whose vertices are the set of indexes associated with each edge of the linearized digraph. Define an edge $(l_1, i, j) \rightarrow (l_2, i, j + 1)$ if there is a sequence $(p, i, j) \stackrel{l_1}{\Rightarrow} (q, i, j + 1) \stackrel{l_2}{\Rightarrow} (r, i, j + 2)$ in the linearized digraph.

If $(p,i,j) \stackrel{l_1}{\Rightarrow} (q,i,j+1) \stackrel{l_2}{\Rightarrow} (r,i,j+2)$ and $(p',i',j') \stackrel{l_1}{\Rightarrow} (q',i',j'+1) \stackrel{l_2}{\Rightarrow} (r',i',j'+2)$ we have two distinct edges $(l_1,i,j) \rightarrow (l_2,i,j+1)$ and $(l_1,i',j') \rightarrow (l_2,i',j'+1)$, spite labels are the same.

We will not overload our manuscript. Distinct edges are originated from distinct columns and places. From now on we omit all extra notation and drop the pair column location.

Now, set the scene for building nested digraphs associated with each vertex (a label). We expect that a k^{th} -interaction to generate an output **satisfiable**. We set the necessary definitions.

Definition 3.19. Let Lab be a Digraph and v a vertex of Lab. An edgelabeled digraph for v is a closed rooted digraph whose edges are labeled and whose root is v. We label the edges with a set of vertices, that is, there is a mapping from the set of edges of Gr(v) into sets of vertices.

$$G(v) = \langle E_v, V_v, labels_v : E_v \mapsto \mathcal{P}(E_v) \rangle$$

The mapping labels_v associate to each edge a set of vertices and labels(v) is called the label of the vertex v.

We require, moreover, that all maximal sequences in G length p have length p.

Define the basic operations over *labeled digraphs* we perform to obtain a simpler digraph and, without altering the state of the closed digraph concerning the question there are compatible antichains or not.

Definition 3.20. Given two labeled digraphs whose root is v,

$$Gr_v = \langle V_v, E_v, labelsv : V_v \mapsto \mathcal{P}(E_v) \rangle$$

and

$$Gr'_v = \langle V'_v, E'_v, labelsv' : E'_v \mapsto \mathcal{P}(E_v) \rangle$$

their union, $Gr_v \cup Gr'_v$ is given by

$$\langle V \cup V', E \cup E', labels''_v : E \cup E' \mapsto \mathcal{P}(V \cup V') \rangle$$

where $labels''_v : E \cup E' \mapsto \mathcal{P}(V \cup V')$ is given by

 $\begin{aligned} labels''_v &= label(v), \text{ if } v \in E_v \setminus E'_v \\ labels''_v(v) &= label'(v), \text{ if } v \in E'_v \setminus E_v \\ labels''_v(v) &= label(v) \cup label'(v), \text{ if } v \in E_v \cap E'_v \end{aligned}$

Their intersection is given by,

$$\langle E_v \cap E'_v, E_v \cap E'_v, labels'' : E_v \cap E'_v \mapsto \mathcal{P}(E_v \cap E'_v) \rangle$$

where labels" is given by labels"(v) = label(v) \cap label'(v), if $v \in E_v \cap E'_v$.

Definition 3.21 (Nested Digraph). A rooted edge-labeled digraph, $Gr_v = \langle V_v, E_v, labelsv : E_v \mapsto \mathcal{P}(E_v) \rangle$ is a nested digraph of length p if Gr_v is the union of a number s of linear digraphs of a fixed dept $p \ge 1$,

$$v \leftarrow a_i^2 \leftarrow a_i^3 \leftarrow \dots \leftarrow a_i^p$$

so that for each linear digraph, we have $label(a_i^{i-1} \leftarrow a_i^i) = \{a_i^p, a_i^{p-1}, \dots, a_i^i\}, 1 \le i \le s.$

Notice that not necessarily vertices at each level j, $1 \le j \le p$ are distinct but, at each distinct level, the vertex are pairwise distinct.

A compatible nested digraph of length p is the union of linear digraphs of length p whose labels are compatible sets.

Lemma 3.22. The union of two nested digraphs of depth p, G(v) and G(v)' is a nested digraph.

The intersection of two nested digraphs of depth p is not necessarily a nested digraph but we can select *the* maximal nested digraph contained in the intersection. Uniqueness follows from the fact that, given two nested maximal digraphs, G(v) and G(v)', then the union of a nested digraph is nested and, therefore, we cannot have two distinct maximal nested digraphs contained in $G(v) \cap G(v)'$.

The maximal nested digraph contained in the intersection of two nested digraphs, G(v) and G(v)', contains the set of all compatible antichains contained in both G(v) and G(v)'.

Definition 3.23. Let G(v) and G(v)' be two nested digraphs of depth p. Denote by maxG(v) the Maximal Nested Subdigraph contained in $G(v) \cap G(v)'$.

Write an algorithm to obtain maxG(v), the maximal nested digraph contained in an intersection of two nested digraphs. The idea behind the two algorithms is, first, from the lower level to the last level, to select nested linear branches. Thereafter, select nested linear branches of level k.

We present two algorithms we perform to obtain the maximal nested digraph contained in the intersection of two nested digraphs that share the same root.

Consider two nested digraphs and their intersection given by, respectively,

$$G(v)' = \langle V'(v), E'(v), label'(v) : E'(v) \mapsto \mathcal{P}(V(v)) \rangle$$

$$G(v)'' = \langle V''(v), E''(v), label''(v) : E''(v) \mapsto \mathcal{P}(V(v)) \rangle$$

$$G(v) = G(v)' \cap G(v)'' = \langle V(v), E(v), label(v) : E(v) \mapsto \mathcal{P}(V(v)) \rangle$$

There is an algorithm to write maxG(v). Written below into two pseudo codes,

Pseudocode 3.24. Given a digraph $G(v) = \langle V, E, label \rangle$, intersection of two nested digraphs, using the first pseudocode obtain a new rooted digraph in which, using the second pseudocode we obtain maxG(v).

We denote the vertices of the intersection by v_i^m , where m is a level and i is the position of the vertex in the level m and $v = v_1^0$.

First Pseudocode: From $m \coloneqq 0$, step 1, to k - 2, do:

For all edges $u_i^m \leftarrow v_j^{m+1}$, consider the vertices that form an ascending sequence of length 2 with $u_i^m \leftarrow v_j^{m+1}$, say, the set $\{w_1^{m+2}, \ldots, w_l^{m+2}\}$. For s

ranging from 1 to l, consider as provisional labels,

$$lbl_{i,j,s}(w_s^{m+2} \to v_j^{m+1}) = label(w_s^{m+2} \to v_j^{m+1}) \cap label(u_j^{m+1} \to v_i^m)$$

Define,

$$label'(u_j^{m+1} \to v_i^m) = \cup_s lbl_{i,j,s}(w_l^{m+2} \to v_j^{m+1}) \cup \{v_j^{m+1}\}$$

Once we defined all labels, $label'(u_j^{m+1} \rightarrow v_i^m)$, define

$$label(w_s^{m+2} \to v_j^{m+1}) = \cup_i lbl_{i,j,s}(w_s^{m+2} \to v_j^{m+1})$$

and, finally, $label(u_j^{m+1} \rightarrow v_i^m) \leftarrow label'(u_j^{m+1} \rightarrow v_i^m)$.

The final result of applying the first pseudocode is the union of linear digraphs ordered by \supseteq .

Second Pseudocode: The second pseudocode is summarized below. We search backwards, for m := k STEP - 1 UNTIL 2.

For all edges $(u_i^m \to v_j^{m-1})$ so that $u_i^m \in label(u_i^m \to v_j^{m-1})$, define $G_{u_i^m v_j^{m-1}}$ as the union of all linear sequences

$$w_{r_k}^k \to w_{r_{k-1}}^{k-1} \to \dots \to u_i^m \to v_j^{m-1} \to \dots \to v$$

where $\{u_i^m, v_j^{m-1}\}$ is contained in all labels of degree lower than m-1. For all e in $Gr_{u_i^m v_i^{m-1}}$, define,

$$label(e) = (label(e) \setminus \{u_p^m, v_q^{m-1} | p \neq i AND q \neq j\}$$

Define G(v) as the union of all digraphs $Gr_{u_i^m v_j^{m-1}}$ and proceed to the preceding level.

In the second procedure, we rule out all edges whose labels do not comply with the rules of nested digraphs. Indeed, for all edge $v_i^k \rightarrow v_j^{k-1}$ so that $v_i^k \in label(v_i^k \rightarrow v_j^{k-1})$, select all linear digraphs whose label contain $\{v_i^k, v_j^{k-1}\}$ and let $label_{v_i^k, v_j^{k-1}}(e)$ be $label(e) \setminus \{v | v \neq v_i^k \ OR \ v_j^{k-1}\}$. Let G(v) be the union of all linear digraphs for all edges $v_i^k \rightarrow v_j^{k-1}$. Obtain the union of all nested linear sequences at level k. As we descend levels, obtain a nested digraph.

We show that we obtain a nested digraph that is the maximal nested digraph contained in the intersection of two nested digraphs that share the same root.

Proposition 3.25 (maxG(v)). Using Pseudocodes 3.24, given two nested digraphs, G(v)' and G(v)'', obtain maxG(v).

Proof. Define,

$$\begin{split} lbl_{i,j,s}(w_s^{m+2} \to v_j^{m+1}) &= label(w_s^{m+2} \to v_j^{m+1}) \cap label(u_j^{m+1} \to v_i^m) \\ label(w_s^{m+2} \to v_j^{m+1}) &= \cup_i lbl_{i,j,m}(w_s^{m+2} \to v_j^{m+1}) \\ label(u_j^{m+1} \to v_i^m) &= \cup_i lbl(w_l^{m+2} \to v_j^{m+1}) \cup \{v_j^{m+1}\} \end{split}$$

Deduce from the previous sentences that $label(u_j^{m+1} \rightarrow v_i^m)$ is the union, ranging over all s, of $lbl(w_l^{m+2} \rightarrow v_j^{m+1})$ and $label(w_s^{m+2} \rightarrow v_j^{m+1})$ is the union of all temporally labels $lbl(w_l^{m+2} \rightarrow v_j^{m+1})$ and, thus, we obtain, using the first algorithm, the union of linear digraphs whose labels are ordered by \subseteq as follows,

$$u_i^m \to v_i^{m-1} \to \dots \to v$$

with $label(u_i^r \to v_i^{r-1}) \subseteq label(u_i^{r-1} \to v_i^{mr-2}).$

Using the second pseudocode, each $G_{u_i^m v_j^{m-1}}$ is built after a selection of all linear chains so that $\{u_i^m, v_j^{m-1}\}$ is contained in the label of all edges of dept lower than m-1. $G_{u_i^m v_j^{m-1}}$ is modified by ruling out all labels of the form $\{w_s^r | (r = m \ IMPLIES \ s \neq i) \ AND \ (r = m-1 \ IMPLIES \ s \neq j\}$. In this way, we keep only the linear arrays of depth k so that it its edges of length less than m contain $\{u_i^m, v_j^{m-1}\}$ in their label, and, in that way, $G_{u_i^m v_j^{m-1}}$ agree to the rules of a nested digraph regarding the edges in a level lower than m-1.

Define G(v) as the union of all $G_{u_i^m v_j^{m-1}}$. Conclude that, as we work backward, from k = m, each step negative, and conclude, as we work at steps -1 that we obtain a nested digraph.

The search for maxG(v) is polynomially bounded, as we show in the analysis of computational boundaries in Proposition 5.6.

After writing the *Linearized Digraph*, we must filter all compatible antichains. We write a nested compatible digraph associated with each vertex v.

These digraphs associated with root v_j^m are recursively created in iteration with each vertex v_i^n , for 1 to m - 1, for $1 \le i \le k_m$. A non-empty final result generates an output *there are compatible antichains* and, otherwise, *there are no compatible antichains*. Any path in the digraph is an antichain in the linearized digraph and, reciprocally, any antichain is a path in the Linearized digraph. Write a nested digraph associated with each **edge** of this digraph from the Linearized Digraph. Edges will play the role of vertices and the arrows connecting the vertices are labeled. We will obtain labeled digraphs whose labels are the pointers of compatible antichains. We will, from the first column act from the second column until depth k^{th} and obtain a nested digraph, whose root is the vertex associated with the v_i^r vertex, $2 \le r \le m$ and $1 \le i \le l_r$.

In the h^{th} iteration, we associate to each vertex v_i^h , $h + 1 \le r \le m$ and $1 \le i \le l_h$, a nested digraph of depth h.

The search for maxG(v) is polynomially bounded, as we show in the analysis of computational boundaries in Proposition 5.6.

Procedure 3.26 (Linear Digraphs). Given a linearized digraph endowed with p branches, $Br = \{B_1, \ldots, B_k\}$, for all $1 < i \leq l$, we will write a set of nested digraphs for each vertex e_j^t , $1 < t \leq k$ and $1 \leq j \leq l_t$. The k^{th} step generates either:

Empty digraphs and an output, there are no compatible antichains or Non empty digraphs and an output, there are compatible antichains.

Step 1: For all e_i^1 in the first branch, Br_1 , for all e_j^r , $2 \le r \le l$, if e_i^1 and e_j^r are compatible, write the digraph,

$$G(e_{j}^{r}, e_{i}^{1}) = \langle \{e_{i}^{1}, e_{j}^{r}\}, \{e_{j}^{r} \rightarrow e_{i}^{1}\}, (label^{1})_{j}^{r}(e_{j}^{r} \rightarrow e_{j}^{1}) = \{e_{i}^{1}\} \rangle$$

Define $G(e_i^r)$ as the union of all $G(e_i^r, e^{1i}), 1 \le i \le k_1$.

All vertices in the columns $i, 2 \leq i \leq k$ are form an edge with the compatible vertices in the first column.

Step r: Suppose we performed all steps from 1 to r - 1 and obtained nested digraphs.

Let $G(v_j^s) = \langle V_{v_j^s}, E_{v_j^s}, label_{v_j^s} \rangle$ be nested rooted digraph of depth r-1, associated to the vertices v_j^s , $r \leq s \leq k$, $1 \leq j \leq l_s$.

For all $G(v_i^r)$, $1 \le i \le l_r$, for all v_j^s , s > r, $1 \le j \le k_s$, if v_i^r , and v_j^s are compatible, replace the (root) vertex v_i^r in $G(v_i^r)$ by v_j^s . Obtain the digraph $G(v_i^r/v_j^s)$.

- 1. Let $G(v_i^s, v_i^r)^{\prime\prime\prime}$ be the intersection of $G(v_i^r/v_i^s)$ and $G(v_i^s)$;
- 2. Let $maxG(v_j^s, v_i^r) = \langle (V_j''^s)_i^r, (E_j''^s)_i^r, (label_j'^s)_i^r \rangle$ be the maximal nested digraph contained in $G(v_j^s, v_i^r)''';$
- 3. Define $G(v_j^s, v_i^r)' = \langle (V_j'^s)_i^r, (E_j'^s)_i^r, label_0^{sjri} \rangle$ as

$$\begin{array}{ll} (V_j'^s)_i^r = & (V_j''^s)_i^r \cup \{v_i^r\} \\ (E_j'^s)_i^r = & (E_j''^s)_i^r \cup \{v \leftarrow v_i^r | v \text{ belongs to level } r-1\} \\ (label_j^s)_i^r(e) = & label'v_j^s(e) \cup \{v_i^r\} \\ (label_j^s)_i^r(v_i^r \rightarrow v) = & \{v_i^r\} \end{array}$$

Define $G(v_i^s)$ as the union of all $G(v_i^s, v_i^r)'$.

Call the set of nested antichains to the set of digraphs obtained at each vertex of level p.

Theorem 3.27. For all vertex v, the digraph G(v), 3.26, is a compatible nested digraph.

Proof. In the first step, the sequences have the form $e \to v_i^1$, if both edges are compatible. The label associated to the sequences $e \to v_i^1$ is v_i^1 . The union of such digraphs is a compatible nested digraph.

At the step r, we replace the vertex $G(v_{ri})$ by v_{sj} , intersect $G(v_i^r/v_j^s)$ and $G(v_j^s)$. By construction, $max(G(v_i^r/v_j^s) \cap G(v_j^s))$ is the biggest compatible nested digraph contained in $G(v_i^r/v_j^s) \cap G(v_j^s)$.

The digraph $G(v_j^s, v_i^r)'$ was obtained by adding the vertex v_i^r , compatible with all vertices in $G(v_i^r/v_j^s) \cap G(v_j^s)$.

Lastly, the union all $G(v_j^s, v_i^r)$ is compatible because it is the union of compatible digraphs.

Finally, highlight that we reached our claim. We show that the nested digraphs, if non-empty, contain all compatible antichains.

Definition 3.28. Call a maximal nested path over a nested digraph to any maximal path

 $v^k \to v^{k-1} \to \dots \to v^2 \to v$

so that for all $1 \leq i \leq k-2$, $label(v^i, v^{i+1}) \subseteq label(v^{i+1}, v^{i+2})$.

Identify the set of all maximal nested paths over a nested digraph with the set of maximal compatible antichains. Conclude,

Theorem 3.29. A pivoted **3-sat** Ψ is unsatisfiable if and only if the k^{th} iteration of nested antichains are empty.

4 A Pivoted Strong Version of a 3-sat Formula

Arbitrarily choose pairs of literals $S^0 = \{p_{01}, \neg p_{01}, \dots, p_{0k}, \neg p_{0k}\}$ and factorize Ψ .

$$(p_{01} \vee S_{p_{01}}) \wedge (\neg p_{01} \vee S_{\neg p_{01}}) \wedge \dots \wedge (p_{0k} \vee S_{p_{0k}}) \wedge (\neg p_{0k} \vee S_{\neg p_{0k}}) \wedge S_3^1$$

so that none of the literals of S^0 appears in any formula in the set of formulas $\{S_{p_{01}}, S_{\neg p_{01}}, \ldots, S_{p_{0k}}, S_{\neg p_{0k}}, S_3^1\}$.

Successively, do the partitions:

$$S_{3}^{1} \equiv (p_{11} \lor S_{p_{11}}) \land (\neg p_{11} \lor S_{\neg p_{11}}) \land \dots \land (p_{1k_{1}} \lor S_{p_{1k_{1}}}) \land (\neg p_{1k_{1}} \lor S_{\neg p_{1k_{1}}}) \land (\neg p_{1k_{1}} \lor S_{\neg p_{1k_{1}}}) \land (S_{3}^{2} \equiv (p_{21} \lor S_{p_{21}}) \land (\neg p_{21} \lor S_{\neg p_{21}}) \land \dots \land (p_{2k_{2}} \lor S_{p_{2k_{2}}}) \land (\neg p_{2k_{2}} \lor S_{\neg p_{2k_{2}}}) \land (S_{3}^{3} \equiv (p_{h1} \lor S_{p_{h1}}) \land (\neg p_{h1} \lor S_{\neg p_{h1}}) \land \dots \land (p_{hk_{h}} \lor S_{p_{hk_{h}}}) \land (\neg p_{hk_{h}} \lor S_{\neg p_{hk_{h}}}) \land (\neg p_{hk_{h}} \lor S_{\neg p_{hk_{h}}}) \land (S_{3} \equiv (p_{2k_{2}} \lor S_{\neg p_{2k_{2}}}) \land (S_{3} \models (p_{2k_{2}} \lor S_{\neg p_{2k_{2}}}) \land (S_{3} \land (p_{2$$

where S_{T} is either a **2** – **sat** formula or the formula T .

Finally, obtain the below partition:



We show that there is a modified pivoted **3-sat**, Ψ_T so that Ψ_T is unsatisfiable if and only if Ψ is unsatisfiable.

Let

 $\{r_{21}, \neg r_{21}, \ldots, r_{1k_1}, \neg r_{1k_1}, \ldots, r_{h1}, \neg r_{h1}, \ldots, r_{hk_h}, \neg r_{h_h}\}$ be a set of literals disjoint from the set $\mathsf{Letter}(\Psi)$. Let Ψ_T be,

$$\begin{array}{l} (p_{11} \lor S_{p_{11}}) \land (\neg p_{11} \lor S_{\neg p_{11}}) \land \cdots \land (p_{1k_1} \lor S_{p_{1k_1}}) \land (\neg p_{1k_1} \lor S_{\neg p_{1k_1}}) \land \\ (r_{21} \lor (S_{p_{21}} \land \neg p_{21})) \land (\neg r_{21} \lor (S_{\neg p_{21}} \land p_{21})) \land \cdots \land \\ (r_{2k_2} \lor (S_{p_{2k_2}} \land \neg p_{2k_2})) \land (\neg r_{2k_2} \lor (S_{\neg p_{2k_2}} \land p_{2k_2})) \land \cdots \land \\ (r_{h1} \lor (S_{p_{h1}} \land \neg p_{h1})) \land (\neg r_{h1} \lor (S_{\neg p_{h1}} \land p_{h1})) \land \cdots \land \\ (r_{hk_b} \lor (S_{p_{b1}} \land \neg p_{h1})) \land (\neg r_{hk_b} \lor (S_{\neg p_{b1}} \land p_{h1})) \land (t \lor S_{\intercal}) \land (\neg t \lor S_{\intercal}) \end{aligned}$$

Write Ψ as its factorized version,

$$\begin{array}{l} (p_{11} \lor S_{p_{11}}) \land (\neg p_{11} \lor S_{\neg p_{11}}) \land \dots \land (p_{1k_1} \lor S_{p_{1k_1}}) \land (\neg p_{1k_1} \lor S_{\neg p_{1k_1}}) \\ (p_{21} \lor S_{p_{21}}) \land (\neg p_{21} \lor S_{\neg p_{21}}) \land \dots \land (p_{2k_2} \lor S_{p_{2k_2}}) \land (\neg p_{2k_2} \lor S_{\neg p_{2k_2}}) \\ \land (\neg p_{2k_2} \lor S_{\neg p_{2k_2}}) \\ \dots \\ (p_{h1} \lor S_{p_{h1}}) \land (\neg p_{h1} \lor S_{\neg p_{h1}}) \land \dots \land (p_{hk_h} \lor S_{p_{hk_h}}) \land (\neg p_{hk_h} \lor S_{\neg p_{hk_h}}) \\ \land S_{\top} \end{array}$$

We show that Ψ_T is unsatisfiable if and only if Ψ is unsatisfiable.

Let Σ be the set of all mappings from $\{11, \ldots, 1k_1, \ldots, h1, \ldots, hk_h\}$ onto $\{True, false\}$. Note that Σ has $2^{k_1+\cdots+k_h}$ elements.

For $\sigma \in \sigma$, for all conjugated pair $\{p_{uv}, \neg p_{uv}\}$, $1 \leq u \leq h$ and $1 \leq v \leq k_v$, let $\epsilon_{uv} = p_{uv}$, if $\sigma(p_{uv}) = True$ and, otherwise, $\epsilon_{uv} = \neg p_{uv}$. Let v_{uv} be r_{uv} if $\epsilon_{uv} = \neg p_{uv}$ and, otherwise $v_{uv} = \neg r_{uv}$. Take the notation F/σ as the substituition of all literals in F by its value over σ .

Using the valuation σ , we have, respectively, for Ψ and Ψ_T ,

$$S_{\neg\epsilon_{11}}/\sigma \wedge \cdots \wedge S_{\neg\epsilon_{1k_{1}}}/\sigma \wedge S_{\neg\epsilon_{21}}/\sigma \wedge \cdots \wedge S_{\neg\epsilon_{2k_{2}}}/\sigma \wedge \cdots \wedge S_{\neg\epsilon_{2k_{2}}}/\sigma \wedge \cdots \wedge S_{\neg\epsilon_{h_{1}}}/\sigma \wedge \cdots \wedge S_{\neg\epsilon_{h_{h_{h}}}}/\sigma \wedge S_{\top}$$

$$S_{\neg\epsilon_{11}}/\sigma \wedge \cdots \wedge S_{\neg\epsilon_{1k_{1}}}/\sigma \wedge \neg v_{21} \wedge (v_{21} \vee S_{\neg\epsilon_{21}}/\sigma) \wedge \cdots \wedge \gamma v_{2k_{2}} \wedge (v_{2k_{2}} \vee S_{\neg\epsilon_{2k_{2}}}/\sigma) \wedge \neg v_{h1} \wedge (v_{h1} \vee S_{\neg\epsilon_{h1}}/\sigma) \wedge \cdots \wedge \gamma v_{hk_{h}} \wedge (v_{hk_{h}} \vee S_{\neg\epsilon_{h1}}/\sigma) \wedge S_{\top}$$

Whether at least one $\neg v_{lm}$ is *false* then, Ψ_T is false,

$$S_{\neg\epsilon_{11}}/\sigma \wedge \cdots \wedge S_{\neg\epsilon_{1k_{1}}}/\sigma \wedge \neg \upsilon_{21} \wedge (\upsilon_{21} \vee S_{\neg\epsilon_{21}}/\sigma) \wedge \cdots \wedge \\ \neg \upsilon_{2k_{2}} \wedge (\upsilon_{2k_{2}} \vee S_{\neg\epsilon_{2k_{2}}}/\sigma) \wedge \neg \upsilon_{h1} \wedge (\upsilon_{h1} \vee S_{\neg\epsilon_{h1}}/\sigma) \wedge \cdots \wedge \\ \neg \upsilon_{hk_{h}} \wedge (\upsilon_{hk_{h}}/\sigma \vee S_{\neg\epsilon_{h1}}/\sigma) \wedge S_{\mathsf{T}}$$

Otherwise, all $\neg v_{lm}$ is **true** and Ψ_T is

$$S_{\neg\epsilon_{11}}/\sigma \wedge \cdots \wedge S_{\neg\epsilon_{1k_{1}}}/\sigma \wedge S_{\neg\epsilon_{21}}/\sigma \wedge \cdots \wedge S_{\neg\epsilon_{2k_{2}}}/\sigma \wedge S_{\neg\epsilon_{h1}}/\sigma \wedge \cdots \wedge S_{\neg\epsilon_{h1}}/\sigma \wedge S_{\top}$$

and, under this conitions of valuation, Ψ is unsatisfiable if and only if Ψ_T is unsatisfiable.

5 Bounds on Computation

Open this section with the study of the bounds in Space, $\mathbf{SPACE}(f(n))$ and in time, $\mathcal{O}(f(n))$. Both bounds ensure that we do not trade space by time or vice-versa in our considerations.

Given a card \mathcal{C} , we follow the polynomially bounded steps

- 1. Write the cylindrical digraph;
- 2. Write the closed digraphs;
- 3. Write the linearized digraph;
- 4. Write the maximal nested digraphs. A special care has to be taken on writing the maximal nested digraph contained in an intersection.

We prove less straightforward bounds in time and space.

Lemma 5.1. The cylindrical digraph, $\mathbb{C}lndr = \langle V, E, label \rangle$ is written in polynomial time and space.

Proof. The size of $\mathbb{C}lndr$ -graph is given by

- 1. The size of V, the set of vertices is the size of literals, L;
- 2. The size of E is bounded by the square of the number of literals, $L |E| \leq |L|^2$. Indeed, the arrows are bounded by the number of arrows in a polyhedron with |L| sides.

There is an algorithm polynomially bounded in time to write a closed digraph. We describe the polynomial in time search for nonempty intervals $[a, \neg a]$. Recall that no loops are allowed. Indeed, if we write a loop, that means that if an incompatible combination $Inc = ij_1, \ldots, ij_r$, any combination that contains Inc is likewise incompatible.

Divide the search algorithm into two procedures,

- 1. Write [a, q[, the subdigraph of CSD of all sequences connected to a (no specified end, just source, a);
- 2. If $[a, q[\neq \emptyset, \text{ from } [a, q[, \text{ write } [a, \neg a], \text{ the subdigraph of paths between } a \text{ and } \neg a \text{ contained in } [a, q[.$

Part I, write an Interval [a, q[

Input cylindrical digraph
$$Cyl = \langle V, E, label \rangle$$
.
 $[a,q[= \langle \{a\}, \emptyset, \emptyset \rangle = \langle V_{[a,q[}, E_{[a,q[}, label_{[a,q[}) \rangle \\ E_{aux} = E \smallsetminus E_{[a,q[} \rangle \end{pmatrix} = \langle V_{[a,q[}, L_{[a,q[}, label_{[a,q[}) \rangle \end{pmatrix}$
while $E_{aux} \neq \emptyset$ do
 $E_{[a,q[} \leftarrow E_{[a,q[} \cup \{c \Rightarrow b|c \in V_{[a,q[} AND \ c \Rightarrow b \in E \rangle \\ V_{[a,q[} \leftarrow V_{[a,q[} \cup \{b|\exists c \in V_{[a,q[}(c \Rightarrow b \in E_{aux})) \} \\ label_{[a,q[} \leftarrow label_{[a,q[} \cup \{label(c \Rightarrow b)|\exists c \in V_{[a,q[}(c \Rightarrow b \in E_{aux})) \} \\ E_{aux} = E \smallsetminus E_{[a,q[}$
end while

end while

Once we obtain $[a,q[=\langle V_{[a,q[},E_{[a,q[},label_{[a,q[}\rangle), \text{ if } \neg a \in V_{[a,q[}, \text{ we perform})$ the above algorithm to [a, q], taking care to search over arrows $b \Rightarrow \neg a$ and obtain $[a, \neg a]$.

The set of digraphs $[a, \neg a]$ is bounded by the number of vertexes in the cylindrical digraph and each $[a, \neg a]$ has its set of vertices and labels bounded by the number of vertices and labels in the cylindrical digraph and, therefore, we wrote no more than |E| digraphs that are bounded by |E|vertices and $|E|^2$ edges.

Observation 5.2. Once we work iteratively with $E \\ E_{[a,q]}$, we avoid sequences of the form,

 $r \Rightarrow q_1 \Rightarrow \cdots \Rightarrow r$

that is, we do not have loops.

Let us scrutinize the size of the closed digraph.

Lemma 5.3. The set of closed digraphs is the union of at most |L| digraphs with $|L|^2$ vertices and $|L|^2$ edges.

Proof. The set of necessarily true pairs has, at most, the size of |L|, the number of literals. Each interval $[\neg p, p]$ has its size bounded by the size of the closed digraphs, that is, at most |L| vertices and $|L|^2$ edges.

Obtain the union of, at most, |L| digraphs with |L| vertices and $|L|^2$ edges, **SPACE**($|L|^3$).

Lemma 5.4. The search for the set of closed digraphs is polynomial in time and space.

Proof. We perform at most |L| searches, one to each interval $[p, \neg p]$ for $p \in L$. Our search goes over the number of vertices

$$egin{array}{l} |V| - 1 \ |V| - 1 - r_1 \ dots \ |V| - 1 - r_1 \ dots \ |V| - 1 - r_1 - \cdots - r_q \end{array}$$

under the conditioning $1 + r_1 + \cdots + r_q = V$, which leads to a maximum $r_1 = \cdots = r_q = V/q$ and the search goes to a top of $|V|^2$ searches over |V| vertices and the search is bonded by $|V|^3$ both in time and space.

Our process continues analyzing the effort to Lift, Multiply Branches and the Addition of Label. After that, we must analyze the whole process of writing the nested digraphs associated with each edge.

Proposition 5.5. Given a closed digraph, the operations of Lifting, Adding Labels and Multiplying Branches are linearly bounded.

Proof. Lifting together with Multiplication of branches multiply the number of any branching a accordingly to the maximum of branches in Up(a) or Down(a).

Adding Label adds several new labels to, at most the number of edges in the closed digraph its addition is bounded by the number of branching and Multiplication of branches is bounded by the maximum number of branches.

Proposition 5.6. Given two nested rooted digraphs, the operation of writing the maximal nested digraph contained in their intersection is linearly bounded.

Proof. In the first algorithm, the union of linear digraphs whose labels are ordered by \subseteq . We operate over a maximum of $|V|^2$ edges in a level m and each edge can interact with $|V|^2$ edges in a level m + 1. Obtain at most $|V|^2 \times |V|^2$ interactions over at most |V| levels, that is, a maximum of $|V|^5$ interactions.

The worst bound in the above operation will cost us a visit in at most |V| edges where we use simple operations to obtain $lbl(v_j^{m+1} \rightarrow v_i^m)$ and the intersection of the labels (in the intersection) of $label(v_j^{m+1} \rightarrow v_i^m)$ and $label(v_j^{m+2} \rightarrow v_j^{m*1})$,

In the second pseudocode, we write, for any edge $u_i^m \to v_j^{m-1}$ the tree of all edges starting with edges $v_j^{m-1} \to v_s^{m-2}$ that have $\{u_i^m, v_j^{m-1} \text{ in their}$ label. Proceed in descending order until reaching the root. Prune edges that don't reach the root, that is, edges of the form $v \to w$ so that there is no edge $w \to u$.

Each edge visits, in a maximum of $|V|^2$ edges. We have, in the worse case, are bounded by the number of closed digraphs, an use of dimension $|L|^3$. so, we search $|V|^5$ in time.

6 Examples

Example 6.1. Consider the CSD,



We do not focus on the compatibility among vertices. It is not relevant for this analysis.

The task of writing the chains is clearly expsize, as we show below

p_1	p_2														
p_3	p_3	p_3	p_3	p_4	p_4	p_4	p_4	p_3	p_3	p_3	p_3	p_4	p_4	p_4	p_4
p_5	p_5	p_6	p_6	p_5	p_5	p_6	p_6	p_5	p_5	p_6	p_6	p_5	p_5	p_6	p_6
p_7	p_8	p_7	p_8	p_7	p_8	p_7	p_8	p_7	p_8	p_7	p_8	p_7	p_8	p_7	p_8
	Solve	usin	g ou	r me	thod.	Mu	ltiply	root	s,						



Add labels, obtain,



Lifting,



Add labels, obtain,



Lifting,

$p_1 \Longrightarrow$			
			$b \neg c \neg a \neg b \neg d$
p_3	p_4	p_3 Z	p_4
$\neg ac$	$\neg bd$	$\neg bc$	$\neg ad$
p_5	p_6	p_6	p_5
П	п	п	н
ac	bd	bc	ad
v	¥	¥	•
p_7	p_7	p_8	p_8

Lifting,

p_1	p_1	p_2	p_2
$\neg a \neg b \neg c$	$\neg a \neg b \neg d$		
p_3	p_4	p_3	p_4
$\neg ac$	$\neg bd$	$\neg bc$	$\neg ad$
p_5	p_6	p_6	p_5
II ac ▼	bd	bc	ad •
p_7	p_7	p_8	p_8

The only possible compatible combinations, accordingly to compatibility of labels we added, besides computing original compatibility, is,

 $\begin{array}{l} p_5 \Rightarrow p_7, p_5 \Rightarrow p_8, p_6 \Rightarrow p_7, p_6 \Rightarrow p_8 \\ p_5 \Rightarrow p_7, p_4 \Rightarrow p_6, p_3 \Rightarrow p_6, p_5 \Rightarrow p_8 \\ p_3 \Rightarrow p_5, p_6 \Rightarrow p_7, p_6 \Rightarrow p_8, p_4 \Rightarrow p_5 \\ p_3 \Rightarrow p_5, p_4 \Rightarrow p_6, p_3 \Rightarrow p_6, p_4 \Rightarrow p_5 \\ p_1 \Rightarrow p_3, p_1 \Rightarrow p_4, p_2 \Rightarrow p_3, p_2 \Rightarrow p_4 \end{array}$

Example 6.2. Consider the two pivoted formulas,

$$\Psi_{1} = a_{1} \vee ((p_{1} \vee q_{1}) \wedge (p_{1} \vee \neg q_{1})) \wedge \neg a_{1} \vee ((p_{1} \vee q_{1}) \wedge (p_{1} \vee \neg q_{1})) \wedge \\ a_{2} \vee ((\neg p_{1} \vee q_{2}) \wedge (\neg p_{1} \vee \neg q_{2})) \wedge \neg a_{2} \vee ((p_{1} \vee q_{2}) \wedge (p_{1} \vee \neg q_{2})) \\ \Psi_{2} = a_{1} \vee ((p_{1} \vee q_{1}) \wedge (\neg p_{1} \vee q_{2})) \wedge \neg a_{1} \vee ((p_{1} \vee q_{1}) \wedge (\neg p_{1} \vee q_{2})) \wedge \\ a_{2} \vee ((p_{1} \vee \neg q_{1}) \wedge (\neg p_{1} \vee \neg q_{2})) \wedge \neg a_{2} \vee ((p_{1} \vee \neg q_{1}) \wedge (\neg p_{1} \vee q_{2}))$$

The formulas Ψ_1 and Ψ_2 are, respectively, unsatisfiable and satisfiable. Write a fragment of the closed digraph,

$$q_{1} \overset{q_{1}}{\underset{C}{\overset{e}{\underset{p_{1}}{\overset{p_{1}}{\underset{p_{1}}{\overset{e}{\underset{p_{1}}{\overset{p_{1}}{\underset{p_{1}}{\overset{p_{1}}{\underset{p_{1}}{\overset{p_{1}}{\underset{p_{1}}{\overset{p_{1}}{\underset{p_{1}}{\overset{p_{1}}{\underset{p_{1}}{\overset{p_{1}}{\underset{p_{1}}{\overset{p_{1}}{\underset{p_{1}}{\underset{p_{1}}{\overset{p_{1}}{\underset{p_{1}}{\overset{p_{1}}{\underset{p_{1}}}{\underset{p_{1}}{\underset{p_{1}}}{\underset{p_{1}}}{\underset{p_{1}}{\underset{p_{1}}{\underset{p_{1}}{p_{1}}{\underset{p_{1}}{\underset{p_{1}}{\underset{p_{1}}{\underset{p_{1}}{p$$

In the case of Ψ_1 , the labels in A, B, C and D are $a_1, \neg a_1$ and E, F, G and H, are $a_2, \neg a_2$.

In the case of Ψ_2 , the labels in are shown below, $A = a_1, \neg a_1, B = a_2, \neg a_2, C = a_2, \neg a_2, D = a_1, \neg a_1 E = a_1, \neg a_1, a_2, \neg a_2, F = a_2 G = a_2 H = a_1, \neg a_1, \neg a_2.$

After lifting and adding labels, we obtain,

$\neg p_1$	$\neg p_1$
Ш	11
$A \neg r$	$B\neg r$
V	V
q_1	$\neg q_1$
Π	Π
$C \neg r$	$D\neg r$
V	V
p_1	p_1
Π	П
Er	Fr
V	V
q_2	$\neg q_2$
Π	П
Gr	Hr
V	▼
$\neg p_1$	$\neg p_1$

Case Ψ_2 , there are compatible combinations and, in the case of Ψ_1 , there is no compatible combination. We can infer Ψ_1 is not satisfiable and, in the case of Ψ_2 , the conclusion is based in writing all combinations.

References

- Cook S.A. Reckhow A.R. The relative efficiency of propositional proof systems. J. of Symbolic Logic, 44(1):36–50, 1979.
- [2] B. Arora, S. Barak. Computational Complexity: A Modern Approach. Cambridge University Press, 2009.
- [3] Plass Aspvall and Tarjan. A linear time algorithm for testing the truth of certain quantified boolean formulas. *Information Processing Letters*, pages 121–123, 1979.
- [4] Papadimitriou C. H. Computational Complexity. Addison-Wesley, 1994.
- [5] Juris Hartmanis and Richard Edwin Stearns. On the computational complexity of algorithms. *Transactions of the American Mathematical Society*, 117:285–306, 1965.
- [6] Stockmeyer L. J. Classifying the computational complexity problems. The Journal of Symbolic Logic, 52 No 1:1–43, 1987.

- [7] Leonid Levin. Universal sorting problems. problems of information transmission. *Problemy Peredachi Informatsii*, 9:265–266, 1973.
- [8] Leonid Levin. universal search problems. Problemy Peredachi Informatsii, 6:265-266, 1984.
- [9] A. R. Meyer and L. J. Stockmeyer. The equivalence problem for regular expressions with squaring requires exponential space. In 13th Annual Symposium on Switching and Automata Theory (swat 1972), pages 125– 129, 1972.
- [10] Rabin M. O. Degree of difficult of computing a functionand a partial ordering of recursive sets. Technical report, Jerusalem, 1960.
- [11] Michael Oser Rabin. Speed of computation and classification of recursive sets. In *Third Convention Sci. Soc.*, Israel, pages 1–2, 1959.
- [12] Cook S.A. The complexity of theorem proving procedures. Proc 3rd STOC, 1971.
- [13] A. M. Turing. On computable numbers, with an application to the entscheidungsproblem. *Proceedings of the London Mathematical Soci*ety, s2-42(1):230-265, 1937.
- [14] Maria Angela Weiss. A polynomial decision for 3-sat. 2022. arXiv:2208.12598.

Index

Ch(e), 8 $Ch\mathcal{P} \mapsto \mathcal{CSD}, 8$ $Down_a, 15$ Gr(B), 14 $Gr(B \star R) = \langle E_{B \star R}, E_{B \star R}, label \rangle, 13$ Gr(R), 14Linclsd, 17 Proj(e'), 17 $Proj: Linclsd \rightarrow CSD, 17$ $Up_a, 15$ Σ , 18 Θ , 18 $\mathbb{C}lndr = \langle V, E, label \rangle, 5$ CSD, 7 $\mathcal{NEC}, 7$ $\mathcal{P}, 7$ maxG(v), 23branch, 13 3-sat formula, 4 **2-sat** formula, 3 addition of the new pair of literals to the set of branching, 15 antichain, 9 branching, 12 chain, 7 chains, 7 clause, 4 Companion Digraph, 21 compatible choice, 8 compatible nested digraph of length p, 23compatible set of entries, 8 complete 3-sat, 5 cylindrical digraph, 5

edge-labeled digraph for v, 22 entails, 9 entries, 4 entry, 6 expansion to \mathcal{P} , 8 incompatible, 8 intersection of digraphs, $Gr_v \cap Gr'_v$, 23interval $[p,q], \tilde{\gamma}$ involves, 18 label of the vertex v, 22 Lifting by B and R, 13 Lifting of Gr by B, 14 Lifting of Gr by R, 14 linear interval, 13 Linearized Closed Digraph, 17 maximal nested path over a nested digraph, 28 maximal branch, 12 Multiplication of a branch br, 14 necessarily true literals, 7 necessarily true literals, 7nested digraph of length p, 23 path, 6Pivot, 4 Pivoted 3-sat formula, 1, 4 root interval, 13 roots, 12 satisfiable, 3

set of closed digraph, 7 set of entries of $\neg a_i$, 4 set of entries of a_i , 4 set of nested antichains, 27

tops, 12

union of digraphs, 22 unsatisfiable, 3