Ortho-unit polygons can be guarded with at most $\lfloor \frac{n-4}{8} \rfloor$ guards

J.M. Díaz-Báñez * P. $Horn^{\dagger}$ M.A. Lopez [‡]

O. Solé-Pi [∥] A. Stevens^{**} J. Urrutia ^{††}

N. Marín[§]

A. Ramírez-Vigueras[¶]

August 30, 2022

Abstract

An orthogonal polygon is called an ortho-unit polygon if its vertices have integer coordinates, and all of its edges have length one. In this paper we prove that any ortho-unit polygon with $n \ge 12$ vertices can be guarded with at most $\lfloor \frac{n-4}{8} \rfloor$ guards.

1 Introduction

A (simple) polygon P is a sequence of n vertices v_1, \ldots, v_n and n edges $v_1v_2, v_2v_3, \ldots, v_{n-1}v_n, v_nv_1$ such that no pair of non consecutive edges share a point. In this paper, and following standard notation in the Art Gallery literature, we will refer to P as the region enclosed by the union of the vertices and edges of P; the boundary of P, denoted as ∂P , consists of the edges and vertices of P. Following O'Rourke's terminology [14], we will assume that $\partial P \subset P$. A point $p \in P$ guards a point $q \in P$ if the line segment pq joining them is contained in P. A set S of points in P guards P if any point in P is guarded by at least one point in S. A polygon is called an *orthogonal polygon* if its edges are horizontal or vertical line segments.

In this paper we will study guarding problems for *ortho-unit polygons*, that is orthogonal polygons all of whose edges have length one. Our main result is that $\lfloor \frac{n-4}{8} \rfloor$ points are always sufficient and sometimes necessary to guard any ortho-unit polygon.

**Department of Computer Science, University of Denver.

^{*}Departamento de Matemática Aplicada II, Universidad de Sevilla, Spain. Partially supported by Ministerio de Ciencia e Innovación CIN/AEI/10.13039/501100011033/ (PID2020-114154RB-I00) and European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement #734922.

[†]Department of Mathematics, University of Denver. Supported in part by Simons Collaboration Grant 525039

[‡]Department of Computer Science, University of Denver, mario.lopez@du.edu. Supported in part by a University of Denver John Evans Research Award.

[§]Posgrado en Ciencia e Ingeniería de la Computación, Universidad Nacional Autónoma de México. nestaly@ciencias.unam.mx. ORCID: 0000-0002-0222-3254. Supported by SEP-CONACyT of Mexico.

[¶]Instituto de Matemáticas, UNAM. adriana.rv@im.unam.mx. Partially supported by PAPIIT IN105221, Programa de Apoyo a la Investigación e Innovación Tecnologíca UNAM.

[|]Departamento de Matemáticas, Facultad de Ciencias, UNAM. oriolandreu@ciencias.unam.mx.

^{††}Instituto de Matemáticas, Universidad Nacional Autónoma de México, urrutia@matem.unam.mx. ORCID: 0000-0002-4158-5979. Supported in part by PAPIIT IN102117 Programa de Apoyo a la Investigación e Innovación Tecnológica, UNAM.

2 Previous work

The study of *guarding* or illumination problems of polygons on the plane is a classic area of research in computational geometry. In 1973 V. Klee posed the problem of determining the minimum number of guards needed to guard any art gallery whose boundary is represented by a polygon (Hosenberg 1976 [9]). Klee's problem was solved by Chvátal [4], who proved that $\lfloor \frac{n}{3} \rfloor$ guards are always sufficient and sometimes necessary. A short and elegant proof was later given by Fisk [6] using triangulations of polygons.




Figure 1: (a) is a family of polygons in which $\lfloor \frac{n}{3} \rfloor$ guards are required. (b) is a family of orthogonal polygons in which $\lfloor \frac{n}{4} \rfloor$ guards are required.

For orthogonal polygons, Kahn, Klawa and Kleitman's [10] proved that $\lfloor \frac{n}{4} \rfloor$ points are always sufficient to guard any orthogonal polygon with *n* vertices. This bound is also tight. Since then a plethora of papers studying several variations to Klee's original question have appeared; see [14, 15, 18].

Orthogonal polygons have been widely studied, among other reasons, since they model in a more realistic way traditional buildings, which are better modeled by orthogonal polygons. To our knowledge there are at least six different proofs of the Orthogonal Art Gallery Theorem, e.g. see [7, 13, 16, 17]. Hofman and Kriegel [8] proved that any orthogonal polygon with holes can always be guarded with $\lfloor \frac{n}{3} \rfloor$ vertex guards. Bjorling-Sachs proved that $\lfloor \frac{3n+4}{6} \rfloor$ edge guards are always sufficient and sometimes necessary to guard an orthogonal polygon [3]. Katz and Roisman [12] studied the problem of guarding the vertices of an orthogonal polygon is NP-hard.

The guarding problem for polyominoes, orthogonal polygons that are obtained by joining sets of unit squares, was studied by Biedl et al. [2]. They proved that $\lfloor \frac{m+1}{3} \rfloor$ point guards are always sufficient and sometimes necessary to guard a polyomino with area m, possibly with holes. Katz and Morgenstern [11] studied the problem of guarding an orthogonal art gallery with cameras that slide along horizontal or vertical straight tracks. They proved that if only horizontal tracks are allowed, then finding the minimum number of tracks needed can be found in polynomial time. Durocher et al. [5] extended Katz and Morgenstern results by proving that finding a set of tracks that minimizes the sum of their lengths can also be solved in polynomial time, even for orthogonal polygons with holes.

Biedl et al. [1] studied the problem of guarding orthogonal polygons with sliding k-transmitters, a point guard that slides along an axis line segment s. A k transmitter can see s point $p \in P$ if the perpendicular line segment joining it to S intersects the boundary of P at most k times.

The interested reader can find more information in a book by O'Rourke [14] and two survey papers, by Shermer [15] and Urrutia [18].

3 Guarding ortho-unit polygons 1

In this section we prove our main result, namely:

Theorem 1. Any ortho-unit polygon with $n \ge 12$ vertices can be guarded with at most $\lfloor \frac{n-4}{8} \rfloor$ guards. Moreover, $\lfloor \frac{n-4}{8} \rfloor$ guards is sometimes necessary.






Figure 2: (a) is a short 'macuahuitl' polygon, with 28 sides, requiring 3 guards. (b) is an illustration of extending a 'macuahuitl' polygon. This requires 8 additional sides and one additional guard. Both polygons require $\frac{n-4}{8}$ guards.

Suppose P is an ortho-unit polygon with $n \ge 12$ vertices. In order to prove that $\lfloor \frac{n-4}{8} \rfloor$ guards suffice to guard P we first introduce an auxiliary structure to help us find appropriate guard positions, the *column tree*.

A vertical (resp. horizontal) cut of P is the extension of a vertical (resp. horizontal) edge of P at a reflex vertex towards the interior of P. Because P is ortho-unit, each vertical cut connects two reflex vertices. As observed by O'Rourke in [14], there are $\frac{n-4}{2}$ reflex vertices in any orthogonal polygon and hence there are $\frac{n}{4} - 1$ vertical cuts. They, in turn, partition P into $\frac{n}{4}$ one wide rectangles, which we call columns. Note that the top and bottom of each column are horizontal unit length edges.





(a) Partition of the polygon into columns. Columns labeled a, c, d, j, k, l are called teeth.




Figure 3: When two columns share a border in figure (a), their corresponding vertices share an edge in figure (b).

The column tree of P, denoted by T, is the tree whose vertices are the $\frac{n}{4}$ columns of P, where two vertices of T are joined by an edge if they are separated by a vertical cut. Note that a guard placed on a vertical cut will guard both incident columns. A simple observation, then, is that a perfect matching in T would yield a set of $\frac{n}{8}$ guards which guard all of P: each of the edges in the matching corresponds to a vertical cut, and placing a guard on each of these vertical cuts would result in a set of guards which guard the entire polygon. Of course, T will typically not contain a perfect matching but this observation is our starting point, nonetheless.

The following proposition, which follows easily by induction, reveals that leaves of T are the barrier to finding a perfect matching. We state it in a form which will be convenient to us later in the proof.

Proposition 2. Suppose T is a tree, and $x \in V(T)$ is a leaf. Then there is a matching in T

which saturates x and all interior vertices of T.

Because P is ortho-unit, leaves in T correspond to columns in P that are one unit high, and have edges to their left or right. We call such structures *teeth* in P; teeth can be either left- or right-facing depending on whether they have edges to their right or left side, respectively. As a second auxiliary structure, we create the *tooth graph* H of P as follows: the vertices of H are the teeth of P. Two teeth t_1 and t_2 are connected by an edge in H if the horizontal cut from either the top or bottom edge of t_1 coincides with a horizontal cut from the top or bottom edge of t_2 . Equivalently, two teeth are connected by an edge if a single guard can simultaneously view both.

Lemma 3. The components of H consist of paths and isolated vertices.

Proof. The maximum degree of a tooth in H is clearly two as there are only two horizontal cuts incident to it, one each coming from its top and bottom edge. Since the teeth have height one, either *both* the top and bottom horizontal cuts of two teeth coincide, or the top cut of one tooth coincides with the bottom cut of its neighbor. This immediately rules out cycles, and so each component is a path.

An *interior edge* of T is an edge of T which is not incident to a leaf. Each interior edge corresponds to a vertical cut which is *not* incident to a tooth. We say that an interior edge of T can guard a tooth if a guard can be placed on the corresponding vertical cut in such a way that it guards the tooth.

The following lemma is the key to efficiently guarding the teeth of the polygon.

Lemma 4. Suppose P is an ortho-unit polygon. Let O denote the set of odd components of the tooth graph H of P. Then there is a matching between O and the interior edges of T which saturates O and so that each matched interior edge can guard a leaf in the corresponding matched component.






Figure 4: Above is a matching of teeth. We see that c and g are matched, as are b and e and, a and d. f is unmatched.

Proof. Components in O are either isolated teeth or odd length paths of teeth. Note that the odd length paths can be arranged vertically, and there is a top-most and bottom-most edge of the top and bottom teeth of the path.

We build a bipartite graph where one partite set consists of components in O and the other partite set consists of interior edges of T as follows. Fix a component x in O. This component has a top-most and bottom-most horizontal edge (if the component consists of only one tooth, then these are the top and bottom edge of that tooth, otherwise they lie in the top and bottom tooth respectively. Consider the horizontal cut defined by the top-most edge; this intersects Pagain at a reflex vertex, r. Because this is the topmost edge in a component of H, r does not lie on a tooth. Hence, the vertical cut through r corresponds to an internal edge e of T. We connect the component x to the internal edge e, and similarly connect x to the internal edge obtained in a similar way using the bottom edge.





(a) A matching of internal edges and teeth using Hall's Theorem.

(b) A representation of the matching in the column tree. Bold edges are unmatched.


The internal edges have maximum degree two, their potential neighbors being determined by the horizontal cuts through the reflex vertices at the top and bottom. Meanwhile, the odd components have degree *exactly* two, unless the top and bottom of the component connect to the same interior edge (and hence the component is the only neighbor of the edge.) The desired matching, then, is an easy consequence of Hall's theorem. That the edge guards a tooth it is matched to is a consequence of the fact that the horizontal cut coming from one of the reflex vertices ending the vertical cut corresponds with a horizontal cut coming from the tooth. \Box

We think of the matching guaranteed by a Lemma 4 as a matching between leaves of T (teeth) and internal edges so that one of the endpoints of each odd component of H is matched.

Now we are ready to turn to the meat of the proof of Theorem 1. For our ortho-unit polygon P, consider a matching guaranteed by Lemma 4. As several matchings are considered in the proof, we call this one the \mathcal{L} -matching. While the \mathcal{L} -matching the proof provides may match teeth with internal edges far from them, it is easier to deal with a 'closer' matching. To that end, we consider a matching satisfying the conclusion of the lemma that minimizes the sum of distances from the teeth (leaves) to the matched internal edges. Such a matching has the property that all edges between a tooth and its matched internal edge are also matched (as otherwise there is a 'better' matching.) Fix such a minimal \mathcal{L} -matching; we will work with it for the remainder of the proof.

Place guards on the vertical cuts corresponding to \mathcal{L} -matched internal edges in such a way that they also guard their matched tooth. Let \mathcal{F}' be the forest spanned by \mathcal{L} -matched internal edges in T, and \mathcal{F} be the forest obtained by removing all vertices in \mathcal{F}' along with the \mathcal{L} -matched teeth. The vertices in \mathcal{F} , then, correspond to columns that still need to be guarded. Denote by H' the graph obtained by removing all \mathcal{L} -matched teeth from the tooth graph H. Note that H' consists solely of even paths, and hence contains a perfect matching, which we call the \mathcal{T} -matching.

Lemma 5. Let X be a set of internal edges which span a component of \mathcal{F}' . Then the guards placed on the corresponding vertical cuts guard a total of 2|X|+1 columns of P, including teeth.



 $\begin{array}{c} & & & \\ & & \\ d & e & f \\ & & \\ & & \\ & & \\ & & \\ & & \\ a & c \end{array}$

(a) A matching which minimized the sum of distances from the teeth to the matched internal edges.

(b) A representation of the matching in the column tree. Edges shown bold are unmatched.

Proof. Each guard sees both columns incident to the vertical cut it sits on. Since the internal edges span a tree whose vertices are these columns, this is precisely |X|+1 columns. Furthermore each guard also guards the tooth \mathcal{L} -matched to its internal edge. Since all edges are internal, these are disjoint from the columns already counted and 2|X| + 1 columns are guarded in total.

Now we need to guard the remaining columns corresponding to vertices in \mathcal{F} . To this end, we fix a root in T. If \mathcal{F}' is non-empty, we choose the root to be a leaf \mathcal{L} -matched to an internal edge in \mathcal{F}' . If all components in H are even, then \mathcal{F}' is empty; in this case, we choose the root to be an arbitrary leaf. We now define, for every component of \mathcal{F} , its root to be the vertex in the component closest to the root of T.

Let ℓ be a leaf of a component of \mathcal{F} . We say ℓ blames a component C of \mathcal{F}' if ℓ is the first vertex outside of C on the shortest path from C to the root in T. From the definition, one immediately has

Lemma 6. No component C of \mathcal{F}' is blamed by more than one leaf ℓ .

On the other hand,

Lemma 7. Suppose ℓ is a leaf or isolated vertex in \mathcal{F} that is not a leaf in T. If ℓ is isolated or not the root of its component, then ℓ blames some component C in \mathcal{F}' .

Proof. Fix a vertex ℓ satisfying the hypothesis of the lemma. We claim that ℓ is incident to multiple internal edges of T. If T were a star, then it is easy to see that P is a vertically oriented double comb, and actually in this case the center vertex is not a leaf in \mathcal{F} (which in the case is all of T, as the components of H in this case are all edges.) Thus ℓ is certainly incident to at least one internal edge of T.

If ℓ were only incident to one internal edge of T – which is not \mathcal{L} -matched to any tooth, as ℓ is in \mathcal{F} and not \mathcal{F}' – then it must also be adjacent to some collection of leaves. Since ℓ is a leaf in \mathcal{F} these leaves must be \mathcal{L} -matched. Then the corresponding tooth could be guarded by the internal edge incident to ℓ . Our choice of \mathcal{L} -matching then rules out this case, as we could replace the \mathcal{L} -matched internal edge of some tooth by the internal edge incident to ℓ and reduce the total distance of the \mathcal{L} -matching.

Thus ℓ is incident to at least two internal edges of T. One of these is not in \mathcal{F} and, since ℓ is either isolated in \mathcal{F} or not the root of its component, not on the shortest path to the root. The other end of this edge is a vertex which is in \mathcal{F}' , and ℓ blames that vertex's component as desired.




Figure 7: Suppose the column tree is rooted at a. The red, blue, yellow and green subtreees are the components of \mathcal{F}' . The vertices f, g, s and t are in \mathcal{F} . While t blames the red component, g blames both the green and blue components. The yellow component remains unblamed.

We are now ready to complete the proof.

Proof of Theorem 1. It remains to guard the vertices of \mathcal{F} . For each non-trivial component, we apply Proposition 2 to obtain a matching in \mathcal{F} , the \mathcal{F} -matching, which covers all interior vertices of the component and the root of each component (taking that root to be x in Proposition 2 if the root is a leaf). We place guards on each of the vertical cuts corresponding to these matched edges. At this point, we have placed guards on vertical cuts the edges of \mathcal{F}' and the \mathcal{F} -matched edges of \mathcal{F} . The only unobserved columns now correspond to isolated vertices and unmatched leaves in \mathcal{F} .

For each unmatched leaf or isolated vertex in \mathcal{F} that is not a leaf in T we place a guard dedicated to guarding the associated column.

The remaining leaves correspond to teeth in P and are handled as follows: fix a perfect \mathcal{T} -matching in H', the tooth graph consisting of teeth that were not matched in the first step. For each edge of this \mathcal{T} -matching, we add guards as follows:

- 1. If both teeth are saturated in the *F*-matching we do nothing, as both teeth are already guarded.
- 2. If one tooth is saturated in the \mathcal{F} -matching but one is not, note that the guard can be placed in such a way that it actually guards *both* teeth simultaneously, while still guarding the columns it was originally responsible for.
- 3. If neither tooth is saturated in the *F*-matching, place a single guard responsible for both teeth.

By this construction all columns are guarded. Furthermore, on average each guard is responsible for at least two unique columns. This follows as the only guards initially assigned a single column are those which, by Lemma 7 blame some component of \mathcal{F}' . Per Lemma 5, the guards guarding the columns of a component of \mathcal{F}' guard, in the aggregate, an extra column beyond the claimed average of two each. Assigning this 'extra' column to the guard only assigned a single column which blames that component gives the claimed average.

This suffices to prove that at most n/8 guards can guard P. To prove that, actually, at most (n-4)/8 guards suffice, we proceed as follows.

If \mathcal{F}' is non-empty, note that the root of T is incident to an internal vertex x of T that by our choice of matching is in \mathcal{F}' . The component of x is not blamed by any leaf and hence the 'extra' column is never assigned. Thus, if g guards are placed 2g + 1 is a lower bound for the number of distinct columns guarded so that $2g + 1 \le \frac{n}{4}$; rearranging $g \le \frac{n-4}{8}$ as desired.

If \mathcal{F}' is empty, which occurs if all components of H are even, there is an 'extra' column assigned to a guard if, when considering the \mathcal{T} -matching, case (b) above ever occurs.

If case (b) never occurs, take an alternating path in the matching on \mathcal{F} starting from some leaf ℓ in T. This ends at some other leaf of T, ℓ' .

If ℓ and ℓ' are not paired in the \mathcal{T} -matching with each other, interchange the edges of the alternating path to obtain a new matching. For this matching it is easy to see that case (b) will occur and the matching satisfies all other desired conditions.

If ℓ and ℓ' are paired in the \mathcal{T} -matching, ensure that we are in case (c) above, possibly interchanging the edges of the alternating path to obtain a new \mathcal{F} -matching. Note that the path between ℓ and ℓ' in T then contains a matched internal edge, and a guard can be placed on this internal edge in such a way that it sees not only the incident columns but also guards *both* the teeth corresponding to both ℓ and ℓ' .

This assignment of guards to columns ensures that there is at least one more than twice the number of guards columns guarded, and hence yields the desired bound.

4 Integral Polygons

An orthogonal polygon is an *integral orthogonal polygon* if all of its edges have integer length. The lower bound on the number of guards needed to guard this family of polygons is given by the integral version of the well-known orthogonal comb polygon which requires n/4 guards. See Figure 8 for an example.




Figure 8: An integral polygon P of perimeter N which requires N/6 guards. The elements of a guard set for P are shown in red.

We can prove, via an unenlightening and complicated proof (not included in this paper), that integral orthogonal polygons of perimeter N can be guarded using at most $\lfloor N/5 \rfloor$ guards. Nevertheless, we conjecture that N/6 guards are always sufficient:

Conjecture 8. Let P be an integral orthogonal polygon of perimeter N. Then P can always be guarded with at most |N/6| guards.

References

- T. Biedl, T. M. Chan, S. Lee, S. Mehrabi, F. Montecchiani, H. Vosoughpour, and Z. Yu. Guarding orthogonal art galleries with sliding k-transmitters: Hardness and approximation. *Algorithmica*, 81(1):69–97, 2019.
- [2] T. Biedl, M. T. Irfan, J. Iwerks, J. Kim, and J. S. Mitchell. The art gallery theorem for polyominoes. Discrete & Computational Geometry, 48(3):711–720, 2012.
- [3] I. Bjorling-Sachs. Edge guards in rectilinear polygons. Computational Geometry, 11(2):111– 123, 1998.
- [4] V. Chvátal. A combinatorial theorem in plane geometry. Journal of Combinatorial Theory, Series B, 18(1):39–41, 1975.

- [5] S. Durocher, O. Filtser, R. Fraser, A. D. Mehrabi, and S. Mehrabi. Guarding orthogonal art galleries with sliding cameras. *Computational Geometry*, 65:12–26, 2017.
- S. Fisk. A short proof of Chvátal's watchman theorem. Journal of Combinatorial Theory, Series B, 24(3):374, 1978.
- [7] E. Györi. A short proof of the rectilinear art gallery theorem. SIAM Journal on Algebraic Discrete Methods, 7(3):452–454, 1986.
- [8] F. Hoffmann and K. Kriegel. A graph-coloring result and its consequences for polygonguarding problems. SIAM Journal on Discrete Mathematics, 9(2):210–224, 1996.
- [9] R. Honsberger. Mathematical gems ii, mathematical association of america, dolciani math, 1976.
- [10] J. Kahn, M. Klawe, and D. Kleitman. Traditional galleries require fewer watchmen. SIAM Journal on Algebraic Discrete Methods, 4(2):194–206, 1983.
- [11] M. J. Katz and G. Morgenstern. Guarding orthogonal art galleries with sliding cameras. International Journal of Computational Geometry & Applications, 21(02):241–250, 2011.
- [12] M. J. Katz and G. S. Roisman. On guarding the vertices of rectilinear domains. Computational Geometry, 39(3):219–228, 2008.
- J. O'Rourke. An alternate proof of the rectilinear art gallery theorem. Journal of Geometry, 21(1):118–130, 1983.
- [14] J. O'Rourke. Art Gallery Theorems and Algorithms, volume 57. Oxford University Press, Inc., New York, NY, USA, 1987.
- [15] T. C. Shermer. Recent results in art galleries (geometry). Proceedings of the IEEE, 80(9):1384–1399, 1992.
- [16] G. T. Toussaint. Computational morphology: A Computational geometric approach to the analysis of form. Elsevier, 2014.
- [17] J. Urrutia. Sixth proof of the orthogonal art gallery theorem. Technical Report TR-97-03, Department of Computer Science, University of Ottawa, 1997.
- [18] J. Urrutia. Art gallery and illumination problems. In J.-R. S. Urrutia, editor, Handbook of Computational Geometry, pages 973 – 1027. North-Holland, Amsterdam, 2000.