# CONVERGENCE OF NONEQUILIBRIUM LANGEVIN DYNAMICS FOR PLANAR FLOWS

# MATTHEW DOBSON AND ABDEL KADER GERALDO

Department of Mathematics, University of Massachusetts Amherst, Amherst, MA 01003

# CONTENTS

1. Introduction	1
2. Reformulation of NELD in Lagrangian coordinates	3
2.1. Remapping the Unit Cell	3
2.2. Remapped coordinates	4
2.3. NELD with LE and KR PBCs in the Lagrangian coordinates	6
3. Ergodicity of NELD under Planar Flow	7
3.1. Fokker-Planck Equation of NELD in Eulerian and Lagrangian Coordinates	7
3.2. Convergence of NELD to a Limit Cycle	9
4. Conclusion	16
Acknowledgements	16
References	17

ABSTRACT. We prove that incompressible two dimensional nonequilibrium Langevin dynamics (NELD) converges exponentially fast to a steady-state limit cycle. We use automorphism remapping periodic boundary conditions (PBCs) techniques such as Lees-Edwards PBCs and Kraynik-Reinelt PBCs to treat respectively shear flow and planar elongational flow. After rewriting NELD in Lagrangian coordinates, the convergence is shown using a technique similar to [ R. Joubaud, G. A. Pavliotis, and G. Stoltz,2014].

# 1. INTRODUCTION

A wide range of nonequilibrium molecular dynamics (NEMD) techniques [5,7] are used in the study of molecular fluids under steady flow, and some recent applications can be found in [1–4,11–21]. Here we study the exponential convergence of the probability density of Nonequilibrium Langevin dynamics (NELD) under incompressible two dimensional flows such as shear flow and planar elongational flow with spatial periodic boundary conditions (PBCs). We consider a molecular system corresponding to the micro-scale motion of a fluid with local strain rate  $\nabla \mathbf{u}$  and denote the steady background flow matrix of the molecular system by  $A = \nabla \mathbf{u} \in \mathbb{R}^{3\times 3}$ . The coordinates of the simulation box are given by three linearly independent column vectors coming from the origin, and we write them in a matrix

$$L_t = \begin{bmatrix} \mathbf{v}_t^1 & \mathbf{v}_t^2 & \mathbf{v}_t^3 \end{bmatrix} \in \mathbb{R}^{3 \times 3}, \qquad t \ge 0.$$

*E-mail address*: dobson@math.umass.edu.

Date: October 28, 2022.

The initial simulation box is given by  $L_0$  and the lattice deforms with the background flow according to the equation

$$L_t = e^{tA} L_0.$$

If  $L_0$  is not chosen appropriately, the simulation box can become extremely stretched with degenerate geometry. For example, in the elongational flow case, if the compression is parallel to one of the edges of the simulation box, then the box will become degenerate to a point where a particle and its image become arbitrarily close. Thus in order to perform a long simulation, we consider specialized PBCs which consist of using a lattice automorphism represented as a  $3 \times 3$  integer matrix with determinant one to remap the simulation box at various times during the simulation. These types of PBCs were first used in the shear flow case by Lees and Edwards (LE) [9] and were then later extended to the planar elongational flow case by Kraynik and Reinelt (KR) [8]. The analog of these types of PBCs which treat three dimensional flows such as uniaxial flow, biaxial flow, and generalized three-dimensional diagonalizable flow can be found in [6, 10, 32].

The NELD equation is derived in [22,23]. We express it in terms of the relative momentum  $\widetilde{\mathbf{p}}$  as

$$\begin{cases} d\widetilde{\mathbf{q}} &= (\widetilde{\mathbf{p}} + A\widetilde{\mathbf{q}})dt, \\ d\widetilde{\mathbf{p}} &= -\nabla V(\widetilde{\mathbf{q}})dt - \gamma \widetilde{\mathbf{p}}dt + \sigma dW, \end{cases}$$
(1)

where  $\sigma^2 = \frac{2\gamma}{\beta}$  is the fluctuation coefficient, with  $\beta$  as the inverse temperature, and V is the potential. We assume that the gradient of the potential is finite. The position and the momentum of the particles are denoted respectively by  $(\tilde{\mathbf{q}}, \tilde{\mathbf{p}}) \in \tilde{\mathcal{L}}_t^d \times \mathbb{R}^{3d}$ , where the set

$$\widetilde{\mathcal{L}}_t = \{ L_t \mathbf{x} \mid \mathbf{x} \in \mathbb{T}^3 \}$$
(2)

defines the time dependent simulation box. Note that when there is no background flow, (1) becomes equilibrium Langevin Dynamics

$$\begin{cases} d\mathbf{q} &= \mathbf{p}dt, \\ d\mathbf{p} &= -\nabla V(\mathbf{q})dt - \gamma \mathbf{p}dt + \sigma dW, \end{cases}$$

with  $(\mathbf{q}, \mathbf{p}) \in \mathcal{L}_0^d \times \mathbb{R}^{3d}$ .

It has been shown (see for instance [24–26, 29–31]) that under suitable conditions, the equilibrium Langevin Dynamics is ergodic with respect to the Boltzmann-Gibbs distribution

$$\nu(\mathbf{q}, \mathbf{p}) d\mathbf{q} d\mathbf{p} = \frac{1}{Z} e^{-\beta H(\mathbf{q}, \mathbf{p})} d\mathbf{q} d\mathbf{p}, \quad Z = \int_{\mathcal{L}_0^d \times \mathbb{R}^{3d}} e^{-\beta H(\mathbf{q}, \mathbf{p})} d\mathbf{q} d\mathbf{p},$$

where Z is the normalization constant, and H is the Hamiltonian of the system given by

$$H(\mathbf{q}, \mathbf{p}) = \frac{1}{2} \langle \mathbf{p}, \mathbf{p} \rangle + V(\mathbf{q}).$$

However, convergence to a limiting measure has not been established for NELD under moving domains. In this paper, we show the existence, uniqueness, and exponential convergence of the NELD to a limit cycle following the work done in [27]. In Section 2, we formulate the NELD in Eulerian and Lagrangian coordinates and the main result of the paper is in Section 3, where we prove the convergence of the NELD to a probability density function in Proposition 1.

### 2. Reformulation of NELD in Lagrangian coordinates

In this section, we rewrite the NELD equation (1) in Lagrangian coordinates. We then define the remapped Eulerian domain under shear flow and planar elongational flow using the LE and KR PBCs respectively in Section 2.1. In order to distinguish from the remapped coordinates, we refer to the original coordinate systems as "absolute Eulerian coordinates" and "absolute Lagrangian coordinates." We then derive the NELD in the remapped coordinates in Section 2.3. We start by considering the change of variables from the absolute Eulerian to Lagrangian coordinates

$$\begin{cases} \mathbf{q} = e^{-tA} \widetilde{\mathbf{q}}, \quad \mathbf{q} \in \mathcal{L}_0^d, \ \widetilde{\mathbf{q}} \in \widetilde{\mathcal{L}}_t^d \\ \mathbf{p} = e^{-tA} \widetilde{\mathbf{p}}, \quad \mathbf{p} \in \mathbb{R}^{3d}. \end{cases}$$
(3)

Computing the time derivative of the position in (3), and using (1), we have

$$d\mathbf{q} = e^{-tA} \left( d\widetilde{\mathbf{q}} - A\widetilde{\mathbf{q}}dt \right) = \mathbf{p}dt,$$

taking the time derivative of  $\mathbf{p}$  gives

$$d\mathbf{p} = e^{-tA} \left( d\widetilde{\mathbf{p}} - A\widetilde{\mathbf{p}}dt \right) = -e^{-tA} \nabla V(e^{tA}\mathbf{q}) dt - \Gamma \mathbf{p}dt + \sigma e^{-tA} dW,$$

where  $\Gamma = (\gamma + A)$ . Thus, the NELD equation in Lagrangian coordinates is written as

$$\begin{cases} d\mathbf{q} &= \mathbf{p}dt, \\ d\mathbf{p} &= -e^{-tA}\nabla V(e^{tA}\mathbf{q})dt - \Gamma \mathbf{p}dt + \sigma e^{-tA}dW, \end{cases}$$

Before we derive the NELD equation in remapped Lagrangian coordinates under LE and KR PBCs, we give some background on those PBCs.

2.1. Remapping the Unit Cell. We start by defining the remapped Eulerian domain under the shear flow followed by the planar elongational flow case.

2.1.1. Shear Case. We denote the background matrix of the shear flow by

$$A = \begin{bmatrix} 0 & \epsilon & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \ \epsilon \in \mathbb{R}^*$$

At a time t, the basis vectors for the simulation box are the columns of the matrix

$$L_t = \begin{bmatrix} 1 & t\epsilon & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} L_0 \text{ where } L_0 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Since  $L_t$  is highly sheared as t becomes large, the interparticle interaction computation becomes more difficult. We can prevent this anomaly by applying the LE PBCs which consists in multiplying  $L_t$  by the lattice automorphism matrix

$$M^{k} = \begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{k} = \begin{bmatrix} 1 & -k & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad k \in \mathbb{Z},$$

to get the remapped simulation box lattice

$$L_t M^k = \begin{bmatrix} 1 & t\epsilon - k & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, k \in \mathbb{Z}.$$

Since M is an integer matrix with determinant equal to one  $(M \in SL(3,\mathbb{Z}))$ , the lattice basis vectors in  $L_t$  and  $L_t M^k$  generate the same lattice. By choosing  $k = -\lfloor t\epsilon \rceil$ , where  $\lfloor x \rceil$  denotes x rounded to nearest integer, we ensure that the stretch is at most half of the simulation box. Then we observe that the stretch matrix is time-periodic with the period  $T = \frac{1}{\epsilon}$ :

$$\begin{bmatrix} 0 & t\epsilon - \lfloor t\epsilon \rceil & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} t \end{bmatrix} \begin{bmatrix} 0 & \epsilon & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} t \end{bmatrix} A, \text{ where } \begin{bmatrix} t \end{bmatrix} \equiv t \mod T.$$
(4)

This implies that the particle position belongs to remapped Eulerian domain

$$\widehat{\mathcal{L}}_t = \{ e^{[t]A} L_0 \mathbf{x} | \mathbf{x} \in \mathbb{T}^3 \}, \text{ where } \mathbb{T}^3 = \mathbb{R}^3 \backslash \mathbb{Z}^3.$$
(5)

In the Section 2.2, we analyse the particles remapped position in  $\widehat{\mathcal{L}}_t$  and in the unit cell.

2.1.2. *Planar Elongational Flow case.* We consider the planar elongational flow (PEF) case with background flow matrix given by

$$A = \begin{bmatrix} \epsilon & 0 & 0 \\ 0 & -\epsilon & 0 \\ 0 & 0 & 0 \end{bmatrix}, \ \epsilon \in \mathbb{R}^*,$$

which means that the simulation box elongates in the x direction and shrinks in the y direction of the standard coordinate plane. As t goes to infinity, a particle and its image can become arbitrarily close if an edge of the simulation box is aligned with the y coordinate. This would lead to a breakdown in the simulation. We prevent this issue by applying the KR PBCs, which consists in carefully choosing the alignment of the initial simulation box and remapping the simulation box with a matrix  $M \in SL(3, \mathbb{Z})$ . We choose M such that it is diagonalizable with eigenvalues of the form

$$MS = S\Lambda, \quad \Lambda = \begin{bmatrix} \lambda & 0 & 0\\ 0 & \lambda^{-1} & 0\\ 0 & 0 & 1 \end{bmatrix}, \quad \lambda > 0, \quad \lambda \neq 1.$$

We choose the initial lattice  $L_0 = S^{-1}$  rather than the standard coordinate directions, and this will prevent particle images from becoming too close. If we remap the lattice basis vectors by applying  $M^k$ , we get

$$L_t M^k = e^{tA} L_0 M^k = e^{t\epsilon D} \Lambda^n S^{-1} = e^{(t\epsilon + k\eta)D} S^{-1}, \text{ where } D = \begin{bmatrix} 1 & 0 & 0\\ 0 & -1 & 0\\ 0 & 0 & 0 \end{bmatrix}, \quad \eta = \log(\lambda).$$

Letting  $T = \frac{\eta}{\epsilon}$ , the stretched matrix [t]A and the position domain of the particles are also respectively expressed in the periodic form as in (4) and (5).

2.2. **Remapped coordinates.** Note that under the PBCs, the remapping of the simulation box is followed by remapping the particle positions to lie within the simulation box. In order to include this remapping in the dynamics, we write down the remapping function for both particle positions and momenta in both Eulerian and Lagrangian coordinate systems.

2.2.1. Remapping particle positions in Eulerian coordinates. Here, we define the function which remaps the particle positions from the absolute Eulerian domain  $\tilde{L}_t$  (2) to the remapped Eulerian domain  $\hat{L}_t$  (5). This function chooses the image particle that lives within the unit cell for the remapped lattice. We start by defining the modulus operation applied to each vector component

$$\mathbf{g}(\mathbf{x}) \equiv \mathbf{x} \mod 1$$
, where  $\mathbf{x} \in \mathbb{R}^3$ .

We then compute  $e^{-[t]A}\tilde{\mathbf{q}}$ , which maps the particles back to a point in time when an integer number of periods have occurred. Multiplying by  $L_0^{-1}$  expresses the particle position in lattice coordinates, where the coordinates corresponding to unit cell is given by  $[0, 1]^3$ , but the particle coordinates may be outside this cube due to the cell's deformation. Applying the modulus operation  $\mathbf{g}$  finds the coordinates of images within the unit cell, and then we map back to Eulerian space by multiplying by  $e^{[t]A}L_0$ , which gives

$$\widehat{\mathbf{g}}_t : \widetilde{\mathcal{L}}_t \to \widehat{\mathcal{L}}_t \qquad \widehat{\mathbf{q}} = \widehat{\mathbf{g}}_t(\widetilde{\mathbf{q}}) = e^{[t]A} L_0 \, \mathbf{g} \left( L_0^{-1} e^{-[t]A} \widetilde{\mathbf{q}} \right)$$

2.2.2. *Remapping particle positions in Lagrangian coordinates.* We use the Eulerian remapping function to remap the Lagrangian coordinates, as shown in the following diagram:

$$\begin{array}{c} \widetilde{\mathcal{L}}_t \xrightarrow{\widehat{\mathbf{g}}_t} \widehat{\mathcal{L}}_t \\ e^{tA} \uparrow & \uparrow e^{[t]A} \\ \mathcal{L}_0 \xrightarrow{\overline{\mathbf{g}}_t} \mathcal{L}_0 \end{array}$$

We map from absolute Lagrangian coordinates to Eulerian coordinates using (3), perform the remapping  $\hat{\mathbf{g}}_t$ , and then returning to Lagrangian coordinates:

$$\overline{\mathbf{g}}_t : \mathcal{L}_0 \to \mathcal{L}_0 \qquad \overline{\mathbf{q}} = \overline{\mathbf{g}}_t(\mathbf{q}) = e^{-[t]A} \,\widehat{\mathbf{g}}_t \left( e^{tA} \mathbf{q} \right)$$

2.2.3. Remapping particle momenta. We now remap the momentum of the particles from the absolute to the remapped domains. We start by taking the time derivative of  $\overline{\mathbf{q}}$ , we get

$$d\overline{\mathbf{q}} = d\overline{\mathbf{g}}_t(\mathbf{q}) = e^{\left\lfloor \frac{t}{T} \right\rfloor TA} \mathbf{p} dt = \overline{\mathbf{p}} dt,$$

where we have defined the remapped momentum

$$\overline{\mathbf{h}}_t : \mathbb{R}^{3d} \to \mathbb{R}^{3d} \qquad \overline{\mathbf{p}} = \overline{\mathbf{h}}_t(\mathbf{p}) = e^{\left\lfloor \frac{t}{T} \right\rfloor T A} \mathbf{p}.$$
(6)

We chose the above mapping based on the form of the position equation  $d\mathbf{\bar{q}} = \mathbf{\bar{p}}dt$ , and we now show that this choice leads to consistency of the NELD equations in remapped Eulerian coordinates. In remapped Eulerian coordinates, the domain is time periodic but the coefficients of the NELD are not changed by the remapping. We denote the position in the remapped Eulerian domain by

$$\widehat{\mathbf{q}} = e^{[t]A}\overline{\mathbf{q}}, \quad \widehat{\mathbf{q}} \in \widehat{\mathcal{L}}_t,$$

and then by taking the time derivative of  $\hat{\mathbf{q}}$  over a periodic time  $\theta = [t] \in [0,T)$ , we get

$$d\widehat{\mathbf{q}} = e^{\theta A} \left( d\overline{\mathbf{q}} + A\overline{\mathbf{q}}d\theta \right) = e^{\theta A} (\overline{\mathbf{p}} + A\overline{\mathbf{q}})d\theta = (\widehat{\mathbf{p}} + A\widehat{\mathbf{q}})d\theta$$

where we have defined the relative momentum of the particles in the remapped Eulerian coordinates by

$$\widehat{\mathbf{p}} = e^{[t]A}\overline{\mathbf{p}}.$$

Since  $\tilde{\mathbf{p}} = e^{tA}\mathbf{p}$  by (3), and by using (6), the relative momentum of the particles in the Eulerian domain are the same, as

$$\widehat{\mathbf{p}} = e^{[t]A}\overline{\mathbf{p}} = e^{tA}e^{-\left\lfloor \frac{t}{T} \right\rfloor TA}\overline{\mathbf{p}} = e^{tA}\mathbf{p} = \widetilde{\mathbf{p}}.$$

Thus the NELD as a function of  $(\widehat{\mathbf{q}}_{kT+\theta}, \widehat{\mathbf{p}}_{kT+\theta}) \in \widehat{\mathcal{L}}_t^d \times \mathbb{R}^{3d}$  which is written as

$$\begin{cases} d\widehat{\mathbf{q}}_{kT+\theta} &= \left(\widehat{\mathbf{p}}_{kT+\theta} + A\widehat{\mathbf{q}}_{kT+\theta}\right)d\theta\\ d\widehat{\mathbf{p}}_{kT+\theta} &= -\nabla V(\widehat{\mathbf{q}}_{kT+\theta})d\theta - \gamma\widehat{\mathbf{p}}_{kT+\theta}d\theta + \sigma dW_{kT+\theta}, \end{cases}$$

has the same coefficients as (1).

Now let us define the momentum remapping function in the Eulerian domain by

$$\widehat{\mathbf{h}}_t : \mathbb{R}^{3d} \to \mathbb{R}^{3d} \quad \widehat{\mathbf{h}}_t(\widetilde{\mathbf{p}}) = \widetilde{\mathbf{p}}$$

We summarize the results in this section using the following definitions:

**Definition 1.** The remapping function from the absolute to the remapped Lagrangian coordinates is given by

$$\mathfrak{R}_t: \mathcal{L}_0^d \times \mathbb{R}^{3d} \to \mathcal{L}_0^d \times \mathbb{R}^{3d} \qquad \mathfrak{R}_t((\mathbf{q}_t, \mathbf{p}_t)) = \left(\overline{\mathbf{g}}_t(\mathbf{q}_t), \overline{\mathbf{h}}_t(\mathbf{p}_t)\right)$$

**Definition 2.** The remapping function from the absolute to the remapped Eulerian coordinates is given by

$$\widetilde{\mathfrak{R}}_t: \widetilde{\mathcal{L}}_t^d \times \mathbb{R}^{3d} \to \widehat{\mathcal{L}}_t^d \times \mathbb{R}^{3d} \qquad \widetilde{\mathfrak{R}}_t((\widetilde{\mathbf{q}}_t, \widetilde{\mathbf{p}}_t)) = \big(\widehat{\mathbf{g}}_t(\widetilde{\mathbf{q}}_t), \widehat{\mathbf{h}}_t(\widetilde{\mathbf{p}}_t)\big).$$
(7)

Using the above definitions, and the linear map

$$\Phi_t : \mathcal{L}_0^d \times \mathbb{R}^{3d} \to \widehat{\mathcal{L}}_t^d \times \mathbb{R}^{3d} \qquad \Phi_t(\overline{\mathbf{X}}_t) = \begin{bmatrix} e^{[t]A} & 0\\ 0 & e^{[t]A} \end{bmatrix} \overline{\mathbf{X}}_t, \tag{8}$$

$$\widetilde{\Phi}_t : \mathcal{L}_0^d \times \mathbb{R}^{3d} \to \widetilde{\mathcal{L}}_t^d \times \mathbb{R}^{3d} \qquad \widetilde{\Phi}_t(\mathbf{X}_t) = \begin{bmatrix} e^{tA} & 0\\ 0 & e^{tA} \end{bmatrix} \mathbf{X}_t, \tag{9}$$

we have the commutative diagram as following:

$$\begin{aligned} \widetilde{\mathcal{L}}_t^d \times \mathbb{R}^{3d} & \xrightarrow{\mathfrak{R}_t} \quad \widehat{\mathcal{L}}_t^d \times \mathbb{R}^{3d} \\ \widetilde{\Phi}_t & \uparrow & \uparrow \\ \mathcal{L}_0^d \times \mathbb{R}^{3d} & \xrightarrow{\mathfrak{R}_t} \quad \mathcal{L}_0^d \times \mathbb{R}^{3d} \end{aligned}$$

The remapping function from the remapped Lagrangian to the remapped Eulerian domain is given by

$$\Phi_t = \widetilde{\mathfrak{R}}_t \circ \widetilde{\Phi}_t \circ \mathfrak{R}_t^{-1}. \tag{10}$$

2.3. **NELD with LE and KR PBCs in the Lagrangian coordinates.** Using the function  $\mathfrak{R}_t$ , we derive the NELD in the remapped Lagrangian domain. The coefficients of the NELD are time periodic, and we use this periodicity in the analysis that follows. In fact, we use  $\mathfrak{R}_t$  to consider the change of variables

$$\begin{cases} \overline{\mathbf{q}}_{kT+\theta} = \overline{\mathbf{g}}_k(\mathbf{q}_{kT+\theta}), \\ \overline{\mathbf{p}}_{kT+\theta} = e^{kTA}\mathbf{p}_{kT+\theta}, \end{cases}$$

where  $\theta = [t] \in [0, T)$  and  $t \in [kT, (k+1)T)$ . We use  $\overline{\mathbf{g}}_k$  to define  $\overline{\mathbf{g}}_t$  for  $t \in [kT, (k+1)T)$ . Changing variables in the position equation of the NELD gives

$$\frac{d}{d\theta}\overline{\mathbf{q}}_{kT+\theta} = \frac{d}{d\theta}\overline{\mathbf{g}}_k(\mathbf{q}_{kT+\theta}) = \nabla_{\mathbf{q}}\overline{\mathbf{g}}_k(\mathbf{q}_{kT+\theta})\frac{d}{d\theta}\mathbf{q}_{kT+\theta} = e^{kTA}\mathbf{p}_{kT+\theta} = \overline{\mathbf{p}}_{kT+\theta},$$

while in the momentum, we have

$$d\overline{\mathbf{p}}_{kT+\theta} = e^{kTA} d\mathbf{p}_{kT+\theta} = -e^{-\theta A} (\nabla V \circ \widetilde{\mathbf{g}}_k) (e^{\theta A} \overline{\mathbf{q}}_{kT+\theta}) d\theta - \Gamma \overline{\mathbf{p}}_{kT+\theta} d\theta + \sigma e^{-\theta A} dW_{kT+\theta}.$$

Thus the NELD as a function of  $(\overline{\mathbf{q}}_{kT+\theta}, \overline{\mathbf{p}}_{kT+\theta}) \in \mathcal{L}_0^d \times \mathbb{R}^{3d}$  is given by

$$\begin{cases} d\overline{\mathbf{q}}_{kT+\theta} &= \overline{\mathbf{p}}_{kT+\theta} d\theta, \\ d\overline{\mathbf{p}}_{kT+\theta} &= -e^{-\theta A} \nabla V \left( e^{\theta A} \overline{\mathbf{q}}_{kT+\theta} \right) d\theta - \Gamma \overline{\mathbf{p}}_{kT+\theta} d\theta + \sigma e^{-\theta A} dW_{kT+\theta}. \end{cases}$$
(11)

# 3. Ergodicity of NELD under Planar Flow

We start this section by deriving the forward and backward Kolmogorov equation of the NELD in Section 3.1, then in Section 3.2 we prove the main result of the paper, the convergence of the NELD to a limit cycle.

3.1. Fokker-Planck Equation of NELD in Eulerian and Lagrangian Coordinates. We now derive the forward and backward Kolmogorov equations of NELD in Eulerian and Lagrangian coordinates. In the absolute Eulerian coordinates, the NELD equations of motion are rewritten as

$$\begin{cases} d\widetilde{\mathbf{X}}_t = \widetilde{\mathbf{b}}(\widetilde{\mathbf{X}}_t)dt + \widetilde{\Sigma}dW_t, & \widetilde{\mathbf{X}}_t \in \widetilde{\mathcal{L}}_t^d \times \mathbb{R}^{3d}, \\ \widetilde{\mathbf{X}}_t = \begin{bmatrix} \widetilde{\mathbf{q}} \\ \widetilde{\mathbf{p}} \end{bmatrix}, & \widetilde{\mathbf{b}}(\widetilde{\mathbf{X}}_t) = \begin{bmatrix} \widetilde{\mathbf{p}} + A\widetilde{\mathbf{q}} \\ (-\nabla V(\widetilde{\mathbf{q}}) - \gamma \widetilde{\mathbf{p}} \end{bmatrix}, & \widetilde{\Sigma} = \begin{bmatrix} 0 & 0 \\ 0 & \sigma \end{bmatrix}.$$
(12)

Given any smooth function  $\widetilde{\mathbf{f}} \in C^{\infty}(\widetilde{\mathcal{L}}_t^d \times \mathbb{R}^{3d}; \mathbb{R})$  Itô's lemma says that

$$d\widetilde{\mathbf{f}}(\widetilde{\mathbf{X}}_t) = \left(\widetilde{U}_t \widetilde{\mathbf{f}}\right)(\widetilde{\mathbf{X}}_t) dt + \left\langle \nabla \widetilde{\mathbf{f}}(\widetilde{\mathbf{X}}_t), \widetilde{\Sigma} dW \right\rangle, \tag{13}$$

where

$$\widetilde{U}_t = \left\langle \widetilde{\mathbf{p}} + A \widetilde{\mathbf{q}}, \nabla_{\widetilde{\mathbf{q}}} \right\rangle + \left\langle -\nabla V(\widetilde{\mathbf{q}}), \nabla_{\widetilde{\mathbf{p}}} \right\rangle - \gamma \left\langle \widetilde{\mathbf{p}}, \nabla_{\widetilde{\mathbf{p}}} \right\rangle + \frac{1}{2} \sigma \sigma^T : \nabla^2.$$

The symbol : and  $\langle \cdot, \cdot \rangle$  denote the Frobenius inner product. Moreover, we derive the generator  $U_t$  of the NELD in the absolute Lagrangian coordinates as follows: rewriting (11) in the absolute Lagrangian coordinates as

$$\begin{cases} d\mathbf{X}_t = \mathbf{b}(\mathbf{X}_t, t)dt + \Sigma(t)dW_t, & \mathbf{X}_t \in \mathcal{L}_0^d \times \mathbb{R}^{3d} \\ \mathbf{X}_t = \begin{bmatrix} \mathbf{q} \\ \mathbf{p} \end{bmatrix}, & \mathbf{b}(\mathbf{X}_t, t) = \begin{bmatrix} \mathbf{p} \\ -e^{-tA}\nabla V(e^{tA}\mathbf{q}) - \Gamma\mathbf{p} \end{bmatrix}, & \Sigma(t) = \begin{bmatrix} 0 & 0 \\ 0 & \sigma e^{-tA} \end{bmatrix}. \tag{14}$$

When applying Itô's lemma, we compute (13) using the change of variables as

$$d(\widetilde{\mathbf{f}} \circ \widetilde{\Phi}_t)(\mathbf{X}_t) = (\partial_t + U_t)(\widetilde{\mathbf{f}} \circ \widetilde{\Phi}_t)(\mathbf{X}_t)dt + \left\langle \nabla_{\mathbf{X}}(\widetilde{\mathbf{f}} \circ \widetilde{\Phi}_t)(\mathbf{X}_t), \Sigma(t)dW \right\rangle,$$

where

$$U_t = \langle \mathbf{p}, \nabla_{\mathbf{q}} \cdot \rangle - \left\langle e^{-tA} \nabla V(e^{tA} \mathbf{q}), \nabla_{\mathbf{p}} \cdot \right\rangle - \left\langle \Gamma \mathbf{p}, \nabla_{\mathbf{p}} \cdot \right\rangle + \frac{1}{2} (\sigma e^{-tA}) (\sigma e^{-tA})^T : \nabla^2.$$

Now let us write the strong solution of (12) as

$$\widetilde{\mathbf{X}}_t - \widetilde{\mathbf{X}}_s = \int_s^t \widetilde{\mathbf{b}}(\widetilde{\mathbf{X}}_u) du + \widetilde{\Sigma} dW_u$$

and define the density transition function from one state to another in the continuous-time Markov chain  $\widehat{\mathbf{X}}_t = \widetilde{\mathfrak{R}}_t(\widetilde{\mathbf{X}}_t)$  by

$$\widehat{\mathcal{P}}_t(\widehat{\mathbf{y}}, \widehat{B}_t) = \mathbb{P}\big(\widehat{\mathbf{X}}_t \in \widehat{B}_t \big| \widehat{\mathbf{X}}_s = \widehat{\mathbf{y}}\big) = \int_{\widehat{B}_t} \widehat{\psi}(t, \widehat{\mathbf{x}} \big| s, \widehat{\mathbf{y}}) d\widehat{\mathbf{x}}, \text{ where } \widehat{\psi}(t, \widehat{\mathbf{x}} \big| t, \widehat{\mathbf{y}}) = \delta(\widehat{\mathbf{x}} - \widehat{\mathbf{y}}),$$

 $\forall \widehat{\mathbf{x}} \in \widehat{\mathcal{L}}_t^d \times \mathbb{R}^{3d}, \ \widehat{B}_t \in \mathcal{B}(\widehat{\mathcal{L}}_t^d \times \mathbb{R}^{3d}).$  Here  $\mathcal{B}(\widehat{\mathcal{L}}_t^d \times \mathbb{R}^{3d})$  denotes the Börel  $\sigma$ -algebra on  $\widehat{\mathcal{L}}_t^d \times \mathbb{R}^{3d}.$ We transform to the remapped Eulerian coordinates using  $\widetilde{\mathfrak{R}}_t$ . Using (10), we define

$$\mathbf{f}:\widehat{\mathcal{L}}_t^d\times\mathbb{R}^{3d}\to\mathbb{R}\quad\mathbf{f}=\widetilde{\mathbf{f}}\circ\widetilde{\mathfrak{R}}_t^{-1},$$

such that

$$\mathbf{f} \circ \Phi_t \circ \mathfrak{R}_t = \widetilde{\mathbf{f}} \circ \widetilde{\mathfrak{R}}_t^{-1} \circ \Phi_t \circ \mathfrak{R}_t = \widetilde{\mathbf{f}} \circ \widetilde{\Phi}_t : \mathcal{L}_0^d \times \mathbb{R}^{3d} \to \mathbb{R}$$

Then in the remapped Eulerian domain  $\widehat{\mathcal{L}}_t$ , we define the expectation of  $\mathbf{f} : \widehat{\mathcal{L}}_t^d \times \mathbb{R}^{3d} \to \mathbb{R}$  with respect to the probability density function above by

$$\phi(s, \widehat{\mathbf{y}}) = \mathbb{E}^{s, \mathbf{y}} \mathbf{f}(\widehat{\mathbf{X}}_t) = \int_{\widehat{\mathcal{L}}_t^d \times \mathbb{R}^{3d}} \mathbf{f}(\widehat{\mathbf{x}}) \psi(t, \widehat{\mathbf{x}} | s, \widehat{\mathbf{y}}) d\widehat{\mathbf{x}}.$$
 (15)

Similarly, in the absolute Lagrangian domain, let us write the strong solution of (14) as

$$\mathbf{X}_t - \mathbf{X}_s = \int_s^t \mathbf{b}(\mathbf{X}_u) du + \Sigma(u) dW_u,$$

and define the density transition function from one state to another in the continuous-time Markov chain  $\overline{\mathbf{X}}_t = \mathfrak{R}_t(\mathbf{X}_t)$  by

$$\overline{\mathcal{P}}_t(\overline{\mathbf{y}},\overline{B}) = \mathbb{P}\big(\overline{\mathbf{X}}_t \in \overline{B} \big| \overline{\mathbf{X}}_s = \overline{\mathbf{y}}\big) = \int_{\overline{B}} \psi(t,\overline{\mathbf{x}} \big| s,\overline{\mathbf{y}}) d\overline{\mathbf{x}}, \text{ where } \psi(t,\overline{\mathbf{x}} \,| t,\overline{\mathbf{y}}) = \delta(\overline{\mathbf{x}} - \overline{\mathbf{y}})$$

 $\forall \overline{\mathbf{x}} \in \mathcal{L}_0^d \times \mathbb{R}^{3d}, \ \overline{B} \in \mathcal{B}(\mathcal{L}_0^d \times \mathbb{R}^{3d})$ . We define a measurable function  $\mathbf{f} \circ \Phi_t : \mathcal{L}_0^d \times \mathbb{R}^{3d} \to \mathbb{R}$ . Using the change of variables (8), we can rewrite the expectation (15) with respect to the probability density function above as

$$(\phi \circ \Phi_s)(s, \overline{\mathbf{y}}) = \mathbb{E}^{s, \mathbf{y}}(\mathbf{f} \circ \Phi_t)(\overline{\mathbf{X}}_t) = \int_{\mathcal{L}_0^d \times \mathbb{R}^{3d}} (\mathbf{f} \circ \Phi_t)(\overline{\mathbf{x}}) \psi(t, \overline{\mathbf{x}} | s, \overline{\mathbf{y}}) d\overline{\mathbf{x}}, \quad s \le t.$$

Now we derive the backward Kolmogorov equation for the NELD in following lemma:

Lemma 1. [37, Theorem 6.1] The backward Kolmogorov equation for the NELD is

$$\partial_s \psi(t, \overline{\mathbf{x}} | s, \overline{\mathbf{y}}) + (U_s \psi)(t, \overline{\mathbf{x}} | s, \overline{\mathbf{y}}) = 0, \quad where \quad \psi(t, \overline{\mathbf{x}} | s, \overline{\mathbf{y}})|_{t=s} = \delta(\overline{\mathbf{x}} - \overline{\mathbf{y}}), \\ \partial_s \widehat{\psi}(t, \widehat{\mathbf{x}} | s, \widehat{\mathbf{y}}) + (\widehat{U}_s \widehat{\psi})(t, \widehat{\mathbf{x}} | s, \widehat{\mathbf{y}}) = 0, \quad where \quad \widehat{\psi}(t, \widehat{\mathbf{x}}, | s, \widehat{\mathbf{y}})\Big|_{t=s} = \delta(\widehat{\mathbf{x}} - \widehat{\mathbf{y}}), \quad s < t.$$

The forward Kolmogorov equation of the NELD is given in the following lemma:

**Lemma 2.** For an operator G, let us denote the adjoint operator with respect to Lebesgue measure by  $G^{\dagger}$ . The forward Kolmogorov equation of the NELD is

$$\left(-\partial_t\psi + U_t^{\dagger}\psi\right)(t,\overline{\mathbf{x}}|s,\overline{\mathbf{y}}) = 0 \text{ and } \left(-\partial_t\widehat{\psi} + \widehat{U}_t^{\dagger}\widehat{\psi}\right)(t,\widehat{\mathbf{x}}|s,\widehat{\mathbf{y}}) = 0.$$
(16)

*Proof.* Using the adjoint property, we have in the remapped Lagrangian and Eulerian domain respectively

$$\int_{\mathcal{L}_0^d \times \mathbb{R}^{3d}} \left( \partial_t + U_t \right) \left( \mathbf{f} \circ \Phi_t \right) (\overline{\mathbf{x}}) \psi(t, \overline{\mathbf{x}} | s, \overline{\mathbf{y}}) d\overline{\mathbf{x}} = \int_{\mathcal{L}_0^d \times \mathbb{R}^{3d}} \left( \mathbf{f} \circ \Phi_t \right) (\overline{\mathbf{x}}) \left( -\partial_t + U_t^{\dagger} \right) \psi(t, \overline{\mathbf{x}} | s, \overline{\mathbf{y}}) d\overline{\mathbf{x}},$$

and

$$\int_{\widehat{\mathcal{L}}_t^d \times \mathbb{R}^{3d}} \left( \partial_t + \widehat{U}_t \right) \mathbf{f}(\widehat{\mathbf{x}}) \widehat{\psi}(t, \widehat{\mathbf{x}} \big| s, \widehat{\mathbf{y}}) d\widehat{\mathbf{x}} = \int_{\widehat{\mathcal{L}}_t^d \times \mathbb{R}^{3d}} \mathbf{f}(\widehat{\mathbf{x}}) \left( -\partial_t + \widehat{U}_t^{\dagger} \right) \widehat{\psi}(t, \widehat{\mathbf{x}} \big| s, \widehat{\mathbf{y}}) d\widehat{\mathbf{x}}$$

Using the previous Lemma, the forward Kolmogorov equation of the NELD (16) follows.  $\Box$ 

Note that the probability density of  $\widehat{\mathbf{X}}_t$ 

$$\nu(t,\widehat{\mathbf{x}}) = \int_{\widehat{\mathcal{L}}_t^d \times \mathbb{R}^{3d}} \widehat{\psi}(t,\widehat{\mathbf{x}} | s, \widehat{\mathbf{y}}) \nu(s,\widehat{\mathbf{y}}) d\widehat{\mathbf{y}} = \int_{\mathcal{L}_0^d \times \mathbb{R}^{3d}} \psi(t,\overline{\mathbf{x}} | s,\overline{\mathbf{y}}) (\nu \circ \Phi_s)(s,\overline{\mathbf{y}}) d\overline{\mathbf{y}}$$

satisfies the forward Kolmogorov equation, thus we denote probability density of  $\overline{\mathbf{X}}_t$  by

$$\left(\nu \circ \Phi_t\right)(t, \overline{\mathbf{x}}) = \int_{\mathcal{L}_0^d \times \mathbb{R}^{3d}} \psi(t, \overline{\mathbf{x}} \big| 0, \overline{\mathbf{y}}) \nu_0(\overline{\mathbf{y}}) d\overline{\mathbf{y}} = \int_{\widehat{\mathcal{L}}_t^d \times \mathbb{R}^{3d}} \widehat{\psi}(t, \widehat{\mathbf{x}} \big| 0, \overline{\mathbf{y}}) \nu_0(\overline{\mathbf{y}}) d\overline{\mathbf{y}},$$

where  $\nu_0 = \nu(0, \overline{\mathbf{x}})$ . Moreover, the backward evolution from  $(\phi \circ \Phi_t)(t, \overline{\mathbf{x}})$  to  $(\phi \circ \Phi_s)(s, \overline{\mathbf{y}})$  satisfies

$$(\phi \circ \Phi_s)(s, \overline{\mathbf{y}}) = \int_{\mathcal{L}_0^d \times \mathbb{R}^{3d}} (\phi \circ \Phi_t)(t, \overline{\mathbf{x}}) \psi(t, \overline{\mathbf{x}} | s, \overline{\mathbf{y}}) d\overline{\mathbf{x}}.$$

3.2. Convergence of NELD to a Limit Cycle. We show in this section that the Markov process  $\overline{\mathbf{X}}_t$  converges to a limit cycle in the remapped Lagrangian domain, and with the following lemma that the convergence of  $\overline{\mathbf{X}}_t$  to a limit cycle implies the convergence of  $\widehat{\mathbf{X}}_t$  to a limit cycle in the remapped Eulerian domain as well.

**Lemma 3.** Let us assume that the Markov process  $\overline{\mathbf{X}}_t$  converges uniformly to a limit cycle  $\psi$  in the remapped Lagrangian domain and that  $\psi$  is smooth and positive. Then the Markov process  $\widehat{\mathbf{X}}_t$  converges uniformly to a probability density function  $\widehat{\psi}$  in the remapped Eulerian domain.

*Proof.* Since the density  $\psi$  is smooth and positive by assumption, we have

$$\begin{split} \mathbb{E}^{s,\mathbf{y}}(\mathbf{f}\circ\Phi_t)(\overline{\mathbf{X}}_t) &= \int_{\mathcal{L}_0^d\times\mathbb{R}^{3d}} (\mathbf{f}\circ\Phi_t)(\overline{\mathbf{x}})\psi(t,\overline{\mathbf{x}}\big|s,\overline{\mathbf{y}})d\overline{\mathbf{x}} \\ &= \int_{\widehat{\mathcal{L}}_t^d\times\mathbb{R}^{3d}} \mathbf{f}(\widehat{\mathbf{x}})(\psi\circ\Phi_t^{-1})(t,\widehat{\mathbf{x}}\big|s,\widehat{\mathbf{y}})d\widehat{\mathbf{x}} \\ &= \int_{\widehat{\mathcal{L}}_t^d\times\mathbb{R}^{3d}} \mathbf{f}(\widehat{\mathbf{x}})\widehat{\psi}(t,\widehat{\mathbf{x}}\big|s,\widehat{\mathbf{y}})d\widehat{\mathbf{x}}, \end{split}$$

where  $\widehat{\psi} = \psi \circ \Phi_t^{-1}$ . In addition,  $\widehat{\psi}(t, \widehat{\mathbf{x}} | \cdot)$  is smooth and positive as  $\psi(t, \overline{\mathbf{x}} | \cdot)$  is.

Now, let us consider  $\overline{\mathbf{x}}_k = (Q_k = \overline{\mathbf{q}}_{kT}, P_k = \overline{\mathbf{p}}_{kT}) \in \mathcal{L}_0^d \times \mathbb{R}^{3d}$ , so that  $(\overline{\mathbf{x}}_k)_{k\geq 0}$  denotes a discrete Markov chain constructed from the particle coordinates at the start of each each period, where  $\overline{\mathbf{x}}_0 = (Q_0, P_0)$  is the initial coordinate of the time-inhomogeneous process  $\overline{\mathbf{x}}_t = (\overline{\mathbf{q}}_t, \overline{\mathbf{p}}_t)$ . Then, we define

$$(\mathcal{U}_T \mathbf{f})(\overline{\mathbf{q}}, \overline{\mathbf{p}}) = \mathbb{E}\Big(\mathbf{f}(Q_{k+1}, P_{k+1})|(Q_k, P_k) = (\overline{\mathbf{q}}, \overline{\mathbf{p}})\Big),$$

the discrete generator of the Markov chain. In addition, we consider the Lyapunov function

$$\mathcal{K}_n(\overline{\mathbf{q}}, \overline{\mathbf{p}}) = 1 + \|\overline{\mathbf{p}}\|^{2n}, n \ge 1$$
(17)

with the associated weighted  $L^{\infty}$  norms defined by

$$\|\mathbf{g}\|_{L^{\infty}_{\mathcal{K}_n}} = \left\| \frac{\mathbf{g}}{\mathcal{K}_n} \right\|_{L^{\infty}}, \quad \mathbf{g}(\overline{\mathbf{q}}, \overline{\mathbf{p}}) \in \mathcal{L}^d_0 imes \mathbb{R}^{3d},$$

and the corresponding  $\|\mathbf{g}\|_{L^{\infty}(L^{\infty}_{\mathcal{K}_{n}})}$  norms defined by

$$\|\mathbf{g}\|_{L^{\infty}(L^{\infty}_{\mathcal{K}_n})} = \sup_{\theta \in T\mathbb{T}} \|\mathbf{g}(\theta)\|_{L^{\infty}_{\mathcal{K}_n}}, \quad \mathbf{g}(\theta, \overline{\mathbf{q}}, \overline{\mathbf{p}}) \in T\mathbb{T} \times \mathcal{L}^d_0 \times \mathbb{R}^{3d}.$$

In the remainder of this section, we prove the following propositions:

**Proposition 1.** (Uniform convergence to a limit cycle) For n > 1, there exists a unique probability measure  $(\nu \circ \Phi_{\theta})(\theta, \overline{\mathbf{q}}, \overline{\mathbf{p}})$  on  $T\mathbb{T} \times \mathcal{L}_{0}^{d} \times \mathbb{R}^{3d}$  and constants  $C_{n}, \lambda_{n}$  such that, for any initial distribution  $(\mathbf{q}_{0}, \mathbf{p}_{0})$ , we have:

(1) Exponential convergence

$$\forall \mathbf{f} \in L^{\infty}(L^{\infty}_{\mathcal{K}_n}), \quad \left| \mathbb{E} \left( (\mathbf{f} \circ \Phi_t)(\overline{\mathbf{q}}_t, \overline{\mathbf{p}}_t) \right) - \overline{\mathbf{f}}([t]) \right| \le C_n e^{-\lambda_n t} \left\| \mathbf{f} - \overline{\mathbf{f}}([t]) \right\|_{L^{\infty}(L^{\infty}_{\mathcal{K}_n})},$$

where, for  $\theta \in T\mathbb{T}$ , the spatial average of **f** reads

$$\overline{\mathbf{f}}(\theta) = \int_{\mathcal{L}_0^d \times \mathbb{R}^{3d}} \left( \mathbf{f} \circ \Phi_\theta \right) (\overline{\mathbf{q}}, \overline{\mathbf{p}}) \left( \nu \circ \Phi_\theta \right) (\theta, \overline{\mathbf{q}}, \overline{\mathbf{p}}) d\overline{\mathbf{q}} d\overline{\mathbf{p}}.$$
(18)

(2) The invariant distribution  $(\nu \circ \Phi_{\theta})(\theta, \overline{\mathbf{q}}, \overline{\mathbf{p}})$  is smooth, positive, and satisfies the Fokker-Planck equation

$$(-\partial_{\theta} + U_{\theta}^{\dagger}) \big( \nu \circ \Phi_{\theta} \big) = 0, \quad \int_{T\mathbb{T} \times \mathcal{L}_{0}^{d} \times \mathbb{R}^{3d}} \big( \nu \circ \Phi_{\theta} \big) (\theta, \overline{\mathbf{q}}, \overline{\mathbf{p}}) d\theta d\overline{\mathbf{q}} d\overline{\mathbf{p}} = T$$

(3) The invariant distribution  $(\nu \circ \Phi_{\theta})(\theta, \overline{\mathbf{q}}, \overline{\mathbf{p}})$  has finite moments of order 2n uniform in the time variable

$$\theta \in T\mathbb{T}, \quad \int_{\mathcal{L}_0^d \times \mathbb{R}^{3d}} \mathcal{K}_n(\overline{\mathbf{q}}, \overline{\mathbf{p}}) \big( \nu \circ \Phi_\theta \big)(\theta, \overline{\mathbf{q}}, \overline{\mathbf{p}}) d\overline{\mathbf{q}} d\overline{\mathbf{p}} \le R_n < \infty$$

and has uniform marginals in the time variable:

$$\overline{\nu}(\theta) = \int_{\mathcal{L}_0^d \times \mathbb{R}^{3d}} \left( \nu \circ \Phi_\theta \right) (\theta, \overline{\mathbf{q}}, \overline{\mathbf{p}}) d\overline{\mathbf{q}} d\overline{\mathbf{p}} = 1.$$

The proof of Proposition 1 is completed once we show the smoothness and positivity of the transition probability (Section 3.2.1), the uniformity of the Lyapunov condition and the uniform minorization conditions of the generator of the Markov chain (Section 3.2.2). Then, we derive the convergence in the Law of Large Numbers of the Markov chain in Section 3.2.3.

Using the above result, we derive the following Proposition:

**Proposition 2.** Let us consider  $\mathbf{f} \in L^{\infty}(L^{\infty}_{\mathcal{K}_n})$ . From all initial positions, we have

$$\frac{1}{t} \int_0^t (\mathbf{f} \circ \Phi_s)(\overline{\mathbf{q}}_s, \overline{\mathbf{p}}_s) ds \xrightarrow[t \to +\infty]{} \int_{\mathcal{L}_0^d \times \mathbb{R}^{3d}} (\mathbf{f} \circ \Phi_t)(\overline{\mathbf{q}}, \overline{\mathbf{p}})(\nu \circ \Phi_t)(t, \overline{\mathbf{q}}, \overline{\mathbf{p}}) d\overline{\mathbf{q}} d\overline{\mathbf{p}} \quad a.s..$$
(19)

3.2.1. Smoothness and positivity of the transition probability. First, we show the smoothness of the transition kernel  $\nu_t(\cdot)$  in the remapped Eulerian coordinates using [33, Lemma 22.2.5], then we show its positivity using [31].

Let  $H_s^{loc}$  denote the local Sobolev space of index s. We will use the following Lemma:

**Lemma 4.** [33, Lemma 22.2.5] If  $\hat{U}_t^{\dagger}$  is hypoelliptic then

 $\widehat{U}_t^{\dagger}g = h \text{ and } h \in H_s^{loc} \text{ at } (\widehat{\mathbf{x}}, \widehat{\mathbf{y}}) \in (\widehat{\mathcal{L}}_t^d \times \mathbb{R}^{3d}, \widehat{\mathcal{L}}_t^d \times \mathbb{R}^{3d}) \implies g \in H_{s+\epsilon}^{loc} \text{ at } (\widehat{\mathbf{x}}, \widehat{\mathbf{y}}).$ 

Then it follows that:

**Corollary 1.** Let us assume that  $\widehat{U}_t^{\dagger}$  is hypoelliptic and that there exists  $\nu_t$  such that  $\left(-\partial_t + \widehat{U}_t^{\dagger}\right)\nu_t = 0.$ 

Then  $\nu_t(\cdot) \in C^{\infty}$ .

*Proof.* We observe that if  $\hat{U}_t^{\dagger}$  is hypoelliptic then, Lemma 4 shows that  $\nu_t(\cdot) \in C^{\infty}$ .  $\Box$ 

We finish the first part of the section by showing that  $\widehat{U}_t$  is hypoelliptic in the following Lemma:

**Lemma 5.**  $\hat{U}_t, \hat{U}_t^{\dagger}$  are hypoelliptic.

*Proof.* We rewrite the generator of the NELD in the time-periodic domain as follows:

$$\widehat{U}_t = \widehat{\mathcal{X}}_0 + \frac{1}{2} \sum_{i=1}^d \widehat{\mathcal{X}}_i, \text{ for } 1 \le i \le d, \text{ and}$$

$$\widehat{\mathcal{X}}_{0} = \left\langle \widehat{\mathbf{p}} + A\widehat{\mathbf{q}}, \nabla_{\widehat{\mathbf{q}}} \right\rangle + \left\langle -\nabla V(\widehat{\mathbf{q}}), \nabla_{\widehat{\mathbf{p}}} \right\rangle - \gamma \left\langle \widehat{\mathbf{p}}, \nabla_{\widehat{\mathbf{p}}} \right\rangle, \ \widehat{\mathcal{X}}_{i} = \sqrt{\frac{2\gamma}{\beta}} \partial_{\widehat{\mathbf{p}}_{i}}.$$

Then, we define  $\mathfrak{L}(\widehat{\mathcal{X}}_0, \ldots, \widehat{\mathcal{X}}_d)$ , the Lie algebra of the family of the vectorial space operators  $(\widehat{\mathcal{X}}_0, \ldots, \widehat{\mathcal{X}}_d) \in \operatorname{Span}(\widehat{\mathcal{X}}_0, \ldots, \widehat{\mathcal{X}}_d)$  satisfying the stability property:

$$B \in \mathfrak{L}(\widehat{\mathcal{X}}_0, \dots, \widehat{\mathcal{X}}_d) \implies [B, \widehat{\mathcal{X}}_i] \in \mathfrak{L}(\widehat{\mathcal{X}}_0, \dots, \widehat{\mathcal{X}}_d), \quad i = 0, \dots, d,$$

where the Lie bracket between two operators  ${\mathscr C}$  and  ${\mathscr D}$  is

$$[\mathscr{C},\mathscr{D}] = \mathscr{C}\mathscr{D} - \mathscr{D}\mathscr{C}.$$

Since for every point  $(\widehat{\mathbf{q}}, \widehat{\mathbf{p}}) \in \widehat{\mathcal{L}}_t^d \times \mathbb{R}^{3d}$ , we have

$$[\widehat{\mathcal{X}}_i, \widehat{\mathcal{X}}_0] = \sqrt{\frac{\gamma}{2\beta}} (\partial_{\widehat{\mathbf{q}}_i} + \gamma) \partial_{\widehat{\mathbf{p}}_i}, \quad \forall i \in \{1 \dots d\},$$

evaluated at  $(\widehat{\mathbf{q}}_0, \widehat{\mathbf{p}}_0)$  span  $\mathbb{R}^{3d}$ , it follows that  $\widehat{U}_t$  and  $\widehat{U}_t^{\dagger}$  are hypoelliptic using [33, Theorem 1.1]. Then it follows that  $\widehat{U}_t$  and  $\widehat{U}_t^{\dagger}$  are hypoelliptic as well.

Now let us prove that the generator in the remapped Lagrangian coordinates has a positive probability density. We consider the following Lemma:

**Lemma 6.**  $U_t$  has a positive transition kernel.

*Proof.* For t > 0 and two points  $(\overline{\mathbf{q}}_0, \overline{\mathbf{p}}_0)$  and  $(\overline{\mathbf{q}}_t, \overline{\mathbf{p}}_t)$ , let us consider  $\varphi(t)$  be any  $\mathcal{C}^2$  path in  $\mathcal{L}_0^d \times \mathbb{R}^{3d}$  which satisfies  $\varphi(0) = \overline{\mathbf{q}}_0$ ,  $\varphi(t) = \overline{\mathbf{q}}_t$ ,  $\varphi'(0) = \overline{\mathbf{p}}_0$ , and  $\varphi'(t) = \overline{\mathbf{p}}_t$ . Then we can rewrite the NELD equation as

$$\mathcal{V}_t = \sqrt{\frac{2\beta}{\gamma}} e^{[t]A} \Big( \ddot{\varphi}_t + \nabla V(e^{[t]A}\varphi_t) + \Gamma \dot{\varphi}_t \Big),$$

where  $\mathcal{V}_t$  is a smooth control. Thus  $(\varphi_t, \dot{\varphi}_t)$  is a solution of the control system so that,  $U_t$  drives the system from  $(\overline{\mathbf{q}}_0, \overline{\mathbf{p}}_0)$  to  $(\overline{\mathbf{q}}_t, \overline{\mathbf{p}}_t)$ . This implies that the support of the transition kernel  $\mathcal{A}_t(\overline{\mathbf{q}}, \overline{\mathbf{p}}) = \mathcal{L}_0^d \times \mathbb{R}^{3d}, \forall s > 0, \ (\overline{\mathbf{q}}, \overline{\mathbf{p}}) \in \mathcal{L}_0^d \times \mathbb{R}^{3d}$ . Thus by [31, Corollary 6.2], it follows that the transition kernel is positive.

Now, let us denote  $\mathcal{B}_{\delta}(x)$  the open ball of raduis  $\delta$  centered at x. We summarize the results of this section in the following Corollary:

**Corollary 2.** The Markov process  $\overline{\mathbf{x}}$  with transition kernel  $\overline{\mathcal{P}}_t(\overline{\mathbf{x}}, \overline{B})$  satisfies, for some fixed compact set  $C \in \mathcal{B}(\mathcal{L}_0^d \times \mathbb{R}^{3d})$ , the following:

• for some  $z^* \in int(C)$  there is, for any  $\delta > 0$ , a  $t_1 = t_1(\delta) \in T\mathbb{T}$  such that

$$\overline{\mathcal{P}}_{t_1}(\overline{\mathbf{x}}, \mathcal{B}_{\delta}(z^*)) > 0, \quad \forall \overline{\mathbf{x}} \in C$$

• for  $t \in T\mathbb{T}$  the transition kernel possesses a density  $\psi(\overline{\mathbf{x}}, \overline{\mathbf{y}})$  precisely

$$\overline{\mathcal{P}}_t(\overline{\mathbf{x}},\overline{B}) = \int_{\overline{B}} \psi(\overline{\mathbf{x}},\overline{\mathbf{y}}) d\overline{\mathbf{y}}, \ \forall \overline{\mathbf{x}} \in C, \ \overline{B} \in \mathcal{B}(\mathcal{L}_0^d \times \mathbb{R}^{3d}) \cap \mathcal{B}(C).$$

and  $\psi(\overline{\mathbf{x}}, \overline{\mathbf{y}})$  is jointly continuous in  $(\overline{\mathbf{x}}, \overline{\mathbf{y}}) \in C \times C$ .

*Proof.* The proof of the first argument is based on the positivity of the transition kernel from Lemma 6 and the second is based on the smoothness of density from Corollary 1.  $\Box$ 

We use the above Corollary in the next section to show that the Lyanpunov function satisfies the minorization condition.

3.2.2. The Invariant Measure of the Discrete Process. The convergence of the Markov chain  $(Q_{k+1}, P_{k+1})$  is based on the uniform Lyapunov condition [28, Assumption 1] and the uniform minorization condition [28, Assumption 2] that we prove in the following two Lemmas.

**Lemma 7.** (Uniform Lyapunov condition) There exists  $a_n \in [0,1)$  and  $b_n > 0$  such that

$$\mathcal{U}_T \mathcal{K}_n \le a_n \mathcal{K}_n + b_n,\tag{20}$$

for  $\mathcal{K}_n$  defined in (17).

*Proof.* First, let us recall the NELD in the remapped Lagrangian coordinates under the remappings as follows:

$$\begin{cases} d\overline{\mathbf{q}}_{kT+\theta} &= \overline{\mathbf{p}}_{kT+\theta} d\theta, \\ d\overline{\mathbf{p}}_{kT+\theta} &= -e^{-\theta A} \nabla V \left( e^{\theta A} \overline{\mathbf{q}}_{kT+\theta} \right) d\theta - \Gamma \overline{\mathbf{p}}_{kT+\theta} d\theta + \sigma e^{-\theta A} dW_{kT+\theta}, \end{cases}$$
(21)

where  $(\theta, \overline{\mathbf{q}}_{kT+\theta}, \overline{\mathbf{p}}_{kT+\theta}) \in T\mathbb{T} \times \mathcal{L}_0^d \times \mathbb{R}^{3d}$ , and  $\Gamma = (\gamma + A)$ . Multiplying the second equation of (21) by the integrating factor  $e^{\Gamma\theta}$ , we get we have

$$d\left(e^{\Gamma\theta}\overline{\mathbf{p}}_{kT+\theta}\right) = e^{\gamma\theta} (-\nabla V(e^{\theta A}\overline{\mathbf{q}}_{kT+\theta})d\theta + \sigma dW_{kT+\theta}).$$
(22)

We integrate (22) over a period to get the evolution of  $\overline{\mathbf{p}}$  up to, but not including the remapping, finding

$$e^{\Gamma T} P_{k+1}^{-} - P_{k} = \int_{0}^{T} e^{\gamma \theta} (-\nabla V (e^{\theta A} \overline{\mathbf{q}}_{kT+\theta}) d\theta + \sigma dW_{kT+\theta}).$$

We multiply the above equation by  $e^{-\Gamma T}$  to get

$$P_{k+1}^{-} = e^{-\Gamma T} (P_k + \mathscr{F}_k + \mathscr{G}_k),$$

where

$$\mathscr{F}_k = \int_0^T e^{\gamma \theta} \nabla V(e^{\theta A} \overline{\mathbf{q}}_{kT+\theta}) d\theta, \quad \mathscr{G}_k = \sqrt{\frac{2\gamma}{\beta}} \int_0^T e^{\gamma \theta} dW_{kT+\theta}.$$

We apply the remapping to the momentum to obtain

$$P_{k+1} = e^{TA} P_{k+1}^{-} = e^{-\gamma T} (P_k + \mathscr{F}_k + \mathscr{G}_k).$$

Letting  $\alpha = e^{-\gamma T}$ , we have  $0 \leq \alpha < 1$ . Also, note that  $\mathscr{G}_k$  has mean zero and covariance  $\frac{1}{\beta}(e^{2\gamma T}-1)$  as

$$\mathbb{E}\left[\left|\mathscr{G}_{k}^{2}\right|\mathcal{F}_{k}\right] = \frac{1}{\beta}\mathbb{E}\left[\left|2\gamma\left(\int_{0}^{T}e^{\gamma\theta}\right)dW_{kT+\theta}\right|^{2}\left|\mathcal{F}_{k}\right]\right] = \frac{1}{\beta}\left(e^{2\gamma T}-1\right).$$

Since  $\left\| e^{\gamma \theta} \nabla V \left( e^{\theta A} \overline{\mathbf{q}}_{kT+\theta} \right) \right\|_2 \leq C$ , we have

$$\left\|\mathscr{F}_{k}\right\|_{2} \leq \alpha \left\|e^{-AT}\right\|_{2} TC < \infty.$$

Using the previous results, we get

$$\begin{aligned} |P_{k+1}||_{2}^{2} &\leq \alpha^{2} \, \|P_{k} + \mathscr{F}_{k} + \mathscr{G}_{k}\|_{2}^{2} \leq \alpha^{2} (\|P_{k}\|_{2}^{2} + \|\mathscr{F}_{k} + \mathscr{G}_{k}\|_{2}^{2} + 2 \, \langle P_{k}, \mathscr{F}_{k} + \mathscr{G}_{k} \rangle) \\ &\leq \alpha^{2} ((1+\mu) \, \|P_{k}\|_{2}^{2} + \left(2 + \frac{1}{\mu}\right) \, \|\mathscr{F}_{k}\|_{2}^{2} + 2 \, \langle P_{k}, \mathscr{G}_{k} \rangle + 2 \, \|\mathscr{G}_{k}\|_{2}^{2}), \end{aligned}$$

where we choose  $\mu$  s.t.  $\alpha^2(1+\mu) < 1$ . Since  $\mathbb{E}\left[ \langle P_k, \mathscr{G}_k \rangle | \mathcal{F}_k \right] = 0$ , taking the expectation of  $\mathcal{K}_2$ , we get

$$\mathbb{E}\Big[\left|\mathcal{K}_2(P_{k+1},Q_{k+1})\right|\mathcal{F}_k\Big] \leq \alpha^2(1+\mu)\mathcal{K}_2(P_k,Q_k) + C_\mu,$$

where  $C_{\mu} = \left(2 + \frac{1}{4\mu}\right) \|\mathscr{F}_k\|_2^2 + 2 \|\mathscr{G}_k\|_2^2$ . This leads us to find a bound on the expectation of  $\mathcal{K}_n$  as

$$\mathbb{E}\Big[\left|\mathcal{K}_{n}(P_{k+1},Q_{k+1})\right|\mathcal{F}_{k}\Big] \leq \left(\alpha^{2}(1+\mu)\mathcal{K}_{2}(P_{k},Q_{k}) + C_{\mu}\right)^{n} \leq \alpha^{2n}(1+\mu)^{n} \|P_{k}\|_{2}^{2n} + \mathfrak{p}(|P_{k}|),$$

where  $\mathfrak{p}$  is a polynomial of degree at most 2(n-1), with positive coefficient. Since we have

$$\frac{\mathfrak{p}(|P_k|)}{\|P_k\|_2^{2n}} \to 0, \text{ as } \|P_k\|_2 \to \infty \implies \mathfrak{p}(|P_k|) = \delta \|P_k\|_2^{2n} + C_\delta,$$

and for  $\delta$  small, finally we get the uniform bound

$$\mathbb{E}\Big[\left|\mathcal{K}_n(P_{k+1},Q_{k+1})\right|\mathcal{F}_k\Big] \leq a \, \|P_k\|_2^{2n} + C_{\delta}.$$

where  $a = \alpha^{2n} (1 + \mu)^n + \delta < 1$ .

**Lemma 8.** (Uniform minorization condition) Fix any  $p_{max} > 0$ , then there exists a probability measure  $\vartheta \in \mathcal{L}_0^d \times \mathbb{R}^{3d}$  and a constant  $\kappa$  such that,

$$\forall \overline{B} \in \mathcal{B}(\mathcal{L}_0^d \times \mathbb{R}^{3d}), \quad \mathbb{P}\Big((P_{k+1}, Q_{k+1}) \in \overline{B} \mid \|P_k\|_2 \le p_{max}\Big) \ge \kappa \vartheta(\overline{B}).$$

The proof is essentially based on the arguments from [25, 31] which uses the continuity property of the Markov process, the irregularity and positivity of the transition kernel. Before we start the proof, we consider the following Lemma:

**Lemma 9.** [25, Lemma 2.3] If the Markov process  $\overline{\mathbf{x}}$  satisfies the assumption in Corollary 2, then there is a choice of  $t \in T\mathbb{T}$ , an  $\kappa \geq 0$ , and a probability measure  $\vartheta$ , with  $\vartheta(C^c) = 0$  and  $\vartheta(C) = 1$ , such that

$$\overline{\mathcal{P}}(\overline{\mathbf{x}},\overline{B}) \ge \kappa \vartheta(\overline{B}), \quad \forall \overline{B} \in \mathcal{B}(\mathcal{L}_0^d \times \mathbb{R}^{3d}), \ \overline{\mathbf{x}} \in C.$$
(23)

Then the proof of Lemma 8 is following:

*Proof.* Since the discrete chain  $(P_{k+1}, Q_{k+1}) \in \overline{B}$  satisfies the assumption in Corollary 2, then using the above Lemma, we have the expected result.  $\Box$ 

Using the previous Lemmas, we state the following uniform convergence result for the sampled chain  $(Q_k, P_k)$  from [28]:

**Theorem 1.** [28, Theorem 1.2] If  $\mathcal{U}_T$  satisfies the Lyapunov condition as in Lemma 7 and the minorization condition as in Lemma 8, then  $\mathcal{U}_T$  admits a unique invariant measure  $\nu$ such that for  $C_n, \lambda_n > 0$ ,

$$\overline{\mathbf{f}} = \int_{\mathcal{L}_0^d \times \mathbb{R}^{3d}} \mathbf{f}(\overline{\mathbf{q}}, \overline{\mathbf{p}}) \nu(\overline{\mathbf{q}}, \overline{\mathbf{p}}) d\overline{\mathbf{q}} d\overline{\mathbf{p}},$$
$$\left\| \mathcal{U}_T^k \mathbf{f} - \overline{\mathbf{f}} \right\|_{L_{\mathcal{K}_n}^\infty} \le C_n e^{-\lambda_n kT} \left\| \mathbf{f} - \overline{\mathbf{f}} \right\|_{L_{\mathcal{K}_n}^\infty}.$$
(24)

Then, we derive the convergence of the continuous process  $(\overline{\mathbf{q}}_t, \overline{\mathbf{p}}_t)$  in the following Lemma:

**Lemma 10.** The Markov process  $(\overline{\mathbf{q}}_t, \overline{\mathbf{p}}_t)$  converges exponentially to the limit cycle  $\nu \circ \Phi_t$ :

$$\left| \mathbb{E}^{s,\mathbf{y}} \Big( \mathbf{f} \circ \Phi_t \Big) (\overline{\mathbf{x}}_t) - \overline{\mathbf{f}}([t]) \right| \le C_n e^{-\lambda_n t} \left\| \mathbf{f} - \overline{\mathbf{f}}([t]) \right\|_{L^{\infty}_{\mathcal{K}_n}} \Big( 1 + \mathcal{K}_n(\mathbf{y}) \Big), \quad \mathbf{y} = \mathbf{x}_0,$$

where  $\overline{\mathbf{f}}([t])$  is defined in (18).

*Proof.* We work in  $(\widehat{\mathbf{q}}, \widehat{\mathbf{p}})$  variables to simplify the proof. We use an argument from [25,36] and the result from Theorem 1 to show that the Markov process  $(\widehat{\mathbf{q}}_{kT+\theta}, \widehat{\mathbf{p}}_{kT+\theta}) \in \widehat{\mathcal{L}}_t^d \times \mathbb{R}^{3d}$  converges exponentially to a limit cycle. The result implies immediately that the Markov process  $(\mathbf{q}_{kT+\theta}, \mathbf{p}_{kT+\theta}) \in \mathcal{L}_0^d \times \mathbb{R}^{3d}$  converges as well. We start by using the result from Lemma 7 and Lemma 8, and derive from Theorem 1 that

$$\left|\mathbb{E}^{s,\mathbf{y}}\mathbf{f}(\widehat{\mathbf{x}}_{kT}) - \overline{\mathbf{f}}\right| \le C_n e^{-\lambda_n kT} \left\|\mathbf{f} - \overline{\mathbf{f}}\right\|_{L^{\infty}_{\mathcal{K}_n}} \mathcal{K}_n(\mathbf{y}), \quad \overline{\mathbf{f}} = \int_{\widehat{\mathcal{L}}^d_{\theta} \times \mathbb{R}^{3d}} \mathbf{f}(\widehat{\mathbf{q}}, \widehat{\mathbf{p}}) \nu(\widehat{\mathbf{q}}, \widehat{\mathbf{p}}) d\widehat{\mathbf{q}} d\widehat{\mathbf{p}},$$

where we rewrite (24) in remapped Eulerian coordinates. Conditioning on  $\mathcal{F}_{\theta}$ , we have

$$\left|\mathbb{E}^{s,\mathbf{y}}\mathbf{f}(\widehat{\mathbf{x}}_{kT+\theta}) - \overline{\mathbf{f}}(\theta)\right| \le C_n e^{-\lambda_n kT} \left\|\mathbf{f} - \overline{\mathbf{f}}(\theta)\right\|_{L^{\infty}_{\mathcal{K}_n}} \mathbb{E}^{s,\mathbf{y}}\mathcal{K}_n(\widehat{\mathbf{x}}_{\theta}).$$
(25)

We compute an upper bound on  $\mathbb{E}^{s,\mathbf{y}}\mathcal{K}_n(\widehat{\mathbf{x}}_{\theta})$  as follows: using the Itô's lemma, we get

$$d\mathbf{f}(\widehat{\mathbf{X}}_t) = \left(\widehat{U}_t \mathbf{f}\right)(\widehat{\mathbf{X}}_t) dt + \left\langle \nabla \mathbf{f}(\widehat{\mathbf{X}}_t), \widetilde{\Sigma} dW \right\rangle,$$

where

$$\widehat{U}_t = \left\langle \widehat{\mathbf{p}} + A\widehat{\mathbf{q}}, \nabla_{\widehat{\mathbf{q}}} \right\rangle + \left\langle -\nabla V(\widehat{\mathbf{q}}), \nabla_{\widehat{\mathbf{p}}} \right\rangle - \gamma \left\langle \widehat{\mathbf{p}}, \nabla_{\widehat{\mathbf{p}}} \right\rangle + \frac{1}{2}\sigma\sigma^T : \nabla^2$$

Then, it follows that

$$\widehat{U}_{t}\mathcal{K}_{n}(\widehat{\mathbf{q}},\widehat{\mathbf{p}}) = -n \|\widehat{\mathbf{p}}\|_{2}^{n-2} \langle \nabla V(\widehat{\mathbf{q}}),\widehat{\mathbf{p}}\rangle - n\gamma \Big( \langle \widehat{\mathbf{p}},\widehat{\mathbf{p}}\rangle - \frac{n+d-2}{\beta} \Big) \|\widehat{\mathbf{p}}\|_{2}^{n-2} \\ \leq -n\gamma \|\widehat{\mathbf{p}}\|_{2}^{n} + n \|\nabla V(\widehat{\mathbf{q}})\|_{2} \|\widehat{\mathbf{p}}\|_{2}^{n-1} + n\gamma \frac{n+d-2}{\beta} \|\widehat{\mathbf{p}}\|_{2}^{n-2}.$$

Thus, there exists  $\widehat{a}_n, \widehat{b}_n \geq 0$  such that

$$\widehat{U}_t \mathcal{K}_n \leq -\widehat{a}_n \mathcal{K}_n + \widehat{b}_n, \quad \widehat{a}_n = n\gamma, \quad \text{as } \lim_{|\widehat{\mathbf{q}},\widehat{\mathbf{p}}| \to \infty} \frac{\widehat{U}_t \mathcal{K}_n}{\mathcal{K}_n} \leq -\widehat{a}_n,$$

and it follows that

$$d\mathcal{K}_n(\widehat{\mathbf{x}}_{kT+\theta}) \leq (-\widehat{a}_n\mathcal{K}_n + \widehat{b}_n)d\theta + \text{Martingle}.$$

Then, we get an upper bound on  $\mathbb{E}^{s,\mathbf{y}}\mathcal{K}_n(\widehat{\mathbf{x}}_{\theta})$  by using Grönwall's inequality:

$$\mathbb{E}^{s,\mathbf{y}}\mathcal{K}_n(\widehat{\mathbf{x}}_{\theta}) \leq e^{-\widehat{a}_n\theta}\mathcal{K}_n(\mathbf{y}) + \frac{\widehat{b}_n}{\widehat{a}_n} \left(1 - e^{-\widehat{a}_n\theta}\right) \leq e^{-\widehat{a}_n\theta}\mathcal{K}_n(\mathbf{y}) + \frac{\widehat{b}_n}{\widehat{a}_n}.$$

Plugging the latter result in (25), we have

$$\left|\mathbb{E}^{s,\mathbf{y}}\mathbf{f}(\widehat{\mathbf{x}}_{kT+\theta}) - \overline{\mathbf{f}}(\theta)\right| \le C_n e^{-\lambda_n kT} \left\|\mathbf{f} - \overline{\mathbf{f}}(\theta)\right\|_{L^{\infty}_{\mathcal{K}_n}} \left(e^{-\widehat{a}_n \theta} \mathcal{K}_n(\mathbf{y}) + \frac{b_n}{\widehat{a}_n}\right).$$

Defining  $\lambda_n$  by  $e^{-\lambda_n} = \widehat{a}_n^{\frac{1}{T}}$ , we obtain the expected result by redefining  $C_n \to \left(1 + \frac{\widehat{b}_n}{\widehat{a}_n} e^{\lambda_n T}\right)$ :  $\left|\mathbb{E}^{s,\mathbf{y}}\mathbf{f}(\widehat{\mathbf{x}}_{kT+\theta}) - \overline{\mathbf{f}}(\theta)\right| \leq C_n e^{-\lambda_n (kT+\theta)} \left\|\mathbf{f} - \overline{\mathbf{f}}(\theta)\right\|_{L^{\infty}_{\mathcal{K}_n}} \left(1 + \mathcal{K}_n(\mathbf{y})\right).$ 

3.2.3. Convergence in Law of Large Numbers for  $(Q_k, P_k)$ . We use Lemmas from previous section and mainly [34] to show that  $(Q_k, P_k)$  is positive Harris recurrent chain. Thus, the Law of Large Number holds:

**Proposition 3.** (Law of Large Numbers for the sampled chain) For any  $\mathbf{f} \in L^{\infty}_{\mathcal{K}_n}$ ,

$$\frac{1}{N}\sum_{k=1}^{N}\mathbf{f}(Q_k, P_k) \xrightarrow[N \to +\infty]{} \int_{\mathcal{L}_0^d \times \mathbb{R}^{3d}} \mathbf{f}(\overline{\mathbf{q}}, \overline{\mathbf{p}})\nu(\overline{\mathbf{q}}, \overline{\mathbf{p}})d\overline{\mathbf{q}}d\overline{\mathbf{p}} \quad a.s.,$$

for all the initial conditions  $(Q_0, P_0)$ .

*Proof.* Corollary 1 from the previous Section shows that the chain  $(Q_k, P_k)$  is irreducible with respect to the Lebesgues measure. Thus, the result from [35, Corollary 1] based on [34, Page 199] provides that the chain  $(Q_k, P_k)$  is Harris recurrent. In addition as  $(\nu \circ \Phi_\theta)(\theta, \overline{\mathbf{q}}, \overline{\mathbf{p}})$  is positive using Lemma 6, it follows from [34, Theorem 17.0.1] that the Law of large Numbers holds.

3.2.4. Convergence to the Limit Cycle. We consider the following Lemma:

**Lemma 11.** The Markov process  $(\overline{\mathbf{q}}_t, \overline{\mathbf{p}}_t)$  converges exponentially to the limit cycle  $\nu \circ \Phi_t$  from any initial configuration  $(\overline{\mathbf{q}}_0, \overline{\mathbf{p}}_0)$ .

*Proof.* The result from Corollary 1 and Lemma 6 show that the chain is irreducible with respect to the Lebesgues measure. In addition, the process  $(\overline{\mathbf{q}}_t, \overline{\mathbf{p}}_t)$  is Harris recurrent using the result from [35, Corollary 1] based on [34, Page 199]. Thus, the exponential convergence of the chain  $(\overline{\mathbf{q}}_t, \overline{\mathbf{p}}_t)$  to an invariant measure  $(\nu \circ \Phi_t)$  defined in Lemma 10, hold for all initial positions. Finally, using [34, Theorem 17.0.1], it follows that the Law of large Numbers defined in (19) holds as well.

Now we use the above result to prove Proposition 1.

*Proof of Proposition 1.* The result from Lemma 11 shows part 1 of Proposition 1, exponential convergence to the limit cycle of the NELD. We prove part 2 of Proposition 1 using Lemma 6 and the forward Kolmogorov equation of the NELD as follows: since  $\nu$  is a positive probability density function, Lemma 2 gives

$$(-\partial_{\theta} + U_{\theta}^{\dagger})(\nu \circ \Phi_{\theta}) = 0.$$

The last part of Proposition 1 is shown by integrating the equation (20) with respect  $\nu$ . In fact, iterating (20) gives

$$\mathbb{E}^{s,\mathbf{y}}\mathcal{K}_n(\widehat{\mathbf{x}}_{kT}) \le e^{-a_nkT}\mathcal{K}_n(\mathbf{y}) + \frac{b_n}{1-a_n} \implies \int_{\mathcal{L}_0^d \times \mathbb{R}^{3d}} \mathcal{K}_n(\nu \circ \Phi_\theta) \le \frac{b_n}{1-a_n}.$$

Now, we derive the proof of Proposition 2.

Proof of Proposition 2. The result from the previous proposition shows that the Markov process  $(\overline{\mathbf{q}}_t, \overline{\mathbf{p}}_t)$  converges exponentially to  $(\nu \circ \Phi_t)(t, \overline{\mathbf{q}}, \overline{\mathbf{p}})$  for all initial positions. In addition since the chain  $(\overline{\mathbf{q}}_t, \overline{\mathbf{p}}_t)$  is Harris recurrent ([35, Corollary 1] based on [34, Page 199]) and  $(\nu \circ \Phi_t)(t, \overline{\mathbf{q}}, \overline{\mathbf{p}})$  is positive using Lemma 6, it follows from [34, Theorem 17.0.1] that the Law of large Numbers holds.

# 4. CONCLUSION

In this paper, we study the ergodicity of NELD under shear flow and planar elongational flow using respectively LE and KR Periodic boundary conditions. This is essentially formulated in Proposition 1 where, after showing existence and uniqueness of the limit cycle using a Lyapunov function and a minorization condition, we established the exponential convergence of the Markov chain generated by the NELD equation given all the initial conditions. It will be interesting to establish the convergence analysis for the three dimensional diagonalizable incompressible flow cases using the R-KR [32] algorithm or the GenKR [6,10] algorithm. However advanced analysis will be needed since the latter PBCs's geometry is not as simple as the current's cases studied.

### Acknowledgements

The authors would like to thank Gabriel Stoltz for helpful comments on an early version of the manuscript.

### References

- [1] Lang, Philipp S. and Obermayer, Benedikt and Frey, Erwin Dynamics of a semiflexible polymer or polymer ring in shear flow. *Phys. Rev. E*, Feb 2014, 10.1103/PhysRevE.89.022606.
- [2] O'Connor, Thomas C. and Ge, Ting and Rubinstein, Michael and Grest, Gary S.. Topological Linking Drives Anomalous Thickening of Ring Polymers in Weak Extensional Flows. *Phys. Rev. Lett.*, Jan 2020, 10.1103/PhysRevLett.124.027801.
- [3] O'Connor, Thomas C. and Alvarez, Nicolas J. and Robbins, Mark O. Relating Chain Conformations to Extensional Stress in Entangled Polymer Melts. *Phys. Rev. Lett.*, Jul 2018, 10.1103/Phys-RevLett.121.047801.
- [4] Baranyai, András and Cummings, Peter T. Steady state simulation of planar elongation flow by nonequilibrium molecular dynamics. *The Journal of Chemical Physics*.
- [5] Todd, Billy D. and Daivis, Peter J. . Nonequilibrium Molecular Dynamics: Theory, Algorithms and Applications.
- [6] Thomas A. Hunt. Periodic boundary conditions for the simulation of uniaxial extensional flow of arbitrary duration, *Molecular Simulation*
- [7] Evans, Denis J. and Morriss, Gary P. Statistical mechanics of nonequilibrium liquids. ANU E Press, Canberra
- [8] A.M. Kraynik and D.A. Reinelt Extensional motions of spatially periodic lattices
- [9] A W Lees and S F Edwards The computer study of transport processes under extreme conditions. J. Phys. C Solid State, 10.1088/0022-3719/5/15/006.
- [10] Matthew Dobson. Periodic boundary conditions for long-time nonequilibrium molecular dynamics simulations of incompressible flows. The Journal of Chemical Physics, 10.1063/1.4901276.
- [11] Xu, Wen-Sheng and Carrillo, Jan-Michael Y. and Lam, Christopher N. and Sumpter, Bobby G. and Wang, Yangyang. Molecular Dynamics Investigation of the Relaxation Mechanism of Entangled Polymers after a Large Step Deformation. ACS Macro Letters, 10.1021/acsmacrolett.7b00900.
- [12] P.J. Daivis and M.L. Matin and B.D. Todd . Nonlinear shear and elongational rheology of model polymer melts by non-equilibrium molecular dynamics. *Journal of Non-Newtonian Fluid Mechanics*,10.1016/S0377-0257(03)00011-9.
- [13] Todd,B. D. and Daivis, Peter J. The stability of nonequilibrium molecular dynamics simulations of elongational flows. *The Journal of Chemical Physics*, 10.1063/1.480642.
- [14] Todd, B. D. and Daivis, Peter J. Nonequilibrium Molecular Dynamics Simulations of Planar Elongational Flow with Spatially and Temporally Periodic Boundary Conditions. *Phys. Rev. Lett.*, 10.1103/PhysRevLett.81.1118.
- [15] Oliveira, AnaSofia and Ciccotti, Giovanni and Haider, Shozeb and Mulholland, Adrian, Dynamical nonequilibrium molecular dynamics reveals the structural basis for allostery and signal propagation in biomolecular systems, *The European Physical Journal B*, July 2021,10.1140/epjb/s10051-021-00157-0
- [16] Nishioka, Akihiro and Takahashi, Tatsuhiro and Masubuchi, Yuichi and Takimoto, Jun-ichi and Koyama, Kiyohito. Description of uniaxial, biaxial, and planar elongational viscosities of polystyrene melt by the K-BKZ model. Journal of Non-Newtonian Fluid Mechanics, May 2000, 10.1016/S0377-0257(99)00047-6
- [17] Nicholson, David A. and Rutledge, Gregory C. . Molecular simulation of flow-enhanced nucleation in n-eicosane melts under steady shear and uniaxial extension, *The Journal of Chemical Physics*, 10.1063/1.4972894.
- [18] Templeton, Clark and Elber, R. and Ferrario, Mauro and Ciccotti, Giovanni. A new boundary driven NEMD scheme for heat and particle diffusion in binary mixtures. *Molecular Physics*, 10.1080/00268976.2021.1892849.
- [19] Menzel, A. G. and Daivis, P. J. and Todd, B. D. Equilibrium and nonequilibrium molecular dynamics methods to compute the first normal stress coefficient of a model polymer solution. *Phys. Rev. Fluids*, Aug 2020, 10.1103/PhysRevFluids.5.084201.
- [20] Ewen, James and Heyes, D. and Dini, Daniele, Advances in nonequilibrium molecular dynamics simulations of lubricants and additives. *Friction*, February 2018 10.1007/s40544-018-0207-9
- [21] Li, Zhen and Xiong, Shiyun and Sievers, Charles and Hu, Yue and Fan, Zheyong and Wei, Ning and Bao, Hua and Chen, Shunda and Donadio, Davide and Ala-Nissila, Tapio . Influence of thermostatting on nonequilibrium molecular dynamics simulations of heat conduction in solids. *The Journal of Chemical Physics*, 10.1063/1.5132543.

- [22] Dobson, Matthew and Legoll, Frédéric and Lelièvre, Tony and Stoltz, Gabriel, Derivation of Langevin dynamics in a nonzero background flow field. ESAIM: M2AN, 10.1051/m2an/2013077
- [23] M.G. McPhie and P.J. Daivis and I.K. Snook and J. Ennis and D.J. Evans Generalized Langevin equation for nonequilibrium systems. *Physica A*.
- [24] Talay, D.. Stochastic Hamiltonian Systems: Exponential Convergence to the Invariant Measure, and Discretization by the Implicit Euler Scheme. *Markov Processes and Related Fields*
- [25] J.C. Mattingly and A.M. Stuart and D.J. Higham . Ergodicity for SDEs and approximations: locally Lipschitz vector fields and degenerate noise. *Stochastic Processes and their Applications*, 10.1016/S0304-4149(02)00150-3.
- [26] Cancès, Eric and Legoll, Frédéric and Stoltz, Gabriel, Theoretical and numerical comparison of some sampling methods for molecular dynamics. ESAIM: M2AN,10.1051/m2an:2007014.
- [27] Joubaud, R. and Pavliotis, G. A. and Stoltz, G. Langevin Dynamics with Space-Time Periodic Nonequilibrium Forcing. *Journal of Statistical Physics*, 10.1007/s10955-014-1118-4.
- [28] Hairer, Martin and Mattingly, Jonathan Yet Another Look at Harris' Ergodic Theorem for Markov Chains.
- [29] Leimkuhler, Benedict and Matthews, Charles and Stoltz, Gabriel. The computation of averages from equilibrium and nonequilibrium Langevin molecular dynamics. *IMA Journal of Numerical Analysis*, Jan 2015, 10.1093/imanum/dru056.
- [30] Lelièvre, Tony and Stoltz, Gabriel. Partial differential equations and stochastic methods in molecular dynamics. Acta Numerica, 10.1017/S0962492916000039.
- [31] Rey-Bellet, Luc. Ergodic Properties of Markov Processes. Lecture Notes in Mathematics.
- [32] Matthew Dobson and Abdel Kader Geraldo. Simple Periodic Boundary Conditions for Molecular Simulation of Uniaxial Flow.
- [33] Hörmander, L.. The Analysis of Linear Partial Differential Operators III: Pseudo-Differential Operators. ZAMM - Journal of Applied Mathematics and Mechanics / Zeitschrift f
  ür Angewandte Mathematik und Mechanik.
- [34] Meyn, Sean and Tweedie, Richard L. and Glynn, Peter W. Markov Chains and Stochastic Stability,10.1017/CBO9780511626630, Cambridge University Press.
- [35] Luke Tierney. Markov Chains for Exploring Posterior Distributions, The Annals of Statistics, 10.1214/aos/1176325750.
- [36] Sean P. Meyn and R. L. Tweedie. Stability of Markovian Processes III: Foster-Lyapunov Criteria for Continuous-Time Processes. *Advances in Applied Probability*.
- [37] Friedman, A. Stochastic Differential Equations and Applications. Vol. 1. Probability and mathematical statistics series, vol. 28.