

Inevitability of knowing less than nothing

Gilad Gour,¹ Mark M. Wilde,² Sarah Brandsen,³ and Isabelle Jianing Geng¹

¹*Department of Mathematics and Statistics, Institute for Quantum Science and Technology,
University of Calgary, Alberta, Canada T2N 1N4*

²*School of Electrical and Computer Engineering,
Cornell University, Ithaca, New York 14850, USA*

³*Department of Physics, Duke University, Durham, North Carolina 27708, USA*

(Dated: August 31, 2022)

A colloquial interpretation of entropy is that it is the knowledge gained upon learning the outcome of a random experiment. Conditional entropy is then interpreted as the knowledge gained upon learning the outcome of one random experiment after learning the outcome of another, possibly statistically dependent, random experiment. In the classical world, entropy and conditional entropy take on non-negative values, consistent with the intuition that one has regarding the aforementioned interpretations. However, for certain entangled states, one obtains negative values when evaluating commonly accepted and information-theoretically justified formulas for the quantum conditional entropy, leading to the confounding conclusion that one can know less than nothing in the quantum world. Here, we introduce a physically motivated framework for defining quantum conditional entropy, based on two simple postulates inspired by the second law of thermodynamics (non-decrease of entropy) and extensivity of entropy, and we argue that all plausible definitions of quantum conditional entropy should respect these two postulates. We then prove that all plausible quantum conditional entropies take on negative values for certain entangled states, so that it is inevitable that one can know less than nothing in the quantum world. All of our arguments are based on constructions of physical processes that respect the first postulate, the one inspired by the second law of thermodynamics.

CONTENTS

I. Introduction	1	C. Quantum conditional majorization reduces to classical conditional majorization	12
A. Summary of results	2	D. Proofs of properties of quantum conditional entropy	13
B. Notation	2	E. A subclass of conditional entropies	14
II. Uncertainty and entropy	3	1. Proof of Theorem 4	15
A. Uncertainty measures	3	2. Proof of Theorem 1	17
B. Entropy	4	F. Proof of Theorem 2	17
III. Conditional uncertainty and entropy	5	1. Conditional min-entropy is a conditional entropy	18
A. Conditional uncertainty measures	5	G. Example of non-negative weakly additive monotonic function	18
B. Conditional entropy	6		
IV. Main result: Inevitability of negative quantum conditional entropy	7		
V. Conditional entropy from relative entropy	7		
VI. Conclusion	8		
Acknowledgments	8		
References	8		
A. Proofs of properties of entropy	10		
B. Choi matrix formulations	10		
1. Choi matrix formulation of conditional unitality	11		
2. Choi matrix formulation of locally balanced channels	12		

I. INTRODUCTION

Quantum information science is famously known for having features that are not present in classical information science. Among these, perhaps the feature that distinguishes it the most from the classical case is that our knowledge of the state of two particles can be greater than our knowledge of the individual states of the particles. Indeed, for an Einstein–Podolsky–Rosen (EPR) state $|\Phi^{(2)}\rangle := (|00\rangle + |11\rangle)/\sqrt{2}$ [1], said to be entangled, our knowledge of the joint state of the two particles is complete. However, the state of an individual particle is described by an unbiased probabilistic ensemble (i.e., $|0\rangle$ or $|1\rangle$ with equal probability), and we thus have incomplete knowledge of the state of an individual particle [2].

Given this peculiar situation, one of the earliest puzzles in quantum information science was to devise a quantum generalization of the concept of conditional entropy. In the classical case, conditional entropy is meant to capture the uncertainty about the state of one particle given access to a second particle that is potentially statistically dependent on the first [3]. If there is no statistical dependence whatsoever, then the conditional entropy reduces to the usual entropy. However, if there is dependence, then there is a reduction in the uncertainty of the first particle when given access to the second. When generalizing the definition of conditional entropy in a straightforward way to the quantum case (by means of this reduction), one finds that quantum conditional entropy is negative when evaluated for the EPR state mentioned above [4]. Since conditional entropy is never negative in classical information science, this situation represents a radical departure from the classical case and has been popularly described as “knowing less than nothing” [5].

In quantum information theory [6–10], the traditional approach to resolving such puzzles is to devise a physically meaningful task for which an information quantity is an optimal rate for accomplishing the task. For the case of quantum conditional entropy, this approach was successfully applied and led to the introduction of the quantum state merging protocol [11, 12]. In this protocol, the goal is for one party A to use entanglement and classical communication to transfer her share A of a joint state to another party B . If the conditional entropy $H(A|B)$ is non-negative, then the protocol consumes entanglement to accomplish the task, at a rate given by $H(A|B)$ EPR states per copy of the original state. Whereas, if the conditional entropy $H(A|B)$ is negative, then the protocol generates entanglement at a rate of $H(A|B)$ EPR states per copy of the original state, while transferring A to B . This protocol inspired further developments regarding conditional entropy in the context of the uncertainty principle [13] and thermodynamics [14], in which its negativity plays an essential role.

Although these works brought great insight to the peculiar situation mentioned above, they mainly focus on a particular definition of the conditional entropy, based on the formula from [4]. It is also well known that there are an infinite number of ways to define quantum conditional entropy, with particular examples being the conditional min-entropy [15], conditional max-entropy [16], conditional Rényi entropies [17–19], and those based more broadly on generalized divergences (see, e.g., [10]). Several of these alternate definitions have physical meanings as well [16, 20]. However, all of these definitions are based on particular mathematical formulas for quantum conditional entropy that hitherto have not been derived on the basis of simple physical principles. As such, the aforementioned contributions do not address the question of why any plausible quantum conditional entropy can be negative when defining it in the most general way possible, which is consistent with some simple physical principles.

A. Summary of results

In this paper, it is our aim to address this pressing foundational question, and we do so by introducing a physically motivated framework for defining quantum conditional entropy, based on two simple postulates. The two postulates are related to the second law of thermodynamics and the extensivity of entropy, widely accepted in scenarios of physical interest. This novel framework is essentially the simplest way of defining any plausible quantum conditional entropy, and one of our main conclusions is that any such quantum conditional entropy must take on a negative value when evaluated for the EPR state. Following the popular terminology introduced in [5], we can summarize this finding as the “inevitability of knowing less than nothing” in the quantum world. As such, our approach answers the aforementioned question, i.e., why the quantum conditional entropy can be negative. We additionally note here that this finding is in stark contrast to the conclusion, also presented here, that any plausible unconditional quantum entropy takes on a non-negative value for all quantum states (here quantum entropy is defined by a similar physically motivated framework that generalizes the classical case in [21]).

As additional findings, we show that all plausible quantum conditional entropies cannot be smaller than the quantum conditional min-entropy, thus justifying once and for all the name of the latter quantity as the smallest plausible quantum conditional entropy. We also establish a logical equivalence between the non-negativity of all quantum conditional entropies and the well known reduction criterion [22] from entanglement theory. As a corollary, it follows that all separable states have non-negative quantum conditional entropy.

In what follows, we present details of our claims. We first generalize the classical entropy framework from [21] in order to define quantum entropy, and then we introduce our framework for defining quantum conditional entropy. After that, we indicate several properties of quantum conditional entropy that follow from the two postulates, before giving arguments for our main results. All arguments for our main results are based on constructions of physical processes that respect the first aforementioned postulate for quantum conditional entropy. The only background needed to understand the rest of our paper is basic knowledge of density operators and quantum channels (physical processes), available in various textbooks on quantum information [6–10].

B. Notation

Throughout our paper, we employ standard notation used in quantum information [6–10]. We use symbols like ρ , σ , ω to refer to quantum states (density operators) and $\mathfrak{D}(A)$ to denote the set of density operators acting on a Hilbert space A . We also identify a quantum system A with its corresponding Hilbert space A . Let $\mathbf{u}_A := I_A/|A|$

denote the uniform state of a system A , where I_A is the identity operator acting on A and $|A|$ denotes the dimension of A . We use system labels to denote bipartite states as $\rho_{AB}, \sigma_{AB}, \omega_{AB} \in \mathfrak{D}(AB)$, and $\rho_A = \text{Tr}_B[\rho_{AB}]$ denotes the marginal state after performing a partial trace over system B . A quantum channel $\mathcal{N}_{A \rightarrow B}$ is a completely positive and trace-preserving map that takes an input system A to an output system B . If the input and output systems are the same, then we simply write \mathcal{N}_A . A particular kind of quantum channel is an isometric channel, typically denoted by $\mathcal{U}_{A \rightarrow B}$ or $\mathcal{V}_{A \rightarrow B}$, and is realized as $\mathcal{U}_{A \rightarrow B}(\cdot) = U_{A \rightarrow B}(\cdot)(U_{A \rightarrow B})^\dagger$, where $U_{A \rightarrow B}$ an isometry (i.e., it satisfies $(U_{A \rightarrow B})^\dagger U_{A \rightarrow B} = I_A$).

II. UNCERTAINTY AND ENTROPY

A. Uncertainty measures

Let us first define what it means for a function to be an uncertainty measure, following the approach of [21] for the classical case. What we expect for any measure of uncertainty is that it does not decrease under the action of a mixing operation. Thus, we demand that an uncertainty measure is a function that is monotone with respect to a set of mixing operations, similar to how one variant of the second law of thermodynamics states that the entropy of an isolated system left to spontaneous evolution does not decrease. This approach is also consistent with that taken in quantum resource theories when defining resource monotones [23], in turn inspired by the second law of thermodynamics.

In the classical case of finite and discrete probability distributions, such a mixing operation is described by a random relabeling of values. For example, for a six-sided die described by a probability distribution, one can randomly relabel or permute the numbers on the faces of the die to realize a mixing operation, and the outcome of the die toss becomes more difficult to guess (and intuitively, more uncertain). Furthermore, the odds of winning a game of chance are lower after performing a mixing operation [24], providing a precise way of measuring uncertainty.

Under such mixing operations, the uniform distribution remains invariant. Furthermore, our intuition is that the uniform distribution should take the largest value of an uncertainty measure for all probability distributions of the same size, given that it is an unbiased distribution with equally likely outcomes and thus the unique distribution for which it is the most difficult to guess which outcome will occur (more generally, for which one is the least likely to win a game of chance [24]). Given everything stated above, one thus defines a function to be an uncertainty measure in the classical case if it is not equal to the zero function (i.e., a function that trivially takes all inputs to the value zero) and if it does not decrease under the action of a mixing operation [21].

In order to generalize this notion to the quantum case,

the key previously stated property that we use to define a mixing operation is the preservation of the uniform state \mathbf{u} . In this way, it is guaranteed that a mixing operation does not increase the value of an uncertainty measure for the uniform state; rather, it remains constant. Indeed, such a state is already intuitively maximally uncertain, as argued above, and so it cannot be any more uncertain, among all states of a fixed dimension. At the very least, the mixing operations we consider for defining an uncertainty measure should be quantum channels because all physical processes are described by quantum channels. The additional property of preserving the uniform state thus constrains the mixing operations to be unital channels (i.e., channels that preserve the identity operator). Formally, recall that a channel \mathcal{N}_A is unital if $\mathcal{N}_A(I_A) = I_A$, or equivalently, if $\mathcal{N}_A(\mathbf{u}_A) = \mathbf{u}_A$ [6–10]. We thus define a function of quantum states to be an uncertainty measure if it is not equal to the zero function and if it does not decrease under the action of a unital channel \mathcal{N}_A , i.e.,

$$f(\mathcal{N}_A(\rho_A)) \geq f(\rho_A) \quad (1)$$

for every state ρ_A and unital channel \mathcal{N}_A .

A state ρ_A majorizes another state σ_A , denoted by $\rho_A \succ \sigma_A$, if the following inequalities hold for all $k \in \{1, \dots, |A|\}$:

$$\sum_{i=1}^k \lambda_i^\downarrow(\rho_A) \geq \sum_{i=1}^k \lambda_i^\downarrow(\sigma_A), \quad (2)$$

where $\lambda_i^\downarrow(\omega_A)$ denotes the components of the vector of eigenvalues of a density operator ω_A sorted in decreasing order. It is well known that ρ_A majorizes σ_A if and only if there exists a unital channel \mathcal{N}_A such that $\sigma_A = \mathcal{N}_A(\rho_A)$ [25–27] (see also [8, Theorem 4.32]). Thus, our definition of an uncertainty measure f implies that f reverses the majorization pre-order, i.e.,

$$\rho_A \succ \sigma_A \Rightarrow f(\rho_A) \leq f(\sigma_A). \quad (3)$$

It is ideal and natural to generalize the definition of an uncertainty measure just slightly in order to compare the uncertainties of states of different dimensions. Suppose that we have a state ρ_A with an uncertainty value given by $f(\rho_A)$ for some uncertainty measure f . Now consider the density matrix $\rho_{A'} := \rho_A \oplus \mathbf{0}$, where $\mathbf{0}$ denotes the zero square matrix, so that $\rho_{A'}$ represents an embedding of ρ_A into a larger system A' . There is no reason that the uncertainty measure should change at all under this embedding, because, intuitively, the extra matrix entries corresponding to $\mathbf{0}$ neither increase nor decrease uncertainty. Indeed, when flipping a die on a table, one could add an extra degree of freedom corresponding to whether the table is upside down or right side up, and under normal circumstances, the table will never be upside down, so that the uncertainty of the die toss will be unaffected by this embedding. As such, we impose the additional constraint that the uncertainty measure f satisfies the

equality $f(\rho_A) = f(\rho_{A'})$ for every state ρ_A and every embedding into a larger system A' .

In this spirit, we now generalize the standard definition of majorization for quantum states from [25–27] (see also [8]). Given two states ρ_A and $\sigma_{A'}$, we say that ρ_A majorizes $\sigma_{A'}$, and write

$$\rho_A \succ \sigma_{A'} \quad (4)$$

if either $|A'| \geq |A|$, there exists an isometric channel $\mathcal{V}_{A \rightarrow A'}$, and a unital channel \mathcal{N}_A , such that

$$\sigma_{A'} = (\mathcal{V}_{A \rightarrow A'} \circ \mathcal{N}_A)(\rho_A), \quad (5)$$

or $|A| \geq |A'|$, there exists an isometric channel $\mathcal{U}_{A' \rightarrow A}$, and a unital channel \mathcal{N}_A , such that

$$\mathcal{U}_{A' \rightarrow A}(\sigma_{A'}) = \mathcal{N}_A(\rho_A). \quad (6)$$

The isometric channels introduced in the definition of majorization above enable us to compare states of different dimensions, as desired. Also, note that majorization between two probability vectors of the same dimension can be extended to vectors of different dimensions in this way (it is a standard approach to add zero components to the vector with the smaller dimension so that the resulting vectors have equal dimension). As discussed in the previous paragraph, such an extension is motivated by the fact that uncertainty measures should not change under embeddings.

In conclusion, our final definition of an uncertainty measure is a function f that is not equal to the zero function and reverses the majorization pre-order from (4); i.e., for all states ρ_A and $\sigma_{A'}$,

$$\rho_A \succ \sigma_{A'} \Rightarrow f(\rho_A) \leq f(\sigma_{A'}). \quad (7)$$

B. Entropy

Following the approach of [21] for the classical case, an uncertainty measure is not necessarily an entropy. To be an entropy, we also require that the uncertainty measure possess an extensivity or additivity property. That is, f is an entropy if it is an uncertainty measure and it is additive on tensor-product states; i.e., for all states ρ_A and $\sigma_{A'}$,

$$f(\rho_A \otimes \sigma_{A'}) = f(\rho_A) + f(\sigma_{A'}). \quad (8)$$

The additivity postulate of entropy is motivated by the basic expectation that the disorder of isolated, non-interacting systems should be the sum of the individual disorders. Indeed, if two physical systems are separate from each other, not having interacted in any way whatsoever previously, we expect that the disorder characterizing the overall system should simply be the sum of the individual disorders (i.e., there would not be a meaningful alternative way of quantifying the overall disorder in this scenario).

The motivation for additivity of entropy is also related to the second law of thermodynamics, in particular, the Clausius and Kelvin–Planck statements. In those statements, cyclic processes are considered, in which a system of interest undergoes a thermodynamic transition, while all other systems, including the environment, heat baths, etc., all return to their original state. In recent developments, using a quantum information-theoretic approach to small-scale thermodynamics [28], these cyclic processes were identified as catalytic processes. That is, consider a thermodynamical evolution of a physical system A , in a state ρ_A , to a physical system A' in a state $\sigma_{A'}$. The thermal machine that corresponds to this transition includes all the other systems (e.g., heat baths, environment, etc.) that can be represented with an additional system M in a state τ_M . Therefore, for cyclic processes, the thermodynamical transition takes the form

$$\rho_A \otimes \tau_M \rightarrow \sigma_{A'} \otimes \tau_M. \quad (9)$$

In this context, the second law not only states that the entropy of system A is no greater than that of system A' , but also that this property holds if and only if the entropy of the joint state $\rho_A \otimes \tau_M$ increases (or remains unchanged) in such a thermodynamic cyclic process that returns τ_M intact. Clearly, if the entropy is measured with an additive function (under tensor products) then the entropy of ρ_A is never greater than that of $\sigma_{A'}$ if and only if the same relation holds between $\rho_A \otimes \tau_M$ and $\sigma_{A'} \otimes \tau_M$. Thus, the second law of thermodynamics also motivates the additivity postulate for entropy.

With all of this reasoning in place, we now provide a formal definition of entropy: a function

$$\mathbf{H} : \bigcup_A \mathfrak{D}(A) \rightarrow \mathbb{R} \quad (10)$$

is a quantum entropy if it is not equal to the zero function and satisfies the following two postulates for all systems A and A' and states ρ_A and $\sigma_{A'}$:

1. Monotonicity:

$$\rho_A \succ \sigma_{A'} \Rightarrow \mathbf{H}(\rho_A) \leq \mathbf{H}(\sigma_{A'}). \quad (11)$$

2. Additivity:

$$\mathbf{H}(\rho_A \otimes \sigma_{A'}) = \mathbf{H}(\rho_A) + \mathbf{H}(\sigma_{A'}). \quad (12)$$

As stated previously, the requirement that an entropy not be equal to the zero function is to avoid a trivial situation in which the entropy of every state takes on the value zero.

By employing the two basic postulates of entropy and the same reasoning given in [21, Lemma 3], some important properties follow (see Appendix A for proofs). The entropy of every state ρ is non-negative, i.e., $\mathbf{H}(\rho) \geq 0$, and it is equal to zero for every pure state $|\psi\rangle\langle\psi|$, i.e., $\mathbf{H}(|\psi\rangle\langle\psi|) = 0$. For every state σ of a fixed dimension d , the entropy does not exceed the entropy of the

uniform state of dimension d , i.e., $\mathbf{H}(\mathbf{u}^{(d)}) \geq \mathbf{H}(\sigma)$. Finally, the entropy of the uniform state of dimension two is strictly positive, i.e., $\mathbf{H}(\mathbf{u}^{(2)}) > 0$. This allows for setting the normalization of entropy to be in units of bits, such that $\mathbf{H}(\mathbf{u}^{(2)}) = 1$, and henceforth, we thus set the normalization of entropy in this way. It then follows that $\mathbf{H}(\mathbf{u}^{(d)}) = \log_2 d$.

III. CONDITIONAL UNCERTAINTY AND ENTROPY

A. Conditional uncertainty measures

Our goal now is to define what it means for a function of a bipartite state ρ_{AB} to be a measure of conditional uncertainty. Inspired by the criterion of monotonicity under mixing operations in the previous case of unconditional uncertainty, we are interested in identifying a criterion for a bipartite channel $\mathcal{N}_{AB \rightarrow A'B'}$ to be a conditional mixing operation. For the unconditional case, we ultimately stated that a quantum channel is a mixing operation if it preserves the uniform state, and this choice led to identifying unital channels as the mixing operations that define an uncertainty measure. The reason for this, as previously stated, is that the uniform state is the least informative state, it thus gives the least likelihood of winning a game of chance, and should thus be a state of maximal uncertainty (for which a mixing operation cannot increase uncertainty).

For the setting of conditional uncertainty, it remains the case that the uniform state $\mathbf{u}_{AB} = \mathbf{u}_A \otimes \mathbf{u}_B$ is the least informative state, for reasons similar to those given before. However, for conditional uncertainty, we are interested in the uncertainty of system A in the presence of system B . That is, the system B can be used as an aid or side information to help win a game of chance performed on the system A . With this in mind, a state of the form $\mathbf{u}_A \otimes \rho_B$ is still least informative, with ρ_B an arbitrary state, because the system B is independent of system A and thus cannot help at all in winning a game of chance performed on A . At the same time, the state of system A is still uniform and thus the least informative. As such, we identify the set of states having the form $\mathbf{u}_A \otimes \rho_B$, where ρ_B is an arbitrary state of system B , as being the set of states with maximum conditional uncertainty.

Thus, we have argued that one requirement for a function to be a measure of conditional uncertainty is that it should not decrease under the action of a bipartite channel that preserves the set of states with maximum conditional uncertainty. We call such channels conditionally unital channels because they generalize unital channels. Formally, a bipartite channel $\mathcal{N}_{AB \rightarrow AB'}$ is a conditionally unital channel if for every $\rho_B \in \mathfrak{D}(B)$, there exists a state $\sigma_{B'} \in \mathfrak{D}(B')$ such that

$$\mathcal{N}_{AB \rightarrow AB'}(\mathbf{u}_A \otimes \rho_B) = \mathbf{u}_A \otimes \sigma_{B'}. \quad (13)$$

See Appendix B 1 for an equivalent formulation of this




FIG. 1. Depiction of the definition of a semi-causal channel \mathcal{N} , where the black square represents the discarding channel.

condition in terms of the Choi matrix of $\mathcal{N}_{AB \rightarrow AB'}$. We note here that conditionally unital channels were independently defined in our companion paper [29] and in [30]. See also [15, Lemma 3.1.12] for an implicit definition of sub-conditionally-unital channels.

In the setting of conditional uncertainty, it is important to consider a further constraint on the class of transformations allowed. Intuitively, conditional uncertainty measures the uncertainty of system A when one is given access to system B . Evidently, by allowing transformations in which information can leak from system A to system B , Bob could gain information that could decrease the uncertainty associated with system A . Thus, it is natural to assume that channels that do not decrease conditional uncertainty do not allow for communication or signaling from Alice to Bob. This requirement is also motivated from a cryptographic perspective, in which conditional uncertainty measures the amount of uncertainty that an eavesdropper has about another system that is intended to be secure [31] (with respect to our system labeling, the eavesdropper would have system B and the secure system would be A). With this motivation in mind, let us recall the definition of $A \not\rightarrow B$ semi-causal channels [32]. A bipartite channel $\mathcal{N}_{AB \rightarrow AB'}$ is $A \not\rightarrow B'$ semi-causal if for every channel \mathcal{M}_A

$$\mathcal{N}_{AB \rightarrow B'} \circ \mathcal{M}_A = \mathcal{N}_{AB \rightarrow B'}, \quad (14)$$

where

$$\mathcal{N}_{AB \rightarrow B'} := \text{Tr}_A \circ \mathcal{N}_{AB \rightarrow AB'}. \quad (15)$$

See Figure 1 for a depiction of the semi-causality requirement.

The authors of [33] proved that every semi-causal channel can be written as a local channel on B that feeds an output (corresponding to a quantum system R) into a channel acting only on A and R . See also [34] and Figure 2 for a visual depiction. As such, for these channels, information can only flow from B to A , and it never leaks from A to B . Formally, if $\mathcal{N}_{AB \rightarrow AB'}$ is an $A \not\rightarrow B'$ semi-causal channel, then there exists a quantum system R , a quantum channel $\mathcal{E}_{AR \rightarrow A}$, and an isometric channel $\mathcal{F}_{B \rightarrow RB'}$ such that

$$\mathcal{N}_{AB \rightarrow AB'} = \mathcal{E}_{RA \rightarrow A} \circ \mathcal{F}_{B \rightarrow RB'}. \quad (16)$$

We use the term locally balanced to describe a channel $\mathcal{N}_{AB \rightarrow AB'}$ that is both conditionally unital and $A \not\rightarrow B'$




FIG. 2. Eq. (16) asserts that a semi-causal bipartite channel $\mathcal{N}_{AB \rightarrow AB'}$ can be implemented in the depicted fashion.

semi-causal. We use this term because the properties of such a channel are invariant under the exchange of the A input and A output systems. This is most readily observed using the formalism of Choi matrices and we discuss it further in Appendix B2. Importantly, we use locally balanced channels to define measures of conditional uncertainty.

Generalizing how unital channels define the majorization pre-order on quantum states, we now define a conditional majorization pre-order on bipartite quantum states, based on locally balanced channels. Given two bipartite states ρ_{AB} and $\sigma_{A'B'}$, we say that ρ_{AB} conditionally majorizes $\sigma_{A'B'}$ with respect to system A , and write

$$\rho_{AB} \succ_A \sigma_{A'B'} \quad (17)$$

if either $|A'| \geq |A|$, there exists an isometric channel $\mathcal{V}_{A \rightarrow A'}$, and a locally balanced channel $\mathcal{N}_{AB \rightarrow AB'}$, such that

$$\sigma_{A'B'} = \mathcal{V}_{A \rightarrow A'} \circ \mathcal{N}_{AB \rightarrow AB'}(\rho_{AB}), \quad (18)$$

or $|A| \geq |A'|$, there exists an isometric channel $\mathcal{U}_{A' \rightarrow A}$, and a locally balanced channel $\mathcal{N}_{AB \rightarrow AB'}$, such that

$$\mathcal{U}_{A' \rightarrow A}(\sigma_{A'B'}) = \mathcal{N}_{AB \rightarrow AB'}(\rho_{AB}). \quad (19)$$

The isometric channels introduced in the definition above enable us to compare bipartite states of different dimensions, as was the case before with the generalized definition of majorization in (4)–(6). We note here that conditional majorization, as defined here, reduces to classical conditional majorization, as defined in [35], when the states involved are classical (see Appendix C for a proof).

We define a function f to be a conditional uncertainty measure if it is not equal to the zero function and if it reverses the conditional majorization pre-order, i.e.,

$$\rho_{AB} \succ_A \sigma_{A'B'} \Rightarrow f(\rho_{AB}) \leq f(\sigma_{A'B'}). \quad (20)$$

B. Conditional entropy

Having defined measures of conditional uncertainty, we now follow the approach from before and define a function to be a conditional entropy if it is not only a condi-

tional uncertainty measure but also additive on tensor-product states. Our motivation for the additivity postulate is the same as before. Formally, a function

$$\mathbf{H} : \bigcup_{A,B} \mathfrak{D}(AB) \rightarrow \mathbb{R} \quad (21)$$

is a quantum conditional entropy if it is not equal to the zero function and satisfies the following postulates for all systems A, B, A', B' and states ρ_{AB} and $\sigma_{A'B'}$:

1. Monotonicity:

$$\rho_{AB} \succ_A \sigma_{A'B'} \Rightarrow \mathbf{H}(A|B)_\rho \leq \mathbf{H}(A'|B')_\sigma. \quad (22)$$

2. Additivity:

$$\mathbf{H}(AA'|BB')_{\rho \otimes \sigma} = \mathbf{H}(A|B)_\rho + \mathbf{H}(A'|B')_\sigma. \quad (23)$$

As indicated in the definition, a conditional entropy \mathbf{H} is defined for all density matrices in all finite dimensions. As a very special case, if $\rho \in \mathfrak{D}(AB)$ and B is a trivial system for which $|B| = 1$, then we write

$$\mathbf{H}(A|B)_\rho = \mathbf{H}(A)_\rho, \quad (24)$$

where $\mathbf{H}(A)_\rho$ is an entropy, as defined before.

Here we state some basic properties of quantum conditional entropy before presenting our main result in the next section (see Appendix D for proofs). First, conditional entropy reduces to entropy when evaluated on a tensor-product state. That is, for every product state $\omega_A \otimes \tau_B$, the following equality holds:

$$\mathbf{H}(A|B)_{\omega \otimes \tau} = \mathbf{H}(A)_\omega. \quad (25)$$

Next, conditional entropy is invariant under the action of local isometric channels. That is, for a bipartite state ρ_{AB} and isometric channels $\mathcal{U}_{A \rightarrow A'}$ and $\mathcal{V}_{B \rightarrow B'}$,

$$\mathbf{H}(A|B)_\rho = \mathbf{H}(A'|B')_\omega, \quad (26)$$

where $\omega_{AB'} := (\mathcal{U}_{A \rightarrow A'} \otimes \mathcal{V}_{B \rightarrow B'}) (\rho_{AB})$. Finally, consider the product state $\mathbf{u}_A^{(2)} \otimes \rho_B$. Similar to what we found previously for unconditional entropy, the following inequality holds

$$\mathbf{H}(A|B)_{\mathbf{u}^{(2)} \otimes \rho} = \mathbf{H}(A)_{\mathbf{u}^{(2)}} > 0. \quad (27)$$

This strict inequality allows us to set a normalization factor for conditional entropy. To be consistent with the normalization convention for unconditional entropy, we set $\mathbf{H}(A|B)_{\mathbf{u}^{(2)} \otimes \rho} = 1$, which in turn implies that

$$\mathbf{H}(A|B)_{\mathbf{u}^{(d)} \otimes \rho} = \log_2 d, \quad (28)$$

where the state being evaluated is $\mathbf{u}_A^{(d)} \otimes \rho_B$.

IV. MAIN RESULT: INEVITABILITY OF NEGATIVE QUANTUM CONDITIONAL ENTROPY

We now have almost everything set up to state our main result. Before doing so, let us define the conditional min-entropy [15] of a bipartite state ρ_{AB} as

$$H_{\min}(A|B)_\rho := - \inf_{\lambda \geq 0} \log_2 \{ \lambda : \rho_{AB} \leq \lambda I_A \otimes \rho_B \}, \quad (29)$$

The conditional min-entropy is indeed a conditional entropy according to (22)–(23) (see Appendix F 1 for a proof). This quantity was previously given the name conditional min-entropy because it is known to be the least among all Rényi conditional entropies [15, 17, 18]. As part of our main result, we strengthen this observation by proving that all plausible quantum conditional entropies are not smaller than the conditional min-entropy; that is, the conditional min-entropy is the least plausible.

The following theorem (proved in Appendix E 2) states that every plausible quantum conditional entropy is bounded from below by the conditional min-entropy and also that this lower bound is saturated for the maximally entangled state. As a consequence, every plausible conditional entropy takes on a negative value for the maximally entangled state.

Theorem 1 *Let \mathbf{H} be a quantum conditional entropy, as defined in (21)–(23). Then, for every state ρ_{AB} ,*

$$\mathbf{H}(A|B)_\rho \geq H_{\min}(A|B)_\rho. \quad (30)$$

Equality is attained in (30) if ρ_{AB} is equal to a maximally entangled state $\Phi_{A'B'}^{(k)}$ by the action of local isometric channels. Thus,

$$\mathbf{H}(A|B)_{\Phi^{(k)}} = -\log_2 k \quad (31)$$

for every conditional entropy \mathbf{H} .

The following statement is a direct consequence of Theorem 1, indicating that every plausible conditional entropy being non-negative is equivalent to the reduction criterion [22], well known in entanglement theory. Thus, Corollary 1 provides a direct link between every conditional entropy (including the conditional min-entropy) and the reduction criterion.

Corollary 1 *Let ρ_{AB} be a bipartite state. Then, the following are equivalent:*

1. *For every conditional entropy \mathbf{H} ,*

$$\mathbf{H}(A|B)_\rho \geq 0. \quad (32)$$

2. *The reduction criterion holds: $I_A \otimes \rho_B \geq \rho_{AB}$.*

The proof of Corollary 1 is immediate from Theorem 1. Indeed, if the first condition holds, then the inequality $H_{\min}(A|B)_\rho \geq 0$ holds in particular (since $H_{\min}(A|B)_\rho$

is a conditional entropy), and the second condition is then a consequence of the definition in (29). Now suppose that the second condition holds. This then means that there exists $\lambda \leq 1$ such that $\rho_{AB} \leq \lambda I_A \otimes \rho_B$. According to the definition in (29), it follows that $H_{\min}(A|B)_\rho \geq 0$ and then Theorem 1 implies the first condition.

Suppose that a state ρ_{AB} is separable [36], meaning that it can be written as a probabilistic mixture of product states (i.e., $\rho_{AB} = \sum_x p(x) \sigma_A^x \otimes \tau_B^x$ for some probability distribution $\{p(x)\}_x$ and sets $\{\sigma_A^x\}_x$ and $\{\tau_B^x\}_x$ of states). Then it follows that the second condition in Corollary 1 holds [22], so that every conditional entropy is non-negative for all separable states.

Although all plausible conditional entropies are non-negative on all separable states, this does not mean that the conditional entropies are only non-negative on separable states. In fact, some entangled states have non-negative conditional entropy on *all* choices of conditional entropy functions. In light of this, Corollary 1 is helpful because it provides a simple sufficient criterion for a bipartite state to have non-negative quantum conditional entropy.

In Appendix E, we discuss a subclass of conditional entropies and report results that are similar to those given in Theorem 1 and Corollary 1. The difference is that the lower bound in (30) is expressed in terms of a different definition of conditional min-entropy and the equivalent condition in Corollary 1 becomes a variation of the reduction criterion, rather than the reduction criterion itself.

V. CONDITIONAL ENTROPY FROM RELATIVE ENTROPY

In this section, we prove that one can construct a quantum conditional entropy from a quantum relative entropy, based on two simple postulates that define a quantum relative entropy [21, 37]. Before giving this construction, let us first recall the definition of a quantum relative entropy [21, 37]. Formally, a function

$$\mathbf{D} : \bigcup_A \{ \mathfrak{D}(A) \times \mathfrak{D}(A) \} \rightarrow \mathbb{R}_+ \cup \{ \infty \} \quad (33)$$

is a quantum relative entropy if it is not equal to the zero function and satisfies the following postulates for all systems A and B , all states $\rho_A, \sigma_A, \omega_B, \tau_B$, and every channel $\mathcal{N}_{A \rightarrow B}$:

1. *Monotonicity:*

$$\mathbf{D}(\rho_A \| \sigma_A) \geq \mathbf{D}(\mathcal{N}_{A \rightarrow B}(\rho_A) \| \mathcal{N}_{A \rightarrow B}(\sigma_A)). \quad (34)$$

2. *Additivity:*

$$\mathbf{D}(\rho_A \otimes \omega_B \| \sigma_A \otimes \tau_B) = \mathbf{D}(\rho_A \| \sigma_A) + \mathbf{D}(\omega_B \| \tau_B). \quad (35)$$

Now we state our final result (see Appendix F for a proof):

Theorem 2 *Let \mathbf{D} be a quantum relative entropy. Then the function*

$$\log_2 |A| - \mathbf{D}(\rho_{AB} \| \mathbf{u}_A \otimes \rho_B) \quad (36)$$

is a quantum conditional entropy.

Suitable relative entropies include the Umegaki (standard) [38], Belavkin–Staszewski [39], Petz–Rényi [40, 41], sandwiched Rényi [18, 42], α -z [43], and geometric Rényi [44, 45] relative entropies, for which we point to [10] for a summary. The construction in (36), for generating a conditional entropy from a relative entropy, has been known for some time, but the difference with our result here is that we prove, starting from the simple postulates for relative entropy in (33)–(35), that the function in (36) satisfies the two simple postulates for conditional entropy in (21)–(23).

VI. CONCLUSION

In summary, we have established a physically motivated, minimal framework for defining conditional entropies, based on two simple postulates (non-decrease under locally balanced channels and additivity on tensor-product states). We then explored the consequences of these postulates, which include reduction to entropy for tensor-product states, invariance under local isometric channels, a lower bound in terms of the conditional min-entropy, and taking on a negative value for the maximally entangled state. We consider the latter property to be the most profound result of this work, and all of our arguments rely on constructing physical processes that respect the first postulate of non-decrease under locally balanced channels.

One may wonder if it might be possible to replace the additivity postulate of conditional entropy with a weaker criterion. For example, suppose we replace the additivity postulate (23) with a weaker postulate of the form

$$\mathbf{H}(A^n | B^n)_{\rho^{\otimes n}} = n\mathbf{H}(A | B)_\rho \quad \forall n \in \mathbb{N}. \quad (37)$$

In Appendix G we show that there exist examples of functions that satisfy both the monotonicity postulate and

the above weak additivity, but take a non-negative value on *all* states. Therefore, the negativity of conditional-entropies (as defined in this paper) on the maximally entangled state is closely related to the (strong) additivity postulate (23).

Every relative entropy \mathbf{D} , can be used as in (36) to define a conditional entropy. One can also attempt to define another type of conditional entropy of the form

$$\log_2 |A| - \min_{\sigma \in \mathfrak{D}(B)} \mathbf{D}(\rho_{AB} \| \mathbf{u}_A \otimes \sigma_B). \quad (38)$$

While the function above behaves monotonically under conditional unital channels, and therefore satisfies the monotonicity postulate (22) it is not clear if this function is additive under tensor products. It is known to be additive if \mathbf{D} is taken to be the Petz Rényi relative entropy [46, Lemma 3] or the sandwiched Rényi relative entropy [47], but it is not known if it is additive in general.

From its definition, the set of all conditional entropies is convex. Hence, not every conditional entropy has to take the form of dimension minus divergence. For example, if \mathbf{D} is one of the quantum Rényi relative entropies then every convex combination of the functions (36) and (38) is also a conditional entropy. It is an interesting open problem if there exists conditional entropies that cannot be expressed in terms of divergences. Another interesting open problem is whether or not all quantum conditional entropies are upper bounded by the conditional max-entropy. We leave these open problems to future investigations.

ACKNOWLEDGMENTS

The authors thank Henry Pfister for helpful discussions. GG and IG acknowledge support from the Natural Sciences and Engineering Research Council of Canada (NSERC). SB acknowledges support from the National Science Foundation (NSF) under Grant Nos. 1908730 and 1910571. MMW acknowledges support from the NSF under Grant No. 1907615. Any opinions, findings, conclusions, and recommendations expressed in this material are those of the authors and do not necessarily reflect the views of these sponsors.

-
- [1] Albert Einstein, Boris Podolsky, and Nathan Rosen. Can quantum-mechanical description of physical reality be considered complete? *Physical Review*, 47:777–780, 1935.
 - [2] Erwin Schrödinger. Discussion of probability relations between separated systems. *Proceedings of the Cambridge Philosophical Society*, 31:555–563, 1935.
 - [3] Claude E. Shannon. A mathematical theory of communication. *The Bell System Technical Journal*, 27(3):379–423, 1948.
 - [4] Nicolas J. Cerf and Christoph Adami. Negative entropy and information in quantum mechanics. *Physical Review Letters*, 79(26):5194–5197, December 1997.
 - [5] Ian Turner. Scientist knows less than nothing. *Bristol Evening Post*, August 2005. Screenshot available at <https://twitter.com/markwilde/status/1558125303569846278>.
 - [6] Masahito Hayashi. *Quantum Information Theory: Mathematical Foundation*. Springer, second edition, 2017.
 - [7] Mark M. Wilde. *Quantum Information Theory*. Cambridge University Press, second edition, 2017.

- arXiv:1106.1445.
- [8] John Watrous. *The Theory of Quantum Information*. Cambridge University Press, 2018.
- [9] Alexander S. Holevo. *Quantum Systems, Channels, Information: A Mathematical Introduction*. Walter de Gruyter, second edition, 2019.
- [10] Sumeet Khatri and Mark M. Wilde. *Principles of Quantum Communication Theory: A Modern Approach*. November 2020. arXiv:2011.04672v1.
- [11] Michał Horodecki, Jonathan Oppenheim, and Andreas Winter. Partial quantum information. *Nature*, 436(7051):673–676, August 2005.
- [12] Michał Horodecki, Jonathan Oppenheim, and Andreas Winter. Quantum state merging and negative information. *Communications in Mathematical Physics*, 269(1):107–136, January 2007. arXiv:quant-ph/0512247.
- [13] Mario Berta, Matthias Christandl, Roger Colbeck, Joseph M. Renes, and Renato Renner. The uncertainty principle in the presence of quantum memory. *Nature Physics*, 6:659–662, 2010. arXiv:0909.0950.
- [14] Lídia del Rio, Johan Åberg, Renato Renner, Oscar Dahlsten, and Vlatko Vedral. The thermodynamic meaning of negative entropy. *Nature*, 474(7349):61–63, 2011.
- [15] Renato Renner. *Security of Quantum Key Distribution*. PhD thesis, ETH Zurich, September 2005. arXiv:quant-ph/0512258.
- [16] Robert Koening, Renato Renner, and Christian Schaffner. The operational meaning of min- and max-entropy. *IEEE Transactions on Information Theory*, 55(9):4337–4347, September 2009. arXiv:0807.1338.
- [17] Marco Tomamichel, Roger Colbeck, and Renato Renner. A fully quantum asymptotic equipartition property. *IEEE Transactions on Information Theory*, 55(12):5840–5847, December 2009. arXiv:0811.1221.
- [18] Martin Müller-Lennert, Frédéric Dupuis, Oleg Szehr, Serge Fehr, and Marco Tomamichel. On quantum Rényi entropies: a new generalization and some properties. *Journal of Mathematical Physics*, 54(12):122203, December 2013. arXiv:1306.3142.
- [19] Marco Tomamichel, Mario Berta, and Masahito Hayashi. Relating different quantum generalizations of the conditional Rényi entropy. *Journal of Mathematical Physics*, 55(8):082206, August 2014. arXiv:1311.3887.
- [20] Masahito Hayashi and Marco Tomamichel. Correlation detection and an operational interpretation of the Rényi mutual information. *Journal of Mathematical Physics*, 57(10):102201, October 2016.
- [21] Gilad Gour and Marco Tomamichel. Entropy and relative entropy from information-theoretic principles. *IEEE Transactions on Information Theory*, 67(10):6313–6327, 2021.
- [22] Michał Horodecki and Paweł Horodecki. Reduction criterion of separability and limits for a class of distillation protocols. *Physical Review A*, 59(6):4206–4216, June 1999.
- [23] Eric Chitambar and Gilad Gour. Quantum resource theories. *Reviews of Modern Physics*, 91(2):025001, April 2019.
- [24] Sarah Brandsen, Isabelle Jianing Geng, and Gilad Gour. What is entropy? a perspective from games of chance. *Physical Review E*, 105(2):024117, February 2022.
- [25] Armin Uhlmann. Sätze über dichtematrizen. *Wissenschaftliche Zeitschrift der Karl-Marx-Universität Leipzig. Mathematisch-naturwissenschaftliche Reihe*, 20(4/5):633–653, 1971.
- [26] Armin Uhlmann. Endlich-dimensionale dichtematrizen i. *Wissenschaftliche Zeitschrift der Karl-Marx-Universität Leipzig. Mathematisch-naturwissenschaftliche Reihe*, 21(4):421–452, 1972.
- [27] Armin Uhlmann. Endlich-dimensionale dichtematrizen ii. *Wissenschaftliche Zeitschrift der Karl-Marx-Universität Leipzig Mathematisch-naturwissenschaftliche Reihe*, 22(2):139–177, 1973.
- [28] Fernando G. S. L. Brandão, Michał Horodecki, Nelly Ng, Jonathan Oppenheim, and Stephanie Wehner. The second laws of quantum thermodynamics. *Proceedings of the National Academy of Sciences*, 112(11):3275–3279, March 2015. arXiv:1305.5278.
- [29] Sarah Brandsen, Isabelle J. Geng, Mark M. Wilde, and Gilad Gour. Quantum conditional entropy from information-theoretic principles. October 2021. arXiv:2110.15330.
- [30] Mahathi Vempati, Saumya Shah, Nirman Ganguly, and Indranil Chakrabarty. A-unital Operations and Quantum Conditional Entropy. *Quantum*, 6:641, February 2022.
- [31] Christopher Portmann and Renato Renner. Security in quantum cryptography. *Reviews of Modern Physics*, 94(2):025008, June 2022.
- [32] David Beckman, Daniel Gottesman, Michael A. Nielsen, and John Preskill. Causal and localizable quantum operations. *Physical Review A*, 64(5):052309, October 2001.
- [33] T. Eggeling, D. Schlingemann, and Reinhard F. Werner. Semicausal operations are semilocalizable. *Europhysics Letters*, 57(6):782–788, March 2002. arXiv:quant-ph/0104027.
- [34] Marco Piani, Michał Horodecki, Paweł Horodecki, and Ryszard Horodecki. Properties of quantum nonsignaling boxes. *Physical Review A*, 74(1):012305, July 2006.
- [35] Gilad Gour, A. Grudka, M. Horodecki, W. Klobus, J. Lodyga, and V. Narasimhachar. The conditional uncertainty principle. *Physical Review A*, 97:042130, 2018.
- [36] Reinhard F. Werner. Quantum states with Einstein-Podolsky-Rosen correlations admitting a hidden-variable model. *Physical Review A*, 40(8):4277–4281, October 1989.
- [37] Gilad Gour and Marco Tomamichel. Optimal extensions of resource measures and their applications. *Physical Review A*, 102(6):062401, December 2020.
- [38] Hisaharu Umegaki. Conditional expectations in an operator algebra IV (entropy and information). *Kodai Mathematical Seminar Reports*, 14(2):59–85, 1962.
- [39] V. P. Belavkin and P. Staszewski. C^* -algebraic generalization of relative entropy and entropy. *Annales de l’I.H.P. Physique théorique*, 37(1):51–58, 1982.
- [40] Dénes Petz. Quasi-entropies for States of a von Neumann Algebra. *Publications of the Research Institute for Mathematical Sciences*, 21(4):787–800, 1985.
- [41] Dénes Petz. Quasi-entropies for finite quantum systems. *Reports in Mathematical Physics*, 23:57–65, 1986.
- [42] Mark M. Wilde, Andreas Winter, and Dong Yang. Strong converse for the classical capacity of entanglement-breaking and Hadamard channels via a sandwiched Rényi relative entropy. *Communications in Mathematical Physics*, 331(2):593–622, October 2014. arXiv:1306.1586.
- [43] Koenraad M. R. Audenaert and Nilanjana Datta. α - z -Rényi relative entropies. *Journal of Mathematical Physics*, 56(2):022202, February 2015.
- [44] Keiji Matsumoto. A new quantum version of f -

divergence. 2013. arXiv:1311.4722.

- [45] Keiji Matsumoto. A new quantum version of f-divergence. In Masanao Ozawa, Jeremy Butterfield, Hans Halvorson, Miklós Rédei, Yuichiro Kitajima, and Francesco Buscemi, editors, *Reality and Measurement in Algebraic Quantum Theory*, volume 261, pages 229–273, Singapore, 2018. Springer Singapore. Series Title: Springer Proceedings in Mathematics & Statistics.
- [46] Naresh Sharma and Naqeeb Ahmad Warsi. Fundamental bound on the reliability of quantum information transmission. *Phys. Rev. Lett.*, 110:080501, Feb 2013.
- [47] Salman Beigi. Sandwiched Rényi divergence satisfies data processing inequality. *Journal of Mathematical Physics*, 54:122202, December 2013.
- [48] Xin Wang. *Semidefinite optimization for quantum information*. PhD thesis, University of Technology Sydney, Centre for Quantum Software and Information, Faculty of Engineering and Information Technology, July 2018. <http://hdl.handle.net/10453/127996>.
- [49] Gilad Gour. Comparison of quantum channels by superchannels. *IEEE Transactions on Information Theory*, 65(9):5880–5904, 2019.
- [50] Robert Salzmann, Nilanjana Datta, Gilad Gour, Xin Wang, and Mark M. Wilde. Symmetric distinguishability as a quantum resource. *New Journal of Physics*, 23:083016, August 2021. arXiv:2102.12512.
- [51] Robert König and Renato Renner. Sampling of min-entropy relative to quantum knowledge. *IEEE Transactions on Information Theory*, 57(7):4760–4787, July 2011.
- [52] Gilad Gour. *Resources of the Quantum World (Volume 1)*. 2023.
- [53] Nilanjana Datta. Min- and max-relative entropies and a new entanglement monotone. *IEEE Transactions on Information Theory*, 55(6):2816–2826, June 2009. arXiv:0803.2770.

Appendix A: Proofs of properties of entropy

Let us employ the postulates in (10)–(12) to derive some basic properties of every entropy. The proofs given here are similar to those given for the classical case in [21, Lemma 3]. First, let us recall an often overlooked fact, that the number 1 is a trivial quantum state of dimension one. Indeed, the underlying Hilbert space is of dimension one, so that the only possible density “operator” for this space is the number 1. We now prove that the entropy of this state is equal to zero. Since, by assumption, the entropy is not equal to the zero function, there exists a state ρ such that $\mathbf{H}(\rho) \neq 0$. Then consider that

$$\mathbf{H}(\rho) = \mathbf{H}(\rho \otimes 1) = \mathbf{H}(\rho) + \mathbf{H}(1) \quad \Rightarrow \quad \mathbf{H}(1) = 0. \quad (\text{A1})$$

The first equality follows as a trivial assertion, because $\rho = \rho \otimes 1$. The next equality is a consequence of the additivity postulate. The conclusion that $\mathbf{H}(1) = 0$ then follows because $\mathbf{H}(\rho) \neq 0$.

It follows that the entropy of every pure state $|\psi\rangle\langle\psi|$ is equal to zero, as a consequence of (A1). Indeed, the state vector $|\psi\rangle$ is an isometry from the trivial Hilbert space of dimension one to the Hilbert space that contains $|\psi\rangle$.

As such, it follows that $1 \succ |\psi\rangle\langle\psi|$ and $1 \prec |\psi\rangle\langle\psi|$ according to the definitions in (4)–(6) and (10)–(12). So we conclude that

$$\mathbf{H}(|\psi\rangle\langle\psi|) = 0 \quad (\text{A2})$$

for every pure state $|\psi\rangle\langle\psi|$.

Next we conclude that the entropy of every state is non-negative. This follows because there exists a unital channel taking a pure state $|\psi\rangle\langle\psi|$ to an arbitrary state ρ , so that $|\psi\rangle\langle\psi| \succ \rho$. Indeed, let $\rho = \sum_x p_x |\psi_x\rangle\langle\psi_x|$ be a pure state decomposition of ρ . Let U_x be a unitary that transforms $|\psi\rangle$ to $|\psi_x\rangle$. Thus, $(\cdot) \rightarrow \sum_x p_x U_x(\cdot)U_x^\dagger$ is a unital channel that transforms $|\psi\rangle\langle\psi|$ to ρ . By applying (11) and (A2), we conclude that

$$\mathbf{H}(\rho) \geq 0 \quad (\text{A3})$$

for every state ρ .

For a fixed dimension d , the uniform state $\mathbf{u}^{(d)} = I/d$ is the state of maximal entropy, which follows from (11) and because every state ρ can be taken to $\mathbf{u}^{(d)}$ by means of the unital channel $(\cdot) \rightarrow \text{Tr}[(\cdot)]\mathbf{u}^{(d)}$. Thus,

$$\mathbf{H}(\mathbf{u}^{(d)}) \geq \mathbf{H}(\rho) \quad (\text{A4})$$

for every state ρ of dimension d .

Let us prove that $\mathbf{H}(\mathbf{u}^{(2)}) > 0$. By the inequality in (A3) and the assumption that entropy is not equal to the zero function, there exists a state ρ of some dimension d such that $\mathbf{H}(\rho) > 0$. By applying (A4), we conclude that $\mathbf{H}(\mathbf{u}^{(d)}) \geq \mathbf{H}(\rho)$. By taking n sufficiently large (i.e., such that $2^n > d$), we conclude that $\mathbf{u}^{(d)} \succ (\mathbf{u}^{(2)})^{\otimes n}$, which implies that $\mathbf{H}((\mathbf{u}^{(2)})^{\otimes n}) > \mathbf{H}(\mathbf{u}^{(d)})$. Finally, by applying the additivity postulate, we conclude that $n\mathbf{H}(\mathbf{u}^{(2)}) = \mathbf{H}((\mathbf{u}^{(2)})^{\otimes n})$. Combining all inequalities, we conclude that $\mathbf{H}(\mathbf{u}^{(2)}) > 0$.

After establishing the normalization convention that $\mathbf{H}(\mathbf{u}^{(2)}) = 1$, the proof that $\mathbf{H}(\mathbf{u}^{(d)}) = \log_2 d$ is the same as that given in the proof of [21, Lemma 3].

Appendix B: Choi matrix formulations

In this appendix, we provide Choi matrix formulations of conditionally unital channels and locally balanced channels. These formulations are useful conceptually and also for numerical optimization if one wishes to use semi-definite programming to optimize over these sets of channels. See [8, 10, 48] for examples and discussions of semi-definite programming in quantum information.

Before giving these formulations, let us first recall the definition of the Choi matrix. The Choi matrix J_{CD}^M of a linear map $\mathcal{M}_{C \rightarrow D}$ is defined by its action on the maximally entangled operator, as follows:

$$J_{CD}^M = \sum_{i,j=1}^{|C|} |i\rangle\langle j| \otimes \mathcal{M}(|i\rangle\langle j|). \quad (\text{B1})$$

In general, the matrix J_{CD}^M is the Choi matrix of a trace-preserving map if and only if

$$\mathrm{Tr}_D[J_{CD}^M] = I_C, \quad (\text{B2})$$

and it is the Choi matrix of a completely positive map if and only if

$$J_{CD}^M \geq 0. \quad (\text{B3})$$

Thus, J_{CD}^M is the Choi matrix of a quantum channel if and only if (B2) and (B3) both hold.

1. Choi matrix formulation of conditional unitality

Here we prove that a conditionally unital channel has an equivalent formulation in terms of its Choi matrix. We then see that a channel being conditionally unital represents a semi-definite constraint on its Choi matrix, which is convenient for optimizing over such channels using semi-definite programming.

Lemma 1 *A channel $\mathcal{N}_{AB \rightarrow \tilde{A}B'}$ is conditionally unital if and only if its Choi matrix $J_{AB\tilde{A}B'}^N$ satisfies*

$$J_{B\tilde{A}B'}^N = J_{BB'}^N \otimes \mathbf{u}_{\tilde{A}}, \quad (\text{B4})$$

where

$$J_{B\tilde{A}B'}^N := \mathrm{Tr}_A[J_{AB\tilde{A}B'}^N], \quad (\text{B5})$$

$$J_{BB'}^N := \mathrm{Tr}_{A\tilde{A}}[J_{AB\tilde{A}B'}^N], \quad (\text{B6})$$

are marginals of $J_{AB\tilde{A}B'}^N$ and we employ this kind of shorthand in what follows.

Proof. We begin by proving that the channel $\mathcal{N}_{AB \rightarrow \tilde{A}B'}$ is conditionally unital if its Choi matrix has the form in (B4). To this end, let $\rho_B \in \mathfrak{D}(B)$ and consider that

$$\begin{aligned} \mathcal{N}(\mathbf{u}_A \otimes \rho_B) &= \mathrm{Tr}_{AB} \left[J_{AB\tilde{A}B'}^N \left(\mathbf{u}_A \otimes (\rho_B)^T \otimes I_{\tilde{A}B'} \right) \right] \\ &= \frac{1}{|A|} \mathrm{Tr}_B \left[J_{B\tilde{A}B'}^N \left((\rho_B)^T \otimes I_{\tilde{A}B'} \right) \right] \end{aligned} \quad (\text{B7})$$

$$= \frac{1}{|A|} \mathrm{Tr}_B \left[J_{B\tilde{A}B'}^N \left((\rho_B)^T \otimes I_{\tilde{A}B'} \right) \right] \quad (\text{B8})$$

$$= \frac{1}{|A|} \mathrm{Tr}_B \left[\mathbf{u}_{\tilde{A}} \otimes J_{BB'}^N \left((\rho_B)^T \otimes I_{\tilde{A}B'} \right) \right] \quad (\text{B9})$$

$$= \mathbf{u}_{\tilde{A}} \otimes \frac{1}{|A|} \mathrm{Tr}_B \left[J_{BB'}^N \left((\rho_B)^T \otimes I_{B'} \right) \right]. \quad (\text{B10})$$

The first equality is a well known identity in quantum information, indicating how the action of a channel on an input state can be understood in terms of a propagation formula using its Choi matrix (see Eq. (2.66) of [8]). The third equality above follows from applying (B4), which holds by assumption. It then follows that $\frac{1}{|A|} \mathrm{Tr}_B \left[J_{BB'}^N \left((\rho_B)^T \otimes I_{B'} \right) \right]$ is a state on the output system B' (i.e., the choice for the state $\sigma_{B'}$ in (13)) because $\frac{1}{|A|} J_{BB'}^N$ is the Choi operator of the channel

$$\tau_B \rightarrow (\mathrm{Tr}_{\tilde{A}} \circ \mathcal{N}_{AB \rightarrow \tilde{A}B'}) (\mathbf{u}_A \otimes \tau_B). \quad (\text{B11})$$

We now prove that the Choi matrix of $\mathcal{N}_{AB \rightarrow \tilde{A}B'}$ has the form in (B4) if $\mathcal{N}_{AB \rightarrow \tilde{A}B'}$ is conditionally unital. Recall from (13) that the defining property of a conditionally unital channel is that for every state ρ_B , there exists a state $\sigma_{B'}$ such that

$$\mathcal{N}_{AB \rightarrow \tilde{A}B'}(\mathbf{u}_A \otimes \rho_B) = \mathbf{u}_{\tilde{A}} \otimes \sigma_{B'}. \quad (\text{B12})$$

By taking the trace over \tilde{A} on both sides of the equation above, we find that

$$\sigma_{B'} = \mathcal{N}_{AB \rightarrow B'}(\mathbf{u}_A \otimes \rho_B) \quad (\text{B13})$$

$$= \mathrm{Tr}_{AB} \left[J_{AB\tilde{A}B'}^N \left(\mathbf{u}_A \otimes (\rho_B)^T \otimes I_{B'} \right) \right] \quad (\text{B14})$$

$$= \frac{1}{|A|} \mathrm{Tr}_B \left[J_{BB'}^N \left((\rho_B)^T \otimes I_{B'} \right) \right]. \quad (\text{B15})$$

On the other hand, observe that

$$\begin{aligned} \mathcal{N}_{AB \rightarrow \tilde{A}B'}(\mathbf{u}_A \otimes \rho_B) &= \mathrm{Tr}_{AB} \left[J_{AB\tilde{A}B'}^N \left(\mathbf{u}_A \otimes (\rho_B)^T \otimes I_{\tilde{A}B'} \right) \right] \\ &= \frac{1}{|A|} \mathrm{Tr}_B \left[J_{B\tilde{A}B'}^N \left((\rho_B)^T \otimes I_{\tilde{A}B'} \right) \right]. \end{aligned} \quad (\text{B16})$$

$$= \frac{1}{|A|} \mathrm{Tr}_B \left[J_{B\tilde{A}B'}^N \left((\rho_B)^T \otimes I_{\tilde{A}B'} \right) \right]. \quad (\text{B17})$$

Therefore, from the two expressions above for $\sigma_{B'}$ and $\mathcal{N}(\mathbf{u}_A \otimes \rho_B)$ we conclude that (B12) can be expressed as

$$\begin{aligned} \mathrm{Tr}_B \left[J_{B\tilde{A}B'}^N \left((\rho_B)^T \otimes I_{\tilde{A}B'} \right) \right] &= \mathrm{Tr}_B \left[\mathbf{u}_{\tilde{A}} \otimes J_{BB'}^N \left((\rho_B)^T \otimes I_{\tilde{A}B'} \right) \right], \end{aligned} \quad (\text{B18})$$

holding for all $\rho_B \in \mathfrak{D}(B)$. Let

$$\eta_{B\tilde{A}B'} := J_{B\tilde{A}B'}^N - \mathbf{u}_{\tilde{A}} \otimes J_{BB'}^N, \quad (\text{B19})$$

and observe that (B18) can be written as

$$\mathrm{Tr}_B \left[\eta_{B\tilde{A}B'} \left((\rho_B)^T \otimes I_{\tilde{A}B'} \right) \right] = 0 \quad \forall \rho_B \in \mathfrak{D}(B). \quad (\text{B20})$$

Due to the existence of bases of density operators that span the space of linear operators acting on B (see Example 2.7 of [8]), we conclude from (B20) that for every Hermitian operator $X_B \in \mathfrak{L}(B)$, we have

$$\mathrm{Tr}_B [\eta_{B\tilde{A}B'} (X_B \otimes I_{\tilde{A}B'})] = 0. \quad (\text{B21})$$

Note that by multiplying both sides of the equation above by an arbitrary element $Y_{\tilde{A}B'} \in \mathfrak{L}(\tilde{A}B')$ and taking the trace, we get that

$$\mathrm{Tr} [\eta_{B\tilde{A}B'} (X_B \otimes Y_{\tilde{A}B'})] = 0. \quad (\text{B22})$$

Since the equation above holds for all $X_B \in \mathfrak{L}(B)$ and all $Y_{\tilde{A}B'} \in \mathfrak{L}(\tilde{A}B')$ it also holds for all linear combinations of matrices of the form $X_B \otimes Y_{\tilde{A}B'}$. Since matrices of the form $X_B \otimes Y_{\tilde{A}B'}$ span the whole space $\mathfrak{L}(B\tilde{A}B')$ we conclude that

$$\mathrm{Tr} [\eta_{B\tilde{A}B'} W_{B\tilde{A}B'}] = 0 \quad \forall W_{B\tilde{A}B'} \in \mathfrak{L}(B\tilde{A}B'). \quad (\text{B23})$$

Therefore, we conclude that $\eta_{B\tilde{A}B'} = 0$, which is equivalent to (B4). ■

2. Choi matrix formulation of locally balanced channels

Here we provide a Choi matrix formulation for locally balanced channels. Before doing so, let us first recall [34, Eq. (40)], which indicates how semi-causality can be expressed in terms of the Choi matrix.

Lemma 2 ([34, 49]) *The channel $\mathcal{N}_{AB \rightarrow AB'}$ is $A \not\rightarrow B'$ semi-causal if and only if the marginals of its Choi matrix $J_{AB\bar{A}B'}^{\mathcal{N}}$ satisfy*

$$J_{ABB'}^{\mathcal{N}} = \mathbf{u}_A \otimes J_{BB'}^{\mathcal{N}}. \quad (\text{B24})$$

As a direct corollary of (B2)–(B3) and Lemmas 1 and 2, we have the following simple semi-definite characterization of a locally balanced channel in terms of its Choi matrix.

Corollary 2 *A matrix $J_{AB\bar{A}B'}^{\mathcal{N}}$ is the Choi matrix of a locally balanced channel $\mathcal{N}_{AB \rightarrow \bar{A}B'}$ if and only if*

$$J_{AB\bar{A}B'}^{\mathcal{N}} \geq 0, \quad (\text{B25})$$

$$\text{Tr}_{\bar{A}B'}[J_{AB\bar{A}B'}^{\mathcal{N}}] = I_{AB}, \quad (\text{B26})$$

$$J_{ABB'}^{\mathcal{N}} = \mathbf{u}_A \otimes J_{BB'}^{\mathcal{N}}, \quad (\text{B27})$$

$$J_{B\bar{A}B'}^{\mathcal{N}} = J_{BB'}^{\mathcal{N}} \otimes \mathbf{u}_{\bar{A}}. \quad (\text{B28})$$

Corollary 2 simplifies the task of optimizing an objective function over the set of locally balanced channels by making use of semi-definite programming. Corollary 2 also clarifies why we have named these channels locally balanced. By inspecting (B27)–(B28), we see that the properties of these channels are invariant under the exchange of the input system A and the output system \bar{A} .

Appendix C: Quantum conditional majorization reduces to classical conditional majorization

In [35], a pre-order \succ_C between two joint probability distributions (i.e., classical states) denotes *classical conditional majorization*. In this appendix, we prove that the approach to conditional majorization defined in Section III A is a generalization of classical conditional majorization. To this end, we demonstrate that quantum conditional majorization is equivalent to the classical conditional majorization when we restrict the underlying bipartite quantum states to classical states.

Suppose the system $A := X$ is classical. We call a channel $\mathcal{M}_{XB \rightarrow XB'}$ *conditionally doubly stochastic* (CDS for short) if it has the following form:

$$\mathcal{M}_{XB \rightarrow XB'} = \sum_j \mathcal{D}_{X \rightarrow X}^{(j)} \otimes \mathcal{F}_{B \rightarrow B'}^{(j)}, \quad (\text{C1})$$

where each $\mathcal{D}^{(j)}$ is a classical doubly stochastic channel and each $\mathcal{F}_{B \rightarrow B'}^{(j)}$ is a completely positive (CP) map such

that $\sum_j \mathcal{F}_{B \rightarrow B'}^{(j)}$ is a quantum channel. CDS maps with classical B and B' systems were introduced and studied in [35], and CDS maps with quantum B and B' systems were introduced and studied in [50].

Definition 1 ([35]) *Let ρ_{XY} and $\sigma_{XY'}$ be classical bipartite states such that $A = X$, $B = Y$, and $B' = Y'$ are all classical systems. We say that ρ_{XY} conditionally majorizes $\sigma_{XY'}$ and write $\sigma_{XY'} \succ_C \rho_{XY}$ if there exists a CDS channel $\mathcal{N}_{XY \rightarrow XY'}$ such that $\sigma_{XY'} = \mathcal{N}_{XY \rightarrow XY'}(\rho_{XY})$.*

Lemma 3 *Suppose all the systems involved are classical with $A := X$, $B := Y$, and $B' := Y'$. Then every CDS channel is locally balanced.*

Proof. Recall that if $\mathcal{M}_{XY \rightarrow XY'}$ is a CDS channel, then there exists a set of classical doubly stochastic channels $\{\mathcal{D}^{(j)}\}_j$ and a quantum instrument $\{\mathcal{F}_{Y \rightarrow Y'}^{(j)}\}_j$ such that

$$\mathcal{M}_{AB \rightarrow AB'} = \sum_j \mathcal{D}_A^{(j)} \otimes \mathcal{F}_{B \rightarrow B'}^{(j)}. \quad (\text{C2})$$

Then for every input of the form $\mathbf{u}_A \otimes \sigma_B$, it follows that

$$\begin{aligned} \mathcal{M}_{AB \rightarrow AB'}(\mathbf{u}_A \otimes \sigma_B) &= \sum_j \mathcal{D}_A^{(j)}(\mathbf{u}_A) \otimes \mathcal{F}_{B \rightarrow B'}^{(j)}(\sigma_B) \\ &= \mathbf{u}_A \otimes \sum_j \mathcal{F}_{B \rightarrow B'}^{(j)}(\sigma_B) \end{aligned} \quad (\text{C3})$$

and therefore \mathcal{M} is conditionally unital. (Note that the third line of the above follows from the fact that $\mathcal{D}^{(j)}$ is doubly stochastic, so that $\mathcal{D}^{(j)}(\mathbf{u}_A) = \mathbf{u}_A$ for all j .)

Finally, we show that \mathcal{M} is semi-causal. Let \mathcal{T}_A be an arbitrary classical channel. Then consider that

$$\text{Tr}_A[\mathcal{M} \circ \mathcal{T}_A] = \text{Tr}_A \left[\sum_j (\mathcal{D}_A^{(j)} \circ \mathcal{T}_A) \otimes \mathcal{F}_{B \rightarrow B'}^{(j)} \right] \quad (\text{C4})$$

$$= \sum_j \text{Tr}_A \left[\mathcal{D}_A^{(j)} \circ \mathcal{T}_A \right] \otimes \mathcal{F}_{B \rightarrow B'}^{(j)} \quad (\text{C5})$$

$$= \sum_j \text{Tr}_A \left[\mathcal{D}_A^{(j)} \right] \otimes \mathcal{F}_{B \rightarrow B'}^{(j)} \quad (\text{C6})$$

$$= \text{Tr}_A[\mathcal{M}_{AB \rightarrow AB'}]. \quad (\text{C7})$$

Given that \mathcal{M} is conditionally unital and semi-causal, then the statement of the lemma follows. ■

Theorem 3 *Let ρ_{XY} and $\sigma_{XY'}$ be classical bipartite states such that $A = X$, $B = Y$, and $B' = Y'$ are all classical systems. Then $\sigma_{XY'} \succ_X \rho_{XY}$ if and only if $\sigma_{XY'} \succ_C \rho_{XY}$, where \succ_C is the classical conditional majorization defined in [35].*

Proof. It immediately follows from Lemma 3 that $\rho_{XY} \succ_C \sigma_{XY'}$ implies $\rho_{XY} \succ_X \sigma_{XY'}$.

To show the other direction, let us suppose that $\rho_{XY} \succ_X \sigma_{XY'}$ as defined in (17)–(19). That is, there exists a locally balanced channel $\mathcal{N}_{XY \rightarrow XY'}$ such that

$$\sigma_{XY'} = \mathcal{N}_{XY \rightarrow XY'}(\rho_{XY}). \quad (\text{C8})$$

Since \mathcal{N} is semi-causal, it has the form

$$\mathcal{N}_{XY \rightarrow XY'} = \sum_{j=1}^k \mathcal{E}_{X \rightarrow X}^{(j)} \otimes \mathcal{F}_{Y \rightarrow Y'}^{(j)}, \quad (\text{C9})$$

where for each $j \in [K]$, the map $\mathcal{E}_{X \rightarrow X}^{(j)}$ is a channel and $\mathcal{F}_{Y \rightarrow Y'}^{(j)}$ is a completely positive map with $\sum_j \mathcal{F}_{Y \rightarrow Y'}^{(j)}$ a channel. Hence,

$$\sigma_{XY'} = \sum_{j=1}^k \mathcal{E}_{X \rightarrow X}^{(j)} \otimes \mathcal{F}_{Y \rightarrow Y'}^{(j)}(\rho_{XY}). \quad (\text{C10})$$

Since ρ and σ are classical, we can make the identification $\rho_{XY} \cong P$ and $\sigma_{XY'} \cong Q$, where P and Q are probability matrices. Thus, (C10) can be expressed as

$$Q = \sum_{j=1}^k E^{(j)} P R^{(j)}, \quad (\text{C11})$$

where for each $j \in [k]$, the matrix $E^{(j)}$ is the transition matrix of $\mathcal{E}^{(j)}$, and $R^{(j)}$ is the transpose of the transition matrix of $\mathcal{F}^{(j)}$. Hence, each $E^{(j)}$ is column stochastic and $R := \sum_{j=1}^k R^{(j)}$ is row stochastic. Let $\{\mathbf{p}_y\}_{y \in [n]}$ be the columns of P , and observe that in component form, the above relation can be expressed as

$$q_{xw} = \sum_{j=1}^k \sum_{y=1}^n (E^{(j)} \mathbf{p}_y)_x r_{y_0 w}^{(j)}. \quad (\text{C12})$$

So far we only used the constraint that \mathcal{N} is semi-causal. The other constraint that \mathcal{N} is conditionally unital implies that $q_{yw} = \frac{1}{m} q_w$ if $p_{xy} = \frac{1}{m} p_y$. Therefore, for simplicity we take $p_{xy} = \frac{1}{m} \delta_y y_0$ for some fixed $y_0 \in [n]$. Then the above equation becomes

$$\frac{1}{m} q_w = \sum_j (E^{(j)} \mathbf{u})_x r_{y_0 w}^{(j)}. \quad (\text{C13})$$

Summing over x on both sides of the equation, and using the fact that each $E^{(j)}$ is column stochastic, which implies that $E^{(j)} \mathbf{u}$ is a probability vector, we have

$$q_w = \sum_j r_{y_0 w}^{(j)}. \quad (\text{C14})$$

Note that we assumed that y_0 was fixed, and therefore, as a result, the equation above does not imply that q_w is independent of y_0 if we allow y_0 to vary. In other words, each value of y_0 corresponds to a different P matrix and

therefore possibly resulting in a different Q matrix. For simplicity, we rename y_0 as y and denote

$$q_w = t_{yw} := \sum_j r_{yw}^{(j)}. \quad (\text{C15})$$

By definition, the matrix $T = (t_{yw})$ is row-stochastic, and with the above notations, we can express (C13) as

$$\mathbf{u} = \sum_j \frac{r_{yw}^{(j)}}{t_{yw}} E^{(j)} \mathbf{u} \quad y \in [n], w \in [\ell]. \quad (\text{C16})$$

Therefore, the matrices

$$D_{(y,w)} := \sum_j \frac{r_{yw}^{(j)}}{t_{yw}} E^{(j)} \quad y \in [n], w \in [\ell] \quad (\text{C17})$$

are all doubly stochastic. To see this, observe that for each y and w , the set $\left\{ \frac{r_{yw}^{(j)}}{t_{yw}} \right\}_j$ forms a probability distribution over j , and the fact that each $E^{(j)}$ is column stochastic implies that $D_{(y,w)}$ is column stochastic. Moreover, the relation $\mathbf{u} = D_{(y,w)} \mathbf{u}$ implies that each $D_{(y,w)}$ is row stochastic. Finally, for general probability matrices $P = [\mathbf{p}_1, \dots, \mathbf{p}_n]$ and $Q = [\mathbf{q}_1, \dots, \mathbf{q}_{n'}]$, we conclude that (C12) is equivalent to

$$\mathbf{q}_w = \sum_{y=1}^n t_{yw} D_{(y,w)} \mathbf{p}_y \quad \forall w \in [n']. \quad (\text{C18})$$

The above relation is precisely the condition given in [35, Lemma 3]. Therefore, we conclude that $\rho_{XY} \succ_X \sigma_{XY'}$ implies classical conditional majorization. ■

Appendix D: Proofs of properties of quantum conditional entropy

Suppose that $\rho_{AB} = \omega_A \otimes \tau_B$ is a product state. Let \mathcal{E}_B be the completely randomizing channel acting on system B , which traces out system B and replaces with the uniform state \mathbf{u}_B . Then it follows that $\mathcal{I}_A \otimes \mathcal{E}_B$ is a conditionally unital channel and that

$$\mathcal{I}_A \otimes \mathcal{E}_B(\omega_A \otimes \tau_B) = \omega_A \otimes \mathbf{u}_B. \quad (\text{D1})$$

Moreover, let \mathcal{R}_B be the replacement channel on system B , which takes every quantum state to a fixed state τ_B . Then, $\mathcal{I}_A \otimes \mathcal{R}_B$ is conditionally unital and

$$\mathcal{I}_A \otimes \mathcal{R}_B(\omega_A \otimes \mathbf{u}_B) = \omega_A \otimes \tau_B. \quad (\text{D2})$$

Thus,

$$\omega_A \otimes \tau_B \succ_A \omega_A \otimes \mathbf{u}_B \quad \text{and} \quad \omega_A \otimes \mathbf{u}_B \succ_A \omega_A \otimes \tau_B, \quad (\text{D3})$$

which implies, by the monotonicity property of the conditional entropy \mathbf{H} , that

$$\mathbf{H}(A|B)_{\omega_A \otimes \tau_B} = \mathbf{H}(A|B)_{\omega_A \otimes \mathbf{u}_B}. \quad (\text{D4})$$

Thus, $\mathbf{H}(A|B)_{\omega_A \otimes \tau_B}$ depends on ω_A only. Moreover, the function $\omega_A \mapsto \mathbf{H}(A|B)_{\omega_A \otimes \tau_B}$ satisfies the two postulates of entropy in (11)–(12) and therefore is an entropy of ω_A . Note that the same statement also holds when system $|B\rangle$ is trivial, i.e., when $|B| = 1$.

We now prove that $\mathbf{H}(A|B)$ is invariant under local isometric channels. It follows immediately from the definition of quantum conditional entropy (specifically, (17)–(19) and (22)) that it is invariant under local isometric channels acting on system A . So we prove invariance under isometric channels acting on system B . The inequality

$$\mathbf{H}(A|B)_\rho \leq \mathbf{H}(A|B')_\omega \quad (\text{D5})$$

follows because $\mathcal{V}_{B \rightarrow B'}$ is a conditionally unital channel. Let τ_B be a state. The reverse inequality follows because

$$\mathcal{R}_{B' \rightarrow B}^\mathcal{V}(\cdot) := \mathcal{V}^\dagger(\cdot) + \text{Tr}[(\text{id}_{B'} - \mathcal{V}^\dagger)(\cdot)]\tau_B \quad (\text{D6})$$

is a channel that is a left inverse of $\mathcal{V}_{B \rightarrow B'}$ [7, Section 4.6.3]. Thus, $\mathcal{R}_{B' \rightarrow B}^\mathcal{V}$ is a conditionally unital channel, from which we conclude that

$$\mathbf{H}(A|B')_\omega \leq \mathbf{H}(A|B)_{\mathcal{R}^\mathcal{V}(\omega)} = \mathbf{H}(A|B)_\rho. \quad (\text{D7})$$

This concludes the proof.

Appendix E: A subclass of conditional entropies

Here we consider a subclass of conditional entropies in which the postulate in (22) is replaced by monotonicity under conditionally unital channels. Since the set of conditionally unital channels contains the set of locally balanced channels, every function that is a conditional entropy according to this alternative definition is also a conditional entropy \mathbf{H} , as defined in (21)–(23). We will later see that this approach is consistent and sufficient for deriving various properties of conditional entropy.

Let us begin by defining another version of a conditional majorization pre-order on bipartite quantum states, based on conditionally unital channels exclusively. We call this approach relaxed conditional majorization. Given two bipartite states ρ_{AB} and $\sigma_{A'B'}$, we say that ρ_{AB} conditionally majorizes $\sigma_{A'B'}$ in the relaxed sense and with respect to system A , and write

$$\rho_{AB} \succ^A \sigma_{A'B'} \quad (\text{E1})$$

if either $|A'| \geq |A|$, there exists an isometric channel $\mathcal{V}_{A \rightarrow A'}$, and a conditionally unital channel $\mathcal{N}_{AB \rightarrow AB'}$, such that

$$\sigma_{A'B'} = \mathcal{V}_{A \rightarrow A'} \circ \mathcal{N}_{AB \rightarrow AB'}(\rho_{AB}), \quad (\text{E2})$$

or $|A| \geq |A'|$, there exists an isometric channel $\mathcal{U}_{A' \rightarrow A}$, and a conditionally unital channel $\mathcal{N}_{AB \rightarrow AB'}$, such that

$$\mathcal{U}_{A' \rightarrow A}(\sigma_{A'B'}) = \mathcal{N}_{AB \rightarrow AB'}(\rho_{AB}). \quad (\text{E3})$$

We define a function f to be a conditional uncertainty measure in the relaxed sense if it is not equal to the zero function and if it reverses the relaxed conditional majorization pre-order, i.e.,

$$\rho_{AB} \succ^A \sigma_{A'B'} \Rightarrow f(\rho_{AB}) \leq f(\sigma_{A'B'}). \quad (\text{E4})$$

We now define this other notion of conditional entropy. Formally, a function

$$\mathbf{H}^\blacktriangle : \bigcup_{A,B} \mathfrak{D}(AB) \rightarrow \mathbb{R} \quad (\text{E5})$$

is a quantum conditional entropy if it is not equal to the zero function and satisfies the following postulates for all systems A, B, A', B' and states ρ_{AB} and $\sigma_{A'B'}$:

1. Monotonicity:

$$\rho_{AB} \succ^A \sigma_{A'B'} \Rightarrow \mathbf{H}^\blacktriangle(A|B)_\rho \leq \mathbf{H}^\blacktriangle(A'|B')_\sigma. \quad (\text{E6})$$

2. Additivity:

$$\mathbf{H}^\blacktriangle(AA'|BB')_{\rho \otimes \sigma} = \mathbf{H}^\blacktriangle(A|B)_\rho + \mathbf{H}^\blacktriangle(A'|B')_\sigma. \quad (\text{E7})$$

Let us now recall a different definition of conditional min-entropy from [15]

$$H_{\min}^\uparrow(A|B)_\rho := \sup_{\sigma_B \in \mathfrak{D}(B)} H_{\min}(\rho_{AB}|\sigma_B), \quad (\text{E8})$$

where

$$H_{\min}(\rho_{AB}|\sigma_B) := - \inf_{\lambda \geq 0} \log_2 \{ \lambda : \rho_{AB} \leq \lambda I_A \otimes \sigma_B \}. \quad (\text{E9})$$

Note that $H_{\min}^\uparrow(A|B)_\rho$ is a conditional entropy $\mathbf{H}^\blacktriangle$ because it is additive and [15, Lemma 3.1.12] implies that $H_{\min}^\uparrow(A|B)_\rho$ is non-decreasing under conditionally unital channels.

We then have the following findings, analogous to those from Theorem 1 and Corollary 1, respectively.

Theorem 4 *Let $\mathbf{H}^\blacktriangle$ be a quantum conditional entropy, as defined in (E5)–(E7). Then, for every state ρ_{AB} ,*

$$\mathbf{H}^\blacktriangle(A|B)_\rho \geq H_{\min}^\uparrow(A|B)_\rho. \quad (\text{E10})$$

Equality is attained in (E10) if ρ_{AB} is equal to a maximally entangled state $\Phi_{A'B'}^{(k)}$, by the action of local isometric channels. Thus,

$$\mathbf{H}^\blacktriangle(A|B)_{\Phi^{(k)}} = - \log_2 k \quad (\text{E11})$$

for every conditional entropy $\mathbf{H}^\blacktriangle$.

Corollary 3 *Let ρ_{AB} be a bipartite state. Then, the following are equivalent:*

1. *For every conditional entropy $\mathbf{H}^\blacktriangle$,*

$$\mathbf{H}^\blacktriangle(A|B)_\rho \geq 0. \quad (\text{E12})$$

2. There exists a state σ_B such that $I_A \otimes \sigma_B \geq \rho_{AB}$.

Proof. The proof of Corollary 3 is immediate from Theorem 4. Indeed, if the first condition holds, then the inequality $H_{\min}^\dagger(A|B)_\rho \geq 0$ holds in particular (since $H_{\min}^\dagger(A|B)_\rho$ is a conditional entropy), and the second condition is then a consequence of the definition in (E8). Now suppose that the second condition holds. This then means that there exists $\lambda \leq 1$ and a state σ_B such that $\rho_{AB} \leq \lambda I_A \otimes \sigma_B$. According to the definition in (E8), it follows that $H_{\min}^\dagger(A|B)_\rho \geq 0$ and then Theorem 4 implies the first condition. ■

We have opted to emphasize the conditional entropy \mathbf{H} over $\mathbf{H}^\blacktriangle$ in the main text because

1. locally balanced channels have a strong motivation in terms of no information leakage from Alice to Bob, as would be expected for a measure of conditional uncertainty,
2. conditional majorization defined in this way reduces to classical conditional majorization, as defined in [35], for the classical case
3. the conditional min-entropy in (29) is in fact the smallest conditional entropy among all conditional entropies consistent with both sets of postulates,
4. every conditional entropy defined in this other way is in fact a conditional entropy \mathbf{H} , because every function that is monotone under all conditionally unital channels is also monotone under the subset of all locally balanced channels.

1. Proof of Theorem 4

Let σ_B be a state that achieves the optimal value of $H_{\min}^\dagger(A|B)_\rho$, in the definition in (E8).

Let us first consider the case when $H_{\min}^\dagger(A|B)_\rho \geq 0$. Set

$$k := |A|, \quad (\text{E13})$$

$$m := \left\lceil 2^{H_{\min}^\dagger(A|B)_\rho} \right\rceil, \quad (\text{E14})$$

and let X be a classical system with dimension k . By the assumption that $H_{\min}^\dagger(A|B)_\rho \geq 0$, and given the dimension bound $H_{\min}^\dagger(A|B)_\rho \leq \log_2 k$ [51, Lemma 5.2], it follows that $m \in [k]$.

Without loss of generality, we can suppose that $k > m$; if it is not the case, then we can embed the system A of ρ_{AB} in a larger Hilbert space, so that $H_{\min}^\dagger(A|B)_\rho$ stays constant but k increases. Set Π_X to be a projection onto an m -dimensional subspace of X (i.e., set $\Pi_X := \sum_{x=1}^m |x\rangle\langle x|_X$, where $\{|x\rangle\}_x$ is an orthonormal

basis specifying the classical system X), and define $\tau_{AB}^{(m)}$ as follows:

$$\tau_{AB}^{(m)} := \frac{I_A \otimes \sigma_B - m\rho_{AB}}{k - m}. \quad (\text{E15})$$

The operator $\tau_{AB}^{(m)}$ is positive semi-definite because $k - m > 0$ and

$$\begin{aligned} & \frac{1}{m} I_A \otimes \sigma_B - \rho_{AB} \\ & \geq 2^{-H_{\min}^\dagger(A|B)_\rho} I_A \otimes \sigma_B - \rho_{AB} \end{aligned} \quad (\text{E16})$$

$$= 2^{D_{\max}(\rho_{AB} \| I_A \otimes \sigma_B)} I_A \otimes \sigma_B - \rho_{AB} \quad (\text{E17})$$

$$\geq 0. \quad (\text{E18})$$

Also, $\tau_{AB}^{(m)}$ has trace equal to one and is thus a state. Let us now define the following measure-and-prepare channel:

$$\begin{aligned} \mathcal{N}_{X \rightarrow AB}(\omega_X) & := \text{Tr}[\Pi_X \omega_X] \rho_{AB} \\ & \quad + \text{Tr}[(I_X - \Pi_X) \omega_X] \tau_{AB}^{(m)}. \end{aligned} \quad (\text{E19})$$

Indeed, the action of this channel is to perform a measurement according to the POVM $\{\Pi_X, I_X - \Pi_X\}$ and prepare the state ρ_{AB} if the first outcome is obtained and the state $\tau_{AB}^{(m)}$ if the second outcome is obtained. It should be understood that Alice has access to the input system X .

This channel is conditionally unital because

$$\begin{aligned} \mathcal{N}_{X \rightarrow AB}(I_X) & \\ & = \text{Tr}[\Pi_X I_X] \rho_{AB} + \text{Tr}[(I_X - \Pi_X) I_X] \tau_{AB}^{(m)} \end{aligned} \quad (\text{E20})$$

$$= m\rho_{AB} + (k - m) \tau_{AB}^{(m)} \quad (\text{E21})$$

$$= I_A \otimes \sigma_B. \quad (\text{E22})$$

Also, if we input the uniform state $\mathbf{u}_X^\Pi := \Pi_X/m$ on the subspace onto which Π_X projects, the output is given by

$$\mathcal{N}_{X \rightarrow AB}(\mathbf{u}_X^\Pi) = \rho_{AB}. \quad (\text{E23})$$

Putting these observations together, we conclude that

$$\log_2 \left[2^{H_{\min}^\dagger(A|B)_\rho} \right] = \mathbf{H}^\blacktriangle(X)_{\mathbf{u}^\Pi} \quad (\text{E24})$$

$$\leq \mathbf{H}^\blacktriangle(A|B)_{\mathcal{N}(\mathbf{u}^\Pi)} \quad (\text{E25})$$

$$= \mathbf{H}^\blacktriangle(A|B)_\rho. \quad (\text{E26})$$

The first equality follows because the entropy of a uniform state is equal to the logarithm of its rank and from the definition of m in (E14). The inequality follows from the fact that $\mathcal{N}_{X \rightarrow AB}$ is conditionally unital (see (E20)–(E22)) and $\mathbf{H}^\blacktriangle$ is a quantum conditional entropy, thus obeying the first postulate in (E6). The final equality follows from (E23).

Therefore, since all conditional entropies are additive

for tensor-product states, we conclude that

$$\mathbf{H}^\bullet(A|B)_\rho = \lim_{n \rightarrow \infty} \frac{1}{n} \mathbf{H}^\bullet(A^n|B^n)_{\rho^{\otimes n}} \quad (\text{E27})$$

$$\geq \lim_{n \rightarrow \infty} \frac{1}{n} \log \left[2^{H_{\min}^\dagger(A^n|B^n)_{\rho^{\otimes n}}} \right] \quad (\text{E28})$$

$$= \lim_{n \rightarrow \infty} \frac{1}{n} \log \left[2^{n H_{\min}^\dagger(A|B)_\rho} \right] \quad (\text{E29})$$

$$= H_{\min}^\dagger(A|B)_\rho. \quad (\text{E30})$$

The inequality follows from (E24)–(E26). The second equality follows from additivity, and the final equality from the assumption that $H_{\min}^\dagger(A|B)_\rho \geq 0$.

Now let us consider the case when $H_{\min}^\dagger(A|B)_\rho < 0$. Let

$$t := 2^{-H_{\min}^\dagger(A|B)_\rho}, \quad (\text{E31})$$

$$k' := \lceil t \rceil, \quad (\text{E32})$$

$$k := |A|. \quad (\text{E33})$$

Observe that $t > 1$ by assumption. Also, observe that t is the smallest real satisfying $\rho_{AB} \leq t I_A \otimes \sigma_B$. Let A' be a system of dimension k' , and let X be a classical system of dimension $|X| := k k'$. Now define

$$\tau_{AB}^{(k,k')} := \frac{k' I_A \otimes \sigma_B - \rho_{AB}}{k k' - 1}, \quad (\text{E34})$$

and observe that $\tau_{AB}^{(k,k')}$ is positive semi-definite because $k k' > 1$ and

$$\begin{aligned} & k' I_A \otimes \sigma_B - \rho_{AB} \\ & \geq 2^{-H_{\min}^\dagger(A|B)_\rho} I_A \otimes \sigma_B - \rho_{AB} \end{aligned} \quad (\text{E35})$$

$$= 2^{D_{\max}(\rho_{AB} \| I_A \otimes \sigma_B)} I_A \otimes \sigma_B - \rho_{AB} \quad (\text{E36})$$

$$\geq 0. \quad (\text{E37})$$

Since $\tau_{AB}^{(k,k')}$ has trace equal to one, it follows that it is a state. We then construct the following channel:

$$\begin{aligned} \mathcal{N}_{X \rightarrow AA'B}(\omega_X) & := \\ & \left(\text{Tr}[|1\rangle\langle 1|_X \omega_X] \rho_{AB} + \text{Tr}[\Pi_X \omega_X] \tau_{AB}^{(k,k')} \right) \otimes \mathbf{u}_{A'}, \end{aligned} \quad (\text{E38})$$

where

$$\Pi_X := I_X - |1\rangle\langle 1|_X, \quad (\text{E39})$$

so that $\text{Tr}[\Pi_X] = k k' - 1$. Also, it should be understood that Alice has access to the X input system. Consider that

$$\mathcal{N}_{X \rightarrow AA'B}(|1\rangle\langle 1|_X) = \rho_{AB} \otimes \mathbf{u}_{A'}. \quad (\text{E40})$$

Furthermore, the channel is conditionally unital because

$$\begin{aligned} & \mathcal{N}_{X \rightarrow AA'B}(I_X) \\ & = \left(\text{Tr}[|1\rangle\langle 1|_X I_X] \rho_{AB} + \text{Tr}[\Pi_X I_X] \tau_{AB}^{(k,k')} \right) \otimes \mathbf{u}_{A'} \end{aligned} \quad (\text{E41})$$

$$= \left(\rho_{AB} + (k k' - 1) \tau_{AB}^{(k,k')} \right) \otimes \mathbf{u}_{A'} \quad (\text{E42})$$

$$= k' I_A \otimes \sigma_B \otimes \mathbf{u}_{A'} \quad (\text{E43})$$

$$= I_A \otimes \sigma_B \otimes I_{A'}. \quad (\text{E44})$$

Then it follows that

$$0 = \mathbf{H}^\bullet(X)_{|1\rangle\langle 1|} \quad (\text{E45})$$

$$\leq \mathbf{H}^\bullet(AA'|B)_{\mathcal{N}(|1\rangle\langle 1|)} \quad (\text{E46})$$

$$= \mathbf{H}^\bullet(AA'|B)_{\rho \otimes \mathbf{u}} \quad (\text{E47})$$

$$= \mathbf{H}^\bullet(A|B)_\rho + \mathbf{H}^\bullet(A')_{\mathbf{u}} \quad (\text{E48})$$

$$= \mathbf{H}^\bullet(A|B)_\rho + \log_2 \left[2^{-H_{\min}^\dagger(A|B)_\rho} \right]. \quad (\text{E49})$$

The first equality follows because the entropy of a pure state is equal to zero. The inequality follows because $\mathcal{N}_{X \rightarrow AA'B}$ is a conditionally unital channel and \mathbf{H}^\bullet is a quantum conditional entropy, thus obeying the first postulate in (E6). The second equality follows from (E40). The penultimate equality follows from the additivity postulate of quantum conditional entropy. The final equality follows because the entropy of the uniform state $\mathbf{u}_{A'}$ is equal to the logarithm of its rank, which is in turn equal to $\left[2^{-H_{\min}^\dagger(A|B)_\rho} \right]$.

Now, since $\mathbf{H}^\bullet(A|B)_\rho$ is additive for tensor-product states, we find that

$$\mathbf{H}^\bullet(A|B)_\rho = \lim_{n \rightarrow \infty} \frac{1}{n} \mathbf{H}^\bullet(A^n|B^n)_{\rho^{\otimes n}} \quad (\text{E50})$$

$$\geq \lim_{n \rightarrow \infty} -\frac{1}{n} \log \left[2^{-H_{\min}^\dagger(A^n|B^n)_{\rho^{\otimes n}}} \right] \quad (\text{E51})$$

$$= \lim_{n \rightarrow \infty} -\frac{1}{n} \log \left[2^{-n H_{\min}^\dagger(A|B)_\rho} \right] \quad (\text{E52})$$

$$= \lim_{n \rightarrow \infty} -\frac{1}{n} \log \lceil t^n \rceil \quad (\text{E53})$$

$$\geq \lim_{n \rightarrow \infty} -\frac{1}{n} \log (t^n + 1) \quad (\text{E54})$$

$$= -\log t = H_{\min}^\dagger(A|B)_\rho. \quad (\text{E55})$$

The first inequality follows from (E45)–(E49). The second equality follows from additivity, and the final equality from the assumption that $t > 1$. This completes the proof of the lower bound.

It is left to prove the equality on maximally entangled states. Since conditional entropy is invariant under local isometries we can assume without loss of generality that $k := |A| = |B|$. Now, consider the state $\Phi_{AB}^{(k)} \otimes \mathbf{u}_{\tilde{A}}$ where $|\tilde{A}| = k$, and observe that by introducing a trivial system B_2 such that $|B_2| = 1$ we get

$$\begin{aligned} & \mathbf{H}^\bullet(A\tilde{A}|B)_{\Phi_{AB}^{(k)} \otimes \mathbf{u}_{\tilde{A}}} \\ & = \mathbf{H}^\bullet(A\tilde{A}|BB_2)_{\Phi_{AB}^{(k)} \otimes \mathbf{u}_{\tilde{A}B_2}} \end{aligned} \quad (\text{E56})$$

$$= \mathbf{H}^\bullet(A|B)_{\Phi^{(k)}} + \mathbf{H}^\bullet(\tilde{A}|B_2)_{\mathbf{u}_{\tilde{A}B_2}} \quad (\text{E57})$$

$$= \mathbf{H}^\bullet(A|B)_{\Phi^{(k)}} + \mathbf{H}^\bullet(\mathbf{u}_{\tilde{A}}) \quad (\text{E58})$$

$$= \mathbf{H}^\bullet(A|B)_{\Phi^{(k)}} + \log k, \quad (\text{E59})$$

where we used the additivity property of conditional entropy and the fact that the entropy of the uniform state $\mathbf{u}_{\tilde{A}}$ is $\log_2 k$. It is therefore sufficient to show that

$$\mathbf{H}^\bullet(A\tilde{A}|B)_{\Phi_{AB}^{(k)} \otimes \mathbf{u}_{\tilde{A}}} \leq 0 \quad (\text{E60})$$

since the above equation implies that

$$\mathbf{H}^\mathbf{A}(A|B)_{\Phi^{(k)}} \leq -\log k = H_{\min}^\uparrow(A|B)_{\Phi^{(k)}}. \quad (\text{E61})$$

For this purpose, let X be a classical system of dimension $|X| = k^2$ and let $\mathcal{N}_{A\bar{A}B \rightarrow X}$ a quantum channel defined by

$$\mathcal{N}_{A\bar{A}B \rightarrow X} := \mathcal{E}_{AB \rightarrow X} \circ \text{Tr}_{\bar{A}}, \quad (\text{E62})$$

where

$$\begin{aligned} \mathcal{E}_{AB \rightarrow X}(\omega_{AB}) &:= \text{Tr}[\Phi_{AB}^k \omega_{AB}] |1\rangle\langle 1|_X \\ &+ \text{Tr}[(I_{AB} - \Phi_{AB}^{(k)}) \omega_{AB}] \frac{I_X - |1\rangle\langle 1|_X}{k^2 - 1}. \end{aligned} \quad (\text{E63})$$

This channel is conditionally unital because

$$\begin{aligned} \mathcal{N}_{A\bar{A}B \rightarrow X}(I_A \otimes I_{\bar{A}} \otimes \sigma_B) \\ = (\mathcal{E}_{AB \rightarrow X} \circ \text{Tr}_{\bar{A}})(I_A \otimes I_{\bar{A}} \otimes \sigma_B) \end{aligned} \quad (\text{E64})$$

$$= k \mathcal{E}_{AB \rightarrow X}(I_A \otimes \sigma_B) \quad (\text{E65})$$

$$= k \left(\begin{array}{c} \text{Tr}[\Phi_{AB}^{(k)}(I_A \otimes \sigma_B)] |1\rangle\langle 1|_X \\ + \text{Tr}[(I_{AB} - \Phi_{AB}^{(k)})(I_A \otimes \sigma_B)] \frac{I_X - |1\rangle\langle 1|_X}{k^2 - 1} \end{array} \right) \quad (\text{E66})$$

$$= k \left(\frac{1}{k} |1\rangle\langle 1|_X + \left(k - \frac{1}{k}\right) \frac{I_X - |1\rangle\langle 1|_X}{k^2 - 1} \right) \quad (\text{E67})$$

$$= |1\rangle\langle 1|_X + (k^2 - 1) \frac{I_X - |1\rangle\langle 1|_X}{k^2 - 1} \quad (\text{E68})$$

$$= I_X. \quad (\text{E69})$$

Also, consider that

$$\mathcal{N}_{A\bar{A}B \rightarrow X}(\Phi_{AB}^{(k)} \otimes \mathbf{u}_{\bar{A}}) = |1\rangle\langle 1|_X. \quad (\text{E70})$$

Then

$$\mathbf{H}^\mathbf{A}(A|B)_{\Phi^{(k)}} + \log k = \mathbf{H}^\mathbf{A}(A\bar{A}|B)_{\Phi^{(k)} \otimes \mathbf{u}} \quad (\text{E71})$$

$$\leq \mathbf{H}^\mathbf{A}(X)_{\mathcal{N}(\Phi^{(k)} \otimes \mathbf{u})} \quad (\text{E72})$$

$$= \mathbf{H}^\mathbf{A}(X)_{|1\rangle\langle 1|} \quad (\text{E73})$$

$$= 0. \quad (\text{E74})$$

The first equality follows from (E56)–(E59). The inequality follows because the channel $\mathcal{N}_{A\bar{A}B \rightarrow X}$ is conditionally unital. The second equality follows from (E70) and the last because the entropy of every pure state is equal to zero. This concludes the proof.

2. Proof of Theorem 1

Here the idea is essentially the same as that given in Appendix E1, except that we need to construct channels that are locally balanced (i.e., both conditionally unital and $A \not\rightarrow B$ semi-causal).

First, throughout, we make the following replacement in the proof:

$$H_{\min}^\uparrow(A|B)_\rho \rightarrow H_{\min}(A|B)_\rho. \quad (\text{E75})$$

To handle the case when $H_{\min}(A|B)_\rho \geq 0$, we employ the same channel in (E19), with the exception that we set

$$\tau_{AB}^{(m)} := \frac{I_A \otimes \rho_B - m\rho_{AB}}{k - m}. \quad (\text{E76})$$

All of the steps in (E13)–(E30) go through with the replacements in (E75) and (E76). We now verify that the $A \not\rightarrow B$ semi-causal constraint holds for this channel. Consider, for the modified channel, that

$$\begin{aligned} &\text{Tr}_A[\mathcal{N}_{X \rightarrow AB}(\omega_X)] \\ &= \text{Tr}[\Pi_X \omega_X] \text{Tr}_A[\rho_{AB}] + \text{Tr}[(I_X - \Pi_X) \omega_X] \text{Tr}_A[\tau_{AB}^{(m)}] \\ &= \text{Tr}[\Pi_X \omega_X] \rho_B + \text{Tr}[(I_X - \Pi_X) \omega_X] \rho_B \\ &= \rho_B. \end{aligned} \quad (\text{E77})$$

Thus, the output state of B has no dependence on the input state ω_X when system A is traced out, so that the semi-causal constraint holds.

Now let us consider the case when $H_{\min}(A|B)_\rho < 0$. We employ the same channel in (E38), with the exception that we set

$$\tau_{AB}^{(k,k')} := \frac{k' I_A \otimes \rho_B - \rho_{AB}}{kk' - 1}. \quad (\text{E78})$$

Then all of the steps in (E31)–(E55) go through, with the replacements in (E75) and (E78). We now verify that the $A \not\rightarrow B$ semi-causal constraint holds for this channel. Consider, for the modified channel, that

$$\begin{aligned} &\text{Tr}_{AA'}[\mathcal{N}_{X \rightarrow AA'B}(\omega_X)] \\ &= \left(\text{Tr}[|1\rangle\langle 1|_X \omega_X] \text{Tr}_A[\rho_{AB}] + \text{Tr}[\Pi_X \omega_X] \text{Tr}_A[\tau_{AB}^{(k,k')}] \right) \\ &\quad \otimes \text{Tr}_{A'}[\mathbf{u}_{A'}] \end{aligned} \quad (\text{E79})$$

$$= \text{Tr}[|1\rangle\langle 1|_X \omega_X] \rho_B + \text{Tr}[\Pi_X \omega_X] \rho_B \quad (\text{E80})$$

$$= \rho_B. \quad (\text{E81})$$

Thus, the output state of B has no dependence on the input state ω_X when system A is traced out, so that the semi-causal constraint holds. This establishes the inequality in (E10).

Finally, we consider the equality in (E11). The argument here is actually exactly the same as in (E56)–(E74), except using the replacement in (E75). We just need to verify that the channel in (E62) is $A \not\rightarrow B$ semi-causal. However, the only output is system X , which belongs to Alice. Thus, after tracing it out, there is no remaining system and the channel is then semi-causal.

Appendix F: Proof of Theorem 2

First, we demonstrate that \mathbf{H} satisfies the monotonicity property of conditional entropy. Let $\mathcal{N}_{AB \rightarrow A'B'}$ be a locally balanced channel, and consider the bipartite density matrix ρ_{AB} . We begin by considering the case where

$A = A'$ and

$$\mathbf{H}(A|B)_{\mathcal{N}(\rho)} = \log |A| - \mathbf{D}(\mathcal{N}(\rho_{AB}) \parallel \mathbf{u}_A \otimes \text{Tr}_A[\mathcal{N}(\rho_{AB})]) \quad (\text{F1})$$

and from using the fact that \mathcal{N} is semi-causal and conditionally unital, we find that

$$\begin{aligned} \mathbf{u}_A \otimes \text{Tr}_A[\mathcal{N}(\rho_{AB})] &= \mathbf{u}_A \otimes \text{Tr}_A[\mathcal{N}(\mathbf{u}_A \otimes \rho_B)] \quad (\text{F2}) \\ &= \mathcal{N}(\mathbf{u}_A \otimes \rho_B), \quad (\text{F3}) \end{aligned}$$

from which we have

$$\begin{aligned} \mathbf{H}(A|B)_{\mathcal{N}(\rho)} &= \log |A| - \mathbf{D}(\mathcal{N}(\rho_{AB}) \parallel \mathcal{N}(\mathbf{u}_A \otimes \rho_B)) \\ &\geq \log |A| - \mathbf{D}(\rho_{AB} \parallel \mathbf{u}_A \otimes \rho_B), \quad (\text{F4}) \end{aligned}$$

where the last line follows from the data-processing inequality.

We also need to prove that it is invariant under the action of a local isometric channel acting on system A . Let $\mathcal{V}_{A \rightarrow A'}$ be an arbitrary isometric channel, and let C represent a Hilbert space such that $|C| = |A'| - |A|$. Then

$$\mathbf{H}(A|B)_{\mathcal{V}(\rho)} = \log |A'| - \mathbf{D}(\mathcal{V}_{A \rightarrow A'}(\rho_{AB}) \parallel \mathbf{u}_{A'} \otimes \rho_B). \quad (\text{F5})$$

Clearly, if \mathcal{V} is a unitary channel such that $A \simeq A'$, then $\mathbf{H}(A'|B)_{\mathcal{V}(\rho)} = \mathbf{H}(A|B)_\rho$. Then without loss of generality, we assume that

$$\mathcal{V}_{A \rightarrow A'}(\rho_{AB}) = \rho_{AB} \oplus \mathbf{0}_{CB} \quad (\text{F6})$$

$$= \begin{pmatrix} \rho_{AB} & \mathbf{0} \\ \mathbf{0} & \mathbf{0}_{CB} \end{pmatrix}, \quad (\text{F7})$$

because the conditional entropy is invariant under local unitaries. We can additionally rewrite

$$\mathbf{u}_{A'} = t\mathbf{u}_A \oplus (1-t)\mathbf{u}_C, \quad (\text{F8})$$

where $t := \frac{|A|}{|A'|}$. From this, it follows that

$$\begin{aligned} &\mathbf{D}(\mathcal{V}_{A \rightarrow A'}(\rho_{AB}) \parallel \mathbf{u}_{A'} \otimes \rho_B) \\ &= \mathbf{D}(\rho_{AB} \oplus \mathbf{0}_{CB} \parallel (t\mathbf{u}_A \otimes \rho_B) \oplus ((1-t)\mathbf{u}_C \otimes \rho_B)) \\ &= \mathbf{D}(\rho_{AB} \parallel t\mathbf{u}_A \otimes \rho_B) \\ &= \mathbf{D}(\rho_{AB} \parallel \mathbf{u}_A \otimes \rho_B) - \log(t), \quad (\text{F9}) \end{aligned}$$

where the second equality follows from a property of relative entropy proved in [52]. Thus, upon substitution, we have that

$$\begin{aligned} \mathbf{H}(A'|B)_{\mathcal{V}(\rho)} &= \log |A'| - \mathbf{D}(\rho_{AB} \parallel \mathbf{u}_A \otimes \rho_B) + \log(t) \\ &= \mathbf{H}(A|B)_\rho. \quad (\text{F10}) \end{aligned}$$

Finally, additivity of \mathbf{H} follows immediately from additivity of \mathbf{D} .

1. Conditional min-entropy is a conditional entropy

Here we justify that the conditional min-entropy is indeed a conditional entropy according to the definition in (21)–(23). Indeed, we use the definition of conditional min-entropy in (29) to rewrite it in terms of the max-relative entropy $D_{\max}(\rho \parallel \sigma)$ as

$$H_{\min}(A|B)_\rho = \log_2 |A| - D_{\max}(\rho_{AB} \parallel \mathbf{u}_A \otimes \rho_B), \quad (\text{F11})$$

where the max-relative entropy of states ρ and σ is defined as [53]:

$$D_{\max}(\rho \parallel \sigma) := \log_2 \inf_{\lambda \geq 0} \{\rho \leq \lambda \sigma\}. \quad (\text{F12})$$

Then we apply Theorem 2.

Appendix G: Example of non-negative weakly additive monotonic function

In this section we prove that there exists a non-zero function that satisfies the following three properties:

1. Monotonicity under conditional majorization; i.e., satisfying (22).
2. Weak additivity of the form (37).
3. Non-negative.

The third property above demonstrates that the negativity of conditional entropy is closely related to the full additivity postulate. The construction of the example is based on a technique developed in [37] for the extension of resource monotones from one domain to another.

Let \mathbb{H} be a classical conditional-entropy. In [37] it was shown that any such function has two optimal extensions to the quantum domain. Here we are interested in the following extension. For any $\rho \in \mathfrak{D}(AB)$ we define

$$f(A|B)_\rho := \inf_{\rho_{AB} \succ_A \sigma_{XY}} \mathbb{H}(X|Y)_\sigma, \quad (\text{G1})$$

where the infimum above is over all classical systems X and Y and all classical states σ_{XY} with the property that $\rho_{AB} \succ_A \sigma_{XY}$. The function f has the following key properties [37]:

1. Reduction: For all classical states $\rho \in \mathfrak{D}(XY)$

$$f(X|Y)_\rho = \mathbb{H}(X|Y)_\rho. \quad (\text{G2})$$

2. Monotonicity: For all $\rho \in \mathfrak{D}(AB)$ and $\omega \in \mathfrak{D}(A'B')$ with $\rho_{AB} \succ_A \omega_{A'B'}$

$$f(A'|B')_\omega \geq f(A|B)_\rho. \quad (\text{G3})$$

3. Subadditivity: The function f is subadditive under tensor products.

4. Positivity: The function f is non-negative.

The last property follows from the definition in (G1). Since f is subadditive, the limit

$$g(A|B)_\rho := \lim_{n \rightarrow \infty} \frac{1}{n} f(A^n|B^n)_{\rho^{\otimes n}} \quad (\text{G4})$$

exists. Therefore, the function g is weakly additive.

Moreover, the function g satisfies the first two properties above, namely, reduction to \mathbb{H} on classical states and behaves monotonically under conditional majorization. Observe that the reduction of g to \mathbb{H} on classical states implies that it is not equal to the zero function since \mathbb{H} is not the zero function. Since g is non-negative it cannot be a conditional entropy. This means that g cannot be fully additive, as this is the only property that does not follow directly from its definition.