

A CHARACTERISATION OF THE SUBSPACE OF RADIALY SYMMETRIC FUNCTIONS IN SOBOLEV SPACES

MATTHIAS OSTERMANN

ABSTRACT. This article studies the classical Sobolev norm on the subspace of radially symmetric functions. An equivalent norm is presented and an isomorphism between the radial subspace and a weighted Sobolev space is established for all non-negative integer orders in all space dimensions.

1. INTRODUCTION

Sobolev spaces are fundamental in the modern analysis of partial differential equations. For $p \geq 1$ and integers $k \geq 0$ the *Sobolev space* $W^{k,p}(\mathbb{B}_r^d)$ is realized by the completion of the space $C^\infty(\overline{\mathbb{B}_r^d})$ of smooth functions on the open ball $\mathbb{B}_r^d \subset \mathbb{R}^d$ of radius $r > 0$ centred about $0 \in \mathbb{R}^d$, whose derivatives of all orders are continuous up to the boundary, with respect to the *classical Sobolev norm*

$$\|\varphi\|_{W^{k,p}(\mathbb{B}_r^d)} = \left(\sum_{0 \leq |\alpha| \leq k} \|\partial^\alpha \varphi\|_{L^p(\mathbb{B}_r^d)}^p \right)^{\frac{1}{p}}$$

for $\varphi \in C^\infty(\overline{\mathbb{B}_r^d})$. On \mathbb{R}^d , the Sobolev norm may be arranged in terms of *classical homogeneous Sobolev norms*

$$\|\varphi\|_{\dot{W}^{k,p}(\mathbb{R}^d)} = \left(\sum_{|\alpha|=k} \|\partial^\alpha \varphi\|_{L^p(\mathbb{R}^d)}^p \right)^{\frac{1}{p}}$$

for $\varphi \in C_c^\infty(\mathbb{R}^d)$. The homogeneous Sobolev space $\dot{W}^{k,p}(\mathbb{R}^d)$ is the completion of the space $C_c^\infty(\mathbb{R}^d)$ of compactly supported smooth functions with respect to the norm $\|\cdot\|_{\dot{W}^{k,p}(\mathbb{R}^d)}$. In many situations, the presence of symmetries leads to function spaces that possess those symmetries. We shall be dealing with radial symmetry and consider the radial subspaces

$$W_{\text{rad}}^{k,p}(\mathbb{B}_r^d) \subset W^{k,p}(\mathbb{B}_r^d) \quad \text{and} \quad \dot{W}_{\text{rad}}^{k,p}(\mathbb{R}^d) \subset \dot{W}^{k,p}(\mathbb{R}^d)$$

which arise from the completion of the spaces of smooth radial functions $C_{\text{rad}}^\infty(\overline{\mathbb{B}_r^d}) \subset C^\infty(\overline{\mathbb{B}_r^d})$ and $C_{\text{rad}}^\infty(\mathbb{R}^d) \cap C_c^\infty(\mathbb{R}^d)$ endowed with $\|\cdot\|_{W^{k,p}(\mathbb{B}_r^d)}$ and $\|\cdot\|_{\dot{W}^{k,p}(\mathbb{R}^d)}$, respectively. The spaces obtained by completion agree with the usual definition in terms of weak derivatives, see e.g. [1, 7]. Also, recall that any smooth radial function φ is represented by a smooth even function f_φ , which in turn is the composition of a smooth function \tilde{f}_φ with the square function, yielding two canonical representations

$$\varphi(x) = f_\varphi(\rho) \quad \text{and} \quad \varphi(x) = \tilde{f}_\varphi(\rho^2), \quad \rho = |x|,$$

and vice versa, see section 2. From differentiating these equations, it is apparent that Sobolev norms will be measured in terms of *weighted Sobolev norms*, which we define for a collection $w = (w_j)_{j=0}^k$ of continuous and almost everywhere positive weight functions $w_j : [0, r] \rightarrow [0, \infty)$ by

$$\|f\|_{W^{k,p}((0,r),w)} = \left(\sum_{j=0}^k \left\| f^{(j)} w_j^{\frac{1}{p}} \right\|_{L^p(0,r)}^p \right)^{\frac{1}{p}}$$

for $f \in C^\infty([0, r])$. The associated completion as well as homogeneous versions are defined analogously as above. However, since derivatives of the Euclidean length $|\cdot|$ become increasingly singular towards the origin, partial derivatives of radial functions expressed in terms of

This work has been supported by the Vienna School of Mathematics (VSM).

derivatives of their radial representative suffer term-wise artificial coordinate singularities. This prevents the equivalence between radial Sobolev norms and weighted Sobolev norms as presented by D. G. De Figueiredo, E. M. Dos Santos and O. H. Miyagaki in [10] for $0 \leq k < d/2 + 1$ to remain valid for high derivatives. We show that this spurious problem is avoided by replacing classical derivatives by

$$D_\rho f(\rho) := \rho^{-1} f'(\rho), \quad \rho > 0,$$

for $f \in C_e^\infty([0, r])$. Here,

$$C_e^\infty([0, r]) = \{f \in C^\infty([0, r]) \mid \forall n \in \mathbb{Z}_{\geq 0} : f^{(2n+1)}(0) = 0\}$$

is the space of smooth even functions, on which the above operation can be extended to all $\rho \in [0, r]$ to give a derivation

$$D : C_e^\infty([0, r]) \rightarrow C_e^\infty([0, r]), \quad f \mapsto (\rho \mapsto D_\rho f(\rho)). \quad (1)$$

This, then, leads to a generalization of [10, Theorem 2.3] to all radial Sobolev spaces of non-negative integer order in all space dimensions and thereby settles an open characterisation raised by W. Sickel, L. Skrzypczak and J. Vybiral in [14, Section 2.1.3, Remark 6].

Theorem 1. *Let $d, k \in \mathbb{Z}_{\geq 0}$ and $p, r \in \mathbb{R}$ with $d \geq 2$, $p \geq 1$, $r > 0$.*

(1) *The equivalence of norms*

$$\|\varphi\|_{W^{k,p}(\mathbb{B}_r^d)} \simeq \sum_{j=0}^k \left\| \left(\cdot \right)^{\frac{d-1}{p}+j} D^j f_\varphi \right\|_{L^p(0,r)} \simeq \sum_{j=0}^k \left\| \left(\cdot \right)^{\frac{d-2}{2p}+\frac{j}{2}} \tilde{f}_\varphi^{(j)} \right\|_{L^p(0,r^2)}$$

holds for all $\varphi \in C_{\text{rad}}^\infty(\overline{\mathbb{B}_r^d})$. In particular,

$$W_{\text{rad}}^{k,p}(\mathbb{B}_r^d) \quad \text{and} \quad W^{k,p}\left((0, r^2), w\right),$$

with weight $w = (w_j)_{j=0}^k$ given by $w_j = \left(\cdot \right)^{\frac{d-2+pi}{2}}$, are isomorphic as normed linear spaces.

(2) *The equivalence of norms*

$$\|\varphi\|_{\dot{W}^{k,p}(\mathbb{R}^d)} \simeq \left\| \left| \cdot \right|^{\frac{d-1}{p}+k} D^k f_\varphi \right\|_{L^p(\mathbb{R})} \simeq \left\| \left| \cdot \right|^{\frac{d-2}{2p}+\frac{k}{2}} \tilde{f}_\varphi^{(k)} \right\|_{L^p(\mathbb{R})}$$

holds for all $\varphi \in C_{\text{rad}}^\infty(\mathbb{R}^d) \cap C_c^\infty(\mathbb{R}^d)$. In particular,

$$\dot{W}_{\text{rad}}^{k,p}(\mathbb{R}^d) \quad \text{and} \quad \dot{W}^{k,p}\left(\mathbb{R}, \left| \cdot \right|^{\frac{d-2+pk}{2}}\right)$$

are isomorphic as normed linear spaces.

Pioneering radial lemmas for Sobolev spaces are the famous inequality by W. Strauss [16] and extensions thereof by W. M. Ni [12] and H. Berestycki and P. L. Lions [3]. By now, there are many versions of radial lemmas that deal with compact or weighted Sobolev embeddings with applications primarily to variational problems. D. G. De Figueiredo, E. M. Dos Santos and O. H. Miyagaki [10] gave an embedding of the radial subspace into a weighted Sobolev space which turns out to be an isomorphism precisely if $0 \leq k < d/2 + 1$. A similar isomorphism given by I. Glogić in the case $p = 2$ can be found in [9], who recently also proved a Strauss-type inequality for partial derivatives of radial functions, see [8, Appendix B]. Among the study of regularity and decay properties of radial functions in more general Besov and Lizorkin-Triebel spaces via atomic decomposition and differences, W. Sickel, L. Skrzypczak and J. Vybiral [14, 13, 15] also discussed characterisations of radial Sobolev spaces without having complete descriptions for higher order Sobolev spaces. To mention another different approach based on an Emden-Fowler transform, R. Musina proved in [11] dilation invariant inequalities where isomorphisms between Sobolev spaces on the line and a weighted Sobolev space were established.

If the domain is the whole Euclidean space, Sobolev norms can be given equivalently in terms of the Fourier transform. A new characterisation of Sobolev spaces on \mathbb{R}^d in terms of a so-called square function is given by R. Alabern, J. Mateu and J. Verdera in [2]. Remarkably, in odd space dimensions, properties of Bessel functions can be exploited in the radial Fourier transform in order to characterise the Sobolev norm as a one-dimensional Sobolev norm of a weighted radial representative, see the research by P. Biernat, R. Donninger, B. Schörkhuber and the author [4, 5]. This characterisation also holds on balls by means of Sobolev extension techniques, also see [6]. If the space dimension is even, this approach fails because the radial Fourier transform is not expressible as a Fourier transform on the line.

We want to point out, as noted in [8], that our elementary characterisation of Sobolev norms of radial functions will be of major importance for progress in the study of stability of self-similar blowup in nonlinear wave equations.

2. SMOOTH RADIAL FUNCTIONS

We say that a function $\varphi : \mathbb{B}_r^d \rightarrow \mathbb{C}$ is *radial* if for any $R \in SO(d)$

$$\varphi(Rx) = \varphi(x)$$

for all $x \in \mathbb{B}_r^d$. In this section we elucidate the relation between a radial function and its radial profile. We begin with an elementary fact which is attributed to H. Whitney [17].

Lemma 1. *A function $f : [0, r] \rightarrow \mathbb{C}$ belongs to $C_e^\infty([0, r])$ if and only if there is a function $\tilde{f} \in C^\infty([0, r])$ such that*

$$f(\rho) = \tilde{f}(\rho^2)$$

for all $\rho \in [0, r]$.

Proof. “ \Rightarrow ”: Let $f \in C_e^\infty([0, r])$. Define $\tilde{f} \in C^\infty((0, r^2])$ by $\tilde{f}(\varrho) = f(\sqrt{\varrho})$. We claim that for all $n \geq 1$

$$\tilde{f}^{(n)}(\varrho^2) = \frac{1}{2^{2n-1}(n-1)!} \int_0^1 (1-t^2)^{n-1} f^{(2n)}(t\varrho) dt, \quad \varrho > 0. \quad (2)$$

Indeed, differentiating the identity $\tilde{f}(\varrho^2) = f(\varrho)$ and using the fundamental theorem of calculus yields

$$\tilde{f}'(\varrho^2) = \frac{1}{2} \int_0^1 f''(t\varrho) dt.$$

Assuming that the claim (2) holds for an arbitrary but fixed $n \geq 1$, differentiating it and performing an integration by parts yields

$$\begin{aligned} 2\varrho \tilde{f}^{(n+1)}(\varrho^2) &= \frac{1}{2^{2n-1}(n-1)!} \int_0^1 (1-t^2)^{n-1} t f^{(2n+1)}(t\varrho) dt \\ &= -\frac{1}{2^{2n-1}(n-1)!} \int_0^1 \frac{1}{2n} \partial_t (1-t^2)^n f^{(2n+1)}(t\varrho) dt \\ &= \frac{\varrho}{2^{2n}n!} \int_0^1 (1-t^2)^n f^{(2(n+1))}(t\varrho) dt \end{aligned}$$

and hence the claim is proved. Thus $\lim_{\varrho \rightarrow 0} \tilde{f}^{(n)}(\varrho)$ exists for all $n \geq 1$, so $\tilde{f} \in C^\infty([0, r^2])$.

“ \Leftarrow ”: If $\tilde{f} \in C^\infty([0, r^2])$, then the function f defined by $f(\rho) = \tilde{f}(\rho^2)$ is smooth and even. \square

We arrive at a characterisation of smooth radial functions.

Lemma 2. *A function $\varphi : \mathbb{B}_r^d \rightarrow \mathbb{C}$ belongs to $C_{\text{rad}}^\infty(\overline{\mathbb{B}_r^d})$ if and only if there is a function $f \in C_e^\infty([0, r])$ such that*

$$\varphi(x) = f(|x|)$$

for all $x \in \mathbb{B}_r^d$.

Proof. “ \Rightarrow ”: Assume $\varphi \in C_{\text{rad}}^\infty(\overline{\mathbb{B}_r^d})$. Define $f \in C^\infty([0, r])$ by $f(\rho) = \varphi(\rho e_d)$, where $e_d \in \mathbb{R}^d$ is the d -th standard basis vector. Then f is a smooth and even function such that $\varphi(x) = f(|x|)$.

“ \Leftarrow ”: Conversely, suppose $f \in C_c^\infty([0, r])$ and define $\varphi(x) = f(|x|)$. Then φ is radial and smooth by lemma 1. \square

The above discussion clearly applies to functions $\varphi : \mathbb{R}^d \rightarrow \mathbb{C}$ too.

3. HARDY-TYPE INTEGRAL INEQUALITIES

Inequalities of Hardy-type allow to trade a singular weight for a derivative within L^p -norms.

Lemma 3. *Let $p, r, s \in \mathbb{R}$ with $p \geq 1$, $r > 0$ and $s > -\frac{1}{p}$.*

(1) *The estimate*

$$\int_0^r x^{ps} |f(x)|^p dx \lesssim |f(r)|^p + \int_0^r x^{p(s+1)} |f'(x)|^p dx$$

holds for all $f \in C^1([0, r])$.

(2) *The estimate*

$$\int_0^\infty x^{ps} |f(x)|^p dx \lesssim \int_0^\infty x^{p(s+1)} |f'(x)|^p dx$$

holds for all $f \in C_c^1(\mathbb{R})$.

Proof. If $p > 1$ we have by the product rule

$$x^{ps} |f(x)|^p = \partial_x \left(\frac{x^{ps+1}}{ps+1} |f(x)|^p \right) - \text{Re} \left(x^{(p-1)s} |f(x)|^{p-1} \frac{f(x)}{|f(x)|} \frac{p}{ps+1} x^{s+1} \overline{f'(x)} \right)$$

at each $x \in (0, r)$, which implies with Young's inequality the differential inequality

$$x^{ps} |f(x)|^p \leq \partial_x \left(\frac{p}{ps+1} x^{ps+1} |f(x)|^p \right) + \left(\frac{p}{ps+1} \right)^p x^{p(s+1)} |f'(x)|^p. \quad (3)$$

If $p = 1$ both identities hold in the sense of absolutely continuous functions. This yields integrated over the respective interval the asserted inequalities. \square

The boundary term in lemma 3 is controlled by the following Sobolev inequality.

Lemma 4. *Let $p, r, s \in \mathbb{R}$ with $p \geq 1$, $r > 0$ and $s > -1$. The estimate*

$$|f(r)|^p \lesssim \int_0^r x^{ps} |f(x)|^p dx + \int_0^r x^{p(s+1)} |f'(x)|^p dx$$

holds for all $f \in C^1([0, r])$.

Proof. The fundamental theorem of calculus and Jensen's integral inequality imply

$$\begin{aligned} |f(r)|^p &= \left| \frac{1}{r^{s+1}} \int_0^r \partial_x \left(x^{s+1} f(x) \right) dx \right|^p \\ &\leq \frac{1}{r^{ps+1}} \int_0^r \left| (s+1)x^s f(x) + x^{s+1} f'(x) \right|^p dx \\ &\leq \frac{2^{p-1}(s+1)^p}{r^{ps+1}} \int_0^r x^{ps} |f(x)|^p dx + \frac{2^{p-1}}{r^{ps+1}} \int_0^r x^{p(s+1)} |f'(x)|^p dx \end{aligned}$$

which is the desired inequality. \square

4. PROOF OF THEOREM 1

The key to theorem 1 is to expand partial derivatives of smooth radial functions in terms of the action of the derivation (1) on their radial representative. In view of lemmas 1 and 2 we will see that this derivation provides a natural differential operator for smooth radial functions. Next to multi-index notation, we introduce *d-index notation*, i.e., for $j \in \mathbb{Z}_{\geq 0}$ we call $I = (i_1, \dots, i_j) \in \{1, \dots, d\}^j$ a *d-index of length j* and define

$$(\partial_I f)(x) := \partial_{x^I} f(x) := \partial_{x^{i_1}} \dots \partial_{x^{i_j}} f(x), \quad x = (x^1, \dots, x^d) \in \mathbb{R}^d,$$

for $f \in C^\infty(\mathbb{R}^d)$.

Proof of Theorem 1. “ \lesssim ”: We start noting with a transformation of the integral

$$\|f(|\cdot|)\|_{L^p(\mathbb{B}_r^d)} \simeq \left\| \left(\cdot \right)^{\frac{d-1}{p}} f \right\|_{L^p(0,r)} \quad (4)$$

for all $f \in C_c^\infty([0, r])$. Now, fix $k \in \mathbb{Z}_{\geq 0}$ and let $0 \leq j \leq k$. Let $I = (i_1, \dots, i_j) \in \{1, \dots, d\}^j$ be a *d-index* of length j . By induction, we get

$$\partial_{x^I} f(|x|) = \sum_{n=\lceil \frac{j}{2} \rceil}^j p_I^n(x) (D^n f)(|x|), \quad (5)$$

where

$$p_I^n(x) := p_n^I(x) := \frac{1}{(j-n)!} \Delta_x^{j-n} (x^{i_1} \dots x^{i_j}), \quad n = \left\lfloor \frac{j}{2} \right\rfloor, \dots, j, \quad x \in \mathbb{R}^d, \quad (6)$$

are homogeneous polynomial functions of degree $2n - j$. Here, Δ^{j-n} are $(j-n)$ applications of the Laplace operator Δ . This implies with an application of lemma 3 in combination with lemma 4

$$\begin{aligned} \|\partial_I f(|\cdot|)\|_{L^p(\mathbb{B}_r^d)} &\lesssim \sum_{n=\lceil \frac{j}{2} \rceil}^j \left\| \left(\cdot \right)^{\frac{d-1}{p}+2n-j} D^n f \right\|_{L^p(0,r)} \\ &\lesssim \sum_{n=\lceil \frac{j}{2} \rceil}^{j-1} |(D^n f)(r)| + \left\| \left(\cdot \right)^{\frac{d-1}{p}+j} D^j f \right\|_{L^p(0,r)} \\ &\lesssim \sum_{n=\lceil \frac{j}{2} \rceil}^j \left\| \left(\cdot \right)^{\frac{d-1}{p}+n} D^n f \right\|_{L^p(0,r)} \end{aligned}$$

for all $f \in C_c^\infty([0, r])$. Summation over all multi-indices yields the first bound. On \mathbb{R}^d , one uses item 2 of lemma 3 instead.

“ \gtrsim ”: Upon contracting eq. (5) with $p_m^I(\frac{x}{|x|})$ over all *d-indices* $I \in \{1, \dots, d\}^j$ for each $m = \left\lfloor \frac{j}{2} \right\rfloor, \dots, j$, we infer the system of equations

$$\sum_{I \in \{1, \dots, d\}^j} p_m^I\left(\frac{x}{|x|}\right) \partial_{x^I} f(|x|) = \sum_{n=\lceil \frac{j}{2} \rceil}^j \left(\sum_{I \in \{1, \dots, d\}^j} p_m^I\left(\frac{x}{|x|}\right) p_I^n\left(\frac{x}{|x|}\right) \right) |x|^{2n-j} (D^n f)(|x|). \quad (7)$$

Since the Laplace operator is invariant under rotations $R \in SO(d)$, i.e.,

$$(\Delta \varphi) \circ R = \Delta(\varphi \circ R),$$

we infer from eq. (6)

$$p_n^I(Rx) = \sum_{I' \in \{1, \dots, d\}^j} R^{i_1}_{i'_1} \dots R^{i_j}_{i'_j} p_n^{I'}(x)$$

for all $R = [R_n^m]_{m,n=1}^d \in SO(d)$ and thus the symmetric coefficient matrix

$$g_m^n := \sum_{I \in \{1, \dots, d\}^j} p_{m+\lceil \frac{j}{2} \rceil - 1}^I(\frac{x}{|x|}) p_I^{n+\lceil \frac{j}{2} \rceil - 1}(\frac{x}{|x|}), \quad m, n = 1, \dots, \left\lfloor \frac{j}{2} \right\rfloor + 1, \quad (8)$$

is the Gram matrix of the vectors

$$v^i = \left(p_I^{i+\lceil \frac{j}{2} \rceil - 1}(e_d) \right)_{I \in \{1, \dots, d\}^j} \in \mathbb{R}^{d^j}, \quad i = 1, \dots, \left\lfloor \frac{j}{2} \right\rfloor + 1. \quad (9)$$

The Gram matrix (8) is invertible if and only if the vectors (9) are linearly independent. To prove the latter, let $\lambda_1, \dots, \lambda_{\lfloor \frac{j}{2} \rfloor + 1} \in \mathbb{R}$ such that

$$\sum_{i=1}^{\lfloor \frac{j}{2} \rfloor + 1} \lambda_i v^i = 0.$$

Now we exploit the structural properties of the polynomials given in eq. (6) further. First, consider in the above linear combination the entry I determined by the indices

$$i_\ell = d \quad \text{if } \ell \leq 2 \left\lfloor \frac{j}{2} \right\rfloor - j \quad \text{and} \quad i_\ell = 1 \quad \text{if } \ell > 2 \left\lfloor \frac{j}{2} \right\rfloor - j$$

so that

$$p_I^n(e_d) \neq 0 \quad \text{if } n = \left\lfloor \frac{j}{2} \right\rfloor \quad \text{and} \quad p_I^n(e_d) = 0 \quad \text{if } n > \left\lfloor \frac{j}{2} \right\rfloor.$$

This forces $\lambda_1 = 0$. To proceed, assume that $\lambda_1, \dots, \lambda_{i-1} = 0$ and choose the entry I with indices

$$i_\ell = d \quad \text{if } \ell \leq 2 \left(\left\lfloor \frac{j}{2} \right\rfloor + i - 1 \right) - j \quad \text{and} \quad i_\ell = 1 \quad \text{if } \ell > 2 \left(\left\lfloor \frac{j}{2} \right\rfloor + i - 1 \right) - j$$

so that

$$p_I^n(e_d) \neq 0 \quad \text{if } n = \left\lfloor \frac{j}{2} \right\rfloor + i - 1 \quad \text{and} \quad p_I^n(e_d) = 0 \quad \text{if } n > \left\lfloor \frac{j}{2} \right\rfloor + i - 1.$$

Hence, $\lambda_i = 0$ follows. From this procedure we infer $\lambda_1 = \dots = \lambda_{\lfloor \frac{j}{2} \rfloor + 1} = 0$ and therefore the system (7) can be inverted. This gives with Hölder's inequality the pointwise estimate

$$\begin{aligned} \left| |x|^j (D^j f)(|x|) \right| &\lesssim \sum_{m=\lceil \frac{j}{2} \rceil}^j \sum_{I \in \{1, \dots, d\}^j} \left| p_m^I(\frac{x}{|x|}) \partial_{x^I} f(|x|) \right| \\ &\lesssim \left(\sum_{I \in \{1, \dots, d\}^j} |\partial_{x^I} f(|x|)|^p \right)^{\frac{1}{p}} \end{aligned}$$

for all $f \in C_e^\infty([0, r])$ and all $x \in \overline{\mathbb{B}_r^d}$, which yields the other side of the inequality.

Finally, the map

$$C^\infty([0, r^2]) \rightarrow C_{\text{rad}}^\infty(\overline{\mathbb{B}_r^d}), \quad \tilde{f} \mapsto \tilde{f}(|\cdot|^2),$$

is bijective with inverse

$$C_{\text{rad}}^\infty(\overline{\mathbb{B}_r^d}) \rightarrow C^\infty([0, r^2]), \quad \varphi \mapsto \tilde{f}_\varphi, \quad \text{where} \quad \varphi = \tilde{f}_\varphi(|\cdot|^2)$$

according to lemmas 1 and 2. Then, if f_φ is the radial representative from lemma 2,

$$(D^j f_\varphi)(\rho) = 2^j \tilde{f}_\varphi^{(j)}(\rho^2)$$

and the above implies the equivalence of norms

$$\|\varphi\|_{W^{k,p}(\mathbb{B}_r^d)} \simeq \sum_{j=0}^k \left\| (\cdot)^{\frac{d-2}{2p} + \frac{j}{2}} \tilde{f}_\varphi^{(j)} \right\|_{L^p(0, r^2)}$$

for all $\varphi \in C_{\text{rad}}^\infty(\overline{\mathbb{B}_r^d})$. Thus the bijection extends to an isomorphism of normed linear spaces. The proof on \mathbb{R}^d is similar. \square

Acknowledgment. The author thanks Roland Donn timer and Irfan Glogić for helpful discussions and useful comments on a first draft of this paper.

REFERENCES

- [1] Robert A. Adams and John J. F. Fournier. *Sobolev spaces*, volume 140 of *Pure and Applied Mathematics (Amsterdam)*. Elsevier/Academic Press, Amsterdam, second edition, 2003.
- [2] Roc Alabern, Joan Mateu, and Joan Verdera. A new characterization of Sobolev spaces on \mathbb{R}^n . *Math. Ann.*, 354(2):589–626, 2012.
- [3] H. Berestycki and P.-L. Lions. Nonlinear scalar field equations. I. Existence of a ground state. *Arch. Rational Mech. Anal.*, 82(4):313–345, 1983.
- [4] Paweł Biernat, Roland Donn timer, and Birgit Schörkhuber. Hyperboloidal similarity coordinates and a globally stable blowup profile for supercritical wave maps. *Int. Math. Res. Not. IMRN*, 2021(21):16530–16591, 2021.
- [5] Roland Donn timer and Matthias Ostermann. A globally stable self-similar blowup profile in energy supercritical yang-mills theory. *arXiv preprint arXiv:2108.13668*, 2021.
- [6] Roland Donn timer and Birgit Schörkhuber. Stable blowup for wave equations in odd space dimensions. *Ann. Inst. H. Poincaré C Anal. Non Linéaire*, 34(5):1181–1213, 2017.
- [7] Lawrence C. Evans. *Partial differential equations*, volume 19 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, second edition, 2010.
- [8] Irfan Glogić. Globally stable blowup profile for supercritical wave maps in all dimensions. *arXiv preprint arXiv:2207.06952*, 2022.
- [9] Irfan Glogić. Stable blowup for the supercritical hyperbolic Yang-Mills equations. *Adv. Math.*, 408:Paper No. 108633, 2022.
- [10] Djairo Guedes de Figueiredo, Ederson Moreira dos Santos, and Olímpio Hiroshi Miyagaki. Sobolev spaces of symmetric functions and applications. *J. Funct. Anal.*, 261(12):3735–3770, 2011.
- [11] Roberta Musina. Weighted Sobolev spaces of radially symmetric functions. *Ann. Mat. Pura Appl. (4)*, 193(6):1629–1659, 2014.
- [12] Wei Ming Ni. A nonlinear Dirichlet problem on the unit ball and its applications. *Indiana Univ. Math. J.*, 31(6):801–807, 1982.
- [13] Winfried Sickel and Leszek Skrzypczak. On the interplay of regularity and decay in case of radial functions II. Homogeneous spaces. *J. Fourier Anal. Appl.*, 18(3):548–582, 2012.
- [14] Winfried Sickel, Leszek Skrzypczak, and Jan Vybíral. On the interplay of regularity and decay in case of radial functions I. Inhomogeneous spaces. *Commun. Contemp. Math.*, 14(1):1250005, 60, 2012.
- [15] Winfried Sickel, Leszek Skrzypczak, and Jan Vybíral. The characterization of radial subspaces of Besov and Lizorkin-Triebel spaces by differences. In *Function spaces X*, volume 102 of *Banach Center Publ.*, pages 197–214. Polish Acad. Sci. Inst. Math., Warsaw, 2014.
- [16] Walter A. Strauss. Existence of solitary waves in higher dimensions. *Comm. Math. Phys.*, 55(2):149–162, 1977.
- [17] Hassler Whitney. Differentiable even functions. *Duke Math. J.*, 10:159–160, 1943.

UNIVERSITY OF VIENNA, FACULTY OF MATHEMATICS, OSKAR-MORGENSTERN-PLATZ 1, 1090 VIENNA, AUSTRIA

Email address: `matthias.ostermann@univie.ac.at`