

Statistical mechanics of deep learning beyond the infinite-width limit

S. Ariosto,^{1,2} R. Pacelli,³ M. Pastore,⁴ F. Ginelli,^{1,2} M. Gherardi,^{5,2} and P. Rotondo^{5,2}

¹*Dipartimento di Scienza e Alta Tecnologia and Center for Nonlinear and Complex Systems, Università degli Studi dell’Insubria, Via Valleggio 11, 22100 Como, Italy*

²*I.N.F.N. Sezione di Milano, Via Celoria 16, 20133 Milano, Italy*

³*Dipartimento di Scienza Applicata e Tecnologia, Politecnico di Torino, 10129 Torino, Italy*

⁴*Université Paris-Saclay, CNRS, LPTMS, 91405 Orsay, France*

⁵*Università degli Studi di Milano, Via Celoria 16, 20133 Milano, Italy*

Decades-long literature testifies to the success of statistical mechanics at clarifying fundamental aspects of deep learning. Yet the ultimate goal remains elusive: we lack a complete theoretical framework to predict practically relevant scores, such as the train and test accuracy, from knowledge of the training data. Huge simplifications arise in the infinite-width limit, where the number of units N_ℓ in each hidden layer far exceeds the number P of training examples. This idealisation, however, blatantly departs from the reality of deep learning practice, where training sets are larger than the widths of the networks. Here, we show one way to overcome these limitations. The partition function for fully-connected architectures, which encodes information about the trained models, can be computed analytically with the toolset of statistical mechanics. The computation holds in the “thermodynamic limit” where both N_ℓ and P are large and their ratio $\alpha_\ell = P/N_\ell$, which vanishes in the infinite-width limit, is now finite and generic. This advance allows us to obtain (i) a closed formula for the generalisation error associated to a regression task in a one-hidden layer network with finite α_ℓ ; (ii) an expression of the partition function (technically, via an “effective action”) for fully-connected architectures with arbitrary number of hidden layers, in terms of a finite number of degrees of freedom (technically, “order parameters”); (iii) a demonstration that the Gaussian processes arising in the infinite-width limit should be replaced by Student-t processes; (iv) a simple analytical criterion to predict, for a given training set, whether finite-width networks (with ReLU activations) achieve better test accuracy than infinite-width ones. As exemplified by these results, our theory provides a starting point to tackle the problem of generalisation in realistic regimes of deep learning.

I. INTRODUCTION

The rise of deep learning, made possible by the progress in computing technology and eventually triggered by the astonishing successes in computer vision [1], was foreshadowed by decades of research in the second half of the 20th century [2], notably in the field of statistical physics [3]. Yet, the sheer rate of technological progress — state-of-the-art deep neural networks (DNN) often achieve super-human performance — has outpaced our ability to put it on firm theoretical ground. For instance, a convincing explanation of why these architectures are able to generalise in the overparametrised regime, where the number of learnable parameters far exceeds the size of the training set, is yet to be found [4, 5].

Two of the ingredients at the basis of the success of DNN models — their many-layers structure and the fact that they implement a non-linear transformation of the inputs — provide also the main technical challenges for theorists. Indeed, disentangling these two ingredients stimulated some of the most recent progress in the field: the observation that in the infinite-width limit deep models are equivalent to Gaussian processes [6–11] and the study of their lazy training regime under gradient descent [12–14] made it possible to establish a connection between deep learning and kernel methods. Such a breakthrough granted a description of the generalisation performance of DNNs in the infinite-width limit [15, 16], by means of statistical physics techniques originally pio-

neered in Ref. [17] for the study of support vector machines [18].

Despite these advances, there is agreement that a more complete theory should address deep learning beyond the infinite-width limit [19, 20]. Indeed, realistic neural networks operate in a qualitatively different regime, where the number of training examples exceeds the width of the largest layer. Modeling the finite-width regime in the thermodynamic limit, where the number of degrees of freedom diverges and the tools of statistical mechanics are most effective, amounts to taking the asymptotic limit where both the size of the training set P and the number of units in each hidden layer N_ℓ ($\ell = 1, \dots, L$ with L being the depth of the network) tend to infinity with their ratios $\alpha_\ell = P/N_\ell$ fixed. It is worth stressing that from this choice it follows that such a network must work in the overparametrised regime.

Another fruitful line of research, in the direction of overcoming the limitations of the infinite-width limit, sacrifices the non-linear nature of the network by considering a deep *linear* input-output mapping: even if the resulting architecture lacks the expressive power [21–28] of the same model with non-linearities, the multi-layer structure maintains the learning problem non-convex, while amenable to analytical investigation [29, 30].

Very recently, Li and Sompolinsky [31] proposed a method to analytically evaluate properties of deep finite-width linear networks (e.g., their generalisation error) trained on a given training set. The method, called back-

propagating kernel renormalisation (BPKR), is based on a statistical physics approach: the posterior distribution of the network weights is evaluated exactly via successive integration of the weights in each layer (starting from the output layer and proceeding backwards through the network). Interestingly, the authors of Ref. [31] observe that a heuristic extension of the BPKR to the non-linear case with rectified linear units (ReLU) fits well the results from numerical simulations in a wide range of parameters. However, the more relevant case of generic non-linear DNNs remains an open problem, despite some recent notable attempts to address it [32–35].

In statistical mechanics, the partition function is the central object encoding the properties of the system in the thermodynamic limit. In this work we address the analytical computation of the partition function of a fully-connected, multi-layer, finite-width, non-linear neural network, as a function of the training set. Technically, the computation amounts to integrating out an extensive number of degrees of freedom (the weights of the network), thus landing on an expression that involves only a finite number (proportional to the depth L) of integrals over a set of quantities, termed order parameters in statistical physics, to be evaluated by the saddle point method. The derivation is exact in the case of one-hidden-layer (1HL, that is $L = 1$) networks under mild hypotheses on the correlation matrix of the inputs and on the form of the non-linearity. It relies on the Breuer-Major (BM) theorem [36, 37], a generalised central limit theorem (CLT).

In the general case of an architecture with L hidden fully-connected layers, we show that the distribution of the pre-activations at each layer ℓ is a mixture of Gaussians that depends on ℓ parameters. Notably, the back-propagating integration performed in [31] is not a viable option as soon as non-linearities are added to the model. We introduce a forward-propagating method to carry out nested integrations starting from the input layer. The result depends on an assumption that is similar, at least in spirit, to the Gaussian equivalence principle employed for random and generic feature models [38–44].

From these developments, we are able to obtain quantitative predictions for observables such as the generalisation error of the network, in the overparametrised regime where perfect training can be achieved, i.e., below the interpolation threshold. Our results have an intriguing interpretation from the point of view of stochastic processes: we show that in the case of finite α_ℓ the Gaussian process arising in the infinite-width limit of Bayesian neural networks [9] should be generalised to a Student’s t stochastic process [45]. Moreover, we show that there is a crucial difference in choosing odd or non-odd activation functions and that the theory is more complex in the latter case. As a first application of the theory in the generic non-odd case, we establish a simple criterion to predict whether it is convenient, in terms of generalisation performance, to employ a finite-width deep neural network with ReLU non-linearity over its infinite-width

version.

II. SETTING OF THE LEARNING PROBLEM AND NOTATION

In the following we consider deep neural networks with L fully-connected hidden layers, where the pre-activations of each layer $h_{i_\ell}^{(\ell)}$ ($i_\ell = 1, \dots, N_\ell$; $\ell = 1, \dots, L$) are given recursively as a non-linear function of the pre-activations at the previous layer $h_{i_{\ell-1}}^{(\ell-1)}$ ($i_{\ell-1} = 1, \dots, N_{\ell-1}$):

$$h_{i_\ell}^{(\ell)} = \frac{1}{\sqrt{N_{\ell-1}}} \sum_{i_{\ell-1}=1}^{N_{\ell-1}} W_{i_\ell i_{\ell-1}}^{(\ell)} \sigma\left(h_{i_{\ell-1}}^{(\ell-1)}\right) + b_{i_\ell}^{(\ell)}, \quad (1)$$

$$h_{i_1}^{(1)} = \frac{1}{\sqrt{N_0}} \sum_{i_0=1}^{N_0} W_{i_1 i_0}^{(1)} x_{i_0} + b_{i_1}^{(1)} \quad (2)$$

where $W^{(\ell)}$ and $b^{(\ell)}$ are respectively the weights and the biases of the ℓ -th layer, whereas the input layer has dimension N_0 (the input data dimension). σ is a non-linear activation function and it is common to each layer. Since we are interested in regression problems we add one last readout layer and we define the function implemented by the deep neural network as:

$$f_{\text{DNN}}(\mathbf{x}) = \frac{1}{\sqrt{N_L}} \sum_{i_L=1}^{N_L} v_{i_L} \sigma\left[h_{i_L}^{(L)}(\mathbf{x})\right], \quad (3)$$

where \mathbf{v} is the vector of weights of the last layer.

We consider a supervised learning problem with training set $\mathcal{T}_P = \{\mathbf{x}^\mu, y^\mu\}_{\mu=1}^P$, where each $\mathbf{x}^\mu \in \mathbb{R}^D$ and the corresponding labels $y^\mu \in \mathbb{R}$. Without loss of generality, we can assume that the training set is drawn from an input/output probability distribution $P(\mathbf{x}, y)$ and that its probability density $\mu(\mathbf{x}, y)$ does exist.

In this manuscript we restrict our analysis to quadratic loss functions:

$$\mathcal{L} = \frac{1}{2} \sum_{\mu=1}^P [y^\mu - f_{\text{DNN}}(\mathbf{x}^\mu)]^2 + \mathcal{L}_{\text{reg}}, \quad (4)$$

$$\mathcal{L}_{\text{reg}} = \frac{\lambda}{2\beta} \sum_{i_L=1}^{N_L} v_{i_L}^2 + \frac{1}{2\beta} \sum_{\ell=1}^L \lambda^{(\ell)} \|W^{(\ell)}\|^2, \quad (5)$$

where L^2 regularisations have been added for each layer to the bare loss function, $\|\cdot\|$ is the standard Frobenius norm defined for matrices and $\beta = 1/T$ is the inverse temperature parameter, at this level just a scaling factor for \mathcal{L}_{reg} .

As a standard practice in statistical mechanics of deep learning, we define the partition function of the problem as:

$$Z = \int \mathcal{D}\theta e^{-\beta \mathcal{L}(\theta)}. \quad (6)$$

where the symbol $\int \mathcal{D}\theta$ indicates the collective integration over the weights of the network, $\theta = \{\{W^{(\ell)}\}, \{b^{(\ell)}\}, v\}$. This choice enforces minimization of the training error for $\beta \rightarrow \infty$. Since we have explicitly included regularization terms into the loss function, we do not need to regularize the measure $\mathcal{D}\theta$, now biased toward weights with small norms. We notice however that our choice of scaling \mathcal{L}_{reg} by $1/\beta$ has a natural Bayesian learning interpretation: the Gibbs probability $P_\beta(\theta) = Z^{-1} e^{-\beta \mathcal{L}(\theta)}$ associated with the partition function in Eq. (6) is the posterior distribution of the weights after training, whereas the Gaussian regularization is a prior equivalent to assuming that weights at initialization have been drawn from a Gaussian distribution (that is often the case in practical applications) with variances equal to $1/\lambda^{(\ell)}$ (hidden layers) and $1/\lambda$ (readout layer). In our work we choose to adopt this scaling convention, which is the same employed in Ref. [31].

Computing the average training error at a given inverse temperature β simply amounts to evaluating:

$$\langle \epsilon_t \rangle = \frac{1}{P} \int \mathcal{D}\theta [\mathcal{L}(\theta) - \mathcal{L}_{\text{reg}}(\theta)] \frac{e^{-\beta \mathcal{L}(\theta)}}{Z}, \quad (7)$$

whereas the average test error over a new (unseen) example (\mathbf{x}^0, y^0) is given by:

$$\langle \epsilon_g(\mathbf{x}^0, y^0) \rangle = \frac{1}{2} \int \mathcal{D}\theta [y^0 - f_{\text{DNN}}(\mathbf{x}^0)]^2 \frac{e^{-\beta \mathcal{L}(\theta)}}{Z}. \quad (8)$$

Training and test errors represent two special observables, but more in general, for an arbitrary observable O we have:

$$\langle O \rangle = \int \mathcal{D}\theta O(\theta) \frac{e^{-\beta \mathcal{L}(\theta)}}{Z}. \quad (9)$$

In the next sections, we show how to reduce the computation of the partition function in Eq. (6) from an integral over an extensive number of variables to a simple saddle-point integral over a finite number of order parameters. We start from the simple case of one hidden layer fully-connected architectures and we later generalise the result to arbitrary deep architectures with L hidden layers.

III. EXACT PARTITION FUNCTION FOR ONE HIDDEN LAYER NEURAL NETWORKS IN THE ASYMPTOTIC LIMIT

In the case of one hidden layer architectures, we are able to show that the final result is exact, since the key approximation that allows to reduce the complexity of the computation is justified by a generalised central limit theorem due to Breuer and Major [36]. For this reason, we briefly state this theorem without proof. A modern approach to this topic can be found in the recent work by Nourdin, Peccati and Podolsky [37].

A. The Breuer-Major theorem

The Breuer-Major theorem deals with the following sequence of random variables:

$$S_N = \frac{1}{\sqrt{N}} \sum_{i=1}^N c_i F(x_i) \quad N \geq 1. \quad (10)$$

Clearly, if the distribution of the vector $\mathbf{x} = (x_1, \dots, x_N)$ is factorized over its coordinates, i.e. $p(\mathbf{x}) = \prod_i p(x_i)$ and $F(x) = x$, the random variable $S = \lim_{N \rightarrow \infty} S_N$ is normal distributed as long as the mean $\mathbb{E}(x_i) = 0$, the variance $\mathbb{E}(x_i^2)$ is finite and the c_i 's satisfy the so-called Lindeberg's condition. This is also true whenever F is a well-behaved non-linearity.

The Breuer-Major theorem essentially extends this result to generic gaussian processes (GPs), providing sufficient conditions on the covariance matrix of the GP and on the non-linearity F that guarantee convergence of S_N to the normal distribution. We report here the modern statement of the theorem given in Ref. [37].

We consider a stationary (unidimensional) GP $x = (x_k)_{k \in \mathbb{Z}}$. Stationarity –which is not essential and will be replaced by a weaker condition in the following– amounts to require that the covariance of the process $C_{ij} = \mathbb{E}(x_i x_j)$ is a function of the difference $i - j$, i.e. $C_{ij} = C(i - j)$. The only technical condition to be imposed on the non-linear function F is to have well-defined Hermite rank R . The Hermite rank is the smallest positive integer that appears in the decomposition of F over the Hermite polynomials:

$$F(x) = \sum_{k=R}^{\infty} f_k \text{He}_k(x), \quad (11)$$

where $\text{He}_k(x)$ is the k -th Hermite polynomial and f_k the coefficient of the expansion. For many reasonable activation functions F , $R = 1$.

Theorem 1 (Breuer and Major, 1983) *Let $x = (x_k)_{k \in \mathbb{Z}}$ be a stationary unidimensional gaussian process with covariance $C(i - j)$. Let $\mathbb{E}[F(x_1)] = 0$ and $\mathbb{E}[F^2(x_1)] < \infty$ and assume that the function F has Hermite rank $R \geq 1$. Suppose that:*

$$\sum_{j \in \mathbb{Z}} |C_{1j}|^R < \infty. \quad (12)$$

Then $\sigma^2 := \mathbb{E}[F(x_1)^2] + 2 \sum_{j=1}^{\infty} \mathbb{E}[F(x_1)F(x_j)]$ is finite. Moreover, one has that the sequence of random variables

$$S_N = \frac{1}{\sqrt{N}} \sum_{i=1}^N F(x_i) \quad N \geq 1 \quad (13)$$

converges in distribution to $\mathcal{N}(0, \sigma^2)$, i.e. to a Gaussian distribution with zero mean and variance σ^2 .

It is worth stressing that for our scopes we will need a slightly stronger statement than the one just mentioned: in fact, in our calculation the covariance will not be stationary. However, it has been shown, already in the original reference [36], that the hypothesis of stationarity can be weakened and replaced with a requirement of uniform convergence of the elements of the covariance, namely:

$$\sum_{j \in \mathbb{Z}} |C_{ij}|^R < B_0 \quad \forall i \in \mathbb{Z}, \quad (14)$$

where B_0 is a positive finite constant. As a matter of fact, we will also need to consider a more general sequence of random variables, where each term of the sum is weighted by a factor $c_i \neq 1$. For standard CLTs, this does not spoil the convergence to the normal distribution as long as the c_i 's satisfy the so-called Lindeberg's condition. Here we assume that the same occurs in the Breuer-Major setting.

B. Sketch of the calculation

We now discuss the salient aspects of the calculation. The starting point is the following partition function:

$$Z = \int \prod_{i_1}^{N_1} dv_{i_1} \prod_{i_1, i_0}^{N_1, N_0} dw_{i_1 i_0} \exp \left\{ -\frac{\lambda}{2} \sum_{i_1}^{N_1} v_{i_1}^2 - \frac{\lambda_1}{2} \|w\|^2 - \frac{\beta}{2} \sum_{\mu} \left[y^{\mu} - \frac{1}{\sqrt{N_1}} \sum_{i_1}^{N_1} v_{i_1} \sigma \left(\sum_{i_0}^{N_0} \frac{w_{i_1, i_0} x_{i_0}^{\mu}}{\sqrt{N_0}} \right) \right]^2 \right\}. \quad (15)$$

where $w = W^{(1)}$ and we took $b^{(1)} = 0$ without loss generality¹.

Notice once again that we have made the choice of including the regularization terms in the probability measure over the weights. The first step is to decouple the weights of the different layers in the loss function. This can be done including standard identities built over two families of Dirac deltas, one for the pre-activations of the hidden layer and one for the output of the network:

$$1 = \int \prod_{\mu}^P ds^{\mu} \delta \left[s^{\mu} - \frac{1}{\sqrt{N_1}} \sum_{i_1}^{N_1} v_{i_1} \sigma(h_{i_1}^{\mu}) \right], \quad (16)$$

$$1 = \int \prod_{\mu}^P \prod_{i_1}^{N_1} dh_{i_1}^{\mu} \delta \left(h_{i_1}^{\mu} - \frac{1}{\sqrt{N_0}} \sum_{i_0}^{N_0} w_{i_1 i_0} x_{i_0}^{\mu} \right). \quad (17)$$

By using a standard Fourier representation of these deltas, which introduces the conjugate variables $\bar{h}_{i_1}^{\mu}$ and

\bar{s}^{μ} , we can separate the integrals on internal weights from those on external weights and obtain:

$$Z = \int \prod_{\mu}^P \frac{ds^{\mu} d\bar{s}^{\mu}}{2\pi} e^{-\frac{\beta}{2} \sum_{\mu} (y^{\mu} - s^{\mu})^2 + i \sum_{\mu} s^{\mu} \bar{s}^{\mu}} \times \int \prod_{\mu, i_1}^{P, N_1} \frac{dh_{i_1}^{\mu} d\bar{h}_{i_1}^{\mu}}{2\pi} e^{i \sum_{i_1}^{N_1} \sum_{\mu} h_{i_1}^{\mu} \bar{h}_{i_1}^{\mu}} \times \int \prod_{i_1, i_0}^{N_1, N_0} dw_{i_1 i_0} e^{-\frac{\lambda_1}{2} \|w\|^2 - i \sum_{i_1}^{N_1} \bar{h}_{i_1}^{\mu} \sum_{i_0}^{N_0} w_{i_1 i_0} x_{i_0}^{\mu} / \sqrt{N_0}} \times \int \prod_{i_1}^{N_1} dv_{i_1} e^{-\frac{\lambda}{2} \sum_{i_1}^{N_1} v_{i_1}^2 - i \sum_{\mu} \bar{s}^{\mu} \sum_{i_1}^{N_1} v_{i_1} \sigma(h_{i_1}^{\mu}) / \sqrt{N_1}}. \quad (18)$$

The integrals on the weights are Gaussian, and solving them yields quadratic terms in the conjugate variables

$$Z = \int \prod_{\mu}^P \frac{ds^{\mu} d\bar{s}^{\mu}}{2\pi} e^{-\frac{\beta}{2} \sum_{\mu} (y^{\mu} - s^{\mu})^2 + i \sum_{\mu} s^{\mu} \bar{s}^{\mu}} \times \left\{ \int \prod_{\mu}^P \frac{dh^{\mu} d\bar{h}^{\mu}}{2\pi} e^{i \sum_{\mu} h^{\mu} \bar{h}^{\mu} - \frac{1}{2\lambda N_1} [\sum_{\mu} \bar{s}^{\mu} \sigma(h^{\mu})]^2} \times e^{-\frac{1}{2\lambda_1 N_0} \sum_{i_0}^{N_0} (\sum_{\mu} \bar{h}^{\mu} x_{i_0}^{\mu})^2} \right\}^{N_1}, \quad (19)$$

where we used the fact that the integrals on $h_{i_1}^{\mu}$ and $\bar{h}_{i_1}^{\mu}$ can be factorized on the index i_1 . The integral over the \bar{h}^{μ} is Gaussian and can be solved:

$$\int \prod_{\mu}^P \frac{d\bar{h}^{\mu}}{2\pi} e^{i \sum_{\mu} h^{\mu} \bar{h}^{\mu} - \frac{1}{2\lambda_1 N_0} \sum_{i_0}^{N_0} (\sum_{\mu} \bar{h}^{\mu} x_{i_0}^{\mu})^2} = P_1(\{h^{\mu}\}), \quad (20)$$

where

$$P_1(\{h^{\mu}\}) = \frac{e^{-\frac{1}{2} \sum_{\mu, \nu} h^{\mu} C_{\mu, \nu}^{-1} h^{\nu}}}{\sqrt{(2\pi)^P \det C}}, \quad C_{\mu\nu} = \frac{1}{\lambda_1 N_0} \sum_{i_0}^{N_0} x_{i_0}^{\mu} x_{i_0}^{\nu}. \quad (21)$$

We notice that this last step requires the covariance matrix C to be invertible. While this is false whenever $P > N_0$, we can easily cope with this issue by adding a small diagonal term to the covariance matrix. One can explicitly check that the final result does not depend on the choice of this extra regularization.

To deal with the integral over h^{μ} we can include a further Dirac delta identity for the random variable $q = 1/\sqrt{\lambda N_1} \sum_{\mu} \bar{s}^{\mu} \sigma(h^{\mu})$. This leaves us with the problem of finding the probability density $P(q)$. In the limit $P, N_1 \rightarrow \infty$ and $\alpha_1 = P/N_1$ fixed, this is exactly the same setting of the Breuer-Major theorem. As such, it is sufficient that both the (regularized) covariance C and the activation function σ satisfy the hypothesis of the theorem to guarantee that the probability distribution

¹ One can map a system with non-zero biases in a zero-bias one increasing by one the dimensions of the input and of the activations at each layer. The original biases are then trivially mapped in the extra weights of the augmented system.

$P(q)$ converges in distribution to a Gaussian:

$$P(q) = \int d^P h P_1(\{h^\mu\}) \delta \left[q - \frac{1}{\sqrt{\lambda N_1}} \sum_{\mu} \bar{s}^\mu \sigma(h^\mu) \right],$$

$$P(q) \rightarrow \mathcal{N}_q(0, Q). \quad (22)$$

with variance

$$Q(\bar{s}, C) = \frac{1}{\lambda N_1} \sum_{\mu, \nu} \bar{s}^\mu \left[\int d^P h P_1(\{h^\rho\}) \sigma(h^\mu) \sigma(h^\nu) \right] \bar{s}^\nu$$

$$= \frac{1}{\lambda N_1} \sum_{\mu, \nu} \bar{s}^\mu K_{\mu\nu}(C) \bar{s}^\nu. \quad (23)$$

Here we have assumed that the variable q has zero mean, a condition verified as long as

$$\int d^P h P_1(\{h^\mu\}) \sigma(h^\nu) = 0, \quad (24)$$

that is for σ odd; for a more general derivation, relevant for non-odd activation function such as ReLU, see Sec. V.

Each element of the kernel matrix $K_{\mu\nu}(C)$ can be easily reduced from a P -dimensional integral to a simpler two-dimensional one:

$$K_{\mu\nu}(C) = \int \frac{dt_1 dt_2}{\sqrt{(2\pi)^2 \det \tilde{C}}} e^{-\frac{1}{2} \mathbf{t}^T \tilde{C}^{-1} \mathbf{t}} \sigma(t_1) \sigma(t_2), \quad (25)$$

where $\mathbf{t} = (t_1, t_2)^T$ and

$$\tilde{C} = \begin{pmatrix} C_{\mu\mu} & C_{\mu\nu} \\ C_{\mu\nu} & C_{\nu\nu} \end{pmatrix}. \quad (26)$$

is the reduced 2×2 input covariance matrix. It is worth noticing that the kernel we find here is the so-called *neural network gaussian process* (NNGP) kernel. It differs from the neural tangent kernel (NTK) that is found in the infinite-width limit of networks trained under gradient descent [46]. The fact that the infinite-width limit of a Bayesian neural network differs from the one obtained from gradient descent is indeed known and discussed in literature [14].

Now we can integrate over the variable q and obtain:

$$\left[\int \frac{dq e^{-\frac{q^2}{2} - \frac{q^2}{2Q(\bar{s}, C)}}}{\sqrt{2\pi Q(\bar{s}, C)}} \right]^{\frac{N_1}{2}} = [Q(\bar{s}, C) + 1]^{-\frac{N_1}{2}}. \quad (27)$$

By direct inspection of the definition of Q , Eq. (23), it is straightforward to notice that as long as $N_1 \rightarrow \infty$ and $N_1 \gg P$, the dependence on N_1 disappears and we get:

$$[Q(\bar{s}, C) + 1]^{-\frac{N_1}{2}} \rightarrow e^{-\frac{1}{2\lambda} \sum_{\mu, \nu} \bar{s}^\mu K_{\mu\nu}(C) \bar{s}^\nu}. \quad (28)$$

Eq. (27) has a very natural interpretation in view of the NNGP literature. Indeed, the \bar{s}^μ appearing there are

the conjugate variables of the s^μ appearing in the loss function through their definition (16). This means that Eq. (27) is a statement about the probability distribution of

$$s^\mu = \frac{1}{\sqrt{N_1}} \sum_i^{N_1} v_{i1} \sigma(h_{i1}^\mu), \quad (29)$$

to be used to average the Gibbs weight in the partition function. For N_1 large and P finite, these variables are jointly multivariate Gaussian distributed according to the central limit theorem, as noted for example in [9]: this limit corresponds indeed to our Eq. (28) and is the cornerstone of the mapping of a infinite-width Bayesian neural network to a Gaussian process. This is however no more the case when P is comparable to N_1 : Eq. (27), derived under the hypothesis of the Breuer-Major theorem, means that the variables \bar{s}^μ are distributed according to a multivariate Student's t -distribution [45, 47–49],

$$p(\{\bar{s}^\mu\}) \sim \left(1 + \frac{1}{\lambda N_1} \sum_{\mu, \nu} \bar{s}^\mu K_{\mu\nu}(C) \bar{s}^\nu \right)^{-\frac{N_1}{2}}. \quad (30)$$

The need of considering Student's t processes as a generalisation of NNGPs has been noted already in the case of different priors on the distribution of the last layer's weights [50]. The reason why this kind of process arises in the case we are considering here can be understood with an heuristic argument: when N_1 and P are of the same order, we cannot take the limit $N_1 \rightarrow \infty$ before $P \rightarrow \infty$, and so we need to use the empirical covariance of the variables (29) instead of their true one in estimating their probability distribution. A more precise characterization of these neural network Student's t processes (NNTPs) and the regime where they arise represent interesting topics for future work.

C. Effective saddle-point action

In the general case of finite $\alpha_1 = P/N_1$, we are only left with the integrals in s^μ and \bar{s}^μ . To solve them it is convenient to introduce one final Dirac delta identity:

$$1 = \int dQ \delta \left[Q - \frac{1}{\lambda N_1} \sum_{\mu, \nu} \bar{s}^\mu K(C)_{\mu\nu} \bar{s}^\nu \right], \quad (31)$$

where $Q \geq -1$ is now an integration variable and not a function of \bar{s} , so that we have removed the explicit dependence on $\sqrt{Q(\bar{s}, C) + 1}$ in the partition function. Finally, the integrals in s^μ and \bar{s}^μ are Gaussian once another integral representation of the delta via a conjugate variable \bar{Q} is inserted. This allows us to get a final effective action of solely two variables:

$$Z = \int dQ d\bar{Q} \exp \left[-\frac{N_1}{2} S(Q, \bar{Q}) \right] \quad (32)$$

where the action S is given by:

$$S = Q\bar{Q} + \log(1 + Q) + \frac{1}{N_1} \text{Tr} \log \beta \left[\frac{\mathbb{1}}{\beta} - \frac{\bar{Q}K(C)}{\lambda} \right] + \frac{1}{N_1} y^\top \left[\frac{\mathbb{1}}{\beta} - \frac{\bar{Q}K(C)}{\lambda} \right]^{-1} y \quad (33)$$

and we have introduced a vectorial notation for the output $y^\top = (y^1, y^2, \dots, y^P)$. This is the first main result of our work: in the thermodynamic limit $N_1, P \rightarrow \infty$ with $\alpha_1 = P/N_1$ fixed, we have been able to show that the computation of the partition function for one hidden layer architectures can be reduced to a two-variables saddle-point integral. The saddle-point equations $\partial_Q S = 0$, $\partial_{\bar{Q}} S = 0$ read, respectively,

$$\bar{Q} = -\frac{1}{1 + Q}, \quad (34)$$

$$Q = \frac{1}{N_1} \text{Tr} \left[\left(\frac{\mathbb{1}}{\beta} - \frac{\bar{Q}K}{\lambda} \right)^{-1} \frac{K}{\lambda} \right] - \frac{1}{N_1} y^\top \left(\frac{\mathbb{1}}{\beta} - \frac{\bar{Q}K}{\lambda} \right)^{-2} \frac{K}{\lambda} y. \quad (35)$$

It is easy to show that $Q^* = 0$ and $\bar{Q}^* = -1$ is the only solution at any temperature in the infinite-width limit $N_1 \gg P$, since the last two term in the action (33) are negligible.

D. Validity of the derivation in the heavily overparametrised limit

For the moment, we have explicitly stated that our derivation holds in the asymptotic limit $P, N_1 \rightarrow \infty$ with $\alpha_1 = P/N_1$ finite, but we have not discussed any additional constraint that may come by the request that the (regularized) covariance matrix C satisfies the hypothesis of the Breuer-Major theorem. The only stringent condition to verify is Eq. (14), that is

$$\sum_{\mu=1}^P |C_{\mu\nu}|^R < B_0 \quad \forall \nu = 1, \dots, P, \quad (36)$$

where B_0 is a given finite constant and R the Hermite rank of the activation function σ . In the case of inputs \mathbf{x} with i.i.d. standard Gaussian coordinates, $C_{\mu\nu}$ is a Wishart random matrix with off-diagonal entries of order $1/\sqrt{N_0}$ and random signs: after taking the absolute value, the sum in Eq (36) is of order $P(N_0)^{-R/2}$. For $R = 1$, this scaling is suggesting that we need to work at finite ratio $P/\sqrt{N_0}$ to comply with the hypothesis of the theorem. For $R \geq 2$ instead we can safely work at finite $\alpha_0 = P/N_0$. Numerical evidence presented in the following seems to suggest that our derivation, and in particular Eq. (22), are valid for values of P much larger also in the case of activation functions with Hermite rank

$R = 1$. A mathematical formalization of this observation is beyond the scope of the present paper.

Here we limit ourselves to notice that as long as $\alpha_0 = O(1)$, the non-linear kernel K has only strictly positive eigenvalues (this can be checked explicitly in those cases where the analytical kernel is known [38]). As such, we expect that the average training error at zero temperature will be exactly zero, i.e. we are in the regime where the network is capable to perfectly fit the training data.

E. Average training error and predictors statistics

We have found an explicit formula for the action that can be used to find the averages of the observables. In particular we are interested in the training and generalisation errors. The first can be easily found performing the derivative over β of the logarithm of the partition function. By doing so, from Eq. (7) we obtain:

$$\langle \epsilon_t \rangle = \frac{1}{2P} y^\top \left[\mathbb{1} - \frac{\beta \bar{Q}^* K(C)}{\lambda} \right]^{-2} y - \frac{1}{2P} \text{Tr} \left\{ \left[\mathbb{1} - \frac{\beta \bar{Q}^* K(C)}{\lambda} \right]^{-1} \frac{\bar{Q}^* K}{\lambda} \right\}. \quad (37)$$

where \bar{Q}^* denotes the solution of the saddle point equation (35).

The second observable we are interested in is the generalisation error (8). We can proceed along the same lines of the calculation performed in Sec. III B introducing, other than the variables s^μ, h_i^μ defined by (16), (17), additional variables s^0, h_i^0 that describe output and pre-activations of the new test example. We thus get:

$$\langle \epsilon_g(\mathbf{x}^0, y^0) \rangle = \frac{1}{Z} \int \frac{ds^0 d\bar{s}^0}{2\pi} \int \prod_{\mu=1}^P \frac{ds^\mu d\bar{s}^\mu}{2\pi} (y^0 - s^0)^2 \times e^{-\frac{\beta}{2} \sum_{\mu=1}^P (y^\mu - s^\mu)^2 + i \sum_{\mu=1}^P s^\mu \bar{s}^\mu + i s^0 \bar{s}^0} \times \left[1 + \frac{1}{\lambda N_1} \left(\sum_{\mu, \nu=1}^P \bar{s}^\mu K_{\mu\nu} \bar{s}^\nu + 2\bar{s}^0 \sum_{\mu=1}^P \bar{s}^\mu \kappa_\mu(\mathbf{x}^0) + (\bar{s}^0)^2 \kappa_0(\mathbf{x}^0) \right) \right]^{-\frac{N_1}{2}}, \quad (38)$$

where κ_μ and κ_0 are respectively the train-test and the test-test kernel integrals defined as in (23) when the covariance matrix involves the test input, namely:

$$\kappa_\mu = \int \frac{dt_1 dt_2}{\sqrt{(2\pi)^2 \det \tilde{C}_\mu}} e^{-\frac{1}{2} \mathbf{t}^\top \tilde{C}_\mu^{-1} \mathbf{t}} \sigma(t_1) \sigma(t_2), \quad (39)$$

$$\kappa_0 = \int \frac{dt}{\sqrt{2\pi C_{00}}} e^{-\frac{t^2}{2C_{00}}} \sigma(t)^2, \quad (40)$$

where

$$\tilde{C}_\mu = \begin{pmatrix} C_{\mu\mu} & C_{\mu 0} \\ C_{\mu 0} & C_{00} \end{pmatrix}, \quad C_{\mu 0} = \frac{1}{\lambda_1 N_0} \sum_{i_0}^{N_0} x_{i_0}^\mu x_{i_0}^0, \quad (41)$$

$$C_{00} = \frac{1}{\lambda_1 N_0} \sum_{i_0}^{N_0} (x_{i_0}^0)^2.$$

Now we can introduce the order parameters Q and \bar{Q} via Eq. (31) and their Fourier representation and perform the integration over all the s^μ , \bar{s}^μ and over the \bar{s}^0 . Doing so yields a single integral in s^0 and integrals on Q and \bar{Q} .

$$\langle \epsilon_g(\mathbf{x}^0, y^0) \rangle = \frac{1}{Z} \int \frac{dQ d\bar{Q}}{2\pi} e^{-\frac{N_1}{2} S(Q, \bar{Q})} \times \int \frac{ds^0 (y^0 - s^0)^2}{\sqrt{2\pi\sigma^2}} e^{-\frac{(s^0 - \Gamma)^2}{2\sigma^2}}, \quad (42)$$

with

$$\Gamma = -\frac{\bar{Q}}{\lambda} \sum_{\mu\nu} \kappa_\mu(\mathbf{x}^0) \left(\frac{1}{\beta} - \frac{\bar{Q}}{\lambda} K \right)_{\mu\nu}^{-1} y_\nu, \quad (43)$$

$$\sigma^2 = -\frac{\bar{Q}}{\lambda} \left[\kappa_0(\mathbf{x}^0) + \frac{\bar{Q}}{\lambda} \sum_{\mu\nu} \kappa_\mu(\mathbf{x}^0) \left(\frac{1}{\beta} - \frac{\bar{Q}}{\lambda} K \right)_{\mu\nu}^{-1} \kappa_\nu(\mathbf{x}^0) \right] \quad (44)$$

We can then unfold the easy integrals in s^0 and evaluate the result on the saddle point solution. Finally we get:

$$\langle \epsilon_g(\mathbf{x}^0, y^0) \rangle = (y^0 - \Gamma)^2 + \sigma^2. \quad (45)$$

It is worth noticing that the test error decomposes in the sum of two terms corresponding to the usual bias-variance decomposition of the generalisation error.

F. Exact solution of the saddle-point equations in the zero temperature limit

The saddle-point equations in (35) considerably simplify in the zero temperature limit ($\beta \rightarrow \infty$). In particular, using the fact that the kernel K has only positive eigenvalues (in the asymptotic regime α_1, α_0 finite), we get:

$$\bar{Q} = -\frac{1}{1+Q}, \quad Q = -\frac{\alpha_1}{\bar{Q}} - \frac{\alpha_1}{\bar{Q}^2} \frac{1}{P} y^T \left(\frac{K}{\lambda} \right)^{-1} y. \quad (46)$$

Given the condition $Q \geq -1$, the unique exact solution for \bar{Q} is non-positive and reads:

$$\bar{Q}^* = \frac{(\alpha_1 - 1) - \sqrt{(\alpha_1 - 1)^2 + 4\alpha_1 \frac{1}{P} y^T \left(\frac{K}{\lambda} \right)^{-1} y}}{2}. \quad (47)$$

In the zero temperature limit, we can also check that the average training error is identically zero, as it is expected since we are working in the heavily over-parametrised limit. The test error, on the other hand, is given by:

$$\langle \epsilon_g(\mathbf{x}^0, y^0) \rangle = \left[y^0 - \sum_{\mu\nu} \kappa_\mu(\mathbf{x}^0) K_{\mu\nu}^{-1} y_\nu \right]^2 - \frac{\bar{Q}^*}{\lambda} \left[\kappa_0(\mathbf{x}^0) - \sum_{\mu\nu} \kappa_\mu(\mathbf{x}^0) K_{\mu\nu}^{-1} \kappa_\nu(\mathbf{x}^0) \right]. \quad (48)$$

Notice that the dependence on α_1 (and thus on the hidden layer width) only influences the test error through \bar{Q}^* . More precisely, the bias part (the first term on the r.h.s. of Eq. (48)) is independent of α_1 and only depends on the training set (and thus implicitly on P, N_0 and possibly any additional parameter that defines the input/output probability distribution) and on the activation function of the network via the infinite-width kernel. The variance (i.e. the second term of r.h.s. of Eq. (48)), on the other hand, is proportional to $-\bar{Q}^*$ through a positive and training set dependent term.

Since the term proportional to the inverse kernel K is intensive and at fixed α_0 , \bar{Q}^* is a monotonically increasing/decreasing function of α_1 depending on whether $\frac{1}{P} y^T \left(\frac{K}{\lambda} \right)^{-1} y$ is smaller/larger than 1. That is, whether the output scale $P^{-1} y^T K^{-1} y$ is smaller/larger than the last layer initialisation variance λ^{-1} [31]. We conclude that the influence of the hidden layer width on the variance part of the test error is monotonic with a sign controlled by the ration between the output scale and the readout variance.

We also wish to stress that Eq. (48) allows to evaluate the generalisation error for any given set of data, and can thus be used to estimate optimal globally connected architectures for practical tasks on known datasets, at least in the regime of vanishing training error.

In Fig. 1 we compare the results of a numerical experiment (where the output y is provided by a linear teacher) with the theory. Numerical experiments are performed using stochastic gradient descent (SGD) with batches of small size and constant learning rate, since it is known [51] that such an algorithm can approximate Bayesian posterior inference. We have employed $\sigma(x) = \text{erf}(x)$ as activation function, since its NNGP kernel is known analytically [52] and with our notation it is given by:

$$K_{\mu\nu}^{\text{erf}}(C) = \frac{2}{\pi} \arcsin \left(\frac{2C_{\mu\nu}}{\sqrt{(1+2C_{\mu\mu})(1+2C_{\nu\nu})}} \right). \quad (49)$$

The agreement with the finite width theory is rather good for both the regimes of small and large variances in the last layer weights initialisation (left and center panel of Fig. 1 respectively), confirming the validity of our theory in the regime $P \sim N_0$.




FIG. 1. (Left and center panel) Test loss of a one hidden layer architecture (with erf activation) as a function of the training set size P . Both the training set and the test sample are drawn from a synthetic data distribution where the output is given by a linear teacher $y = \mathbf{t} \cdot \mathbf{x} / \sqrt{N_0}$ and \mathbf{x}, \mathbf{t} are i.i.d. random variables sampled from a normal distribution with zero mean and unit variance. Numerical experiments are compared with the theory at finite width for small ($1/\lambda = 1$) and large ($1/\lambda = \sqrt{3}$) variance of the initialisation of the weights of the last layer. (Right panel) Data collapse of the test curves (as a function of α_1) at different N_1, N_0 , but fixed aspect ratio N_1/N_0 . Data have been averaged over 50 different realisation of the initial weights at fixed instances of the dataset.

Furthermore, we note that in the case of a linear teacher no further extensive variables other than P, N_1 and N_0 may appear in Eq. (48); we conclude that the formula for the generalisation error only depends on α_0 and α_1 . This scaling prediction is numerically checked in the right panel Fig. 1 of by verifying the collapse of test curves (as a function of α_1) at different N_1, N_0 , but fixed aspect ratio α_1/α_0 .

In order to test our theory on a realistic dataset, we further performed a regression experiment [44] with the MNIST dataset. Our results, reported in Fig. 2, show the power of our theory in capturing the behavior of one-hidden layer networks at finite width. The grey band represents the variability obtained when we run experiments at different learning rates.

IV. GENERALISATION TO DEEP NEURAL NETWORKS WITH L HIDDEN LAYERS

In the following we will consider the more generic case of deep neural networks with L fully-connected hidden layers, as introduced in Section II.

We are interested in computing the partition function of the DNN. In the same spirit of the one hidden layer calculation, we introduce L sets of auxiliary variables $h_{i_\ell}^\mu$ (where $i_\ell = 1, \dots, N_\ell$) that are equal to the pre-activations at each layer. The strategy to perform the calculation is to show that the probability distribution of the pre-activations at each layer $P_\ell(\{h_{i_\ell}^\mu\})$ can be computed recursively, starting from the input layer. We notice that this is conceptually different from the backpropagating kernel renormalisation introduced in Ref. [31]. It is still a kernel renormalisation, but forward-propagating, and represents a generalisation to NNTPs of the kernel recurrence arising in NNGPs [9]. In practice, our approach amounts to a systematic, layer-by-layer description of the pre-activation statistics by the Student's t




FIG. 2. Test loss of a one-hidden layer neural network with N_1 hidden units and erf activation trained on a regression task with $P = 500$ patterns of the MNIST dataset. The theory (green) quantitatively captures the finite-width performance of the network. The infinite-width limit (orange line) is also reported for comparison. The grey band represents two standard deviations of the variability of the generalisation performance when we run experiments at different learning rates.

distribution that we have rigorously show to appear in the one hidden layer case. This can be seen as a quantitative correction to the standard Gaussian statistics that is recovered in the infinite width limit.

A. Iterative scheme for weights integration

Let us start by integrating the weights of the first layer. This defines a probability distribution over the

pre-activations of the first layer via:

$$P_1(\{h_{i_1}^\mu\}) = \int \mathcal{D}W^{(1)} \prod_{i_1, \mu} \delta \left(h_{i_1}^\mu - \frac{1}{\sqrt{N_0}} \sum_{i_0=1}^{N_0} W_{i_1 i_0}^{(1)} x_{i_0}^\mu \right) \\ = \prod_{i_1=1}^{N_1} \frac{e^{-\frac{1}{2} \sum_{\mu\nu} h_{i_1}^\mu C_{\mu\nu}^{-1} h_{i_1}^\nu}}{\sqrt{(2\pi)^P \det C}}. \quad (50)$$

This result is straightforward and it is valid for any N_0 , P and N_1 , since the prior for the weights is Gaussian. Notice that $C_{\mu\nu} = 1/(N_0\lambda_1) \sum_{i_0} x_{i_0}^\mu x_{i_0}^\nu$. At the second layer we have:

$$P_2(\{h_{i_2}^\mu\}) = \int \mathcal{D}W^{(2)} \mathcal{D}h_1 P_1(\{h_{i_1}^\mu\}) \\ \times \prod_{i_2, \mu} \delta \left(h_{i_2}^\mu - \frac{1}{\sqrt{N_1}} \sum_{i_1=1}^{N_1} W_{i_2 i_1}^{(2)} \sigma(h_{i_1}^\mu) \right) \quad (51)$$

We now introduce conjugate variables $\bar{h}_{i_2}^\mu$ to the activation of the second layer and the calculation proceeds as in the case of one hidden layer architectures. To make analytical progress we need to make two fundamental approximations: (i) assuming that the set of random variables $q_{i_2} = 1/(\sqrt{N_1}\lambda_2) \sum_{\mu} \bar{h}_{i_2}^\mu \sigma(h^\mu)$, where $\mathbf{h} \sim \mathcal{N}(0, C)$, is Gaussian-distributed; (ii) neglecting correlations between different pre-activations of the second hidden layer (technically this amounts to perform a first-order expansion of a determinant that appears in the calculation). In conclusion we get:

$$P_2(\{h_{i_2}^\mu\}) = \int dQ_1 d\bar{Q}_1 e^{-\frac{N_1}{2} (Q_1 \bar{Q}_1 + \log(1+Q_1))} \\ \times \prod_{i_2=1}^{N_2} \frac{e^{-\frac{1}{2} \sum_{\mu\nu} h_{i_2}^\mu (-\bar{Q}_1 K(C)/\lambda_2)_{\mu\nu}^{-1} h_{i_2}^\nu}}{\sqrt{(2\pi)^P \det(-\bar{Q}_1 K(C)/\lambda_2)}}, \quad (52)$$

where $K(C)$ is defined by Eq. (23) and $Q_\ell \geq -1$ for essentially the same reasons as in the one hidden layer case. Notice that except for the integration over the two variables Q_1 and \bar{Q}_1 , this is the same as the probability distribution of the one hidden layer system (21) if we replace C with $-\bar{Q}_1 K(C)/\lambda_2$. This reasoning can be easily iterated across layers and gives:

$$P_L(\{h_{i_L}^\mu\}) = \int \prod_{\ell=1}^{L-1} dQ_\ell d\bar{Q}_\ell e^{-\sum_{\ell=1}^{L-1} \frac{N_\ell}{2} [Q_\ell \bar{Q}_\ell + \log(1+Q_\ell)]} \\ \times \prod_{i_L=1}^{N_L} \frac{e^{-\frac{1}{2} \sum_{\mu\nu} h_{i_L}^\mu (K_{L-1}^{(R)}(\{\bar{Q}_\ell\}))_{\mu\nu}^{-1} h_{i_L}^\nu}}{\sqrt{(2\pi)^P \det(K_{L-1}^{(R)}(\{\bar{Q}_\ell\}))}}, \quad (53)$$

where $K_\ell^{(R)}(\{\bar{Q}_\ell\})$ is a renormalised kernel that satisfies the following recurrence relation:

$$K_\ell^{(R)}(\{\bar{Q}_\ell\}) = -\bar{Q}_\ell / \lambda_{\ell+1} K \circ [K_{\ell-1}^{(R)}(\{\bar{Q}_\ell\})], \quad K_0 = C, \quad (54)$$

and we stress that each $K_\ell^{(R)}$ depends on the variables $Q_1, \dots, Q_{\ell-1}$ only. On the basis of the comment we reported at the end of Sec. III B, we can say that this recursion is the generalisation to NNTPs of the one arising in the contexts of NNGPs [9], signal propagation in deep networks [53] and compositional kernels [54]. For completeness we stress that the recurrence relation for the infinite-width kernel K_L is given by Eq. (54) with $\bar{Q}_\ell = -1 \forall \ell = 1, \dots, L$.

In conclusion, the partition function for deep learning is given by a $2L$ -dimensional integral:

$$Z_{\text{DNN}} = \int \prod_{\ell=1}^L dQ_\ell d\bar{Q}_\ell e^{-\frac{N_\ell}{2} S_{\text{DNN}}(\{Q_\ell, \bar{Q}_\ell\})}, \quad (55)$$

where the effective deep learning action is given by:

$$S_{\text{DNN}} = \sum_{\ell=1}^L \frac{N_\ell}{N_L} [Q_\ell \bar{Q}_\ell + \log(1+Q_\ell)] \\ + \frac{1}{N_L} \text{Tr} \log \beta \left(\frac{\mathbb{1}}{\beta} + K_L^{(R)}(\{\bar{Q}_\ell\}) \right) \\ + \frac{1}{N_L} y^T \left(\frac{\mathbb{1}}{\beta} + K_L^{(R)}(\{\bar{Q}_\ell\}) \right)^{-1} y. \quad (56)$$

This action shares the same structure of the one found in section III for the special case of one hidden layer, with the difference that for L hidden layers, the recursive nature of the derivation introduces additional order parameters that are nested in the definition of the kernel K_L . Furthermore, since our derivation applies to layers of arbitrary size N_ℓ , the action also depends on the aspect ratios α_ℓ .

The derivative of the action w.r.t. the variables Q_ℓ is straightforward and gives:

$$\bar{Q}_\ell = -\frac{1}{1+Q_\ell} \quad \forall \ell = 1, \dots, L. \quad (57)$$

The derivation w.r.t. to the variables \bar{Q}_ℓ with $\ell \neq L$ is more complicated. For a generic kernel it does involve the application of the chain rule to the non-linear kernel function K_L . In the next sections we consider two cases where simplifications arise. These special cases correspond to kernels K such that $K(\alpha C) = \alpha^s K(C)$, where α is any positive scalar and $s \geq 0$ is an integer. It turns out that $s = 0$ is realized by the sign activation function, whereas $s = 1$ holds for piece-wise linear activations such as ReLU or Leaky-ReLU.

The computation of the generalisation error over a new example (\mathbf{x}^0, y^0) gives:

$$\langle \epsilon_g(\mathbf{x}^0, y^0) \rangle = (y^0 - \Gamma_L)^2 + \sigma_L^2 \quad (58)$$

where

$$\Gamma_L = \sum_{\mu\nu} \kappa_{L\mu}^{(R)} \left(\frac{\mathbb{1}}{\beta} + K_L^{(R)}(\{\bar{Q}_\ell\}) \right)_{\mu\nu}^{-1} y_\nu, \quad (59)$$

$$\sigma_L^2 = \kappa_{L0}^{(R)} - \sum_{\mu\nu} \kappa_{L\mu}^{(R)} \left(\frac{\mathbb{1}}{\beta} + K_L^{(R)}(\{\bar{Q}_\ell\}) \right)_{\mu\nu}^{-1} \kappa_{L\nu}^{(R)} \quad (60)$$

and $\kappa_{L\mu}^{(R)}$, $\kappa_{L0}^{(R)}$ are recursive kernels that generalise the train-test and test-test kernels (39)-(40). They are defined starting from Eq. (54) where the kernel K is now evaluated with the covariance matrix C involving train-test or test-test points. Note that L -hidden layers generalisation error is essentially found replacing the one hidden layer kernel with its recursive generalisation (54).

B. Saddle-point equations for scale independent kernels of the form $K(\alpha C) = K(C)$ ($\alpha > 0$)

In the case of sign activation function, it is straightforward to show that the behavior under scalar multiplication of the kernel $K_L^{(R)}(C)$ follows from the property $\text{sign}(\alpha x) = \text{sign}(x)$. It turns out that in this special case the effective action for deep learning considerably simplifies, since the non-linear dependence of $K_L^{(R)}$ on the variables $\{\bar{Q}_\ell\}_{\ell \neq L}$ disappears. This allows to solve the saddle-point equations exactly. In particular:

$$Q_\ell^* = 0, \quad \bar{Q}_\ell^* = -1 \quad \forall \ell = 1, \dots, L-1, \quad (61)$$

whereas the functional form of the solution for \bar{Q}_L is the same as in the one-hidden layer case (in the zero temperature limit):

$$\bar{Q}_L^* = \frac{(\alpha_L - 1) - \sqrt{(\alpha_L - 1)^2 + 4\alpha_L \frac{1}{P} y^T K_L^{-1} y}}{2}. \quad (62)$$

In practice, such a solution shows that deep architectures with sign activation (that are problematic to employ in practice since it is challenging to backpropagate derivatives) essentially behave as one hidden layer neural networks at finite width and the only marker of the depth L is retained in the infinite-width kernel K_L .

C. Saddle-point equations for piecewise linear kernels of the form $K(\alpha C) = \alpha K(C)$

The linear behavior of the kernel under scalar multiplication follows for ReLU and leaky ReLU activation function from the property $\text{ReLU}(\alpha x) = \alpha \text{ReLU}(x)$. It turns out that in this case the effective action for deep learning reads:

$$\begin{aligned} S_{\text{DNN}}(\{Q_\ell, \bar{Q}_\ell\}) &= \sum_{\ell=1}^L \frac{N_\ell}{N_L} [Q_\ell \bar{Q}_\ell + \log(1 + Q_\ell)] \\ &+ \frac{1}{N_L} \text{Tr} \log \beta \left(\frac{1}{\beta} + \left(\prod_{\ell=1}^L -\bar{Q}_\ell \right) K_L(C) \right) \\ &+ \frac{1}{N_L} y^T \left(\frac{1}{\beta} + \left(\prod_{\ell=1}^L -\bar{Q}_\ell \right) K_L(C) \right)^{-1} y. \quad (63) \end{aligned}$$

Exactly as for the one hidden layer case, the saddle-point equations simplify in the zero temperature limit

and under the assumption that the L -hidden layers kernel K_L has only positive eigenvalues:

$$-Q_\ell \bar{Q}_\ell - \alpha_\ell + \frac{\alpha_\ell}{(-\prod_{\ell_1} \bar{Q}_{\ell_1})} \frac{1}{P} y^T K_L^{-1} y = 0 \quad (64)$$

for all $\ell = 1, \dots, L$.

Notice that if $\alpha_\ell = \alpha$ for all $\ell = 1, \dots, L$, it is easy to show that the only solution must satisfy $Q_\ell^* = Q^*$ for all ℓ and we recover the heuristic mean field theory proposed in Ref. [31]. The reason for this equivalence is obvious: the authors of [31] found the heuristic mean field theory for ReLU activation by replacing the linear kernel with the corresponding NNGP kernel, noticing that the ReLU kernel transforms as the linear one under multiplication by a scalar. Our derivation shows that this replacement is not correct for general activation functions (see for instance the case of sign activation previously discussed), but it is possible in this particular case.

V. GENERALISING THE DEEP LEARNING EFFECTIVE ACTION TO NON-ODD ACTIVATION FUNCTIONS

In this section we show how the theory can be generalised in the case of non-odd activation functions. In fact, up to this point, our derivation assumed that the integral of the activation function over a Gaussian with zero mean is zero, i.e. the activation function is odd ($\sigma(x) = -\sigma(-x)$). The goal of this section is to show that removing such hypothesis modifies the effective action in the asymptotic limit. Since ReLU activation belongs to this more general case, the findings of this section imply that Li-Sompolinsky heuristic theory [31] should be modified as well. As for the rest of the manuscript, we start by considering one hidden layer architectures and we later extend the result to L hidden layers.

The crucial difference w.r.t. the case studied in section III is that if the activation function is not odd, the random variable

$$q = \frac{1}{\sqrt{N_1 \lambda}} \sum_{\mu=1}^P \bar{s}^\mu \sigma(h^\mu) \quad (65)$$

has now non-zero mean. In particular:

$$\langle q \rangle_{P(q)} = \frac{1}{\sqrt{N_1 \lambda}} \sum_{\mu=1}^P \bar{s}^\mu m^\mu, \quad (66)$$

$$m^\mu = \int \frac{dt}{\sqrt{2\pi C_{\mu\mu}}} e^{-t^2/(2C_{\mu\mu})} \sigma(t). \quad (67)$$

A straightforward calculation shows that the result for non-odd activations is found by performing the replace-

ment:

$$-\frac{\bar{Q}}{\lambda}K \rightarrow K^{(R)}(Q, \bar{Q}) = -\frac{\bar{Q}}{\lambda}K + \frac{(\bar{Q} + \frac{1}{1+\bar{Q}})}{\lambda}K^{(1)}, \quad (68)$$

$$K_{\mu\nu}^{(1)} = m^\mu m^\nu \quad (69)$$

in the effective action given in Eq. (33). As such, the one-hidden layer action for non-odd activation functions reads:

$$\begin{aligned} S_{\text{1HL}} &= Q\bar{Q} + \log(1+Q) + \\ &+ \frac{1}{N_1} \text{Tr} \log \beta \left[\frac{\mathbb{1}}{\beta} + K^{(R)}(Q, \bar{Q}) \right] + \\ &+ \frac{1}{N_1} y^\top \left[\frac{\mathbb{1}}{\beta} + K^{(R)}(Q, \bar{Q}) \right]^{-1} y \end{aligned} \quad (70)$$

It is worth noticing that while in the zero mean case there was a simple relations between Q and \bar{Q} at any temperature, we now lose that property and the saddle-point equations are not exactly solvable anymore, not even in the zero temperature limit. On the contrary, one can check that in the infinite-width limit we recover the previous result $\bar{Q} = -1$, $Q = 0$ and the rank one matrix $K^{(1)}$ does not contribute to the generalisation error, since it does always appear in combination with the scalar $\bar{Q} + 1/(1+Q)$ that vanishes in the infinite-width limit.

The generalisation error can be written in terms of the renormalised kernel $K^{(R)}(Q, \bar{Q})$:

$$\begin{aligned} \langle \epsilon_g(\mathbf{x}^0, y^0) \rangle &= \left\{ y^0 - \sum_{\mu\nu} \kappa_\mu^{(R)} [(K^{(R)})^{-1}]_{\mu\nu} y_\nu, \right\}^2 \\ &+ \left\{ \kappa_0^{(R)} - \sum_{\mu\nu} \kappa_\mu^{(R)} [(K^{(R)})^{-1}]_{\mu\nu} \kappa_\nu^{(R)} \right\}, \end{aligned} \quad (71)$$

where $\kappa_\mu^{(R)}(\mathbf{x}^0)$ and $\kappa_0^{(R)}(\mathbf{x}^0)$ are computed on the test set input \mathbf{x}^0 as done in Section III. Differently from the case of odd activation function, now the test error depends on both variables Q and \bar{Q} , which are determined by solving the corresponding saddle-point equations.

Let us move to the derivation of the effective action for L hidden layers. As for the derivation with odd activation function, the key step is to understand how the joint probability of the pre-activations at layer ℓ is linked to the one at layer $\ell - 1$. While in the odd activation case, the key observation was that P_2 is related to P_1 by the replacement $C \rightarrow -\bar{Q}_1 K(C)/\lambda_2$ (see Eq. (52)), here we find that the correct replacement is $C \rightarrow -\bar{Q}_1 K(C)/\lambda_2 + (\bar{Q}_1 + 1/(1+Q_1))K^{(1)}/\lambda_2$. Differently from the odd activation case, where the kernel at layer L was only depending on the variables $\{Q_\ell\}$, here we find that the recurrence is given in terms of the $\{Q_\ell\}$ as well. In conclusion, this produces a more unpleasant action where all the $\{Q_\ell, \bar{Q}_\ell\}$ are coupled via the nested non-linear expression of the kernel. The explicit recurrence relation for non-odd activation functions is given

by:

$$\begin{aligned} K_\ell^{(R)} &= -\frac{\bar{Q}_\ell}{\lambda_{\ell+1}}K \circ [K_{\ell-1}^{(R)}] + \frac{(\bar{Q}_\ell + \frac{1}{1+\bar{Q}_\ell})}{\lambda_{\ell+1}}K^{(1)} \circ [K_{\ell-1}^{(R)}] \\ K_0^{(R)} &= C \end{aligned} \quad (72)$$

and the effective saddle-point action reads:

$$\begin{aligned} S_{\text{DNN}} &= \sum_{\ell=1}^L \frac{N_\ell}{N_L} [Q_\ell \bar{Q}_\ell + \log(1+Q_\ell)] \\ &+ \frac{1}{N_L} \text{Tr} \log \beta \left(\frac{\mathbb{1}}{\beta} + K_L^{(R)}(\{Q_\ell, \bar{Q}_\ell\}) \right) \\ &+ \frac{1}{N_L} y^T \left(\frac{\mathbb{1}}{\beta} + K_L^{(R)}(\{Q_\ell, \bar{Q}_\ell\}) \right)^{-1} y. \end{aligned} \quad (73)$$

In view of the above considerations, it should be now clear that the heuristic Li-Sompolinsky theory (re-derived in the previous section) amounts to disregard all the additional terms $K^{(1)}$ that arise from the approach presented in this section. We leave the numerical investigation of this more complex finite-width theory for a future work. Here we derive the first perturbative correction to the infinite-width limit ($\alpha_\ell = 0$) of Eq. (73). This computation provides an intriguing prediction for the finite-width performance of deep networks that can be checked in experiments. It is worth pointing out that the regime in which real-world networks work is the finite-width one, thus this examination provides insight in a practically relevant setting.

A straightforward Taylor expansion around the infinite-width limit $\alpha_\ell = \alpha = 0 \forall \ell = 1, \dots, L$ shows that the first correction to the test loss $\Delta\epsilon_g$ is proportional to:

$$\Delta\epsilon_g \propto \alpha \left(\frac{1}{P} y^T K_L^{-1} y - 1 \right). \quad (74)$$

This means that there exists a simple scalar observable that determines whether the finite-width deep neural network will outperform its infinite-width counterpart:

$$s_L = \frac{1}{P} y^T K_L^{-1} y. \quad (75)$$

In particular, we expect the finite-width network to outperform its infinite-width counterpart whenever $s_L < 1$. In panel (a) of Fig. V we check this prediction for deep architectures with ReLU activation on a regression task with the MNIST dataset. In the experiment we set $P = 500$ and $N_\ell = N = 3000$, so that we are close to the infinite-width limit and we consider the test loss (generalisation error) as a function of the number of hidden layers L . For $L < 5$, $s_L < 1$ and the finite-width network improves the infinite-width prediction, whereas for $L \geq 5$, $s_L > 1$ and the correction to the infinite-width performance is positive. Notice that s_L quickly diverges to infinity as the number of hidden layers L grows. The




FIG. 3. (a) Generalisation error of a deep neural network with L hidden layers and ReLU activation trained on a regression task with $P = 500$ patterns from the MNIST dataset (the size of the hidden layers is $N_\ell = N = 3000$) [56]. The plot shows that there exists a simple criterion that determines whether the finite-width network will outperform its infinite-width counterpart. As long as $s_L = y^T K_L^{-1} y / P$ is smaller than one (left of the vertical dashed line), we expect it to be convenient to employ finite-width architectures. In this particular case, this occurs for $L < 5$. The grey band represents three standard deviations w.r.t. experiments with different learning rates. (b) Behaviour of the ReLU kernel as the number of layers L is increased. The figures represent the matrix K_L at different values of L starting from $P = 5$ randomly chosen MNIST datapoints and iterating the recurrence (72) at infinite width ($Q_\ell = 0, \bar{Q}_\ell = -1$) using the analytical form of the ReLU kernel (see, for example, [55]).

reason for this is simply that the ReLU NNGP kernel K_L develops at least one zero eigenvalue as $L \rightarrow \infty$. This ultimately occurs because each element of the matrix K_L converges to 1 as L grows (see panel (b) of Fig. V), as one can easily check by looking at the explicit recurrence relation for the ReLU kernel [55].

VI. DISCUSSION

In our work we have described a strategy to investigate the statistical mechanics of deep neural networks beyond the infinite-width limit, that is in the finite asymptotic regime $P, N_\ell \rightarrow \infty$ at $\alpha_\ell = P/N_\ell > 0$ as opposed to the infinite-width $\alpha_\ell = 0$. In the one-hidden layer case, our evaluation of the saddle-point action is exact in the above thermodynamic limit. As such, we do not expect any additional corrections to the result, at least in the asymptotic regime. In particular, we have found a closed expression for the generalisation error that in principle provides a Bayesian estimator of the generalisation capabilities of fully-connected architectures for any given

empirical dataset, provided that the chosen architecture is capable of perfectly fitting the training set (vanishing training error).

For the case of arbitrarily deep networks, we are not able to formally prove that our result is exact. However, since the key assumption to perform the evaluation is given, it should be possible, at least in principle, to take systematically into account non-Gaussian corrections to the saddle-point action to check whether these are relevant or not for the theory at finite width [57]. Our approximated theory, however, can still be useful and instructive in various contexts: it provides us with a simple analytical tool to predict whether finite-width networks with ReLU activation functions will achieve better test accuracy than infinite-width ones. In the future we plan to investigate whether a similar criterion can be found for other commonly used activation functions. It will also be of great interest to understand to what extent the more general theory presented in section V captures the learning curves of deep architectures. We are currently working on a numerical routine capable of finding the solutions of the $2L$ coupled saddle-point equations that arise in the case non-odd activation functions.

From the mathematical perspective, our results can be interpreted as a generalisation to Student's t processes of the NNGPs arising in the infinite-width limit of Bayesian neural networks. We find this observation very promising. The precise characterization of this mapping and its limits of validity represent a research line for future investigation.

Notably, our theory predicts that a kernel limit should also appear at finite width and depth L that approaches infinity. This could be checked, for instance, considering the isotropic limit $\alpha_\ell = \alpha \forall \ell$ and ReLU activation. Here one can numerically solve the saddle-point equation for \bar{Q} at large L and verify that $\bar{Q} \rightarrow -1$ for all α , as shown in Fig. 4 (right panel). As such, also in this limit we expect an equivalence with a kernel theory with kernel given by $K_\infty(C)$.

It is fair to stress that our theory only describes the equilibrium regime of zero train loss, so that our analysis should not apply in the regime $P/N_1 \gg 1$. Interestingly, numerical simulations performed with one hidden layer architectures of varying width and random training labels show that the train loss follows a universal behavior [58] w.r.t. α_1 (see Fig. 4, left panel) also in the regime where the DNN is not capable to perfectly fit the data. It would be desirable to develop a theory that also captures this phase.

Another interesting aspect to understand is the degree to which this mean-field static analysis can be extended beyond equilibrium in order to assess the full training dynamics; such a theory would indeed make it possible to investigate the performance of the many (often heuristics) learning algorithms employed to train deep neural networks.

We conclude by pointing out that our theory can be extended to convolutional layers of weights, replacing neu-




FIG. 4. (Left panel) Training loss of different one hidden layer architectures trained on a completely random task (i.e. both the inputs x and outputs y are i.i.d. random variables sampled from a normal distribution with zero mean and unit variance) as a function of α_1 . At the moment we can not capture this universal phenomenon with our theory, which only describes the overparametrised limit where the training error is exactly zero. (Right panel) The numerical evaluation of the solution \bar{Q} is shown in the case of ReLU activation function and isotropic network $\alpha_\ell = \alpha \forall \ell$, for different depths L . The parameter \bar{Q} quickly approaches -1 for all α , suggesting that also finite-width DNNs converge to a kernel limit if the depth $L \rightarrow \infty$.

rons with channels, as done in the infinite-width case in Ref. [11].

ACKNOWLEDGEMENTS

M. P. is supported by a grant from the Simons Foundation (grant No. 454941, S. Franz). P. R. acknowledges funding from the Fellini program under the H2020-MSCA-COFUND action, Grant Agreement No. 754496, INFN (IT). The authors would like to thank Silvio Franz,

Luca Molinari, Fabian Aguirre-López and the Computing Sciences group at Bocconi University in Milan for discussions and suggestions.

DATA AND CODE AVAILABILITY

Data and code employed are publicly available at https://github.com/rosalba-p/deepL_beyond_infwidth [56].

-
- [1] A. Krizhevsky, I. Sutskever, and G. E. Hinton, Imagenet classification with deep convolutional neural networks, *Commun. ACM* **60**, 84–90 (2017).
 - [2] I. Goodfellow, Y. Bengio, and A. Courville, *Deep Learning* (MIT Press, 2016).
 - [3] A. Engel and C. Van den Broeck, *Statistical Mechanics of Learning* (Cambridge University Press, 2001).
 - [4] C. Zhang, S. Bengio, M. Hardt, B. Recht, and O. Vinyals, Understanding deep learning (still) requires rethinking generalization, *Commun. ACM* **64**, 107–115 (2021).
 - [5] Y. Bahri, J. Kadmon, J. Pennington, S. S. Schoenholz, J. Sohl-Dickstein, and S. Ganguli, Statistical mechanics of deep learning, *Annual Review of Condensed Matter Physics* **11**, 501 (2020).
 - [6] R. M. Neal, Priors for infinite networks, in *Bayesian Learning for Neural Networks* (Springer New York, New York, NY, 1996) pp. 29–53.
 - [7] C. Williams, Computing with infinite networks, in *Advances in Neural Information Processing Systems*, Vol. 9, edited by M. Mozer, M. Jordan, and T. Petsche (MIT Press, 1996).
 - [8] A. G. de G. Matthews, J. Hron, M. Rowland, R. E. Turner, and Z. Ghahramani, Gaussian process behaviour in wide deep neural networks, in *International Conference on Learning Representations* (2018).
 - [9] J. Lee, J. Sohl-dickstein, J. Pennington, R. Novak, S. Schoenholz, and Y. Bahri, Deep neural networks as gaussian processes, in *International Conference on Learning Representations* (2018).
 - [10] A. Garriga-Alonso, C. E. Rasmussen, and L. Aitchison, Deep convolutional networks as shallow gaussian processes, in *International Conference on Learning Representations* (2019).
 - [11] R. Novak, L. Xiao, Y. Bahri, J. Lee, G. Yang, D. A. Abolafia, J. Pennington, and J. Sohl-dickstein, Bayesian deep convolutional networks with many channels are gaussian processes, in *International Conference on Learning Representations* (2019).

- [12] A. Jacot, F. Gabriel, and C. Hongler, Neural tangent kernel: Convergence and generalization in neural networks, in *Advances in Neural Information Processing Systems*, Vol. 31, edited by S. Bengio, H. Wallach, H. Larochelle, K. Grauman, N. Cesa-Bianchi, and R. Garnett (Curran Associates, Inc., 2018).
- [13] L. Chizat, E. Oyallon, and F. Bach, On lazy training in differentiable programming, in *Advances in Neural Information Processing Systems*, Vol. 32, edited by H. Wallach, H. Larochelle, A. Beygelzimer, F. d'Alché-Buc, E. Fox, and R. Garnett (Curran Associates, Inc., 2019).
- [14] J. Lee, L. Xiao, S. Schoenholz, Y. Bahri, R. Novak, J. Sohl-Dickstein, and J. Pennington, Wide neural networks of any depth evolve as linear models under gradient descent, in *Advances in Neural Information Processing Systems*, Vol. 32, edited by H. Wallach, H. Larochelle, A. Beygelzimer, F. d'Alché-Buc, E. Fox, and R. Garnett (Curran Associates, Inc., 2019).
- [15] B. Bordelon, A. Canatar, and C. Pehlevan, Spectrum dependent learning curves in kernel regression and wide neural networks, in *Proceedings of the 37th International Conference on Machine Learning*, Proceedings of Machine Learning Research, Vol. 119, edited by H. D. III and A. Singh (PMLR, 2020) pp. 1024–1034.
- [16] A. Canatar, B. Bordelon, and C. Pehlevan, Spectral bias and task-model alignment explain generalization in kernel regression and infinitely wide neural networks, *Nature communications* **12**, 1 (2021).
- [17] R. Dietrich, M. Opper, and H. Sompolinsky, Statistical mechanics of support vector networks, *Phys. Rev. Lett.* **82**, 2975 (1999).
- [18] C. Cortes and V. Vapnik, Support-vector networks, *Machine Learning* **20**, 273 (1995).
- [19] M. Seleznova and G. Kutyniok, Neural tangent kernel beyond the infinite-width limit: Effects of depth and initialization, arXiv preprint arXiv:2206.00553 (2022).
- [20] N. Vyas, Y. Bansal, and N. Preetum, Limitations of the ntk for understanding generalization in deep learning, arXiv preprint arXiv:2206.10012 (2022).
- [21] Y. Bengio and O. Delalleau, On the expressive power of deep architectures, in *International conference on algorithmic learning theory* (Springer, 2011) pp. 18–36.
- [22] P. L. Bartlett, N. Harvey, C. Liaw, and A. Mehrabian, Nearly-tight vc-dimension and pseudodimension bounds for piecewise linear neural networks, *The Journal of Machine Learning Research* **20**, 2285 (2019).
- [23] P. Rotondo, M. C. Lagomarsino, and M. Gherardi, Counting the learnable functions of geometrically structured data, *Phys. Rev. Research* **2**, 023169 (2020).
- [24] P. Rotondo, M. Pastore, and M. Gherardi, Beyond the storage capacity: Data-driven satisfiability transition, *Phys. Rev. Lett.* **125**, 120601 (2020).
- [25] M. Pastore, P. Rotondo, V. Erba, and M. Gherardi, Statistical learning theory of structured data, *Phys. Rev. E* **102**, 032119 (2020).
- [26] M. Gherardi, Solvable model for the linear separability of structured data, *Entropy* **23**, 10.3390/e23030305 (2021).
- [27] M. Pastore, Critical properties of the SAT/UNSAT transitions in the classification problem of structured data, *Journal of Statistical Mechanics: Theory and Experiment* **2021**, 113301 (2021).
- [28] F. Aguirre-López, M. Pastore, and S. Franz, Satisfiability transition in asymmetric neural networks, *Journal of Physics A: Mathematical and Theoretical* **55**, 305001 (2022).
- [29] A. M. Saxe, J. L. McClelland, and S. Ganguli, Exact solutions to the nonlinear dynamics of learning in deep linear neural networks, in *2nd International Conference on Learning Representations, ICLR 2014, Banff, AB, Canada, April 14-16, 2014, Conference Track Proceedings*, edited by Y. Bengio and Y. LeCun (2014).
- [30] A. M. Saxe, J. L. McClelland, and S. Ganguli, A mathematical theory of semantic development in deep neural networks, *Proceedings of the National Academy of Sciences* **116**, 11537 (2019), <https://www.pnas.org/doi/pdf/10.1073/pnas.1820226116>.
- [31] Q. Li and H. Sompolinsky, Statistical mechanics of deep linear neural networks: The backpropagating kernel renormalization, *Phys. Rev. X* **11**, 031059 (2021).
- [32] J. Zavatore-Veth, A. Canatar, B. Ruben, and C. Pehlevan, Asymptotics of representation learning in finite bayesian neural networks, in *Advances in Neural Information Processing Systems*, Vol. 34, edited by M. Ranzato, A. Beygelzimer, Y. Dauphin, P. Liang, and J. W. Vaughan (Curran Associates, Inc., 2021) pp. 24765–24777.
- [33] G. Naveh and Z. Ringel, A self consistent theory of gaussian processes captures feature learning effects in finite cnns, in *Advances in Neural Information Processing Systems*, Vol. 34, edited by M. Ranzato, A. Beygelzimer, Y. Dauphin, P. Liang, and J. W. Vaughan (Curran Associates, Inc., 2021) pp. 21352–21364.
- [34] I. Seroussi and Z. Ringel, Separation of scales and a thermodynamic description of feature learning in some cnns, arXiv preprint arXiv:2112.15383 (2021).
- [35] J. A. Zavatore-Veth, W. L. Tong, and C. Pehlevan, Contrasting random and learned features in deep bayesian linear regression, *Phys. Rev. E* **105**, 064118 (2022).
- [36] P. Breuer and P. Major, Central limit theorems for nonlinear functionals of gaussian fields, *Journal of Multivariate Analysis* **13**, 425 (1983).
- [37] I. Nourdin, G. Peccati, and M. Podolskij, *Quantitative Breuer-Major theorems* (2010).
- [38] F. Gerace, B. Loureiro, F. Krzakala, M. Mezard, and L. Zdeborova, Generalisation error in learning with random features and the hidden manifold model, in *Proceedings of the 37th International Conference on Machine Learning*, Proceedings of Machine Learning Research, Vol. 119, edited by H. D. III and A. Singh (PMLR, 2020) pp. 3452–3462.
- [39] B. Loureiro, C. Gerbelot, H. Cui, S. Goldt, F. Krzakala, M. Mezard, and L. Zdeborová, Learning curves of generic features maps for realistic datasets with a teacher-student model, *Advances in Neural Information Processing Systems* **34** (2021).
- [40] S. Goldt, B. Loureiro, G. Reeves, F. Krzakala, M. Mézard, and L. Zdeborová, The gaussian equivalence of generative models for learning with shallow neural networks, arXiv preprint arXiv:2006.14709 (2020).
- [41] E. Dobriban and S. Wager, High-dimensional asymptotics of prediction: ridge regression and classification, *The Annals of Statistics* **46**, 247 (2018).
- [42] S. Mei and A. Montanari, The generalization error of random features regression: Precise asymptotics and the double descent curve, *Communications on Pure and Applied Mathematics* (2019).
- [43] B. Ghorbani, S. Mei, T. Misiakiewicz, and A. Montanari, Linearized two-layers neural networks in high dimension,

- The Annals of Statistics **49**, 1029 (2021).
- [44] S. Ariosto, R. Pacelli, F. Ginelli, M. Gherardi, and P. Rotondo, Universal mean-field upper bound for the generalization gap of deep neural networks, *Phys. Rev. E* **105**, 064309 (2022).
- [45] A. Shah, A. Wilson, and Z. Ghahramani, Student-t Processes as Alternatives to Gaussian Processes, in *Proceedings of the Seventeenth International Conference on Artificial Intelligence and Statistics*, Proceedings of Machine Learning Research, Vol. 33, edited by S. Kaski and J. Corander (PMLR, Reykjavik, Iceland, 2014) pp. 877–885.
- [46] J. Lee, S. Schoenholz, J. Pennington, B. Adlam, L. Xiao, R. Novak, and J. Sohl-Dickstein, Finite versus infinite neural networks: an empirical study, in *Advances in Neural Information Processing Systems*, Vol. 33, edited by H. Larochelle, M. Ranzato, R. Hadsell, M. Balcan, and H. Lin (Curran Associates, Inc., 2020) pp. 15156–15172.
- [47] A. C. C. Coolen, M. Sheikh, A. Mozeika, F. Aguirre-López, and F. Antenucci, Replica analysis of overfitting in generalized linear regression models, *Journal of Physics A: Mathematical and Theoretical* **53**, 365001 (2020).
- [48] A. Mozeika, M. Sheikh, F. Aguirre-López, F. Antenucci, and A. C. C. Coolen, Exact results on high-dimensional linear regression via statistical physics, *Phys. Rev. E* **103**, 042142 (2021).
- [49] Y. Uchiyama, H. Oka, and A. Nono, Student’s t-process regression on the space of probability density functions, *Proceedings of the ISCIE International Symposium on Stochastic Systems Theory and its Applications* **2021**, 1 (2021).
- [50] H. Lee, E. Yun, H. Yang, and J. Lee, Scale mixtures of neural network gaussian processes, in *International Conference on Learning Representations* (2022).
- [51] S. Mandt, M. D. Hoffman, and D. M. Blei, Stochastic gradient descent as approximate bayesian inference, *Journal of Machine Learning Research* **18**, 1 (2017).
- [52] G. Pang, L. Yang, and G. E. Karniadakis, Neural-net-induced gaussian process regression for function approximation and pde solution, *Journal of Computational Physics* **384**, 270 (2019).
- [53] B. Poole, S. Lahiri, M. Raghu, J. Sohl-Dickstein, and S. Ganguli, Exponential expressivity in deep neural networks through transient chaos, in *Advances in Neural Information Processing Systems*, Vol. 29, edited by D. Lee, M. Sugiyama, U. Luxburg, I. Guyon, and R. Garnett (Curran Associates, Inc., 2016).
- [54] Y. Cho and L. Saul, Kernel methods for deep learning, in *Advances in Neural Information Processing Systems*, Vol. 22, edited by Y. Bengio, D. Schuurmans, J. Lafferty, C. Williams, and A. Culotta (Curran Associates, Inc., 2009).
- [55] A. Bietti and J. Mairal, On the inductive bias of neural tangent kernels, in *Advances in Neural Information Processing Systems*, Vol. 32, edited by H. Wallach, H. Larochelle, A. Beygelzimer, F. d'Alché-Buc, E. Fox, and R. Garnett (Curran Associates, Inc., 2019).
- [56] https://github.com/rosalba-p/deepL_beyond_infwidth, (GitHub, 2023).
- [57] D. A. Roberts, S. Yaida, and B. Hanin, *The Principles of Deep Learning Theory* (Cambridge University Press, 2022) <https://deeplearningtheory.com>, arXiv:2106.10165 [cs.LG].
- [58] F. Gerace, F. Krzakala, B. Loureiro, L. Stephan, and L. Zdeborová, Gaussian universality of linear classifiers with random labels in high-dimension, arXiv preprint arXiv:2205.13303 (2022).